Broad and differential animal ACE2 receptor usage by SARS-CoV-2

Xuesen Zhao[1, 2#]*, Danying Chen[1, 2#], Robert Szabla[3], Mei Zheng[1, 2], Guoli Li[1, 2],

Pengcheng Du[1, 2], Shuangli Zheng[1, 2], Xinglin Li[1, 2], Chuan Song[1, 2], Rui Li[1, 2], Ju-Tao

Guo[4], Murray Junop[3], Hui Zeng[1, 2]*, Hanxin Lin[5]*

[1]Institute of Infectious disease, Beijing Ditan Hospital, Capital Medical University,

Beijing 100015, China.

[2]Beijing Key Laboratory of Emerging Infectious Disease, Beijing 100015, China.

[3]Department of Biochemistry, Western University, 1151 Richmond Street, London,

Ontario, Canada.

[4]Baruch S. Blumberg Institute, Hepatitis B Foundation, 3805 Old Easton Road,

Doylestown, PA 18902. USA.

[5]Department of Pathology and Laboratory Medicine, Western University, 1151

Richmond Street, London, Ontario, Canada.

Running Title: Broad animal ACE2 receptor usage by SARS-CoV-2

* Corresponding author's e-mail address:

Xuesen Zhao, zhaoxuesen@ccmu.edu.cn

Hui Zeng, zenghui@ccmu.edu.cn

Hanxin Lin, hanxin.lin@lhsc.on.ca

[#] These authors are co-first authors.

ABSTRACT

The COVID-19 pandemic has caused an unprecedented global public health and

economy crisis. The origin and emergence of its causal agent, SARS-CoV-2, in the

human population remains mysterious, although bat and pangolin were proposed to be

the natural reservoirs. Strikingly, comparing to the SARS-CoV-2-like CoVs identified in

bats and pangolins, SARS-CoV-2 harbors a polybasic furin cleavage site in its spike (S)

glycoprotein. SARS-CoV-2 uses human ACE2 as its receptor to infect cells. Receptor

recognition by the S protein is the major determinant of host range, tissue tropism, and

pathogenesis of coronaviruses. In an effort to search for the potential intermediate or

amplifying animal hosts of SARS-CoV-2, we examined receptor activity of ACE2 from

14 mammal species and found that ACE2 from multiple species can support the

infectious entry of lentiviral particles pseudotyped with the wild-type or furin cleavage

site deficient S protein of SARS-CoV-2. ACE2 of human/rhesus monkey and rat/mouse

exhibited the highest and lowest receptor activity, respectively. Among the remaining

species, ACE2 from rabbit and pangolin strongly bound to the S1 subunit of

SARS-CoV-2 S protein and efficiently supported the pseudotyped virus infection. These

findings have important implications for understanding potential natural reservoirs,

zoonotic transmission, human-to-animal transmission, and use of animal models.

Key words: SARS-CoV-2; animal ACE2; receptor; entry; furin cleavage; animal hosts

Introduction

Coronavirus disease 2019 (COVID-19) was first identified in Dec. 2019 in the city of Wuhan, China [1], and has since spread worldwide, causing ~2.3 million infected and around 160,000 fatalities as of April 18th, 2020 (https://coronavirus.jhu.edu/map.html). These numbers are still growing rapidly. The global COVID-19 pandemic has caused an unprecedented public health and economy crisis.

COVID-19 is caused by a novel coronavirus, Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2; initially named as 2019-nCoV) [2,3]. The origin of SARS-CoV-2 and its emergence in the human population remain mysterious. Many of the early cases were linked to the Huanan seafood and wild animal market in Wuhan city, raising the possibility of zoonotic origin [4]. Sequencing analyses showed that the genome of SARS-CoV-2 shares 79.5%, 89.1%, 93.3%, and 96.2% nucleotide sequence identity with that of human SARS-CoV, bat coronavirus (CoV) ZC45, bat CoV RmYN02, and bat CoV RaTG13, respectively, suggesting that SARS-CoV-2 probably has bat origins [2,3,5]. This finding is not surprising as bats are notorious for serving as the natural reservoir for many emerging zoonotic viral pathogens, including two other deadly human coronaviruses, SARS-CoV and Middle East respiratory syndrome coronavirus (MERS-CoV), which caused global outbreak during 2002-2003 and 2012-2015, respectively [6,7].

Although SARS-CoV-2 may have originated from bats, bat CoVs are unlikely to jump directly to humans due to the general ecological separation. Other mammal species may have been served as intermediate or amplifying hosts where the progenitor

virus acquires critical mutations for efficient zoonotic transmission to human. This has

been seen in the emergence of SARS-CoV and MERS-CoV where palm civet and

dromedary camel act as the respective intermediate host [7]. The Huanan seafood and

wild animal market in Wuhan city would otherwise be a unique place to trace any

potential animal source; however, soon after the disease outbreak, the market was

closed and all the wild animals were cleared, making this task very challenging or even

impossible. As an alternative, wide screening of wild animals becomes imperative.

Several recent studies identified multiple SARS-COV-2-like CoVs (SL-CoVs) from

smuggled Malayan pangolins in China. These pangolin CoVs (PCoV) form two

phylogenetic lineages, PCoV-GX and PCoV-GD [8-11]. In particular, lineage PCoV-GD

was found to carry a nearly identical receptor-binding motif (RBM) in the spike (S)

protein to that of SARS-CoV-2 (Fig.1). However, the genome of these pangolin

SL-CoVs share only 85.5%-92.4% nucleotide identities with that of SARS-CoV-2. This

is in contrast to SARS-CoV and MERS-CoV where CoVs isolated form the

intermediate host palm civet and dromedary camel share 99.6% and 99.9% % genome

sequence identities with their human counterpart, respectively [12,13]. Therefore, pangolins

tested in these studies are not the direct intermediate host for SARS-CoV-2. Whether or

not SARS-CoV-2 came from other pangolins or other wild animal species remains to be

determined.

Receptor recognition by the viral S protein is the major determinant of host range,

cell, tissue tropism, and pathogenesis of coronaviruses [14]. The S protein of

SARS-CoV-2 is a type I membrane glycoprotein, which can be cleaved to S1 and S2

subunit during biogenesis at the polybasic furin cleavage site (RRAR) (Fig.1) [15-17].

Previous studies have shown that furin cleavage is not essential for coronavirus-cell

membrane fusion, but enhances cell-to-cell fusion [18-22], increases the fitness of sequence

variant within the quasispecies population of bovine CoV [23], and convert avirulent avian

influenza virus isolate to a highly pathogenic isolate [24]. Interestingly, this cleavage site

is not present in the S protein of SARS-CoV, bat SL-CoVs or pangolin SL-CoVs

identified so far [5,15]. During cell entry, S1 binds to the cellular receptor, subsequently

triggering a cascade of events leading to S2-mediated membrane fusion between host

cells and coronavirus particles [25]. S1 protein contains an independently folded domain

called the receptor binding domain (RBD), which harbors an RBM that is primarily

involved in contact with receptor (Fig. 1). Human ACE2 (hACE2) has been identified as

the cellular receptor for both SARS-CoV-2 [3,15,17,26] and SARS-CoV [27]. In addition to

hACE2, ACE2 from horseshoe bat (*Rhinolophus alcyone*) was found to support cell

entry of SARS-CoV-2 S-mediated VSV-based pseudotyped virus [15]. By using infectious

virus it has also been shown that ACE2 from Chinese horseshoe bat (*Rhinolophus*

sinicus), civet and swine, but not mouse, could serve as functional receptors [3]. However,

in this infection system, the entry step was coupled with other steps during virus life

cycle, i.e. viral genome replication, translation, virion assembly and budding, and thus

the receptor activity of these animal ACE2 orthologs were not directly investigated.

In an effort to search for potential animal hosts, we examined the receptor activity

of ACE2 from 14 mammal species, including human, rhesus monkey, Chinese

horseshoe bat (Rs bat), Mexican free-tailed bat (Tb bat), rat, mouse, palm civet, raccoon

dog, ferret badger, hog badger, canine, feline, rabbit, and pangolin for SARS-CoV-2 and

a mutant virus lacking the furin cleavage site in the S protein. Our results show that

multiple animal ACE2 could serve as receptors for SARS-CoV-2 and the SARS-CoV-2

mutant. ACE2 of human/rhesus monkey and rat/mouse exhibited the highest and lowest

receptor activity, respectively, with the other 10 ACE2s exhibiting intermediate activity.

The implications of our findings were discussed in terms of the natural reservoir,

zoonotic transmission, human-to-animal transmission, animal health, and animal model.

Results

Human ACE2 serves as a functional receptor for SARS-CoV-2

To examine the receptor activity of human ACE2 (hACE2) for SARS-CoV-2, we

first established a HIV-based pseudotyped virus entry system. This system has been

widely used in studies of coronavirus entry. To improve the expression level of S protein

and the yield of pseudotyped virus, a codon-optimized S gene based on the sequence of

isolate Wuhan-Hu-1 [2] was synthesized and used for production of pseudotyped virus as

previously described for other human coronaviruses (HCoVs), including SARS-CoV,

MERS-CoV, NL63, 229E, and OC43 [28,29]. The pseudotyped virus was then used to

infect 293T cells transfected with either empty vector, or a plasmid expressing APN

(receptor for HCoV-229E), DDP4 (receptor for MERS-CoV), ACE1 or hACE2. Two

days post-infection, the luciferase activity was measured. As shown in Fig. 2A, only

hACE2 was able to efficiently support virus entry. The entry of SARS-CoV-2, but not

influenza virus A (IVA) or HCoV-43, was blocked by antibody against hACE2 in a

dose-dependent manner (Fig. 2B). We also performed a syncytia formation assay to

assess the membrane fusion triggered by hACE2-S binding. As shown in Fig. 1C,

syncytia formation was only seen for cells expressing hACE2, but not hACE1, mixed

with cells expressing the S protein of SARS-CoV-2 or SARS-CoV. These results

confirm that hACE2 is the *bone fide* entry receptor for SARS-CoV-2.

Multiple animal ACE2 orthologs serve as receptors for SARS-CoV-2 and

SARS-CoV-2 mutant with S protein lacking the furin cleavage site

To test if other animal ACE2 orthologs can also be used as receptor for

SARS-CoV-2, we cloned or synthesized ACE2 from rhesus monkey, Chinese horseshoe

bat (Rs bat), Mexican free-tailed bat (Tb bat), rat, mouse, palm civet, raccoon dog, ferret

badger, hog badger, canine, feline, rabbit, and pangolin. These animals were chosen as

being either the proposed natural hosts for SARS-CoV-2 (bat, pangolin) [3,10],

intermediate hosts for SARS-CoV (civet, raccoon) [12], common animal model (rat,

mouse, monkey), or household pets (canine, feline, rabbit). These ACE2 molecules were

transiently expressed in 293T cells (Fig.3A), which were then infected with

pseudotyped virus of SARS-CoV-2 (SARS-CoV-2pp). The luciferase activity was

measured and normalized to hACE2 (Fig. 3B). The results showed that (1) ACE2 of

human and rhesus monkey were the most efficient receptors; (2) ACE2 of rat and mouse

barely supported virus entry (<10% of hACE2); (3) the receptor activities of the other

10 animal ACE2s were between human/monkey and rat/mouse. Among these, ACE2 of

canine, feline, rabbit and pangolin could support virus entry at levels >50% of hACE2.

To examine receptor binding ability, we performed immunoprecipitation (IP)

analysis by using both S1 and receptor binding domain (RBD) as probe. Among the 14

different ACE2s tested, ACE2 from human, monkey, feline, rabbit and pangolin

exhibited significant and consistent association with S1 and RBD (Fig.3C). Importantly,

these ACE2s correspond to the group of ACE2s that supported the most efficient virus

entry (Fig.3B).

A striking difference between SARS-CoV-2 and animal SL-CoVs is the presence

of a polybasic furin cleavage site at the S1/S2 boundary of the S protein (Fig.1). Here,

we generated a SARS-CoV-2 S gene mutant with the furin cleavage site deleted to

mimic the bat SL-CoV CZ45. This S mutant has been previously demonstrated to

express a full-length non-cleaved S protein during biogenesis in cells [30]. Pseudotyped

virus with this mutant S protein was produced and used to infect ACE2-transfected

293T cells. Similar or slightly higher efficiency were observed for the mutant S

protein-mediated pseudoviral infection in cells transfected with all the animal ACE2s,

except for mouse, rat and civet where the mutant S protein mediated a slight lower

efficiency of infection. Interesting, pangolin ACE2 was now as efficient as hACE2 for

supporting mutant virus entry (Fig.3B).

We also tested the receptor usage of these 14 ACE2 by SARS-CoV (Fig.3D). The

results indicated that ACE2 of Rs bat and rat were the poorest receptors (<20% of

hACE2), while the other ACE2s could support SARS-CoV entry at levels >50% of

hACE2. Interestingly, ACE2 of rabbit and pangolin were even more efficient than

hACE2 for supporting SARS-CoV entry. Together, these results demonstrated that

SARS-CoV-2 and its mutant virus lacking furin cleavage site, as well as SARS-CoV,

could use multiple animal ACE2s as receptor.

Molecular basis of different ACE2 receptor activities

To help understand the molecular basis of different ACE2 receptor activities, we

first examined the overall sequence variation between these ACE2s. For this purpose,

we constructed a phylogenetic tree based on the nucleotide sequences of ACE2s (Fig.4).

Interestingly, the phylogenetic clustering of ACE2s is correlated with their abilities to

support SARS-CoV-2 entry. For example, ACE2s in subclade IIA (human, rhesus

monkey and rabbit) and IIB (rat and mouse) were the most efficient and poorest receptor,

respectively, while ACE2s in clade I (from the remaining animals) were intermediate

between subclades IIA and IIB. This correlation suggests that sequence variations that

define for speciation are responsible for observed differences in receptor activity.

Next, based on the published crystal structures of hACE2-RBD complex we

compared amino acid sequences of ACE2 receptors, focusing on 23 critical residues in

close contact with RBD of SARS-CoV-2 [16,31,32] (Fig. 5). Two obvious patterns were

observed. First, hACE2 and rhesus monkey ACE2 are identical at all critical residues

for RBD interaction. This explains why rhesus monkey ACE2 supported virus entry as

efficient as hACE2 (Fig.3B). Second, since rat and mouse merely support virus entry,

the three substitutions (D30N, Y83F and K353H) that are only seen in rat and mouse

ACE2s may be the key.

To further explain the different receptor activities, we used homology-based

structure modeling to analyze the effect of residue substitutions at the atomic level.

Structure models of 14 ACE2s were generated based on the crystal structure of

SARS-CoV-2 RBD/ACE2 complex [16]. The effects of critical residue substitutions were

analyzed and are summarized in Table S1. Overall, the predicted effects of residue

substitutions in ACE2s were consistent with corresponding receptor activities. ACE2 of

rodents and bats are presented as examples of this analysis (Fig.6).

First, we examined the rodent-unique substitutions D30N, Y83F and K353H as

they may play a key role in rat and mouse ACE2 inactivity. In humans the residues at

all three of these positions directly contact the RBD via hydrogen bonds. D30 contacts

K417, Y83 contacts N487, and K353 appears to be at the center of a hydrogen bond

network spanning seven RBD residues (Y449, G496, Q498, T500, N501, G502, Y505)

and eight ACE2 residues (D38, Y41, Q42, N330, K353, G354, D355, R357). The D30N,

Y83F and K353H substitutions are all predicted to disrupt these interactions in rat and

mouse ACE2 (Figure 6). This is consistent with previous reports which pinpoint K353

as an important hotspot for both SARS-CoV-2 [16] and SARS-CoV [33] binding. It has been

experimentally demonstrated that introduction of K353H into hACE2 significantly

reduces binding to SARS-CoV S1; in contrast, introduction of H353K into rat ACE2

significantly increases binding to SARS-CoV S1 [34]. Our homology models indicate that

other residue substitutions may also be contributing to the low viral entry activity in

mouse and rat ACE2. Substitutions Q24N, Q27S, M82N, Q325P and E329T in rat

ACE2, and L79T, M82S and E329A in mouse ACE2, are all predicted to disrupt

interactions with RBD residues (Fig. 6 and Table S1).

Both Bat ACE2s are also inefficient receptors for viral entry (Fig.3B). Since the

profile of residues at the receptor/RBD interface is significantly different from rat and

mouse ACE2, we examined other bat-specific residue substitutions that may be

contributing to receptor dysfunction. There are 8 and 10 critical residue substitutions in

the Rs bat and Tb bat ACE2s, respectively (Fig.5). Among these, we examined the

substitutions at positions Y41, H34 and E329 as they are only seen in bat ACE2s. The

Y41H substitution in both bat ACE2s appears to be disrupting the same H-bond network

that was disrupted by K353H in rat and mouse ACE2. Although Y41 is not as centrally

located in the H-bond network as K353, it directly contacts N501 from the RBD, which is

the same residue that is stabilized by K353. A second interaction which appears to be

disrupted in only bat ACE2s occurs at position H34. In humans, H34 forms a H-bond

with Y453 from the RBD, which is broken through a H34T substitution in bat ACE2s.

Finally, the bat-unique substitution E329N appears to be disrupting H-bonds connecting

two ACE2 residues (E329, Q325) and two RBD residues (N439, Q506). In Tb bat ACE2,

all connections in the H-bond network are disrupted by the single E329N substitution,

however the H-bond network is predicted to be restored by an additional substitution,

Q325E in Rs bat. In addition, other residue substitutions, i.e. T27M and M82N in Rs bat,

and D30Q and L79H in Tb bat, are also disruptive (Fig. 6).

These results reveal that the poor and low receptor activity of rodent ACEs and bat

ACE2 are resulted from a broken interaction network by a key residue substitution, i.e.

K353H in rodents and Y41H in bats, and additive disruptive effects by multiple residue

substitutions.

Discussion

In this study, we examined the receptor activity of 14 ACE2 orthologues. The

results suggested that wild type and mutant SARS-CoV-2 lacking the furin cleavage site

in S protein could use ACE2 from a broad range of animal species to enter host cells.

Below we discuss the implication of our findings in terms of natural reservoir, zoonotic

transmission, human-to-animal transmission, animal health, and animal model.

Implication in natural reservoirs and zoonotic transmission

Among the 14 ACE2s tested here, hACE2 and rhesus monkey ACE2 are the most

efficient receptors, suggesting that SARS-CoV-2 has already been well adapted to

humans. In addition, ACE2s of other animals, except mouse and rat, could also support

SARS-CoV-2 entry (Fig.3B). Although these data were obtained by using HIV1-based

pseudotyped virus, for ACE2 of Rs bat, civet, and mouse, the data is consistent with *in*

vitro infection data using infectious virus [3]. Receptor usage by coronaviruses has been

well known to be a major determinant of host range, tissue tropism, and pathogenesis

[14,35,36]. It is therefore reasonable to assume that SARS-CoV-2 would be able to infect all

these animals. As a matter of fact, several *in vivo* infection and seroconversion studies

have confirmed that SARS-CoV-2 can infect rhesus monkey [37], feline, ferret, and canine

[38,39]. Our findings are also in line with the concordance between ACE2 receptor usage

by SARS-CoV pseudotyped virus and susceptibility of the animals to SARS-CoV

infection. As shown in Fig.3D, ACE2 of rhesus monkey, mouse, civet, ferret badger,

raccoon and feline could support SARS-CoV pseudotyped virus entry; concordantly, all

these animals are susceptible to native SARS-CoV virus infection [12,40-43].

Among all those wild animals that are potentially infected by SARS-CoV-2, bat

and pangolin have already been proposed to be the natural reservoirs as closely related

SL-CoVs have been identified in bats [2,3,5] and pangolins [8-11]. A recent study has shown

that bat SL-CoV RaTG13 could use hACE2 as receptor, consistent with the presence of

several favorable hACE2-binding residues (aa 455, 482-486) in the receptor binding

motif (RBM) of the S protein (Fig.1) [16]. For pangolin SL-CoVs, lineage PCoV-GD has

only one non-critical amino acid substitution (Q483H) in the RBM when compared to

SARS-CoV-2 (Fig.1) [10]. Therefore, PCoV-GD most likely can also use hACE2 and

other animal ACE2s as functional receptors.

We also tested the receptor usage by a SARS-CoV-2 mutant that lacks the furin

cleavage site at the S1/S2 boundary. Our result showed that the mutant virus behaved

similarly to the wt virus. Namely, the entry of mutant virus could also be supported by

those animal ACE2s that supported the entry of wt virus. This result is similar to another

study that used the S gene mutant but in a MLV-based pseudotyped virus system [17] and

the role of furin cleavage during coronavirus infection. Furin cleavage is not essential

for coronavirus-cell membrane fusion, but enhances cell-to-cell fusion [18-21,44]. This

could provide certain level of advantage during infection. For example, in the

quasispecies population of bovine CoV, a minor sequence variant with a polybasic

furin-like cleavage site in the S2 subunit quickly dominated the population even after a

single passage in cells [23]. However, by using pseudotyped virus system, which is a

single cycle infection system, we may not see the advantage. Still, ours result

unequivocally showed that SARS-CoV-2 without this cleavage site could use multiple

animal ACE2s as receptors to enter cells. As there is a need to continuously search for

potential intermediate hosts for SARS-CoV, results presented here can help significantly

narrow down the scope of potential targets.

Collectively, our results highlight the potential of these wild animals to serve as

natural reservoirs or intermediate hosts for SARS-CoV-2 and its progenitor, the risk of

zoonotic transmission of animal SL-CoVs to human, and the necessity of virus

surveillance in wild animals.

Implication in human-to-animal transmission and animal health

Among those animal species tested here, canine and feline are of special concern as

they are often raised as companion pets. Our data indicate that ACE2 of canine and

feline could support SARS-CoV-2 pseudotyped virus entry quite efficiently (>50% of

hACE2, Fig.3B), raising the alarming possibility of virus transmission from infected

human to these pets or potentially vice versa. As a matter of fact, there was a recent

report that a Pomeranian dog in Hong Kong tested weakly positive for SARS-CoV-2

while maintaining an asymptomatic state. The genome of the virus isolated from this

dog has only three nucleotide changes compared to the virus isolated from two infected

persons living in the same household, suggesting that this dog probably acquired the

virus from the infected owners [45]. Our results are further supported by two additional

studies. One study showed that both dog and cat were susceptible to SARS-CoV-2

infection. While the virus replicated poorly in dogs, it replicated efficiently in cats and

was able to transmit to unaffected cats that were housed with them [38]. The other study

revealed that 14.7% of cat sera samples collected in Wuhan city after the outbreak were

positive for antibody against SARS-CoV-2, demonstrating that many cats were infected

during the outbreak, most likely from infected humans in close contact [39]. Domestic cats

are also susceptible to SARS-CoV infection [40] and human-to-cat transmission was

evident during the SARS-CoV outbreak in 2003 in Hong Kong [46]. These findings were

also in agreement with our results that ACE2 of cat and dog could serve as receptor for

SARS-CoV (Fig.3D).

As described above, it seems that dogs are not as susceptible as cats to

SARS-CoV-2 [38,45]. Interestingly, this is in agreement with results from IP analysis that

showed cat ACE2 could bind to S1 or RBD more efficiently than dog ACE2 (Fig.3C).

Structural models further suggest that, at those critical RBD-binding residues, dog and

cat ACE2 share 4 substitutions (Q24L, D30E, D38E, and M82T), while dog ACE2 has

an additional substitution, H34Y (Fig.5). Based on structural modeling, both Q24L and

M82T are predicted to be disruptive, while both D30E and D38E are tolerable (Table

S1). H34Y in dog ACE2 is predicted to disrupt the hydrogen bond with Y453 of RBD

(Table S1). These atomic interactions explain why dog ACE2 binds to S1 or RBD less

efficiently compared to cat ACE2, and both are less efficient than human ACE2.

In addition to cat and dog, rabbits are also often raised as household pets. Our

results indicate that rabbit ACE2 is an efficient receptor (Fig. 3B and 3C), suggesting

that rabbit may be more susceptible to SARS-CoV-2 infection than cat.

Currently, there is no evidence that infected pets can transmit the virus back to

human; however, this may be possible and should be investigated. Out of an abundance

of caution it would be best when one is infected to have both human and pets

quarantined, and the pets tested as well.

Implication in animal model

Animal models are essential for study of pathogenesis, vaccinology and

therapeutics of viral pathogens. Rodents are probably the most common and amenable

animal models because of low cost, easy handling, defined genetics, and the possibility

of scalability [47]. However, our results showed that both mouse and rat ACE2 are poor

receptors for SARS-CoV-2 (Fig.3B and 3C), suggesting that they are probably resistant

to infection. Actually, this has been verified by using infectious SARS-CoV-2 to infect

mouse ACE2-transfected cells [3] or mice [48]. Genetically engineered mice expressing

hACE2 were previously developed as an animal model for SARS-CoV [49]. This model

has been tested recently for SARS-CoV-2, and found to be susceptible to SARS-CoV-2

infection and development of interstitial pneumonia [48], a common clinical feature of

COVID-19 patients [50]. hACE2-transgenic mice therefore represent useful animal

models. However, because of the high demand, and discontinuance due to the

disappearance of SARS-CoV in the human population after 2004, it is expected that this

mouse model will be in short supply [51]. Alternative methods should be sought to

develop a mouse-adapted SARS-CoV-2 strain. Mouse-adapted SARS-CoV strains were

developed by serial passage of virus in mice [52,53]. However, this method may not work

for SARS-CoV-2 as mouse ACE2 still supports some entry for SARS-CoV (Fig.3D),

but not SARS-CoV-2. An alternative way to make a mouse-adapted SARS-CoV-2 strain

could be achieved by rational design of the S gene. Based on the structural model, we

know that receptor dysfunction of mouse ACE2 is due to disruptive D30N, L79T, M82S,

Y83F, E329A and K353H substitutions (Fig.5, Fig.6 and Table S1). Therefore, by

specifically introducing mutations into the RBM of S gene it may be possible to restore

or at least partly restore interactions with these ACE2 substitutions. Consequently, the

engineered virus may be able to efficiently infect wildtype mice.

To date, several animals (i.e. rhesus monkey, ferret, dog, cat, pig, chicken and duck)

have been examined as potential animal models for SARS-CoV-2 [37,38]. Although the

rhesus monkey, ferret and cat may seem to be the promising candidates, none of them

are perfect in terms of recapitulation of typical clinical features in COVID-19 patients.

Therefore, multiple animal models may be needed. Our results indicate that rabbit

ACE2 is a more efficient receptors than other animal ACE2s for both SARS-CoV-2 and

SARS-CoV (Fig.3). Therefore, it may be worthy assessing rabbit as a useful animal

model for further studies.

Methods

Cell lines and antibodies

293T cells and Lenti-X 293T cells were cultured in Dulbecco's modified Eagle's

medium (DMEM; Gibco) [54]. All growth medium was supplemented with 10% fetal

bovine serum (FBS), 110 mg/L sodium pyruvate, and 4.5 g/L D-glucose. β-actin

antibody and C9 antibody were purchased from Sigma (A2228) and SANTA CRUZ

(sc-57432), respectively. A polyclonal antibody against human ACE2 and anti-IDE

polyclonal antibody were purchased from R&D Systems (catalog No. AF933 and

AF2496, respectively).

Construction of ACE2 plasmids

ACE2 of human (*Homo sapiens*), civet (*Paguma larvata*), and rat (*Rattus*

norvegicus) were cloned into a modified pcDNA3.1-cmyc/C9 vector (Invitrogen) as

previously described [34,55]. ACE2 protein expressed from this vector has a c-myc tag at

the N-terminus and a C9 tag at the C-terminus. An Age I site was engineered right

downstream of the signal peptide sequence (nt. 1-54) of ACE2. ACE2 protein expressed

from this vector has a c-myc tag at the N-terminus and a C9 tag at the C-terminus.

ACE2 of Chinese ferret badger (*Melogale moschata*), raccoon dog (*Nyctereutes*

procyonoides), Mexican free-tailed bat (*Tadarida brasiliensis*), rhesus monkey (*Macaca*

mulatta), hog badger (*Arctonyx collaris*), New Zealand white rabbit (*Oryctolagus*

cuniculus), domestic cat (*Felis catus*) and domestic dog (*Canis lupus familiaris*) were

cloned into Age I/Kpn I-digested pcDNA3.1-cmyc-C9 vector previously (Hanxin Lin,

Ph. D Thesis Dissertation. "Molecular interaction between the spike protein of human

coronavirus NL63 and ACE2 receptor" McMaster University, Health Science

Library.https://discovery.mcmaster.ca/iii/encore/record/C__Rb2023203__SMolecular%

20interaction%20between%20the%20spike%20protein%20of%20human%20coronavir

us%20NL63%20and%20ACE2%20receptor%20Lw%3D%3D%20by%20Hanxin%20Li

n__Orightresult__U__X4?lang=eng&suite=def). The nucleotide sequence of ACE2 of

Chinese horseshoe bat (*Rhinolophus sinicus*, Rs, GenBank No. KC881004.1) and

pangolin (*Manis javanica*, GenBank No. XM_017650263.1) were synthesized and

cloned into pcDNA3.1-N-myc/C-C9 vector.

Construction of plasmids expressing S, S1 and RBD of SARS-CoV-2

The nucleotide sequence of SARS-CoV-2 S gene was retrieved from NCBI

database (isolate Wuhan-Hu-1, GenBank No. MN908947). According the method

described by Gregory J. Babcock et al [56], the codon-optimized S gene was synthesized,

and cloned into pCAGGS vector. The SARS-CoV-2 S gene mutant without the furin

cleavage site at the S1/S2 boundary was generated by an overlapping PCR-based

method as previously described [30]. The S1 subunit (aa 14-685) and RBD (aa 331-524)

were cloned into a soluble protein expression vector, pSecTag2/Hygro-Ig vector, which

contains human IgG Fc fragment and mouse Ig *k*-chain leader sequence [57]. The protein

expressed is soluble and has a human IgG-Fc tag.

Western blot assay

As previously described, the expression of ACE2-C9, S1-Ig, and RBD-Ig fusion

proteins were examined by western-blot [57]. Briefly, lysates or culture supernatants of

293T cells transfected with plasmid encoding ACE2 orthologs and S1-Ig or RBD-Ig

were collected, boiled for 10 min, and then resolved by 4~12% SDS-PAGE. A PVDF

membrane containing the proteins transferred from SDS-PAGE was blocked with

blocking buffer (5% nonfat dry milk in TBS) for 1h at room temperature and probed

with primary antibody overnight at 4 °C. The blot was washed three times with washing

buffer (0.05% tween-20 in TBS), followed by incubation with secondary antibody for

1h at room temperature. After three-time washes, the proteins bounded with antibodies

were imaged with the Li-Cor Odyssey system. (Li-Cor Biotechnology).

Immunoprecipitation (IP) assay.

The association between Ig-fused S1 protein or RBD protein and ACE2 protein

with C9 tag was measured by IP according to a previously described method [30]. Briefly,

HEK293T cells were transfected with plasmid encoding ACE2 with Lipofectamine

2000 (Invitrogen). 48 h post-transfection, the transfected 293T cells were harvested and

lysed in PBS buffer containing 0.3 % *n*-decyl-β-D-maltopyranoside (DDM, Anatrace).

Cell lysates were incubated with Protein A/G PLUS Agarose (Santa Cruz, sc-2003)

together with 4μg of S1-Ig or RBD-Ig. Protein A/G agarose were washed three times in

TBS/1% Ttriton-X100, resolved by SDS–PAGE, and detected by western blot using

anti-C9 monoclonal antibody.

Production of pseudotyped virus

Following the standard protocol of calcium phosphate transfection, Lenti-X cells in

10-cm plate were co-transfected by 20μg of HIV-luc and 10μg of CoVs spike gene

plasmid. At 48h post-transfection, 15 ml supernatant was collected and passed through a

0.45um pore size PES filter. The purified virus was titrated with Lenti-X p24 Rapid

Titer Assay (Takara Bio, Cat. No. 632200). The virus was stored at -80 °C for future

use.

Virus entry assay

Each well of 293T cells in a 96-well plate was transfected with 0.1μg of ACE2

plasmid DNA following the standard protocol of Lipofectamine 2000 (Invitrogen). At

48 hours post-transfection, 150 μl of p24-normalized (10 ng) of pseudotype virus was

added into each well and incubated at 37°C for 3 hours. The virus was then removed,

and 250 µl of fresh medium was added into each well for further incubation. Two days

post-infection, the medium was removed and the cells were lysed with 30 µl/well of 1×

cell lysis buffer (Promega) for 15 min, followed by adding 50 µl/well of luciferase

substrate (Promega). The firefly luciferase activities were measured by luminometry in

a TopCounter (PerkinElmer). For each ACE2, four wells were tested in a single

experiment, and at least three repeat experiments were carried out. The luciferase

activity was expressed as relative light unit (RLU) and normalized to human ACE2 for

plotting.

Syncytial formation assay

293T cells with approximately 90% confluent on 12-well plate were transfected

with 1.6µg of plasmid DNA encoding viral S gene or ACE2. At 24 h post-transfection,

293T cells expressing the S protein were mixed at a 1:1 ratio with 293T cells expressing

ACE2 and plated on 12-well plate. Multinucleated syncytia were observed 24 h after the

cells were mixed.

Sequence analysis

Multiple alignments of nucleotide or amino acid sequences of the spike gene of

coronaviruses and ACE2 orthologs were performed using Clustal X [58]. Phylogenetic

tree was constructed based on the nucleotide sequences of animal ACE2 using the

neighbor-joining algorithm implemented in MEGA X. The tree is drawn to scale with

branch lengths in the same units as those of the evolutionary distances used to infer the

phylogenetic tree. Evaluation of statistical confidence in nodes was based on 1000

bootstrap replicates. Branches with <50% bootstrap value were collapsed. Platypus

ACE2 (*Ornithorhynchus anatinus*, GenBank No. XM_001515547) was used as an

outgroup.

Homology-based structural modeling.

Human ACE2 (PDB: 6VW1) in the bound conformation was extracted from the

SARS-CoV-2 RBD/ hACE2 complex and used as a template for homology modeling

[16]. ACE2 Homology models were generated using the one-to-one threading algorithm of

Phyre2 [59]. The models were then aligned and compared to the intact SARS-CoV-2

RBD/ ACE2 complex in PyMOL (The PyMOL Molecular Graphics System, Version 2.0

Schrödinger, LLC).

FUNDING

This work was supported by grants from the National Natural Science Foundation of

China (81772173 and 81971916) and National Science and Technology Mega-Project of

China (2018ZX10301-408-002) to X Zhao and from the Canadian Institutes of Health

Research (MOP-89903 to MSJ).

  Figure and Figure legends

Fig. 1. Schematic diagram of domain structures and critical ACE2-binding residues of the spike (S) protein of SARS-CoV-2. Signal: signal peptide. NTD: N-terminal domain. RBD: receptor-binding domain. RBM: receptor-binding motif. FP: fusion peptide. TM: transmembrane domain. CT: cytoplasmic tail. The S protein is cleaved into S1 and S2 subunit during biogenesis at the polybasic furin cleavage site (RRAR↓), which is not present in SARS-CoV and other animal SARS-CoV-2-like CoVs. The S1 subunit is required for binding to ACE2 receptor, while the S2 subunit containing a FP mediates membrane fusion. In SARS-CoV-2, the S1 contains NTD and an independently folded domain known as RBD, which harbors a region called receptor binding motif (RBM), that are primarily in contact with receptor. The most critical hACE2-binding residues in the RBM of several SARS-CoV-2-related CoVs are highlighted in yellow and referred from the crystal structure of RBD-hACE2 complex (Shang et al, Nature). PCoV-GX: pangolin CoV isolate GX-PL4. PCoV-GD: pangolin CoV isolate MP789. The only difference in the RBM between PCoV-GD and SARS-CoV-2 is Q498H (underlined). The GenBank No. for these CoVs is: SARS-CoV-2 (isolate Wuhan-Hu-1, MN908947), SARS-CoV (isolate Tor2, NC_004718.3), bat-ZC45 (MG772933.1), bat-RaTG13 (MN996532.1), PCoV-GX (isolate P4L, MT040333.1), PCoV-GD (isolate MP789, MT084071.1).

503
504
Fig. 2. Human ACE2 served as receptor for SARS-CoV-2. (A) ACE2 supported
HIV-Luc-based pseudotyped virus entry. 293T cells were transfected with empty vector
pcDNA3.1, APN (receptor for HCoV-229E), DDP4 (receptor for MERS-CoV), ACE1
or ACE2. At 48 h post transfection, the cells were infected by SARS-CoV-2 S protein
pseudotyped virus (SARS-CoV-2pp). At 48 h post infection, luciferase activity was
measured. (B) Human ACE2 antibody inhibited virus entry at a dose-dependent manner.
293T cells were transfected with ACE2. At 48 h post transfection, the cells were
pre-incubated with indicated concentration of hACE2 antibody or control antibody

(anti-IDE) for 1 h, and then infected by pseudotyped virus of SARS-CoV-2, Influenza virus A (IAVpp) or human coronavirus (HCoV) OC43 (HCoV-OC43pp) in the presence of indicated concentration of hACE2 antibody or control antibody (anti-IDE) for another 3 h, then the virus and antibodies were removed. At 48 h post infection, luciferase activity was measured and normalized to the control antibody for SARS-CoV-2pp. Error bars reveal the standard deviation of the means from four biological repeats. (C) Syncytia formation assay. 293T cells transfected with a plasmid the expressing S protein of SARS-CoV-2 or SARS-CoV were mixed at a 1:1 ratio with those cells transfected with a plasmid expressing ACE1 or ACE2. Twenty-four hours later, syncytia formation was recorded.

Fig. 3. Multiple ACE2 orthologues served as receptors for SARS-CoV-2. (A) Transient expression of ACE2 orthologs in 293T cells. The cell lysates were detected by western blot assay, using an anti-C9 monoclonal antibody. (B) HIV-Luc-based pseudotyped virus entry. 293T cells were transfected with ACE2s orthologs. At 48 h post transfection, the cells were infected by the pseudotyped virus of wildtype SARS-CoV-2 or mutant ΔFurin. At 48 h post infection, luciferase activity was measured and normalized to human ACE2, respectively. Error bars reveal the standard deviation of the means from four biological repeats. (C) IP assay. The upper panel showed the input of ACE2 protein with C9 tag, S1 and RBD with IgG tag. The lower panel showed the ACE2 pulled down by S1-Ig or RBD-Ig fusion protein.

Fig. 4. Phylogenetic clustering of ACE2s correlates with their receptor activities.
Upper panel: phylogram tree of 14 ACE2s. The tree was constructed based on
nucleotide sequences using the Neighbor-joining method implemented in program
MEGA X. The percentage of replicate trees in which the associated taxa clustered
together in the bootstrap test (1000 replicates) are shown next to the branches. The tree
was rooted by ACE2 of platypus (*Ornithorhynchus anatinus*). The taxonomic orders
where these animals are classified are shown on the right-hand side of the tree. **Lower**
panel: a heat bar summarizing the relative levels of pseudotyped virus entry supported
by different animal ACE2s.

ACE2	19	24	27	28	30	31	34	35	37	38	41	42	45	79	82	83	325	329	330	353	354	355	357
Human	S	Q	T	F	D	K	H	E	E	D	Y	Q	L	L	M	Y	Q	E	N	K	G	D	R
Monkey	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rs bat	-	E	M	-	-	-	T	K	-	-	H	-	-	-	N	-	E	N	-	-	-	-	-
Tb bat	-	E	I	-	Q	R	T	-	-	E	H	-	-	H	R	-	-	N	-	-	-	-	-
Rat	-	K	S	-	N	-	Q	-	-	-	-	-	-	-	I	N	F	P	T	-	H	-	-
Mouse	-	N	-	-	N	N	Q	-	-	-	-	-	-	-	T	S	F	-	A	-	H	-	-
Civet	-	L	-	-	E	T	Y	-	Q	E	-	-	V	-	T	-	-	-	-	-	D	-	-
Raccoon	-	L	-	-	E	-	Y	-	-	E	-	-	-	-	T	-	-	-	-	R	-	-	-
Ferret badger	-	L	-	-	E	-	-	-	-	E	-	-	-	H	T	-	E	Q	-	-	R	-	-
Hog badger	-	L	-	-	E	-	R	-	-	E	-	-	-	H	T	-	E	K	-	-	H	-	-
Canine	-	L	-	-	E	-	Y	-	-	E	-	-	-	-	T	-	-	-	-	-	-	-	-
Feline	-	L	-	-	E	-	-	-	-	E	-	-	-	-	T	-	-	-	-	-	-	-	-
Rabbit	-	L	-	-	E	-	Q	-	-	-	-	-	-	-	T	-	-	-	-	-	-	-	-
Pangolin	-	E	-	-	E	-	S	-	-	E	-	-	-	I	N	-	-	-	-	-	H	-	-

Fig. 5. Critical RBD-binding residues in ACE2 orthologs. Upper panel: 23 RBD-binding residues at the contact interface between hACE2 and RBD of SARS-CoV-2. Human ACE2 (PDB: 6VW1) in the bound conformation was extracted from the SARS-CoV-2 RBD/ACE2 complex and used as a template for homology modeling [16]. **Low panel**: Critical RBD-binding residues in ACE2 orthologs. Residue substitutions highlighted in red and orange are those unique to both mouse and rat ACE2s and both bats, respectively. The rest of residue substitutions are highlighted in yellow.

Fig. 6. Structural models of key residue substitutions in ACE2 of mouse, rat and bats. Human ACE2 (PDB: 6VW1) in the bound conformation was extracted from the SARS-CoV-2 RBD/ACE2 complex and used as a template for homology modeling [16]. ACE2 Homology models were generated using the one-to-one threading algorithm of Phyre2 [59]. The models were then aligned and compared to the intact SARS-CoV-2 RBD/ ACE2 complex in PyMOL.

Table S1. Predicted effect of clinical residue substitutions in ACE2 orthologs on the interaction with RBD of SARS-CoV-2

Human ACE2		Interaction			Predicted status of interaction in ACE2 homologs (Disrupted/ Enhanced/ Tolerated/ Unsure/ No residue change)												
Index	Residue	Molecule	Residue	Type	Monkey	BatRs	BatTb	Rat	Mouse	Civet	Raccoon	Ferret	HogBadger	Dog	Cat	Rabbit	Pangolin
19	S	ACE2	Q24	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
24	Q	RBD	N487	H-bond	-	T	T	D	T	D	D	D	D	D	D	D	T
		ACE2	S12	H-bond	-	D	D	T	T	D	D	D	D	D	D	D	D
27	T	RBD	Y489	VdW	-	D	T	D	-	-	-	-	-	-	-	-	-
		RBD	A475	VdW	-	D	T	D	-	-	-	-	-	-	-	-	-
		RBD	F456	VdW	-	D	T	D	-	-	-	-	-	-	-	-	-
28	F	ACE2	Y83	π	-	-	-	-	-	-	-	-	-	-	-	-	-
30	D	RBD	K417	H-bond	-	-	D	D	D	T	T	T	T	T	T	T	T
31	K	RBD	Q493	H-bond	-	-	T	-	D	D	-	-	-	-	-	-	-
34	H	RBD	Y453	H-bond	-	D	D	T	T	D	D	-	T	D	-	T	D
35	E	RBD	Q493	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
37	E	ACE2	R393	H-bond	-	-	-	-	-	D	-	-	-	-	-	-	-
		RBD	Y505	VdW	-	-	-	-	-	E	-	-	-	-	-	-	-
38	D	ACE2	Q42	H-bond	-	-	-	-	-	T	T	T	T	T	T	-	T
		ACE2	K353	H-bond	-	-	-	-	-	T	T	T	T	T	T	-	T
		RBD	Q498	H-bond	-	-	-	-	-	T	T	T	T	T	T	-	T
		RBD	Y449	VdW	-	-	-	-	-	T	T	T	T	T	T	-	T
41	Y	RBD	Q498	π	-	D	D	-	-	-	-	-	-	-	-	-	-
		RBD	T500	VdW	-	D	D	-	-	-	-	-	-	-	-	-	-
		RBD	N501	H-bond	-	D	D	-	-	-	-	-	-	-	-	-	-
		ACE2	R357	H-bond	-	D	D	-	-	-	-	-	-	-	-	-	-
		ACE2	K353	π	-	D	D	-	-	-	-	-	-	-	-	-	-
42	Q	RBD	Y449	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	D38	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
45	L	RBD	Q498	VdW	-	-	-	-	-	T	-	-	-	-	-	-	T
79	L	RBD	F486	Hydrophobic	-	-	D	T	D	D	-	D	D	-	-	-	D
82	M	RBD	F486	π	-	D	U	D	D	D	D	D	D	D	D	D	D
83	Y	RBD	F486	π	-	-	-	D	D	-	-	-	-	-	-	-	-
		RBD	N487	H-bond	-	-	-	D	D	-	-	-	-	-	-	-	-
		ACE2	F28	π	-	-	-	D	D	-	-	-	-	-	-	-	-
325	Q	ACE2	E329	H-bond	-	T	D	D	D	-	-	T	T	-	-	-	-
329	E	RBD	Q506	H-bond	-	T	D	D	D	-	-	T	T	-	-	-	-
		RBD	N439	H-bond	-	T	D	D	D	-	-	T	T	-	-	-	-
		ACE2	Q325	H-bond	-	T	D	D	D	-	-	T	T	-	-	-	-
330	N	ACE2	R357	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	D355	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
353	K	RBD	G496	H-bond	-	-	-	D	D	-	T	-	-	-	-	-	-
		RBD	G502	H-bond	-	-	-	T	T	-	T	-	-	-	-	-	-
		RBD	Y505	π	-	-	-	D	D	-	T	-	-	-	-	-	-
		RDB	Q498	H-bond	-	-	-	D	D	-	T	-	-	-	-	-	-
		RBD	N501	H-bond	-	-	-	D	D	-	T	-	-	-	-	-	-
		ACE2	D38	H-bond	-	-	-	D	D	-	T	-	-	-	-	-	-
		ACE2	Y41	π	-	-	-	T	T	-	T	-	-	-	-	-	-
354	G	RBD	G502	H-bond	-	-	-	-	-	T	-	T	T	-	-	-	T
355	D	RBD	T500	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	N330	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	R357	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
357	R	RBD	T500	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	N330	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	Y41	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-
		ACE2	D355	H-bond	-	-	-	-	-	-	-	-	-	-	-	-	-

H-bond: hydrogen bond; Vdw: Van der Waals force; D: disruptive; T: tolerated; U: unsure; E: enhanced.

The effects of residues substitution were predicted by homologous-based modeling analyses based on the crystal structure of SARS-CoV-2 RBD/hACE2 complex (Ref.16)

References

1   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*
382, 727-733, doi:10.1056/NEJMoa2001017 (2020).

2   Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**,
265-269, doi:10.1038/s41586-020-2008-3 (2020).

3   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin.
Nature **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

4   Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus:
implications for virus origins and receptor binding. *Lancet* **395**, 565-574,
doi:10.1016/S0140-6736(20)30251-8 (2020).

5   Hong Zhou, X. C., Tao Hu, Juan Li, Hao Song, Yanran Liu, Peihan Wang, Di Liu, Jing Yang,
Edward C. Holmes, Alice C. Hughes, Yuhai Bi, Weifeng Shi. A novel bat coronavirus reveals
natural insertions at the S1/S2 cleavage site of the Spike protein and a possible recombinant
origin of HCoV-19. *BioRxiv* **https://doi.org/10.1101/2020.03.02.974139** (2020).

6   Hayman, D. T. Bats as Viral Reservoirs. *Annu Rev Virol* **3**, 77-99,
doi:10.1146/annurev-virology-110615-042203 (2016).

7   Cui, J., Li, F. & Shi, Z. L. Origin and evolution of pathogenic coronaviruses. *Nat Rev Microbiol*
17, 181-192, doi:10.1038/s41579-018-0118-9 (2019).

8   Zhang, T., Wu, Q. & Zhang, Z. Probable Pangolin Origin of SARS-CoV-2 Associated with the
COVID-19 Outbreak. *Curr Biol*, S0960-9822(0920)30360-30362,
doi:10.1016/j.cub.2020.03.022 (2020).

9   Xiao, K. P. *et al.* Isolation and Characterization of 2019-nCoV-like Coronavirus from Malayan
Pangolins. *BioRxiv* **https://doi.org/10.1101/2020.02.17.951335**,
doi:https://doi.org/10.1101/2020.02.17.951335 (2020).

10   Lam, T. T.-Y. *et al.* Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins.
Nature **https://doi.org/10.1101/2020.02.13.945485**,
doi:https://doi.org/10.1101/2020.02.13.945485 (2020).

11   Liu, P., Chen, W. & Chen, J. P. Viral Metagenomics Revealed Sendai Virus and Coronavirus
Infection of Malayan Pangolins (Manis javanica). *Viruses* **11**, doi:10.3390/v11110979 (2019).

12   Guan, Y. *et al.* Isolation and characterization of viruses related to the SARS coronavirus from
animals in southern China. *Science* **302**, 276-278 (2003).

13   Hemida, M. G. *et al.* MERS coronavirus in dromedary camel herd, Saudi Arabia. *Emerg Infect*
Dis **20**, 1231-1234, doi:10.3201/eid2007.140571 (2014).

14   Li, F. Receptor recognition and cross-species infections of SARS coronavirus. *Antiviral Res* **100**,
246-254, doi:10.1016/j.antiviral.2013.08.014 (2013).

15   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked
by a Clinically Proven Protease Inhibitor. *Cell* **10.1016/j.cell.2020.02.052**,
doi:10.1016/j.cell.2020.02.052 (2020).

16   Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature*
10.1038/s41586-020-2179-y, doi:10.1038/s41586-020-2179-y (2020).

17   Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein.
Cell **10.1016/j.cell.2020.02.058**, doi:10.1016/j.cell.2020.02.058 (2020).
18   Bos, E. C., Heijnen, L., Luytjes, W. & Spaan, W. J. Mutational analysis of the murine
coronavirus spike protein: effect on cell-to-cell fusion. *Virology* **214**, 453-463,
doi:S0042-6822(85)70056-6 [pii]
10.1006/viro.1995.0056 (1995).
19   de Haan, C. A., Stadler, K., Godeke, G. J., Bosch, B. J. & Rottier, P. J. Cleavage inhibition of the
murine coronavirus spike protein by a furin-like enzyme affects cell-cell but not virus-cell fusion.
J Virol **78**, 6048-6054, doi:10.1128/JVI.78.11.6048-6054.2004
78/11/6048 [pii] (2004).
20   Stauber, R., Pfleiderera, M. & Siddell, S. Proteolytic cleavage of the murine coronavirus surface
glycoprotein is not required for fusion activity. *J Gen Virol* **74 (Pt 2)**, 183-191 (1993).
21   Taguchi, F. Fusion formation by the uncleaved spike protein of murine coronavirus JHMV
variant cl-2. *J Virol* **67**, 1195-1202 (1993).
22   Millet, J. K. & Whittaker, G. R. Host cell proteases: Critical determinants of coronavirus tropism
and pathogenesis. *Virus Res* **202**, 120-134, doi:10.1016/j.virusres.2014.11.021 (2015).
23   Borucki, M. K. *et al.* The role of viral population diversity in adaptation of bovine coronavirus to
new host environments. *PLoS One* **8**, e52752, doi:10.1371/journal.pone.0052752 (2013).
24   Ito, T. *et al.* Generation of a highly pathogenic avian influenza A virus from an avirulent field
isolate   by   passaging   in   chickens.   *Journal   of   virology*   **75**,   4439-4443,
doi:10.1128/JVI.75.9.4439-4443.2001 (2001).
25   Gallagher, T. M. & Buchmeier, M. J. Coronavirus spike proteins in viral entry and pathogenesis.
Virology **279**, 371-374, doi:10.1006/viro.2000.0757S0042-6822(00)90757-8 [pii] (2001).
26   Letko, M., Marzi, A. & Munster, V. Functional assessment of cell entry and receptor usage for
SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* **10.1038/s41564-020-0688-y**,
doi:10.1038/s41564-020-0688-y (2020).
27   Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus.
Nature **426**, 450-454 (2003).
28   Chen, D. *et al.* GILT restricts the cellular entry mediated by the envelope glycoproteins of
SARS-CoV, Ebola virus and Lassa fever virus. *Emerg Microbes Infect* **8**, 1511-1523,
doi:10.1080/22221751.2019.1677446 (2019).
29   Zhao, X. *et al.* Identification of Residues Controlling Restriction versus Enhancing Activities of
IFITM   Proteins   on   Entry   of   Human   Coronaviruses.   *Journal   of   virology*   **92**,
doi:10.1128/JVI.01535-17 (2018).
30   Lin, H. X. *et al.* Identification of residues in the receptor-binding domain (RBD) of the spike
protein of human coronavirus NL63 that are critical for the RBD-ACE2 receptor interaction. *The*
Journal of general virology **89**, 1015-1024, doi:10.1099/vir.0.83331-0 (2008).
31   Jun Lan, J. G., Jinfang Yu, Sisi Shan, Huan Zhou, Shilong Fan, Qi Zhang, Xuanling Shi, Qisheng
Wang, Linqi Zhang, Xinquan Wang. Crystal structure of the 2019-nCoV spike receptor-binding
domain bound with the ACE2 receptor. *BioRxiv* **doi: https://doi.org/10.1101/2020.02.19.956235**
(2020).
32   Yan, R. *et al.* Structural basis for the recognition of the SARS-CoV-2 by full-length human
ACE2. *Science* **10.1126/science.abb2762**, doi:10.1126/science.abb2762 (2020).
33   Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike
receptor-binding domain complexed with receptor. *Science* **309**, 1864-1868 (2005).

34   Li, W. *et al.* Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2.

716   *EMBO J* **24**, 1634-1643, doi:7600640 [pii]10.1038/sj.emboj.7600640 (2005).

717   35   Li, F. Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol* **3**,

718   237-261, doi:10.1146/annurev-virology-110615-042301 (2016).

719   36   Perlman, S. & Netland, J. Coronaviruses post-SARS: update on replication and pathogenesis.

720   *Nat Rev Microbiol* **7**, 439-450, doi:10.1038/nrmicro2147 (2009).

721   37   Wei Deng, L. B., Hong Gao, Zhiguang Xiang, Yajin Qu, Zhiqi Song, Shunran Gong, Jiayi Liu,

722   Jiangning Liu, Pin Yu, Feifei Qi, Yanfeng Xu, Fengli Li, Chong Xiao, Qi Lv, Jing Xue, Qiang

723   Wei, Mingya Liu, Guanpeng Wang, Shunyi Wang, Haisheng Yu, Xing Liu, Wenjie Zhao, Yunlin

724   Han, Chuan Qin. Rhesus macaques can be effectively infected with SARS-CoV-2 via ocular

725   conjunctival route. *BioRxiv* **doi: https://doi.org/10.1101/2020.03.13.990036** (2020).

726   38   Jianzhong Shi, Z. W., Gongxun Zhong, Huanliang Yang, Chong Wang, Renqiang Liu, Xijun He,

727   Lei Shuai, Ziruo Sun, Yubo Zhao, Libin Liang, Pengfei Cui, Jinliang Wang, Xianfeng Zhang,

728   Yuntao Guan, Hualan Chen, and Zhigao Bu. Susceptibility of ferrets, cats, dogs, and different

729   domestic animals to SARS-coronavirus-2 *BioRxiv* **doi:**

730   **https://doi.org/10.1101/2020.03.30.015347** (2020).

731   39   Qiang Zhang, H. Z., Kun Huang, Yong Yang, Xianfeng Hui, Jindong Gao, Xinglin He, Chengfei

732   Li, Wenxiao Gong, Yufei Zhang, Cheng Peng, Xiaoxiao Gao, Huanchun Chen, Zhong Zou,

733   Zhengli Shi, Meilin Jin. SARS-CoV-2 neutralizing serum antibodies in cats: a serological

734   investigation. *BioRxiv* **doi: https://doi.org/10.1101/2020.04.01.021196** (2020).

735   40   Martina, B. E. *et al.* Virology: SARS virus infection of cats and ferrets. *Nature* **425**, 915 (2003).

736   41   Qin, C. *et al.* An animal model of SARS produced by infection of Macaca mulatta with SARS

737   coronavirus. *J Pathol* **206**, 251-259 (2005).

738   42   Rowe, T. *et al.* Macaque model for severe acute respiratory syndrome. *J Virol* **78**, 11401-11404

739   (2004).

740   43   Roberts, A. *et al.* Animal models and vaccines for SARS-CoV infection. *Virus Res* **133**, 20-32,

741   doi:10.1016/j.virusres.2007.03.025 (2008).

742   44   Follis, K. E., York, J. & Nunberg, J. H. Furin cleavage of the SARS coronavirus spike

743   glycoprotein enhances cell-cell fusion but does not affect virion entry. *Virology* **350**, 358-369

744   (2006).

745   45   Thomas HC Sit, C. J. B., Sin Ming Ip, Karina WS Tam, Pierra YT Law, Esther MW To, Veronica

746   YT Yu, Leslie D Sims, Dominic NC Tsang, Daniel KW Chu, Ranawaka APM Perera, Leo LM

747   Poon, Malik Peiris. Canine SARS-CoV-2 infection. *Research Square*

748   **DOI:10.21203/rs.3.rs-18713/v1** (2020).

749   46   Lun, Z. R. & Qu, L. H. Animal-to-human SARS-associated coronavirus transmission? *Emerg*

750   *Infect Dis* **10**, 959, doi:10.3201/eid1005.040022 (2004).

751   47   Safronetz, D., Geisbert, T. W. & Feldmann, H. Animal models for highly pathogenic emerging

752   viruses. *Curr Opin Virol* **3**, 205-209, doi:10.1016/j.coviro.2013.01.001 (2013).

753   48   Linlin Bao, W. D., Baoying Huang, Hong Gao, Lili Ren, Qiang Wei, Pin Yu, Yanfeng Xu,

754   Jiangning Liu, Feifei Qi, Yajin Qu, Wenling Wang, Fengdi Li, Qi Lv, Jing Xue, Shuran Gong,

755   Mingya Liu, Guanpeng Wang, Shunyi Wang, Linna Zhao, Peipei Liu, Li Zhao, Fei Ye, Huijuan

756   Wang, Weimin Zhou, Na Zhu, Wei Zhen, Xiaojuan Zhang, Zhiqi Song, Li Guo, Lan Chen,

757   Conghui Wang, Ying Wang, Xinming Wang, Yan Xiao, Qiangming Sun, Hongqi Liu, Fanli Zhu,

758   Chunxia Ma, Lingmei Yan, Mengli Yang, Jun Han, Wenbo Xu, Wenjie Tan, Xiaozhong Peng, Qi

759   Jin, Guizhen Wu, Chuan Qin. The Pathogenicity of 2019 Novel Coronavirus in hACE2

Transgenic Mice. *BioRxiv* **doi: https://doi.org/10.1101/2020.02.07.939389** (2020).

McCray, P. B., Jr. *et al.* Lethal infection of K18-hACE2 mice infected with severe acute respiratory syndrome coronavirus. *Journal of virology* **81**, 813-821, doi:10.1128/JVI.02012-06 (2007).

Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **10.1016/S0140-6736(20)30183-5**, doi:10.1016/S0140-6736(20)30183-5 (2020).

Callaway, E. Labs rush to study coronavirus in transgenic animals - some are in short supply. *Nature* **579**, 183, doi:10.1038/d41586-020-00698-x (2020).

Day, C. W. *et al.* A new mouse-adapted strain of SARS-CoV as a lethal model for evaluating antiviral agents in vitro and in vivo. *Virology* **395**, 210-222, doi:10.1016/j.virol.2009.09.023 (2009).

Roberts, A. *et al.* A mouse-adapted SARS-coronavirus causes disease and mortality in BALB/c mice. *PLoS Pathog* **3**, e5, doi:10.1371/journal.ppat.0030005 (2007).

Jiang, D. *et al.* Identification of three interferon-inducible cellular enzymes that inhibit the replication of hepatitis C virus. *J Virol* **82**, 1665-1678, doi:JVI.02113-07 [pii]10.1128/JVI.02113-07 (2008).

Li, W. *et al.* Efficient replication of severe acute respiratory syndrome coronavirus in mouse cells is limited by murine angiotensin-converting enzyme 2. *J Virol* **78**, 11429-11433 (2004).

Babcock, G. J., Esshaki, D. J., Thomas, W. D., Jr. & Ambrosino, D. M. Amino acids 270 to 510 of the severe acute respiratory syndrome coronavirus spike protein are required for interaction with receptor. *J Virol* **78**, 4552-4560 (2004).

Lin, H. X. *et al.* Identification of residues in the receptor-binding domain (RBD) of the spike protein of human coronavirus NL63 that are critical for the RBD-ACE2 receptor interaction. *J Gen Virol* **89**, 1015-1024, doi:10.1099/vir.0.83331-0 (2008).

Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673-4680 (1994).

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858, doi:10.1038/nprot.2015.053 (2015).