Jacalin domain-containing protein *Os*SalT interacts with *Os*DREB2A and *Os*NAC1 to impart drought stress tolerance *in planta*

Salman Sahid [1, 2, †], Chandan Roy [1, †], Dibyendu Shee [1], Riddhi Datta [2] *, Soumitra Paul [1] *

[1]*Molecular Plant Physiology Laboratory, Department of Botany, University of Calcutta, 35 Ballygunge Circular Road, Kolkata-700019, West Bengal, India*

[2]*Department of Botany, Dr. A. P. J. Abdul Kalam Government College, Plot BA1, Action area 1, New Town, Kolkata-700156, West Bengal, India*

*Authors for correspondence:

Email: riddhi.bot@gmail.com, psoumitra@ymail.com; Telephone No: +919433084074, +919748605305; Fax No: +9133 2461 4849

ORCID IDs: Riddhi Datta: 0000-0001-7490-190X; Soumitra Paul: 0000-0001-7560-5512

[†] Equal Contribution

Key message:

Ectopic expression of *Os*SalT significantly enhances drought tolerance in tobacco. It interacts with *Os*DREB2A and *Os*NAC1 augmenting drought-responsive gene expression to impart stress tolerance *in planta*.

Declarations

Funding

This work was supported by Department of Science and Technology and Biotechnology, Government of West Bengal, India [BT(Budget)/RD-29/2016].

Conflict of interest

The authors declare no conflict of interest.

Acknowledgement

We thank the central instrumentation facilities of Department of Botany, University of Calcutta and Department of Botany, Dr. A. P. J. Abdul Kalam Government College as well as the confocal microscopic facility of DBT-IPLS, Department of Biochemistry, University of Calcutta. We thank Prof. Jörg Kudla (University of Munster, Germany) for providing the *pVYNE* and *pVYCE* vectors and Prof. Prabod Trivedi (CSIR-National Botanical Research Institute, India) for kindly sharing the *Agrobacterium tumefaciens* GV3101 strain.

Author Contributions

RD and SP conceived and designed the research plan; SS and CR performed most of the experiments; DS performed the *in silico* analysis; RD and SP analyzed the data; SS drafted the manuscript; RD and SP supervised and complemented the writing.

Abstract

With the changing climatic conditions, drought has become one of the most threatening abiotic stress factors that adversely affect rice cultivation and productivity. Although the involvement of the jacalin domain-containing protein, *Os*SalT, has been reported in drought and salinity tolerance, its functional mechanism still remains largely unexplored. In this study, expression of the *OsSalT* gene was found to be positively correlated with the drought tolerance potential with its higher transcript abundance in the tolerant *indica* rice cultivar, Vandana and lower abundance in the susceptible cultivar, MTU1010. Moreover, the ectopic expression of *Os*SalT in tobacco imparted drought stress tolerance in the transgenic lines. The transgenic lines exhibited significantly improved growth and higher osmolyte accumulation over the wild-type (WT) plants together with the induction in the *Os*SalT expression under drought stress. Fascinatingly, the yeast two-hybrid and bimolecular fluorescence complementation (BiFC) analyses confirmed the interaction of *Os*SalT protein with two interesting transcription factors (TFs), *Os*NAC1 and OsDREB2A. *In silico* analysis further revealed that the *Os*SalT protein interacted with the regulatory domain of *Os*DREB2A and the C-terminal domain of *Os*NAC1 leading to their activation and induction of their downstream drought-responsive genes. Together, this study unravels a novel model for *Os*SalT-mediated regulation of drought tolerance in plants.

Keywords: Drought stress, *Os*SalT, jacalin, lectin, DREB2A, NAC1.

Introduction

Drought is among the most important abiotic stress factors that affect crop productivity resulting in significant yield losses (Hu and Xiang 2014). The changing climatic conditions like global warming and insufficiency of proper rainfall adversely affect plant growth and development compromising productivity. Plants have evolved a highly sophisticated adaptive mechanism to combat drought stress. In-depth analysis of drought tolerance mechanism suggests an intricate crosstalk between different signalling cascades in plants. The drought stress response in plants is regulated by two distinct signalling pathways - the abscisic acid (ABA)-dependent and ABA-independent pathways. Several important transcription factors (TFs) such as *myelocytomatosis oncogene* (*MYC*), *myeloblastosis oncogene* (*MYB*), *basic leucine zipper* (*bZIP*), *NAM, ATAF, and CUC* (*NAC*) and *dehydration responsive element binding* (*DREB*) function in both the pathways and are highly accumulated in response to drought stress. These TFs are known to modulate downstream osmotic stress-responsive genes to render stress tolerance (Nakashima et al. 2009; Atkinson and Urwin 2012).

Lectins are a family of carbohydrate-binding proteins that specifically recognize sugar moieties. Lectins are known to play diverse roles in plant defence, in addition to its involvement in various other physiological phenomena. For example, lectin proteins were found to be induced in response to heat shock in cell suspension cultures (Spadoro-Tank and Etzler 1988; Shakirova et al. 1996). Again, transgenic Thai rice cultivars ectopically expressing snowdrop lectin exhibited resistance to sap-sucking insects (Tinjuangjun et al. 2000). ASAL, a novel lectin from garlic, provided resistance against phloem-limited viruses in rice (Saha et al. 2006). In addition, a large number of pattern recognition receptors in plants contain lectin domains that are essential for pathogen recognition and defence (Kaku et al. 2006; Bellande et al. 2017).

Plant lectins usually constitute a heterogeneous group that has been classified into 7 subfamilies based on their structure and evolutionary relatedness (van Damme et al. 1998). However, a later study has recognized 12 families among which 4 families encompassed the newly identified lectin members (Jiang et al. 2010). Among them, the jacalin-related lectin (JRL) subfamily is characterized by the presence of a conserved disaccharide-binding jacalin domain that was first identified in jackfruit (*Artocarpus integrifolia*) (de Azevedo Moreira and Ainouz 1981). The JRL subfamily comprises of at least 30 JRL proteins in rice distributed over its 9 out of 12 chromosomes. Among them 16 members contain only one

jacalin domain and are called merojacalins. Thirteen members are chimeric jacalins containing at least one of the NB_ARC, Pkinase, dirigent or peptidase domains, in addition to the jacalin domain. The remaining one member has 3 repeats of the jacalin domain and is known as holojacalin (Yi-juan et al. 2018). The JRL proteins can again be categorized into two groups: the galactose-binding JRLs (gJRLs) with vacuolar localization and the mannose-binding JRLs (mJRLs) with nucleo-cytoplasmic localization (Van Damme et al. 2004; Lannoo and Van Damme 2014).

Members of the JRL subfamily have been demonstrated to play a crucial role in plant defence. Rice lectins containing mannose-binding jacalin domain have been demonstrated to play a role in salinity stress (Zhang et al. 2000). A map-based cloning strategy revealed that the *Jacalin-type Lectin Required for Potexvirus Resistance 1* (*JAX1*) gene confers resistance to *Plantago asiatica* mosaic virus infection in *Arabidopsis* (Yamaji et al. 2012). Chimeric JRLs have been reported to enhance tolerance against pathogen invasion, insect attack as well as abiotic stress in monocots (Song et al. 2014). Another jacalin like lectin protein, *Os*JAC1 imparted a high level of resistance against *Magnaporthe* infection in rice (Weidenbach et al. 2016).

An interesting JRL subfamily member, the *Os*SalT protein, is a 14.5 kDa mannose-binding lectin which comprises a single jacalin domain. It has been identified to accumulate in the root and sheath tissues of rice in response to salinity and drought stress (Claes et al. 1990; Filho et al. 2003). Transgenic rice plants overexpressing *OsSalT* gene have been shown to significantly improve salinity tolerance (He et al. 2016). In addition, a proteomic study has revealed that *Os*SalT was up-accumulated in rice roots under drought stress conditions (Paul et al. 2015). These pieces of evidence suggest a possible role of *Os*SalT protein in drought tolerance. However, the mechanism of how it imparts drought tolerance in plants has not been investigated so far.

In the present study, we have aimed to decipher the molecular mechanism of the drought-responsive role of *Os*SalT protein. Ectopic expression of *OsSalT* gene in tobacco significantly enhanced drought tolerance. We further identified that the *Os*SalT protein interacts with *Os*DREB2A and *Os*NAC1 TFs which induce the downstream drought-responsive gene expression thus imparting stress tolerance in plants.

Materials and Methods

Plant material, growth condition and stress assay

Seeds of *indica* rice cultivars, Vandana and MTU1010, were procured from the Chinsurah Rice Research Station, West Bengal, India. Seeds were germinated and grown in the net house under optimum conditions. The rice cultivars were normally irrigated for several weeks after transplanting. For stress assay, 100 days old plants were exposed to drought by the withdrawal of watering for 7 days until the soil moisture content reached 45 % as standardized before (Paul et al. 2015).

Seeds of tobacco (*Nicotiana tabaccum* L. *cv. Xanthi*) were germinated and grown in Murashige and Skoog (MS) medium under 16 h light/ 8 h dark photoperiod as standardized before (Murashige and Skoog 1962; Ghanta et al. 2013). For stress assay, 60 days old soil-grown plants were subjected to drought stress by complete withdrawal of watering for 5 days until soil moisture content reached 45 %.

Morphological analysis

Different agronomic parameters such as plant height, percentage of brown leaves, percentage of rolled leaves, shoot and root dry weight were recorded for the two rice cultivars under control and drought conditions. For tobacco, plant height, percentage of yellow leaves, percentage of rolled leaves, shoot and root dry weight were recorded for wild-type (WT) and transgenic (OX) lines.

Biochemical analysis

For estimation of proline content, 0.2 g leaf tissue was collected from each sample and extracted with ethanol. The supernatant was mixed with 1% ninhydrin solution (in 60% acetic acid) and 20% ethanol and incubated at 95 °C for 30 min. The optical density was measured in spectrophotometer (Hitachi) at 595 nm (Woodrow et al. 2016). The leaf samples were homogenized in 0.05 % toluene and glycine betaine content was estimated following Grieve and Grattan (1983). Soluble sugar content was estimated according to Dubois (1956). Briefly, samples were homogenized in double distilled water, boiled at 100 °C for 30 min and centrifuged at 5000 rpm for 5 min. The supernatant was mixed with 9% phenol and 98% sulfuric acid, incubated at room temperature for 30 min and optical density was recorded at 485 nm. Total chlorophyll was extracted with 80 % acetone and quantified according to Lichtenthaler (1987).

RNA Isolation and q-RT PCR

Total RNA was isolated following Trizol method. Complementary DNA was synthesized using iScript cDNA synthesis kit (Bio-rad) following the manufacturer's instructions. The qPCR analysis was performed in CFX96 Real-time PCR Detection System (Bio-rad) by using iTaq Universal SYBR Green Supermix (Bio-rad) and gene-specific primers (Table S1). *Actin* was used as the reference gene.

Cloning, vector construction and raising of transgenic lines

The *OsSalT* gene was cloned into pGEMT-Easy vector (Promega) followed by sub-cloning into the *EcoRI* and *BamHI* RE sites of pEGAD vector under control of *CaMV35S* promoter. The resulting construct, *CaMV35S::OsSalT-GFP*, was transformed into tobacco using *Agrobacterium*-mediated (*Agrobacterium tumefaciens* GV3101) leaf disc method as standardized before (Ghanta et al. 2013). The putative transgenic plants were screened through genomic DNA PCR with *bar* gene-specific primer. The positive transgenic lines were maintained up to T_2 generation which was used for further experiments. The ectopic expression of the transgene was confirmed by qRT-PCR analysis using gene-specific primers.

Yeast two hybrid assay

Yeast two hybrid assay was performed using Matchmaker Gold Yeast Two-Hybrid System (Clontech). The *OsSalT* gene was cloned in-frame into the *EcoRI* and *BamHI* RE sites of the yeast vector, pGBKT7 to obtain a *BD-OsSalT* construct as bait. To study the protein-protein interaction of *OsSalT*, a cDNA library, ligated to pGADT7-*Rec* as prey, was prepared from rice roots under drought stress using Make Your Own "Mate and Plate" Library System (Clontech). The constructs were co-transformed into Y2H Gold yeast strain and screened on DDO (SD/-Leu/-Trp) and QDO (SD/–Ade/–His/–Leu/–Trp) plates. This was followed by a QDO/X/A (SD/–Ade/–His/–Leu/–Trp/X-α-Galactosidase/Aureobasidin-A) selection to test interaction. To identify the interacting protein partner(s), selected colonies were analysed by PCR followed by sequencing according to the manufacturer's instruction. The *AD-T-antigen* and *BD-p53* interaction was used as a positive control.

Bimolecular fluorescence complementation (BiFC) assay

The *OsSalT* gene was cloned into the *35S::pVYCE-cVenus* vector between *SpeI* and *BamHI* RE sites to obtain *35S::OsSalT-cVenus* construct. The interacting partners, *OsDREB2A* and

OsNAC1 were cloned into *35S::pVYNE-nVenus* vector between the same RE sites to obtain *35S::OsDREB2A-nVenus* and *35S::OsNAC1-nVenus* constructs (Waadt et al. 2008). These constructs were introduced into the onion epidermal cells in pairwise combinations according to Yang et al. (2014). The *35S::OsSalT-cVenus* construct with empty *35S::pVYNE-nVenus* vector was used as negative control. Samples were examined under laser scanning confocal microscope (Olympus FV1000-IX81) using excitation wavelength of 514 nm.

Fold prediction and homology modelling of *Os*SalT, *Os*DREB2A and *Os*NAC1 proteins

The *Os*SalT (Uniprot ID: Q0JMY8), *Os*DREB2A (Uniprot ID: Q0JQF7) and *Os*NAC1 (Uniprot ID: Q8H0I5) sequences were subjected to PSIPRED (Buchan and Jones 2019), PHYRE2 (Kelley et al. 2015) and ROBETTA (Song et al. 2013) for fold prediction and homology modelling. The 3D structures of the protein sequences were filtered or energy minimized by YASARA server (Krieger et al. 2009). Models were further validated by PROCHECK (Laskowski et al. 1996) using the Ramachandran plot. According to Ramachandran plot, the residues of the disordered region was further refined by using MODELLER based ModLoop (Fiser and Sali 2003) webserver (only the loop regions of a 3D model were refined). The final models were re-validated by PROCHECK.

Molecular docking analysis

The best-fitted model of *Os*DREB2A and *Os*NAC1 were used for molecular docking with *Os*SalT through protein-protein docking server PatchDock (Schneidman-Duhovny et al. 2005) and GRAMM-X (Tovchigrechko and Vakser 2006). Top 100 dock solutions were generated by the PatchDock web server. Solutions were further refined and rescored using the FireDock docking refinement server (Mashiach et al. 2008). The best dock solution was selected according to its lowest global energy. Approximate ΔG of binding of best selected docked solutions was calculated by using the PDBePISA server (Krissinel and Henrick 2007).

Sequence analysis and prediction of ubiquitination and SUMOylation sites

The protein sequence for *Os*DREB2A was analyzed manually for the presence of NRD domain on the basis of existing reports (Mizoi et al. 2019). The DNA-binding domain residues were retrieved from NCBI database. Ubiquitination sites were predicted using UbiSite prediction server (http://csb.cse.yzu.edu.tw/UbiSite/prediction.php) with specificity

level high. For SUMOylation consensus site prediction, GPS-SUMO web server (Ren et al. 2009; Zhao et al. 2014) was used with specificity level set as high.

Statistical Analysis:

All experiments were performed in three independent biological replicates and data were represented as mean ± standard error of mean (SEM). Differences in morphological parameters, metabolite contents and transcript abundances among genotypes and treatments were assessed by two-way ANOVA followed by Sidak's multiple comparison tests using the GraphPad Prism version 8.0.0 (GraphPad Software, San Diego, California USA, www.graphpad.com). The statistical significance at P ≤ 0.05 to identify the difference between the two sets of data was considered.

Results

Expression of *OsSalT* gene positively co-related with the drought tolerance potential in two rice cultivars

Two *indica* rice cultivars, Vandana and MTU1010 were analyzed for their drought tolerance potential. Several morphological parameters such as plant height, percentage of rolled and brown leaves, and root and shoot dry weights were estimated under control and drought conditions. No significant alteration in plant height and shoot dry weight was observed in response to drought stress in both the varieties. The root dry weight increased in Vandana under drought stress conditions while no alteration was observed for MTU1010 roots. The percentage of rolled and brown leaves was, however, significantly lower in the case of Vandana in comparison to MTU1010 under drought stress conditions (Fig. 1).

To determine the effect of drought stress on the two cultivars, different osmotic stress-related metabolites such as proline, soluble sugars, glycine betaine, and chlorophyll contents were estimated. The total chlorophyll content decreased in response to drought stress in both the cultivars and the content was considerably lower in MTU1010 in comparison to Vandana. The proline and glycine betaine contents increased in response to drought stress in both the cultivars in a similar pattern. In case of soluble sugars, however, the content was more steeply increased in case of Vandana (97.29 mg/g FW to 158.8 mg/g FW) than in MTU1010 (55.7 mg/g FW to 92.49 mg/g FW) in response to drought stress (Fig. 2A-D). Together, these morphological and biochemical analyses indicate that Vandana is significantly more tolerant to drought stress in comparison to MTU1010.

Furthermore, the expression of *OsSalT* along with *OsOsmotin*, a marker gene for drought stress, was analyzed in both Vandana and MTU1010 under drought stress. It was observed that the relative transcript abundance of *OsSalT* gene was increased by 15.61 fold in Vandana and by 5.61 fold in MTU1010 in response to drought stress. In the case of the *OsOsmotin* gene, the relative transcript abundance was significantly higher in Vandana in comparison to MTU1010 under both control and drought conditions (Fig. 2E-F). These observations suggested a positive correlation of *OsSalT* expression with the drought tolerance potential in rice.

Ectopic expression of *OsSalT* gene improved the drought tolerance in tobacco

To further validate the involvement of the *OsSalT* gene in imparting drought tolerance, the gene was ectopically expressed in tobacco. Among the 17 putative transformants generated, 11 lines were found to be positive (Fig. S1). Three best transgenic lines (OX1, OX2, and OX3) with the highest expression of *OsSalT* were considered for further experiments. Morphological analysis revealed no variations in these OX lines as compared to the WT and the vector control (VC) plants.

Stress assay was performed to assess the drought tolerance potential of the OX lines. It was observed that all the three OX lines were significantly more tolerant to drought stress in comparison to the WT and VC plants. The morphological analysis demonstrated that there was no alteration in shoot dry weight in the OX, VC and WT plants under control condition. However, the shoot dry weight was considerably higher in the case of the OX lines in comparison to the VC and WT plants under drought stress. The percentage of wilted leaves and yellow leaves were much higher in WT and VC plants than the OX lines (Fig. 3A-D).

The accumulation of several osmotic stress-related metabolite contents under drought stress was also investigated. It was observed that the drop in the total chlorophyll content under drought stress was much steeper in case of the WT and VC plants in comparison to the OX lines. Accumulation of other metabolites like proline, glycine betaine and soluble sugars were also induced under drought stress condition. However, the induction was significantly higher in the OX lines than the WT and VC plants in response to drought stress (Fig. S2).

Besides, the expression of *OsSalT* in the WT, VC and OX lines were analyzed. As expected no expression was detected in the WT and VC plants. In the case of the OX1, OX2 and OX3 lines, the *OsSalT* expression was induced by 6.26, 4.47 and 6.76 folds respectively in

response to drought stress (Fig. 3E). Together, these observations confirmed that the ectopic expression of *OsSalT* gene significantly improved drought stress tolerance in tobacco.

OsSalT protein interacts with OsDREB2A and OsNAC1 to impart drought tolerance

To explore the mechanism of how *OsSalT* imparts drought tolerance, the interacting protein partners were identified using yeast two-hybrid assay. A rice root library constructed under drought stress condition was used for this study. Out of 9 recombinant cDNA clones, 3 non-redundant putative interacting proteins partners were identified. Interestingly, two proteins, *Os*DREB2A and *Os*NAC1, were identified as positive independent interacting partners of *Os*SalT protein (Fig. 4A). The co-expression of *BD-OsSalT* with *AD-OsDREB2A* and *AD-OsNAC1* exhibited strong activation of X-α-galactosidase and aureobasidin A in the QDO/X/A medium. The interaction of *BD-p53* with *AD-T-antigen* was used as a positive control.

Moreover, the interaction between *Os*SalT with *Os*DREB2A and *Os*NAC1 was confirmed by BiFC analysis (Fig. 4B). When the onion epidermal cells were co-transfected with the *35S::OsSalT-cVenus* and *35S::OsDREB2A-nVenus* constructs, strong yellow fluorescence was observed in the nucleus thus confirming a nuclear interaction. In the case of the *35S::OsSalT-cVenus* and *35S::OsNAC1-nVenus* constructs, yellow fluorescence was also detected in the nucleus indicating a nuclear interaction. However, no fluorescence was detected when the *35S::OsSalT-cVenus* construct was used with the empty *35S::pVYNE-nVenus* vector. These results confirmed the physical interaction of the *Os*SalT protein with *Os*DREB2A and *Os*NAC1 to regulate drought tolerance in plants.

***In silico* analysis confirmed the physical interaction of OsSalT protein with OsDREB2A and OsNAC1**

The 3D structures for *Os*SalT, *Os*DREB2A and *Os*NAC1 proteins were generated by homology modelling followed by energy minimization. The quality of the models was assessed using PROCHECK server. Ramachandan plot of the complete structures revealed that 91.2 %, 90.3% and 87.6 % residues were present in the most favoured regions in case of the *Os*SalT, *Os*DREB2A and *Os*NAC1 proteins respectively. None of the residues were present in the disallowed region (Fig. S3-S8). Protein-protein interaction for *Os*SalT-*Os*DREB2A and *Os*SalT-*Os*NAC1 was studied *in silico* through molecular docking using PatchDock and GRAMM-X servers. The docking results confirmed physical interaction of

the *Os*SalT protein with *Os*DREB2A and *Os*NAC1 proteins (Fig. 5-6). Sequence analysis along with the molecular docking data indicated that *Os*NAC1 protein possesses its DNA-binding domain at the N-terminal end while it interacts with the *Os*SalT protein at its C-terminal domain (Fig. S9). The docking analysis further revealed that the *Os*SalT protein binds to a regulatory region of the *Os*DREB2A protein without blocking its DNA-binding domain (residues 75-132) (Fig. 5, S10). This interaction presumably activates the *Os*DREB2A protein under drought stress thus enabling it to induce the transcription of the DREB-responsive genes.

It has been reported that the NRD domain of most DREB proteins contain an acidic sequence containing a stretch of at least five serine/threonine residue cluster (Mizoi et al. 2019). Analysis of the protein sequence revealed that *Os*DREB2A protein, unlike *Os*DREB2B and most DREB proteins in other plants, lacks a well-defined NRD regulatory region indicating that it may possess unique regulatory sites. Next, its structure was analyzed for probable ubiquitination and SUMOylation sites. It was observed that the protein possess three probable ubiquitination sites at K184, K185 and K199 (Fig. S10). In addition, a SUMOylation consensus site at K199 had been predicted which indicated a probable regulation via ubiquitination-SUMOylation switch (Fig. S10). It was further observed that this K199 does not overlap with both the DNA-binding as well as the *Os*SalT-interacting sites. These observations suggested a unique regulatory mechanism for *Os*DREB2A protein in rice.

Expression of *DREB2A* and *NAC1* responsive genes were enhanced in transgenic tobacco plants (OX) under drought stress

To further confirm the *Os*SalT-mediated activation of DREB2A and NAC1 proteins, expression of several stress-responsive genes were analyzed under drought condition in the OX plants. It is known that the transcriptions of the *Dehydrin* and *Late Embryogenesis Abundant* (LEA) genes are induced by DREB2A while the *Phosphatidylinositol-3-kinase* (*PI3K*) and *Sucrose phosphate synthase* (*SPS*) are NAC1-responsive genes (Lata and Prasad 2011; Saad et al. 2013). Therefore, these genes were selected in this analysis. In addition, *Responsive to dehydration 22* (*rd22*) and *Alcohol dehydrogenase 1* (*ADH1*) genes, which function in DREB2A and NAC1 independent manner, were selected as negative control (Abe et al. 2003). Interestingly, it was observed that the expression of *Dehydrin*, *LEA*, *PI3K* and *SPS* genes were significantly enhanced in the OX plants under drought stress as compared to the WT plants (Fig. 7). On the contrary, expression of both *rd22* as well as *ADH1* genes was

not significantly altered. This confirmed that *Os*SalT activates both DREB2A and NAC1 proteins that in turn results in the induction of the downstream drought-responsive genes ultimately leading to stress tolerance.

Discussion

In plants, lectin proteins are widely known to be involved in various biotic and abiotic stress responses. Jacalin is a mannose-binding lectin protein that serves an important role in regulating blast disease and osmotic stress tolerance in rice (Zhang et al. 2000; Song et al. 2014). In this study, we have dissected the role of *Os*SalT, a jacalin domain containing lectin protein, in drought stress tolerance in plants. Earlier, it had been identified that the expression of *OsSalT* was augmented in rice in response to salt stress. The *OsSalT* overexpressing rice plants also displayed enhanced osmotic stress tolerance (Filho et al. 2003; He et al. 2016). However, how this *Os*SalT protein provides osmotic stress tolerance in plants remains elusive so far. Here, we have explored the molecular mechanism of *Os*SalT protein in regulating drought stress tolerance *in planta*.

To identify the drought stress-responsive role of *Os*SalT protein, its expression was analyzed in response to drought stress in a drought tolerant (Vandana) and a drought susceptible (MTU1010) *indica* rice cultivar. The expression of the *OsSalT* gene was found to be significantly higher in the drought-tolerant cultivar in comparison to the susceptible one under drought stress. This observation suggested a direct correlation of the *OsSalT* expression with the degree of drought tolerance in rice. Furthermore, the ectopic expression of *OsSalT* gene provided enhanced drought tolerance in tobacco. The OX plants displayed an improved wilting percentage and less chlorophyll degradation. The higher accumulation of several osmotically active compounds has long been associated with osmotic stress tolerance potential in plants (Chen and Murata 2002). In the present study, the OX lines exhibited an elevated accumulation of proline, glycine betaine and soluble sugar contents under drought stress which indicated their better adaptability over the WT plants.

With this confirmation, we aimed to decipher the molecular mechanism of how the *Os*SalT protein imparts drought tolerance in plants. It was identified that the *Os*SalT is a nucleo-cytoplasmic protein that interacts with two interesting protein partners, *Os*DREB2A and *Os*NAC1 in rice roots under drought stress conditions. It has been widely demonstrated that both of these protein partners function as important TFs and can regulate the expression of different drought and thermotolerance-associated genes (Liu et al. 1998; Dubouzet et al.

2003; Saad et al. 2013). It was further observed that its interaction with both *Os*DREB2A and *Os*NAC1 occurs in the nucleus.

Emerging evidences suggested that NAC family comprises of a large group of plant-specific TFs named upon 3 protein members, NAM, APAF1-2, and CUC2. They play a pivotal role in several abiotic stress responses in plants (Aida et al. 1997). An array of NAC family members have been identified from different plant species including rice. In rice, 151 NAC family genes were reported while most of them were characterized as stress-responsive NAC genes (sNAC) (Nuruzzaman et al. 2010). Recently, the drought tolerance activity of *OsNAC1* has been confirmed in bread wheat (Saad et al. 2013). In the present study, it was identified that the jacalin domain-containing protein, *Os*SalT, can activate the *Os*NAC1 protein in rice in response to drought stress. The *Os*NAC1 protein was found to possesses an N-terminal DNA binding domain which binds to several drought- or salinity-responsive genes (Aida et al. 1997). On the other hand, the *Os*SalT protein was found to interact with its C-terminal end to modulate its function. This observation can be corroborated with the previous report that all NAC proteins contain a highly conserved N-terminal DNA-binding domain and a variable C-terminal regulatory region (Kikuchi et al. 2000; Ooka et al. 2003). This study delineates a probable mechanism of *Os*NAC1 regulation in response to drought stress in rice.

Previous investigations highlighted that several *DREB* subfamily members regulate osmotic stress tolerance in plants (Dubouzet et al. 2003; Lata and Prasad 2011). It has been demonstrated that these DREB proteins are specifically regulated via modulation of their protein stability. Although the induction of *Arabidopsis DREB2A* gene expression was found to be significant under drought stress, the activation of downstream signalling was completely dependent on the stability of the DREB2A protein. This protein stability has been reported to be maintained by phosphorylation and ubiquitination at a regulatory NRD domain (Shinozaki and Yamaguchi-Shinozaki 2000; Sakuma et al. 2002; Yamaguchi-Shinozaki and Shinozaki 2006; Mizoi et al. 2019). Recently, it has been showed that SUMOylation at the K163 amino acid residue close to the NRD domain inhibited the ubiquitination during stress and stabilised the protein thus imparting thermotolerance in *Arabidopsis* (Wang et al. 2020). However, the mechanism of its activation during stress still remains unanswered. In addition, most of the studies dissecting the DREB2A regulation have been restricted to *Arabidopsis* and no detailed activation mechanism for *Os*DREB2A protein during drought stress has been undertaken.

Interestingly, the *Os*DREB2A protein in rice, unlike the *Os*DREB2B, does not possess any conserved NRD domain and the activation mechanism has not been studied so far. This motivated us to explore the *Os*DREB2A structure through *in silico* analysis. It was observed that the protein still possesses three probable ubiquitination sites at K184, K185 and K199 among which K199 is also a SUMOylation consensus site. This observation suggested that in spite of lacking a conserved NRD domain, the *Os*DREB2A protein can still be regulated by an ubiquitination-SUMOylation switch. The balance between these two phenomena maintains the *Os*DREB2A protein accumulation in the cell to a basal level under control condition while quickly augmenting its level under drought stress.

In addition, *Os*SalT interacted with *Os*DREB2A protein at a regulatory domain, other than the DNA-binding domain and the K199 residue. This presumably activated the protein to regulate its downstream drought stress-responsive genes. *Os*DREB2A is a TF that can bind to the dehydrin responsive protein (DRE) motif of different drought stress-responsive genes like *Dehydrin* and *LEA* to regulate drought stress tolerance (Lata and Prasad 2011). In this study, it was also observed that the DREB2A responsive genes, *Dehydrin* and *LEA* were significantly up-regulated in the tobacco OX lines suggesting that *Os*SalT presumably activates the DREB2A protein in tobacco which in turn activates the downstream drought-responsive genes. In fact, a proteomic study of *Os*DREB2A overexpressing rice plants identified up-accumulation of the *Os*SalT protein in roots under drought stress which corroborates with the present observation (Paul et al. 2015).

In summary, the present investigation has demonstrated the functional role of *Os*SalT protein in imparting drought stress tolerance via its interaction with *Os*DREB2A and *Os*NAC1 proteins (Fig. 8). It can be hypothesized that under drought stress, the ubiquitination of *Os*DREB2A is blocked due to SUMOylation of a specific lysine residue. This inhibits *Os*DREB2A degradation thus augmenting its accumulation in the cell. The *Os*SalT protein then interacts with the stabilized *Os*DREB2A. The binding of *Os*SalT protein probably leads to a conformational change in the *Os*DREB2A protein which facilitates its activation under drought stress. It can then bind to the DRE sequence of the drought stress responsive genes leading to their transcriptional activation. On the other hand, drought stress induces the expression of *Os*NAC1. The *Os*SalT protein then interacts with and activates *Os*NAC1 protein which in turn induces the expression of its downstream drought-responsive genes.

Together, the present study has revealed the novel role of a jacalin domain containing *Os*SalT protein in regulating two crucial TFs thus enhancing the drought stress tolerance *in planta*.

Supplementary data

Supplementary Fig. S1. Screening of putative transformants by genomic DNA PCR for *bar* gene. A prominent band at 412 bp indicated positive transgenic lines. L: 1 kb DNA ladder, N: negative control (WT), P: positive control (recombinant plasmid), 1-17: putative transformants.

Supplementary Fig. S2: Biochemical analysis of transgenic tobacco lines ectopically expressing OsSalT gene under drought stress condition. (A) Chlorophyll content, (B) Proline content, (C) Glycine betaine content, and (D) Soluble sugar content. Results were represented as mean±SEM (n=3). Statistical difference between the cultivars under control and drought stress was denoted by asterisks at $p < 0.0001$ (****).

Supplementary Fig. S3: Homology modelling of *Os*SalT protein. (A) Fold prediction using PSIPRED, Surface view of the *Os*SalT protein in (B) side view and (C) top view.

Supplementary Fig. S4: Ramachandran plot for *Os*SalT structure.

Supplementary Fig. S5: Homology modelling of *Os*DREB2A protein. (A) Fold prediction using PSIPRED, Surface view of the *Os*DREB2A protein in (B) side view and (C) top view.

Supplementary Fig. S6: Ramachandran plot for *Os*DREB2A structure.

Supplementary Fig. S7: Homology modelling of *Os*NAC1 protein. (A) Fold prediction using PSIPRED, Surface view of the *Os*NAC1 protein in (B) side view and (C) top view.

Supplementary Fig. S8: Ramachandran plot for *Os*NAC1 structure.

Supplementary Fig. S9: Sequence analysis for *Os*NAC1 protein showing the DNA-binding and OsSalT-interacting sites.

Supplementary Fig. S10: Sequence analysis for *Os*DREB2A protein. (A) Protein sequence showing the DNA-binding, OsSalT-interacting, ubiquitination and SUMOylation consensus sites. (B) Ubiquitination site prediction. (C) SUMOylation site prediction.

Supplementary Table S1: List of primers used.

Reference

Abe H, Urao T, Ito T, Seki M, Shinojaki K, Yamaguchi-Shinojaki K (2003) *Arabidopsis* AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. Plant Cell 15:63-78. https://doi.org/10.1105/tpc.006130

Aida M, Ishida T, Fukaki H, Fujisawa H, Tasaka M (1997) Genes involved in organ separation in *Arabidopsis*: an analysis of the cup-shaped cotyledon mutant. Plant Cell 9:841–857. https://dx.doi.org/10.1105/tpc.9.6.841

Atkinson NJ, Urwin PE (2012) The interaction of plant biotic and abiotic stresses: from genes to the field. J Exp Bot 63:3523–3543. https://doi.org/10.1093/jxb/ers100

Bellande K, Bono JJ, Savelli B, Jamet E, Canut H (2017) Plant lectins and lectin receptor-like kinases: how do they sense the outside? Int J Mol Sci 18:1164. https://doi.org/10.3390/ijms18061164

Buchan DWA, Jones DT (2019) The PSIPRED Protein Analysis Workbench: 20 years on. Nucleic Acids Res 47:W402-W407. https://doi.org/10.1093/nar/gkz297

Chen TH, Murata N (2002) Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. Curr Opin Plant Biol 5:250–257. https://doi.org/10.1016/s1369-5266(02)00255-8

Claes B, Dekeyser R, Villarroel R, Van den Bulcke M, Bauw G, Van Montagu M, Caplanet A (1990) Characterization of a rice gene showing organ-specific expression in response to salt stress and drought. Plant Cell 2:19-27. https://doi.org/10.1105/tpc.2.1.19

de Azevedo Moreira R, Ainouz IL (1981) Lectins from seeds of jack fruit (*Artocarpus integrifolia* L.): isolation and purification of two isolectins from the albumin fraction. Biol Plant 23:186. https://doi.org/10.1007/BF02894883

Dubois M (1956) Colorimetric method for determination of sugars and related substances. Anal Chem 28:350–356. https://doi.org/10.1021/ac60111a017

Dubouzet JG, Sakuma Y, Ito Y, Kasuga M, Dubouzet EG, Miura S, Seki M, Shinozaki K, Yamaguchi☐Shinozakiet K (2003) OsDREB genes in rice, *Oryza sativa* L., encode transcription activators that function in drought☐, high☐salt☐ and cold☐responsive gene expression. Plant J 33:751-763. https://doi.org/10.1046/j.1365-313x.2003.01661.x

Filho GAS, Ferreira BS, Dias JM, Queiroz KS, Branco AT, Bressan-Smith RE, Oliveira JG, Garcia AB (2003) Accumulation of SALT protein in rice plants as a response to environmental stresses. Plant Sci 164:623-628. https://doi.org/10.1016/S0168-9452(03)00014-1

Fiser A, Sali A (2003) ModLoop: automated modeling of loops in protein structures. *Bioinformatics* 19:2500–2501. https://doi.org/10.1093/bioinformatics/btg362

Ghanta S, Datta R, Bhattacharyya D, Sinha R, Kumar D, Hazra S, Mazumdar AB, Chattopadhyay S (2013) Multistep involvement of glutathione with salicylic acid and ethylene to combat environmental stress. J Plant Physiol 171:940-950. https://doi.org/10.1016/j.jplph.2014.03.002

Grieve CM, Grattan SR (1983) Rapid assay for the determination of water soluble quaternary ammonium compounds. Plant Soil 70:303-307. https://doi.org/10.1007/BF02374789

He X, Li L, Xu H, Xi J, Cao X, Xu H, Rong S, Dong Y, Wang C, Chen R, Xu J, Gao X, Xu Z (2016) *Os*JRL, a rice jacalin-related mannose-binding lectin gene, enhances *Escherichia coli* viability under high-salinity stress and improves salinity tolerance of rice. Plant Biol (Stuttg) 19:257–267. https://doi.org/10.1111/plb.12514

Hu H, Xiong L (2014) Genetic engineering and breeding of drought resistant crops. Annu Rev Plant Biol 65:715–741. https://doi.org/10.1146/annurev-arplant-050213-040000

Jiang SY, Ma Z, Ramachandran S (2010) Evolutionary history and stress regulation of the lectin superfamily in higher plants. BMC Evol Biol 10:79. https://doi.org/10.1186/1471-2148-10-79

Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiyama C, Dohmae N, Takio K (2006) Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor. Proc Natl Acad Sci U S A 103:11086–11091. https://doi.org/10.1073/pnas.0508882103

Kelley L, Mezulis S, Yates CM, Wass MN, Sternberg MJE (2015) The Phyre2 web portal for protein modeling, prediction and analysis. Nat Protoc 10:845–858. https://doi.org/10.1038/nprot.2015.053

Kikuchi K, Ueguchi-Tanaka M, Yoshida KT, Nagato Y, Matsusoka M, Hirano HY (2000) Molecular analysis of the NAC gene family in rice. Molecular Genetics and Genomics 262:1047–51. https://doi.org/10.1007/pl00008647

Krieger E, Joo K, Lee J, Lee J, Raman S, Thompson J, Tyka M, Baker D, Karplus K (2009) Improving physical realism, stereochemistry, and side-chain accuracy in homology modeling: Four approaches that performed well in CASP8. Proteins 77 Suppl 9:114-1122. https://doi.org/10.1002/prot.22570

Krissinel E, Henrick K (2007) Inference of macromolecular assemblies from crystalline state. J Mol Biol 372:774–797. https://doi.org/10.1016/j.jmb.2007.05.022

Lannoo N, Van Damme EJMV (2014) Lectin domains at the frontiers of plant defense. Front Plant Sci 5:397. https://dx.doi.org/10.3389%2Ffpls.2014.00397

Laskowski RA, Rullmannn JA, MacArthur MW, Kaptein R, Thornton JM (1996) AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. J Biomol NMR 8:477-486. https://doi.org/10.1007/bf00228148

Lata C, Prasad M (2011) Role of DREBs in regulation of abiotic stress responses in plants. J Exp Bot 62:4731-4748. https://doi.org/10.1093/jxb/err210

Lichtenthaler HK (1987) Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. Method Enzymol 148:350–382. https://doi.org/10.1016/0076-6879(87)48036-1

Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain, separate two cellular signal transduction pathways in drought- and low temperature-responsive gene expression, respectively, in *Arabidopsis*. Plant Cell 10:1391–1406. https://doi.org/10.1105/tpc.10.8.1391

Mashiach E, Schneidman-Duhovny D, Andrusier N, Nussinov R, Wolfson HJ (2008) FireDock: a web server for fast interaction refinement in molecular docking. Nucleic Acids Res 36:W229-32. https://doi.org/10.1093/nar/gkn186

Mizoi J, Kanazawa N, Kidokoro S, Takahashi F, Qin F, Morimoto K, Shinozaki K, Yamaguchi-Shinozaki K (2019) Heat-induced inhibition of phosphorylation of the stress-

protective transcription factor DREB2A promotes thermotolerance of *Arabidopsis thaliana*. J Biol Chem 294:902-917. https://doi.org/10.1074/jbc.RA118.002662

Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. Physiol Plant 15:473-497. https://doi.org/10.1111/j.1399-3054.1962.tb08052.x

Nakashima K, Ito Y, Yamaguchi-Shinozaki K (2009) Transcriptional regulatory networks in response to abiotic stresses in *Arabidopsis* and grasses. Plant Physiol 149:88–95. https://doi.org/10.1104/pp.108.129791

Nuruzzaman M, Manimekalai R, Sharoni AM, Satoh K, Kondoh H, Ooka H, Kikuchi S (2010) Genome-wide analysis of NAC transcription factor family in rice. Gene 465:30–44. https://doi.org/10.1016/j.gene.2010.06.008

Ooka H, Satoh K, Doi K, Nagata T, Otomo Y, Murakami K, Matsubara K, Osato N, Kawai J, Carninci P (2003) Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*. DNA Res 10:239-47. https://doi.org/10.1093/dnares/10.6.239

Paul S, Gayen D, Datta SK, Datta K (2015) Dissecting root proteome of transgenic rice cultivars unravels metabolic alterations and accumulation of novel stress responsive proteins under drought stress. Plant Sci 234:133–143. https://doi.org/10.1016/j.plantsci.2015.02.006

Ren J, Gao X, Jin C, Zhu M, Wang X, Shaw A, Wen L, Yao X, Xue Y (2009) Systematic study of protein sumoylation: Development of a site-specific predictor of SUMOsp 2.0. Proteomics 9:3409-3412. https://doi.org/10.1002/pmic.200800646

Saad AS, Li X, Li HP, Huang T, Gao CS, Chen W, Zhao GY, Liao YC (2013) A rice stress-responsive NAC gene enhances tolerance of transgenic wheat to drought and salt stresses. Plant Sci 203-204:33-40. https://doi.org/10.1016/j.plantsci.2012.12.016

Saha P, Dasgupta I, Das S (2006) A novel approach for developing resistance in rice against phloem limited viruses by antagonizing the phloem feeding hemipteran vectors. Plant Mol Biol 62:735-752. https://doi.org/10.1007/s11103-006-9054-6

Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K (2002) DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. Biochem Biophys Res Commun 3:998-1009. https://doi.org/10.1006/bbrc.2001.6299

Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ (2005) PatchDock and SymmDock: servers for rigid and symmetric docking. Nucleic Acids Res 33:W363-367. https://doi.org/10.1093/nar/gki481

Shakirova F, Bezrukova M, Shayakhmetov I (1996) Effect of temperature shock on the dynamics of abscisic acid and wheat germ agglutinin accumulation in wheat cell culture. Plant Growth Regul 19:85-87. https://doi.org/10.1007/BF00024406

Shinozaki K, Yamaguchi-Shinozaki K (2000) Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. Curr Opin Plant Biol 3:217-223.

Song M, Xu W, Xiang Y, Jia H, Zhang L, Ma Z (2014) Association of jacalin-related lectins with wheat response to stresses revealed by transcriptional profiling. Plant Mol Biol 84:95-110. https://doi.org/10.1007/s11103-013-0121-5

Song Y, DiMaio F, Wang RYR, Kim D, Miles C, Brunette TJ, Thompson J, Baker D (2013) High resolution comparative modeling with RosettaCM. Structure 21:1735-1742. https://doi.org/10.1016/j.str.2013.08.005

Spadoro-Tank JP, Etzler EM (1988) Heath shock enhances the synthesis of a lectin-related protein in *Dolichos biflorus* cell suspension cultures. Plant Physiol 88:1131-1135. https://doi.org/10.1104/pp.88.4.1131

Tinjuangjun P, Loc N, Gatehouse A, Gatehouse J, Christou P (2000) Enhanced insect resistance in Thai rice varieties generated by particle bombardment. Mol Breed 6:391-399. https://doi.org/10.1023/A:1009633703157

Tovchigrechko A, Vakser IA (2006) GRAMM-X public web server for protein-protein docking. Nucleic Acids Res 34:W310-W314. https://doi.org/10.1093/nar/gkl206

Van Damme EJM, Peumans WJ, Barre A, Rouge P (1998) Plant lectins: a composite of several distinct families of structurally and evolutionary related proteins with diverse biological roles. Crit Rev Plant Sci 17:575-692. https://doi.org/10.1080/07352689891304276

Van Damme EJMV, Barre A, Rouge P, Peumans WJ (2004) Cytoplasmic/nuclear plant lectins: a new story. Trends Plant Sci 9:484-489. https://doi.org/10.1016/j.tplants.2004.08.003

Waadt R, Schmidt LK, Lohse M, Hashimoto K, Bock R, Kudla J (2008) Multicolor bimolecular fluorescence complementation reveals simultaneous formation of alternative CBL/CIPK complexes in planta. Plant J 56:505-516. https://doi.org/10.1111/j.1365-313X.2008.03612.x

Wang F, Liu Y, Shi Y, Han D, Wu Y, Ye W, Yang H, Li G, Cui F, Wan S, Lai J, Yang C (2020) SUMOylation stabilizes the transcription factor DREB2A to improve plant thermotolerance. Plant Physiol (In press) https://doi.org/10.1104/pp.20.00080

Weidenbach D, Esch L, Möller C, Hensel G, Kumlehn J, Höfle C, Hückelhoven R, Schaffrath U (2016) Polarized defense against fungal pathogens is mediated by the Jacalin-related lectin domain of modular Poaceae-specific proteins. Mol Plant 9:514–527. https://doi.org/10.1016/j.molp.2015.12.009

Woodrow P, Ciarmiello LF, Annunziata MG, Pacifico S, Iannuzzi F, Mirto A, D'Amelia L, Dell'Aversana E, Piccolella S, Fuggi A, Carillo P (2016) Durum wheat seedling responses to simultaneous high light and salinity involve a fine reconfiguration of amino acids and carbohydrate metabolism. Physiol Plant 159:290-312. https://doi.org/10.1111/ppl.12513

Yamaguchi-Shinozaki K, Shinozaki K (2006) Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. Annu Rev Plant Biol 57:781–803. https://doi.org/10.1146/annurev.arplant.57.032905.105444

Yamaji Y, Maejima K, Ozeki J, Komatsu K, Shiraishi T, Okano Y, Himeno M, Sugawara K, Neriya Y, Minato N, Miura C, Hashimoto M, Namba S (2012) Lectin-mediated resistance impairs plant virus infection at the cellular level. Plant Cell 24:778-793. https://doi.org/10.1105/tpc.111.093658

Yang ZP, Li HL, Guo D, Tang X, Peng SQ (2014) Identification and characterization of the 14-3-3 gene family in *Hevea brasiliensis*. Plant Physiol Biochem 80:121-127. https://doi.org/10.1016/j.plaphy.2014.03.034

Yi-juan H, Zhong ZH, Song LL, Olsson S, Wang ZH, Lu GD (2018) Evolutionary analysis of plant jacalin-related lectins (JRLs) family and expression of rice JRLs in response to *Magnaporthe oryzae*. J Integr Agric 17:1252-1266. https://doi.org/10.1016/S2095-3119(17)61809-4

Zhang W, Peumans WJ, Barre A, Astoul CH, Rovira P, Rougé P, Proost P, Truffa-Bachi P, Jalali AA, Van Damme EJM (2000) Isolation and characterization of a jacalin-related mannose-binding lectin from salt-stressed rice (*Oryza sativa*) plants. Planta 210:970-978. https://doi.org/10.1007/s004250050705

Zhao Q, Xie Y, Zheng Y, Jiang S, Liu W, Mu W, Zhao Y, Xue Y, Ren J (2014) GPS-SUMO: a tool for the prediction of sumoylation sites and SUMO-interaction motifs. Nucleic Acids Res 42:W325-W330. https://dx.doi.org/10.1093/nar/gku383

FIGURE LEGENDS

Fig. 1. Drought stress analysis of two *indica* rice cultivars, Vandana and MTU1010. 100 days old plants were subjected to drought stress for 7 days and different morphological parameters were analyzed. (A) Morphology of Vandana and MTU1010 in response to drought stress, (B) Plant height, (C) Shoot dry weight, (D) Percentage of rolled leaves, (E) Percentage of brown leaves, and (F) Root dry weight. Results were represented as mean±SEM (n=3). Statistical difference between the cultivars under control and drought stress was denoted by asterisks at $p<0.1$ (**), $p<0.01$ (***) and $p<0.001$ (****).

Fig. 2. Biochemical and transcript analysis of Vandana and MTU1010 cultivars in response to drought stress. Samples were collected from control and stressed plants and several biochemical parameters were analyzed. (A) Chlorophyll content, (B) Glycine betaine content, (C) Soluble sugar content, and (D) Proline content. Samples were also used for qRT-PCR analysis to study the relative transcript abundance for (E) *OsOsmotin* and (F) *OsSalT* genes. Results were represented as mean±SEM (n=3). Statistical difference between the cultivars under control and drought stress was denoted by asterisks at $p<0.05$ (*), $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****).

Fig. 3. Response of transgenic tobacco lines ectopically expressing *OsSalT* gene under drought stress condition. The WT, VC and three independent transgenic lines (OX1, OX2 and OX3) were subjected to drought stress for 5 days and analyzed. (A) Morphology, (B) Shoot dry weight, (C) Percentage of yellow leaves and (D) Percentage of wilted leaves. The relative transcript abundance of OsSalT gene (E) was also analyzed. Results were represented as mean±SEM (n=3). Statistical difference between the lines under control and drought stress was denoted by asterisks at $p<0.01$ (**), $p<0.001$ (***) and $p<0.0001$ (****).

Fig. 4. Identification of interacting protein partners of *Os*SalT protein under drought stress. (A) Yeast two-hybrid analysis identified the interaction of *Os*SalT with *Os*DREB2a and *Os*NAC1 proteins. The interaction of p53 protein with T-antigen was used as a positive control. (B) BiFC analysis confirmed the interaction of *Os*SalT protein with *Os*DREB2A in the nucleus and cytoplasm and with *Os*NAC1 in the nucleus only. Venus fluorescence, bright field, and merged images were represented for each set of constructs. Arrows indicate interaction in the nucleus.

Fig. 5. *In silico* analysis for *Os*SalT-*Os*DREB2A interaction. Protein structures for *Os*SalT and *Os*DREB2A were generated through homology modelling. The top views (A,C) and the side views (B,D) of *Os*SalT and *Os*DREB2A structures have been shown. The structures were used for molecular docking analysis. (E) Surface view, (F) Cartoon view, (G) Interphase residues. Black dashed lines indicate polar contacts among the interphase residues. Yellow residues indicate the DNA-binding domain of *Os*DREB2A.

Fig. 6. *In silico* analysis for *Os*SalT-*Os*NAC1 interaction. Protein structure for *Os*NAC1 was generated through homology modelling. The top view (A) and the side view (B) of *Os*NAC1structure have been shown. The structure was used for molecular docking analysis with *Os*SalT protein. (C) Cartoon view, (D) Surface view, (E) Interphase residues. Black dashed lines indicate polar contacts among the interphase residues. Yellow residues indicate the DNA-binding domain of *Os*NAC1.

Fig. 7. Expression of *DREB2A* and *NAC1* responsive genes under drought stress in the WT and transgenic tobacco (OX) plants. The relative transcript abundance of *NtDehydrin, NtLEA, NtSPS, NtPI3K, Ntrd22 and NtADH1* genes were analyzed under drought stress. Results were represented as mean ± SEM (n=3). Statistical difference between the lines under control and drought stress was denoted by asterisks at $p<0.0001$ (****).

Fig. 8. Model for *Os*SalT-mediated regulation of drought stress. Drought stress induces the expression of *Os*SalT in cells. In the mean tine, *Os*NAC1 expression is also induced and its accumulation increases. On the other hand, *Os*DREB2A protein is presumably stabilized by SUMOylation. The *Os*SalT now binds with the stabilized *Os*DREB2A and *Os*NAC1 proteins thus activating them. This results in the transcriptional activation of downstream drought-responsive genes leading to stress tolerance.

A

Bait	Prey	DDO				QDO/X/A			
		10^1	10^{-1}	10^{-2}	10^{-3}	10^1	10^{-1}	10^{-2}	10^{-3}
*Os*SalT	*Os*DREB2A								
*Os*SalT	*Os*NAC1								
p53	T-antigen								

B

	Venus	Brightfield	Merged
35S::OsSalT-cVenus + *35S::OsDREB2A-nVenus*			
35S::OsSalT-cVenus + *35S::OsNAC1-nVenus*			
35S::OsSalT-cVenus + *35S::pVYNE-nVenus*			

Scale: 50 um

A	B	C	D
Side view	Top view	Side view	Top view

Side view

Top view