Effects of genomic and functional diversity on stand-level productivity and performance of

non-native Arabidopsis

Kathryn G. Turner[1,2], Claire M. Lorts[1], Asnake T. Haile[1,3], Jesse R. Lasky[1]

[1]Department of Biology
Pennsylvania State University
USA

[2]Department of Biological Sciences
Idaho State University
USA

[3]Department of Plant Biology and Biodiversity Management
Addis Ababa University
Ethiopia

Correspondence: lasky@psu.edu

Abstract

Biodiversity can affect the properties of groups of organisms, such as ecosystem function and the

persistence of colonizing populations. Genomic data offer a newly available window to diversity,

complementary to other measures like taxonomic or phenotypic diversity. We tested whether

native genetic diversity in field experimental stands of *Arabidopsis thaliana* affected their

aboveground biomass and fecundity in their colonized range. We constructed some stands of

genotypes that we *a priori* predicted would differ in performance or show overyielding. We

found no relationship between genetic diversity and stand total biomass. However, increasing

stand genetic diversity increased fecundity in high resource conditions. Polyculture (multiple

genotype) stands consistently yielded less biomass than expected based on the yields of

component genotypes in monoculture. This under-yielding was strongest in stands with late-

flowering and high biomass genotypes, potentially due to interference competition by these

genotypes. Using a new implementation of association mapping, we identified genetic loci

whose diversity was associated with stand-level yield, revealing a major flowering time locus

associated with under-yielding of polycultures. Our field experiment supports community

ecology studies that find a range of diversity-function relationships. Nevertheless, our results

suggest diversity in colonizing propagule pools can enhance population fitness. Furthermore,

interference competition among genotypes differing in flowering time might limit the advantages

of polyculture.

Keywords: diversity-productivity relationships, evolution, phenology, invasion

INTRODUCTION

The characteristics of groups of organisms, such as their genetic diversity, can affect their persistence and collective functions. The emergent properties of groups of organisms are major topics in agriculture, ecology, and evolutionary biology [1–3]. The mechanisms underlying the benefits of diversity for group-level performance can be divided into at least two major categories [4]. First, greater diversity can reduce the negative effects associated with high density populations or communities. This can occur via diversity in resource use that reduces competition, diversity in susceptibility to enemies that reduces "apparent" competition, or facilitation among diverse individuals. Second, diversity can increase group-level performance via a sampling effect: greater diversity increases the chance that high-performing individuals will be present. By contrast, diversity can also reduce group-level performance. This can occur when more diverse groups tend to have lower performing individuals (e.g. when diversity is driven by immigration of maladapted individuals) [5,6] or when diversity results in interference competition among different types [7]. While diversity effects on ecosystem functions (EF) can act at both population and community levels, they have been mostly studied at the community level [8,9]. However, *intraspecific* diversity can have major impacts on EF, potentially surpassing interspecific effects when a few species dominate biomass [10].

Community ecologists have often hypothesized that more diversity increases ecosystem functions [11,12] while invasion biologists have hypothesized that greater diversity increases the growth rates of colonizing populations [13–15]. These two research programs have been largely independent and largely un-synthesized, but they share much in common (see also related topics in kin and group selection [16]). Both address hypotheses based on the same underlying biological processes: 1) sampling effects, whereby greater diversity increases the chance a group will contain high functioning/performing individuals, 2) insurance effects (a type of sampling effect), whereby greater diversity increases the chance that a group will contain an individual that is high functioning under some potential new environment, or 3) complementarity effects, whereby diversity relieves negative density effects. Invasion biology has long investigated how the number of individuals in a founding group affects colonization success [17,18]. Evidence from many studies suggests that increased genetic diversity, rather than increased numbers of individuals *per se*, causes improved establishment [13], growth rate [13–15], biomass production

[2,19,20], and persistence of colonizing populations, and decreases vulnerability to

environmental change (reviewed in [21]). The effects of diversity on group-level performance

(e.g. average fitness) and EF are likely to be conditional on the environment. For example,

positive facilitative interactions may dominate in extremely harsh environments, leading to

stronger positive diversity effects [22,23], or novel environments might be more limited by

diversity due to a rarity of high-functioning individuals. Empirical studies of genetic diversity in

an invasion context have found it favorable under some conditions but not others [2,24] and

theoretical predictions also conflict [20,25].

Diversity effects on the performance of a group of organisms arise fundamentally due to

the effects of phenotypic diversity [26]. However, the phenotypes that affect group-level

performance are very often unknown. Instead, genetic, taxonomic, or phylogenetic diversity can

be useful proxies. Past studies of intraspecific diversity and EF have used either genotypes of

unknown relatedness [19,20,27], or have used a small number of genetic markers having unclear

connections to functional variation [2,28,29]. By contrast, large numbers of markers across the

genome allow precise estimation of genetic similarity between individuals, and it is now feasible

to measure whole genome diversity in many non-model systems. Furthermore, genetic diversity

effects may be most strongly associated with the specific loci that underly complex phenotypes

associated with resource economics. Genetic variation in complex traits corresponding to life

history and resource use likely involves changes in many loci [30,31], and simple genome-wide

additive models can often predict variation in such traits [32–34]. Additionally, genetic mapping

may be used to identify specific genes and mutations controlling phenotypes whose diversity

affects EF [23,35].

Here we investigate how intraspecific diversity influences productivity and fecundity for

the model plant *Arabidopsis thaliana* across environmental conditions in a field experiment

outside its native range. Arabidopsis is useful for our questions: it is a model for ecological

genomics [36], diverse germplasm with resquenced genomes is available [37], and it exhibits

variation in resource strategies and life history [38,39] suggesting that diversity may impact EF.

Our study builds upon previous work in Arabidopsis [19,23,35] by using ecologically realistic

and relevant field experimental conditions, including a low nutrient well-drained soil treatment,

and by studying stands with up to 20 genotypes. Because Arabidopsis is largely selfing and we

studied a single generation, we set aside the effects of diversity on population performance due

to sexual reproduction, such as heterosis.

Experimental studies of biodiversity effects typically use randomly assembled groups of

genotypes or species, despite existing prior knowledge of potential drivers of compositional and

diversity effects on EF. Here, in addition to randomly assembled groups representing a range of

genotypic diversity, we use knowledge of our system to develop *a priori* selected groups of

genotypes that we predict will exhibit particularly strong differences in EF, groups benefitting

from potential niche complementarity, and groups of genotypes we predict are particularly well-

suited for the experimental site. We use information indicative of local adaptation to the climate

at our site as well as information on genetic diversity in an ecologically important plant trait that

may mediate competition (flowering time) to design collections of genotypes predicted to differ

in performance or predicted to perform well.

METHODS

Plant material

We studied 60 diverse Arabidopsis genotypes from across the native Eurasian range of

Arabidopsis, mainly from the Mediterranean, central Asia, the UK and central Europe (Figure

S1). All genotypes are representatives of natural inbred lines that had their genomes resequenced

[37]. These genotypes were nearly evenly distributed among 7 genetic clusters or admixed

genotypes identified by [37] (though we did not have any Swedish genotypes). We selected these

genotypes based on diversity and seed availability from a common seed production environment

used in previous experiments (Extended Methods S1).

Field experiment

To assess the impact of different levels of genetic diversity in experimentally assembled non-

native populations of Arabidopsis, we measured performance of Arabidopsis stands in a field

common garden at a site (40.8106º N, 77.8472º W) in the non-native range near other established

non-native populations (within ~10 km) but with none at this site (within ~1km). Thus our

experimental stands were likely not influenced by density-associated processes driven by wild

individuals nearby. To assess the environmental context dependency of diversity effects, we
replicated all stand genotype compositions in two soil types: a putatively benign high-resource
potting mixture, and a lower resource sandy soil (described below). Each of the 60 genotypes
were grown in monoculture in three replicate stands for each of the two soil environments, for a
total of 360 monoculture stands (Figure S2).

We began by constructing polycultures based on *a priori* hypotheses for factors likely to
affect productivity. We used five different criteria to generate these 'prediction stands': (1)
genomic diversity among accessions, (2) climate similarity of collection location to our field site,
combined with genomic diversity, (3) early flowering genotypes, (4) late flowering genotypes,
(5) climate similarity of collection location to our study site, combined with flowering time
diversity (see details on the algorithms we used to assemble these stands in Methods S1). Two
stand compositions were chosen for each of the five prediction scenarios at each diversity level
(2, 10, 20), for a total of ten prediction stand compositions for each diversity level. Each of the
prediction stands were planted into both of the two soil environments, for a total of 60 prediction
stands.

After genotypes were assigned to prediction stands, we also generated random
polyculture compositions. In assembling these stands we sought to limit similarity of stands
within a treatment and within stands having the same number of genotypes [19,40]. We
generated polycultures of either 2, 10, or 20 different randomly selected genotypes, with ten
stand replicates per diversity level, for a total of 30 different randomly generated polycultures.
All genotypes were included within the 10 stand replicates, randomly assigned to replicates such
that each replicate had a different assortment of genotypes but had the same diversity level. Each
of these polycultures were planted into both soil environment treatments with 10 replicates of
diversity level each for a total of 60 randomly assigned polyculture stands (see methods on
assigning these stand compositions in Methods S1).

Seeds were initially sown into 20 positions within cow manure pots, with each position
thinned to a single plant, targeting a 20-plant stand (regardless of diversity level). High resource
pots were filled with potting mix and low resource pots with a high proportion of sand. Stands
were germinated in growth chambers and transplanted into the field site on May 24, 2018
(Methods S1).

We measured specific leaf area (SLA) on a representative, healthy leaf from a single

plant in each monoculture stand, and flowering date on all stands. We harvested each individual

plant once it reached maturity. We recorded the total silique (fruit) number and the length of one

representative middle silique. All remaining plants were harvested on July 18-20, as summer

heat stress set in and reproduction largely ceased. We measured aboveground dry biomass on all

plants. Canopy area was similarly measured for each stand just prior to the first harvested plant

using Easy Leaf Area and a red 4 cm^2 square reference [41].

Measures of stand composition and diversity

To estimate genomic diversity of stands, we used published SNPs from resequencing data [37].

We calculated the genetic distance matrix using the R package SNPRelate [42] and then

calculated the mean genetic distance among plants sown in a stand (including zero distance for

individuals of the same genotype). We also estimated phenotypic variation, based on our new

SLA data and published flowering time data [37]. For SLA, we took breeding values of

log(SLA) measured on plants in our monocultures and calculated mean for individual plants in

all stands. Flowering time composition and diversity were respectively estimated using

individuals' breeding value averages of 10ºC and 16ºC published flowering time growth chamber

experiments [37].

Statistical analyses

To gain general insight into the sources of variation in phenotypes and performance, we tested

for treatment effects, estimated broad-sense heritability, and estimated the genomic signature of

performance in monocultures. To test for treatment effects on trait plasticity in monocultures, we

implemented linear mixed models using the R package 'VCA' [43] and included genotype and

block as random effects and soil treatments and plot x-y coordinate as fixed effects. Plastic trait

responses were characterized with this soil fixed effect. We estimated broad sense heritability of

stand-level traits and performance for monocultures using linear models in R where genotypes

were fixed effects, taking R^2 of these as broad sense heritability. We tested for a genomic

signature in fecundity and biomass production using genomic prediction models, where traits are

a function of genotype-specific random effects that are correlated according to the genome-wide

similarity matrix, using the R package 'rrBLUP' [44]. We used out-of-sample prediction and ten-

fold cross validation to estimate the ability of genome-wide similarity to predict performance.

To address our primary research questions, we then tested whether multiple measures of

diversity affected stand level performance, assessed by fecundity, aboveground biomass, and leaf

canopy area. In our analyses, we controlled for variation among stands in the number of plants

surviving transplant or germinating late, with a minimum of 11 plants required for inclusion in

analysis. We used two different counts of plants surviving transplant, on May 29 and June 12,

and took the maximum, but we found results were similar when using the count on either date.

We used linear regression models to test diversity effects on stand-level yield while

accounting for additional sources of error. The covariates were each tested in models including

an interaction with the resource-limitation treatment

$$y_i = \beta_0 + \beta_1 x_i + \beta_{trt} trt_i + \beta_2 x_i trt_i + \varepsilon_i$$

where y_i is the performance measure (biomass or fecundity) of stand i and trt_i is a dummy

variable for resource treatment, and x_i is a covariate of stand i's composition or diversity.

Covariates x tested in separate models included diversity treatment level (1, 2, 10, or 20

genotypes), genomic similarity (mean genetic distance among plants in a stand), stand-level

average of mean flowering time for each genotype, a measure of the stand-mean climate distance

from the field site, and stand-mean SLA breeding value. We similarly tested for stand diversity

and composition effects on fecundity [mean estimated total silique (fruit) length per fruiting

plant in stand].

We calculated the difference between the observed polyculture biomass yields and that

expected based on yields of monocultures and their proportional representation in the

polyculture. This difference is a commonly studied metric in community diversity-ecosystem

function studies [4], and we refer to this quantity as the deviation from monoculture expectation

(DME, this is referred to as the "net diversity effect" by [4]). We tested how the measures of

diversity and composition described above were related to DME, using linear models with a

parallel structure to those used for aboveground biomass and fecundity (described above).

To determine whether our prediction stands were successful at explaining variation in

fecundity and biomass production, we compared different stand types using ANOVA. Different

prediction criteria were each coded as a factor level as were randomly assembled stands. Two-

way ANOVA showed no significant treatment-stand type interaction, thus we focused on stand

type effects in a two-way ANOVA without interactions. We compared each pair of stand types

with post-hoc Tukey's tests with FDR control. Because only two stands predicted to be later

flowering produced fruit, we removed this plot type from our analysis of fecundity in prediction

plots.

To map loci across the Arabidopsis genome where allele frequency and variance were

related to stand biomass and DME, we used an approach akin to genome wide association

mapping. Specifically, we tested how both stand-level per plant biomass and DME (as response

variables) were associated with stand covariates of allele frequency and variance (the latter a

measure of diversity for a biallelic locus), for each SNP. For a biallelic SNP, variance = pq

where p and q are the respective frequencies of the two alleles. We used SNPs from published

whole genome resequencing [37]. We excluded SNPs segregating at <0.05 minor allele

frequency in our panel of 60 genotypes, leading to a total of ~1.9M SNPs. For each combination

of resource treatment (high or low) and performance measure (total yield or DME) we tested for

association with SNP allele frequency or variance. Because background variation across the

genome can contribute to phenotypic variation, we controlled for genome-wide similarity among

stands. Specifically, we calculated genome-wide similarity among stands in either allele

frequency or variance at all SNPs [45]. This builds off the traditional mixed-model approach in

genome-wide associations for phenotypes. We used a linear mixed model that included a random

effect that was correlated among stands according to the genome-wide similarity matrix, using

the EMMA software [45]. We also focused these association scans on SNPs within 3kb of 26 *a*

priori candidate genes for flowering time [46], based on the hypothesis that genes controlling

this trait might influence stand biomass and DME.

RESULTS

Plastic changes in traits and performance, heritability in monoculture

Comparing phenotypes for monoculture stands, SLA did not change significantly across soil

treatments, nor did flowering dates (linear mixed-effects models with genotype random effects,

Table S1). However, both average plant fecundity and biomass per plant in monocultures were

significantly greater in the high resource treatment (Table S1, Figure 1). Date of flowering was

highly heritable (~0.8 broad-sense, depending on metric and treatment), while log(SLA),

fecundity, and biomass had moderate heritability (~0.4-0.5 depending on trait and treatment,

Table S2).

Genomic signature of performance variation

There was a strong effect of genome-wide similarity among genotypes on biomass production in

monoculture in the low resource treatment. Genomic predictions (out-of-sample) were correlated

with actual breeding values for per plant biomass in monoculture in high resource (r = 0.05) and

low resource (r = 0.38) treatments. For example, in the low resource treatment a group of closely

related ecotypes from the Caucasus region (Azerbaijan and Georgia) produced among the most

biomass, while clusters of related genotypes from central Europe or Spain had low productivity.

We found a similar pattern for mean per plant fecundity in monocultures (genomic prediction

accuracy: high resource r = -0.01, low resource r = 0.33).

Figure 1. Reaction norms for aboveground biomass (AGB, left) and neighbor-joining trees
(based on pairwise genetic distance), where higher yielding genotypes are colored in green.
Three letter ISO country codes indicate the origin of each genotype. Two genotypes with poor
germination are not shown.

How did composition and diversity of stands impact stand fecundity and biomass production?

Among all stands (including monocultures and polycultures), the low resource soil treatment

significantly reduced stand fecundity, canopy coverage, and biomass production, as tested in

separate linear models of each response variable (Figure 2, Tables S3-5). We found that a greater

number of genotypes in a stand significantly increased the per plant fecundity in the high

resource treatment (Table S3) but not the low resource treatment, while other measures of

composition and diversity were unrelated with fecundity. We found that higher average breeding

values for flowering time were associated with significantly greater stand biomass in both

treatments, while lower breeding values for SLA was associated with greater stand biomass only

in the high resource treatment (Table S4). None of the several diversity measures (i.e. the other

covariates) we tested were significantly associated with biomass production.

We also tested if "prediction stands" had distinct stand-level fecundity or biomass from

randomly assembled stands (Figure S5). We did not find a significant effect of stand type in a

two-way ANOVA for either plot level fecundity or biomass.

Figure 2. Stand-level biomass and fecundity versus composition and diversity metrics. Each
point represents a stand; red circles indicate stands in the high resource treatment, blue triangles
indicate the low resource treatment. Resource treatment had a significant effect on both biomass
and fecundity (Tables S3-4). Shaded areas represent standard error. (a) Later stand mean
breeding values for flowering time from published experiments [37] significantly increased stand
biomass at harvest. (b) log(SLA) breeding value had a significant negative effect on stand
biomass at harvest in the high resource treatment. (c) Experimental stand diversity level (1, 2, 10,
20 genotypes) significantly increased estimated fecundity (total fruit length) per flowering plant
in the high resource treatment.

How did polyculture productivity deviate from monoculture expectation?

We found that in both treatments, the yield of polyculture stands was not closely correlated with

the expectation based on the yield of their component genotypes in monoculture (high resource, r

= 0.051, p = 0.7138; low resource, r = 0.007, p = 0.9585). Specifically, polycultures tended to

yield less than expected based on the yield of their component genotypes in monoculture. The

deviation from monoculture expectation (DME) was significantly less than zero in the high

resource stands (median = -61.01 mg, wilcoxon test p = 0.0025) and in the low resource stands

(median = -57.13 mg, wilcoxon test p = 0.0158). These results were unchanged (both p < 0.02)

when we generated randomly assembled polycultures to match the number of plants surviving

transplant using genotype-specific transplant survival in monoculture.

To evaluate potential causes of negative DME, we tested correlations between DME and

explanatory factors. One potential cause of negative DME was competitive inhibition of many

smaller plants by late flowering plants whose rosettes grew to overtop the earlier flowering

smaller plants. We found that in the high resource treatment, there was more negative DME in

stands with greater average breeding values for flowering time (mean flowering time; linear

model, flowering time in high resource slope = -0.0064 g/day, p = 0.0167, in low resource slope

= -0.0012 g/day, p = 0.6532) or stands having plants with greater flowering time (90th percentile

of flowering time; high resource slope = -0.0051 g/day, p = 0.0170, in low resource slope = -

0.0006 g/day, p = 0.7883, Table S6). We also found (perhaps unsurprisingly) that biomass in

monoculture of component genotypes was negatively associated with DME in polyculture:

stands with greater median or 90th percentile of genotypes' biomass in monoculture showed

significantly bigger reductions in polyculture yield in both high and low resource treatments

(Table S6). We tested whether DME might be due to genomic similarity, number of genotypes,

or climate of origin of genotypes. We found no strong effects on DME of mean kinship, total tree

diversity, number of genotypes, mean SLA, or mean distance to climate of origin along the first

five PCs (Table S6).

We also tested if "prediction stands" had distinct deviations from monoculture

expectation (DME). We found that there were significant differences in DME among prediction

stand criteria (two-way ANOVA with treatment and stand prediction criteria, including randomly

assembled stands: treatment p = 0.8675, prediction p = 0.0433, only including prediction stands:

treatment p = 0.8258, prediction p = 0.0328). Consistent with the regression of DME on

flowering time on all plots, the differences among our prediction plots were largely driven by

differences between the predicted early (positive diversity effect) and late flowering (negative

diversity effects) stands (Tukey HSD comparison of means, including randomly assembled

stands, only significant difference after multiple comparisons correction is: early flowering - late

flowering predicted stands DME = 0.27 g, FDR adjusted p = 0.0292).

Figure 3. Polyculture deviations from monoculture expectation (DME) in comparison with measures of stand composition (top panels) or stand assembly criteria (bottom panels). (Top panels) Stands with large biomass genotypes, or late flowering genotypes (90th percentiles) also tend to show lower DME (under-yielding), even more so than when measured via mean monoculture biomass or flowering time (Table S5). (Bottom panels) Stands assembled via different criteria show different DME. In particular, stands with genotypes predicted to be more

342   early flowering (EF) had higher DME (over-yielding) compared to late flowering (LF) stands
343   (Tukey HSD, FDR adjusted p = 0.0292). Gdiv = stands assembled based on genomic diversity,
344   CliGdiv = stands assembled based on similar climate of origin to study site but genomic
345   diversity, CliFdiv = stands assembled based on similar climate of origin to study site but
346   flowering time diversity.
347

What genetic loci contribute to diversity effects?

We tested whether per plant biomass or DME in each treatment were associated with allele
frequency or diversity at particular SNPs across the genome. We found that overall, there were
few loci with dramatically stronger (outlier) associations with biomass and DME, but rather
patterns suggest that the genetic basis of variation in biomass and DME were polygenic.
Nevertheless, we found that, in concordance with the association of late-flowering genotypes
with under-yielding polycultures, the top SNP (chr. 2, 9584630 bp, TAIR10, genome-wide
significant at FDR = 0.1, nominal p = 8.1 x 10^{-7}) associated with DME in the high resource
treatment (both allele frequency and variance association scans) was closest to SHORT
VEGETATIVE PHASE (SVP, also known as AGL-22, AT2G22540). SVP is a MADS-box
transcription factor, one of the top loci associated with natural variation in Arabidopsis flowering
time in [37] (Figure 4). Indeed, the minor allele of this SNP that we identified here as having
more positive yields in polycultures compared to expected was also associated with earlier
flowering at 16°C among the 60 genotypes we used from [37] (t-test, t= -2.6687, p = 0.0279). As
the frequency of this early-flowering minor allele increased and the stand variance at this locus
increased, DME also became more positive (Figure 4). We found the 26 flowering time
candidate genes had a number of SNPs with nominally significant associations with biomass and
DME in both treatments. Notably, SPL4 (AT1G53160), HEN2 (AT2G06990), ULP1B
(AT4G00690), and ATH1 (AT4G32980) were significantly associated with DME in low
resource stands (SNP frequency and variance associations, with candidate SNP-wide FDR <
0.05, Table S7).
369

A. Var(allele freq.) associations with DME in high-resource

B. Top SNP for DME in high resource

Figure 4: Associations between DME in the high resource treatment with var(allele frequency) across the genome (A) and at the top associated SNP (B). (A) Linear mixed model genome-wide association with 1.9M SNPs allele frequency variance (var $= pq$) versus deviation from monoculture expectation (DME) in high resource environments. SNPs significant with FDR < 0.1 are shown in red. The top SNP is highlighted with a large red dot in panel A and is featured in panel B (this was also the top SNP for allele frequency associations with DME). Boxplot (panel B) shows variance in allele frequency within polycultures (x-axis) at this SNP (chromosome 2, 9584630 bp) associated with DME of biomass (panel B y-axis) in the high resource environment.

DISCUSSION

Diversity can affect the group-level properties of organisms, including ecosystem functions like

carbon cycling and population trajectories. Diversity effects have been often studied via

observation of natural, uncontrolled systems that vary in species or genetic diversity and testing

whether diversity covaries with group properties [29,47]. In particular, species and higher levels

of diversity have been demonstrated to have a range of effects on ecosystem function via

multiple mechanisms [8,48]. We built on this research to test how intraspecific variation in trait

and genomic composition and diversity affect the productivity and fecundity of an established

non-native species in the field. We found that the diversity of genotypes did not increase biomass

production, but rather polycultures showed lower biomass production than expected, suggesting

interference competition among genotypes. This was associated with low polyculture yield of

stands containing late-flowering genotypes that had high productivity in monoculture. We

identified genetic variation in a known flowering time locus associated with these effects. We

also found a weak but significant increase in fecundity with the number of genotypes in a

treatment, due to unclear mechanisms.

Diversity effects on average individual fecundity

Because of logistical and (in the case of invasive species) bioethical constraints, many studies of

the effect of diversity on population performance focus on the impact on biomass production and

may not assess fecundity directly (see [2,19,20]). While none of the diversity metrics we tested

had an effect on biomass production, we did find a small but significant effect of genotype

number on stand level fecundity in the high resource environment (Fig 2c). The mechanism

underlying this effect is unclear, given the lack of an effect on biomass. This may be due to

sampling effects, specifically where greater diversity increases the chance a group will contain

early flowering and more fecund genotypes. Alternately, interference competition may trigger

earlier reproduction in small (especially early flowering) genotypes to a greater degree than large

(especially late flowering) genotypes, leading to positive diversity-fecundity associations.

Diversity effects on stand-level biomass

We found no evidence that increased genetic diversity led to greater stand-level aboveground

biomass production of Arabidopsis in the field, highlighting the complexity of stand-level

properties in even this model plant. Our findings may be explained by a lack of such genetic

diversity effects in Arabidopsis populations in general, or by technical/logistical constraints that

made us miss such effects. First, diversity effects might not have been observed if positive

effects of diversity (complementarity of resource use or pathogen susceptibility) were opposed

by negative effects (interference competition) [7] or if diversity effects did not exist. Our finding

of reduced polyculture yield compared to expectation from monoculture (DME) suggests

interference competition as one potential explanation. Second, diversity effects might have been

overlooked if they occurred along dimensions that were not measured, such as seed survival or

nutrient use [26]. Diversity effects might also have been overlooked if they occur across

environmental gradients unobserved in our single year, two resource level, study [49,50].

By contrast, we found consistent under-yielding of polycultures of genotypes compared

to the expectations from their biomass in monoculture and relative representation in the

polycultures (deviation from monoculture expectation, DME, also known as "diversity effects")

[4,7]. Generally, negative DMEs are thought to arise from interference competition [7]. This

process could play out in our experiment as follows: when grown with genotypes having

different trait values, individuals exhibit reduced performance compared to in monoculture with

genetically identical individuals, and apparently without much benefit to the performance of the

higher performing genotypes. For example, [51] found that *Pseudomonas flourescens* genotypic

diversity lead to reduced ecosystem function (host plant protection) due to antagonistic

interference (mutual poisoning) among *Pseudomonas* genotypes. Similar patterns have been

reported in plants, where individuals have higher performance among similar genotypes and

reduced performance among different genotypes [24,27]. For example, [24] found reduced

variance in biomass among individuals and higher individual fecundity of *Plantago lanceolata*

when grown among related individuals, which he interpreted as being due to reduced

interference competition in high relatedness stands (though the term "interference competition"

was not used). This might occur if higher performing genotypes reduce access to resources of

lower performing genotypes to a greater degree than those high performing genotypes benefit by

being surrounded by fewer of their own type. Alternatively, higher performing genotypes in

monoculture might be more limited by lower performing genotypes, with little benefit to lower

performing genotypes. We found that polyculture plots containing later flowering and larger

biomass (in monoculture) genotypes were those that had the strongest negative DME, possibly

due to shading of different and early flowering smaller neighbors. Other authors have sometimes

interpreted these patterns as evidence of kin selection [52,53]. The interference competition

explanation for negative DME seems more likely than plants' ability to first recognize kin *per se*

and then reduce interference with related neighbors (cf. [53]).

We demonstrated how genomic data on individuals in experimental stands can be used to
identify genetic loci potentially affecting group-level performance, although we did not identify
effects of whole genome diversity on stand performance. Recent studies using Arabidopsis
grown in pots, combined with linkage [23] or association mapping [35] have demonstrated
complementary genomic approaches using a smaller number of genotypes and individuals per
stand (two genotypes and three-four individuals/pot). Controlled experiments like these and our
present study, combined with group-level performance and genetic data, could be applied to
nearly any system. It is not clear whether such approaches could detect genetic loci operating in
natural uncontrolled systems, where uncontrolled environmental variation plays an important but
often unmeasured role in driving spatial variation in group-level performance. Here, we
implemented a study of association of stand level performance with both allele frequency and
variance in stands. However, note that to distinguish between effects of variance (diversity *per*
se) versus frequency (composition) at a biallelic SNP, a given SNP would need to span a wide
range of allele frequencies among stands. This condition is not likely met for most SNPs given
they have low minor allele frequency, thus allele frequency and variance show a large degree of
covariation, limiting our ability to distinguish the effects of diversity versus allelic composition
at most loci.
466
Phenotypes and genotypes associated with stand-level performance
We found that stand phenotypic composition was associated with stand-level performance.
Specifically, monoculture productivity was associated with certain closely related genotypes
under the low-resource environment (Fig. 1), and biomass production was associated with stands
comprised of low SLA or late-flowering genotypes. Late flowering genotypes invest greater
proportions of resources and time in vegetative growth before transitioning to reproduction. For
this reason, late-flowering genotypes may be more productive in terms of biomass and similarly
they may often possess traits useful for biomass production crops [54]. Here, the late-flowering
genotypes we studied were associated with polycultures that had less biomass than expected
(based on monocultures of component genotypes) in the high resource environment. This
suggests that late-flowering genotypes may have been more likely to be involved in interference
competition when resources were abundant, perhaps because these larger individual plants
limited their competitors from resource capture (e.g. by shading) more than they benefited,

reducing the overall total biomass [24]. Evidence suggests that size asymmetries leading to

shading [55] as well as allelopathy against competitors or other trophic levels [56] can also lead

to such interference competition.

Previous studies on Arabidopsis

Previous studies in the laboratory have shown some evidence that certain combinations of

Arabidopsis genotypes may result in higher stand level performance. In greenhouse populations

of Arabidopsis, both additive and non-additive benefits of up to eight genotypes per stand (from

a pool of 23) were found to increase emergence, biomass, flowering duration, and fecundity [19].

[23] used an Arabidopsis biparental mapping population to identify a single locus causing

overyielding when grown in diversity in pots. Additionally, [35] used pot experiments on pairs of

genotypes in a greenhouse, combined with association mapping on an individual genotype's

association with over- versus under-yielding. Our study was conducted in the field where plants

were subject to a wide range of potential stressors. It is noteworthy that [57] found that natural

stands in Germany with a single genotype tended to have fewer individuals than stands with

multiple genotypes, suggesting potential positive diversity effects in nature.

A priori prediction of stand-level performance

Ecology has been criticized for not being a predictive science. Here we tested our knowledge of

the system with *a priori* sets of stands predicted to have divergent performance. We found that

our predictions were most closely associated with stand-level yield driven by flowering time

differences among genotypes. It is no accident that we chose flowering time as the one trait to

include in our prediction criteria. Flowering time has well-documented links to a major axis of

life history and physiological variation [39,58], and is so intensely studied for agronomic

interest. \We believe the use of phenotyped and genotyped individuals in such experiments are a

way forward for predictive approaches to group performance and diversity.

Conclusions

Biodiversity can affect the properties of groups of organisms and may be important to

understanding the performance and persistence of both natural and introduced populations and

communities. Here we did not identify strong effects of genetic diversity *per se* on the

performance of stands of non-native Arabidopsis populations but we found a pattern of higher

biomass yield when genetic composition was shifted to late-flowering stands. We also identified

that in polyculture, the presence of later-flowering genotypes led to lower-than expected yield,

possibly due to interference competition. Our research here on stand-level effects on

performance in the field for the model plant Arabidopsis suggest a way forward to understanding

the importance of such effects in nature.

Acknowledgments

Connor Campana, Timothy Gilpatrick, Sarah Lucas, and Ken Hull provided assistance with

fieldwork. Victoria DeLeo, Emily Bellis, Lua Lopez, and Jonathan Kizer provided assistance

with sowing seeds. Paras Patel and Crosley Williams assisted with sorting and sowing seeds and

preparing pots. Leland Shawn Burghard assisted with plant care. Megan Vahsen provided useful

discussions on experimental design. This work was supported by a Pennsylvania State University

Eberly Postdoctoral Fellowship and National Science Foundation grants and NSF Idaho EPSCoR

Program award OIA-1757324 to KGT.

References

1. Denison RF. 2012 *Darwinian Agriculture: How Understanding Evolution Can Improve Agriculture*. Princeton University Press.

2. Hughes AR, Stachowicz JJ. 2004 Genetic diversity enhances the resistance of a seagrass ecosystem to disturbance. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 8998–9002. (doi:10.1073/pnas.0402642101)

3. Loreau M. 2010 *From populations to ecosystems: theoretical foundations for a new ecological synthesis*. Princeton University Press. See http://books.google.com/books?id=qIxM7eNj9sUC.

4. Loreau M, Hector A. 2001 Partitioning selection and complementarity in biodiversity experiments. *Nature* **412**, 72–76. (doi:10.1038/35083573)

5. Lasky JR. 2019 Eco-evolutionary community turnover following environmental change. *Evol. Appl.* **12**, 1434–1448. (doi:10.1111/eva.12776)

6. Mouquet N, Loreau M. 2003 Community Patterns in Source-Sink Metacommunities. *Am. Nat.* **162**, 544–557.

7. Loreau M. 1998 Separating Sampling and Other Effects in Biodiversity Experiments. *Oikos* **82**, 600–602. (doi:10.2307/3546381)

8. Cadotte MW. 2013 Experimental evidence that evolutionarily diverse assemblages result in higher productivity. *Proc. Natl. Acad. Sci.* (doi:10.1073/pnas.1301685110)

9. Srivastava DS, Cadotte MW, MacDonald AAM, Marushia RG, Mirotchnick N. 2012 Phylogenetic diversity and the functioning of ecosystems. *Ecol. Lett.* **15**, 637–648. (doi:10.1111/j.1461-0248.2012.01795.x)

10. Reynolds LK, Chan KM, Huynh E, Williams SL, Stachowicz JJ. 2018 Plant genotype identity and diversity interact with mesograzer species diversity to influence detrital consumption in eelgrass meadows. *Oikos* **127**, 327–336. (doi:10.1111/oik.04471)

11. Loreau M. 1998 Biodiversity and ecosystem functioning: A mechanistic model. *Proc. Natl. Acad. Sci.* **95**, 5632–5636.

12. Naeem S, Li S. 1997 Biodiversity enhances ecosystem reliability. *Nature* **390**, 507–509. (doi:10.1038/37348)

13. Szűcs M, Vahsen ML, Melbourne BA, Hoover C, Weiss-Lehman C, Hufbauer RA. 2017 Rapid adaptive evolution in novel environments acts as an architect of population range expansion. *Proc. Natl. Acad. Sci.* **114**, 13501–13506. (doi:10.1073/pnas.1712934114)

14. Hufbauer RA, Rutschmann A, Serrate B, Conchard HV de, Facon B. 2013 Role of propagule pressure in colonization success: disentangling the relative importance of demographic, genetic and habitat effects. *J. Evol. Biol.* **26**, 1691–1699. (doi:10.1111/jeb.12167)

15. Szűcs M, Melbourne BA, Tuff T, Hufbauer RA. 2014 The roles of demography and genetics
in the early stages of colonization. *Proc. R. Soc. B Biol. Sci.* **281**, 20141073.
(doi:10.1098/rspb.2014.1073)

16. Okasha S. 2006 *Evolution and the Levels of Selection*. Clarendon Press.

17. Simberloff D. 2009 The role of propagule pressure in biological invasions. *Annu. Rev. Ecol.*
Evol. Syst. **40**, 81–152.

18. Vahsen ML, Shea K, Hovis CL, Teller BJ, Hufbauer RA. 2018 Prior adaptation, diversity,
and introduction frequency mediate the positive relationship between propagule pressure
and the initial success of founding populations. *Biol. Invasions* **20**, 2451–2459.
(doi:10.1007/s10530-018-1713-4)

19. Crawford KM, Whitney KD. 2010 Population genetic diversity influences colonization
success. *Mol. Ecol.* **19**, 1253–1263. (doi:10.1111/j.1365-294X.2010.04550.x)

20. Drummond EBM, Vellend M. 2012 Genotypic Diversity Effects on the Performance of
Taraxacum officinale Populations Increase with Time and Environmental Favorability. *PLoS*
ONE **7**. (doi:10.1371/journal.pone.0030314)

21. Forsman A, Wennersten L. 2016 Inter-individual variation promotes ecological success of
populations and species: evidence from experimental and comparative studies. *Ecography*
39, 630–648. (doi:10.1111/ecog.01357)

22. Bertness MD, Callaway R. 1994 Positive interactions in communities. *Trends Ecol. Evol.* **9**,
191–193. (doi:10.1016/0169-5347(94)90088-4)

23. Wuest SE, Niklaus PA. 2018 A plant biodiversity effect resolved to a single chromosomal
region. *Nat. Ecol. Evol.* **2**, 1933–1939. (doi:10.1038/s41559-018-0708-y)

24. Tonsor SJ. 1989 Relatedness and Intraspecific Competition in Plantago lanceolata. *Am.*
Nat. **134**, 897–906. (doi:10.1086/285020)

25. Hedge LH, Leung B, O'Connor WA, Johnston EL. 2014 The interacting effects of diversity
and propagule pressure on early colonization and population size. *J. Anim. Ecol.* **83**, 168–
175. (doi:10.1111/1365-2656.12125)

26. Flynn DFB, Mirotchnick N, Jain M, Palmer MI, Naeem S. 2011 Functional and phylogenetic
diversity as predictors of biodiversity–ecosystem-function relationships. *Ecology* **92**, 1573–
1581. (doi:10.1890/10-1245.1)

27. Simonsen AK, Chow T, Stinchcombe JR. 2014 Reduced plant competition among kin can
be explained by Jensen's inequality. *Ecol. Evol.* **4**, 4454–4466. (doi:10.1002/ece3.1312)

28. Fischer DG *et al.* 2017 Tree genetics strongly affect forest productivity, but intraspecific
diversity–productivity relationships do not. *Funct. Ecol.* **31**, 520–529. (doi:10.1111/1365-
2435.12733)

29. Madritch MD, Kingdon Clayton C., Singh Aditya, Mock Karen E., Lindroth Richard L.,
Townsend Philip A. 2014 Imaging spectroscopy links aspen genotype with below-ground

processes at landscape scales. *Philos. Trans. R. Soc. B Biol. Sci.* **369**, 20130194.
(doi:10.1098/rstb.2013.0194)

30. Lasky JR *et al.* 2015 Genome-environment associations in sorghum landraces predict
adaptive traits. *Sci. Adv.* **1**, e1400218. (doi:10.1126/sciadv.1400218)

31. Reynolds M, Tuberosa R. 2008 Translational research impacting on crop productivity in
drought-prone environments. *Genome Stud. Mol. Genet. Ed. Juliette Meaux Maarten*
Koornneef Plant Biotechnol. Ed. Andy Greenl. Jan Leach **11**, 171–179.
(doi:10.1016/j.pbi.2008.02.005)

32. Gienapp P, Fior S, Guillaume F, Lasky JR, Sork VL, Csilléry K. 2017 Genomic Quantitative
Genetics to Study Evolution in the Wild. *Trends Ecol. Evol.* **32**, 897–908.
(doi:10.1016/j.tree.2017.09.004)

33. VanRaden PM, Van Tassell CP, Wiggans GR, Sonstegard TS, Schnabel RD, Taylor JF,
Schenkel FS. 2009 Invited Review: Reliability of genomic predictions for North American
Holstein bulls. *J. Dairy Sci.* **92**, 16–24. (doi:10.3168/jds.2008-1514)

34. Jannink J-L, Lorenz AJ, Iwata H. 2010 Genomic selection in plant breeding: from theory to
practice. *Brief. Funct. Genomics* **9**, 166–177. (doi:10.1093/bfgp/elq001)

35. Wuest SE, Pires ND, Luo S, Vasseur F, Messier J, Grossniklaus U, Niklaus PA. 2019
Increasing plant group productivity through latent genetic variation for cooperation. *bioRxiv* ,
641449. (doi:10.1101/641449)

36. Mitchell-Olds T, Schmitt J. 2006 Genetic mechanisms and evolutionary significance of
natural variation in Arabidopsis. *Nature* **441**, 947–952. (doi:10.1038/nature04878)

37. Alonso-Blanco C *et al.* 2016 1,135 Genomes Reveal the Global Pattern of Polymorphism in
Arabidopsis thaliana. *Cell* **166**, 481–491. (doi:10.1016/j.cell.2016.05.063)

38. Vasseur F, Sartori K, Baron E, Fort F, Kazakou E, Segrestin J, Garnier E, Vile D, Violle C.
2018 Climate as a driver of adaptive variations in ecological strategies in Arabidopsis
thaliana. *Ann. Bot.* **122**, 935–945. (doi:10.1093/aob/mcy165)

39. Lovell JT *et al.* 2013 Pleiotropy of FRIGIDA enhances the potential for multivariate
adaptation. *Proc. R. Soc. B Biol. Sci.* **280**, 20131043. (doi:10.1098/rspb.2013.1043)

40. Huston MA, McBride AC. 2002 Evaluating the relative strengths of biotic versus abiotic
controls on ecosystem processes. *Biodivers. Ecosyst. Funct. Synth. Perspect. Oxf. Univ.*
Press Oxf. , 47–60.

41. Easlon HM, Bloom AJ. 2014 Easy Leaf Area: Automated digital image analysis for rapid and
accurate measurement of leaf area1. *Appl. Plant Sci.* **2**. (doi:10.3732/apps.1400033)

42. Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. 2012 A high-performance
computing toolset for relatedness and principal component analysis of SNP data.
Bioinforma Oxf Engl **28**. (doi:10.1093/bioinformatics/bts606)

43. Schuetzenmeister A, Dufey F. 2018 *VCA: Variance Component Analysis*. See
https://CRAN.R-project.org/package=VCA.

44. Endelman JB. 2011 Ridge Regression and Other Kernels for Genomic Selection with R
Package rrBLUP. *Plant Gen* **4**, 250–255.

45. Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, Eskin E. 2008 Efficient
Control of Population Structure in Model Organism Association Mapping. *Genetics* **178**,
1709–1723. (doi:10.1534/genetics.107.080101)

46. Atwell S *et al.* 2010 Genome-wide association study of 107 phenotypes in Arabidopsis
thaliana inbred lines. *Nature* **465**, 627–631. (doi:10.1038/nature08800)

47. Zhang Y, Chen HYH, Reich PB. 2012 Forest productivity increases with evenness, species
richness and trait variation: a global meta-analysis. *J. Ecol.* **100**, 742–749.
(doi:10.1111/j.1365-2745.2011.01944.x)

48. Hooper DU *et al.* 2012 A global synthesis reveals biodiversity loss as a major driver of
ecosystem change. *Nature* **486**, 105–108. (doi:10.1038/nature11118)

49. Isbell F *et al.* 2015 Biodiversity increases the resistance of ecosystem productivity to climate
extremes. *Nature* **526**, 574–577.

50. Lasky JR, Uriarte M, Boukili VK, Erickson DL, John Kress W, Chazdon RL. 2014 The
relationship between tree biodiversity and biomass dynamics changes with tropical forest
succession. *Ecol. Lett.* **17**, 1158–1167. (doi:10.1111/ele.12322)

51. Becker J, Eisenhauer N, Scheu S, Jousset A. 2012 Increasing antagonistic interactions
cause bacterial communities to collapse at high diversity. *Ecol. Lett.* **15**, 468–474.
(doi:10.1111/j.1461-0248.2012.01759.x)

52. Donohue K. 2003 The Influence of Neighbor Relatedness on Multilevel Selection in the
Great Lakes Sea Rocket. *Am. Nat.* **162**, 77–92. (doi:10.1086/375299)

53. Dudley SA, File AL. 2007 Kin recognition in an annual plant. *Biol. Lett.* **3**, 435–438.
(doi:10.1098/rsbl.2007.0232)

54. Rooney WL, Blumenthal J, Bean B, Mullet JE. 2007 Designing sorghum as a dedicated
bioenergy feedstock. *Biofuels Bioprod. Biorefining* **1**, 147–157. (doi:10.1002/bbb.15)

55. Weiner J, Thomas SC. 1986 Size Variability and Competition in Plant Monocultures. *Oikos*
47, 211–222. (doi:10.2307/3566048)

56. Wardle DA, Nilsson M-C, Gallet C, Zackrisson O. 1998 An ecosystem-level perspective of
allelopathy. *Biol. Rev.* **73**, 305–319. (doi:10.1017/S0006323198005192)

57. Bomblies K, Yant L, Laitinen RA, Kim S-T, Hollister JD, Warthmann N, Fitz J, Weigel D.
2010 Local-Scale Patterns of Genetic Variability, Outcrossing, and Spatial Structure in
Natural Stands of Arabidopsis thaliana. *PLOS Genet.* **6**, e1000890.
(doi:10.1371/journal.pgen.1000890)

58. Mitchell-Olds T. 1996 Genetic Constraints on Life-History Evolution: Quantitative-Trait Loci
Influencing Growth and Flowering in Arabidopsis Thaliana. *Evolution* **50**, 140–145.
(doi:10.1111/j.1558-5646.1996.tb04480.x)

675