Title: Erythropoietin directly remodels the clonal composition of murine hematopoietic multipotent progenitor cells

A.S. Eisele[1], J. Cosgrove[1], A. Magniez[1], E. Tubeuf[1], S. Tenreira Bento[1], C. Conrad[1], F. Cayrac[1], T. Tak[1], A.M. Lyne[1], J. Urbanus[2] and L. Perié[1*].

Affiliations

1. Institut Curie, Université PSL, Sorbonne Université, CNRS UMR168, Laboratoi re Physico Chimie Curie, 75005 Paris, France
2. Netherlands Cancer Institute, Amsterdam, Netherlands

Contact information

Leila Perié, Postal address: UMR168, Institute Curie, 11 rue Pierre et Marie Curie, 75005 Paris, Leila.Perie@curie.fr, +33156246229.

Abstract

The cytokine erythropoietin (EPO) is a potent inducer of erythrocyte development and

one of the most prescribed biopharmaceuticals. The action of EPO on erythroid

progenitor cells is well established, but its direct action on hematopoietic stem and

progenitor cells (HSPCs) is still debated. Here, using cellular barcoding, we traced the

differentiation of hundreds of single murine HSPCs, after *ex vivo* EPO-exposure and

transplantation, in five different hematopoietic cell lineages, and observed the transient

occurrence of high-output Myeloid-Erythroid-megaKaryocyte (MEK)-biased and

Myeloid-B-cell-Dendritic cell (MBDC)-biased clones. Single-cell RNA sequencing

(ScRNAseq) analysis of *ex vivo* EPO-exposed HSPCs revealed that EPO induced the

upregulation of erythroid associated genes in a subset of HSPCs, overlapping with

multipotent progenitor (MPP) 1 and MPP2. Transplantation of Barcoded EPO-

exposed-MPP2 confirmed their enrichment in Myeloid-Erythroid-biased clones.

Collectively, our data show that EPO does act directly on MPP independent of the

niche, and modulates fate by remodeling the clonal composition of the MPP pool.

Introduction

Erythrocytes are the most numerous hematopoietic cells in our body and are constantly

renewed[1]. The major inducer of erythroid cell development in steady state and anemic

conditions is the cytokine EPO[2]. Recombinant EPO is widely used to treat anemia and

is one of the most sold biopharmaceuticals[3]. Previously, EPO was thought to solely

target erythroid-committed progenitors and induce their increased proliferation and

survival via the EPO receptor (EPOR)[4]. Recently, EPO has also been suggested to act

on HSPCs [5–12], but the nature of EPO´s effect on HSPC fate remains unresolved

despite potential adverse side effects during long-term EPO usage in the clinics and

associations of high EPO levels with leukemias[13–15].

It is well established that EPO can induce HSPCs to cycle, as evidenced by a number

of bulk and single cell studies *in vitro* and *in vivo*[5,12,8,9,11]. It's role in modulating HSPC

fate is less clear however, with a lack of studies that functionally assess HSPC fate at

the single cell level *in vivo*, and analyze the direct effect of EPO on HSPCs, and not

the effect of the surrounding niche. More specifically, the upregulation of erythroid

associated genes in HSPC (LSK $CD150^+$ $Flt3^-$ $CD48^-$) in response to *in vivo* EPO has

been observed with bulk transcriptomics[11], suggesting that HSPCs deviate their fate

toward erythroid production. Recent ScRNAseq analysis observed different changes

of lineage-associated gene expressions after *in vivo* EPO-exposure of HSPCs[9,8]. As

changes of gene expression do not necessarily result in cell fate modification[14],

functional validations *in vivo* are necessary. In one study such functional validation was

performed using bulk transplantation of *in vivo* EPO-exposed HSPCs (LSK $CD150^+$

$Flt3^-$)[7]. This study showed increased erythroid production and decreased myeloid cell

production, concluding that EPO deviates the fate of HSPCs in favor of erythroid

production. As EPO has also been shown to target hematopoietic niche cells

(osteoblasts and osteocytes[15,16] endothelial cells[17,16], adipocytes[18,19], and

mesenchymal stem cells[6,20]), it remains however still unclear whether EPO acts directly

on HSPCs, via their environment, or both.

It is now established that HSPCs encompass cells with different long-term

reconstitution capacity after transplantation, as well as a heterogeneous output in

terms of quantity (lineage-bias) and type of cells (lineage-restriction)[21–28]. Recently, the

HSPC compartment was subdivided into long-term hematopoietic stem cells (LT-HSC)

and different multipotent progenitors (MPP) (MPP1-4[29,30]), with variation around the

phenotypic definition[31]. Interestingly, HSPC composition responds to irradiation with

HSC transiently self-renewing less and increasing their production of MPP2-3[31]. In the

case of EPO, conflicting results suggest that either HSC or MPP respond to EPO [8,9,11]

but the difference in HSPC definition and the lack of functional validation make it

difficult to compare these studies. As HSPCs are functionally heterogeneous, and the

current phenotypic definition partially capture this heterogeneity, a single cell in vivo

lineage tracing approach is needed to assess whether EPO can influence HSPC fate

decisions.

To analyze the functional effect of EPO on the differentiation of individual HSPCs (C-

Kit$^+$ Sca1$^+$ CD150$^+$ Flt3$^-$) removing the effect of the niche, we here utilized cellular

barcoding technology which allowed us to trace the progeny of hundreds of single

HSPCs *in vivo*. By analyzing cellular barcodes in five mature hematopoietic lineages

and HSPCs, we observed transient induction of high-output MEK-biased barcode

clones compensated by MBDC-biased clones after ex vivo EPO-exposure. ScRNAseq

of *ex vivo* EPO-exposed HSPCs revealed upregulation of erythroid associated genes

in a subset of the compartment with overlap to gene signatures of MPP1 (C-Kit$^+$ Sca1$^+$

Flt3$^-$ CD150$^+$ CD48$^-$ CD34$^+$), and MPP2 (C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$ CD48$^+$)[30,29] and not

of LT-HSCs (C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$ CD48$^-$ CD34$^-$). Transplantation of barcoded

MPP2 confirmed their enrichment in ME-biased clones in response to EPO. Moreover,

the increased contribution of biased HSPC clones to the mature cell lineages after

EPO-exposure did not match their frequency in HSPCs, indicating that they were

differentiating more than self-renewing, a property associated to multipotent

progenitors. The transient effect of EPO on HSPCs further corroborates an action of

EPO on MPP1/2 rather than LT-HSCs. Altogether our results are consistent with a

model in which perturbations induce clonal remodeling of HSPC contributing to

hematopoiesis, with biased MPPs transiently contributing more than LT-HSC. They

also demonstrate a direct effect of EPO on MPPs after transplantation with implications

for basic HSC research and therapeutic applications in the clinic.

Results

EPO-exposure induces biases in single HSPCs. Given the debate surrounding

which HSPC subset is responding to EPO, we decided to analyzed the direct effect of

EPO on the differentiation of HSPCs defined as C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$

(encompassing LT-HSC (C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$ CD48$^-$ CD34$^-$), MPP1 (C-Kit$^+$

Sca1$^+$ Flt3$^-$ CD150$^+$ CD48$^-$ CD34$^+$), and MPP2 (C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$ CD48$^+$)[30,29])

at the single cell level by cellular barcoding. To this purpose, we generated a new high-

diversity lentiviral barcode library (LG2.2, 18,026 barcodes in reference list), consisting

of random 20 nucleotides sequences positioned adjacent to the green fluorescent

protein (GFP) gene, enabling the tracking of many individual cells in parallel. Using this

LG2.2 library, we labeled single HSPCs (Figure 1- figure supplement 1a) with unique

genetic barcodes as previously described[33], exposed them to EPO (1,000 ng/ml) or

PBS for 16 hours *ex vivo* and transplanted around 2,600 cells (Mean 2,684 cells +/-

175 cells) of which around 10% barcoded cells into irradiated mice (Figure 1a). Note

that HSPCs kept their sorting phenotype after *ex vivo* culture albeit a slight

downregulation of C-Kit[44] and Flt3 (Figure 1 – figure supplement 1f). At day 30 after

transplantation, the earliest timepoint at which HSPCs produce simultaneously

erythroid, myeloid and lymphoid cells[45], barcoded (GFP$^+$) erythroblasts (E; Ter119$^+$

CD44^{+34}), myeloid cells (M; Ter119$^-$ CD19$^-$ CD11c$^-$ CD11b$^+$), and B-cells (B; Ter119$^-$

CD19$^+$) (Figure 1 – figure supplement 1b-c, e) were sorted from the spleen and their

barcode identity assessed through PCR and deep-sequencing. Note that bone and

spleen had similar barcoding profiles (Figure 1 – figure supplement 2). No difference

in chimerism was observed between the EPO and control group in the spleen and

blood, even when mTdTomato/mGFP donor mice were used to better assess the

erythroid lineage (Figure 1b-c). On average, we detected around eighty barcodes per

mouse, of which most were detected in several lineages (Figure 1d). Comparison of

the numbers of barcodes producing each lineage showed that EPO-exposure resulted

in the same number of engrafting and differentiating cells as in control (Figure 1d).

Notably, the number of erythroid restricted cells remained stable in the EPO group as

compared to control (Figure 1e), indicating that the response to EPO is more complex

than a direct instruction of erythroid-restricted HSPCs.

To quantify the effect of EPO on HSPC lineage-biases, barcode-labeled HSPCs were

classified based on the balance of their cellular output in the M, B and E lineages. With

this classification using a 10% threshold, cells classify for example as ME-biased if

they have above 10% of their output in the M and E lineage, and under 10% of their

output in the B lineage (Figure 1e other thresholds in Figure 1 – figure supplement 3b).

Interestingly, application of this classification revealed that although the proportion of

lineage-biased HSPCs in control and EPO group was similar (Figure 1f), their

contribution to the different lineages was increased by EPO-exposure (Figure 1g). In

the control group, balanced HSPCs (MBE) produced the majority of all lineages, as

previously published[46]. In the EPO group, ME- and MB-biased clones produced most

cells of the analyzed lineages (Figure 1g). ME-biased HSPCs produced the majority of

erythroid cells (57% +/- 10%), MB-biased HSPCs produced the majority of B-cells

(58% +/- 36%), and ME- and MB-biased clones contributed the majority of myeloid

cells (MB-biased 45% +/- 38% and ME-biased 20% +/- 13%, together 65% +/- 25%).

To test the significance of this effect, we used a permutation test, which compares the

effect size between control and EPO group to the one of all random groupings of

mice[47]. The contributions of the ME- and of the MB-biased HSPC classes to the

different lineages were significantly different in EPO and control groups (Table 1).

These results were reproduced in an additional experiment (Figure 1 – figure

supplement 3c-f, Supplementary File 1). A lower EPO concentration (160 ng/ml) as

well as an additional single injection of EPO (133 ug/kg) during transplantation gave

similar results (Figure 2, Figure 2 – figure supplement 1a, Table 1). Also at six weeks

post transplantation, similar results were obtained (Figure 1 – figure supplement 4,

Supplementary File 1). In summary, *ex vivo* EPO priming of HSPCs modified the output

balance of HSPCs rather than the number of lineage-restricted and -biased cells.

Balanced clones produced a smaller percentage of the mature cells; ME-biased

HSPCs produced most of the erythroid cells and MB-biased HSPCs produced most of

the B cells.

Contribution of ME- and MB-biased HSPCs to the DC and MkP lineage. To further

characterize the cells produced by the ME-biased and MB-biased HSPCs, we repeated

our experimental setup including the analysis of the megakaryocyte and dendritic cell

(DC) lineages (Figure 3). Megakaryocyte Progenitors (MkP) were chosen as proxy for

the production of platelets which are not suitable for barcode analysis. Barcoded

(GFP$^+$) DCs (DC; Donor Ter119$^-$ CD19$^-$ CD11c$^+$ CD11b$^-$) and MkP (MkP; C-Kit$^+$ Sca-1$^-$

CD150$^+$ CD41$^+$) (Figure 1 – figure supplement 1c-e) were sorted together with M, E,

and B cells, 4 weeks after transplantation of control or EPO-exposed HSPCs (1,000

ng/ml). In both groups, the majority of clones produced also DCs (Figure 3a). In the

control group, balanced HSPCs produced the majority of DCs (65% +/- 9%) (Figure

3b). However, in the EPO group, balanced HSPCs decreased their contribution to the

DC lineage (36% +/-25%) and MB-biased HSPCs significantly increased their

contribution (86% +/- 43% EPO vs 22% +/- 11% control group) (Figure 3b, Table 1),

thus, they were MBDC-biased HSPCs. In contrast the ME-biased HSPCs produced

few DCs in both groups (Figure 3b), indicating that ME-biased HSPCs are restricted

both in their B and DC production compared to the M and E production.

The majority of the MkP production came from the ME-biased HSPCs in both groups

(58% +/- 21% control and 55% +/- 14% EPO group, Figure 3c-d), indicating that ME-

biased HSPCs were also MkP-biased HSPCs (thus MEK-biased). We did not detect a

high contribution of MkP-restricted HSPCs[25,24,48] to the MkP lineage (Figure 3d).

Finally, as high EPO-exposure has been linked to changes in macrophage numbers[49–

56] we analyzed the contribution of control and EPO-exposed HSPCs to the myeloid

lineage in more detail, but could not detect changes in the percentage of the different

myeloid subsets produced (Figure 3 – figure supplement 1a-c).

Effect of EPO on short-term HSPC self-renewal. In light of previous studies which

suggested changes in HSPC proliferation after *in vivo* EPO-exposure[5,12,8,9,11], we next

explored if the short-term self-renewal capacity of HSPCs was impacted. To this end,

we analyzed barcodes in bone marrow HSPCs in addition to the spleen E, M, and B

lineages at week 4 after transplantation of control or EPO-exposed HSPCs (160 and

1,000 ng/ml) (Figure 4). We reasoned that barcodes of HSPCs differentiating and

short-term self-renewing (dividing to give rise to other HSPCs) after transplantation are

detected in both compartments, while detection in only HSPCs or mature lineages

indicates a prevalence of short-term self-renewal or differentiation respectively. Most

of the barcodes detected in HSPCs overlapped with barcodes in the mature cells

(Figure 4b left) in both the control and two EPO groups, showing that most of the

transplanted cells had given rise to other HSPCs and differentiated irrespective of the treatment. Some barcodes were only detected in mature cells (Figure 4b right), indicating that some HSPCs had only differentiated or were below the limit of detection. These HSPCs were equally abundant in the control and two EPO groups (Figure 4b right).

To analyze if different lineage biases correlated to different short-term self-renewal capacity, we analyzed the proportion of biased HSPC classes, as previously defined, within the HSPC compartment (Figure 4c). In the control group, balanced and ME-biased HSPCs contributed most to the HSPC reads (34% +/- 36% MBE and 37% +/- 34% ME-biased HSPCs), while barcodes of MB-biased HSPCs contributed less (15% +/- 32%) (Figure 4c), a trend that has been previously described[57,58,25]. Surprisingly, the pattern of contributions of different biased HSPC subsets to HSPC reads was unchanged in the EPO groups (Figure 4c), implying that the extent of short-term self-renewal was unchanged after *ex vivo* EPO-exposure.

To study if the increased production of cells by the ME- and MB-biased HSPCs to the mature cells observed after *ex vivo* EPO-exposure (Figure 1-3) correlated with short-term self-renewal capacity of HSPCs, we analyzed the contribution of barcodes detected or not in HSPCs to the E, M, and B lineages (Figure 4d). In the control group, the majority of mature cells were derived from barcodes also present in HSPCs. However, in both EPO groups, the contribution of barcodes detected in HSPCs to mature cells was significantly lower (Figure 4d, Table 1) implying that the increased contribution of biased HSPC classes to the mature cell lineages after *ex vivo* EPO-exposure was most likely caused by cells differentiating more than short-term self-renewing.

EPO-exposure induces an erythroid program in a subgroup of HSPCs. To further characterize the effect of EPO-exposure on HSPCs, we performed scRNAseq of barcoded C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$ cells after *ex vivo* culture in medium supplemented with EPO or PBS using the 10X Genomics Chromium platform. 1,706 cells from control and 1,595 cells from the EPO group passed our quality control. To compare the HSPCs injected with non-cultured hematopoietic cells, we generated a reference map of 44,802 C-kit$^+$ cells from[42] and used published signatures as detailed in the supplementary materials and methods[31,43] to annotate this map (Figure 5 – figure supplement 1a,b and e,f). Projection of our single cell data on this map showed that both the control and the EPO-exposed HSPCs similarly overlapped with non-MPP4 LSK cells, according to their sorting phenotype (Figure 5 – figure supplement 1e,f). These results indicate that neither the *ex vivo* culture itself nor the EPO treatment dramatically affected the global identity of the sorted HSPCs.

When comparing the EPO and control group, we found 1,176 differentially expressed genes (Figure 5a and Supplementary File 2) and this number was significantly higher than the number expected due to chance (p-value=0,01) as assessed by permutation testing. Among the most upregulated genes in the EPO-exposed HSPCs, were genes with erythroid association as *Hbb-bs*, *Erdr1*, *Wtap*, *Kmt2d*, or *Nfia*[59], and GATA1 targets (*Abhd2*, *Cbx3*, *Kdelr2*, *Pfas*), cell cycle related genes (*Tubb5*, *Hist1h2ap*) as well as genes previously described to be induced in HSPCs after *in vivo* EPO-exposure, such as *Bmp2k*[6] and *Ifitm1*[8] (Figure 5a). Genes involved in stem cell maintenance, such as *Serpina3g*, *Mecom, Txnip*, *Meis1*, *Pdzk1ip1*[8], *Sqstm1*[60], *Smad7*[61], *Aes*[62], were among the most downregulated genes in the EPO-exposed HSPCs (Figure 5a).

As our cellular barcoding data suggests that single HSPCs differ in their response to EPO, we assessed the heterogeneity of EPO responses at the transcriptomic level. UMAP-based visualization of the data suggested that a subgroup of EPO-exposed cells was transcriptomically distinct (Figure 5b), independently of the number of PCA components and genes used in the analysis (Figure 5 – figure supplement 1c). To test this observation, we defined an EPO response signature based on differentially expressed genes between the EPO and control group. Plotting the expression of the EPO response signature at the single cell level showed that the majority of the transcriptomic differences between the control and EPO group were indeed driven by this small subgroup of cells (Figure 5c). Reasoning that this subgroup contains the cells directly responding to EPO, we defined as EPO-responders, cells in the 90[th] percentile of EPO response signature expression (Figure 5c) for subsequent analysis. Importantly, unsupervised clustering analysis of the data (Figure 5 – figure supplement 2a-b) showed similar results. The genes encoding EPOR, as well as the alternative EPO receptors EphB4, CD131, CRFL3 were equally expressed between the EPO-responders, non-responders and control groups (Figure 5 – figure supplement 1d). Reasoning that the EPO-responders correspond to MEK-biased HSPCs, we also looked for potential MBDC-biased HSPCs but could not detect a subgroup of cells with upregulation of lymphoid associated genes, suggesting that the MBDC-bias is not a direct effect of EPO-exposure but more an indirect effect. In summary, the scRNAseq analysis corroborated our functional barcoding data, showing that a subset of HSPCs can respond directly to EPO stimulation.

EPO-responder HSPCs overlap with MPP1 and MPP2 signatures. As our barcode analysis suggested that the effect of direct EPO-exposure on HSPCs is caused by cells

differentiating more than self-renewing, we next wanted to assess which of the HSPC

subsets are the EPO responders in our scRNAseq dataset. We annotated the UMAP-

based visualization of our data with published signatures of the HSC (dormant HSCs[40]

and LT-HSC[41]), MPP1[40] and MPP2[31] subsets included in our HSPC gate, and

analyzed its overlap with the previously defined EPO-responder and non-responder

cells (Figure 6a,b). Relative to the control group and non-responders of the EPO-

group, the EPO-responders had a reduced expression of HSC gene signatures and

increased expression of MPP1 and MPP2 signatures (Figure 6a,b). An annotation of

the reference map generated from data of[42] likewise showed a low overlap of EPO-

responders with the most quiescent HSC subsets (Figure 6c,d). The independent

analysis using unsupervised clustering further supported this result (figure 5 – figure

supplement 2c,d). All in all, our scRNAseq analysis implied that, in line with our

barcoding results, the HSPCs directly reacting to EPO are most likely multipotent

progenitor cells of the MPP1 and MPP2 subsets.

EPO-exposure induces ME-biases in single MPP2.

To confirm that MPP2 are a subset within HSPCs reacting directly to EPO as predicted

by the scRNAseq analysis, we transplanted barcoded control or EPO-exposed (1,000

ng/ml) MPP2 together with unbarcoded CD48$^-$ HSPCS (C-Kit$^+$ Sca1$^+$ Flt3$^-$ CD150$^+$

CD48$^-$), (Figure 7 – figure supplement 1a) and analyzed their barcoded progeny in the

E, M, and B lineages of the spleen at week 4 after transplantation (Figure 7). We found

an equivalent engraftment as for the entire HSPC compartment and no difference

between the EPO-treated and the control group (Figure 7a-b). Applying the same

classification as in Figure 1 to quantify the effect of EPO on MPP2 lineage-biases, we

observed that, as for the whole HSPC compartment, ME-biased cells contributed more

to the M and E lineages (Figure 7d-e, other threshold in Figure 7 – figure supplement

1c). Similarly to our data on whole HSPC compartment (Figure 1g) the proportion of

the differently biased-MPP2 was similar between control and EPO group (Figure 7c).

This data confirms that the MPP2 population is enriched in HSPCs responding to EPO.

Transient effect of EPO-exposure.

Finally, we reasoned that if EPO directly acts on multipotent progenitors MPP1/2 with

short reconstitution capacity after transplantation rather than long-term repopulating

HSC, then the EPO effect should be transient. To test this hypothesis, we repeated the

experiment and analyzed barcodes in the E, M, and B lineages at 4 months after

transplantation of control or EPO-exposed HSPCs (160 and 1,000 ng/ml) (Fig. 8). In

the control group, as reported before[63], the chimerism at 4 months was higher and the

number of barcodes detected was lower than at one month post-transplantation (Fig.

8c-d and 1b-d). We detected no significant changes in the clonal output of HSPCs

between control and EPO group at this timepoint (Fig. 8e), with the majority of cells in

all lineages produced by balanced HSPCs (Fig. 8a-b), implying that the effect of direct

EPO-exposure on HSPCs is transient. This confirms that the effect of direct EPO-

exposure on HSPCs is likely caused by multipotent progenitor cells with a short

reconstitution capacity after transplantation.

Discussion

EPO is a key regulator of hematopoiesis, and is classically considered to support the

proliferation and survival of erythroid committed progenitors. By analyzing the *in vivo*

fate of hundreds of EPO-stimulated vs untreated transplanted (c-Kit$^+$ Sca1$^+$ Flt3$^-$

CD150$^+$) HSPCs at the single-cell level, we established that EPO can change HSPC

differentiation in the absence of an EPO-stimulated bone marrow microenvironment.

Collectively our results yield 2 important conclusions: (i) EPO has a direct effect on

HSPCs, that is not solely due to the effects of EPO on the surrounding niche and (ii)

EPO directly remodels the clonal composition of HSPC by inducing fate biased MPP

and reducing the output of HSC.

Specifically, we observe that EPO induced MEK-biased (ME) and MBDC-biased (MB)

HSPCs that produced the majority (>60%) of mature hematopoietic cells at four and

six weeks after transplantation. In contrast, balanced HSPCs (MBE) had a reduced

output of mature cells in response to EPO. The increased erythroid-associated gene

signature in a subset of HSPCs after *ex vivo* EPO-exposure suggests that EPO directly

induces high output MEK-biased HSPCs, which is indirectly compensated for by the

occurrence of high-output of MBDC-biased HSPCs to maintain a balanced production

of hematopoietic cells.

These biased clones had a higher propensity to differentiate than to self-renew, and

their response to EPO was transient, suggesting that EPO-responsive cells are multi-

potent progenitors, and not LT-HSCs. This is supported by transcriptomic analysis

showing that EPO-responders express the MPP1/MPP2 gene-signatures.

Transplantation of barcoded EPO-exposed MPP2 confirmed their enrichment in ME-

biased clones in response to EPO.

Similar to studies that assessed the effect of high systemic EPO-exposure [10,9,11,8], we

found that multipotent progenitors, not HSC, are responding to EPO. The occurrence

of myeloid, megakaryocytic and erythroid gene expression in MPP1 after bleeding [9] is

in line with our findings. Previously, long-term EPO exposure in Tg6 transgenic mice

did however not change the in vitro differentiation outcome of MPP2 [11]. Furthermore,

we did not detect a fate deviation toward erythroid production at the expense of myeloid production as seen for in vivo EPO-exposed HSPCs after transplantation[7]. These differences could be due both to the duration and route of EPO exposure as well as the indirect effects of systemic EPO-exposure through other cells, for example from the bone marrow niche.

In addition, our data shows that direct cytokine-stimulation leads to a clonal remodeling of the HSPC compartment, with a transient increase in the contribution of fate-biased MPPs. Without longitudinal barcoding data within the same animal, we cannot distinguish if EPO is transiently changing the fate and outcome of the same HSPCs over time or if EPO is pushing the differentiation of some HSPCs that will be replaced by more balanced and stable HSPCs.

Different studies have suggested that the behavior of transplanted HSPC differ from native HSPC[64,65]. Transplantation seems to favor the long-term output from HSC whereas steady state hematopoiesis is maintained more by MPPs contribute than LT-HSC[64,65,66]. Our work is in line with a model in which MPP are a highly malleable cell population that can rapidly respond to changing demands for new cells, such as transplantation or infection[31].

The direct effect of EPO on MPPs we described here could be one of the factors underlying the development of adverse side effects and co-morbidities during long-term EPO use in the clinics and associations of high EPO levels with leukemias[67,13,68]. To translate these results to the clinics and understand the side effect of EPO treatment, further work is required to determine if HSPCs and erythroid progenitors like CFU-E are responding to the same dose and duration of EPO exposure.

Materials and Methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Strain, strain background (*Mus musculus*)	C57BL/6J CD45.1[+],	Jackson Laboratory	B6.SJL-*Ptprc[a] Pepc[b]*/BoyJ, stock No 002014, B6 Cd45.1	male
Strain, strain background(*Mus musculus*)	C57BL/6J CD45.2[+],	Jackson Laboratory	C57BL/6J, stock No 000664, B6	male
Strain, strain background (*Mus musculus*)	*Rosa26CreER[T2];mT/mG*	Jackson Laboratory	STOCK *Gt(ROSA)26Sor[tm4(ACTB-tdTomato,-EGFP)Luo]*/J, stock No 007576, mT/mG, mTmG	male
Strain, strain background (*Escherichia coli*)	ElectroMAX[TM] Stbl4[TM] Competent Cells	ThermoFisher Scientific	Cat# 11635018	
Recombinant DNA reagent	pRRL-CMV-GFP plasmid[32]	PMID 9765382		Ton Schumacher lab, NKI, Amsterdam
Cell line (*human*)	HEK293T cells	other		Philippe Benaroch lab, Institute Curie, Paris
Recombinant DNA reagent	p8.9-QV	other		Philippe Benaroch lab, Institute Curie, Paris
Recombinant DNA reagent	pVSVG	other		Philippe Benaroch lab, Institute Curie, Paris

Chemical compound, drug	anti-CD117 magnetic beads	Miltenyi	Cat# 130-091-224, **RRID:AB_2753213**	
Chemical compound, drug	Propidium iodide	Sigma	Cat# 81845	
Chemical compound, drug	StemSpanMedium SFEM	STEMCELL Technologies	Cat# 9650	
Chemical compound, drug	Mouse recombinant SCF	STEMCELL Technologies	Cat# 78064.2	
Chemical compound, drug	Eprex, erythropoietin alpha	Janssen		
Chemical compound, drug	anti-biotinylated beads	Miltenyi	Cat# 130090485, **RRID:AB_244365**	
Antibody (rat, monoclonal)	anti-Ter119-biotin	BD Biosciences	Cat# 553672, clone TER119, **RRID:AB_394985**	(1:100)
Antibody (mouse, monoclonal)	anti-cd45.1-PE	BD Biosciences	Cat# 553776, clone A20, **RRID:AB_395044**	(1:100)
Antibody (rat, monoclonal)	anti-Ter119-PE-Cy7	BD Biosciences	Cat# 557853, clone TER119, **RRID:AB_396898**	(1:100)
Antibody (hamster, monoclonal)	anti-cd11c-APC	eBioscience	Cat# 17-0114-82, clone N418, **RRID:AB_469346**	(1:100)
Antibody (rat, monoclonal)	anti-cd19-APC-Cy7	BD Biosciences	Cat# 557655, clone ID3, **RRID:AB_396770**	(1:100)

Antibody (rat, monoclonal)	anti-cd11b-PerCP-Cy5.5	eBiosciences	Cat# 45-0112-82, clone M1/70, **RRID:AB_953558**	(1:100)
Antibody (rat, monoclonal)	anti-cd117-APC	BioLegend	Cat# 105812, clone 2B8, **RRID:AB_313221**	(1:100)
Antibody (rat, monoclonal)	anti-cd135-PE	eBiosciences	Cat# 12 135182, clone A2F10, **RRID:AB_465859**	(100)
Antibody (rat, monoclonal)	anti-cd135-PE-Cy5	Life technologies	Cat# 15_1351_82, clone A2F10, **RRID:AB_494219**	(1:100)
Antibody (rat, monoclonal)	anti-Sca1-PacificBlue	BioLegend	Cat# 122520, clone D7, **RRID:AB_2143237**	(1:200)
Antibody (rat, monoclonal)	anti-cd150-PE-Cy7 (Mouse-monoclonal)	BioLegend	Cat# 115914, clone TC15-12F12.2, **RRID:AB_439797**	(1:100)
Antibody (rat, monoclonal)	anti-cd44-PE	BD Biosciences	Cat# 553134, clone IM7, **RRID:AB_394649**	(100)
Antibody (rat, monoclonal)	anti-cd41-BV510	BD Biosciences	Cat# 740136, clone MVVREG30, **RRID:AB_2739892**	(1:100)
Antibody (rat, monoclonal)	anti-Siglec-F-PE-CF594	BD Biosciences	Cat# 562757, clone E50-2440, **RRID:AB_2687994**	(1:200)
Antibody (rat, monoclonal)	anti-Ly6g-BV510	BioLegend	Cat# 127633. clone 1A8, **RRID:AB_2562937**	(1:200)

Antibody (rat, monoclonal)	anti-cd115-PE	BioLegend	Cat# 135505, clone AFS98, **RRID:AB_1937254**	(1:200)
Antibody (hamster, monoclonal)	anti-cd48-APC-Cy7	BD Biosciences	Cat# 561242 clone HM48-1, **RRID:AB_10644381**	(1:100)
Chemical compound, drug	Viagen Direct PCR Lysis Reagent (cell)	Euromedex	Cat# 301-C	
Chemical compound, drug	Proteinase K Solution RNA grade	Invitrogen	Cat# 25530-049	
Sequence-based reagent	top-LIB	This paper	PCR primer	TGCTGCCGTCAACTAGA ACA
Sequence-based reagent	bot-LIB	This paper	PCR primer	GATCTCGAATCAGGCG CTTA
Sequence-based reagent	PCR2-Read1-plate-index-forward	This paper	PCR primer	ACACTCTTTCCCTACAC GACGCTCTTCCGATCT NNNNCTAGAACACTCG AGATCAG
Sequence-based reagent	PCR2-Read2-reverse	This paper	PCR primer	GTGACTGGAGTTCAGA CGTGTGCTCTTCCGAT CGATCTCGAATCAGGC GCTTA
Sequence-based reagent	PCR3-P5-forward	This paper	PCR primer	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT
Sequence-based reagent	PCR3-P7-sample-index-reverse	This paper	PCR primer	CAAGCAGAAGACGGCATACGAGANNNNNNGT

				GACTGGAG TTCAGA CGTGCTCT TCCGATC
Commercial assay or kit	Agencourt AMPure XP system	Beckman Coulter	Cat# A63881	
Commercial assay or kit	Chromium Single Cell 3' Reagent Kits v2 Chemistry	10X Genomics		
Software, algorithm	R-3.4.0	other		(R Development Core Team (2019) http://www.R -project.org
Software, algorithm	GraphPad Prism version 8.0 for Mac	GraphPad	**RRID:SCR_00 2798**	GraphPad Software, La Jolla California USA, www.graphp ad.com
Software, algorithm	XCALIBR	other		(https://githu b.com/NKI-GCF/xcalibr)
Software, algorithm	Cellranger v3	10X Genomics	**RRID:SCR_01 7344**	
Software, algorithm	Seurat v3	DOI 10.1016/j.cell .2019.05.031	**RRID:SCR_00 7322**	

374 **Mice**

375 Male C57BL/6J CD45.1+, C57BL/6J CD45.2+, and *Rosa26CreER^{T2}*;*mT/mG* mice from

376 Jackson Laboratory or bred at Institute Curie aged between 7-13 weeks were used in

377 all experiments. All procedures were approved by the responsible national ethics

378 committee (APAFIS#10955-201708171446318 v1).

Barcode library, barcode reference list and lentivirus production

The LG2.2 barcode library is composed of a DNA stretch of 180 bp with a 20 bp "N"-stretch. DsDNA was generated by 10 PCR rounds and cloned into the XhoI-EcoRI site of the lentiviral pRRL-CMV-GFP plasmid[32]. Subsequently ElectroMaxStbl4 cells were transformed, and >10,000 colonies picked for amplification by Maxiprep. To create the barcode reference list (https://github.com/PerieTeam/Eisele-et-al.-), barcode plasmids were PCR amplified twice in duplicate and sequenced as described below. Sequencing results were filtered for barcode reference list generation as previously described in[33]. Lentiviruses were produced by transfecting the barcode plasmids and p8.9-QV and pVSVG into HEK293T cells in DMEM-Glutamax (Gibco) supplemented with 10% FCS (Eurobio), 1% MEM NEAA (Sigma), and 1% sodium pyruvate (Gibco) using Polyethyleneimine (Polysciences). Supernatant was 0,45 um filtered, concentrated by 1,5 h ultracentrifugation at 31,000g and frozen at -80°C. HEK293T were tested for mycoplasma contamination every 6 months.

HSPC and MPP2 isolation, barcoding, EPO treatment, and transplantation

Isolation and labeling of cells with the barcoding library was performed as described in[33]. Briefly, after isofluorane aneasthesia and cervical dislocation, bone marrow cells were isolated from femur, tibia and iliac bones of mice by flushing, and C-Kit⁺ cells were enriched with anti-CD117 magnetic beads on the MACS column system (Miltenyi). Cells were stained for C-Kit, Flt3, CD150, Sca-1 (Key Resources Table) Propidium iodide (PI) (Sigma) (1:5,000), and if appropriate CD48. Lineage staining was not performed after C-Kit+ MACS enrichment for transplantation cohorts. For HSPC cohorts, HSPCs (Figure 1 – figure supplement 1a) were sorted and transduced with the barcode library in StemSpanMedium SFEM (STEMCELL Technologies) with 50

ng/ml mSCF (STEMCELL Technologies) through 1,5 h of centrifugation at 300 g and

4,5 h incubation at 37°C to obtain 10% barcoded cells. After transduction, cells were

incubated with human recombinant EPO (Eprex, erythropoietin alpha, Janssen) at a

final concentration of 1,000 or 160 ng/ml or PBS for 16 h at 37°C. After the incubation,

the cells were transplanted by tail vein injection in recipient mice 6 Gy sub-lethally

irradiated 3 hours before on a CIXD irradiator. Mice were allocated to groups of 4-5

mice for each condition randomly without masking. When indicated, cells were injected

together with additional EPO (133 ug/kg). On average 2,600 cells (Mean 2,684 cells

+/- 175 cells) were injected in the tail vein of each mouse. For the MPP2 cohort, MPP2

and CD48$^-$ HSPCs (Figure 7 – figure supplement 1a) were sorted. Both populations

were cultured alike, but only MPP2 were transduced with the barcode library and

treated with 1,000 ng/ml recombinant EPO as described above. After the culture,

barcoded MPP2 and un-barcoded CD48$^-$ HSPCs were mixed at a ratio of 32/45 (to be

as close as possible to the original ratio of both populations in the HSPCs) and

transplanted as described above. A FACSAria^{TM} (BD Biosciences) was used for

sorting. FACSDiva^{TM} software (BD Biosciences) was used for measurements and

FlowJo^{TM} v.10 (TreeStar) for analysis.

Cell progeny isolation for barcode analysis

Spleens were mashed and both blood and spleen cells were separated based on

Ter119 using a biotinylated anti-Ter119 antibody (Key Resources Table) and anti-

biotinylated beads on the MACS column system (Miltenyi). Ter119$^+$ cells were stained

for Ter119 and CD44[34]. Ter119$^-$ cells were stained for CD45.1 CD11b, CD11c, CD19,

and if appropriate CD115, Siglec-F, and Ly6G (Key Resources Table). Bone marrow

cells were flushed from bones and enriched for C-Kit$^+$ cells as above. When

appropriate the C-Kit$^-$ fraction was further separated based on Ter119 and stained as

above. C-Kit[+] cells were stained for C-Kit, Flt3, CD150, Sca-1, and if appropriate CD41

(Key Resources Table), and PI (1:5000) as described above. For analyzed and/or

sorted populations see Figure 1 – figure supplement 1. Populations were only sorted

for mice with an engraftment (donor cells percentage) of above 5% in spleen, bone

and blood.

Lysis, barcode amplification and sequencing

Sorted cells were lysed in 40 µl Viagen Direct PCR Lysis Reagent (cell) (Euromedex)

supplemented with 0,5 mg/ml Proteinase K Solution RNA grade (Invitrogen) at 55°C

for 120 min, 85°C for 30 min, 95°C for 5 min. Samples were then split into two

replicates, and a three-step nested PCR was performed to amplify barcodes and

prepare for sequencing. The first step amplifies barcodes (top-LIB

(5′TGCTGCCGTCAACTAGA ACA-3′) and bot-LIB (5′GATCTCGAATCAGGCGCTTA-

3′)). A second step adds unique 4 bp plate indices as well as Read 1 and 2 Illumina

sequences (PCR2-Read1-plate-index-forward

5'ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNCTAGAACACTCGAGAT

CAG3′ and PCR2-Read2-reverse

5'GTGACTGGAGTTCAGACGTGTGCTCTTCCGAT

CGATCTCGAATCAGGCGCTTA3′). In a third step, P5 and P7 flow cell attachment

sequences and one of 96 sample indices of 7 bp are added (PCR3-P5-forward

5'AATGATA

CGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT3′

and PCR7-P7-sample-index-reverse

5'CAAGCAGAAGACGGCATACGAGANNNNNNNGTGACTGGAGTTCAGA

CGTGCTCTTCCGATC3′) (PCR program: hot start 5 min 95°C, 15 s at 95°C; 30 s at

57.2°C; 30 s at 72°C, 5 min 72°C, 30 (PCR1-2) or 15 cycles (PCR 3)). Both index

sequences (sample and plate) were designed based on[35] such that sequences differed

by at least 2 bp (https://github.com/PerieTeam/Eisele-et-al.-). To avoid lack of diversity

at the beginning of the reads, at least 4 different plate indices were used for each

sequencing run. Primers were ordered desalted, as high-performance liquid

chromatography (HPLC) purified. During lysis and each PCR, a mock control was

added. The DNA amplification by the three PCRs was monitored by the run on a large

2% Agarose gel. PCR3 products for each sample and replicate were pooled, purified

with the Agencourt AMPure XP system (Beckman Coulter), diluted to 5 nM. and

sequenced on a HiSeq system (Illumina) (SR-65bp) at Institute Curie facility with 10%

of PhiX spike-in.

Barcode sequence analysis

Sequencing results were filtered, and barcodes were categorized in progenitor classes

as in[33] and further explained on github (https://github.com/PerieTeam/Eisele-et-al.-).

In brief, sequencing results were analyzed using R-3.4.0 (R Development Core Team

(2019) http://www.R-project.org.), Excel, and GraphPad Prism version 8.0 for Mac

(GraphPad Software, La Jolla California USA, www.graphpad.com). Reads were first

filtered for perfect match to the input index- and common-sequences using XCALIBR

(https://github.com/NKI-GCF/xcalibr) and filtered against the barcode reference list.

Samples were then filtered for containing at least 5000 reads and normalized to 10^5

per sample. Samples with a Pearson correlation between duplicates below 0.9 were

discarded and barcodes present in one of the two replicates were set to zero. Samples

with under 10 barcodes were filtered out, unless indicated in the figure legend. The

mean of the replicates was used for further processing. When the mean percentage of

barcodes shared between different sequencing runs was higher than within the same

sequencing run for mice of a same transduction batch, reads below the read quartile

of the mean percentage of barcodes shared between mice of a same transduction

batch but sequenced on different sequencing runs were set to zero in order to equalize

the barcode sharing between mice transplanted from a same transduction batch in

different sequencing runs to the barcode sharing between mice within each

sequencing run. After filtering, read counts of each barcode in the different cell lineages

were normalized enabling categorization into classes of biased output toward the

analyzed lineages using a threshold of 10% of barcode reads (Other thresholds in

Figure 1 – figure supplement 3c and Figure 7 – figure supplement 1c). Statistics on

barcoding results were performed using a permutation test as in [36]. Significance of flow

cytometry results was assessed using Student's T test. Some mice were excluded from

the analysis due to death before readout, or due to a donor cell engraftment <5%, as

well as the filtering out of mice for which one or more cell subset samples did not pass

the barcode data filtering steps as detailed above.

ScRNAseq and analysis

ScRNAseq was performed using the 10X Genomics platform on one pool of HSPCs

isolated from 8 mice, barcoded and culture with or without EPO for 16h *in vitro* as

described above. Sequencing libraries were prepared using the Chromium Single Cell

3' v2 kit and sequenced on a HiSeq system (Illumina) at Institut Curie NGS facility.

Data was analyzed using Cellranger v3 (10X Genomics), Seurat v3[37] and customized

scripts. Raw sequencing reads were processed using Cellranger. To obtain a

reads/cell/gene count table, reads were mapped to the mouse GRCm38.84 reference

genome. scRNAseq analysis was performed using Seurat[37]. During filtering, Gm, Rik,

and Rp genes were discarded as non-informative genes. Cells with less than 1,000 genes per cell and with a high percentage of mitochondrial genes were removed from downstream analyses. Following our filtering procedures, the average UMI count per cell was 5157, with mitochondrial genes accounting for 5% of this. The average number of genes detected per cell was 2337. Cell cycle annotation using the cyclone method from the scran R package showed that 2938 cells were in G1 phase, 233 cells were in G2M phase, and 127 cells were in S phase. No batch effect was detected between the EPO and no-EPO group, therefore no batch correction was applied. Data normalization was performed using the default Seurat approach and differentially expressed genes were determined using a logistic regression in Seurat. Unsupervised clustering was performed on the significant variable genes using the ten first principle component analysis followed by the non-linear dimensionality reduction technique UMAP[38] (Figure 5 – figure supplement 2). Unsupervised Louvain clustering of the data was performed across a range of resolution parameters and the resolution value that led to the most stable clustering profiles was chosen[39] (Figure 5 – figure supplement 2). Annotation of the clusters was obtained by mapping published signatures using the *AddModuleScore* method of Seurat. The signatures are defined in the following publications: dHSC, and MPP1 signatures were obtained from[40]. The MolO LT-HSC signature was taken from[41] and the MPP2 and 4 signature was taken from[31]. An excel file listing the genes in these signatures is available on github (https://github.com/PerieTeam/Eisele-et-al.-). To identify EPO-responder cells in the EPO group, differential expression analysis was performed between control and EPO groups (Lists of DEGs are available at https://github.com/PerieTeam/Eisele-et-al.-). Subsequently, genes that were differentially expressed (adjusted p-value< 0.05) between EPO and control groups were transformed into an EPO response signature

which when overlaid onto the UMAP based visualization was enriched only in a subset

of the EPO group cells. Briefly this signature was obtained by taking the background

corrected mean expression values of both the up and downregulated genes per cell

as implemented in the *AddModuleScore* method of Seurat. Within each cell these two

signature scores were used to create a composite EPO response score by subtracting

the downregulated response from the upregulated response signature. Cells in the

upper 90[th] percentile with regards to the expression of the EPO response signature

were labelled EPO responders.

To perform supervised cell type annotation, a reference map was generated from a

published single cell sequencing dataset of 44,802 C-Kit[+] cells from[42]. Preprocessing

was performed using a scanpy pipeline[43]. Data was then visualized using the non-

linear non-dimensionality reduction technique UMAP[38]. Annotation of the reference

map was obtained by overlaying published signatures as above using the

AddModuleScore method of Seurat and also known markers as *Flt3*, *slamf1*, and

Gata1 (Figure 5 – figure supplement 1b). For the erythroid progenitors these markers

are *Gata1, Klf1, Epor, Gypa, Hba-a2, Hba-a1* (figure 5 – figure supplement 1a). Cells

were mapped onto the reference map using a k-nearest neighbors mapping approach.

Briefly, for each cell in the query dataset, the nearest neighbors in PCA space of the

reference dataset were determined using the nn2 function of the RANN package and

the mean UMAP 1 and 2 coordinates of the 10 nearest neighbors were taken as the

reference point for the new cell of interest. To benchmark our mapping approach, cells

from an independent dataset of erythroid progenitors [10] were used without additional

pre-processing (Figure 5 – figure supplement 1a).

Data sharing statement

Raw data are available at zenodo DOI 10.5281/zenodo.5645045**.** All codes to filter

and process raw data, as well as filtered data are available at

https://github.com/PerieTeam/Eisele-et-al.-. Contact author is leila.perie@curie.fr.

Acknowledgements

560 We thank Dr. T. Schumacher for discussion and lentiviral library production and Dr. K

561 Duffy for advices on permutation testing. We thank the Institute Curie flow cytometry,

562 next generation sequencing, and animal facility. We thank Fahima Di Federico from

563 the UMR168 BMBC facility for amplifying the barcode plasmid pool.

564
Disclosure of Conflict of interest

566 The authors declare no competing interests.

References

569
1.   Sender R, Fuchs S, Milo R. Are We Really Vastly Outnumbered? Revisiting the
Ratio of Bacterial to Host Cells in Humans. *Cell*. 2016;164(3):337–40.
2.   Richmond TD, Chohan M, Barber DL. Turning cells red: signal transduction
mediated by erythropoietin. *Trends Cell Biol.* 2005;15(3):146–155.
3.   Walsh G. Biopharmaceutical benchmarks 2014. *Nat. Biotechnol.*
2014;32(10):992–1000.
4.   Koury MJ. Tracking erythroid progenitor cells in times of need and times of
plenty. *Exp. Hematol.* 2016;44(8):653–63.
5.   Cheshier SH, Prohaska SS, Weissman IL. The effect of bleeding on
hematopoietic stem cell cycling and self-renewal. *Stem Cells Dev.*
2007;16(5):707–17.
6.   Shiozawa Y, Jung Y, Ziegler AM, et al. Erythropoietin Couples Hematopoiesis
with Bone Formation. *PLoS One*. 2010;5(5):e10853.
7.   Grover A, Mancini E, Moore S, et al. Erythropoietin guides multipotent
hematopoietic progenitor cells toward an erythroid fate. *J. Exp. Med.*
2014;211(2):181–188.
8.   Giladi A, Paul F, Herzog Y, et al. Single-cell characterization of haematopoietic
progenitors and their trajectories in homeostasis and perturbed
haematopoiesis. *Nat. Cell Biol.* 2018;1.
9.   Yang J, Tanaka Y, Seay M, et al. Single cell transcriptomics reveals
unanticipated features of early hematopoietic precursors. *Nucleic Acids Res.*
2017;45(3):1281–1296.
10.   Tusi BK, Wolock SL, Weinreb C, et al. Population snapshots predict early
haematopoietic and erythroid hierarchies. *Nature*. 2018;555(7694):54–60.

11. Singh RP, Grinenko T, Ramasz B, et al. Hematopoietic Stem Cells but Not
Multipotent Progenitors Drive Erythropoiesis during Chronic Erythroid Stress in
EPO Transgenic Mice. *Stem Cell Reports.* 2018;10(6):1908–1919.
12. Dubart A, Feger F, Lacout C, et al. Murine pluripotent hematopoietic
progenitors constitutively expressing a normal erythropoietin receptor
proliferate in response to erythropoietin without preferential erythroid cell
differentiation. *Mol. Cell. Biol.* 1994;14(7):4834–42.
13. Ma W, Kantarjian H, Zhang K, et al. Significant association between
polymorphism of the erythropoietin gene promoter and myelodysplastic
syndrome. *BMC Med. Genet.* 2010;11(1):163.
14. Weinreb C, Rodriguez-Fraticelli A, Camargo FD, Klein AM. Lineage tracing on
transcriptional landscapes links state to fate during differentiation. *Science.*
2020;
15. Li C, Shi C, Kim J, et al. Erythropoietin Promotes Bone Formation through
EphrinB2/EphB4 Signaling. *J. Dent. Res.* 2015;94(3):455–463.
16. Singbrant S, Russell MR, Jovic T, et al. Erythropoietin couples erythropoiesis,
B-lymphopoiesis, and bone homeostasis within the bone marrow
microenvironment. *Blood.* 2011;117(21):.
17. Ito T, Hamazaki Y, Takaori-Kondo A, Minato N. Bone Marrow Endothelial Cells
Induce Immature and Mature B Cell Egress in Response to Erythropoietin. *Cell
Struct. Funct.* 2017;42(2):149–157.
18. American Society of Hematology S, Alvarez JC, Noguchi CT. Blood. American
Society of Hematology; 2017.
19. Zhang Y, Wang L, Dey S, et al. Erythropoietin action in stress response, tissue
maintenance and metabolism. *Int. J. Mol. Sci.* 2014;15(6):10296–333.
20. Tari K, Atashi A, Kaviani S, et al. Erythropoietin induces production of
hepatocyte growth factor from bone marrow mesenchymal stem cells in vitro.
Biologicals. 2017;45:15–19.
21. Dykstra B, Kent D, Bowie M, et al. Long-Term Propagation of Distinct
Hematopoietic Differentiation Programs In Vivo. *Cell Stem Cell.* 2007;1(2):218–
229.
22. Morita Y, Ema H, Nakauchi H. Heterogeneity and hierarchy within the most
primitive hematopoietic stem cell compartment. *J. Exp. Med.* 2010;
23. Oguro H, Ding L, Morrison SJ. SLAM family markers resolve functionally
distinct subpopulations of hematopoietic stem cells and multipotent
progenitors. *Cell Stem Cell.* 2013;13(1):102–16.
24. Sanjuan-Pla A, Macaulay IC, Jensen CT, et al. Platelet-biased stem cells
reside at the apex of the haematopoietic stem-cell hierarchy. *Nature.*
2013;502(7470):232–6.
25. Carrelha J, Meng Y, Kettyle LM, et al. Hierarchically related lineage-restricted
fates of multipotent haematopoietic stem cells. *Nature.* 2018;554(7690):106–
111.
26. Yamamoto R, Morita Y, Ooehara J, et al. Clonal analysis unveils self-renewing
lineage-restricted progenitors generated directly from hematopoietic stem cells.
Cell. 2013;154(5):1112–1126.
27. Lu R, Neff NF, Quake SR, Weissman IL. Tracking single hematopoietic stem
cells in vivo using high-throughput sequencing in conjunction with viral genetic
barcoding. *Nat. Biotechnol.* 2011;29(10):928–33.
28. Verovskaya E, Broekhuis MJC, Zwart E, et al. Heterogeneity of young and
aged murine hematopoietic stem cells revealed by quantitative clonal analysis

using cellular barcoding. *Blood*. 2013;122(4):523–532.

29. Wilson A, Laurenti E, Oser G, et al. Hematopoietic Stem Cells Reversibly
Switch from Dormancy to Self-Renewal during Homeostasis and Repair. *Cell*.
2008;135(6):1118–1129.

30. Cabezas-Wallscheid N, Klimmeck D, Hansson J, et al. Identification of
regulatory networks in HSCs and their immediate progeny via integrated
proteome, transcriptome, and DNA methylome analysis. *Cell Stem Cell*.
2014;15(4):507–522.

31. Pietras EM, Reynaud D, Kang Y-A, et al. Functionally Distinct Subsets of
Lineage-Biased Multipotent Progenitors Control Blood Production in Normal
and Regenerative Conditions. *Cell Stem Cell*. 2015;17(1):35–46.

32. Dull T, Zufferey R, Kelly M, et al. A Third-Generation Lentivirus Vector with a
Conditional Packaging System. *J. Virol.* 1998;72(11):8463.

33. Naik SH, Perié L, Swart E, et al. Diverse and heritable lineage imprinting of
early haematopoietic progenitors. *Nature*. 2013;496(7444):229–232.

34. Chen K, Liu J, Heck S, et al. Resolving the distinct stages in erythroid
differentiation based on dynamic changes in membrane protein expression
during erythropoiesis. *Proc. Natl. Acad. Sci.* 2009;106(41):17413–17418.

35. Faircloth BC, Glenn TC. Not All Sequence Tags Are Created Equal: Designing
and Validating Sequence Identification Tags Robust to Indels. *PLoS One*.
2012;7(8):e42543.

36. Tak T, Prevedello G, Simon G, et al. Hspcs display within-family homogeneity
in differentiation and proliferation despite population heterogeneity. *Elife*.
2021;10:.

37. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of
single-cell gene expression data. *Nat. Biotechnol.* 2015;33(5):495–502.

38. McInnes L, Healy J, Saul N, Großberger L. UMAP: Uniform Manifold
Approximation and Projection. *J. Open Source Softw.* 2018;

39. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. Fast unfolding of
communities in large networks. *J. Stat. Mech. Theory Exp.* 2008;

40. Cabezas-Wallscheid N, Buettner F, Sommerkamp P, et al. Vitamin A-Retinoic
Acid Signaling Regulates Hematopoietic Stem Cell Dormancy. *Cell*. 2017;

41. Wilson NK, Kent DG, Buettner F, et al. Combined Single-Cell Functional and
Gene Expression Analysis Resolves Heterogeneity within Stem Cell
Populations. *Cell Stem Cell*. 2015;

42. Dahlin JS, Hamey FK, Pijuan-Sala B, et al. A single-cell hematopoietic
landscape resolves 8 lineage trajectories and defects in Kit mutant mice.
Blood. 2018;131(21):e1–e11.

43. Wolf FA, Hamey FK, Plass M, et al. PAGA: graph abstraction reconciles
clustering with trajectory inference through a topology preserving map of single
cells. *Genome Biol.* 2019;

44. Matsuoka Y, Sasaki Y, Nakatsuka R, et al. Low level of C-kit expression marks
deeply quiescent murine hematopoietic stem cells. *Stem Cells*.
2011;29(11):1783–1791.

45. Boyer SW, Rajendiran S, Beaudin AE, et al. Clonal and Quantitative In Vivo
Assessment of Hematopoietic Stem Cell Differentiation Reveals Strong
Erythroid Potential of Multipotent Cells. *Stem Cell Reports*. 2019;12(4):801–
815.

46. Perié L, Duffy KR, Kok L, de Boer RJ, Schumacher TN. The Branching Point in
Erythro-Myeloid Differentiation. *Cell*. 2015;163(7):1655–62.

47.   Tak T, Prevedello G, Simon G, et al. Simultaneous tracking of division and
differentiation from individual hematopoietic stem and progenitor cells reveals
within-family homogeneity despite population heterogeneity. *bioRxiv.*
2019;586354.
48.   Rodriguez-Fraticelli AE, Wolock SL, Weinreb CS, et al. Clonal analysis of
lineage fate in native haematopoiesis. *Nat. Publ. Gr.* 2018;553:.
49.   Theurl I, Hilgendorf I, Nairz M, et al. On-demand erythrocyte disposal and iron
recycling requires transient macrophages in the liver. *Nat. Med.*
2016;22(8):945–51.
50.   Wang J, Hayashi Y, Yokota A, et al. Expansion of EPOR-negative
macrophages besides erythroblasts by elevated EPOR signaling in
erythrocytosis mouse models. *Haematologica.* 2018;103(1):40–50.
51.   Gilboa D, Haim-Ohana Y, Deshet-Unger N, et al. Erythropoietin enhances
Kupffer cell number and activity in the challenged liver. *Sci. Rep.*
2017;7(1):10379.
52.   Kuzmac S, Grcevic D, Sucur A, Ivcevic S, Katavic V. Acute hematopoietic
stress in mice is followed by enhanced osteoclast maturation in the bone
marrow microenvironment. *Exp. Hematol.* 2014;42(11):966–975.
53.   Ulyanova T, Phelps SR, Papayannopoulou T. The macrophage contribution to
stress erythropoiesis: when less is enough. *Blood.* 2016;128(13):1756–65.
54.   Mausberg AK, Meyer Zu Hörste G, Dehmel T, et al. Erythropoietin ameliorates
rat experimental autoimmune neuritis by inducing transforming growth factor-β
in macrophages. *PLoS One.* 2011;6(10):e26280.
55.   Liao C, Prabhu KS, Paulson RF. Monocyte derived macrophages expand the
murine stress erythropoietic niche during the recovery from anemia. *Blood.*
2018;blood-2018-06-856831.
56.   Bretz CA, Divoky V, Prchal J, et al. Erythropoietin Signaling Increases
Choroidal Macrophages and Cytokine Expression, and Exacerbates Choroidal
Neovascularization. *Sci. Rep.* 2018;8(1):2161.
57.   Kim S, Kim N, Presson AP, et al. Dynamics of HSPC repopulation in
nonhuman primates revealed by a decade-long clonal-tracking study. *Cell*
Stem Cell. 2014;
58.   Aiuti A, Biasco L, Scaramuzza S, et al. Lentiviral hematopoietic stem cell gene
therapy in patients with wiskott-aldrich syndrome. *Science (80-.).*
2013;341(6148):.
59.   Starnes LM, Sorrentino A, Pelosi E, et al. NFI-A directs the fate of
hematopoietic progenitors to the erythroid or granulocytic lineage and controls
β-globin and G-CSF receptor expression. *Blood.* 2009;114(9):1753–1763.
60.   Meenhuis A, Van Veelen PA, De Looper H, et al. MiR-17/20/93/106 promote
hematopoietic cell expansion by targeting sequestosome 1-regulated pathways
in mice. *Blood.* 2011;118(4):916–925.
61.   Blank U, Karlsson G, Moody JL, et al. Smad7 promotes self-renewal of
hematopoietic stem cells. *Blood.* 2006;
62.   Steffen B, Knop M, Bergholz U, et al. AML1/ETO induces self-renewal in
hematopoietic progenitor cells via the Groucho-related amino-terminal AES
protein. *Blood.* 2011;117(16):4328–4337.
63.   Wu C, Espinoza DA, Koelle SJ, et al. Geographic clonal tracking in macaques
provides insights into HSPC migration and differentiation. *J. Exp. Med.*
2018;215(1):217–232.
64.   Sun J, Ramos A, Chapman B, et al. Clonal dynamics of native haematopoiesis.

Nature. 2014;514(7522):.

65. Busch K, Klapproth K, Barile M, et al. Fundamental properties of unperturbed
haematopoiesis from stem cells in vivo. *Nature*. 2015;518(7540):542–546.

66. Schoedel KB, Morcos MNF, Zerjatke T, et al. The bulk of the hematopoietic
stem cell population is dispensable for murine steady-state and stress
hematopoiesis. *Blood*. 2016;128(19):2285–2296.

67. Rainville N, Jachimowicz E, Wojchowski DM. Targeting EPO and EPO receptor
pathways in anemia and dysregulated erythropoiesis. *Expert Opin. Ther.*
Targets. 2016;20(3):287–301.

68. Halawi R, Cappellini MD, Taher A. A higher prevalence of hematologic
malignancies in patients with thalassemia: Background and culprits. *Am. J.*
Hematol. 2017;92(5):414–416.

Tables

Table 1: Permutation testing of changes in clonality after transplantation of EPO-

exposed HSPCs. Same data as in Figure 1, 2, 3, 4 and Figure 7. HSPCs or MPP2

were cultured with different concentrations of EPO (160 ng/ml or 1,000 ng/ml) for 16h,

and when indicated a soluble dose of EPO (133 ug/kg) was injected together with

barcoded HSPCs at the moment of transplantation. Barcodes in the erythroid (E),

myeloid (M), B-lymphoid (B) lineage, dendritic cell (DC) and HSPCs, were analyzed

four weeks after transplantation and categorized by bias using a 10% threshold. For

the data of Figure 1, 2 and 7, the output of MB and ME classified barcodes to the B,

M, and E lineages was analyzed using a permutation test. For the data of Figure 3, the

output of MBE and MB classified barcodes to the DC lineage was analyzed. For the

data of Figure 4, the output of barcodes present in HSPCs to the B, M, and E lineages

was analyzed using a permutation test. By permutating the mice of control and EPO

groups, the random distribution of this output was generated and compared to the real

output difference between control and EPO group. A p-value was generated as in[47]

Figure	Condition	p-value			
		MB in B	**MB in M**	**ME in E**	**ME in M**
1	**HSPCs 160 ng/ml**	0.02	0.04	0.02	0.04
1, 2	**HSPCs 1000 ng/ml**	0.0075	0.0071	0.0071	0.012
2	**HSPCs 160 ng/ml + inj.**	0.01	0.016	0.012	0.011
2	**HSPCs 1000 ng/ml + inj.**	0.0018	0.0018	0.002	0.0025
7	**MPP2 1000 ng/ml**	0.006	0.008	0.004	0.004
		MBE in DC	**MB in DC**		
3	**HSPCs 1000 ng/ml**	0.07	0.0075		
		HSPC in B	**HSPC in M**	**HSPC in E**	
4	**HSPCs 160 ng/ml**	0.035	0.029	0.029	
	HSPCs 1000 ng/ml	0.008	0.06	0.01	

Figure Legends

Figure 1: High output ME- and MB-biased clones occur after EPO-exposure and transplantation of HSPCs. **a**, HSPCs were sorted from the bone marrow of donor mice, lentivirally barcoded, cultured *ex vivo* with or without 1,000 ng/ml EPO for 16 h and transplanted into sublethally irradiated mice. At week 4 post-transplantation, the erythroid (E), myeloid (M), and B-cells (B) lineages were sorted from the spleen and processed for barcode analysis. **b**, The percentage of donor derived cells (CD45.1$^+$) among the total spleen, myeloid cells (CD11b$^+$) or B-cells (CD19$^+$) in the spleen of control and EPO group. **c**, To better assess chimerism in erythroid cells mTdTomato/mGFP donor mice were used. The fraction of Tom$^+$ cells among erythroid cells (Ter119$^+$) in the spleen and blood in control and EPO group. **d**, Number of barcodes retrieved in the indicated lineages at week 4 after transplantation in control and EPO group. **e**, Triangle plots showing the relative abundance of barcodes (circles) in the E, M, and B lineage with respect to the summed output over the three lineages (size of the circles) for control and EPO group. **f**, Percentage of HSPCs classified by the indicated lineage bias, using a 10% threshold for categorization. **g**, Quantitative contribution of the classes as in f to each lineage. Shown are values from several animals (n= 8 EPO, n= 10 control in b, n= 3 EPO, n=4 control in c spleen, n= 4 EPO, n=8 control in c blood collected over 5 different experiments (d-g) n=5 for control and n=2 for EPO group collected over one experiment). For all bar graphs mean and S.D. between mice are depicted. Statistical significance tested using Mann-Withney U test p=0,05 for (b-c). Statistical significance tested by permutation test for different subsets in g (see Table 1). This Figure has four supplements.

Figure 2: Effect of different EPO concentrations on HSPC clonality after

transplantation. Same protocol as in Figure 1 but HSPCs were cultured with different

concentrations of EPO (160 ng/ml or 1,000 ng/ml) for 16 h, and when indicated a single

dose of EPO (133 ug/kg) was injected together with barcoded cells at the moment of

transplantation. **a**, Triangle plots showing the relative abundance of barcodes (circles)

in the erythroid (E), myeloid (M), and B-lymphoid (B) lineage with respect to the

summed output over the three lineages (size of the circles) for the different

experimental groups as indicated. **b**, The percentage of each lineage produced by the

barcodes categorized by bias using a 10% threshold. Shown are values from several

animals (n=2 for 160 ng/ml, 1,000 ng/ml, and 160 ng/ml + EPO injection, n=4 for 1,000

ng/ml + EPO injection (collected over 4 different experiments)). For all bar graphs

mean and S.D. between mice are depicted. Statistical significance tested by

permutation test for different subsets in b (see Table 1). This figure has one

supplement.

Figure 3: Production of Dendritic Cells (DC) and Megakaryocyte Progenitors

(MkP) by HSPCs after EPO-exposure and transplantation. In addition to the

analysis of barcodes in the erythroid (E), the myeloid (M), and the B-cell (B) lineage,

the DC lineage in spleen and MkP in bone marrow were added. **a**, Percentage of

barcoded HSPCs producing DC in the different HSPC categories (classification as in

Figure 2 based on the M, E, and B lineage only using a 10% threshold. The DC only

category was added). **b**, The percentage of the DC lineage produced by the barcodes

categorized by bias as in a. **c-d**, Representations as in a-b for barcode detection in

MkP. Data is derived from a cohort with detailed myeloid sorting. The myeloid lineage

was merged according to the percentage of total donor myeloid each subset

contributed as in Figure 2 – supplement 1a to allow classification as in a-b based on

the M, E, and B lineage only using a 10% threshold. The MkP only category was added.

Shown are values from several animals (a-b, n=5 for control and n=2 for EPO group,

c-d, n=3 for control and n=1 for EPO group (collected over two experiments)). For all

bar graphs mean and S.D. between mice are depicted. Statistical significance tested

using Mann-Withney U test p=0,05 for (a,c). Statistical significance tested by

permutation test for different subsets in b (see Table 1). This figure has one

supplement.

Figure 4: Overlap of barcodes in HSPCs and mature cells after transplantation

of EPO-exposed HSPCs. Same protocol as in Figure 1 but HSPCs were cultured with

two different concentrations of EPO (160 ng/ml or 1000 ng/ml) for 16 h. In addition,

HSPCs were sorted and subjected to barcode analysis. **a**, The total number of

barcodes found back in HSPCs. **b**, The percentage of barcodes in the mature cell

subsets also detected in HSPCs and the percentage of barcodes in HSPCs also

detected in mature cells. **c**, The percentage of the HSPC lineage contributed by

barcodes categorized by bias as in Figure 2 based on the M, E, and B lineage using a

10% threshold. **d**, The percentage of each lineage produced by the barcodes color

coded for presence (blue) and absence (grey) in HSPCs. Shown are values from

several animals (n=5 for control, n=2 for EPO 160 ng/ml group and n=3 for EPO 1,000

ng/ml group). For all bar graphs mean and S.D. between mice are depicted. Statistical

significance tested using Mann-Withney U test p=0,05 for (a-b). Statistical significance

tested by permutation test for different subsets in d (see Table 1).

Figure 5: Characterization of EPO-exposed HSPCs by scRNAseq. HSPCs were

sorted, barcoded and cultured *ex vivo* with or without 1,000 ng/ml EPO for 16 h, and

analysed by scRNAseq using the 10X Genomics platform. 1,706 cells from control and

1,595 cells from EPO group passed quality control **a**, Volcano plot of log2 fold change

of the differentially expressed genes between control and EPO-exposed cells versus

the adjusted p-value. Genes of interest are annotated. Differentially expressed genes

were used to defined an EPO response signature. **b**, UMAP visualization of the EPO-

exposed and control HSPCs. **c**, The level of expression in the EPO-exposed HSPCs

of the genes in the EPO response signature (top), and definition of the EPO-responder

and non-responder subgroups using the 90th percentile expression of the EPO

response signature from c (bottom). **d**, The expression of the indicated genes in the

control, EPO-responder and non-responder subgroups as defined in c. Genes that are

significantly upregulated in the EPO-responder group, when compared to control and

non-responder groups. Differential expression was assessed using a logistic

regression testing approach, as implemented in Seurat. This figure has two

supplements and correspond to one 10X experiment of a pool of 8 mice.

Figure 6: EPO-responders are multipotent progenitors, not HSCs. Same protocol

as in Figure 5**. a,** Expression of published gene-signatures of HSCs (dormant HSC[40],

molecular overlap (molO) HSC signature[41]) and MPPs (MPP1[40]-2[31]) across the entire

dataset (see Methods) **b**, Expression of the signatures from 6a, across control, non-

responder, and EPO-responder groups as defined in Figure 5c. Statistical comparisons

made using a Kruskal Wallis Test with a Dunns multiple comparisons post-hoc test **c**,

Expression of the molO HSC signature on the published reference map[42]. **d**, Nearest-

neighbor mapping of control, EPO-responder and non-responder cells onto the

published reference map[42].

Figure 7: MPP2 are enriched for ME-biased clones after EPO-exposure and

transplantation. MPP2 and CD48⁻ HSPCs were sorted from the bone marrow of donor

mice, MPP2 were lentivirally barcoded, and both populations cultured *ex vivo* with or

without 1,000 ng/ml EPO for 16 h. After the culture, barcoded MPP2 and un-barcoded

CD48⁻ HSPCs were mixed and transplanted into sublethally irradiated mice. At week

4 post-transplantation, the erythroid (E), myeloid (M), and B-cells (B) lineages were

sorted from the spleen and processed for barcode analysis. **a**, The fraction of donor

cells among the indicated cell types in spleen. **b**, Barcode number retrieved in the

indicated lineage at 4 weeks after transplantation in control and EPO 1,000 ng/ml

group. **c**, Percentage of MPP2s classified using a threshold of 10% in experimental

groups as indicated. **d**, The percentage of each lineage produced by the MPP2

barcodes categorized by bias using a 10% threshold. **e**, Triangle plots showing the

relative abundance of barcodes (circles) in the erythroid (E), myeloid (M), and B-

lymphoid (B) lineage with respect to the summed output over the three lineages (size

of the circles). Shown are data from several mice (n=3 for control and n=4 for EPO

group). For all bar graphs mean and S.D. between mice are depicted. Statistical

significance tested using Mann-Withney U test p=0,05 for (c-e). Statistical significance

tested by permutation test for different subsets in a (see Table 1). This figure has one

supplement.

Figure 8: The effect of EPO on HSPC clonality after transplantation is transient.

Same protocol as in Figure 1, but barcodes in the E, M, and B lineage in spleen of

individual mice sacrificed at month 4 post-transplantation were analyzed. **a**, Triangle

plots showing the relative abundance of barcodes (circles) in the erythroid (E), myeloid

(M), and B-cell (B) lineage with respect to the summed output over the three lineages

(size of the circles) for the different experimental groups as indicated. **b**, The

percentage of each lineage produced by the barcodes categorized by bias using a

10% threshold. **c**, The fraction of donor cells among the indicated cell types in spleen.

d, Barcode number retrieved in the indicated lineage at month 4 after transplantation

in control, EPO 160 ng/ml, and EPO 1,000 ng/ml group. **e**, Percentage of HSPCs

classified using a threshold of 10% in experimental groups as indicated. Shown are

data from several mice. (c, n=5 for control and n=4 for each EPO group, (a-b, d-e) n=6

for control and n=4 for each EPO group (collected over two experiments)). For all bar

graphs mean and S.D. between mice are depicted. Statistical significance tested using

Mann-Withney U test p=0,05 for (c-e).

Figure supplements legends

Figure 1 – figure supplement 1: Gating strategies and HSPC marker expression

after lentiviral transduction and *ex vivo* culture with or without EPO. **a**, HSPCs

were gated as propidium iodide negative single C-Kit$^+$ Sca-1$^+$ Flt3$^-$ CD150$^+$ cells of C-

Kit$^+$ enriched bone marrow cells. **b**, Erythroblast cells were gated as Ter119$^+$ CD44$^+$

FSC^{hi} cells on Ter$^+$ enriched cells. **c**, Gating strategy for B-cells (CD19$^+$ CD11b$^-$),

dendritic cells (CD19$^-$ CD11b$^-$ CD11c$^+$), and myeloid cells (CD119$^-$ CD11c$^-$ CD11b$^+$) on

Ter119$^-$ live single donor cells. **d**, Gating for MkP (C-Kit$^+$ Sca-1$^-$ CD150$^+$ CD41$^+$) from

C-Kit$^+$ enriched bone marrow cells. **e**, Sort gating for GFP$^+$ erythroid, myeloid, B-, and

dendritic cells respectively used for barcoding analysis. **f**, Representative flow

cytometry plots of sorted HSPC pool after 6 h lentiviral transduction and 16 h *ex vivo*

incubation with or without EPO.

Figure 1 – figure supplement 2: Correlations in barcoding profiles of spleen and

bone. HSPCs were sorted from the bone marrow of donor mice, lentivirally barcoded,

cultured *ex vivo* for 16 h and transplanted into sublethally irradiated mice. At week 4

post-transplantation, erythroid (E), B-cells (B), and the myeloid lineage (M) cells

monocytes, eosinophils, neutrophils, and macrophages were sorted from the spleen

and from bone and processed for barcode analysis. The myeloid lineage was merged

according to the percentage of total donor myeloid each subset contributed as in

Supplementary Figure 4a. **a**, Heatmaps showing the output of individual barcodes

(rows) in different samples (columns) as indicated. Data is normalized by cell subset,

log transformed, and clustered by complete linkage using Euclidean distance. No

output is represented in black. **b**, The percentage of barcodes in spleen and bone

detected in the respective other organ. The spearman rank correlation of barcodes in

bone and spleen was for the B-, M- and E-lineage 0.81, 0,69 and 0,7 respectively.

Shown are values from several animals (n= 3). For all bar graphs mean and S.D.

between mice are depicted. Statistical significance tested using Mann-Withney U test

p=0,05 for (b).

Figure 1 – figure supplement 3: Characterization of lineage biases after

transplantation of EPO-exposed HSPCs. **a**, Triangle plots from Figure 1e color

coded by mice. **b**, Quantitative contribution of the classes to each lineage as in Figure

1 using different thresholds of 0%, 5%, 10%, 15% and 20%. **c-f**, Data for an additional

experiment as in Figure 1. **c**, Number of barcodes retrieved in the indicated lineages

at week 4 after transplantation in control and EPO group. **d**, Triangle plots showing the

relative abundance of barcodes (circles) in the erythroid (E), myeloid (M), and B-cell

(B) lineage with respect to the summed output over the three lineages (size of the

circles) for control and EPO group. **e**, Proportion of HSPCs classified in the indicated

lineage bias category, using a 10% classification threshold. **f**, Quantitative contribution

of the classes as in f to each lineage. Shown are values from several animals (a-c, n=5

for control group and n=2 for EPO group (collected over one experiment), d-f, n= 2 for

control, n= 3 for EPO group (collected over one experiment)). For all bar graphs mean

and S.D. between mice are depicted. Statistical significance tested using Mann-

Withney U test p=0,05 for (c, e).

Figure 1 – figure supplement 4: High output ME- and MB-biased HSPCs occur

six weeks after transplantation of EPO-exposed HSPCs. HSPCs were sorted from

the bone marrow of donor mice, lentivirally barcoded, cultured *ex vivo* with or without

1,000 ng/ml EPO for 16 h and transplanted into sublethally irradiated mice. At week 6

post-transplantation, the erythroid (E), myeloid (M), and B-cells (B) lineages were

sorted from the spleen and processed for barcode analysis. Quantitative contribution

of HSPCs classified by the indicated lineage bias, using a 10% threshold for

categorization, to each lineage. Shown are values from several animals (n= 2 EPO, n=

4 control). Mean and S.D. between mice are depicted.

Figure 2 – figure supplement 1: Variability in the effect of different EPO

concentrations on clonality after HSPC transplantation at different timepoints. a,

Triangle plots showing the relative abundance of barcodes (circles) in the erythroid (E),

myeloid (M), and B-cell (B) lineage with respect to the summed output over the three

lineages (size of circles) for control and EPO groups of Figure 2, color coded by mice.

b, Same representation as in a for data of Figure 8.

Figure 3 – figure supplement 1: Production of macrophages, monocytes,

neutrophils, eosinophils, and Megakaryocyte Progenitors (MkP) by HSPCs after

EPO-exposure and transplantation. Same experimental protocol as in Figure 1 but

the myeloid cells were subdivided into monocytes, eosinophils, macrophages and

neutrophils, and MkP were sorted. **a**, Gating for detailed myeloid subsets on myeloid

cells. Monocytes (Mo) were sorted as CD115$^+$ cells, eosinophils (Eo) as CD115$^-$

SiglecF$^+$ Ly6G$^+$, macrophages (Ma) as CD115$^-$ SiglecF$^-$ Ly6G$^-$, and neutrophils (Neu)

as CD115$^-$ SiglecF$^-$ Ly6G$^+$ cells. **b**, The contribution of different cell types to the overall

donor myeloid subset in control and EPO group. **c**, Heatmaps showing the output of

individual barcodes (rows) in different samples (columns) as indicated. Data is

normalized by cell subset, log transformed, and clustered by complete linkage using

Euclidean distance. No output is represented in black. Shown are values from several

animals (b, n=5 for control and n= 3 for EPO group c, n=3 for control and n=2 for EPO

group (collected over two experiments)). For all bar graphs mean and S.D. between

mice are depicted. Statistical significance tested using Mann-Withney U test p=0,05

for (b).

Figure 5 – figure supplement 1: ScRNAseq analysis of control and EPO-exposed

HSPCs a, UMAP projection of the scRNAseq dataset from[42] annotated with *flt3*,

CD150, and *gata1* gene expression. **b**, Projecting of erythroid-biased progenitors

from[10] on UMAP projection of a. **c**, Robustness of the UMAP visualization and

unsupervised clustering of the data in Figure 4c .The amount of variance explained by

each principle component (left) and UMAP-based visualization using 10 PCA (right)

for different number of genes. **d**, The expression of genes encoding known EPO

receptors in each subgroup. **e,** Overview of the reference map using supervised cell

type annotation of the dataset from[42]. On the right hand side we overlay the MPP4

signature defined by[31] onto our reference map to facilitate celltype annotation **f**, HSPCs

were sorted, barcoded and cultured *ex vivo* with or without 1,000 ng/ml EPO for 16 h,

and analysed by scRNAseq using the 10X Genomics platform. Mapping of the

transcriptomes of the 1,706 cells from control and 1,595 cells from EPO group obtained

after quality control onto the reference map using a k-nearest neighbors mapping

approach.

1004

Figure 5 – figure supplement 2: Unsupervised clustering of control and EPO-

exposed HSPCs a, Cluster stability analysis varying the resolution parameter of the

Seurat clustering method. The significant variable genes using 10 PCA were used as

input. **b**, UMAP visualization of the data in a using a clustering resolution of 0.1. with

the proportion of each cluster as in the control, EPO-responder and non-responder

subgroups. **c**, expression of published signatures of established celltypes used to

annotate our clusters. All comparisons in signature expression between clusters were

statistically significant (adjusted p-value < 0.05) as determined by a Kruskal-Wallis

Test and Dunns Post-Hoc analysis. **d**, Nearest neighbor mapping of unsupervised

clusters from b onto the reference map.

1015

Figure 7 – figure supplement 1: Characterization of lineage biases after

transplantation of EPO-exposed MPP2. **a**, MPP2s and CD48$^-$ HSPCs were gated as

propidium iodide negative single C-Kit$^+$ Sca-1$^+$ Flt3$^-$ CD150$^+$ CD48$^+$ (MPP2) and CD48$^-$

(CD48⁻ HSPCs) cells of C-Kit⁺ enriched bone marrow cells. **b**, Triangle plots from

Figure 7b color coded by mice. **c**, Quantitative contribution of the classes to each

lineage as in Figure 7a using different thresholds of 0%, 5%, 10%, 15% and 20%. (a-

c, n=5 for control group and n=2 for EPO group (collected over one experiment).

Shown are values from several animals (n= 3 for control, n= 4 for EPO group (collected

over one experiment)). For all bar graphs mean and S.D. between mice are depicted.

1025

1026 The article has one Supplementary File 1**.**

1027

1028 **Supplementary File legends:**

1029

Supplementary File 1. Permutation testing of changes in clonality after transplantation

of EPO-exposed HSPCs**.** Same data as in Figure 1 figure supplement 4 and Figure 1

figure supplement 3. HSPCs were cultured with EPO (1,000 ng/ml) for 16h. Barcodes

in the erythroid (E), myeloid (M), B-lymphoid (B) lineage, dendritic cell (DC) and

HSPCs, were analyzed four weeks after transplantation and categorized by bias using

a 10% threshold. The output of MB and ME classified barcodes to the B, M, and E

lineages was analyzed using a permutation test. By permutating the mice of control

and EPO groups, the random distribution of this output was generated and compared

to the real output difference between control and EPO group. A p-value was generated

by permutation testing.

1040

a

Gated on Myeloid cells

CD115 / CD11b

Monocyte

Siglec-F / Ly6G

Eosinophil

Neutrophil

Macrophage

b

% of donor myeloid

● No EPO
■ EPO

ns ns ns ns

Eo   Neu   Ma   Mo

c

No EPO

barcodes

Eo   Neu   Ma   Mo   MkP

EPO

barcodes

Eo   Neu   Ma   Mo   MkP

Abundance

0 2 4 6 8 10

a Gated on PI^neg single C-Kit⁺ Sca-1⁺ cells (LSK)

Supplementary File 1: Permutation testing of changes in clonality after transplantation of EPO-exposed HSPCs. Same data as in Figure 1 figure supplement 4 and Figure 1 figure supplement 3. HSPCs were cultured with EPO (1,000 ng/ml) for 16h. Barcodes in the erythroid (E), myeloid (M), B-lymphoid (B) lineage, dendritic cell (DC) and HSPCs, were analyzed four weeks after transplantation and categorized by bias using a 10% threshold. The output of MB and ME classified barcodes to the B, M, and E lineages was analyzed using a permutation test. By permutating the mice of control and EPO groups, the random distribution of this output was generated and compared to the real output difference between control and EPO group. A p-value was generated using permutation testing.

Figure 1 figure supplement	Condition	p-value			
		MB in B	MB in M	ME in E	ME in M
3	HSPCs 1000 ng/ml repeat	0.01	0.0083	0.0087	0.0087
4	HSPCs 1000 ng/ml - 6 weeks timepoint	0.01	0.01	0.0138	0.0185