Achromatic body markings as a widespread visual mechanism across group living
animals

[1]Negro, Juan J.; [2,3]Doña, Jorge; [4]Blázquez, M. del Carmen; [1,5]Rodríguez, Airam; [6]Herbert-
Read, James E., [6]Brooke, M. de L.

1. Estación Biológica de Doñana-CSIC. Avda. Americo Vespucio 26. 41092 Sevilla, Spain

2. Illinois Natural History Survey, Prairie Research Institute, University of Illinois at
Urbana-Champaign, 1816 S. Oak St., Champaign, IL 61820, USA

3. Departamento de Biología Animal, Universidad de Granada, Granada, Spain

4. Centro de Investigaciones Biológicas del Noroeste (CIBNOR) 23096 La Paz, B.C.S.,
México

5. Grupo de Ornitología e Historia Natural de las Islas Canarias, GOHNIC, Canarias, Spain

6. Department of Zoology, University of Cambridge, Downing St, Cambridge, CB2 3EJ,
UK

Corresponding author: Juan J. Negro (negro@ebd.csic.es)

Abstract

Grouping is a widespread form of predator defense, with individuals in groups often
performing evasive collective movements in response to predators' attacks. Individuals
in these groups use behavioral rules to coordinate their movements, with visual cues
about neighbors' positions and orientations informing movement decisions. Although
the exact visual cues individuals use to coordinate their movements with neighbors have
not yet been decoded, some studies have suggested that stripes, lines or other body
patterns may act as conspicuous conveyors of movement information that could
promote coordinated group movement, or promote dazzle camouflage, thereby
confusing predators. We used phylogenetic logistic regressions to test whether the
contrasting achromatic body patterns of four different taxa vulnerable to predation,
including species within two orders of birds (Anseriformes and Charadriiformes), a
suborder of Artiodactyla (the ruminants) and several orders of marine fish
(predominantly Perciformes) were associated with group living. Contrasting body
patterns were significantly more prevalent in social species, and tended to be absent in
solitary species or species less vulnerable to predation. We suggest that body patterns
taking the form of white lines on dark backgrounds, or *vice versa,* provide a widespread
mechanism across taxa that serves either to inform conspecifics of neighbors'
directional movement, or to confuse predators, when moving in groups. Detection and
processing of patterns and of motion in the visual channel is essentially colourblind. That

diverse animal taxa with widely different vision systems (including di-, tri- and tetrachromats) appear to have converged on a similar use of achromatic patterns is therefore expected given signal-detection theory. This hypothesis would explain the convergent evolution of conspicuous coloration patterns as an antipredator mechanism in numerous vertebrate species.

Keywords: stripes, melanin, body patterns, collective movement, antipredator defenses, dazzle

Introduction

Group living characterizes the lives of many animals including flocking birds, schooling fish and ungulate herds. Grouping is often a defensive strategy to reduce the likelihood of individuals being attacked through detection, dilution, confusion, or self-herd effects (Hamilton 1971, Foster and Treherne 1981, Lima 1995, Jeschke and Tollrian 2007). Because isolated individuals are more likely to be targeted by predators (Ioannou 2012, Hogan et al. 2017, Cuthill 2019, Romenskyy et al. 2020), individuals in groups often move to maintain cohesion by escaping from predators in the same direction as other group members (Handegard et al. 2012, Herbert-Read et al. 2015). Many models aimed at explaining such coordinate movements assume animals rely primarily on visual information to infer neighbors' movements and positions (Pita et al. 2016). These assumptions seem appropriate considering that birds are highly visually-oriented animals (Martin 2017, Camacho et al. 2019), and many fishes and mammals similarly rely on vision when interacting with conspecifics and their surroundings over intermediate distances (Guthrie and Muntz 1993).

Because vision appears essential for many species relying on coordinated and cohesive escape from predators, particular visual cues may be used by social animals to promote coordination. Numerous grouping species display conspicuous marks or patterns on their bodies (Figure 1). In mammals and fishes, the majority of which have elongated bodies parallel to the ground, lines often run in parallel to the longitudinal axis of their bodies. In birds, contrasting lines are often displayed in open wings, and markings are often achromatic in design: white lines appear on an otherwise darker (i.e., melanized) plumage. Brooke (1998) found that flocking shorebirds tended to have white wing marks, while solitary species did not, and suggested that these white marks could provide a conspicuous flash to other group members promoting cohesive take-off, or could aid in coordinated movement. Indeed, the escape waves of birds such as dunlins, *Calidris alpina*, where flocks swiftly change color when seen from a distance because of the difference in dorsal –brown- and ventral –white- coloration in the species suggest such visual cues may act to increase coordination within these and other groups (Hemelrijk et al. 2015). In such species, a role of sexual selection as a driver of such body patterns may be dismissed, as these marks are shown year-round by both males and females in otherwise sexually dichromatic species that display both breeding and non-breeding plumages (Brooke, 1998). The presence of stripes in other social taxa, in

particular for the contrasting bands on the bodies of some social mammals, such as
gazelles (Caro and Stankowich 2010), and stripes in different species of fishes (Denton
and Rowe 1998, Seehausen et al. 1999, Kelley et al. 2013), has similarly been debated
but their function remains unclear.

Others have suggested that such body markings, when combined in a mass
moving group, may produce "dazzle" patterns that could reduce either the likelihood of
predators attacking or their attack success rates (Stevens et al. 2008, Hogan et al. 2017,
Cuthill 2019). Contrasting stripes or lines have been hypothesized to provide camouflage
or to function to startle predators when they are suddenly displayed by fleeing prey
(Cuthill 2019 and references therein). Dazzle patterns may make it difficult for a
predator to judge speed and direction of movement of their prey (Hughes et al. 2014).
Although there are no empirical data using or observing wild animals, computer
simulations suggest that striped targets are among the most difficult to capture (Hughes
et al. 2014). Nonetheless, simpler patterns, such as all-grey colorations, are equally
difficult to catch, and thus evidence that stripes play an adaptive role in dazzle
patterning remains inconclusive.

Expanding on Brooke's (1998) original idea for shorebirds, here we separately
analyze coloration data on shorebirds, waterfowl, ruminants and fish, while controlling
for phylogeny, to assess whether social species are more prone to display lateral
markings on their bodies and investigate their possible function. We posit that social
species may use contrasting stripes or bands on their bodies to aid in coordinated
movement or dazzle predators during coordinated movement. According to these
hypotheses, we predicted that for any given taxa, contrasting stripes should be more
prevalent in group-living species than in solitary ones. Here we test this prediction in
species that rely on group cohesion when fleeing from an attacking predator including
herding ungulates, flocking birds and schooling fish.

Material and Methods

Study species

We selected three different vertebrate taxa: two orders of birds (shorebirds
within – Charadriiformes, and waterfowl, – Anseriformes), one suborder of mammals
(Ruminantia), and fishes in the community of the Eastern Tropical Pacific Ocean with
species in the Orders: Anguilliformes, Clupeiformes, Beloniformes, Perciformes,
Scombriformes and Istiophoriformes, all of which include species which have
contrasting white lines on dark bodies (Figure 1), as well as other species with no such
patterns. The chosen taxa included both solitary and social species that often forage or
rest in groups. If approached by a predator, or directly attacked, individuals that detect
the predator increase their speed during escape movements, alerting others of the
predator's attack (Ridley et al. 2010, Ridley et al. 2013).

Charadriiformes

We used the same dataset and the species' ecological attributes previously used
by Brooke (1998) that assessed the occurrence of flash marks in shorebirds (excluding
gulls, auks and allies). Our study included 205 species in this order, although Brooke's
previous study included 210 (we excluded five species because of recent taxonomic
changes). Brooke investigated several ecological variables, such as migration, habitat,
feeding technique and flocking behavior and found that only flocking behavior was
related to the presence of white marks on the wings, rumps and tail. For our analysis,
we included (a) whether the species was mainly migratory or sedentary, as the marks
might help in group forming during migration trips, (b) whether it tended to form flocks
or was mainly solitary, and (c) wing chord length as a proxy for overall size. Our response
variable was presence/absence of white lines on the wings (1 present, 0: absent).

Anseriformes

To capture the difference in wing patch expression that we knew *a priori*
distinguished the smaller ducks from the larger geese and swans (see Figuerola and
Green 2000) we included 148 representative members of all families of waterfowl (this
order comprises 180 species in total). We used a binomial response variable indicating
the presence or absence of white patches on the wings following Hegyi et al. (2008). In
particular, we unified their scores of 1, 2, and 3 into a single score, i.e. 1 = presence of
white patch in the wings, 0 = absence of white patch. For species of geese (not included
in Hegyi et al's (2008) data set), the presence of white wing patch was scored from flying
birds displayed in the identification guide of Madge and Burn (1987), and following a
similar procedure to Hegyi et al. (2008). For species that are predominantly white, like
some swans, dark patches on the wings were scored as presence of patches.

In contrast to the other taxa included in this study, a majority of the species of
Anseriformes are highly social during at least part of their annual cycle. Therefore, for
this Order, we could not distinguish between solitary or social species. As explanatory
variable, therefore we used body mass as a proxy for body size, and a binomial variable
indicating if the species is migratory or resident (again to assess whether marks might
assist group formation during migrations). We predicted that wing marks would be
associated with smaller species (ducks and teals) that are hunted by falcons, whereas
these lines should be largely absent in the larger birds (geese and swans). Given the
sexual dimorphism in many Anseriformes and the absence of sexual differences in wing
patch expression (Hegyi et al. 2008), we selected male body mass to increase variation
(female body mass produced similar results – not shown). Body mass data were taken
from Figuerola and Green (2000). The migratory status of the species was taken from
the distribution maps of the identification guide by Madge and Burn (1987). For partially-
migratory species, we classified the species as migratory if the breeding range
abandoned in winter was larger than the resident breeding area (present all year
around).

Ruminantia

Many species within the Artiodactyla, to which Ruminantia belong, have linear contrasting color patches on their flanks and dorsum, as well as other markings in legs, rump and faces. Caro and Stankowich (2010) previously characterized the presence and absence of contrasting marking in different body parts in 198 species of artiodactyls and investigated their function. Here, we built our own dataset to assess whether the longitudinal bands on the flanks of Ruminantia are related to their degree of sociality. Similar to above, we recorded the presence or absence of longitudinal markings on the bodies of animals as binary response variable, given the definition of a longitudinal marking by Caro and Stankowich (2010) as a band in both flanks contrasting in coloration with the belly below and the dorsum. Social species were defined as the ones forming herds including more than one parent-offspring unit or more than three unrelated individuals. Conversely, solitary species are defined as the ones in which individuals tend to stay alone or just in parent-offspring units. Smaller markings on other body parts (i.e. legs, neck or head) were not considered in the present study, although they may serve a similar function. Body size data were obtained from Smith et al. (2003).

Fishes

We selected representative species of the main Class of fishes, the Actinopterygii. Among them, we chose species from the fish community of the Eastern Tropical Pacific Ocean, following the guide by Allen and Robertson (1998) and the online-actualized version https://biogeodb.stri.si.edu/sftep/en/pages. Because many tropical species of fish can be colorful, we chose species with several patterns, including species that were both banded and un-banded on the sides of their body. Our searching criteria was as follows: First, and following the phylogenetic order of the book, we identified species with some form of stripe, either parallel to the longer axis – horizontal stripes, or down the short axis – vertical stripes. After this initial selection, we went through our database, and for each species with vertical or horizontal stripes, randomly identified another species within the same genus, or genus within the same family, that did not have either of those stripes (plain species in Kelley et al. 2013). In our final database of fishes, most of them (90%) belong to the Perciformes, but with some representatives of additional Orders (Anguilliformes, Clupeiformes, Beloniformes, Scombriformes and Istiophoriformes). Finally, and to align coherently with the other taxa in this study, we only retained primary prey species (defined as species that remain small enough to be eaten by larger fish throughout their lives), and recorded as binary response variables presence/absence of one or more bars or stripes (any color), and their sociality level (i.e., solitary versus schooling).

Phylogenetic analysis

We tested whether body markings (stripes, lines or other markings on the body)
correlated with the evolution of group living, or other attributes such as body mass or
ecological characteristics, using phylogenetic logistic regression (Paradis et al. 2004; Ives
& Garland 2010, 2014; Ives et al. 2011; Ho & Ané, 2014; R Core Team 2019). In
comparative analyses of trait evolution, both "complete" trees (i.e., those that include
unsampled species using polytomy solvers) and "sequenced species only" trees have
their advantages and disadvantages (Upham et al. 2019). First, birth-death polytomy
resolvers used to infer "complete trees" place species onto trees randomly and
independently from the trait's values, thus potentially breaking down natural patterns
of trait phylogenetic structure (Rabosky 2015). Similarly, "sequenced species only" trees,
while not suffering from this problem, contain fewer species that are potentially non-
randomly sampled, risking excluding their trait values from analyses (Upham et al.,
2019). Accordingly, here we followed Upham et al.'s (2019) recommendations and
compared analyses run on samples from both complete and sequences only trees (see
supplementary material).

Overall, we downloaded phylogenetic subsets from broad phylogenetic
syntheses and then summarized those trees by computing single 50% majority-rule
consensus trees using SumTrees v 4.4.0 in DendroPy v4.4.0. (Sukumaran & Holder, 2010,
2015) following Rubolini et al. (2015). For the two bird taxa, we downloaded
phylogenetic subsets from the BirdTree (Jetz et al. 2012, http://birdtree.org). In
particular, we download phylogenetic trees from the Hackett tree distributions (1,000
trees from "All species" + 1,000 trees from "Sequenced species" distributions). For the
Ruminantia taxa, we used the species-level trees of extant Mammalia of Upham et al.
(2019) (1,000 trees from "birth-death node-dated completed trees" + 1,000 trees from
"birth-death node-dated DNA-only" distributions). Lastly, for fish taxa, we downloaded
data from the fish tree of life (100 trees from "all-taxon assembled" distributions plus
the "DNA-only" time-calibrated tree; Rabosky et al. 2018). The final consensuses
"complete trees" were composed of 205 Charadriiformes, 148 Anseriformes, 197
Ruminantia, and 159 fish species, and the final consensuses "sequenced species only"
trees were composed by for 144 Charadriiformes, 137 Anseriformes, 182 Ruminantia,
and 120 fish species.

To assess whether line markings were associated with group-living behavior, we
used phylogenetic logistic regression analyses with the function *phyloglm*
(logistic_MPLE method, 1,000 independent bootstrap replicates, and p-values
computed using Wald tests) in the phylolm R package v.2.6 (Paradis et al. 2004; Ives &
Garland 2010, 2014; Ives et al. 2011; Ho & Ané 2014; R Core Team 2019). We evaluated
model fit with binned residual plots –specifically, whether standard error bands from
these plots contain 95% of the binned residuals– (*binnedplot* function from the arm
v.1.10-1 R package; Gelman & Su 2018) and computing an R^2 statistic suitable for
statistical models with correlated errors, such as the phylogenetic logistic regression

models used here (*R2* function from the rr2 R package; Ives 2019). Then, we used the
fitted coefficients from the phyloglm models to plot the phylogenetic logistic regression
curves using the *plogis* function of the R package "stats."

Results

We found significant associations between body markings and the level of
sociality in all taxa that we investigated, and these results were consistent across
complete and sequenced only trees (see supplementary material). Flocking
Charadriiformes (shorebirds) were more likely to have body markings than non-flocking
species (phyloglm: R^2_{lik} = 0.26, *P* = 0.001; Table 1, Fig 2A). Migratory status or body size
(wing-chord length), however, were not significantly related to the presence or absence
of wing markings (phyloglm: migratory status, *P* = 0.599; body size, *P* = 0.121; Table 1).
In Anseriformes (waterfowl), most species of ducks had white wing stripes on dark
backgrounds, whereas most of the geese did not (see complete tree in Fig. 2B, and DNA-
only tree in Fig. 2BS). In these species, body mass was significantly correlated to the
presence or absence of stripes, with smaller species more likely to have stripes than
larger ones (phyloglm: R^2_{lik} = 0.18, *P* = 0.001; Table 1, Fig. 3B). Similar to the
Charadriiformes, migratory status was not related to the presence or absence of wing
stripes (phyloglm: R^2_{lik} = 0.18, *P* = 0.346; Table 1). A model excluding the three seemingly
outlier data points (i.e., species with body mass > 10,000 g) revealed consistent results
(phyloglm: R^2_{lik} = 0.12; body mass, *P* = 0.003; migratory status, *P* = 0.337).

Social species of ruminants were also more likely to have longitudinal stripes on
the flanks than non-social species (phyloglm: R^2_{lik} = 0.27; *P* = 0.003; Table 1, Fig. 3C). Body
mass, however, was not related to the presence or absence of stripes (phyloglm: body
mass, *P* = 0.133; Table 1).

In fishes, sociality was associated with the presence or absence of horizontal
stripes on the body, with social species more likely to have those body markings
(phyloglm: R^2_{lik} = 0.37; *P* < 0.001; Table 1, Fig. 3D). Vertical lines, however, were
associated in the opposite direction to social species (i.e., social species were less likely
to have vertical lines; phyloglm: vertical lines, R^2_{lik} = 0.16; *P* = 0.009; Table 1).

Discussion

Stripes, lines and markings on the plumage, fur and scales of prey species were
strongly associated with group living across taxa. While different roles for the
contrasting marks on animals' bodies had previously been suggested in independent
studies of shorebirds, waterfowl, ruminants and fishes (Brooke 1998, Hegyi et al. 2003,
Caro and Stankowich 2010, Kelley et al. 2013, Seehausen et al. 1999), evidence for their
importance in social behavior remained inconclusive. In the all-social waterfowl, wing

marks were related to the species' body mass; a strong predictor of predation risk
(Cresswell, 1994, Pomeroy 2006). Given this, and given predation is a strong driver of
group living, we suggest these contrasting lines play an important role in social anti-
predator defenses of at least three different vertebrate classes; mammals, birds and
fishes.

The finding that markings on the body are associated with social behavior is even
more striking considering the visual systems of these taxa are known to vary widely.
However, the combination of adjacent achromatic lines or bands – mostly black/dark
and white (except in some fishes where structural colors also play a role), suggests that
color perception is not necessary for their role. Indeed, detection of patterns and of
motion is achieved by achromatic mechanisms, and thus is essentially colorblind (Osorio
et al. 1999). These different taxa, therefore, appear to have converged on a similar use
of achromatic patterns, as would be expected given signal-detection-theory. Indeed,
white bands with no pigment combined with melanized areas may be generated by
virtually any vertebrate (McGraw 2006). Hence, these markings may be just as efficiently
detected in either di-, tri-, or tetrachromatic species (Bennet and Thery 2007, Bowmaker
1998, Marshall 2000).

While the association between stripes and social behavior appears widespread,
these markings' functional role remains unresolved. Two primary (non-exclusive)
functions have been proposed for these stripes. First, they could be used by conspecifics
to inform neighbors of rapid movement and aid in coordinated movement (Brooke
1998), or second, to reduce predation success through flash, dazzle, or confusion effects
(Stevens et al. 2008, Cuthill 2019). In many cases of birds (including the Charadriiformes
and Anseriformes studied here) wing markings are often only exposed when a bird takes
flight, but remain hidden when the bird folds its wings, with cryptic dorsal plumages
intended for background matching (Cuthill 2019). This sudden display of contrasting
lines has led many authors to consider them primarily as flash or dazzle marks destined
to confuse an approaching predator that might misjudge the speed or direction of the
prey's escape (Brooke 1998). While such an effect could also be achieved by a single
individual exposing its wings during its escape flight, the flashes from multiple
individuals would likely produce a stronger stimulus, and could hence be used by social
animals to reduce per capita predation risk. Indeed, because predators prioritize
distance estimation, their perception of horizontal contours may be disrupted by such
movements, viewing such collectively moving prey as blurred images (Banks et al. 2015).
While such marking may confuse predators from long-range, at short-range on the other
hand, these markings may indicate to other group members the presence of a predator,
and hence achieve more coordinated take-offs (Brooke 1998).

In the Ruminantia, longitudinal (horizontal) stripes were more likely to be
associated with species that formed groups, irrespective of their body size, and
horizontal lines in fishes were associated with shoaling. The roles of lines in both these
taxa has previously been debated. In cichlid fishes, open-water species that engage in
shoaling behavior tend to have longitudinal stripes (Seehausen et al. 1999), but vertical

or horizontal stripes in butterflyfishes are not associated with social behavior, while
diagonal stripes are negatively correlated with fish's shoaling tendencies (Kelley et al.
2013). Our work represents definitive evidence that across a wide phylogenetic tree,
horizontal stripes indeed appear important in social behavior. In contrast to the wing
markings of birds that are only exposed when birds fly, longitudinal lines of mammals
and fish are generally permanently visible. What roles do these markings play, therefore,
in social behavior? Caro and Stankowich (2010) suggested several functions for the
stripes in Ruminantia, including pursuit deterrence, intraspecific signaling and crypsis. In
fishes, some authors have suggested that stripes may disrupt the body shape making it
unrecognizable for predators (Armbruster and Page, 1996) or that stripes may make it
difficult for a predator to focus on a specific individual target (the confusion effect,
McRobert and Brander 1998). Once again, the association of grouping with these lines
suggests that both coordination and confusion effects may play an important role for
these species. Indeed, Denton and Rowe (1998) suggested that stripes in the mackerel
Scomber scombrus help to coordinate shoaling behavior because the way stripes are
perceived changes with body orientation (although they referred to the vertical zebra
stripes of the upper part of the fish body). Further studies on the functional role of
horizontal and vertical stripes are clearly warranted.

Prerequisites for both confusion or coordination hypotheses are that species
should be diurnal, and live in relatively open habitat in order to make the signals visible.
These assumptions appear to be supported in the Anseriformes and Charadriiformes
that often live on open water, marshlands and coastlines, and in the species of
ruminants studied which tend to graze on open grasslands (Caro and Stankowich 2010).
Indeed, even though many mammals are nocturnal, Stoner et al. (2003) noted that
artiodactyls with contrasting lines on their flanks were diurnal and lived in open habitats.
The fishes in our sample similarly live in illuminated shallow waters. And while our result
confirms many social species have stripes and contrasting marks, there are species
which form large moving groups in open environments without any conspicuous
contrasting markings. Indeed, the European starling *Sturnus vulgaris*, that does not
display visually contrasting patterns in the plumage (Svensson et al. 1999), yet displays
some of the most coordinated and collective behaviour in the animal kingdom (Ballerini
et al. 2008a, 2008b, Cavagna et al. 2008a, 2008b, 2010). Clearly, stripes are not a
prerequisite for coordinated and cohesive movement.

Using a comparative method where we accounted for phylogenetic signal, we
found that the contrasting and achromatic lines are more likely to be present across a
broad taxonomic range of social species that are often predated upon. It appears that
these lines may aid in group coordination, or act to increase the confusion effect, both
of which may reduce per capita predation risk. These achromatic patterns even suggest
that communication signals for alerting and recruiting others may even work within
mixed-species groups that are so prevalent among social birds –including ducks and
waders-, mammals –including gazelles- and fish (Goodale et al. 2017). Our work
represents a rare example of aligned phenotypic characteristics across a wide range of
taxonomic diverse animals.
369

Acknowledgements

371 We thank Enrique Figueroa-Luque for his help in building the dataset for
372 mammals, and Mr. C. Swann for the fish picture. AR was supported by Juan de la Cierva
373 programme, Spanish Ministry of Economy, Industry and Competitiveness (IJCI-2015-
374 23913).

Literature Cited

Allen, G.A., Robertson, D.D. (1998) Peces del Pacifico Oriental Tropical. Smithsonian
Tropical Research Institute. Conabio-Sierra Madre-CEMEX. México.

Armbruster JW and Page L.M. (1996) Convergence of a cryptic saddle pattern in benthic
freshwater fishes. *Environ Biol Fishes* 45:249–257.

Ballerini, M., Cabibbo, N., Candelier, R., Cavagna, A., Cisbani, E., *et al.* (2008a) Empirical
investigation of starling flocks: A benchmark study in collective animal behaviour. *Anim*
Behav 76:201–215.

Ballerini, M., Cabibbo, N. Candelier, R., Cavagna, A., Cisbani, E. *et al.* (2008b) Interaction
ruling animal collective behavior depends on topological rather than metric distance:
Evidence from a field study. *Proceedings of the National Academy of Sciences of the*
U.S.A. 105:1232–1237.

Banks, M.S., Sprague, W.S., Schmoll, J., Parnell, J.A., Love, G.D. (2015) Why do animal
eyes have pupils of different shapes? *Sci Adv* 2015;1: e150039.
390

391 Bennet, A., Thery, D. (2007) Avian Color Vision and Coloration: Multidisciplinary
392 Evolutionary Biology. *Am. Nat*. 169, pp. S1–S6.
393

394 Bowmaker, J.K. 1998. Evolution of colour vision in vertebrates. *Eye* 12, 541-547.
395

396 Brooke, M. de L. 1998. Ecological factors influencing the occurrence of 'flash marks' in
397 wading birds. *Funct Ecol* 12, 339–346.
398

399 Camacho, C., Negro, J.J., Redondo, I. *et al.* (2019) Correlates of individual variation in the
400 porphyrin-based fluorescence of red-necked nightjars (*Caprimulgus ruficollis*). *Sci*
401 *Rep* 9**,** 19115. https://doi.org/10.1038/s41598-019-55522-y.
402

Caro, T., Stankowich, T. (2010) The function of contrasting pelage markings in
Artiodactyls. *Behav Ecol* 21:78–84. doi:10.1093/beheco/arp165.

Cavagna, A., Cimarelli, A., Giardina, I., Parisi, G., Santagati, R., et al. (2010) Scale-free
correlations in starling flocks. *Proceedings of the National Academy of Sciences of the*
U.S.A. 107:11865–11870.

Cavagna, A., Giardina, I., Orlandi, A., Parisi, G., Procaccini, A. (2008a) The STARFLAG
handbook on collective animal behaviour. 2: Three-dimensional analysis. *Anim Behav*
76:237–248.

Cavagna, A., Giardina, I., Orlandi, A., Parisi, G., Procaccini, A. *et al.* (2008b) The STARFLAG
handbook on collective animal behaviour. 1: Empirical methods. *Anim Behav* 76:217–
236.

Cresswell, W. (1994) Flocking is an effective anti-predation strategy in redshanks, *Tringa*
totanus. Anim Behav 47: 433-442.

Cuthill, I. (2019) Camouflage. *Journal of Zoology* 308: 75-92.

Denton, E.J., Rowe, D.M. (1998) Bands against stripes on the backs of mackerel, *Scomber*
scombrus L. *Proc R Soc B*. 265:1051– 1058.

Figuerola, J., Green, A.J. (2000) The evolution of sexual dimorphism in relation to mating
patterns, cavity nesting, insularity and sympatry in the Anseriformes. *Funct Ecol* 14:701–
710.

Foster, W. A., Treherne, J. E. (1981). Evidence for the dilution effect in the selfish herd
from fish predation on a marine insect. *Nature*, 293(5832): 466-467.

Gelman, A., S, Y.S. (2018) arm: Data Analysis Using Regression and
Multilevel/Hierarchical Models. R package version 1.10-1. https://CRAN.R-
project.org/package=arm.

Goodale, E., Beauchamp, G., Ruxton, G. (2017). Mixed-species groups of animals.
Academic Press. London, U.K.

Guthrie D.M., Muntz W.R.A. (1993) Role of vision in fish behaviour. In: Pitcher, T.J. (ed.).
Behaviour of teleost fishes, 2nd edition. pp: 89–128. Chapman & Hall, Fish and Fisheries
Series 7. London.

Hamilton, W. D. (1971). Geometry for the selfish herd. *Journal of Theoretical*
Biology, 31(2): 295-311.

Handegard, N. O., Boswell, K. M., Ioannou, C. C., Leblanc, S. P., Tjøstheim, D. B., Couzin,
I. D. (2012). The dynamics of coordinated group hunting and collective information
transfer among schooling prey. *Current Biology*, 22(13), 1213-1217.

Hegyi, G., Garamszegi, L., Eens, M. (2008) The roles of ecological factors and sexual
selection in the evolution of white wing patches in ducks. *Behav Ecol* 19:1208-1216.
doi:10.1093/beheco/arn085.

Hemelrijk, C.K., van Zuidam, L., Hildebrandt, H. (2015) What underlies waves of agitation
in starling flocks. *Behav Ecol Sociobiol* 69:755–764.

Herbert-Read, J. E., Buhl, J., Hu, F., Ward, A. J., Sumpter, D. J. (2015) Initiation and
spread of escape waves within animal groups. *Royal Society Open Science*, 2(4),
140355.

Ho, L.S.T., Ané, C. (2014) A linear-time algorithm for Gaussian and non-Gaussian trait
evolution models. *Systematic Biology*, 63(3), 397-408.

Hogan, B., Cuthill, I.C., Scott-Samuel, N.E. (2017) Dazzle camouflage and the confusion
effect: the influence of varying speed on target tracking. *Anim Behav* 123: 349–353.
456

457 Hughes, A.E., Troscianko, J., Stevens, M. (2014) Motion dazzle and the effects of target
458 patterning on capture success. *BMC Evol Biol* 14: 201.
459

460 Ioannou, C.C., Guttal, V., Couzin, I.D. (2012) Predatory Fish Select for Coordinated
461 Collective Motion in Virtual Prey. *Science*, 337: 1212-1215.
462

Ives, A.R. (2019) R s for Correlated Data: Phylogenetic Models, LMMs, and
GLMMs. *Systematic Biology*, 68(2): 234-251.

Ives, A.R., Garland Jr, T. (2010) Phylogenetic logistic regression for binary dependent
variables. *Systematic Biology*, 59(1), 9-26.

Ives, A.R., Garland, T. (2014). Phylogenetic regression for binary dependent variables. In
Modern phylogenetic comparative methods and their application in evolutionary
biology (pp. 231-261). Springer, Berlin, Heidelberg.

Ives, A.R., Helmus, M.R. (2011). Generalized linear mixed models for phylogenetic
analyses of community structure. *Ecological Monographs*, 81(3), 511-525.

Jeschke, J. M., & Tollrian, R. (2007). Prey swarming: which predators become confused
and why? *Anim Behav*, 74(3), 387-393.

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., Mooers, A.O. (2012) The global diversity
of birds in space and time. *Nature* 491(7424), 444.

Kelley J.L., Fitzpatrick J.L., Merilaita S. (2013) Spots and stripes: ecology and colour
pattern evolution in butterflyfishes. *Proc R Soc* B 280:
20122730.http://dx.doi.org/10.1098/rspb.2012.2730.
479

480 Lima, S. L. (1995) Back to the basics of anti-predatory vigilance: the group-size
481 effect. *Anim Behav*, 49(1), 11-20.
482

Madge S, Burn H. (1987) Wildfowl: an identification guide to the ducks, geese and swans
of the world. London: Helm.

Martin, G.R. (2017) *The Sensory Ecology of Birds*. New York, United States of America,
Oxford University Press.

Marshall, N.J. (2000) Communication and camouflage with the same 'bright' colours in
reef fishes**.** *Phil.Trans. R. Soc. Lond. B* 355:1243-1248.

McGraw, K.J., (2006) Mechanics of melanin-based coloration. In: Hill, G.E., McGraw, K.
(Eds.), Bird Coloration, vol. I. Mechanisms and Measurements. Harvard University Press,
Cambridge MS, pp. 243–294.

McRobert S.P., Bradner J. (1998) The influence of body coloration on shoaling
preferences in fish. *Anim Behav* 56:611–615.

Osorio D.A., Miklosi, A., Gonda, Z. (1999) Visual ecology and perception of coloration
patterns by domestic chicks. *Evolutionary Ecology* 13: 673-689.
496
497 Paradis, E., Claude, J., Strimmer, K. (2004). APE: analyses of phylogenetics and evolution
498 in R language. *Bioinformatics*, 20(2), 289-290.

499 Pita D., Collignon B., Halloy J., Fernández-Juricic E. (2016) Collective behavior in
500 vertebrates: a sensory perspective. *R. Soc. open sci.* **3**: 160377.
501
502 Pomeroy, A.C. (2006) Tradeoffs between food abundance and predation danger in
503 spatial usage of a stopover site by western sandpipers, *Calidris mauri*. *Oikos* 112: 629-
504 637.
505
R Core Team. (2019) A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing; 2012. URL https://www. R-project. org.

Rabosky, D. L. (2015). No substitute for real data: a cautionary note on the use of
phylogenies from birth–death polytomy resolvers for downstream comparative
analyses. *Evolution* 69(12), 3207-3216.

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L. *et al.* (2018) An inverse
latitudinal gradient in speciation rate for marine fishes. *Nature* 559(7714): 392-395.

Ridley, A. R. N. J. Raihani, M., Bell, B.V. (2010) Experimental evidence that sentinel
behavior is affected by risk. *Biol Lett* 6: 445–448.
515
516 Ridley, A.R., Nelson-Flower, M.J., Thompson, A.M. (2013) Is sentinel behaviour safe? An
517 experimental investigation. *Anim Behav* 85: 137-142.
518
Romenskyy, M., Herbert-Read, J. E., Ioannou, C. C., Szorkovszky, A., Ward, A. J.,
Sumpter, D. J. (2020). Quantifying the structure and dynamics of fish shoals under
predation threat in three dimensions. *Behav Ecol*, 31(2), 311-321.

Rubolini, D, Liker, A., Garamszegi, L.G., Møller, A.P., Saino, N. (2015) Using the
BirdTree.org website to obtain robust phylogenies for avian comparative studies: A
primer. *Current Zoology* 61: 959–965.

Seehausen O., Mayhew P.J., van Alphen J.J.M. (1999) Evolution of colour patterns in East
African cichlid fish. *J Evol Biol* 12:514–534.

Smith, F.A., S.K. Lyons, S.K.M. Ernest, K.E. Jones, D.M. Kaufman, *et al.* (2003) Body mass
of late Quaternary mammals. *Ecology* 84:3402 (updated version obtained from senior
author).

Stevens, M., Yule, D.H., Ruxton, G.D. (2008) Dazzle coloration and prey movement. *Proc.*
R. Soc. B 275:2639–2643.
Stoner CJ, Caro TM, Graham CM. (2003) Ecological and behavioral correlates of
coloration in artiodactyls: systematic analyses of conventional hypothesis. *Behav Ecol*
14: 823-840.
535
Sukumaran, J., Holder, M.T. (2010) DendroPy: a Python library for phylogenetic
computing. *Bioinformatics* 26(12), 1569-1571.

Sukumaran, J., Holder, M.T. (2015) SumTrees: Phylogenetic Tree Summarization. 4.4.0.
Available at https://github.com/jeetsukumaran/DendroPy.

Svensson, L., Mullarney, K., Zetterström, D., Grant P.J. (1999) Collins Bird Guide: the
most complete guide to the birds of Britain and Europe, 2nd ed. – Harper Collins.
542
543 Upham, N. S., Esselstyn, J. A., Jetz, W. (2019) Inferring the mammal tree: Species-level
544 sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biology*,
545 17(12), e3000494.

546   Table 1. Summary of phyloglm models.

	Estimate	SE	95% CI	R^2_{lik}	Z	p
Charadriiformes	-	-	-	0.261	-	-
α	0.015	-	0.015, 0.015	-	-	-
Intercept	0.105	0.811	0.185, 0.845	-	0.129	0.897
Sociality	0.773	0.291	0.552, 0.793	-	2.659	0.008
Wing chord	-0.010	0.003	0.497, 0.500	-	-1.550	0.121
Migratory status	-0.025	0.047	0.470, 0.517	-	-0.526	0.599
Anseriformes	-	-	-	0.180	-	-
α	0.016	-	0.016, 0.016	-	-	-
Intercept	1.688	1.013	0.426, 0.975	-	1.666	0.096
Body mass	-0.001	0	0.500, 0.500	-	-3.202	0.001
Migratory status	-0.222	0.236	0.335, 0.56	-	-0.943	0.346
Ruminants	-	-	-	0.273	-	-
α	0.136	-	0.031, 0.743	-	-	-
Intercept	-1.885	0.506	0.053, 0.290	-	-3.728	0
Sociality	1.393	0.475	0.614, 0.911	-	2.936	0.003
Body mass	0	0	0.500, 0.500	-	-1.502	0.133
Fish (longitudinal)	-	-	-	0.369	-	-
α	0.016	-	0.008, 0.133	-	-	-
Intercept	-0.548	0.342	0.228, 0.531	-	-1.602	0.109
Sociality	1.616	0.367	0.710, 0.912	-	4.402	0
Fish (vertical)	-	-	-	0.162	-	-
α	0.046	-	0.024, 0.28	-	-	-
Intercept	0.142	0.232	0.422, 0.645	-	0.612	0.541
Sociality	-0.9	0.346	0.171, 0.444	-	-2.605	0.009

Figure legends

Fig. 1. Species of birds, mammals and fishes with examples of contrasting patterns. From
left to right, and top to bottom: Mallard *Anas platyrhynchos*, including both males and
females; northern shoveler *Spatula clypeata*; common oystercatcher *Haematopus*
ostralegus; black-tailed godwit *Limosa limosa*; springbok *Antidorcas marsupialis*; dorcas
gazelle *Gazella dorcas*. In the fish picture: with longitudinal stripes, spottail grunts
Haemulon maculicauda, and with vertical stripes, Panamic sergeant major *Abudefduf*
troschelii. Credits: birds and mammals; Juan J. Negro. Fish picture: Christopher Swann.

Fig 2. Trait plots depicting the distribution of contrasting stripes (colored traits) across
the phylogeny of shorebirds (Charadriiformes, a, gray), Anseriformes (b, orange),
Ruminants (c, brown), and Fishes (d, blue).

Fig 3. Phylogenetic logistic regression curves for Charadriiformes (a, gray), Anseriformes
(b, orange), Ruminants (c, brown), and Fishes (d, blue). Individual points are horizontally
and vertically jittered to improve visualization.

Figure 1

Figure 2

Figure 2

Figure 3

573