Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*

Wiberg, R. A. W.[1,+,*], Tyukmaeva, V.[2+], Hoikkala, A.[2], Ritchie, M. G.[1] & Kankare, M.[2]

1. Centre for Biological Diversity, School of Biology, University of St Andrews, St Andrews, KY16 9TH, Scotland, UK.

2. Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, 40014, Finland.

+**Present addresses: RAWW** University of Basel, Department of Environmental Sciences, Zoological Institute, Vesalgasse 1, 4051 Basel, Switzerland, **VT** Centre d'Ecologie Fonctionelle et Evolutive, CNRS, Montpellier, France.

* Correspondence: raw.wiberg@protonmail.com

Keywords: Environmental adaptation, genomic divergence, cold tolerance, CTmin, CCRT, cline populations, *D. montana*

ORCID

RAWW	0000-0002-8074-8670
VT	0000-0002-1864-5101
AH	0000-0001-5407-7992
MGR	0000-0001-7913-8675
MK	0000-0003-1541-9050

21 **ABSTRACT**

Detecting signatures of ecological adaptation in comparative genomics is challenging, but analysing population samples with characterised geographic distributions, such as clinal variation,

can help identify genes showing covariation with important ecological variation. Here we analysed patterns of geographic variation in the cold-adapted species *Drosophila montana* across phenotypes, genotypes and environmental conditions and tested for signatures of cold adaptation in population

genomic divergence. We first derived the climatic variables associated with the geographic distribution of 24 populations across two continents to trace the scale of environmental variation experienced by the species, and measured variation in the cold tolerance of the flies of six

populations from different geographic contexts. We then performed pooled whole genome sequencing of these six populations, and used Bayesian methods to identify SNPs where genetic differentiation is associated with both climatic variables and the population phenotypic

measurements, while controlling for effects of demography and population structure. The top candidate SNPs were enriched on the X and 4th chromosomes, and they also lay near genes implicated in other studies of cold tolerance and population divergence in this species and its close

relatives. We conclude that ecological adaptation has contributed to the divergence of *D. montana* populations throughout the genome and in particular on the X and 4th chromosomes, which also showed highest interpopulation F_{ST}. This study demonstrates that ecological selection can drive

genomic divergence at different scales, from candidate genes to chromosome-wide effects.

INTRODUCTION

42 The geographic structure of a species is a result of its phylogeographic history, influenced by past and present dispersal, population demography and selection. Obtaining genome-wide data on genetic polymorphisms across multiple populations of a species is becoming relatively easy, but

45 interpreting the patterns of geographic variation in such data and identifying genes which vary primarily due to selection remains challenging. Often simple 'outlier' approaches using genome scans which measure genetic differentiation such as F_{ST} or Dxy are adopted, but results are difficult

48 to interpret due to confounding between selection, drift and population structure, or genomic features such as inversions and other causes of variation in recombination rate (Noor & Bennet, 2009; Cruickshank & Hahn, 2014; Wolf & Ellegren, 2016; Ravinet et al., 2017). If environmental

51 data is available, we can use associations with such factors to help identify loci where genetic differentiation covaries with this environmental variation. Some genome scan methods can incorporate environmental variation and simultaneously fit effects for covariance with

54 environmental factors, while controlling for effects of population demography (Foll & Gaggiotti

2008; de Villemereuil & Gaggiotti 2015). This approach has successfully identified genetic variation associated with altitude in humans, among other examples (Foll, Gaggiotti, Daub, Vatsiou, & Excoffier, 2014; de Villemereuil & Gaggiotti, 2015; Gautier, 2015) and has become a useful approach to investigate the ecological adaptations underlying population divergence.

Clinal patterns of variation in phenotypes or gene frequencies have a long history of being used to infer selection along ecotones, and analyses of cline shape can sometimes identify loci under direct selection from others showing clinal variation for other reasons such as phylogeographic history (Barton & Gale, 1993). Such studies can be very powerful, especially when independent parallel clines are available. For example, Kolaczkowski, Kern, Holloway & Begun (2011) sampled isofemale lines from extremes of a cline in Australian populations of *D. melanogaster*, and found many genes implicated in clinaly varying phenotypes to show highest differentiation. Also, Bergland, Behrman, O'Brien, Schmidt & Petrov (2014), and Kapun, Fabian, Goudet & Flatt (2016) sampled North American clines in *D. melanogaster* and *D. simulans* over several years and seasons to uncover clinal and seasonal variation at a genome scale. They found that many SNPs showed consistent seasonal fluctuations in allele frequencies throughout the clines, which indicates a regular response to seasonally varying selection pressures (Bergland et al., 2014). On the other hand, Machado et al. (2015) concluded that migration and gene flow play a greater role than adaptation in the overall clinality of genomic variants in *D. simulans* than *D. melanogaster*. While the two species share a significant proportion of the genes showing clinal variation, their differences in overwintering ability, migration and population bottlenecks probably act as additional drivers of differences in patterns of variation between them (Machado et al., 2015). Similar studies of clinal variation in phenotypes and allele frequencies have also been carried out in other insects (e.g. Paolucci, Salis, Vermeulen, Beukeboom, & van de Zande, 2016), plants (e.g. Chen et al., 2012; Bradbury, Smithson, & Krauss, 2013), mammals (e.g. Hoekstra, Drumm, & Nachman, 2004; Carneiro et al., 2013), fish (e.g. Vines et al., 2016), and other organisms (Endler, 1973; Endler, 1977; Takahashi, 2015). However, more studies are needed that simultaneously compare patterns of variation in allele frequencies and potentially causal environmental variation or ecologically important traits while controlling for population structure, as such studies are necessary to determine if adaptation to climate is directly driving patterns of genetic differentiation.

Here we investigated geographic variation at both the phenotypic and genetic levels in *Drosophila montana* samples from two continents. This species has spread around the northern hemisphere (Throckmorton, 1969) and both mtDNA and microsatellite data revealed genetically distinct Finnish and North American populations (Mirol et al., 2007). Moreover, more recent modelling of genome-wide SNP frequencies suggest that the Finnish-North American split happened around 1.75MYA and that of the North American populations shortly after that

(Garlovsky et al., 2020). It is one of the most cold-tolerant *Drosophila* species (Kellerman et al., 2012;) and the basic cold tolerance of *D. montana* flies can increase towards the cold seasons through two mechanisms, photoperiodic reproductive diapause (Vesala & Hoikkala, 2011) and cold-

acclimation induced by a decrease in day length and/or temperature (Vesala, Salminen, Laiho, Hoikkala, & Kankare, 2012; Kauranen et al., 2019). *D. montana* populations have been found to show clinal variation in the critical day length required for diapause induction (CDL; Tyukmaeva,

Salminen, Kankare, Knott, & Hoikkala 2011; Lankinen, Tyukmaeva, & Hoikkala 2013). There is also a correlation between CDL and latitudinally co-varying climatic factors such as the mean temperature of the coldest month (Tyukmaeva, Lankinen, Kinnunen, Kauranen, & Hoikkala, 2020).

In addition, *D. montana* populations from different geographic regions show variation in their courtship cues and mate choice (Routtu et al., 2007; Klappert, Mazzi, Hoikkala, & Ritchie, 2007), which has led to partial reproductive isolation between some distant populations (Jennings, Mazzi,

Ritchie, & Hoikkala, 2011; Jennings, Snook, & Hoikkala, 2014). At the genetic level, differential gene expression studies have identified candidate genes underlying diapause (Kankare, Salminen, Laiho, Vesala, & Hoikkala, 2010; Kankare, Parker, Merisalo, Salminen, & Hoikkala, 2016),

perception of day length (Parker, Ritchie, & Kankare, 2016), and cold acclimation (Parker et al., 2015). Furthermore, a quasi-natural selection experiment for shorter CDL, accompanied by a decrease in cold-tolerance, induced widespread changes in loci with potential roles with these traits

(Kauranen et al., 2019). Finally, population genomic analyses have identified several outlier loci when examining differentiation between North American and European populations (Parker et al., 2018). All this makes *D. montana* an interesting example of nascent speciation, potentially

influenced by ecological adaptation.

        Here we sought to ask to what extent patterns in the genomic divergence of *D. montana* populations across two continents are correlated with climatic variation and phenotypic responses to

114   cold adaptation. We performed pooled whole-genome sequencing (pool-seq) on six different populations and used Bayesian methods to examine the association between genomic differentiation between populations and environmental variables across both continents. We also phenotyped

117   populations for two different cold tolerance measures; critical thermal minimum (CTmin), and chill coma recovery time (CCRT), and investigated the associations between them and the genetic and climatic data. Ultimately, we asked if the genomic loci showing an association between genetic and

120   environmental differentiation also showed association with population differentiation in cold tolerance phenotypes, and examined the possible overlap between the set of genes close to candidate SNPs with sets of candidate genes from previous studies of cold adaptation in *D.*

123   *montana*. If population differentiation is driven by ecological selection then we would predict the

extreme cold adaptation of *D montana* to have left a signature of genomic divergence associated with environmental and phenotypic differentiation across these loci.

MATERIALS AND METHODS

Sample collections and DNA extraction

129   We collected samples of 49-50 wild-caught flies from six *D. montana* populations from a range of latitudes from 66°N to 38°N in the spring of 2013 or 2014. Four of these populations represented a range of latitudes in North America (N. A.), and two populations were from a range in Finland

132   (figure 1; table 1). Samples of wild-caught flies from the six populations were stored in ethanol (the male/female ratio varied across samples; table 1) and DNA of individual flies was extracted using CTAB solution and phenol-chloroform-isoamylalcohol purifications in 2016. Genomic DNA was

135   extracted from individual flies and quantified using Qubit (Thermo Fisher Scientific), and an equal amount of DNA from each individual (50 ng) was pooled into the final sample. Sequencing was performed at the Finnish Functional Genomics Centre in Turku, Finland (www.btk.fi/functional-

138   genomics) on the Illumina HiSeq3000 platform (paired-end reads, read length = 150bp, estimated coverage ~121x).

141   **Phenotyping**

We measured the critical thermal minimum (CTmin) and chill coma recovery time (CCRT) of flies from six populations. Fly samples for these tests were collected for five populations (Seward,

Terrace, Ashford, Crested Butte and Korpilahti) from mass-bred populations that have been maintained in the lab since 2013-2014. For the Oulanka population, flies were collected from three isofemale strains (established in 2014), because the mass-bred population had been contaminated by

another species. The mass-bred populations were originally established from F4 progenies of 20 isofemale strains each (400 flies) and have been maintained in constant temperature (19 ± 1°C) and light (24h of light, to prevent the flies from entering diapause) regimes for about 20-25 generations

before the experiment. All flies were supplied with fresh malt medium in half-pint bottles every week (Lakovaara, 1969). The newly emerged flies were collected using light CO_2 anaesthesia within 24 hours after emergence, separated by sex and placed in malt-vials in the same conditions

until sexual maturity (20-21 days old) and were then used in CTmin and CCRT tests. The same individual flies were first assayed for CTmin and then for CCRT, the flies were not anaesthetised before these tests.

We assayed a total of 328 females and 302 males for CTmin and CCRT. These assays were done in batches of between 22 and 30 flies, and split evenly by sex (21 batches in total). Between 32 and 46 (mean 39) flies per population per sex were tested (for the Oulanka isofemale strains,

between 15 and 39 flies per strain per sex were used). CTmin tests are based on detecting the temperature (CTmin) at which flies lose neuromuscular function and enter reversible state of chill coma (Andersen et al., 2015). In these tests, the flies were placed into tubes sealed with parafilm

and submerged into a 30% glycol-water mixture in Julabo F32-HL chamber. The temperature was decreased at the rate of 0.5°C per minute (from 19°C to -6°C) and CTmin was determined by eye, as the temperature at which a fly was unable to stand on its legs. Immediately after the Ctmin test,

the temperature was set to -6°C and the flies were left in this temperature for 16 hours. Then vials were quickly taken out of the glycol-water bath and the flies' CCRT was determined as the time required for the flies to recover from chill coma and stand on their legs. The ambient room

temperatures was recorded during trials but in an initial analysis including source population and room temperature there was no statistically significant effect of room temperature on CCRT ($F_{5,569}$ = 0.72, p = 0.4), this variable was therefore left out of further analyses.

To investigate population differences in CTmin and CCRT phenotypes we fit General Additive Mixed Models (GAMMs) in R (v. 3.6.3; R Development Core Team, 2020) using the "mgcv" R package (Wood, 2004). The full model included altitude, latitude, and sex as fixed effects

and experimental batch as a random effect. We used a cubic regression spline as the basis smoothing function for both altitude and latitude. The full models are given in the Supplementary Materials.

Bioclimatic variables and population geography

We obtained representative climate data from the WorldClim database (Hijmans, Cameron,

180 Parra, Jones, & Jarvis, 2005) for each *D. montana* population sampled for the pool-seq (see above), as well as for 18 additional populations of this species (Tyukmaeva et al., 2020). We downloaded the climate data and extracted the values corresponding to population coordinates using the R

183 package "raster" (v. 2.5-8; Hijmans et al., 2016). In total this amounts to 55 bioclimatic variables for each population (table S1). To reduce the number of variables in the dataset a principle components analysis (PCA) was performed using the "PCA()" function from the R package

186 "FactoMineR" (v. 1.28; Lê, Josse, & Husson, 2008). Principle components were kept for further analysis if their eigenvalues were > 1. PCA scores for each population were z-transformed using the "scale()" function in base R. Additionally, CTmin and CCRT were summarised to a mean value for

189 each population. In total, this gives four "environmental" variables measured for each population (PC1, PC2, CTmin, and CCRT).

192 **Mapping, SNP calling and genomic analysis**

Quality of raw reads was checked with FASTQC (v. 0.11.5) (Andrews, 2015) and reads were trimmed using trimmomatic (v. 0.32; Bolger, Lohse, & Usadel, 2014; see Supplementary Materials for full trimming parameters). Trimmed reads were mapped to the *D. montana* reference genome (Parker et al., 2018) using BWA mem (v. 0.7.7; Li, 2013) with the default options but keeping only alignments with a mapping quality of > 20 following best practice guidelines for pool-seq (Schlötterer, Tobler, Kofler, & Nolte, 2014). Duplicate alignments were removed with samtools rmdup (v 1.3.1; Li et al., 2009) and regions around indels were re-aligned using picard (v. 1.118, Broad Institute), GATK (v. 3.2-2, McKenna et al., 2010) and samtools. Separate .bam files for each of the sequenced populations were finally merged using bamtools (v. 2.4.0; Barnett, Garrison, Quinlan, Strömberg, & Marth, 2011).

Over 80% of reads were properly mapped and retained in all samples. The mean coverage for Seward samples was nearly twice that of the other samples (figure S1 and figure S2). To remove the potential for this difference to cause artefacts in downstream analyses, the .bam files for Seward were down-sampled to contain 94.1 million reads (the average across the 5 remaining populations). Median empirical coverage was between ~62 and 88x (table S2; figure S1) and much less variable among the populations, allowing common maximum and minimum thresholds to be set based on the aggregate distribution (figure S3). Allele counts for each population at each genomic position were obtained with samtools mpileup (v. 1.3.1; Li et al., 2009) using options to skip indel calling as well as ignoring reads with a mapping quality < 20 and sites with a base quality < 15. This was followed by the heuristic SNP calling software PoolSNP using a minimum count of 5 to call an allele, and a minimum coverage of 37 and a maximum coverage < 95th percentile of the scaffold-wide coverage distribution to call a SNP (Kapun et al., 2019). Even if all these filters were passed, an allele was not considered if its frequency was < 0.001. Finally, we only considered SNPs on scaffolds > 10kb in length. The final set consisted of 2,190,511 biallelic SNPs that could be placed on scaffolds ordered along the chromosomes and were used in downstream analyses (table 2).

To test for an association between the four environmental variables and genetic differentiation we used BayeScEnv (v. 1.1; Foll & Gaggiotti, 2008; de Villemereuil & Gaggiotti, 2015). BayeScEnv fits a model of F_{ST} to population differentiation for each locus, incorporating environmental differentiation as a predictor (included as the parameter *g*) while fitting two locus-specific effects, one for environmental selection, and the other for other processes (demography, other types of selection). It therefore controls for confounding effects of population structure/relatedness in testing for an association with environmental variables. BayeScEnv was run with five pilot runs of 1000 iterations each, followed by a main chain of 4,000 iterations of which 2,000 were discarded as burn-in. Four MCMC chains were run for each analysis to evaluate convergence of the chains to common parameter estimates. Because of the unbalanced number of

228   males (and the resulting variation in ratios of X:Y chromosomes) in the pools, BayeScEnv analyses were performed separately on SNPs that could be assigned to the autosomal linkage groups (chromosomes) and the X chromosome. Raw count data were used for the autosomal data. For X

231   linked SNPs, allele count data were scaled to the known number of X chromosomes in the pool using n_{eff}, the effective sample size taking into account the multiple rounds of binomial sampling inherent to a pool-seq design (Kolaczkowski et al., 2011; Feder, Petrov, & Bergland, 2012). N_{eff}

234   scaled the allele counts at each SNP downwards based on the known number of chromosomes in the pool (see table 1).

    Chains were assessed for convergence with the "coda" R package (v. 0.19-1; Plummer, Best,

237   Cowles, & Vines, 2006). Convergence was reached across the four chains for most analyses and parameters (potential scale reduction factors (PSRFs) of ~1 in a Gelman-Rubin diagnostic test; figures S4-S7), except for analyses of autosomal SNPs and PC2 as the environmental variable

240   which showed mild convergence problems (PSRF = 1.71), although parameter estimates agreed well with the other chains. Thus, this first chain (black line in figures S4-S7) was discounted for all parameters and only estimates from the remaining 3 chains were used. The union of significant

243   SNPs (q-value for the g parameter < 0.05) across these chains were taken as the final candidate SNPs.

    Finally, population genetic statistics (π, and Tajima's D) were computed in windows of 10kb

with a step size of 5kb using methods implemented for pool-seq data (Kapun et al., 2019). Windows contained a mean of 488.2 SNPs (SE ±1.2). These statistics were only computed for scaffolds with a length > 50kb. F_{ST} was computed for each population with the R package "poolfstat" (v. 1.1.1;

Hivert, Leblois, Petit, Gautier, & Vitalis, 2018) by first computing all pairwise values, and then deriving population specific F_{ST} values by averaging across all pairwise values where a population was included. See the supplementary material for pseudocode commands of the key pipeline steps.

We performed GO term enrichment analyses with DAVID (v. 6.8; Huang et al., 2009a, b) and GOwinda (v. 1.12; Kofler & Schlötterer), which accounts for differences in gene length. For DAVID, since the *D. montana* annotation contains information about orthologs in *D. virilis,* we

extracted all genes within 10kb of a candidate SNP with an ortholog in *D. virilis* and submitted them to DAVID. For analyses with GOwinda, because the gene-sets need to be given manually, we obtained gene-sets from FuncAssociate3 (Berriz, Beaver, Cenik, Tasan & Roth 2009). As above, we

considered SNPs within 10kb of a gene (--gene-definition updownstream10000) and performed 1 million simulations to obtain the empirical p-value distribution. Regulatory elements, such as enhancers, and transcription factor binding sites, can occur up to 1Mb up- or downstream from a

target gene in other species (e.g. Maston, Evans & Green. 2006; Chan et al., 2010; Werner, Koshikawa, Williams & Carroll 2010; Pennacchio, Bickmore, Dean, Nobrega, & Bejerano 2013)

but generally lie within 2kb of a gene region in *D. melanogaster* (Arnosti, 2003), thus 10kb represents a compromise.

RESULTS

Cold tolerance measures, bioclimatic variables and population geography

Across individuals, there was no evidence of an association between minimum critical temperature (CTmin; pooled across sexes) and the chill coma recovery time (CCRT; cor = 0.08, p = 0.07). CTmin and CCRT varied among populations according to sex, latitude, and altitude. CTmin was significantly different between sexes (t = 3.14, p = 0.002). Latitude also had a strong effect on CTmin, and the Effective Degrees of Freedom (edf = 1), suggested this was largely a linear effect after accounting for altitude (F = 40.4, p < 0.001; figure 2A). Altitude had a more complex effect (edf = 1.262, F = 24.0, p < 0.001), but with a strong altitude outlier in Crested Butte (table 1), this result should be taken with some caution. Moreover, the overall adjusted R-squared was low at only 0.09. For CCRT, only latitude had a significant effect (edf = 1, F = 17.2, p < 0.001; figure 2B). Taken together, CTmin was lower for populations at higher latitudes as well as for populations at higher altitudes (figure 2A). As one would expect, CTmin showed on average lower values (figure 2A), and CCRTs were shorter (figure 2B), at higher latitudes, meaning that more northern and higher altitude populations show higher cold tolerance.

We performed a Principal Component Analysis (PCA) of the WorldClim climate data for a total of 24 *D. montana* populations, from which we had collected samples and where climate data were available. We wanted to examine whether the populations chosen for the cold tolerance tests and genome sequencing were representative of the range of environmental variation experienced by the species. These analyses identified four Principal Components (PCs) that together explained about 98% of the variation (figure 3A; figure S8). The first two PCs separated the populations roughly by a measure of distance from the coast (PC1) and then by latitude (or altitude) (PC2). PC1 explained ~54% of the variation (figure 3A and B) and loaded heavily on climate and biological variables associated with precipitation and temperature such as "Mean Temperature of Coldest Quarter", "Precipitation of Wettest Month", "Annual Precipitation" (see figure S8 and table S3). Meanwhile, PC2 explained ~23% of the variation (figure 3A and C) and loaded heavily on biological variables that are associated with latitudinal clinality, e.g. "Mean Diurnal [Temperature] Range," and "Isothermality" which is the diurnal range divided by the mean "Annual [Temperature] Range". The remaining PCs (PC3 and PC4) explained about 11.5 and 5% of the variation, respectively and did not capture as much of the climatic variation, and were therefore not considered further. Latitude (Spearman's Rank Correlation: rho = -0.50, p = 0.01) but not altitude (rho = -0.19, p = 0.39) correlated significantly with PC1. However, both altitude (rho = -0.50, p = 0.02) and latitude (rho =

-0.59, p = 0.003) correlated with PC2 (see also figure 3C). Importantly, all these patterns also held if PCA was performed using only the six populations for which genomic data were collected. This suggests that environmental variation across the six populations selected for sequencing reflects variation experienced by all 24 populations. Thus, any relationship between environmental variables and genetic differentiation in the samples selected for pool-seq is likely to reflect true patterns across populations of *D. montana*. For all subsequent analyses, we used the results of the PCA using all 24 populations, and to examine the association between climate and phenotype, we compared these across the six populations. CTmin was positively correlated with PC1 (Pearson's correlation coefficient (cor) = 0.94, p < 0.01) and had a marginally non-significant association with PC2 (cor = 0.75, p = 0.09). However, CCRT showed no relationship with either PC1 (cor = 0.09, p = 0.87) or PC2 (cor = -0.46, p = 0.36) although the small sample sizes (N = 6 in all cases) makes reliable conclusions difficult (see figure S9).

Genomic analyses

The number of SNPs with a significant (q-value < 0.05) association between population-based F_{ST}, the two cold tolerance measures (CCRT and CTmin), and the two PCs of the bioclimatic data varied from 312 (chromosome 3, CCRT) to 2,480 (chromosome 4, PC2) across the chromosomes (table 2). Using PC1 as an environmental variable with BayeScEnv gave a total of 2,976 and 1,528 SNPs with a q-value < 0.05 on the autosomes and on the X chromosome, respectively. Interestingly, the distribution across the chromosomes was not random. Using the distribution of all SNPs to obtain expected counts, there was a significant deviation from expectation (X^2 = 2,906.4, d.f. = 4, p < 0.001). There were many more SNPs than expected on chromosome 4 (1,432 vs. 954) and on the X chromosome (1,528 vs. 526). Results were similar for PC2 with 6,607 and 1,861 SNPs with a q-value < 0.05 on the autosomes and X chromosomes, respectively. Again, there was a significant deviation from the expected distribution of SNPs across the chromosomes (X^2 = 1,681.9, d.f. = 4, p < 0.001) with an overrepresentation on the 4th (2,480 vs. 1,794) and the X chromosomes (1,861 vs. 989). Manhattan plots of the distribution of SNPs across chromosomes are given in figures S10-S13.

We also used average CTmin values per population in similar analyses and found a total of 2,668 and 1,272 SNPs with a q-value < 0.05 on the autosomes and X chromosomes, respectively. The pattern of significant deviations from expected distributions (X^2 = 2,526.2, d.f. = 4, p < 0.001) was also due to an excess on the 4th (1,383 vs. 835) and the X chromosomes (1,272 vs. 460). Similar results were found for CCRT with a total of 2,240 and 1,228 SNPs with a q-value < 0.05, respectively. Once again, there was a significant deviation from the expected distribution of SNPs

(X^2 = 2,825.6, d.f. = 3, p < 0.001) with and excess on the 4th (1,252 vs. 735) and the X chromosomes (1,228 vs. 405).

To more closely examine the loci implicated in the four BayScEnv analyses, we identified genes containing or nearby (within 10kb) the candidate SNPs (table S4). Overall, there is quite a large overlap among these genes with ~39% (1,102 in total) of them being shared by all the four analyses (table S4, figure 4). Some (147, ~13%) of these common genes are novel to *D. montana* (i.e. not annotated in *D. virilis* or in other *Drosophila* spp.) and therefore have no annotation, but 955 have an identifiable *D. virilis* ortholog (table S4). The second largest set are those genes unique to the analysis of PC2 as an environmental covariate (figure 4). We performed functional enrichment analyses with DAVID keeping all annotation clusters with an enrichment score > 1.3 (corresponding to an average corrected p-value of 0.05). This revealed several common categories of genes associated with the climatic variables and population phenotypes (table S5). For example, terms associated with membrane and transmembrane structures, immunoglobulins, HAD hydrolase and nucleotide binding were enriched in most of the variables (table S5). Interestingly, there were also several gene ontology categories that were only enriched in one of the variables, such as glycoside and ATPase hydrolase in CCRT and ion channels and transport, as well as metal binding in PC1 (table S4, table S5). GOwinda analyses revealed significant enrichment of GO terms (after accounting for variation in gene lengths and correcting for multiple testing) only for genes near SNPs associated with variation in CCRT across populations. Interestingly, the term "carbohydrate derivative binding" was identified among the most enriched terms (table S6), which agrees closely with some terms identified using DAVID for the same variables (table S5). Similarly, for PC2, "nucleotide binding" was among the top scoring enriched terms in both GOwinda and DAVID analyses, though this was not significant after controlling for multiple testing in GOwinda (table S5, table S6)

We then compared these loci with genes implicated in previous studies of climatic adaptation in *D. montana* including gene expression studies of traits connected to diapause and cold-tolerance (Kankare et al., 2010; 2016; Parker et al., 2015; 2016). Additionally, several candidate genes have been identified near the most significantly differentiated SNPs among *D. montana* populations from Oulanka (Finland), and from North American populations in Colorado and Vancouver (Parker et al., 2018). Finally, quasi-natural selection experiments identified several genes within 10kb of SNPs responding to selection for a shorter CDL for diapause induction (Kauranen et al., 2019). We tested for an overlap between the total set of genes within 10kb of outlier SNPs from all of the BayScEnv runs (N = 2,694) and the candidate gene sets identified in earlier studies (see table S7 for the gene sets and studies used). We computed a bootstrap distribution of overlaps by sampling 2,694 random genes from the *D. montana* annotation (N = 13,683). For each of the gene sets from previous

studies this was done 100 times and the distribution compared to the empirical overlap. Results are given in figure S10. In all the cases the empirical overlap was greater than expected by chance with empirical p-values < 0.05 (ranging from < 0.001 to 0.014; figure S14). The only gene that was found in all five of the previous studies used and in the comparison here is called *sidestep II* (*side-II*; table S8). Unfortunately, there is no information available about the biological processes or molecular functions connected to it. Moreover, from 44 other genes that were common to four of our previous studies and this study (table S8) most (27) have an ortholog in *D. melanogaster*. These genes have molecular functions such as transmembrane signaling or transporter, acetylcholinesterase, ATP binding, protein serine/threonine kinase, carboxylic ester hydrolase, or Rho guanyl-nucleotide exchange factor activity (Thurmond et al., 2019). Many of the genes are also connected to metal ion, nucleid acid or zinc ion binding (table S8). After identifying information on molecular or biological function and Interpro domains, eventually only five genes remained for which there was no information available (table S8).

Examination of population genetic parameters identified the Crested Butte population as anomalous. The distribution of Tajima's D is centered close to zero in most populations, being slightly more negative in North American populations (figure S15). However, Crested Butte is an outlier with a greatly reduced genome-wide Tajima's D (figure S15). Similarly, diversity (π) is also lower in this population than in other populations. There is no overall relationship between latitude and π (Spearman's rho = 0.14, N = 6, p = 0.8; figure S15) but there is a strong correlation between latitude and Tajima's D which is influenced by this population (with Crested Butte: rho = 0.88, N = 6, p = 0.03; without Crested Butte: rho = 0.8, N = 5, p = 0.13). Although Crested Butte occurs at a much higher altitude (>2800 meters) than other populations neither Tajima's D nor π correlated significantly with altitude (Tajima's D: rho = -0.6, N = 6, p = 0.24, π: rho = -0.6, N = 6, p = 0.24). Furthermore, F_{ST} was similar across all populations and chromosomes with the exception of Crested Butte which remained an outlier with clearly high F_{ST} (table 3). Finally, F_{ST} was always highest on chromosome 4 and the X chromosome, complementing the results seen in BayeScEnv analyses (table 3 *c.f.* table 2) and as expected if ecological selection is influencing genomic divergence.

DISCUSSION

Detecting genomic signatures of climatic adaptation is an important, but challenging, task. Here we use multiple sources of evidence to study ecological adaptation and population divergence in a highly cold tolerant species of *Drosophila*, *D. montana*. This species is characterised by a wide circumpolar distribution extending to high latitudes both in North America and Europe, and to high altitudes in the southern part of its range in the Rocky Mountains of North America. These habitats impose extreme seasonal and climatic selective pressure. We collected bio-climatic data from 24

populations along a latitudinal gradient of about 2,900 km in North America, and six populations from a gradient of 720 km in Finland. We characterised population level cold-tolerance for six populations from these ranges using two methods, critical thermal minimum (CTmin) and chill coma recovery time (CCRT) and show that CTmin were lower and CCRT shorter in higher latitude populations, as one would expect. Thus, northern populations are more cold-tolerant. Finally, we performed pool-seq of these six populations to investigate the association between genomic and environmental differentiation.

The two methods examining cold tolerance gave somewhat different results, as CTmin, but not CCRT, differed significantly between sexes, with females having lower CTmin than males. In an earlier study investigating seasonal changes in *D. montana* CCRTs, only one out of six comparisons showed a significant difference between sexes (Vesala, Salminen, Kostal, Zahradnickova, & Hoikkala, 2012) and Gibert, Moreteau, Petavy, Karan & David (2001) did not detect sex-specific differences in CCRT in any of 84 *Drosophila* species they studied. However, several studies of *D. melanogaster* have detected shorter CCRT in females than in males, suggesting that females are more cold tolerant than males (David et al., 1998; Andersen et al., 2015; Bauerfeind, Kellermann, Moghadam, Loeschcke, & Fischer, 2014), perhaps related to their greater body mass (e.g. Wilder et al., 2010). Consequently, the extent and adaptive significance of sex-specific differences in CCRT in *Drosophila* remains unclear.

We derived principal components to summarise WorldClim climatic variables using data from all the 24 populations. The first principal component (PC1) separated populations roughly by a measure of "distance inland" and loaded heavily on climate and variables associated with precipitation and temperature. These results follow the geographic distribution of the populations, for example, the population with highest values for PC1 is Ashford, which is on the Pacific coast and receives most rain, but also experiences warm summers and mild winters. Principal component 2 (PC2), loaded heavily on bioclimatic variables associated with latitudinal clinality, which also mapped onto the populations intuitively as those with higher values on PC2 also occurred at higher latitudes. Interestingly, CTmin values were positively correlated with PC1, but not with PC2, while CCRT showed no relationship with either of these components. This suggests that CTmin and CCRT measure at least slightly different biochemical or physiological mechanisms (see e.g. MacMillan, Williams, Staples, & Sinclair, 2012; Findsen, Pedersen, Petersen, Nielsen & Overgaard 2013), and could hence be correlated with different climatic variables and also show relatively weak correlation to each other (Andersen et al., 2015). It is also interesting to note that high altitude populations have very similar values on PC1 and PC2 to high latitude populations, most likely reflecting the similar climates in these populations. These similarities in climate are likely also

underlying the similarities in CTmin between the high altitude population (Crested Butte) and the high latitude populations from Finland.

The Bayesian analysis identified SNPs showing an association of genetic differentiation with climatic and phenotypic variation. The extent to which the loci associated with the phenotypes and adaptations to different climatic conditions are shared indicates that these are closely associated in

influencing genome evolution. We found that genes near SNPs showing a significant association between genetic and climatic differentiation overlapped to a large extent with genetic and phenotypic differentiation among populations. The largest intersection set, containing 1,102 genes,

was the one containing genes near SNPs associated with all the four variables examined (PC1, PC2, CTmin and CCRT). However, PC2 loads heavily on bioclimatic variables relating to latitude, and our analysis using PC2 as a covariate has a large number of private genes (see figure 4), suggesting

that there is also a substantial amount of genetic variation underlying adaptations to latitude unrelated to the phenotypic measures we have quantified. Our study is an excellent example of how strong ecological selection may be detected in genomic studies. In particular, because the Bayesian

methods examining both ecological variables and relevant phenotypes gave significant overlap amongst the associated loci, and that these are further associated with more broad genomic differentiation between populations, gives confidence that we are consistently identifying genes

associated with ecological selection.

    Analyses of the functional annotation of these genes strengthens our conclusions that climate driven adaptation is important. Regions near SNPs associated with climatic variables were enriched

for genes previously identified as candidates related to cold tolerance, diapause and responses to changes in day length in *D. montana* (Kankare et al., 2010; 2016; Parker et al., 2015; 2016). Thus, genetic variation across populations of these flies may be largely shaped by differences in

ecological and climatic variation. This ecological specialisation may have also contributed to the divergence of *D. montana* from its relatives. Parker et al., (2018) surveyed the rates of molecular evolution in eleven cold tolerant and non-cold tolerant species of Drosophila. The genes found to be

evolving at faster rates in cold-tolerant species were enriched for many of the same functional categories as in our current study, including e.g. membrane and transmembrane proteins and immunoglobulins (Parker et al., 2018).

The fact that many of the same annotation terms are enriched in clusters for all environmental variables in DAVID analyses, suggests that similar genes or biochemical pathways are involved in these adaptations. Membrane proteins and lipids are an important determinant of

membrane and cuticular permeability at different temperatures, which in turn has an effect on the resistance to desiccation stress in insects (Gibbs, 2002; Stanziano, Sové, Rundle, & Sinclair, 2015). Importantly, there is evidence for a close link between the desiccation stress response and cold

tolerance across species and in *Drosophila* in particular, suggesting an overlap in some of the mechanisms involved (Sinclair, Nelson, Nilson, Roberts, & Gibbs, 2007). Cold-hardy lines of *D. melanogaster* are known to exhibit elevated lipid metabolism, perhaps in order to allow rapid lipid

membrane modification (Williams et al., 2016) in different environmental conditions. Pleckstrin homology (PH) domain was one of the functional clusters found in both the current study and in the species comparison of Parker et al., (2018). This domain is a flexible module of 100-120 amino

acids which interacts with a variety of different ligands, composing a protein-protein interaction platform (Scheffzek & Welti, 2012). As changes in the membrane lipid biochemistry form an integral part of the cold tolerance response, genes associated with PH may assist in homeoviscous

adaptation i.e. alteration of membrane phospholipid composition to maintain fluidity at low temperatures (Sinensky, 1974). Interestingly, the only gene found in all five of our previous studies of cold tolerance in *D. montana* was *sidestep II* (*side-II*). Unfortunately, there is no information

available for the biological processes or molecular functions associated with this gene, but *side-II* has protein features including immunoglobulin and immunoglobulin-like domain superfamily (Thurmond et al., 2019) and could be involved in immunological processes during cold response of

the flies (Parker et al., 2021). Insects are known to produce a diverse range of antimicrobial peptides and proteins as part of their immune activity against viruses, bacteria, fungi and parasites (Mylonakis, Podsiadlowski, Muhammed, & Vilcinskas, 2016) and hence immune responses could

be part of a general stress response in cold tolerance (Sinclair, Ferguson, Salehipour-shirazi, & MacMillan, 2013; Ferguson, Heinrichs, & Sinclair, 2016). We also find confirmation of some enriched GO terms in GOwinda analyses that control for variation in gene length, although overlap

is not substantial. In sum, our results indicate that some of the same biochemical processes that are targeted by selection on larger evolutionary scales (i.e. across species), are also involved in local adaptation for different populations within a species, providing a rare bridge between the processes

of population differentiation and speciation.

At the chromosomal level, we found an over-representation of loci associated with ecological selection on chromosomes X (which corresponds to chromosome 2L in *D. melanogaster*) and 4. It

498 is well-known that X chromosomes can generally evolve quickly due to selection on semi-recessive advantageous loci in the hemizygous sex, and smaller effective population size (Charlesworth, Coyne, & Barton, 1987) and are often most divergent between closely related species (Abbott,

501 Norden, & Hansson, 2017; Ellegren et al., 2012). However, there is no obvious reason to expect faster divergence in chromosome 4. *Drosophila montana* populations have considerable chromosomal polymorphism (e.g. Patterson and Stone, 1952; Stone, Guest, & Wilson, 1960;

504 Morales-Hojas, Päällysaho, Vieira, Hoikkala, & Vieira, 2007) and American populations have been classified into several chromosomal forms on the basis of their geographical origin, size and

chromosome structure (Throckmorton, 1982). Moreover, both the X chromosome and chromosome 4 of *D. montana* are known to harbour polymorphic inversions (Stone et al., 1960; Morales-Hojas et al., 2007) and such inversions have often been found to vary clinally (e.g. Kolaczkowski et al., 2011; Cheng et al., 2012; Kapun et al., 2016). Recent genomic data confirms that there are several inversions in both chromosomes X and 4, but we do not yet have any detailed information on their distributions within populations (Poikela et al., unpublished). The reduced recombination within inversions can independently capture advantageous alleles under selection (Kirkpatrick & Barton, 2006) and divergence is often greater in chromosomes carrying inverted regions and non-colinear regions (Lohse, Clarke, Ritchie, & Etges, 2015) and such regions may divergence more quickly during speciation with gene flow. Indeed, here we found that the X chromosome and chromosome 4 always have the highest levels of F_{ST} across all populations, as expected if the ecological selection on loci on these chromosomes influenced overall patterns of genomic diversion, perhaps due to hitchhiking.

Finally, our findings include intriguing result regarding the Crested Butte population which shows reduced genetic variability and a substantial reduction of Tajima's D relative to the other populations. This population occurs at a very high altitude (>2800 m) and also shows reproductive incompatibilities with other populations (Jennings et al., 2014). It may have also been bottlenecked during its adaptation to this high altitude, so population expansion or selected sweeps could be prevalent within this population (Garlovsky et al., 2020). Whether ecological specialisation is associated with the spread of incompatibilities is an intriguing possibility.

CONCLUSION

Identifying the genetic variation that underlies population divergence and ultimately speciation remains a challenge. Detecting associations between genetic and environmental differentiation at loci across populations has become a popular approach. Here we apply Bayesian methods to detect such loci across populations of *Drosophila montana*, which is an extremely cold-tolerant *Drosophila* species where we expect strong ecological selection. We identify many genes that are associated with both climate variables and population-level cold-tolerance phenotypes. These genes overlap with candidate genes from other studies of variation in cold-tolerance in *D. montana* and were also over-represented on chromosomes X and 4 which may be associated with inversion polymorphisms known to be present in these chromosomes. Our study thus provides a clear example of how strong ecological selection can be detected in genome studies, using Bayesian methods to detect local adaptation in combination with studies of ecologically important variables and phenotypes.

Data Availability

Raw reads have been deposited with NCBI (https://www.ncbi.nlm.nih.gov) under the BioProject

543   accession PRJNA588720.

Acknowledgments

546   This work was supported by a combined Natural Environment Research Council and St

Andrews 600th Anniversary PhD Studentship grant (NE/L501852/1) to RAWW, The Ella and

Georg Ehrnrooth Fellowship to VT, NERC grant (NE/P000592/1) to MGR, and Academy of

549   Finland project 267244 to AH and projects 268214 and 322980 to MK. Bioinformatics and

Computational Biology analyses were supported by the University of St Andrews Bioinformatics

Unit which is funded by a Wellcome Trust ISSF award (grant 105621/Z/14/Z). We would also like

552   to thank R. Tapanainen and J. Kinnunen for assistance in lab work and Oscar E Gaggiotti for advice

on BayeScEnv analyses.

Author contributions

AH, MGR, and MK conceived the study. VT performed the phenotyping experiments. RAWW

performed all analyses. RAWW, MK and MGR led the writing of the MS. All authors contributed to

558   the final form of the manuscript.

REFERENCES

561   [dataset] Wiberg, R. A. W., Tyukmaeva, V., Hoikkala, A., Ritchie, M. G. & Kankare, M.; 2020;
        Pooled sequencing of *Drosophila montana* wild population samples; NCBI BioProject;
        PRJNA588720

564   Abbott, J. K., Norden, A. K., & Hansson, B. (2017). Sex chromosome evolution: Historical insights
        and future perspectives. *Proceedings of the Royal Society B: Biological Sciences*, *284*.
        doi: 10.1098/rspb.2016.2806

567   Andersen, J. L., Manenti, T., Sorensen, J. G., MacMillan, H. A., Loeschcke, V., & Overgaard, J.
        (2015). How to assess Drosophila cold tolerance: Chill coma temperature and lower lethal
        temperature are the best predictors of cold distribution limits. *Functional Ecology, 29*, 55–65.

570   doi: 10.1111/1365-2435.12310

        Andrews, S. (2015). FastQC: A quality control tool for high throughput sequence data. Retrieved
        from http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/

573   Arnosti, D.N. (2003). Analysis and function of transcriptional regulatory elements: Insights from
        Drosophila. Annual Review of Entomology. 48, 579-602. doi:
        10.1146/annurev.ento.48.091801.112749

576   Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P., & Marth, G. T. (2011). Bamtools: A C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics*, *27*, 1691–1692. doi: 10.1093/bioinformatics/btr174

579   Barton, N. H., & Gale, K. S. (1993). Genetic analysis of hybrid zones. In R. G. Harrison (Ed.), *Hybrid Zones and the Evolutionary Process*. (pp. 13–45). Oxford, UK: Oxford University Press.

Bauerfeind, S. S., Kellermann, V., Moghadam, N. N., Loeschcke, V., & Fischer, K. (2014).
582   Temperature and photoperiod affect stress resistance traits in *Drosophila melanogaster*. *Physiological Entomology, 39*, 237–246. doi: 10.1111/phen.12068

Bergland, A. O., Behrman, E. L., O'Brien, K. R., Schmidt, P. S., & Petrov, D. (2014). Genomic
585   evidence of rapid and stable adaptive oscillations over seasonal time scales in Drosophila. *PLoS Genetics, 10*, e1004775. doi: 10.1371/journal.pgen.1004775

Berriz, G.F., Beaver, J.E., Cenik, C., Tasan, M., & Roth, F.P. (2009). Next generation software for
588   functional trend analysis. *Bioinformatics*. *25,* 3043-3044. doi:10.1093/bioinformatics/btp498

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics, 30*, 2114–2120. doi: 10.1093/bioinformatics/btu170

591   Bradbury, D., Smithson, A., & Krauss, S. L. (2013). Signatures of diversifying selection at EST-SSR loci and association with climate in natural Eucalyptus populations. *Molecular Ecology, 22,* 5112–5129. doi: 10.1111/mec.12463

594   Broad Institute. Picard. Retrieved from http://broadinstitute.github.io/picard/

Carneiro, M., Baird, S. J. E., Afonso, S., Ramirez, E., Tarroso, P., Teotónio, H., … Ferrand, N. (2013). Steep clines within a highly permeable genome across a hybrid zone between two
597   subspecies of the European rabbit. *Molecular Ecology, 22*, 2511–2525. doi: 10.1111/mec.12272

Chan, Y.F., Marks, M.E., Jones, F.C., Villarreal, G., Shapiro, M.D., Brady, S.D., Southwick, A.M., Absher, D.M., Grimwood, J., Schmutz, J., Myers, R.M., Petrov, D., Jonsson, B., Schluter, D.,
600   Bell, M.A. & Kingsley, D.M. (2010). Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science*. *327* 302-305. doi: 10.1126/science.1182213

Charlesworth, B., Coyne, J. A., & Barton, N. H. (1987). The relative rates of evolution of sex
603   chromosomes and autosomes. *The American Naturalist, 130*, 113–146.

Chen, J., Källman, T., Ma, X., Gyllenstrand, N., Zaina, G., Morgante, M., … Lascoux, M. (2012). Disentangling the roles of history and local selection in shaping clinal variation of allele
606   frequencies and gene expression in Norway spruce (*Picea abies*). *Genetics, 191*, 865–881. doi: 10.1534/genetics.112.140749

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology, 23*, 3133–3157. doi: 10.1111/mec.12796

David, R., Gibert, P., Pla, E., Petavy, G., Karan, D., & Moreteau, B. (1998). Cold stress tolerance in *Drosophila*: Analysis of chill coma recovery in *D. melanogaster. Journal of Thermal Biology, 23*, 291–299. doi: 10.1016/S0306-4565(98)00020-5

Ellegren, H., Smeds, L., Burri, R., Olason, P. I., Backstrom, N., Kawakami, T., … Wolf, J. B. W. (2012). The genomic landscape of species divergence in *Ficedula* flycatchers. *Nature, 491,* 756–760. doi: 10.1038/nature11584.

Endler, J. A. (1973). Gene flow and differentiation. *Science. 179*, 243–250. doi: 10.1126/science.179.4070.243

Endler, J. A. (1977). *Geographic Variation, Speciation, and Clines.* Princeton, NJ: Princeton University Press.

Feder, A. F., Petrov, D. A., & Bergland, A. O. (2012). LDx: Estimation of linkage disequilibrium from high-throughput pooled resequencing data. *PLoS ONE, 7.* doi: 10.1371/journal.pone.0048588

Ferguson, L. V, Heinrichs, D. E., & Sinclair, B. J. (2016). Paradoxical acclimation responses in the thermal performance of insect immunity. *Oecologia, 181*, 77–85. doi: 10.1007/s00442-015-3529-3526.

Findsen, A., Pedersen, T. H., Petersen, A. G., Nielsen, O. B., & Overgaard, J. (2014). Why do insects enter and recover from chill coma? Low temperature and high extracellular potassium compromise muscle function in *Locusta migratoria. The Journal of experimental biology, 217*, 1297–1306. doi: 10.1242/jeb.098442

Foll, M., & Gaggiotti, O. E. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. *Genetics, 180*, 977–993. doi: 10.1534/genetics.108.092221

Foll, M., Gaggiotti, O. E., Daub, J. T., Vatsiou, A., & Excoffier, L. (2014). Widespread signals of convergent adaptation to high altitude in Asia and America. *American Journal of Human Genetics, 95*, 394–407. doi: 10.1016/j.ajhg.2014.09.002.

Garlovsky, M. D., Yusuf, L. H., Ritchie, M. G. & Snook, R. R. (2020). Within-population sperm competition intensity does not predict asymmetry in conpopulation sperm precedence. *Philosophical Transactions of the Royal Society. 375,* 20200071. doi: 10.1098/rstb.2020.0071

Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with population-specific covariates. *Genetics, 201*, 1555–1579. doi: 10.1534/genetics.115.181453

642   Gibbs, A. G. (2002). Lipid melting points and cuticular permeability: New insights into an old
problem. *Journal of Insect Physiology, 48*, 391–400. doi: 10.1016/s0022-1910(02)00059-8

Gibert, P., Moreteau, B., Petavy, G., Karan, D., & David, J. (2001). Chill-coma tolerance, a major

645   climatic adaptation among *Drosophila* species. *Evolution, 55*,1063–1068. doi: 10.1554/0014-
3820(2001)055[1063:cctamc]2.0.co;2

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution

648   interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*,
1965–1978. doi: 10.1002/joc.1276

Hijmans, R. E., van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J. A., … Wueest, R.

651   (2016). raster: Geographic Data Analysis and Modeling. Retrieved from: https://CRAN.R-
project.org/package=raster

Hivert, V., Leblois, R., Petit, E. J., Gautier, M., & Vitalis, R. (2018). Measuring genetic

654   differentiation from pool-seq data. *Genetics*, *210*, 315–330. doi: 10.1534/genetics.118.300900

Hoekstra, H., Drumm, K., & Nachman, M. (2004). Ecological genetics of adaptive color
polymorphism in pocket mice: Geographic variation in selected and neutral genes. *Evolution, 58*,

657   1329–1341. doi: 10.1111/j.0014-3820.2004.tb01711.x

Huang, D. W., Sherman, B. T. and Lempicki, R. A. (2009a). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nature Protocols. 4*, 44-57. doi:

660   10.1038/nprot.2008.211

Huang, D. W., Sherman, B. T. and Lempicki, R. A. (2009b). Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research 37*, 1-

663   13. doi: 10.1093/nar/gkn923

Jennings, J. H., Mazzi, D., Ritchie, M. G., & Hoikkala, A. (2011). Sexual and postmating
reproductive isolation between allopatric *Drosophila montana* populations suggest speciation

666   potential. *BMC Evolutionary Biology*, *11*. doi: 10.1186/1471-2148-11-68

Jennings, J. H., Snook, R. R., & Hoikkala, A. (2014). Reproductive isolation among allopatric
Drosophila montana populations. *Evolution, 68*, 3095–3108. doi: 10.1111/evo.12535

669   Kankare, M., Salminen, T., Laiho, A., Vesala, L., & Hoikkala, A. (2010). Changes in gene
expression linked with adult reproductive diapause in a northern malt fly species: A candidate
gene microarray study. *BMC Ecology*, *10*. doi: 10.1186/1472-6785-10-3

672   Kankare, M., Parker, D. J., Merisalo, M., Salminen, T. S., & Hoikkala, A. (2016). Transcriptional
differences between diapausing and non-diapausing *D. montana* females reared under the same
photoperiod and temperature. *PLoS ONE, 11,* e0161852. doi: 10.1371/journal.pone.0161852

675 Kapun, M., Fabian, D.K., Goudet, J., & Flatt, T. (2016). Genomic evidence for adaptive inversion clines in *Drosophila melanogaster*. *Molecular Biology and Evolution, 33*, 1317–1336. doi: 10.1093/molbev/msw016

678 Kapun, M., Barrón, M. G., Staubach, F., Vieira, J., Obbard, D. J., Goubert, C., … González, J. (2019). Genomic analysis of European *Drosophila melanogaster* populations on a dense spatial scale reveals longitudinal population structure and continent-wide selection, and unknown DNA

681 viruses. *BioRxiv*. doi: 10.1101/313759

Kauranen, H., Kinnunen, J., Hiillos, A-L., Lankinen, P., Hopkins, D., Wiberg, R. A. W., … Hoikkala, A. (2019). Selection for reproduction under short photoperiods changes diapause-

684 associated traits and induces widespread genomic divergence. *Journal of Experimental Biology, 222*, jeb205831. doi:10.1242/jeb.205831

Kellermann, V., Overgaard, J., Hoffmann, A. A., Flojgaard, C., Svenning, J. C., & Loeschcke, V.

687 (2012). Upper thermal limits of Drosophila are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences of the United States of America, 109*, 16228–16233. doi: 10.1073/pnas.1207553109

690 Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and speciation. *Genetics, 173*, 419–434. doi: 10.1534/genetics.105.047985

Klappert, K., Mazzi, D., Hoikkala, A., & Ritchie, M. G. (2007). Male courtship song and female

693 preference variation between phylogeographically distinct populations of *drosophila montana*. *Evolution, 61*, 1481–1488. doi: 10.1111/j.1558-5646.2007.00125.x

Kofler R., & Schlötterer, C. (2012). GOwinda: Unbiased analysis of gene set enrichment for

696 genome-wide association studies. *Bioinformatics. 28,* 2084-2085. doi: 10.1093/bioinformatics/bts315.

Kolaczkowski, B., Kern, A. D., Holloway, A. K., & Begun, D. J. (2011). Genomic differentiation

699 between temperate and tropical Australian populations of *Drosophila melanogaster*. *Genetics, 187*, 245–260. doi: 10.1534/genetics.110.123059

Lakovaara, S. (1969). Malt as a culture medium for Drosophila species. *Drosophila Information*

702 *Service, 44*, 128.

Lankinen, P., Tyukmaeva, V. I., & Hoikkala, A. (2013). Northern *Drosophila montana* flies show variation both within and between cline populations in the critical day length evoking

705 reproductive diapause. *Journal of Insect Physiology, 59*, 745–751. doi: 10.1016/j.jinsphys.2013.05.006

Lê, S., Josse, J., & Husson, F. (2008). FactoMineR: An R package for multivariate analysis. *Journal*

708 *of Statistical Software, 25*, 1–18. doi: 10.18637/jss.v025.i01

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:1303.3997v1 [q-bio.GN] Retrieved from: https://arxiv.org/abs/1303.3997

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... The 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*, 2078–2079. doi: 10.1093/bioinformatics/btp352

Lohse, K., Clarke, M., Ritchie, M. G., & Etges, W. J. (2015). Genome-wide tests for introgression between cactophilic Drosophila implicate a role of inversions during speciation. *Evolution, 69*, 1178–1190. doi: 10.1111/evo.12650

Machado, H. E., Bergland, A. O., O'Brien, K., Behrman, E. L., Schmidt, P. S., & Petrov, D. (2015). Comparative population genomics of latitudinal variation in *Drosophila simulans* and *Drosophila melanogaster. Molecular Ecology, 25*, 723–740. doi: 10.1111/mec.13446

MacMillan, H. A., Williams, C. M., Staples, J. F., & Sinclair, B. J. (2012). Reestablishment of ion homeostasis during chill-coma recovery in the cricket *Gryllus pennsylvanicus. Proceedings of the National Academy of Sciences of the United States of America, 109*, 20750–20755. doi: 10.1073/pnas.1212788109.

Maston, G.A., Evans, S.K. & Green, M.R. 2006. Transcriptional regulatory elements in the human genome. *Annual Review of Genomics and Human Genetics 7,* 29-59. doi: 10.1146/annurev.genom.7.080505.115623

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., ... DePristo, M. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*, 1297–1303. doi:10.1101/gr.107524.110.

Mirol, P.M., Schäfer, M.A., Orsing, L., Routtu, J., Schlötterer, C., Hoikkala, A. & Butlin, R.K. (2007). Phylogeographic patterns in Drosophila montana. *Molecular Ecology 16,* 1085-1097. doi: 10.1111/j.1365-294X.2006.03215.x

Morales-Hojas, R., Päällysaho, S., Vieira, C. P., Hoikkala, A., & Vieira, J. (2007). Comparative polytene chromosome maps of *D. montana* and *D. virilis*. *Chromosoma, 116*, 21–7.doi: 10.1007/s00412-006-0075-3

Mylonakis, E., Podsiadlowski, L., Muhammed, M., & Vilcinskas, A. (2016). Diversity, evolution and medical applications of insect antimicrobial peptides. *Philosophical Transactions of the Royal Society B: Biological Sciences, 371*, 20150290. doi: 10.1098/rstb.2015.0290

Noor, M. A. F., & Bennett, S. M. (2009). Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity, 103*, 439–444. doi: 10.1038/hdy.2009.151

Parker, D. J., Vesala, L., Ritchie, M. G., Laiho, A., Hoikkala, A., & Kankare, M. (2015). How consistent are the transcriptome changes associated with cold acclimation in two species of the *Drosophila virilis* group? *Heredity, 115*, 13–21. doi: 10.1038/hdy.2015.6

Parker, D. J., Ritchie, M. G., & Kankare, M. (2016). Preparing for Winter: The transcriptomic response associated with different day lengths in *Drosophila montana. G3: Genes, Genomes, Genetics, 6*, 1373–1381. doi: 10.1534/g3.116.027870

Parker, D. J., Wiberg, R. A. W., Trivedi, U., Tyukmaeva, V. I., Gharbi, K., Butlin, R. K., ... Ritchie, M. G. (2018). Inter- and intra-specific genomic divergence in *Drosophila montana* shows evidence for cold adaptation. *Genome Biology and Evolution, 10*, 2086–2101. doi:10.1093/gbe/evy147

Parker D.J., Envall T., Ritchie M.G. & Kankare, M. (2021). Sex-specific responses to cold in a very cold-tolerant, northern *Drosophila* species. *Heredty.* https://doi.org/10.1038/s41437-020-00398-2

Patterson J.T. & Stone W.S. (1952). Evolution in the Genus Drosophila. Macmillan, N.Y

Pennacchio, L.A., Bickmore, W., Dean, A., Nobrega, M.A. & Bejerano, G. 2013. Enhancers: five essential questions. *Nature Reviews Genetics. 14* 288-295. doi: 10.1038/nrg3458

Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: Convergence diagnosis and output analysis for MCMC. *R News, 6*, 7–11.

Paolucci, S., Salis, L., Vermeulen, C. J., Beukeboom, L. W., & van de Zande, L. (2016). QTL analysis of the photoperiodic response and clinal distribution of period alleles in *Nasonia vitripennis. Molecular Ecology, 25*, 4805–4817. doi: 10.1111/mec.13802

R Development Core Team. (2020). R: A language and environment for statistical computing. Retrieved from http://www.r-project.org

Ravinet, M., Faria, R., Butlin, R. K., Galindo, J., Bierne, N., Rafajlovic, M., …Westram, A. M. (2017). Interpreting the genomic landscape of speciation: A road map for finding barriers to gene flow. *Journal of Evolutionary Biology, 30*, 1450–1477. doi: 10.1111/jeb.13047

Routtu, J., Mazzi, D., van der Linde, K., Mirol, P., Butlin, R., & Ritchie, M. G. (2007). The extent of variation in male song, wing and genital characters among allopatric *Drosophila montana* populations. *Journal of Evolutionary Biology, 20*, 1591–1601. doi: 10.1111/j.1420-9101.2007.01323.x

Scheffzek, K., & Welti, S. (2012). Pleckstrin homology (PH) like domains – versatile modules in protein–protein interaction platforms. *FEBS Letters, 586*, 2662–2673. doi: 10.1016/j.febslet.2012.06.006

Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals — mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics, 15,* 749–763. doi: 10.1038/nrg3803

Sinclair, B.J., Ferguson, L. V., Salehipour-shirazi, G., & MacMillan, H. A. (2013). Cross-tolerance and cross-talk in the cold: Relating low temperatures to desiccation and immune stress in insects. *Integrative and Comparative Biology, 53,* 545–556. doi: 10.1093/icb/ict004

Sinclair, B. J., Nelson, S., Nilson, T. L., Roberts, S. P., & Gibbs, A. G. (2007). The effect of selection for desiccation resistance on cold tolerance of *Drosophila melanogaster. Physiological Entomology, 32,* 322–327. doi: 10.1111/j.1365-3032.2007.00585.x

Sinensky, M. (1974). Homeoviscous adaptation–A homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli. Proceedings of the National Academy of Sciences of the United States of America, 71,* 522–525. doi: 10.1073/pnas.71.2.522

Stanziano, J. R., Sové, R. J., Rundle, H. D., & Sinclair, B. J. (2015). Rapid desiccation hardening changes the cuticular hydrocarbon profile of *Drosophila melanogaster. Comparative Biochemistry and Physiology, 180,* 38–42. doi: 10.1016/j.cbpa.2014

Stone, W. S., Guest, W. C., & Wilson, F. D. (1960). The evolutionary implications of the cytological polymorphism and phylogeny of the *virilis* group of Drosophila. *Proceedings of the National Academy of Sciences of the United States of America, 46,* 350–361. doi: 10.1073/pnas.46.3.350

Takahashi, Y. (2015). Mechanisms and tests for geographic clines in genetic polymorphisms. *Population Ecology. 57,* 355–362.

Throckmorton, L. H. (1969). The *virilis* Species Group. *The Genetics and Biology of Drosophila, 3,* 795   (pp. 227–296).

Thurmond, J., Goodman, J. L., Strelets, V. B., Attrill, H., Gramates, L. S., Marygold, S. J., … The FlyBase Consortium. (2019). FlyBase 2.0: The next generation. *Nucleic Acids Research, 47,* 798   759–765. doi: 10.1093/nar/gky1003

Tyukmaeva, V. I., Salminen, T. S., Kankare, M., Knott, E. & Hoikkala, A. (2011). Adaptation to a seasonally varying environment: A strong latitudinal cline in reproductive diapause combined 801   with high gene flow in *Drosophila montana. Ecology and Evolution, 1,* 160–168. doi: 10.1002/ece3.14

Tyukmaeva, V. I., Lankinen, P., Kinnunen, J., Kauranen, H., & Hoikkala, A. (2020). Latitudinal 804   clines in the timing and temperature-sensitivity of photoperiodic reproductive diapause in *Drosophila montana. Ecography, 43,* 1–10. doi: 10.1111/ecog.04892

Vesala, L., & Hoikkala, A. (2011). Effects of photoperiodically induced reproductive diapause and cold hardening on the cold tolerance of Drosophila montana. *Journal of Insect Physiology, 57,* 46–51. doi: 10.1016/j.jinsphys.2010.09.007

Vesala, L., Salminen, T. S., Kostal, V., Zahradnickova, H., & Hoikkala, A. (2012). *Myo*-inositol as a main metabolite in overwintering flies: Seasonal metabolomic profiles and cold stress tolerance in a northern drosophilid fly. *Journal of Experimental Biology, 215*, 2891–2897. doi:10.1242/jeb.069948

Vesala, L., Salminen, T. S., Laiho, A., Hoikkala, A., & Kankare M. (2012). Cold tolerance and cold-induced modulation of gene expression in two Drosophila virilis group species with different distributions. *Insect Molecular Biology, 21*, 107–118. doi: 10.1111/j.1365-2583.2011.01119.x

Vigoder, F. M., Parker, D. J., Cook, N., Tournière, O., Sneddon, T., & Ritchie, M. G. (2016). Inducing Cold-Sensitivity in the Frigophilic Fly *Drosophila montana* by RNAi. *PLoS ONE, 11*, 1–9. doi: 10.1371/journal.pone.0165724

Vines, T. H., Dalziel, A. C., Albert, A.Y. K., Veen, T., Schulte, P. M., & Schluter, D. (2016). Cline coupling and uncoupling in a stickleback hybrid zone. *Evolution. 70,* 1023–1038. doi: 10.1111/evo.12917

de Villemereuil, P., & Gaggiotti, O. E. (2015). A new FST-based method to uncover local adaptation using environmental variables. *Methods in Ecology and Evolution, 6*, 1248–1258. doi: 10.1111/2041-210X.12418.

Werner, T., Koshikawa, S., Williams, T.M. & Carroll, S.B. (2010). Generation of a novel wing colour pattern by the Wingless morphogen. *Nature. 464,* 1143-1148. doi: 10.1038/nature08896

Wilder, S. M., Mayntz, D., Toft, S., Rypstra, A. L., Pilati, A., & Vanni, M. J. (2010). Intraspecific variation in prey quality: A comparison of nutrient presence in prey and nutrient extraction by predators. *Oikos, 119*, 350–358. doi: 10.1111/j.1600-0706.2009.17819.x

Williams, C. M., McCue, M. D., Sunny, N. E., Szejner-Sigal, A., Morgan, T. J., Allison, D. B., & Hahn, D. A. (2016). Cold adaptation increases rates of nutrient flow and metabolic plasticity during cold exposure in Drosophila melanogaster. *Proceedings of the Royal Society B: Biological Sciences, 283*, 20161317. doi: 10.1098/rspb.2016.1317

Wolf, J. B. W., & Ellegren, H. (2016). Making sense of genomic islands of differentiation in light of speciation. *Nature Reviews Genetics*, *18*, 87–100. doi: 10.1038/nrg.2016.133

Wood, S.N. (2004) Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association. 99,* 673-686. doi: 10.1198/016214504000000980

840 **Tables, Figures, and Figure Legends**

Table 1. The sources of genomic samples (coordinates and the name of the nearest town), altitude
843 of the sampling site, the year in which sampling was performed, and the number of males and
females sampled (M/F) for each pool.

Source	Sampling Site	Year	M/F
USA, Alaska	Seward 60°9'N; 149°27'W Altitude 35 m	2013	30/20
Canada, British Columbia	Terrace 54°27'N; 128°34'W Altitude 217 m	2014	22/27
USA, Washington	Ashford, 46°45'N; 121°57'W Altitude 573 m	2013	16/34
USA, Colorado	Crested Butte 38°54'N; 106°57'W Altitude 2900 m	2013	36/13
Finland	Oulanka 66°40'N; 29°20'E Altitude 337 m	2013	25/25
Finland	Korpilahti 62°20'N; 25°34'E Altitude 133 m	2013	27/23

Table 2. Number of SNPs with a q-value < 0.05 for the g parameter on each linkage group (chromosomes) and each environmental variable (see Methods). The total number of SNPs on each linkage group are given in the bottom row.

Environmental Variable	LG2	LG3	LG4	LG5	Autosomes Total	X	Overall Total
PC1	561	485	1,432	498	2,976	1,528	4,504
PC2	1,639	1,191	2,480	1,297	6,607	1,861	8,468
CTmin	502	399	1,383	384	2,668	1,272	3,940
CCRT	360	312	1,252	316	2,240	1,228	3,468
Total	441,381	613,952	463,965	415,312	441,381	255,901	2,190,511

Table 3. Mean (SD) F_{ST} values for each linkage group (chromosome) and population

Chromosome	Oulanka	Korpilahti	Ashford	Seward	Crested Butte	Terrace
LG2	0.12 (0.16)	0.12 (0.16)	0.10 (0.15)	0.10 (0.15)	0.15 (0.21)	0.11 (0.16)
LG3	0.11 (0.16)	0.11 (0.16)	0.09 (0.14)	0.09 (0.14)	0.15 (0.21)	0.10 (0.14)
LG4	0.15 (0.18)	0.15 (0.18)	0.14 (0.18)	0.13 (0.17)	0.19 (0.23)	0.13 (0.17)
LG5	0.12 (0.17)	0.13 (0.17)	0.11 (0.15)	0.11 (0.15)	0.16 (0.22)	0.11 (0.16)
X	0.16 (0.21)	0.16 (0.21)	0.14 (0.20)	0.15 (0.20)	0.19 (0.24)	0.14 (0.19)

Figure 1. Maps of all *D. montana* populations in this study. Panels show population locations from

855 **A** Finland and **B** North America showing the locations of all populations sampled. Labelled, blue circles give the locations of populations sampled for phenotyping and sequencing.

858 **Figure 2.** Population phenotypes. **A** and **B** show the variation in CTmin and CCRT across populations and latitude, respectively. Solid and dashed lines show the predicted values from the partial effect of latitude from the best model (see Results) for males and females, respectively. In **B**,

861 although the points are plotted separately for males and females, the best model only included latitude as a covariate. In both panels the points and error bars give the mean and standard errors, respectively.

Figure 3. Principle Components Analysis. **A** The distribution of all populations along the two first PC axes. The loadings of variables on each axis can be found in table S3. Blue circles give the

867 populations that have been pool-sequenced for this study, red squares and triangles give the other Finnish and North American populations, respectively. **B** and **C** give PCs 1 and 2 as a function of latitude and altitude, respectively. The legend for altitude in both **B** and **C** is given in **C**.

Figure 4. Overlap of genes within 10kb of top SNPs from BayeScEnv analyses for PC1, PC2 CTmin, and CCRT. The total number of genes near significant SNPs for each variable are given by

873 the bars in the bottom left (x-axis labelled "Set Size"). The overlaps are depicted by points connected by lines along the *x*-axis and the height of the bars indicate the number of genes in common (intersection) to the variables connected by dots and lines.

Figure 1

 Figure 2

Figure 3.

Figure 4.

Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*

Wiberg et al.

Pseudocode bioinformatic pipeline with model commands:

1. Trimming

java -jar Trimmomatic.jar PE -phred33 *_R1_001.fastq.gz *_R2_001.fastq.gz *_tqc_R1_pe.fq.gz *_ftqc_R1_se.fq.gz *_tqc_R2_pe.fq.gz *_ftqc_R2_se.fq.gz ILLUMINACLIP:TruSeq3-PE.fa:2:30:10:8:TRUE CROP:145 LEADING:20 TRAILING:20 SLIDINGWINDOW:5:20 MINLEN:100;

2. Mapping

bwa mem -t 5 D_montana_genome_freeze_v1.4.fa *_R1_pe.fq.gz *_R2_pe.fq.gz | samtools view -Sb -q 30 -f 0x02 - > *.bam

samtools rmdup *_srt.bam *_srt_rmdup.bam

java -Xmx2g -jar $gatk -T RealignerTargetCreator -R D_montana_genome_freeze_v1.4.fa -I *_srt_rmdup_srt_rdgrp.bam -o *_srt_rmdup_srt_rdgrp.intervals

java -Xmx4g -jar $gatk -T IndelRealigner -R D_montana_genome_freeze_v1.4.fa -I *_srt_rmdup_srt_rdgrp.bam -targetIntervals *_srt_rmdup_srt_rdgrp.intervals -o *_srt_rmdup_srt_rdgrp_indraln.bam

3. Subsample Seward sample

XX=94100000

cat <(samtools view -H Seward_srt.bam) <(samtools view Seward_srt.bam | shuf -n ${XX}) > Seward_subs.sam

samtools view -bS Seward_subs.sam -o Seward_subs.bam

4. SNP calling

samtools mpileup --max-depth 1000000 --skip-indels --min-MQ 20 --min-BQ 15 --fasta-ref
D_montana_genome_freeze_v1.4.fa Oulanka.bam Korpilahti.bam Ashford.bam Seward.bam
Crested_Butte.bamTerrace.bam > montana_clines.mpileup

PoolSNP.sh mpileup=montana_clines.mpileup output=montana_clines
reference=D_montana_genome_freeze_v1.4.fa
names=Oulanka,Korpilahti,Ashford,Seward,Crested_Butte,Terrace min-cov=37 max-cov=0.95
min-count=5 min-freq=0.001 miss-frac=0.01 base-quality=15 jobs=10

5. BayeScEnv
bayescenv *_all_bscanin.txt -burn 2000 -pilot 1000 -npb 5 -n 2000 -env *.txt -od *_bscanout

6. Gowinda
gowinda --output-file *_topsnps_gowinda.tab --gene-definition updownstream10000 --simulations
1000000 --mode gene --gene-set-file funcassociate_go_associations.txt --snp-file all_snps.tab --
candidate-snp-file *_candidate_snps_file.tab --annotation-file
D_montana_genome_freeze_v1.4_dmel.gtf

Final models of population phenotypes
gamm(

      Ctmin ~ s(latitude, bs="cr", k=5) + s(altitude, bs="cr", k=5) + sex ,

      random = list(exp_batch=~1),

      data=cold_pop_dat

)

gamm(

      CCRT ~ s(latitude, bs="cr", k=5),

      random = list(exp_batch=~1),

      data=cold_pop_dat

)

where the term *exp_batch* was the experimental batch random effect.

Supplementary Tables, Figures and Legends

Table S1. See supplementary .tsv file

Table S2. Mapping statistics from merged .bam files. Statistics for Seward samples are given after downsampling of reads.

Sample	Number of reads	Median coverage
Oulanka	78,261,294	62x
Korpilahti	88,285,498	71x
Ashford	82,318,578	66x
Crested Butte	97,443,044	80x
Seward	92,743,585	76x
Terrace	109,266,003	88x

Table S3. Loadings (correlation coefficients) of the 55 bioclimatic variables on principal components (PCs) 1-4. Correlation coefficients > 0.9 are highlighted in bold

Variable	Variable Name	Dim.1	Dim.2	Dim.3	Dim.4
bio1	Annual Mean Temperature	**0.95**	-0.26	0.15	-0.12
bio2	Mean Diurnal Range (Mean of monthly (max temp - min temp))	-0.11	-0.81	-0.56	0
bio3	Isothermality (BIO2/BIO7) (* 100)	0.62	-0.49	-0.56	-0.05
bio4	Temperature Seasonality (standard deviation *100)	-0.9	-0.17	0.2	0.29
bio5	Max Temperature of Warmest Month	0.33	**-0.92**	0.01	0.13
bio6	Min Temperature of Coldest Month	**0.97**	0.08	0.1	-0.13
bio7	Temperature Annual Range (BIO5-BIO6)	-0.8	-0.53	-0.1	0.19
bio8	Mean Temperature of Wettest Quarter	-0.33	-0.07	0.71	0.21
bio9	Mean Temperature of Driest Quarter	0.85	-0.24	-0.26	-0.14
bio10	Mean Temperature of Warmest Quarter	0.55	-0.62	0.51	0.14
bio11	Mean Temperature of Coldest Quarter	**0.97**	-0.06	0	-0.19
bio12	Annual Precipitation	0.81	0.49	-0.21	0.25
bio13	Precipitation of Wettest Month	0.81	0.44	-0.14	0.35
bio14	Precipitation of Driest Month	0.29	0.66	-0.2	-0.56
bio15	Precipitation Seasonality (Coefficient of Variation)	0.55	0.22	0.28	0.63
bio16	Precipitation of Wettest Quarter	0.81	0.43	-0.16	0.36
bio17	Precipitation of Driest Quarter	0.51	0.66	-0.35	-0.33
bio18	Precipitation of Warmest Quarter	-0.23	**0.92**	0.14	-0.12
bio19	Precipitation of Coldest Quarter	0.84	0.35	-0.25	0.33
tmin1	Minimum Temperature – January	**0.97**	0.08	0.11	-0.13
tmin2	Minimum Temperature – February	**0.97**	0.06	0.07	-0.12
tmin3	Minimum Temperature – March	**0.97**	-0.01	0.18	-0.13
tmin4	Minimum Temperature – April	0.91	-0.02	0.37	-0.1
tmin5	Minimum Temperature – May	0.71	0.02	0.68	0.02
tmin6	Minimum Temperature – June	0.39	0.1	0.9	0.12
tmin7	Minimum Temperature – July	0.36	0.08	0.91	0.06
tmin8	Minimum Temperature – August	0.61	0.13	0.77	0
tmin9	Minimum Temperature – September	0.81	0.19	0.55	-0.04
tmin10	Minimum Temperature – October	0.88	0.15	0.35	-0.18
tmin11	Minimum Temperature – November	**0.93**	0.05	0.23	-0.2

tmin12	Minimum Temperature – December	**0.96**	0.11	0.18	-0.15
tmax1	Maximum Temperature – January	**0.91**	-0.21	-0.15	-0.25
tmax2	Maximum Temperature – February	**0.92**	-0.27	-0.18	-0.19
tmax3	Maximum Temperature – March	**0.91**	-0.38	-0.04	-0.12
tmax4	Maximum Temperature – April	0.82	-0.54	0.08	-0.05
tmax5	Maximum Temperature – May	0.58	-0.73	0.25	0.08
tmax6	Maximum Temperature – June	0.25	-0.89	0.26	0.19
tmax7	Maximum Temperature – July	0.31	**-0.92**	0.01	0.12
tmax8	Maximum Temperature – August	0.5	-0.84	-0.09	0.07
tmax9	Maximum Temperature – September	0.68	-0.7	-0.18	0.02
tmax10	Maximum Temperature – October	0.77	-0.57	-0.23	-0.14
tmax11	Maximum Temperature – November	0.86	-0.36	-0.12	-0.26
tmax12	Maximum Temperature – December	0.91	-0.19	-0.09	-0.28
prec1	Precipitation – January	0.83	0.33	-0.26	0.35
prec2	Precipitation – February	0.83	0.35	-0.27	0.32
prec3	Precipitation – March	0.83	0.29	-0.31	0.34
prec4	Precipitation – April	0.85	0.33	-0.3	0.22
prec5	Precipitation – May	0.78	0.36	-0.35	0.06
prec6	Precipitation – June	0.22	0.78	-0.14	0.11
prec7	Precipitation – July	-0.6	0.7	0.21	-0.18
prec8	Precipitation – August	-0.35	0.82	0.25	-0.15
prec9	Precipitation – September	0.37	0.74	0	-0.18
prec10	Precipitation – October	0.72	0.57	-0.02	0.01
prec11	Precipitation – November	0.86	0.36	-0.18	0.3
prec12	Precipitation – December	0.85	0.35	-0.22	0.31

Table S4. See supplementary .xls format spreadsheet (bscan top SNPs)

Table S5. See supplementary .xls format spreadsheet (DAVID results)

Table S6. See supplementary .xls format spreadsheet (GOwinda results)

Table S7. Previous studies to which the results from the current study are compared.

Reference	Study	Geneset being compared
Kankare et al., 2010	Gene expression change during diapause	Genes that significantly changed their expression in response to diapause
Parker et al., 2015	Gene expression response to cold acclimation	Genes that were DE in either *D. montana* or in both *D. montana* and *D. virilis* in response to cold acclimation
Parker et al., 2016	Gene expression response to different day lengths	Genes that responded in expression to day length in either diapausing or non-diapausing flies.
Kankare et al., 2016	Gene expression differences in diapuasing and non-diapausing females	Genes that were either upregulated or downregulated in diapausing compared to non-diapausing females.
Parker et al., 2018	*D. montana* genome: population differentiation among three populations.	Genes that were within 1kb of outlier SNPs in any of three pairwise Fst outlier scans.
Kauranen et al., 2019	Quasinatural selection under shorter day lengths.	Genes within 10kb of a SNP that changed in allele frequency

Table S8. See supplementary .xls format spreadsheet (genes/overlap between studies)

Figure S1. Coverage distributions of all samples before downsampling of Seward data. Note the different scales of the X-axes.

Figure S2. Coverage distributions of merged samples before downsampling of Seward data. Note the different scales of the X-axes.

Figure S3. Coverage distributions of merged samples after downsampling of Seward data. Note the different scales of the X-axes.

Figure S4. Diagnostic gelman and trace plots for BayeScEnv MCMC chains with CTmin as the environmental variable. Analyses were split into **A** autosomal SNPs and **B** X chromosome SNPs. Gelman plots show the Gelman-Rubin shrink factors that summarise variation in parameter estimates across chains. Only data from the three chosen MCMC chains are included in these plots. Trace plots show parameter values across iterations of MCMC chains. These plots show data from all four chains that were run. See supplementary .pdf files.

Figure S5. Diagnostic plots for BayeScEnv MCMC chains with CCRT as the environmental variable. Analyses were split into **A** autosomal SNPs and **B** X chromosome SNPs. Gelman plots show the Gelman-Rubin shrink factors that summarise variation in parameter estimates across chains. Only data from the three chosen MCMC chains are included in these plots. Trace plots show parameter values across iterations of MCMC chains. These plots show data from all four chains that were run. See supplementary .pdf files.

Figure S6. Diagnostic plots for BayeScEnv MCMC chains with PC1 of the bioclimatic PCA as the environmental variable. Analyses were split into **A** autosomal SNPs and **B** X chromosome SNPs. Gelman plots show the Gelman-Rubin shrink factors that summarise variation in parameter estimates across chains. Only data from the three chosen MCMC chains are included in these plots. Trace plots show parameter values across iterations of MCMC chains. These plots show data from all four chains that were run. See supplementary .pdf files.

Figure S7. Diagnostic plots for BayeScEnv MCMC chains with PC2 of the bioclimatic PCA as the environmental variable. Analyses were split into **A** autosomal SNPs and **B** X chromosome SNPs. Gelman plots show the Gelman-Rubin shrink factors that summarise variation in parameter estimates across chains. Only data from the three chosen MCMC chains are included in these plots. Trace plots show parameter values across iterations of MCMC chains. These plots show data from all four chains that were run. See supplementary .pdf files.

Figure S8. "Scree plot" of principal components 1-23 from a Principle Components Analysis of the 55 bioclimatic variables. Main plot shows the eigenvalues of the principal components, the horizontal dashed line shows the eigenvalue threshold of 1. The inset figure shows the cumulative amount of variance explained by the principal components. The horizontal dashed line gives 98% of the cumulative variance.

Figure S9. Scatter plots of the relationship between the population phenotypes (CTmin and CCRT) and the two chosen principal components (PC1 and PC2). PC values and phenotype values have been z-transformed.

Figure S10. Manhattan plot of the q-values from BayeScEnv for a test of genetic differentiation with PC1. Red points show those SNPs with q-values < 0.05. The horizontal dashed line gives the FDR threshold used. Alternating black and grey points along chromosomes denote different scaffolds. Note that scaffolds can be place on chromosomes in this genome assembly but are not yet ordered.

Figure S11. Manhattan plot of the q-values from BayeScEnv for a test of genetic differentiation with PC2. Red points show those SNPs with q-values < 0.05. The horizontal dashed line gives the FDR threshold used. Alternating black and grey points along chromosomes denote different scaffolds. Note that scaffolds can be place on chromosomes in this genome assembly but are not yet ordered.

Figure S12. Manhattan plot of the q-values from BayeScEnv for a test of genetic differentiation with CTmin. Red points show those SNPs with q-values < 0.05. The horizontal dashed line gives the FDR threshold used. Alternating black and grey points along chromosomes denote different scaffolds. Note that scaffolds can be place on chromosomes in this genome assembly but are not yet ordered.

Figure S13. Manhattan plot of the q-values from BayeScEnv for a test of genetic differentiation with CCRT. Red points show those SNPs with q-values < 0.05. The horizontal dashed line gives the FDR threshold used. Alternating black and grey points along chromosomes denote different scaffolds. Note that scaffolds can be place on chromosomes in this genome assembly but are not yet ordered.

Kankare et al., 2010 (N = 14): N overlaps = 8
Bootstrap Ovarlaps = 2.77 [0, 6], p = 0.0004

Parker et al., 2015 (N = 42): N overlaps = 13
Bootstrap Ovarlaps = 6.28 [2, 11], p = 0.0011

Kankare et al., 2016 (N = 3929): N overlaps = 946
Bootstrap Ovarlaps = 773.14 [732, 814], p < 0.001

Parker et al., 2016 (N = 130): N overlaps = 27
Bootstrap Ovarlaps = 18.72 [11, 26], p = 0.014

Parker et al., 2018 (N = 2114): N overlaps = 629
Bootstrap Ovarlaps = 414.7 [382, 448], p < 0.001

Kauranen et al., 2019 (N = 1751): N overlaps = 402
Bootstrap Ovarlaps = 344.64 [314, 375], p < 0.001

Figure S14 (previous page). Bootstrap distributions, from 10,000 bootstrap samples, of overlaps between the union of genes within 10kb of SNPs associated with PC1, PC2, CTmin, ant CCRT and various previous candidate gene sets (see table S4). The red vertical lines give the empirical overlap value. Values above each figure give the citation to the study, the number of genes in the list considered from each study, the number of overlaps, the mean and 95% confidence intervals of the bootstrap distribution, and the empirical p-value (proportion of bootstrap samples with an overlap greater than the empirical overlap).

Figure S15. A nucleotide diversity (π) and **B** Tajima's D as a function of latitude. Inset labels give the names of the corresponding populations.