

1 Digestive parameters during gestation of Holstein heifers

2 Digestion during pregnancy

3

4 Jéssica Marcela Vieira Pereira^{1¶}, Marcos Inácio Marcondes^{1¶}, Sebastião de Campos

5 Valadares Filho^{1&}, Joel Caton^{2¶}, Anna Luiza Lacerda Sguizzato^{1&}, Alex Lopes Silva^{3&},

6 Julia Travassos da Silva^{1&}, Valber Carlos Lima Moraes^{1&}, Lucas Faria Gomes^{1&}, Polyana

7 Pizzi Rotta^{1¶*}

8 ¹ Department of Animal Science, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil

9 ² Department of Animal Science, North Dakota State University, Fargo, North Dakota, USA

10 ³ Department of Animal Production, Universidade Federal Rural do Rio de Janeiro, Seropédica, Rio de
11 Janeiro, Brazil

12

13 * Corresponding author

14 E-mail: polyana.rotta@ufv.br (PPR)

15

16 ¶These authors contributed equally to this work.

17 &These authors also contributed equally to this work.

19 **Abstract**

20 Our objective was to estimate nutrient intake, ruminal flow, total apparent and ruminal
21 digestibility, rates of passage and digestion, ruminal and omasal pH, blood metabolite
22 concentrations, and body measurements during gestation of Holstein heifers. Eleven
23 pregnant Holstein heifers, 8 of which fitted with a rumen cannula (450 ± 27.6 kg of body
24 weight and 20 ± 3.5 months of initial age) were used. All heifers received the same diet
25 composed of corn silage, soybean meal, corn meal, minerals and vitamins, with a corn
26 silage:concentrate ratio of 50:50 (on a dry matter basis), aiming an average daily gain of
27 approximately 1.0 kg. The sampling periods were established according to the days of
28 gestation: 145, 200, and 255 with a duration of 10 days per period. Total fecal samples
29 were collected to estimate dry matter, organic matter, crude protein, and neutral detergent
30 fiber digestibility. Blood samples were collected to analyze metabolites (non-esterified
31 fatty acids, β -hydroxybutyrate, urea, and glucose). Data were analyzed as a repeated
32 measurements scheme, using MIXED procedure, with differences declared when $P <$
33 0.05. Dry matter intake expressed in kg/day increased from d-145 to d-200, and remaining
34 stable until d-255 of gestation. The same results were observed for organic matter and
35 crude protein intake, increasing 15.0 and 35.8% respectively. In contrast, when dry matter
36 intake was evaluated as % body weight, we observed a decrease of 16.7% from d-200 to
37 d-255. Days of gestation did not influence ruminal flow of dry matter, organic matter,
38 crude protein, and neutral detergent fiber. We observed an increase in the ruminal
39 digestibility of neutral detergent fiber by 20.5%. The apparent total-tract digestibility of
40 dry matter, organic matter, and crude protein changed over days of gestation, with an
41 increase of 11.9, 8.5, and 9.8%, respectively, when comparing d-145 with d-200. The rate
42 of digestion of neutral detergent fiber increased from 2.0 to 3.5% h^{-1} . Glucose levels
43 decreased, while β -hydroxybutyrate and non-esterified fatty acids increased from d-145

44 to d-255. In conclusion, our results demonstrate a reduction in dry matter intake in %
45 body weight due to pregnancy. It also shows an increase in total apparent digestibility
46 through gestation, which imply a greater efficiency of use of nutrients by pregnant
47 animals. Thus, further research is still needed to consolidate such results and to elucidate
48 the mechanism about nutrient usage during the final third of gestation in heifers.

49

50 **Introduction**

51 According to the Hoffman [1], heifers should have the first parturition with at least
52 94% of their mature body weight, considering the fetal weight. Thus, their nutritional plan
53 requires strategies to maintain constant and satisfactory gains until the end of gestation
54 [2]. According to Mohd [3] the advances in reproduction were superior to the nutritional
55 advances of heifers, producing precocious animals that may present low milk yield in the
56 first lactation.

57 Literature lacks information on digestive aspects of ruminal flow, intake, passage and
58 digestion rates of pregnant replacement heifers. However, several studies addressing
59 nutrient fluxes of non-pregnant lactating animals are found in the literature [4, 5].
60 Eventually, mathematical models were developed to increase feed efficiency and to
61 estimate rates of passage, digestion, and to estimate outflow of rumen digesta [6, 7, 8],
62 however even these models were not developed or evaluated with pregnant heifers.

63 Data regarding nutrition of dairy cattle during gestation are focused on lactating dairy
64 cows and are scarce for heifers [2, 9, 10]. It is well known that dry matter intake decreases
65 during the final third of gestation due to the greater size of the gravid uterus and the
66 increased estrogen concentration [11, 12]. This decrease in dry matter intake may reach
67 up to 50% in the last 2 months of gestation in dairy cows [12]. However, it is not known
68 whether dry matter intake in heifers present a similar decrease since they are still growing;

69 such information would be of great use to improve management practices in pregnant
70 heifers.

71 The hypothesis of this study was that the digestive parameters of pregnant heifers
72 would be different from data in the literature for pregnant cows and would also be not
73 altered during pregnancy. Thus, the objectives of this study were to estimate ruminal flow,
74 total and partial digestibility of dry matter and diet constituents, intake, passage and
75 digestion rates, ruminal and omasal pH, blood metabolite concentrations, and body
76 measurements during gestation in Holstein heifers.

77

78 **Materials and methods**

79 **Animals and management**

80 All procedures were previously approved by the Animal Ethics and Welfare
81 Committee of the Federal University of Viçosa according to protocol number 020-2016.

82 Eleven pregnant Holstein heifers, 8 of them fitted with a rumen cannula, with a mean
83 body weight of 450 ± 27.6 kg and mean initial age of 20 ± 3.5 months were used in this
84 experiment. The sampling periods were established according to the days of gestation:
85 145, 200, and 255 ± 4.45 , with duration of 10 days for each period. The animals were
86 allocated to individual stalls of 10 m². Prior to the start of the experiment, heifers
87 underwent a 30-day adaptation period to experimental facilities and conditions. Heifers
88 received the same diet (Table 1) for an average daily gain of 0.98 ± 0.30 kg/day, according
89 to NRC (2001).

90 The roughage:concentrate ratio was 50:50 (dry matter basis) and was delivered daily
91 at 0700 and 1500 h. Animals were allowed 5% of orts on fresh basis. Once a week, corn
92 silage was sampled and dried in a ventilated oven at 55°C for 72 h to adjust the diet dry
93 matter content.

94 **Table 1. Ingredients and chemical composition of feed used in the experimental diet.**

Item	% of DM
Ingredients	
Corn silage	50.0
Soybean meal	25.0
Ground corn	21.6
Limestone	1.07
Sodium bicarbonate	0.75
Magnesium oxide	0.25
Salt	0.50
Vitamins	0.76
Mineral mix ¹	0.02
Chemical composition	
Dry matter	56.3
Organic matter	95.2
Crude protein	17.7
Neutral detergent fiber	32.2
Ether extract	2.7
Non fiber carbohydrates	42.6
Indigestible neutral detergent fiber	10.3

¹Mineral mix composition = 0.83 mg/kg of cobalt sulfate, 38.6 mg/kg of copper sulphate, 0.215 g/kg iron sulphate, 2.39 mg/kg of sodium selenite, 0.167 g/kg of zinc sulfate, and 1.71 mg/kg potassium iodate.

95

96 **Intake and digestibility trial**

97 Dry matter intake was daily measured by weighing offered and refused feedstuffs. To
98 estimate nutrients digestibility, total feces were collected from all heifers for 3
99 consecutive days in first period: 145-148 days of gestation, second period: 200-203 days
100 of gestation, and third period 255-258 days of gestation. After 24 h of collection, the total
101 feces were weighed and approximately 250 g was oven-dried (55°C for 72 h). After each
102 collection period, a proportional sample was composed using samples and data from the
103 3 days of fecal collection, based on fecal dry matter production of each day. Then, feces
104 and feed samples were ground using a mill (Willye, model TE-680, TECNAL, Brazil)
105 with a 1- and 2-mm sieves and stored for further chemical analyses.

106

107 **Omasal and ruminal sampling**

108 A continuous infusion of 5 g/day of Co-EDTA (0.7 g cobalt/day) from the 1st to the 6th
109 day of collection was performed during each period (first period: 145-150 days of
110 gestation, second period: 200-205 days of gestation, and third period 255-260 days of
111 gestation) using 2 peristaltic pumps (model BP-600.4, Colombo, Paraná, Brazil). A total
112 of 8 samples per animal were collected from the omasum at a 9-h interval for 3 days;
113 sampling times were 0200, 0500, 0800, 1100, 1400, 1700, 2000, and 2300 h. The
114 technique, that was developed by Huhtanen [13] and adapted by Leão [14], was used for
115 sampling of the omasal digesta, where approximately 250 mL of digesta was used for a
116 bacterial isolate and 500 mL was used for the ruminal flow estimation [15]. The samples
117 were stored in plastic containers at -20 °C for further analysis.

118 At the end of each experimental period, samples of omasal digesta were thawed at
119 room temperature and one composite was created per animal, following the
120 recommendations of Rotta [15]. Thus, 4 L were obtained to estimate the omasal flow,
121 which was filtered using a 100 µm nylon filter with an open area of 44% (Sefar Nytex
122 100/44; Sefar, Thal, Switzerland), thereby obtaining two phases: liquid particles and solid
123 particles. Thus, the double marker system technique was used to estimate the digesta
124 omasal flow, using cobalt and indigestible neutral detergent fiber as markers. All samples
125 were stored in plastic containers and kept at -80° C for further lyophilization (LP510;
126 Liobras, São Paulo, Brazil). After lyophilization, samples were milled at 2- and 1-mm
127 sieves mills (Willye, model TE-048, TECNAL, Brazil).

128 Ruminal contents were sampled during the same times as omasal sampling, resulting
129 in nine samples. Samples were manually collected at the liquid-solid interface of the
130 rumen, filtered through a 100 µm nylon filter (Sefar Nitex; Sefar, Thal, Switzerland;

131 porosity of 50 μm), and were subjected to pH measurement by using a digital
132 potentiometer (pH-221, Lutron Electronics, Taiwan).

133 On the 7th day of collection of each experimental period, total rumen emptying was
134 carried out 4 h after feeding to estimate the rate of passage and digestion [16]. After
135 removal of all ruminal contents, the digesta was weighed and filtered in a double layer of
136 cheesecloth for separation of solid and liquid fractions, which were then weighed and
137 sampled. After sampling, the digesta was reconstituted and returned to the rumen of the
138 respective animals. On the 9th day, the entire rumen emptying procedure was repeated
139 before morning feeding. The collected samples were lyophilized and ground in a knife
140 mill with 1 mm sieve (Willye, model TE-680, TECNAL, Brazil). Thus, composite
141 samples of the dried solid and liquid fractions after emptying the rumen were obtained
142 and were based on the dry weight of each sample.

143

144 **Biometric Measurements**

145 At the end of each experimental period (first period: 154 days of gestation, second
146 period: 209 days of gestation, and third period 264 days of gestation), body weight,
147 withers height, croup height, body length, and thoracic perimeter were measured while
148 the animals were kept in a standing position within the chute.

149

150 **Blood measurements**

151 Blood samples were collected on the 10th day of each period. The samples were
152 collected via jugular venipuncture by using vacuum tubes with clot activator and gel for
153 serum separation (BD Vacutainer® SST® II Advance®, São Paulo, Brazil) in order to
154 quantify urea, nonesterified fatty acids, and β -hidroxibutirate; an extra tube with EDTA
155 and sodium fluoride (BD Vacutainer® Fluorinated/EDTA, São Paulo, Brazil) was used

156 to quantity plasma concentration of glucose. After the collection, samples were
157 centrifuged at $3600 \times g$ for 20 min, and serum and plasma were immediately frozen at -
158 20°C until further analysis.

159 Bioclin® kits were used to quantity urea (K056) and glucose (K082), nonesterified
160 fatty acids, and β -hidroxibutirate were analyzed by using Randox® kits (FA115 and
161 RB1007, Antrim, United Kingdom). All mentioned analyses were conducted using an
162 automated biochemical analyzer (Mindray, BS200E, Shenzhen, China).

163

164 **Laboratory analysis and calculations**

165 Samples of omasal digesta, feces, feed, and ruminal emptying were analyzed for dry
166 matter, organic matter, and nitrogen ([17]; method 934.01 for dry matter, 930.05 for
167 organic matter, and 981.10 for nitrogen, respectively). The ether extract analysis was
168 performed according to AOAC recommendations ([18]; method 945.16). The neutral
169 detergent fiber was analyzed, without the addition of sodium sulphite, but with the
170 addition of detergent thermostable alpha amylase ([19]; INCT-CA method F-001/1). Ash
171 correction was performed in the neutral detergent fiber residues ([19]; method 942.05).
172 Indigestible neutral detergent fiber was estimated according with Valente [20], in
173 triplicate for omasal digesta (particle phase), fecal samples, and rumen emptying samples.
174 Non-fibrous carbohydrates were calculated according to Van Soest [21], with no
175 correction for urea.

176 The total-tract apparent digestibility was calculated as:

$$177 \text{Digestibility (\%)} = \{(x - y)/x\} \times 100,$$

178 where x (kg/day) and y (kg/day) are the intake and the output in feces of each component,
179 respectively.

180 Ruminal digestibility coefficients were estimated by measuring the difference between
181 the rate of nutrient intake and the flow of the nutrients through the rumen. The calculation
182 of intestinal digestibility was estimated by the concentration of nutrients in omasal digesta
183 and feces. The flow of the omasal digesta was estimated through the reconstitution of
184 digesta technique [22] using the double marker system. Cobalt was used as the liquid
185 phase marker and the indigestible neutral detergent fiber as the particle phase marker. The
186 reconstitution factor was calculated based on the concentrations of the markers during the
187 different phases of the digesta [23].

188 The rates of ingestion (k_i), passage (k_p), and digestion (k_d) were calculated using the
189 pool-and-flux method [16], according to the following models:

$$190 \quad k_i = (\text{intake/rumen pool}) \times 100$$

$$191 \quad k_p = (\text{rumen flow/rumen pool}) \times 100$$

$$192 \quad k_d = k_i - k_p$$

193 Where, k_i = ingestion rate of feed fractions (%/h); intake = feed intake (kg/h); rumen pool
194 = amount of total rumen dry matter (kg); k_p = passage rate of feed fractions (%/h); rumen
195 flow = amount of dry matter or nutrients in the omasum (kg/h); k_d = digestion rate of diet
196 fractions (%/h).

197

198 **Statistical analysis**

199 Data were analyzed as a repeated measurements scheme using the SAS MIXED
200 procedure (SAS Institute Inc., version 9.4) according to the statistical model:

$$201 \quad Y_{ijk} = \mu + DG_i + \delta_{ij} + \epsilon_{ijk},$$

202 where Y_{ijk} = dependent variable, μ is the overall mean, DG_i = fixed effect of days of
203 gestation, δ_{ij} the random error with mean of 0 and a variance $\sigma^2 \delta$, the variance between
204 the days of gestation is equal covariance between repeated measures within animals and

205 ϵ_{ijk} is the model error with a mean of 0 and a variance of τ^2 . Days of gestation was
206 considered as repeated measures, and data from ruminal and omasal pH measurements
207 were also included in the statistical model as time-repeated measures (split-plot of DG).
208 The fixed time effects and the interaction between time and days of gestation were tested
209 by using the appropriate covariance structure. The following covariance matrices were
210 tested: components of variance, composite symmetry, heterogeneous composite
211 symmetry, and first order heterogeneous auto-regression. The selection of the covariance
212 matrix was based on the Akaike-corrected criterion, and the composite symmetry matrix
213 was chosen. Results are presented as least squared means and their standard error of the
214 mean. Differences were declared when $P < 0.05$.

215

216 **Results**

217 **Intake, ruminal flow and digestibility**

218 The dry matter intake (kg/day) increased ($P = 0.03$) at d-200 and d-255 compared to
219 d-145 (Table 2). As for nutrient intake, the same pattern was observed, with an increase
220 of 14.7% ($P < 0.01$) for organic matter, 41.7% ($P < 0.01$) for crude protein, at 145 and
221 200 days of gestation, and for neutral detergent fiber there was no difference. Regarding
222 to dry matter intake in % body weight (Fig 1), there was no difference ($P > 0.05$) between
223 145 to 200 days of gestation, but there was a 16.7% ($P < 0.01$) reduction from 200 to 255
224 days of gestation.

225 Ruminal flows of dry matter, organic matter, crude protein, and neutral detergent fiber
226 were not affected by days of gestation (Table 2). The same pattern was observed for
227 ruminal digestibility of dry matter and organic matter, but neutral detergent fiber
228 increased by 46.6% from 145 to 255 days of gestation ($P = 0.03$). The apparent total-tract

229 digestibility of dry matter, organic matter, and crude protein was altered ($P < 0.01$) by
 230 days of gestation, with an increase of 11.9, 8.5, and 9.8%, respectively, when comparing
 231 the 145 and 200 days of gestation. There was no difference (Table 2) for neutral detergent
 232 fiber digestibility among days of gestation.

233

234 **Table 2. Intake (n = 11), ruminal outflow (n = 8), ruminal (n = 8) and apparent total-**
 235 **tract digestibility (n = 11) of Holstein heifers in different days of gestation.**

Item	Day of gestation			P-value
	145	200	255	
Intake, kg/day				
Dry matter	8.4 ± 0.48 ^b	9.7 ± 0.48 ^a	9.0 ± 0.51 ^{ab}	0.03
Organic matter	7.5 ± 0.36 ^b	8.6 ± 0.36 ^a	8.4 ± 0.38 ^a	< 0.01
Crude protein	1.2 ± 0.76 ^b	1.7 ± 0.76 ^a	1.6 ± 0.08 ^a	< 0.01
Neutral detergent fiber	2.8 ± 0.12	2.7 ± 0.12	2.7 ± 0.13	0.64
Ruminal outflow, kg/day				
Dry matter	5.2 ± 0.28	5.5 ± 0.28	5.2 ± 0.31	0.54
Organic matter	4.3 ± 0.24	4.5 ± 0.24	4.3 ± 0.26	0.57
Crude protein	1.0 ± 0.07	1.1 ± 0.07	1.1 ± 0.08	0.57
Neutral detergent fiber	1.2 ± 0.14	1.4 ± 0.14	1.7 ± 0.16	0.08
Ruminal digestibility, %				
Dry matter	54.3 ± 2.64	55.7 ± 2.64	55.7 ± 3.15	0.75
Organic matter	58.6 ± 2.37	58.8 ± 2.37	59.8 ± 2.63	0.89
Neutral detergent fiber	42.4 ± 4.22 ^b	51.1 ± 4.22 ^{ab}	62.1 ± 4.88 ^a	0.03
Apparent total-tract digestibility, %				
Dry matter	72.1 ± 1.35 ^a	79.6 ± 1.35 ^b	79.9 ± 1.49 ^b	< 0.01
Organic matter	75.0 ± 1.29 ^a	81.4 ± 1.29 ^b	82.5 ± 1.43 ^b	< 0.01
Crude protein	75.2 ± 1.21 ^a	82.7 ± 1.21 ^b	82.7 ± 1.33 ^b	< 0.01
Neutral detergent fiber	67.0 ± 2.35	70.4 ± 2.35	73.9 ± 2.59	0.15

236 ^{a-b}Means within a row with different superscripts differ ($P < 0.05$).

237

238 **Fig 1. Dry matter intake (% BW) in different days of gestation of Holstein heifers**
 239 **(number of animals = 11).** * Indicative of significance in the days of gestation ($P < 0.05$)

240

241 **Digestion kinetics**

242 Rumen pools and kinetics data are shown in Table 4. The values for dry matter k_i , k_p ,
 243 and k_d did not differ among days of gestation. The means of dry matter k_i , k_p , and k_d were
 244 8.0, 4.7, and 3.3 % h⁻¹, respectively. The values for neutral detergent fiber k_i and k_p did
 245 not differ among days of gestation, and the means were 4.3 and 2.0 % h⁻¹, respectively.
 246 The k_d of neutral detergent fiber increased from 2.0 to 3.5 % h⁻¹ ($P = 0.01$).

247

248 **Table 4. Digestion kinetics for Holstein heifers in different days of gestation (n = 8).**

Item	Days of gestation			<i>P</i> -value	
	145	200	255		
Rates, % h ⁻¹					
Dry matter	k_i ¹	7.3 ± 0.65	8.4 ± 0.65	8.3 ± 0.72	0.25
	k_p ²	4.5 ± 0.43	4.8 ± 0.43	4.9 ± 0.47	0.57
	k_d ³	2.8 ± 0.36	3.6 ± 0.36	3.4 ± 0.41	0.24
Neutral detergente fiber	k_i	4.8 ± 0.37	4.5 ± 0.37	5.0 ± 0.41	0.20
	k_p	2.2 ± 0.36	2.3 ± 0.36	1.6 ± 0.42	0.22
	k_d	2.0 ± 0.26 ^b	2.6 ± 0.26 ^{ab}	3.5 ± 0.30 ^a	0.01
Rumen pool sizes, kg					
Dry matter		4.9 ± 0.28	5.0 ± 0.28	4.6 ± 0.30	0.36
Neutral detergente fiber		2.5 ± 0.16	2.2 ± 0.16	2.2 ± 0.18	0.14

249 ¹ k_i = rate of ingestion of feed fractions.

250 ² k_p = passage rate of feed fractions.

251 ³ k_d = digestion rate of feed fractions.

252

253 There was no difference ($P > 0.05$) for the ruminal pool of dry matter and neutral
 254 detergent fiber among days of gestation, thereby presenting with an average ruminal pool
 255 of 3.9 kg for dry matter and 2.3 kg for neutral detergent fiber.

256

257 **Ruminal and omasal pH**

258 There was pH interaction ($P = 0.01$) between collection times and days of gestation
 259 for ruminal fluid and omasal fluid ($P < 0.01$) (Figs 2 and 3). Values for ruminal pH
 260 increased ($P < 0.01$) at 0500 h from 145 to 255 days of gestation, the same pattern was

261 observed at 1700, 2000 and 2300 h ($P < 0.01$) with pH values from 5.5 to 6.11, 5.6 to 5.9
 262 and 5.6 to 6.2, respectively. The omasal pH interaction effect was present at 0500, 0800,
 263 1100 and 1400 h. Values of pH at 0500 increased in 10.8 % from 145 to 200 days of
 264 gestation, and the pH values at 0800, 1100 and 1400 h increased 2.6%, 6.1% and 6.0%
 265 from 145 to 255 days of gestation, respectively.

266

267 **Figure 2. pH of ruminal fluid in different days of gestation of Holstein heifers.**

268 Arrows indicate the time of diet delivered at 0700 and 1500 h (number of animals = 8).

269

270 **Figure 3. pH of omasal fluid in different days of gestation of Holstein heifers.** Arrows

271 indicate the time of diet delivered at 0700 and 1500 h (number of animals = 8).

272

273 **Body measurements**

274 The measurements of the teats were not different ($P > 0.05$) among the evaluated
 275 periods, however, we observed that the anterior teats were 15.1% larger than the posterior
 276 teats (Table 3). The measurements of body weight, withers height, croup height, body
 277 length, and thoracic perimeter presented a difference ($P < 0.01$) among periods that were
 278 evaluated with an increase of 23.7, 4.4, 4.2, 23.1 and 11.1% respectively, when we
 279 compare 145 and 255 days of gestation (Table 3).

280

281 **Table 3. Biometric measurements in pregnant Holstein heifers (n = 11).**

Item	Days of gestation			P-value
	145	200	255	
Body weight, kg	469.9 ± 20.0 ^c	539.5 ± 20.0 ^b	580.9 ± 20.3 ^a	< 0.01
Withers height, m	1.35 ± 0.01 ^c	1.39 ± 0.01 ^b	1.41 ± 0.01 ^a	< 0.01
Croup height, m	1.41 ± 0.02 ^c	1.44 ± 0.02 ^b	1.47 ± 0.02 ^a	< 0.01
Body length, m	1.21 ± 0.04 ^b	1.48 ± 0.04 ^a	1.49 ± 0.04 ^a	< 0.01
Thoracic perimeter, m	1.89 ± 0.06 ^b	2.03 ± 0.06 ^a	2.10 ± 0.06 ^a	< 0.01

282 ^{a-b}Means within a row with different superscripts differ ($P < 0.05$).

283

284 **Blood metabolites**

285 Glucose levels decreased ($P = 0.01$) by approximately 6.9% from 145 to 255 days of
286 gestation and the levels of β -hydroxibutirate increased ($P < 0.01$) in the same period by
287 62.9% (Table 5). The urea and nonesterified fatty acids concentration did not differ ($P >$
288 0.05) among days of gestation.

289

290 **Table 5. Metabolic parameters (mmol/L) of blood samples for Holstein heifers in**
291 **different days of gestation (n = 11).**

Item	Days of gestation			P-value
	145	200	255	
Glucose	4.0 \pm 0.05 ^a	3.9 \pm 0.05 ^{ab}	3.8 \pm 0.06 ^b	< 0.01
Non-esterified fatty acids	0.2 \pm 0.04	0.2 \pm 0.04	0.3 \pm 0.05	0.14
Urea	1.7 \pm 0.12	1.8 \pm 0.12	1.8 \pm 0.14	0.49
β -hydroxibutyrate	0.3 \pm 0.04 ^b	0.4 \pm 0.04 ^a	0.4 \pm 0.04 ^a	< 0.01

292 ^{a-b}Means within a row with different superscripts differ ($P < 0.05$).

293

294 **Discussion**

295 There is a lack of information about dry matter intake changes of growing heifers
296 during the last trimester of gestation [9]. To our knowledge this is the first study aiming
297 to evaluate digestive parameters in pregnant Holstein heifers. When comparing dry matter
298 intake (kg/day) at three days of gestation evaluated, we observed an increase of
299 approximately 15.5% between 145 and 200 days of gestation (1.3 kg/day). This might be
300 explained by the fact that heifers are still growing (increasing body measurements) and
301 had an average daily gain around 1.0 kg which is not the case of mature cows, which have

302 a reduction in dry matter intake during pregnancy [12, 24, 25]. Similarly, nutrient intake
303 also showed the same pattern as dry matter in this study.

304 However, we observed that dry matter intake expressed in % body weight was constant
305 at 145 and 200 days of gestation and had a 16.7% decrease at 255 days of gestation, which
306 corresponds to the 37th week of gestation, approximately. This reduction was also
307 reported by Park [26] (25.6%), and Rotta [12] (22.0%) when evaluating dairy cows.
308 However, the reduction of dry matter intake expressed in % body weight, from dairy cows
309 reported was numerically greater than our study with pregnancy heifers. The reduction of
310 dry matter intake might be influenced by increase or decrease in the concentration of
311 hormones such as estrogen and progesterone, respectively [27], as well as the reduction
312 of rumen physical space at the end of gestation [12], which, according to Forbes [28], is
313 the most limiting factor of consumption. This result implies a greater use of feed by the
314 animal.

315 According to Rotta [12] and Park [26], with advancing gestation the total-tract dry
316 matter, organic matter, crude protein and neutral detergent fiber digestibility, has shown
317 a reduction, which in part can be explained by the increase of passage rate [29].
318 Nevertheless, in the present study, we observed that apparent total-tract dry matter,
319 organic matter, crude protein and neutral detergent fiber digestibility increased and no
320 difference was observed for the k_p in the period of gestation evaluated, which explains
321 the non-reduction of total-tract digestibility.

322 Studies evaluating digestion kinetics during gestation of cows are scarce. In our study
323 the rates of gestation, passage and digestion remained constant, with no differences
324 among the days of gestation, excepted for the neutral detergent fiber k_d . Comparing
325 multiparous cows [30,31] observed an increase in the rate of passage during the
326 pregnancy, with decrease nutrient digestibility [32]. However, the pattern observed for

327 pregnant dairy cows was not observed in this study. Factors such as physiological,
328 gastrointestinal, and feed changes should be considered to explain response rates in
329 addition to feed characteristics (particle size) [26] and might have influenced the lack of
330 difference along the days of gestation when these variables were assessed.

331 Further studies should be performed to better elucidate rumen parameters at late
332 gestation in Holstein heifers, so they may be used as the basis for formulating rations to
333 meet the requirements of this category, since the focus in the studies are based on
334 multiparous lactating cows.

335 Considering the concentration of metabolites in dairy cows, studies using lactating
336 cows have been carried out and limits of several metabolites have been established [33,
337 34, 35]. However, according to Briscic [36], reference values for heifers at the end of
338 gestation have not been studied. In this study, values for glucose and urea were similar to
339 those found by Briscic [36]: minimum concentration of 2.8 mmol/L and maximum of 4.0
340 mmol/L for glucose, and for urea the values were between 1.1 mmol/L and 5.8 mmol/L,
341 thus establishing such values as a reference for heifers during late-gestation.

342 Values of β -hidroxibutirate in the present study reached 0.44 mmol/L, which are 3
343 times lower than the values utilized as threshold for ketosis [37]. The values found for
344 nonesterified fatty acids reached 0.34 mmol/L. According to Overton [33], levels above
345 0.5 mmol/L in the prepartum period indicate a greater probability of developing some
346 metabolic disease. Thus, the heifers in the present study with an average daily gain
347 approximately of 1.0 kg are unlikely to present metabolic problems due to the low levels
348 of β -hidroxibutirate and nonesterified fatty acids.

349

350 **Conclusions**

351 Holstein heifers during the last trimester of pregnancy demonstrate an increase in the
352 intake and apparent total-tract digestibility of dry matter, organic matter, and crude
353 protein from 200 days of gestation. Also, there was a reduction of dry matter intake
354 expressed as a percentage of body weight at the end of gestation by 16.7%, reaching
355 1.49% body weight at 250 days of gestation. The digestion kinetics remained constant.
356 Studies evaluating such factors are scarce in the literature, so further research is still
357 needed to consolidate such results.

358

359 **Acknowledgments**

360 The authors thank the Instituto Nacional de Ciências e Tecnologia de Ciência Animal
361 (INCT-CA), Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG),
362 Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Conselho
363 Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil) for financial
364 support.

365

366 **References**

- 367 1. Hofmman PC. Feeding management of the dairy heifer from 4 months to calving.
368 In: Beede DK, editor. Large Dairy Herd Management. Champaign: American
369 Dairy Science Association; 2017. p. 431-444.
- 370 2. Heinrichs AJ, Zanton GI, Lascano GJ, Jones CM. A 100-Year Review: A century
371 of dairy heifer research. J Dairy Sci. 2017; 100:10173–10188.
372 doi:10.3168/jds.2017-12998.

- 373 3. Mohd N, Steeneveld W, van Werven T, Mourits MCM, Hogeveen H. First-calving
374 age and first-lactation milk production on Dutch dairy farms. *J. Dairy Sci.* 2013;
375 96:981–992. doi:10.3168/jds.2012-5741.
- 376 4. Berends H, van den Borne JJGC, Stockhofe-Zurwieden N, Gilbert MS, Zandstra
377 T, Pellikaan WF, et al. Effects of solid feed level and roughage-to-concentrate ratio
378 on ruminal drinking and passage kinetics of milk replacer, concentrates, and
379 roughage in veal calves. *J Dairy Sci.* 2015; 98: 5621–5629. doi:10.3168/jds.2015-
380 9367.
- 381 5. Ramirez-Ramirez HA, Castillo Lopez E, Jenkins CJR, Aluthge ND, Anderson C,
382 Fernando SC, et al. Reduced-fat dried distillers grains with solubles reduces the
383 risk for milk fat depression and supports milk production and ruminal fermentation
384 in dairy cows. *J Dairy Sci.* 2016; 99: 1912–1928. doi:10.3168/jds.2015-9712.
- 385 6. Allen MS, Linton JAV. in *Vivo Methods To Measure Digestibility and Digestion*
386 *Kinetics of Feed Fractions in the Rumen.* Simpósio Int Sobre Avanços em Técnicas
387 *Pesqu Nutr Ruminantes.* 2007; 72–89.
- 388 7. Gregorini P, Beukes P, Waghorn G, Pacheco D, Hanigan M. Development of an
389 improved representation of rumen digesta outflow in a mechanistic and dynamic
390 model of a dairy cow, Molly. *Ecol Modell.* 2015; 313:293-306. doi:
391 10.1016/j.ecolmodel.2015.06.042.
- 392 8. Bannink A, van Lingen HJ, Ellis JL, France J, Dijkstra J. The contribution of
393 mathematical modeling to understanding dynamic aspects of rumen metabolism.
394 *Front Microbiol.* 2016; 7:1820. doi:10.3389/fmicb.2016.01820.
- 395 9. NRC. *Nutrient requirements of dairy cattle.* 7th ed. Washington, D.C.: National
396 Academic Press; 2001.
- 397 10. Akers RM. Mammary development in calves and heifers. In: Beede DK, editor.

- 398 Large Dairy Herd Management. Champaign: American Dairy Science Association;
399 2017. p. 815-827.
- 400 11. Bertoni G, Trevisi E. Use of the liver activity index and other metabolic variables
401 in the assessment of metabolic health in dairy herds. *Vet Clin North Am - Food*
402 *Anim Pract.* 2013; 29: 413–431. doi:10.1016/j.cvfa.2013.04.004
- 403 12. Rotta PP, Filho SCV, Gionbelli TRS, Costa e Silva LF, Engle TE, Marcondes MI,
404 et al. Effects of day of gestation and feeding regimen in Holstein × Gyr cows: I.
405 Apparent total-tract digestibility, nitrogen balance, and fat deposition. *J Dairy Sci.*
406 2015; 98: 3197–3210. doi:10.3168/jds.2014-8280.
- 407 13. Huhtanen P, Brotz PG, Satter LD. Omasal sampling technique for assessing
408 fermentative digestion in the forestomach of dairy cows. *J. Anim Sci.* 1997;
409 75:1380–1392. doi:10.2527/1997.7551380x.
- 410 14. Leão MI, De Campos Valadares Filho S, Rennó LN, Gonçalves LC, Cecon PR,
411 Azevedo JAG, et al. Consumos e digestibilidades totais e parciais de matéria seca,
412 matéria orgânica, proteína bruta e extrato etéreo em novilhos submetidos a três
413 níveis de ingestão e duas metodologias de coleta de digestas abomasal e omasal.
414 *Rev Bras Zootec.* 2004; 33: 1604–1615. doi:10.1590/s1516-35982004000600028.
- 415 15. Rotta PP, Valadares Filho SC, Detmann E, Costa e Silva LF, Paulino MF,
416 Marcondes MI, et al. Digesta sampling sites and marker methods for estimation of
417 ruminal outflow in bulls fed different proportions of corn silage or sugarcane. *J*
418 *Anim Sci.* 2014; 92:2996-3006. doi: 10.2527/jas.2013-7364.
- 419 16. Allen MS, Linton JAV. In vivo methods to measure digestibility and digestion
420 kinetics of feed fractions in the rumen. *Simpósio Intern Avanços Técnicas Pesqu*
421 *Nutr Rumin.* 2007; 72–89.
- 422 17. AOAC International. Official Methods of Analysis. 17th ed. AOAC International,

- 423 Arlington, VA; 2000.
- 424 18. AOAC International. Official Methods of Analysis. 18th ed. AOAC International,
425 Gaithersburg, MD; 2006.
- 426 19. Detmann E, Souza MA, Valadares Filho SC, Queiroz AC, Berchielli TT, Saiba
427 EOS, et al. Methods for feed analysis-INCT-Animal Science. Visconde do Rio
428 Branco, BR: Suprema Gráfica; 2012.
- 429 20. Valente TNP, Detmann E, Queiroz AC, Valadares Filho SC, Gomes DI, Figueiras
430 JF. Evaluation of ruminal degradation profiles of forages using bags made from
431 different textiles. Rev Bras Zootec. Sociedade Brasileira de Zootecnia; 2011; 40:
432 2565–2573. <https://doi.org/10.1590/S1516-35982011001100039>.
- 433 21. Van Soest PJ. Nutritional Ecology of the Ruminant. 2nd ed. Ithaca, NY: Comstock
434 Publishing Associates; 1994.
- 435 22. Faichney GJ. The use of markers to partition digestion within the gastrointestinal
436 tract of ruminants. In: Digestion and metabolism in the ruminant. Proc. IV Int.
437 Symposium Ruminant and Physiology. Sydney, Australia. 1975; p.277-291.
- 438 23. France J, Siddons RC. Determination of digesta flow by continuous marker
439 infusion. J Theor Biol. 1986; 121:105–120. doi: 10.1016/S0022-
440 5193(86)80031-5.
- 441 24. Shonka BN, Tao S, Dahl GE, Spurlock DM. Genetic regulation of prepartum dry
442 matter intake in Holstein cows. J Dairy Sci. 2015; 98: 8195–8200.
443 doi:10.3168/jds.2015-9675
- 444 25. Salin S, Vanhatalo A, Elo K, Taponen J, Boston RC, Kokkonen T. Effects of
445 dietary energy allowance and decline in dry matter intake during the dry period on
446 responses to glucose and insulin in transition dairy cows. J Dairy Sci. 2017; 100:
447 5266–5280. doi:10.3168/jds.2016-11871.

- 448 26. Park AF, Shirley JE, Titgemeyer EC, DeFrain JM, Cochran RC, Wickersham EE,
449 et al. Characterization of ruminal dynamics in Holstein dairy cows during the
450 periparturient period. *J Anim Physiol Anim Nutr (Berl)*. 2011; 95: 571–582.
451 doi:10.1111/j.1439-0396.2010.01085.x.
- 452 27. Ingvarsten KL, Andersen JB. Integration of metabolism and intake regulation: A
453 review focusing on periparturient animals. *J Dairy Sci*. 2000; 83: 1573–1597.
454 doi:10.3168/jds.S0022-0302(00)75029-6
- 455 28. Forbes JM. The physical relationships of the abdominal organs in the pregnant ewe.
456 *J Agric Sci*. 1968; 70:171–177. doi: 10.1017/S0021859600011023.
- 457 29. Beharka A, Cochran B, Harmon D, Avery T. The influence of stage of pregnancy
458 on digestion characteristics of beef cows. *Kansas State Univ. Prog. Rep*. 1988;
459 539:13–15.
- 460 30. Stanley TA, Cochran RC, Vanzant ES, Harmon DL, Corah LR. Periparturient
461 changes in intake, ruminal capacity, and digestive characteristics in beef cows
462 consuming alfalfa hay. *J Anim Sci*. 1993; 71:788–795. doi:
463 10.2527/1993.713788x.
- 464 31. Reynolds CK, Dürst B, Lupoli B, Humphries DJ, Beaver DE. Visceral tissue mass
465 and rumen volume in dairy cows during the transition from late gestation to early
466 lactation. *J Dairy Sci*. 2004;87: 961–971. doi:10.3168/jds.S0022-0302(04)73240-
467 3.
- 468 32. Tyrrell HF, Moe PW. Effect of intake on digestive efficiency. *J Dairy Sci*. 1975;
469 58:1151-1163. doi: 10.3168/jds.S0022-0302(75)84694-7.
- 470 33. Overton TR, McArt JAA, Nydam DV. A 100-Year Review: Metabolic health
471 indicators and management of dairy cattle. *J Dairy Sci*. 2017; 100: 10398–10417.
472 doi:10.3168/jds.2017-13054.

- 473 34. Ospina PA, Nydam DV, Stokol T, Overton TR. Associations of elevated
474 nonesterified fatty acids and β -hydroxybutyrate concentrations with early lactation
475 reproductive performance and milk production in transition dairy cattle in the
476 northeastern United States. *J Dairy Sci.* 2010; 93: 1596–1603.
477 doi:10.3168/jds.2009-2852.
- 478 35. Cozzi G, Ravarotto L, Gottardo F, Stefani AL, Contiero B, Moro L, et al. Short
479 communication: reference values for blood parameters in Holstein dairy cows:
480 Effects of parity, stage of lactation, and season of production. *J Dairy Sci.* 2011;
481 94:3895–3901. doi: 10.3168/jds.2010-3687.
- 482 36. Brscic M, Cozzi G, Lora I, Stefani AL, Contiero B, Ravarotto L, et al. Short
483 communication: Reference limits for blood analytes in Holstein late-pregnant
484 heifers and dry cows: Effects of parity, days relative to calving, and season. *J Dairy*
485 *Sci.* 2015;98: 7886–7892. doi:10.3168/jds.2015-9345.
- 486 37. Duffield TF, Lissemore KD, McBride BW, Leslie KE. Impact of hyperketonemia
487 in early lactation dairy cows on health and production. *J Dairy Sci.* 2009; 92:571–
488 580. doi: 10.3168/jds.2008-1507.




Figure 1




Figure 2




Figure 3