A fungal member of the microbial phyllosphere antagonizes infection of *Arabidopsis thaliana* by the oomycete pathogen *Albugo laibachii* via a putative secreted hydrolase

Katharina Eitzen[1,2], Priyamedha Sengupta[1], Samuel Kroll[2], Eric Kemen*[2,3], Gunther Doehlemann*[1]

[1] *Institute for Plant Sciences and Cluster of Excellence on Plant Sciences (CEPLAS), University of Cologne, Center for Molecular Biosciences, Zuelpicher Str. 47a, 50674 Cologne, Germany.*

[2] *Max Planck Institute for Plant Breeding Research, Carl-von-Linne-Weg 10, 50829 Köln, Germany.*

[3] *Department of Microbial Interactions, IMIT/ZMBP, University of Tübingen, Tübingen, Germany*

* correspondence to:

Eric Kemen, eric.kemen@uni-tuebingen.de,

Gunther Doehlemann, g.doehlemann@uni-koeln.de

Short title: Microbial antagonism suppressing *Albugo laibachii* infection

Key words: microbial antagonism, plant pathogen, Albugo laibachii, Ustilaginales, Arabidopsis, transcriptomics, effectors

Abstract

In natural habitats, plants are not only challenged by pathogenic organisms, but also extensively colonized by commensal microbes. The network of interactions that these microbes establish with their host and amongst each other has been suggested to contribute to the immune responses of plants against pathogens. In wild *Arabidopsis thaliana* populations, the oomycete pathogen *Albugo laibachii* has been shown to play an influential role in structuring the microbial communities of the leaf surface, known as phyllosphere. In this study, we show that the epiphytic yeast *Moesziomyces albugensis,* a close relative of pathogenic smut fungi, plays an antagonistic role in the microbial phyllosphere of *Arabidopsis thaliana*. In particular, *M. albugensis* prevents infection of *A. thaliana* by *A. laibachii*. Through a combination of transcriptome analysis and reverse genetics we identified a gene encoding a putative GH25 hydrolase as the major effector of the microbial antagonism of *M. albugensis*. Our findings broaden the understanding of microbial interactions within the phyllosphere, provide insights into the evolution of epiphytic basidiomycete yeasts and pave the way for the development of novel biocontrol strategies.

Introduction

Plants are colonized by a wide range of microorganisms. While some microbes enter the plant and establish endophytic interactions with a broad range of outcomes from beneficial to pathogenic, plant surfaces harbor a large variety of microbial organisms. Recent research has focused largely on the importance of the rhizosphere microbiota in nutrient acquisition, protection from pathogens, and boosting overall plant growth and development [1–3]. However, the above ground parts of the plant including the phyllosphere are colonized by diverse groups of microbes that also assist in plant protection and immunity [4,5]. The environment has a greater impact on the microbial communities of the leaf surface than on those below-ground, ultimately influencing their interactions with the host [6].

Scale-free network analysis was performed with the leaf microbial population of *Arabidopsis thaliana* [7]. The majority of the interactions in the network were found to be negative, consistent with the fact that are rather the antagonistic interactions that stabilize a microbial community [8]. Phyllosphere network analysis of *A. thaliana* identified a small number of microbes as "hub" organisms, i.e. influential microbes

which have severe effects on the community structure. The major hub microbe in the *A. thaliana* phyllosphere is the oomycete *Albugo laibachii*, which is a pathogenic symbiont biotrophic of Arabidopsis. This pathogen has been shown to significantly reduce the bacterial diversity of epiphytic and endophytic leaf habitats. Since bacteria generally comprise a large proportion of the phyllosphere microbiome [9], phylogenetic profiling of *A. thaliana* was also directed towards identifying a small group of bacteria that frequently colonize *A. thaliana* leaves. The analysis helped to develop a synthetic community of bacteria for experiments in gnotobiotic plants.

Besides bacteria and oomycetes, the microbiota of the *A. thaliana* leaf also comprises a broad range of fungi. Among those fungi, basidiomycete yeasts are frequently found and the most frequent ones are the epiphytic basidiomycete genus *Dioszegia,* as well as an anamorphic yeast classified as *Pseudozyma. sp.* belonging to Ustilaginales [7]. This order includes many pathogens of important crop plants, for example corn smut and loose smut of oats, barley and wheat are caused by *Ustilago maydis*, *U. avenae*, *U. nuda* and *U. tritici*, respectively. Generally, pathogenic development of smut fungi is linked with sexual recombination and plant infection is only initiated upon mating when two haploid sporidia form a dikaryotic filament [10]. Ustilaginales *Pseudozyma* sp. yeasts, however, are not known to be pathogenic and seem to be devoid of sexual recombination [11] and epiphytically colonize a wide range of habitats because of its rapid rate of asexual reproduction. Phylogenetic reconstruction [12] showed that the smut pathogen of millet, *Moesziomyces bullatus* and four species of *Pseudozyma*, namely *P. antarctica*, *P. aphidis*, *P. parantarctica* and *P. rugulosa*form a monophyletic group. The latter does represent anamorphic and culturable stages of *M. bullatus* and, hence, can be renamed and grouped to this genus. *Moesziomyces* strains have been reported in a number of cases to act as microbial antagonists. *Pseudozyma aphidis* inhibited *Xanthomonas campestris* pv. *vesicatoria*, *X. campestris* pv. *campestris*, *Pseudomonas syringae* pv. *tomato*, *Clavibacter michiganensis*, *Erwinia amylovora*, and *Agrobacterium tumefaciens in-vitro* and also led to the activation of induced defense responses in tomato against the pathogen [13]. It was reported that *P. aphidis* can parasitize the hyphae and spores of *Podosphaeraxanthii* [14]. *Pseudozyma churashimaensis* was reported to induce systemic defense in pepper plants against *X. axonopodis*, Cucumber mosaic virus, Pepper mottle virus, Pepper mild mottle virus, and broad bean wilt virus [15].

In the present study, we explored the antagonistic potential of an anamorphic Ustilaginales yeast within the leaf microbial community of *A. thaliana*. We show that *Moesziomyces albugensis* prevents infection by the oomycete pathogen *A. laibachii.* and identified fungal candidate genes that were upregulated in the presence of *A. laibachii*, when both the microbes were co-inoculated in the host plant. A knockout mutant of one of the candidates, which belongs to the glycoside hydrolase – family 25 (GH25), was found to lose its antagonistic abilities towards *A. laibachii,* providing mechanistic insights into fungal-oomycete antagonism within the phyllosphere microbiota. Functional characterization of GH25 will be an important step towards establishing *M. albugensis* as a suitable biocontrol agent.

Results

In a previous study we isolated a basidiomycetous yeast from *Arabidopsis thaliana* leaves infected with the causal agent of white rust, *Albugo laibachii* [7]. This yeast was tightly associated with *A. laibachii* spore propagation. Even after years of subculturing in the lab and re-inoculation of plants with frozen stocks of *A. laibachii* isolate Nc14, this yeast remained highly abundant in spore isolates. Phylogenetic analyses based on fungal ITS-sequencing identified the yeast as *Pseudozyma sp.* Those yeasts can be found across the family of Ustilaginaceae, being closely related to pathogens of monocots like maize, barley, sugarcane or sorghum (Figure 1A and [16]). Microscopic analyses verified the morphological similarity between the putative *Pseudozyma sp.* and the Ustilaginaceous pathogen *Ustilago maydis*, the causal agent of corn smut (Figure 1B; [16]). Based on phylogenetic similarity to the pathogenic smut *Moesziomyces bullatus* which infects millet, several anamorphic *Pseudozyma* isolates were suggested to be renamed and grouped to *M. bullatus* [12]. Since the *Pseudozyma sp.* that was isolated from *A. laibachii* spores groups into the same cluster, we classified this newly identified species as *Moesziomyces albugensis*.

Based on the identification of *M. albugensis* as a hub microbe in the *Arabidopsis* phyllosphere [7], we tested its interaction with 30 bacterial strains from 17 different species of a synthetic bacterial community (SynCom) of *Arabidopsis* leaves in one-to-one plate assays. This experiment identified seven strains being inhibited by *Moesziomyces,* as indicated by halo formation after 7 days of co-cultivation (Supplementary Figure S1). Interestingly, this inhibition was not seen when the

pathogenic smut fungus *U. maydis* was co-cultivated with the bacteria, indicating a
specific inhibition of the bacteria by *M. albugensis* (Supplementary Figure S1).
The primary hub microbe in the *Arabidopsis* phyllosphere was found to be the
pathogenic oomycete *A. laibachii,* which was isolated in direct association with
Moesziomyces [7]. To test if both species interfere with each other, we deployed a
gnotobiotic plate system and quantified *A. laibachii* infection symptoms on
Arabidopsis. In control experiments, spray inoculation of only *A. laibachii* spores on
Arabidopsis leaves led to about 33% infected leaves at 14 dpi (Figure 2). When the
bacterial SynCom was pre-inoculated on leaves two days before *A. laibachii* spores a
significant reduction of *A. laibachii* infection by about 50% was observed (Figure 2).
However, if *Moesziomyces* was pre-inoculated with the bacterial SynCom, *A.*
laibachii spore production was almost completely abolished. Similarly, the pre-
inoculation of only *M. albugensis* resulted in an almost complete loss of *A. laibachii*
infection, independently of the presence of a bacterial community (Figure 2). This
demonstrates that *M. albugensis* holds a strong antagonistic activity towards *A.*
laibachii, resulting in efficient biocontrol of pathogen infection.

117
The genome of *Moesziomyces albugensis.*
Genome sequence of *M. albugensis* was analyzed by Single Molecule Real-Time
sequencing (Pacific Biosciences, Menlo Park, CA), which lead to 69674 mapped
reads with an accuracy of 87.3% and 8596bp sub-read length. Sequence assembly
using the HGAP-pipeline (Pacific Biosciences) resulted in 31 Contigs with a
N_{50}Contig Length of 705kb. The total length of all contigs results in a predicted
genome size of 18.3Mb (Table 1). Gene prediction for the *M. albugensis* genome
with Augustus [17] identified 6653 protein coding genes, of which 559 carry a
secretion signal. Out of these 559, 380 are predicted to be secreted extracellularly
(i.e. they do not carry membrane domains or cell-wall anchors) (Table 1). The small
genome size and high number of coding genes results in a highly compact genome
structure with only small intergenic regions. These are features similarly found in
several pathogenic smut fungi such as *U. maydis* and *S. reilianum* (Table 1).
Remarkably, both *M. albugensis* and *Anthracocystis flocculosa,* which is another
anamorphic and apathogenic yeast, show a similarly high rate of introns, while the
pathogenic smut fungi have a significantly lower intron frequency (Table 1).

To gain better insight in the genome organization of *M. albugensis,* we compared its
structure with the *U. maydis* genome, which serves as a manually annotated high-
quality reference genome for smut fungi [18]. Out of the 31 *M. albugensis* contigs, 21
show telomeric structures and a high synteny to chromosomes of *U. maydis,* with
three of them displaying major events of chromosomal recombination (Figure 3A).
Interestingly, the *Moesziomyces* contig 2, on which also homologs to pathogenic loci
like the *U. maydis* virulence cluster 2A [18] can be found, contains parts of three
different *U. maydis* chromosomes (Chr. 2, 5, 20). The second recombination event
on contig 6 affects the *U. maydis* leaf-specific virulence factor *see1*, which is required
for tumor formation [19]. This recombination event is also found in the genome of the
maize head smut *S. reilianum*, wherein the *U. maydis* chromosomes 5 and 20
recombined in the promoter region of the *see1* gene (Figure 3B). In this respect it
should be noted that *S. reilianum*, although infecting the same host, does not
produce leaf tumors as *U. maydis* does [20]. Therefore, therefore, it is tempting to
speculate that in both *S. relianum* and *M. albugensis,* this chromosomal
recombination has a negative impact on the expression of the orthologs of the
tumorigenesis effector *see1*.
Also the third major recombination event, affecting *M. albugensis* contig 8, changes
the genomic context genes encoding essential virulence factors in *U. maydis* (*stp1* &
pit1/2), as well as the A mating type locus, which is important for pheromone
perception and recognition of mating partners [21]. Based on the strong antibiotic
activities of *M. albugensis,* we mined the genome of *M. albugensis* for the presence
of secondary metabolite gene clusters. Using AntiSMASH, we were able to predict 13
of such clusters, of which three can be assigned to terpene synthesis, three contain
non-ribosomal peptide synthases and one cluster has a polyketide synthetase as
backbone genes (Supplementary Figure S2A). Interestingly, the secondary
metabolite cluster that is involved in the production of the antimicrobial metabolite
ustilagic acid in other Ustilaginomycetes, is absent in *M. albugensis* (Supplementary
Figure S2B). On the contrary, we could identify three *M. albugensis* specific
metabolite clusters which could potentially be involved in the antibacterial activity of
M. albugensis (Supplementary Figure 2C).
Genome comparison of the related Ustilaginales yeast *A. flocculosa* with *U. maydis*
reveals that this anamorphic strain has lost most of its effector genes, reflecting the
absence of a pathogenic stage in this organism [22]. In contrast, *M. albugensis*

contains 1:1 homologs of several known effectors with a known virulence function in
U. maydis (Table 2). To assess the potential virulence activity of *M. albugensis*
effector homologs, we expressed the homolog of the *U. maydis* core effector Pep1 in
an *U. maydis pep1* deletion strain (SG200Δ01987). This resulted in complete
restoration of *U. maydis* virulence, demonstrating that, when ectopically expressed,
M. albugensispep1 encodes a functional effector (Supplementary Figure S3).
A hallmark of the *U. maydis* genome structure is the presence of large clusters with
effector genes, the expression of which is only induced during plant infection [18]. To
assess the presence of potential virulence clusters in *M. albugensis*, we compared all
U. maydis effector gene clusters to the *M. albugensis* genome, based on homology.
This revealed that the twelve major effector clusters of *U. maydis* are present in *M.*
albugensis. However, while many of the clustered effector genes are duplicated in
pathogenic smut fungi, *M. albugensis* carries only a single copy of each effector
gene. This results in the presence of "short" versions of the *U. maydis* gene effector
clusters (Supplemental Figure S4). This gets particularly obvious for the biggest and
most intensively studied virulence cluster of smut fungi, the effector cluster 19A
[20,23,24]. In *M. albugensis* only three out of the 24 effector genes present in *U.*
maydis are conserved in this cluster (Figure 4). Interestingly, some anamorphic
yeasts like *Kalmanozyma brasiliensis* and *A. flocculosa* completely lost virulence
clusters, while another non-pathogenic member of the Ustilaginales, *Pseudozyma*
hubeiensis, shows an almost complete set of effectors when compared to *U. maydis*
(Figure 4).
190
Genetic characterization of *M. albugensis*
To perform reverse genetics in *M. albugensis,* we established a genetic
transformation system based on protoplast preparation and PEG-mediated DNA
transfer. In preliminary transformation assays, we expressed a cytosolic GFP
reporter-gene under control of the constitutive *o2tef*-Promoter (Figure 5A). For the
generation of knockout strains, a split marker approach was used to avoid ectopic
integrations (Figure 5B). To allow generation of multiple knockouts, we used a
selection marker-recycling system (pFLPexpC) which allows selection marker
excision at each transformation round [25].
We decided to apply the transformation system to study the *M. albugensis* mating
type loci in more detail. Although phylogenetically closely related to *U. hordei,* which

has a bi-polar mating system, *M. albugensis* owns a tetrapolar mating system
whereby both mating type loci are physically not linked. This situation is similar to the
mating type structure in the pathogenic smut *U. maydis* (Figure A). The *a*-locus,
which encodes a pheromone – receptor system that is required for sensing and
fusion of compatible cells, is located on contig 6. The *b*-locus can be found on contig
1. This multiallelic mating locus contains two genes (*b-East* and *b-West*), which code
for a pair of homeodomain transcription factors. Upon mating of compatible cells,
pathogenic and sexual development are triggered by a heterodimeric bE/bW complex
[10]. Since the *M. albugensis* genome is completely equipped with mating type
genes, we first deployed a screen for potential mating partners. To this end, we
screened wild *M. bullatus* isolates to find a suitable mating partner, but we could not
observe any mating event (Supplementary Figure S5). To test if *M. albugensis* is able
to undergo pathogenic differentiation in the absence of mating, we generated a self-
compatible strain (CB1) which carries compatible b-mating alleles: to construct the
CB1 strain, we used compatible alleles of the *b-East* and *b-West* genes of the barley
smut *U. hordei*, a pathogen which is the phylogenetically most closely related to *M.*
albugensis and amenable to reverse genetics. The native *M. albugensis* locus was
replaced by the compatible *U. hordei b-East* and *b-West* gene alleles via homologous
recombination (Figure 6B).
Incubation of the *M. albugensis* CB1 on charcoal plates led to the formation of aerial
hyphae with the characteristic fluffy phenotype of filamentous strains like the self-
compatible, solopathogenic *U. maydis* SG200 strain (Figure 6C). A second
established method to induce filament formation in smuts is on hydrophobic parafilm
[26]. Quantification after 18 hours incubation of *M. albugensis* CB1 on parafilm
resulted in the formation of filaments comparable to those of the *U. maydis* SG200
strain (Figure 6D). While about 17% of *M. albugensis* wild type cells showed
filaments, the CB1 strain with compatible b-genes showed 38% filamentous growth.
While the switch from yeast-like growth to filamentous development is the first step in
the pathogenic development of smut fungi, host penetration is accompanied by the
formation of a terminal swelling of infectious hyphae, termed "appressoria". Induction
of appressoria-formation *in vitro* can be induced by adding 100 µM of the cutin
monomer 16-Hydroxyhexadecanoic acid (HDD) to the fungal cells prior to cell
spraying onto a hydrophobic surface [26]. In absence of HDD, only about 8% of the
U. maydis SG200 cells and 14% of the *M. albugensis* cells formed appressoria on

parafilm 24 hours after spraying (Figure 6E). Addition of 100μM HDD resulted in a
significant induction of appressoria in both *U. maydis* and *M. albugensis*,
demonstrating that *M. albugensis* does hold the genetic repertoire to form infection
structures *in-vitro*. Together, the analysis of the recombinant CB1 strain indicates that
M. albugensis can sense pathogenesis-related surface cues and produce penetration
structures to a similar level as that seen for the pathogenic model organism *U.*
maydis.

Identification of microbe-microbe effector genes by RNA-Seq

To study the transcriptomic response of *M. albugensis* to different biotic interactions,
RNA sequencing was performed. The *M. albugensis* transcriptome was profiled in
five different conditions (Figure 7A; cells in axenic culture versus cells on-planta, on-
planta + SynCom, on-planta + *A. laibachii*, on-planta + SynCom + *A. laibachii*).
Inoculations of *A. thaliana* leaves were performed as described above for *A. laibachii*
infection assays (Figure 7A). For *M. albugensis* RNA preparation, the epiphytic
microbes were peeled from the plant tissue by using liquid latex (see methods
section for details).

The libraries of the 15 samples (five conditions in three biological replicates each)
were generated by using a poly-A enrichment and sequenced on an Illumina
HiSeq4000 platform. The paired end reads were mapped to the *M. albugensis*
genome by using Tophat2 [27]. The analysis revealed that *M. albugensis* cells on *A.*
thaliana leaves (on-planta) downregulated 1300 and upregulated 1580 genes
compared to cells in axenic culture (Figure 7B). A GO-terms analysis revealed that,
among the downregulated genes, 50% were associated with primary metabolism
(Supplementary Figure S6). In the two conditions in which *A. laibachii* was present,
we observed upregulation of 801 genes. Among these genes, 411 genes were
specific to co-incubation of *M. albugensis* with *A. laibachii* and SynCom while 174
were specific to incubation with *A. laibachii* only. A set of 216 genes was shared in
both conditions (Figure 7B).

In presence of *A. laibachii,* mainly metabolism- and translation-dependent genes
were upregulated, which might indicate that *M. albugensis* can access a new nutrient
source in presence of *A. laibachii* (Supplementary Figure S6). Among all *A. laibachii*
– induced *M. albugensis* genes, 25 genes encode proteins carrying a secretion signal
peptide and having no predicted transmembrane domain (Figure 7C). After excluding

proteins being predicted to be located in intracellular organelles, nine candidate genes remained as potential microbe-microbe dependent effectors, i.e. *M. albugensis* genes which are induced by *A. laibachii*, show no or low expression in axenic culture and encode for putative secreted proteins (Figure 7C). Interestingly, four of these genes encode putative glycoside hydrolases. Furthermore, two genes encode putative peptidases, one gene likely encodes an alkaline phosphatase and two encode uncharacterized proteins (Figure 7C).

To directly test the eventual antagonistic function of those genes towards *A. laibachii*, we selected the two predicted glycoside hydrolases-encoding genes *g5* & *g2490* (GH43 & GH25) and the gene encoding the uncharacterized protein *g5755* for gene deletion in *M. albugensis.* The respective mutant strains were tested in stress assays to assess, whether the gene deletions resulted in general growth defects. Wild type and mutant *M. albugensis* strains were exposed to different stress conditions including osmotic stress (sorbitol, NaCl), cell wall stress (calcofluor, congored) and oxidative stress (H_2O_2). Overall, in none of the tested conditions we observed a growth defect of the deletion mutants in comparison to wild type *M. albugensis* (Supplementary Figure_S8). To test an eventual impact of the deleted genes in the antagonism of the two microbes, the *M. albugensis* deletion strains were each pre-inoculated on *A. thaliana* leaves prior to *A. laibachii* infection. Deletion of *g5* resulted in a significant but yet marginal increase of *A. laibachii* disease symptoms, while deletion of *g5755* had no effect on *A. laibachii*. We therefore considered these two genes being not important for the antagonism of *M. albugensis* towards *A. laibachii.* Strikingly, the *M. albugensis Δg2490* strain almost completely lost its biocontrol activity towards *A. laibachii.* This phenotype was reproduced by two independents *g2490* deletion strains (Figure 8A). To check if this dramatic loss of microbial antagonism is specific to the deletion of *g2490*, in-locus genetic complementation of strain *Δg2490_1* was performed via homologous recombination. The resulting strain *M. albugensis Δg2490/compl* regained the ability to suppress *A. laibachii* infection, confirming that the observed phenotype specifically resulted from the deletion of the *g2490* gene (Figure 8B*). Together, these results demonstrate that the biocontrol of the pathogenic oomycete *A. laibachii* by the basidiomycete yeast *M. albugensis* is determined by the secretion of a previously uncharacterized GH25 enzyme, which is transcriptionally activated specifically when both microbes are co-colonizing the *A. thaliana* leaf surface.

Discussion

Healthy plants in natural habitats are extensively colonized by microbes, therefore it has been hypothesized that the immune system and the microbiota may instruct each other beyond the simple co-evolutionary arms race between plants and pathogens [28]. Community members as individuals or in a community context have been reported to confer extended immune functions to their plant host. Root endophytic bacteria for example were found to protect *A. thaliana* and stabilize the microbial community by competing with filamentous eukaryotes [29]. A large inhibitory interaction network was found in the leaf microbiome of *A. thaliana* and used genome mining to identify over 1000 predicted natural product biosynthetic gene clusters (BGCs) [30]. In addition, the bacterium *Brevibacillus* sp. leaf 182 was found to inhibit half of the 200 strains isolated from *A. thaliana* phyllosphere. Further analysis revealed that *Brevibacillus* sp. leaf 182 produces a trans-acyltransferase polyketide synthase-derived antibiotic, macrobrevin along with other putative polyketide synthases [30].

In this study, we describe the role of the basidiomycete yeast *M. albugensis,* which we previously co-isolated with the oomycete pathogen *A. laibachii* and now characterized as an antagonistic driver in the *A. thaliana* phyllosphere. *A. laibachii* inhibits *in-vitro* growth of seven members of a bacterial leaf SynCom and, most strikingly, strongly suppresses disease progression and reproduction of the pathogenic oomycete *A. laibachii* on *A. thaliana*. *M. albugensis* is a member of the Ustilaginales, which had previously been classified into the group of pathogenic smut fungi of the *Moesziomyces bullatus* species [11]. Our genome analysis identified the anamorphic yeasts *M. rugulosus, M. aphidis* and *M. antarcticus*, which had previously been classified as "*Pseudozyma spec.*", as the closest relatives of *M. albugensis*. Anamorphic Ustilaginales yeasts are long known and have been used for biotechnological applications and also biocontrol [31]. Mannosylerythritol lipids produced by *M. antarcticus* are known to act as biosurfactants and are of great interest for pharmaceutical applications [32,33]. Glycolipids like flocculosin produced by *A. flocculosa* or ustilagic acid characterized in the smut fungus *U. maydis*, have antifungal activity. Those compounds destabilize the membrane of different fungi and thus serve as biocontrol agents against powdery mildews or grey mould [34–36].

We identified 13 potential secondary metabolite gene clusters in *M. albugensis,* including non-ribosomal peptide synthases and polyketide synthetase. Interaction among microbes within the same habitat is believed to have given rise to a variety of secondary metabolites [37,38]. The presence of *Streptomyces rapamycinicus* was shown to activate an otherwise silent polyketide synthase gene cluster, *fgnA,* in *Aspergillus fumigatus.* The resultant compound proved to be a potent fungal metabolite that inhibited the germination of *S. rapamycinicus* spores [39]. Therefore, secondary metabolite gene clusters and their corresponding products may confer a competitive advantage to fungi over the bacteria that reside in the same environment. What is still under debate is the relation of anamorphic yeasts with the related pathogenic smuts. Many smut fungi, including the model species *U. maydis* are dimorphic organisms. In their saprophytic phase they grow as haploid non-pathogenic yeast cells. Only on appropriate host surfaces, haploid cells switch to filamentous growth and expression of pathogenicity-related genes is only activated upon mating in the filamentous dikaryon. A prime prerequisite for pathogenic development is therefore the ability of mating [40,41]. Our genome analysis identified a tetrapolar mating system with a complete set of mating genes in *M. albugensis.* Looking more closely on the phylogeny of different mating genes it appears that all sequenced *Moesziomyces* strains have the same pheromone receptor type (Supplementary Figure S7). Together with our unsuccessful mating assays, this suggests that all sequenced strains of this species have the same mating type and, therefore, are unable to mate. Mating type bias after spore germination was reported for *Ustilago bromivora,* which leads to a haplo-lethal allele linked to the MAT-2 locus [42]. In this case, an intratetrad mating event rescues pathogenicity in nature as the second mating partner is not viable after spore germination. Together with the observation that anamorphic *Moesziomyces* yeasts are ubiquitous in nature, one could hypothesize that these fungi are highly competitive in their haploid form and antagonism might have led to the selection of one viable mating type. This eventually adapted to the epiphytic life style.

Transcriptome analysis showed that epiphytic growth of *M. albugensis* on *A. thaliana* leads to massive transcriptional changes particularly in primary metabolism, which might reflect adaptation to the nutritional situation on the plant surface. Moreover, *M. albugensis* showed specific transcriptional responses to a bacterial community, as well as to *A. laibachii* when being co-inoculated on plant leaves. Presence of *A.*

laibachii resulted in the induction of primary metabolism and biosynthesis pathways,
which might reflect enhanced growth of *M. albugensis* in the presence of *A. laibachii*.
A set of *M. albugensis* genes encoding secreted hydrolases was induced by *A.*
laibachii and one of these genes which encodes a putative GH25 hydrolase with
similarity to *Chalaropsis* type lysozymes appeared to be essential for the biocontrol of
A. laibachii. Initially discovered in the fungi *Chalaropsis* sp., this group of proteins is
largely present in bacteria as well as phages for example the germination specific
muramidase from *Clostridium perfringens* S40 [43]. The bacterial muramidase,
cellosyl from *Streptomyces coelicolor* [44] also belongs to the *Chalaropsis*type of
lysozyme. These proteins are proposed to cleave the β-1,4-glycosidic bond between
N-acetylmuramic acid (NAM) and N-acetylglucosamine (NAG) in the bacterial
peptidoglycan. Specifically, the β -1,4-N,6-O-diacetylmuramidase activity allows the
Chalaropsis type lysozyme to degrade the cell wall of *Staphylococcus aureus*, in
contrast to the commercially available Hen egg-white lysozyme (HEWL) [44]. The
structure of glycoside hydrolase family 25 from *Aspergillus fumigates* was
characterized and the presence of N-terminal signal peptide was considered to
indicate an extracellular secretion of the protein with possible antimicrobial properties
[45]. The role of the secreted hydrolase in the fungal kingdom is not completely
explored yet. The presence of such hydrolases has in many cases been
hypothesized to be associated with hyperparasitism of fungi parasitizing fungi [46] or
oomycetes parasitizing oomycetes [47]. Our results might therefore indicate a cross
kingdom hyperparasitism event between a fungus and an oomycete. Previous work
on microbial communities has indicated that negative interactions stabilize microbial
communities. Hyperparasitism is such a negative interaction with a strong eco-
evolutionary effect on pathogen-host interactions and therefore on community
stability [48]. *M. albugensis* might therefore regulate *A. laibachii* epidemics and
reduce disease severity. As *A. laibachii* has been shown to reduce microbial diversity
[7], *M. albugensis* might increase diversity through hyperparasitism of *A. laibachii*. At
the same time this increased diversity might have caused the need for more
secondary metabolites to evolve in the *M. albugensis* genome to defend against
niche competitors. Through its close association with *A. laibachii*, *M. albugensis*
could be a key regulator of the *A. thaliana* microbial diversity and therefore relevant
for plant health beyond the regulation of *A. laibachii* infection.

In conclusion, the secreted hydrolase we identified as main factor of *A. laibachii* inhibition has great potential to act as antimicrobial agent. The isolated compound is not only valuable per se in an ecological context. It can further lay the grounds for exploring other microbial bioactive compounds that mediate inter-species and inter-kingdom crosstalk. A main goal of our future studies will be to understand on the mechanistic level, how the GH-25 suppresses *A. laibachii*, and at which developmental step the oomycete infection is blocked. Since the GH-25 enzyme is well conserved amongst Ustilaginales including pathogenic species, it will also be tempting to elucidate whether the species-specific antagonism identified here is broadly conserved among Ustilaginales fungi and oomycetes. Functional investigation of these interactions can provide meaningful insights as to why certain yeasts prefer to colonize specific environments. At the same time, it will be worth exploring how the basidiomycete yeasts influence the bacterial major colonizers of the phyllosphere.

Material and Methods

Strains and growth conditions

M. albugensis wildtype strain was isolated from *A. laibachii* infected *A. thaliana* leaves [7]. Wild-type *M. albugensis* (at 22 degrees) and *U. maydis* (at 28 degrees) strains were grown in liquid YEPSlight medium and maintained on Potato dextrose agar plates. King's B medium was used for culturing Syn Com bacterial members at 22 degrees. All the strains were grown in a rotary shaker at 200rpm. All the recipes for medium and solutions can be found in Supplementary Table S2. Stress assays for fungi: wildtype and mutant strains of *M. albugensis* grown to an optical density (600 nm) of 0.6-0.8 were centrifuged at 3500rpm for 10 minutes and suspended in sterile water to reach an OD of 1.0. Next, a dilution series from 10^0 to 10-4 was prepared in sterile H2O. In the end, 5 µl of each dilution were spotted on CM plates supplemented with the indicated stress agents. The plates were incubated for 2 days at 22°C. Confrontation assays: at first, *M. albugensis* and SynCom bacterial strains were grown to an O.D of 0.8-1. *M. albugensis* cultures (10ul) were dropped in four quadrants of a Potato Dextrose Agar plate, previously spread with a bacterial culture. Plates were incubated for 2-4 days at 22°C.

Transformation of *M. albugensis*

Strains were grown in YEPSL at 22°C in a rotary shaker at 200rpm until an O.D. of

0.6 was reached and centrifuged for 15 mins at 3500rpm. The cells were washed in

20 ml of SCS (Table S2), and further centrifuged for 10 minutes at 3000rpm, before

being treated with 3ml SCS solution with 20mg/ml of Glucanex (Lysing Enzyme from

Trichoderma harzianum, # L1412, Sigma). After 20 minutes of incubation at room

temperature, as cell wall lysis was occurred, cold SCS was added to the mix and

protoplasts spun down for 10 minutes at 2400rpm. They were then washed twice with

SCS and resuspended with 10 ml STC (SupplementaryTable S2) to be centrifuged at

2000rpm for 10 minutes. Finally, the pellet was dissolved in 500 µl STC, and stored

in aliquots of 50 µl at -80°C. 5µg of plasmid DNA along with 15 µg Heparin was

added to 50 µl protoplasts. After incubation on ice for 10 minutes, STC/40%PEG

(500 µl) was added to it and mixed gently by pipetting up and down; this step was

followed by another 15 minutes on ice. The transformation mix was added to 10 ml of

molten regeneration (reg) agar and poured over a layer of already solidified reg agar

containing appropriate antibiotic solution. For the bottom layer, we used 400 µg/ml

Hygromycin/ 8 µg/ml Carboxin/ 300 µg/ml nourseothricin (NAT).

Knockout mutant generation and plasmid construction

Plasmids were cloned using *Escherichia coli* DH5α cells (Invitrogen, Karlsruhe,

Germany). Construction of deletion mutants was performed by homologous

recombination; the 5' and 3' flanking regions of the target genes were amplified and

ligated to an antibiotic resistance cassette [49]. The ligated fragment was

subsequently transformed into *M. albugensis*. Homologous integration of the target

gene was verified via PCR on the antibiotic resistant colonies. Oligonucleotide pairs

for knockout generation and verification can be found in Supplementary Table S3.

PCR amplification was done using Phusion© DNA polymerase (Thermo Scientific,

Bonn, Germany), following the manufacturer's instructions, with 100 ng of genomic

DNA or cDNA as template. Nucleic acids were purified from 1% TAE agarose gels

using Macherey-Nagel™ NucleoSpin™ Gel and PCR Clean-up Kit.

Mating assay and generation of the self-compatible *M. albugensis* strain CB1

Haploid strains of *M. albugensis* were grown in liquid cultures, mixed and drops

arranged on PD-plates with charcoal to induce filament formation. Plate with the

haploid *U. maydis* strains FB1 and FB2 and the solopathogenic strain SG200 served
as internal control.

The complete b-locus of the solopathogenic *U. hordei* strain DS200 was amplified
(Figure S2) and inserted into the *M. albugensis* b-locus by homologous
recombination. The strain obtained, known as compatible b1 (CB1) was tested
positive by amplification of the right border and left border areas with primers specific
for the genomic locus and for the plasmid region. Additionally, two primers specific
for the *M. albugensis bE*and *bW* genes were chosen to amplify parts of the native
locus. To induce filament and appressoria formation in vitro we used a
Moesziomyces YEPSL culture at OD_{600} 0.6-0.8. The cells were diluted to an OD_{600} of
0.2 in 2% YEPSL (for appressoria formation 100µM 16-hydroxyhexadecanoic acid
(Sigma-Aldrich) or 1% ethanol was added) and sprayed the yeast like cells on
parafilm which mimics the hydrophobic plant surface. After 18h incubation at 100%
humidity the number of cells grown as filaments (or generating appressoria) was
determined relative to the total number of total cells by using a light microscope.

Arabidopsis thaliana leaf infections

Sterilized *Arabidopsis thaliana* seeds were subjected to cold treatment for 7 days and
sown on 1/2 strength Murashige Skoog (MS) medium (Supplementary Table S2).
The MS plates are directly transferred to growth chambers having 22°C on a short-
day period (8 h light) with (33-40%) humidity and grown for 4 weeks before
inoculation. Overnight liquid cultures of *M. albugensis* and SynCom bacterial strains
were grown to an OD_{600} of 0.6. The cultures were spun down at 3500rpm for 10
minutes and the pellets dissolved in $MgCl_2$. 500µl of each culture was evenly sprayed
on three-week old *A. thaliana* seedlings using airbrush guns. Two days later, a spore
solution of *A. laibachii* was then sprayed on the seedlings following the protocol of
Ruhe et al. [50]. Two weeks later, the disease symptoms on the leaves were scored
as a percentage between infected and non-infected leaves.

Nucleic acid methods

RNA-Extraction of Latex-peeled samples: Four weeks old *A. thaliana* plants were
fixed between two fingers and liquid latex was applied to the leaf surface by using a
small brush. The latex was dried using the cold air option of a hair dryer, carefully
peeled off with a thin tweezer and immediately frozen in liquid nitrogen. Afterwards,

the frozen latex pieces were grinded with liquid nitrogen and the RNA was isolated by using Trizol® Reagent (Invitrogen, Karlsruhe, Germany) according to the manufacturer's instructions. Turbo DNA-Free™ Kit (Ambion, life technologies™, Carlsbad, California, USA) was used to remove any DNA contamination in the extracted RNA. Synthesis of cDNA was performed using First Strand cDNA Synthesis Kit (Thermo Fischer scientific, Waltham, Massachusetts, USA) according to recommended instruction starting with a concentration of 10µg RNA. QIAprep Mini Plasmid Prep Kit (QIAGEN, Venlo, Netherlands) was used for isolation of plasmid DNA from bacteria after the principle of alkaline lysis. Genomic DNA was isolated using phenol-chloroform extraction protocol [18].

RT-qPCR oligonucleotide pairs were designed with Primer3 Plus. The oligonucleotide pairs were at first tested for efficiency using a dilution series of genomic DNA. The reaction was performed in a Bio-Rad iCycler system using the following conditions: 2 min at 95 °C, followed by 45 cycles of 30 s at 95 °C, 30 s at 61 °C and generation of melting curve between 65°C to 95°C.

Bioinformatics and computational data analysis

Sequence assembly of *M. albugensis* strains was performed using the HGAP pipeline (Pacific Biosciences). *M. albugensis* genome was annotated with the Augustus software tool. Secretome was investigated using SignalP4.0. Analysis of functional domains in the secreted proteins was done by Inter-Pro Scan. AntiSmash was used to predict potential secondary metabolite clusters. RNA sequencing was done at the CCG- Cologne Center for Genomics by using a poly-A enrichment on an Illumina HiSeq4000 platform. The achieved paired end reads were mapped to the *M. albugensis* and *A. thaliana* TAIR10 genome by using Tophat2 [27]. RNA-Seq reads of *M. albugensis* axenic cultures were used to generate exon and intron hints and to start a second annotation with Augustus. Heat-maps were performed using the heatmap.2 function of the package gplots (version 3.0.1) in r-studio (R version 3.5.1). An analysis of variance (ANOVA) model was used for pairwise comparison of the conditions, with Tukey's HSD test to determine significant differences among them (P values <0.05).

Data availability

Genome information and RNA sequencing have been submitted to NCBI Genbank

and are available under the following links:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE148670

Acknowledgments

This work was funded through by the Deutsche Forschungsgemeinschaft (DFG,

German Research Foundation) under Germany's Excellence Strategy EXC-2048/1,

Project ID 390686111, and the DFG priority program SPP2125 "DECRyPT". We are

grateful to Marco Thines for generously providing *M. bullatus* wild type strains. We

thank Libera Lo Presti for critically reading the manuscript and helpful comments and

suggestion.

References

1. Ritpitakphong U, Falquet L, Vimoltust A, Berger A, Métraux JP, L'Haridon F.
The microbiome of the leaf surface of Arabidopsis protects against a fungal
pathogen. New Phytol. 2016. doi:10.1111/nph.13808

2. Walker TS, Bais HP, Grotewold E, Vivanco JM. Root exudation and
rhizosphere biology. Plant Physiology. 2003. doi:10.1104/pp.102.019661

3. Bulgarelli D, Schlaeppi K, Spaepen S, van Themaat EVL, Schulze-Lefert P.
Structure and Functions of the Bacterial Microbiota of Plants. Annu Rev Plant
Biol. 2013. doi:10.1146/annurev-arplant-050312-120106

4. Busby PE, Peay KG, Newcombe G. Common foliar fungi of Populus
trichocarpa modify Melampsora rust disease severity. New Phytol. 2016.
doi:10.1111/nph.13742

5. Mikiciński A, Sobiczewski P, Puławska J, Maciorowski R. Control of fire blight
(Erwinia amylovora) by a novel strain 49M of Pseudomonas graminis from the
phyllosphere of apple (Malus spp.). Eur J Plant Pathol. 2016.
doi:10.1007/s10658-015-0837-y

6. Stone BWG, Weingarten EA, Jackson CR. The Role of the Phyllosphere
Microbiome in Plant Health and Function. Annual Plant Reviews online. 2018.
doi:10.1002/9781119312994.apr0614

7. Agler MT, Ruhe J, Kroll S, Morhenn C, Kim ST, Weigel D, et al. Microbial Hub
Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. PLoS Biol.
2016. doi:10.1371/journal.pbio.1002352

8. Coyte KZ, Schluter J, Foster KR. The ecology of the mCoyte, K.Z., Schluter, J.
& Foster, K.R. (2015) The ecology of the microbiome: Networks, competition,
and stability. Science, 350, 663–666.icrobiome: Networks, competition, and

stability. Science (80-). 2015. doi:10.1126/science.aad2602

9. Vorholt JA. Microbial life in the phyllosphere. Nature Reviews Microbiology.
2012. doi:10.1038/nrmicro2910

10. Brefort T, Doehlemann G, Mendoza-Mendoza A, Reissmann S, Djamei A,
Kahmann R. Ustilago maydis as a Pathogen . Annu Rev Phytopathol. 2009.
doi:10.1146/annurev-phyto-080508-081923

11. Kruse J, Doehlemann G, Kemen E, Thines M. Asexual and sexual morphs of
Moesziomyces revisited. IMA Fungus. 2017.
doi:10.5598/imafungus.2017.08.01.09

12. Wang QM, Begerow D, Groenewald M, Liu XZ, Theelen B, Bai FY, et al.
Multigene phylogeny and taxonomic revision of yeasts and related fungi in the
Ustilaginomycotina. Stud Mycol. 2015. doi:10.1016/j.simyco.2015.10.004

13. Barda O, Shalev O, Alster S, Buxdorf K, Gafni A, Levy M. Pseudozyma aphidis
induces salicylic-acid-independent resistance to Clavibacter michiganensis in
tomato plants. Plant Dis. 2015. doi:10.1094/PDIS-04-14-0377-RE

14. Gafni A, Calderon CE, Harris R, Buxdorf K, Dafa-Berger A, Zeilinger-Reichert
E, et al. Biological control of the cucurbit powdery mildew pathogen
Podosphaera xanthii by means of the epiphytic fungus Pseudozyma aphidis
and parasitism as a mode of action. Front Plant Sci. 2015.
doi:10.3389/fpls.2015.00132

15. Lee G, Lee SH, Kim KM, Ryu CM. Foliar application of the leaf-colonizing yeast
Pseudozyma churashimaensis elicits systemic defense of pepper against
bacterial and viral pathogens. Sci Rep. 2017. doi:10.1038/srep39432

16. Zuo W, Ökmen B, Depotter JRL, Ebert MK, Redkar A, Misas Villamil J, et al.
Molecular Interactions Between Smut Fungi and Their Host Plants. Annu Rev
Phytopathol. 2019. doi:10.1146/annurev-phyto-082718-100139

17. Stanke M, Steinkamp R, Waack S, Morgenstern B. AUGUSTUS: A web server
for gene finding in eukaryotes. Nucleic Acids Res. 2004.
doi:10.1093/nar/gkh379

18. Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, et al. Insights
from the genome of the biotrophic fungal plant pathogen Ustilago maydis.
Nature. 2006. doi:10.1038/nature05248

19. Redkar A, Hoser R, Schilling L, Zechmann B, Krzymowska M, Walbot V, et al.
A secreted effector protein of Ustilago maydis guides maize leaf cells to form
tumors. Plant Cell. 2015. doi:10.1105/tpc.114.131086

20. Schirawski J, Mannhaupt G, Münch K, Brefort T, Schipper K, Doehlemann G,
et al. Pathogenicity determinants in smut fungi revealed by genome
comparison. Science (80-). 2010. doi:10.1126/science.1195330

21. Bölker M, Urban M, Kahmann R. The a mating type locus of U. maydis
specifies cell signaling components. Cell. 1992. doi:10.1016/0092-
8674(92)90182-C

22. Lefebvre F, Joly DL, Labbé C, Teichmann B, Linning R, Belzile F, et al. The
transition from a phytopathogenic smut ancestor to an anamorphic biocontrol

agent deciphered by comparative whole-genome analysis. Plant Cell. 2013.
doi:10.1105/tpc.113.113969

23.　Brefort T, Tanaka S, Neidig N, Doehlemann G, Vincon V, Kahmann R.
Characterization of the Largest Effector Gene Cluster of Ustilago maydis. PLoS
Pathog. 2014. doi:10.1371/journal.ppat.1003866

24.　Dutheil JY, Mannhaupt G, Schweizer G, Sieber CMK, Münsterkötter M,
Güldener U, et al. A tale of genome compartmentalization: The evolution of
virulence clusters in smut fungi. Genome Biol Evol. 2016.
doi:10.1093/gbe/evw026

25.　Khrunyk Y, Münch K, Schipper K, Lupas AN, Kahmann R. The use of FLP-
mediated recombination for the functional analysis of an effector gene family in
the biotrophic smut fungus Ustilago maydis. New Phytol. 2010.
doi:10.1111/j.1469-8137.2010.03413.x

26.　Mendoza-Mendoza A, Berndt P, Djamei A, Weise C, Linne U, Marahiel M, et
al. Physical-chemical plant-derived signals induce differentiation in Ustilago
maydis. Mol Microbiol. 2009. doi:10.1111/j.1365-2958.2008.06567.x

27.　Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2:
Accurate alignment of transcriptomes in the presence of insertions, deletions
and gene fusions. Genome Biol. 2013. doi:10.1186/gb-2013-14-4-r36

28.　Vannier N, Agler M, Hacquard S. Microbiota-mediated disease resistance in
plants. PLoS Pathog. 2019. doi:10.1371/journal.ppat.1007740

29.　Durán P, Thiergart T, Garrido-Oter R, Agler M, Kemen E, Schulze-Lefert P, et
al. Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival.
Cell. 2018. doi:10.1016/j.cell.2018.10.020

30.　Helfrich EJN, Vogel CM, Ueoka R, Schäfer M, Ryffel F, Müller DB, et al.
Bipartite interactions, antibiotic production and biosynthetic potential of the
Arabidopsis leaf microbiome. Nat Microbiol. 2018. doi:10.1038/s41564-018-
0200-0

31.　Boekhout T. Pseudozyma Bandoni emend: Boekhout (1985) and a comparison
with the yeast state of Ustilago maydis (De Candolle) Corda (1842). The
Yeasts. 2011. doi:10.1016/B978-0-444-52149-1.00153-1

32.　Kitamoto D, Akiba S, Hioki C, Tabuchi T. Extracellular accumulation of
mannosylerythritol lipids by a strain of candida antarctica. Agric Biol Chem.
1990. doi:10.1080/00021369.1990.10869918

33.　Morita T, Konishi M, Fukuoka T, Imura T, Kitamoto D. Microbial conversion of
glycerol into glycolipid biosurfactants, mannosylerythritol lipids, by a
basidiomycete yeast, Pseudozyma antarctica JCM 10317T. J Biosci Bioeng.
2007. doi:10.1263/jbb.104.78

34.　Cheng Y, McNally DJ, Labbé C, Voyer N, Belzile F, Bélanger RR. Insertional
mutagenesis of a fungal biocontrol agent led to discovery of a rare cellobiose
lipid with antifungal activity. Appl Environ Microbiol. 2003.
doi:10.1128/AEM.69.5.2595-2602.2003

35.　Mimee B, Labbé C, Pelletier R, Bélanger RR. Antifungal activity of flocculosin,

a novel glycolipid isolated from Pseudozyma flocculosa. Antimicrob Agents
Chemother. 2005. doi:10.1128/AAC.49.4.1597-1599.2005

36.   Teichmann B, Linne U, Hewald S, Marahiel MA, Bölker M. A biosynthetic gene
cluster for a secreted cellobiose lipid with antifungal activity from Ustilago
maydis. Mol Microbiol. 2007. doi:10.1111/j.1365-2958.2007.05941.x

37.   Schroeckh V, Scherlach K, Nützmann HW, Shelest E, Schmidt-Heck W,
Schuemann J, et al. Intimate bacterial-fungal interaction triggers biosynthesis
of archetypal polyketides in Aspergillus nidulans. Proc Natl Acad Sci U S A.
2009. doi:10.1073/pnas.0901870106

38.   Rutledge PJ, Challis GL. Discovery of microbial natural products by activation
of silent biosynthetic gene clusters. Nature Reviews Microbiology. 2015.
doi:10.1038/nrmicro3496

39.   Stroe MC, Netzker T, Scherlach K, Krüger T, Hertweck C, Valiante V, et al.
Targeted induction of a silent fungal gene cluster encoding the bacteria-specific
germination inhibitor fumigermin. Elife. 2020. doi:10.7554/eLife.52541

40.   Bölker M. Ustilago maydis - A valuable model system for the study of fungal
dimorphism and virulence. Microbiology. 2001. doi:10.1099/00221287-147-6-
1395

41.   Nadal M, Gold SE. The autophagy genes atg8 and atg1 affect morphogenesis
and pathogenicity in Ustilago maydis. Mol Plant Pathol. 2010.
doi:10.1111/j.1364-3703.2010.00620.x

42.   Rabe F, Bosch J, Stirnberg A, Guse T, Bauer L, Seitner D, et al. A complete
toolset for the study of Ustilago bromivora and Brachypodium sp. as a fungal-
temperate grass pathosystem. Elife. 2016. doi:10.7554/eLife.20522

43.   Chen Y, Miyata S, Makino S, Moriyama R. Molecular characterization of a
germination-specific muramidase from Clostridium perfringens S40 spores and
nucleotide sequence of the corresponding gene. J Bacteriol. 1997.
doi:10.1128/jb.179.10.3181-3187.1997

44.   Rau A, Hogg T, Marquardt R, Hilgenfeld R. A new lysozyme fold. Crystal
structure of the muramidase from Streptomyces coelicolor at 1.65 Å resolution.
J Biol Chem. 2001. doi:10.1074/jbc.M102591200

45.   Korczynska JE, Danielsen S, Schagerlöf U, Turkenburg JP, Davies GJ, Wilson
KS, et al. The structure of a family GH25 lysozyme from Aspergillus fumigatus.
Acta Crystallogr Sect F Struct Biol Cryst Commun. 2010.
doi:10.1107/S1744309110025601

46.   Hyde KD, Xu J, Rapior S, Jeewon R, Lumyong S, Niego AGT, et al. The
amazing potential of fungi: 50 ways we can exploit fungi industrially. Fungal
Diversity. 2019. doi:10.1007/s13225-019-00430-9

47.   Horner NR, Grenville-Briggs LJ, van West P. The oomycete Pythium
oligandrum expresses putative effectors during mycoparasitism of
Phytophthora infestans and is amenable to transformation. Fungal Biol. 2012.
doi:10.1016/j.funbio.2011.09.004

48.   Parratt SR, Laine AL. The role of hyperparasitism in microbial pathogen

ecology and evolution. ISME Journal. 2016. doi:10.1038/ismej.2015.247

49. Kämper J. A PCR-based system for highly efficient generation of gene replacement mutants in Ustilago maydis. Mol Genet Genomics. 2004. doi:10.1007/s00438-003-0962-8

50. Ruhe J, Agler MT, Placzek A, Kramer K, Finkemeier I, Kemen EM. Obligate biotroph pathogens of the genus albugo are better adapted to active host defense compared to niche competitors. Front Plant Sci. 2016. doi:10.3389/fpls.2016.00820

51. Doehlemann G, Van Der Linde K, Aßmann D, Schwammbach D, Hof A, Mohanty A, et al. Pep1, a secreted effector protein of Ustilago maydis, is required for successful invasion of plant cells. PLoS Pathog. 2009. doi:10.1371/journal.ppat.1000290

52. Schilling L, Matei A, Redkar A, Walbot V, Doehlemann G. Virulence of the maize smut Ustilago maydis is shaped by organ-specific effectors. Mol Plant Pathol. 2014. doi:10.1111/mpp.12133

53. Seitner D, Uhse S, Gallei M, Djamei A. The core effector Cce1 is required for early infection of maize by Ustilago maydis. Mol Plant Pathol. 2018. doi:10.1111/mpp.12698

54. Krombach S, Reissmann S, Kreibich S, Bochen F, Kahmann R. Virulence function of the Ustilago maydis sterol carrier protein 2. New Phytol. 2018. doi:10.1111/nph.15268

55. Schipper K. Charakterisierung eines Ustilago maydis Genclusters, das f€ur drei neuartige sekretierte Effektoren kodiert. PhDitle. PhD thesis. 2009.

56. Doehlemann G, Reissmann S, Aßmann D, Fleckenstein M, Kahmann R. Two linked genes encoding a secreted effector and a membrane protein are essential for Ustilago maydis-induced tumour formation. Mol Microbiol. 2011. doi:10.1111/j.1365-2958.2011.07728.x

57. Ma LS, Wang L, Trippel C, Mendoza-Mendoza A, Ullmann S, Moretti M, et al. The Ustilago maydis repetitive effector Rsp3 blocks the antifungal activity of mannose-binding maize proteins. Nat Commun. 2018. doi:10.1038/s41467-018-04149-0

58. Djamei A, Schipper K, Rabe F, Ghosh A, Vincon V, Kahnt J, et al. Metabolic priming by a secreted fungal effector. Nature. 2011. doi:10.1038/nature10454

59. Ökmen B, Kemmerich B, Hilbig D, Wemhöner R, Aschenbroich J, Perrar A, et al. Dual function of a secreted fungalysin metalloprotease in Ustilago maydis. New Phytol. 2018. doi:10.1111/nph.15265

Supporting information captions

Figure S1: Biocontrol activity of *M. albugensis,* but not *U. maydis,* against bacterial SynCom members [7]. Inhibition by *Moesziomyces* can be seen as a characteristic halo after 48hrs of co-incubation.

Figure S2: (A) Predicted secondary metabolite clusters in the genome of *M. albugensis*. Most clusters have unpredictable functions, three belong to the type of terpene or nonribosomal peptide synthetase types and one is a polyketide synthetase cluster type I. (B) The gene cluster encoding for production of ustilagic acid,a well-studied secondary metabolite of smut fungi [36], is not present in the genome of *M. albugensis*. (C) Out of the 13 predicted secondary metabolite clusters, three are unique to *M. albugensis*. Cluster 2 is predicted to encode a terpene, cluster 8 is a cluster of unknown function and cluster 10 is predicted as NRPS cluster. Core biosynthetic genes are highlighted in red, additional biosynthetic genes in yellow and transport-related genes in blue, based on AntiSMASH predictions.

Figure S3: Protein alignment of the core effector Pep1 (Hemetsberger et al, 2015) from different Ustilaginomycetes. (A) Pep1 regions important for functionality are present in all the aligned sequences. Deletion of the *pep1* gene (UMAG_01987) in *U. maydis* leads to complete loss of virulence, which can be restored by complementing the deletion mutant with the *M. albugensis pep1* gene (Ma1682). Infection of maize leaves was done as described in [19]. Disease symptoms were scored at 12 days post infection in three independent biological replicates. n= number of plants infected

Figure S4: Comparison of known virulence clusters [18] between *U. maydis* and *M. albugensis*. Numbers are gene numbers (*UMAG_NUMBER* for *U. maydis*; *gNUMBER* for *M. albugensis).*

Figure S5: Mating assays of *M. albugensis* and different *M. bullatus* isolates. Mixing

of haploid *U. maydis* strains FB1 & FB2 and the solopathogenic strain SG200 served

as a positive control for mating on charcoal plates, wherein filamentous growth is

indicated by white, fluffy appearance of colonies. Haploid wild type strains without

mating partner don't show a fluffy phenotype. For all combinations of *Moesziomyces*

strains, no mating event resulting in filamentous growth on charcoal plates was

observed.

Figure S6: (A) Sequence distribution of Gene Ontology terms. 60% of all the genes

that are downregulated in *M. albugensis* on plant compared to axenic culture growth

can be assigned to GO-terms related to metabolism and cell cycle. (B) In contrast,

presence of *A. laibachii* leads to transcriptional activation of metabolic processes.

52% of all GO-terms associated with genes upregulated in presence of *A. laibachii*

are related to metabolic processes.

Figure S7: A molecular phylogenetic analysis using maximum likelihood estimation

and based on pheromone receptor protein sequences similarity. *M. albugensis*

protein sequence clusters together with type 1 pheromone receptors of other

Ustilaginomycetes.

Figure S8: Stress assays of *M. albugensis* wild type and knockout mutants of genes

g5, g5755 and g2490, respectively, on CM medium and 2% Glucose (A)

supplemented with different stress agents (B: 100 µg/ml Calcofluor; C: 150 µg/ml

Calcoflour; D: 1 mM H_2O_2; E:45 µg/ml Congored; F: 1 M NaCl; G: 1 M Sorbitol). The

strains were dropped on the different CM plates in a dilution series from 10^0 to 10^{-4}

Table S1: *Moesziomyces albugensis* gene expression data

Table S2: Growth media and buffers used in this study

Table S3: PCR primers used in this study

 Figure Legends

Figure 1: Position of *M. albugensis* in the family of Ustilaginaceae. (A) molecular

phylogenetic analysis by maximum likelihood method based on fungal ITS

sequences and showing grouping of *M. albugensis* with the millet pathogen *M.*

bullatus. Pathogenic (green) and apathogenic species can be found across the

phylogenetic tree. (B) Comparison of the morphology of *M. albugensis* (bottom) and

U. maydis (top) haploid cells grown in YEPSLight medium to OD_{600}0.6.

Figure 2: Infection assay of *A. laibachii* on *A. thaliana*. Addition of a bacterial

SynCom reduces the infection symptoms of *A. laibachii* at 14dpi. Those symptoms

can be almost abolished by spraying *M. albugensis* to the plant, independently of the

presence of the bacterial community. An analysis of variance (ANOVA) model was

used for pairwise comparison of the conditions, with Tukey's HSD test to determine

significant differences among them (P values <0.05).

Figure 3:Circos comparison of *M. albugensis* and *U. maydis* chromosome structure

(A). We highlighted potential secondary metabolite clusters, secreted proteins and

gene predictions on both strands (+/-). (B) Homology based comparisons identified

three chromosomal recombination events, which affects the *M. albugensis* contigs 2,

6 & 8.

Figure 4: Structure of the largest virulence cluster (Cluster 19A) in pathogenic and

non-pathogenic smut fungi. Colors indicate genes with homology to each other:

Related gene families are indicated in orange, yellow, blue, green and brown,

whereas unique effector genes are shown in grey. Genes encoding proteins without

a secretion signal are shown in white [23].

Figure 5: Genetic transformation of *M. albugensis.* (A) Stable transformants that

express cytosolic GFP could be obtained by generating protoplasts with Glucanex

and ectopically integrating linear DNA-fragments into the genome via PEG-mediated

transformation. (B) Overview of the split-marker approach that was used to generate
deletion mutants via homologous recombination.

Figure 6: The self-compatible *M. albugensis* strain CB1 (A) *M. albugensis* mating
type genes, unlike the ones of *U. hordei*, can be found on two different chromosomes
similar to the tetrapolar mating type system of *U. maydis*. (B) To generate a self-
compatible strain (CB1), the b-mating genes of *U.hordei* were integrated at the native
M. albugensis b-locus. (C) Unlike the *M. albugensis* wild type strain (top left), strain
CB1 (bottom left) shows a fluffy phenotype on charcoal plates and filamentous
growth. *U. maydis* haploid F1 strain (top right) and self-compatible SG200 strain
(bottom right) were used as negative and positive control, respectively. (D,E)
Induction of filamentation and appressoria formation in strain CB1. After incubation
on a hydrophobic surface, both, filament and appressoria formation in strain CB1
were significantly different (* Chi-Square Test for Independence – p = 0,0001) when
compared to *M. albugensis* wild type and similar to the level of the self-compatible *U.*
maydis strain SG00. *U. maydis* haploid F2 strain was used as negative control [18].
Scale bar: 20µm.

Figure 7: Transcriptome analysis of *M. albugensis* (A) Experimental setup used for
the transcriptomic (RNA-Sequencing) analysis in *M. albugensis*. (B) Venn diagrams
showing differential regulated *M. albugensis* genes after spraying of haploid cells
onto the *A. thaliana* leaf surface. A total number of 801 genes were upregulated in
response to *A. laibachii* in presence and absence of bacterial SynCom. 216 of the
801 genes were upregulated in both conditions. (C) Hierarchical clustering of the 27
A. laibachii - induced *M. albugensis* genes that are predicted to encode secreted
proteins. Of these genes, nine were selected as candidate microbe-microbe effector
genes, based on their transcriptional upregulation and prediction to encode for
extracellularly localized proteins.

Figure 8: A reverse-genetic approach to identify the *M. albugensis* gene which is
responsible for the suppression of *A. laibachii* infection. (A) Three candidate microbe-
microbe effector genes (*g5*, *g5755* &*g2490*) were deleted in *M. albugensis* and

deletion strains were individually inoculated on *A. thaliana* together with *A. laibachii*.

Inoculation of two independent *g2490 null* strains *(Δg2490_1; Δg2490_2)* resulted in

significant and almost complete loss the biocontrol activity of *M. albugensis*. While

deletion of g5 resulted in a marginal reduction of disease symptoms at 14 days post

infection, deletion of g5755 had no effect on *A. laibachii.* (B) Genetic

complementation of the *g2490* deletion restores the biocontrol activity to wild type

levels.

Tables

Table 1: Comparison of Genomes and genomic features of known pathogenic and apathogenic Ustilaginomycetes.

	M. albugensis	*U. bromivora*	*S. scitamineum*	*S. reilianum*	*U. maydis*	*U. hordei*	*M. pennsylvanicum*	*A. flocculosa*
Assembly statistics								
Total contig length (Mb)	18.3		19.5	18.2	19.7	20.7	19.2	23.2
Total scaffold length (Mb)		20.5	19.6	18.4	19.8	21.15	19.2	23.3
Average base coverage	50x	154x	30x	20x	10x	25x	339x	28x
N50 contig (kb)	705.1		37.6	50.3	127.4	48.7	43.4	38.6
N50 scaffold length (kb)		877	759.2	738.5	817.8	307.7	121.7	919.9
Chromosomes	21	23		23	23	23		
GC-content (%)	60.9	52.4	54.4	59.7	54	52	50.9	65.1
Coding (%)	62.8	54.4	57.8	62.6	56.3	54.3	54	66.3
Coding Sequence								
Percentage CDS (%)	69.5	59.8	62	65.9	61.1	57.5	56.6	54.3
Average gene size (bp)	1935	1699	1819	1858	1836	1708	1734	2097
Average gene density (gene/kb)	0.36	0.35	0.34	0.37	0.34	0.34	0.33	0.30
Protein-coding genes	6653	7233	6693	6648	6786	7113	6279	6877
Exons	11645	11154	10214	9776	9783	10907	9278	19318
Average exon size (bp)	1091	1101	1191	1221	1230	1107	527	658
Exons/gene	1.75	1.5	1.5	1.47	1.44	1.53	1.48	2.8
tRNA genes	150	133	116	96	111	110	126	176
Noncoding sequence								
Introns	9333	3921	3521	3103	2997	3161	2999	12427
Introns/gene	1.40	0.54	0.53	0.47	0.44	0.44	0.48	1.81
Average intron length (base)	163	163	130.1	144	142	141	191.4	141
Average intergenic distance (bp)	769	1054	1114	929	1127	1186	1328	1273
Secretome								
Protein with signal peptide	559		622	632	625	538	419	622
Secreted without TMD	380				467			737
- with known domain	260				264			554

[24,42]

Table 2: *M. albugensis* proteins homologous to *U. maydis* effector genes with known virulence function.

Name	Homologue	Query cover	E-value	Identity (%)	*U. maydis* knockout phenotype	Reference
g1653	UMAG_01987 (Pep1)	82%	3-e56	60.96	complete loss of tumor formation - blocked in early stages of infection	[51]
g1828	UMAG_01829 (Afu1)	99%	0.0	71.57	organ specific effector - reduced virulence in seedling leaves	[52]
g2626	UMAG_12197 (Cce1)	98%	1e-48	60.16	complete loss of tumor formation - blocked in early stages of infection	[53]
g2765	UMAG_11938 (Scp2)	100%	1e-73	93.44	Reduced in virulence	[54]
g2910	UMAG_02475 (Stp1)	32%	3e-42	60.71	complete loss of tumor formation - blocked in early stages of infection	[55]
g3652	UMAG_02239 (See1)	43%	9e-11	54.90	organ specific effector - reduced virulence in seedling leaves	[19]
g3113	UMAG_01375 (Pit2)	*	*	*	complete loss of tumor formation - blocked in early stages of infection	[56]
g3279	UMAG_03274 (Rsp3)	10%	5e-20	70.11	strong attenuation of virulence – reduced tumor size and number	[57]
g5296	UMAG_05731 (Cmu1)	98%	3e-70	43.84	Reducedvirulence	[58]
g6183	UMAG_06098 (Fly1)	100%	0.0	81.85	Reducedvirulence	[59]
g5835	UMAG_05302 (Tin2)	87%	8e-24	37.81	Minor impact on tumor formation – reduced anthocyanin biosynthesis	[23]

Figure 1

A

B

U. maydis FB2

M. albugensis

Figure 1: Position of *M. albugensis* in the family of Ustilaginaceae. (A) molecular phylogenetic analysis by maximum likelihood method based on fungal ITS sequences and showing grouping of *M. albugensis* with the millet pathogen *M. bullatus*. Pathogenic (green) and apathogenic species can be found across the phylogenetic tree. (B) Comparison of the morphology of *M. albugensis* (bottom) and *U. maydis* (top) haploid cells grown in YEPSLight medium to OD_{600}0.6.

Figure 2

Figure 2: Infection assay of *A. laibachii* on *A. thaliana*. Addition of a bacterial SynCom reduces the infection symptoms of *A. laibachii* at 14dpi. Those symptoms can be almost abolished by spraying *M. albugensis* to the plant, independently of the presence of the bacterial community. An analysis of variance (ANOVA) model was used for pairwise comparison of the conditions, with Tukey's HSD test to determine significant differences among them (P values <0.05).

Figure 3

Figure 3: Circos comparison of *M. albugensis* and *U. maydis* chromosome structure (A). We highlighted potential secondary metabolite clusters, secreted proteins and gene predictions on both strands (+/-). (B) Homology based comparisons identified three chromosomal recombination events, which affects the *M. albugensis* contigs 2, 6 & 8.

Figure 4

Figure 4: Structure of the largest virulence cluster (Cluster 19A) in pathogenic and non-pathogenic smut fungi. Colors indicate genes with homology to each other: Related gene families are indicated in orange, yellow, blue, green and brown, whereas unique effector genes are shown in grey. Genes encoding proteins without a secretion signal are shown in white [23].

Figure 5

Figure 5: Genetic transformation of *M. albugensis.* (A) Stable transformants that express cytosolic GFP could be obtained by generating protoplasts with Glucanex and ectopically integrating linear DNA-fragments into the genome via PEG-mediated transformation. (B) Overview of the split-marker approach that was used to generate deletion mutants via homologous recombination.

Figure 6

Figure 6: The self-compatible *M. albugensis* strain CB1 (A) *M. albugensis* mating type genes, unlike the ones of *U. hordei*, can be found on two different chromosomes similar to the tetrapolar mating type system of *U. maydis*. (B) To generate a self-compatible strain (CB1), the b-mating genes of *U.hordei* were integrated at the native *M. albugensis* b-locus. (C) Unlike the *M. albugensis* wild type strain (top left), strain CB1 (bottom left) shows a fluffy phenotype on charcoal plates and filamentous growth. *U. maydis* haploid F1 strain (top right) and self-compatible SG200 strain (bottom right) were used as negative and positive control, respectively. (D,E) Induction of filamentation and appressoria formation in strain CB1. After incubation on a hydrophobic surface, both, filament and appressoria formation in strain CB1 were significantly different (* Chi-Square Test for Independence – p = 0,0001) when compared to *M. albugensis* wild type and similar to the level of the self-compatible *U. maydis* strain SG00. *U. maydis* haploid F2 strain was used as negative control [18]. Scale bar: 20µm.

Figure 7

Figure 7: Transcriptome analysis of *M. albugensis* (A) Experimental setup used for the transcriptomic (RNA-Sequencing) analysis in *M. albugensis*. (B) Venn diagrams showing differential regulated *M. albugensis* genes after spraying of haploid cells onto the *A. thaliana* leaf surface. A total number of 801 genes were upregulated in response to *A. laibachii* in presence and absence of bacterial SynCom. 216 of the 801 genes were upregulated in both conditions. (C) Hierarchical clustering of the 27 *A. laibachii* - induced *M. albugensis* genes that are predicted to encode secreted proteins. Of these genes, nine were selected as candidate microbe-microbe effector genes, based on their transcriptional upregulation and prediction to encode for extracellularly localized proteins.

Figure 8

Figure 8: A reverse-genetic approach to identify the *M. albugensis* gene which is responsible for the suppression of *A. laibachii* infection. (A) Three candidate microbe-microbe effector genes (*g5, g5755* &*g2490*) were deleted in *M. albugensis* and deletion strains were individually inoculated on *A. thaliana* together with *A. laibachii*. Inoculation of two independent *g2490 null* strains *(Δg2490_1; Δg2490_2)* resulted in significant and almost complete loss the biocontrol activity of *M. albugensis*. While deletion of g5 resulted in a marginal reduction of disease symptoms at 14 days post infection, deletion of g5755 had no effect on *A. laibachii.* (B) Genetic complementation of the *g2490* deletion restores the biocontrol activity to wild type levels.