Synonymous mutations and the molecular evolution of SARS-Cov-2 origins

Hongru Wang[1], Lenore Pipes[1] and Rasmus Nielsen[1, 2,3*]

[1] Department of Integrative Biology, UC Berkeley, Berkeley, CA 94707, USA.

[2]Department of Statistics, UC Berkeley, Berkeley, CA 94707, USA.

[3]Globe Institute, University of Copenhagen, 1350 København K, Denmark.

*Address: 4098 Valley Life Sciences Building, Department of Integrative Biology, UC Berkeley. Berkeley, CA 94707. rasmus_nielsen@berkeley.edu

Abstract

Human severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is most closely

related, by average genetic distance, to two coronaviruses isolated from bats, RaTG13 and

RmYN02. However, there is a segment of high amino acid similarity between human SARS-

CoV-2 and a pangolin isolated strain, GD410721, in the receptor binding domain (RBD) of

the spike protein, a pattern that can be caused by either recombination or by convergent

amino acid evolution driven by natural selection. We perform a detailed analysis of the

synonymous divergence, which is less likely to be affected by selection than amino acid

divergence, between human SARS-CoV-2 and related strains. We show that the

synonymous divergence between the bat derived viruses and SARS-CoV-2 is larger than

between GD410721 and SARS-CoV-2 in the RBD, providing strong additional support for

the recombination hypothesis. However, the synonymous divergence between pangolin

strain and SARS-CoV-2 is also relatively high, which is not consistent with a recent

recombination between them, instead it suggests a recombination into RaTG13. We also

find a 14-fold increase in the d_N/d_S ratio from the lineage leading to SARS-CoV-2 to the

strains of the current pandemic, suggesting that the vast majority of non-synonymous

mutations currently segregating within the human strains have a negative impact on viral

fitness. Finally, we estimate that the time to the most recent common ancestor of SARS-

CoV-2 and RaTG13 or RmYN02 based on synonymous divergence, is 51.71 years (95%

C.I., 28.11-75.31) and 37.02 years (95% C.I., 18.19-55.85), respectively.

Introduction

The Covid19 pandemic is perhaps the biggest public health and economic threat that the world

has faced for decades (Li, Guan, et al. 2020; Wu, et al. 2020; Zhou, Yang, et al. 2020). It is

caused by a coronavirus (Lu, et al. 2020; Zhang and Holmes 2020), Severe acute respiratory

syndrome coronavirus 2 (SARS-CoV-2), an RNA virus with a 29,903 bp genome consisting of

four major structural genes (Wu, et al. 2020; Zhou, Yang, et al. 2020). Of particular relevance to

this study is the *spike* protein which is responsible for binding to the primary receptor for the

virus, angiotensin-converting enzyme 2 (*ACE2)* (Wan, et al. 2020; Wu, et al. 2020; Zhou, Yang,

et al. 2020).

Human SARS-CoV-2 is related to a coronavirus (RaTG13) isolated from the bat

Rhinolophus affinis from Yunnan province of China (Zhou, Yang, et al. 2020). RaTG13 and the

human strain reference sequence (Genbank accession number MN996532) are 96.2% identical

and it was first argued that, throughout the genome, RaTG13 is the closest relative to human

SARS-CoV-2 (Zhou, Yang, et al. 2020). And RaTG13 and SARS-CoV-2 were 91.02% and

90.55% identical, respectively, to coronaviruses isolated from Malayan pangolins (Pangolin-

CoV) seized at the Guangdong customs of China, which therefore form a close outgroup to the

SARS-CoV-2+RaTG13 clade (Zhang, et al. 2020). Furthermore, five key amino acids in the

receptor-binding domain (RBD) of *spike* were identical between SARS-CoV-2 and Pangolin-

CoV, but differed between those two strains and RaTG13 (Zhang, et al. 2020). Xiao et al

assembled and analysed a full-length Pangolin-CoV genome sequence, showing that the

receptor-binding domain of its S protein differs from the SARS-CoV-2 by only one noncritical

amino acid (Xiao, et al. 2020). Similar observations were made using Pangolin-CoV strains

found in Malayan pangolin samples seized by the Guangxi customs of China (Lam, et al. 2020).

Additionally, it is shown that when analyzing a window of length 582bp in the RBD,

nonsynonymous mutations support a phylogenetic tree with SARS-CoV-2 and Pangolin-CoV as

sister-groups, while synonymous mutations do not (Lam, et al. 2020). They discuss two possible

explanations for their results, one which includes recombination and another which includes

selection-driven convergent evolution. Independent analysis also support SARS-CoV-2 obtains

the receptor binding motif through recombination from a donor related to this Pangolin-CoV

strain (Li, Giorgi, et al. 2020). Detailed phylogenetic analysis on sub-regions across the S

protein showed that it is the RaTG13 sequence that show exceptionally divergent pattern in the

RBD region, they instead argued a recombination occurred into RaTG13 from an unknown

divergent source (Boni, et al. 2020). This would explain the amino acid similarity between

SARS-CoV-2 and Pangolin-CoV in the RBD as an ancestral trait that has been lost (by

recombination) in RaTG13. Using a phylogenetic analysis they also dated the RaTG13 and

SARS-CoV-2 divergence to be between 40 to 70 years. Recently, Zhou et al. discovered a viral

strain, RmYN02 from the bat *Rhinolophus malayanus*, with a reported 97.2% identity in the

ORF1ab gene but with only 61.3% sequence similarity to SARS-CoV-2 in the RBD (Zhou, Chen,

et al. 2020). Moreover, the RmYN02 strain also harbors multiple amino acid insertions at the

S1/S2 cleavage site in the spike protein (Zhou, Chen, et al. 2020).

To analyze the history of these sequences further, we here focus on patterns of

synonymous divergence, which has received less focus, but also is less likely to be affected by

selection than amino acid divergence. We develop a bias corrected estimator of synonymous

divergence specific for SARS-CoV-2 and related strains, and analyze divergence using both

sliding windows and a whole-genome approach between SARS-CoV-2 and related viral strains.

Materials and methods

BLAST searches: Sequences for blast databases were downloaded on March 26, 2020 from the

following sources: EMBL nucleotide libraries for virus

(ftp://ftp.ebi.ac.uk/pub/databases/embl/release/std), NCBI Virus Genomes

(ftp://ftp.ncbi.nlm.nih.gov/genomes/Viruses), NCBI Virus Genbank Entries

(ftp://ftp.ncbi.nlm.nih.gov/genomes/genbank/viral/), NCBI Influenza Genomes

(ftp://ftp.ncbi.nlm.nih.gov/genomes/INFLUENZA/), all Whole Genome Shotgun

(https://www.ncbi.nlm.nih.gov/genbank/wgs/) assemblies under taxonomy ID 10239, along with

GISAID Epiflu and EpiCoV databases. Recently published sequences from the Myanmar bat

samples (Valitutto, et al. 2020) were also added to the database. Blast databases were created

using the default parameters for makeblastdb. Blast searches were performed using blastn

(Altschul, et al. 1990) with parameters "-word_size 7 -reward 1 -penalty -3" and all other

parameters as the default settings. All the blast hits to different Guangdong pangolin viral strain

sequences were merged as one hit, and the blast hits to different Guangxi pangolin viral strain

sequences were also merged.

Alignment: To obtain an in-frame alignment of the genomes, we first identified the coding

sequences of each viral strain using independent pairwise alignments with the coding

sequences of the SARS-CoV-2 (Wuhan-Hu-1) genome. The genome alignments were

performed using MAFFT (v7.450) (Katoh and Standley 2013) with parameters "--maxiterate

1000 --localpair". The coding sequences of each gene were aligned using PRANK (Loytynoja

2014) (v.170427) with parameters "-codon -F". Finally, the alignments for all genes were

concatenated following their genomic order. ORF1a was excluded since its sequence is a

subset of ORF1ab.

Recombination detection: We detected possible recombination events across the genome using

a combination of 7 alogorithms, RDP (Martin and Rybicki 2000), GENECONV (Padidam, et al.

1999), Bootscan (Salminen, et al. 1995), Maxchi (Smith 1992), Chimaera (Posada and Crandall

2001), SiSscan (Gibbs, et al. 2000), and 3Seq (Boni, et al. 2007) implemented in RDP5

program (Martin, et al. 2015) (version Beta 5.5) and then considered the recombination signals

that were supported by at least two methods. We note that these 7 methods are all based on

inferring recombination using the same type of evidence, and concordance between the

methods cannot be interpreted as validation of the recombination signal. However, we will also

use phylogenetic methods and methods based on relative sequence divergence to further

investigate the putative recombination signals. The analysis was performed on the multiple

sequence alignment consisting of the five viral strains. All regions showing recombination

signals (Supplementary Table 5) were removed in subsequent analyses from all strains when

stating that recombination regions were removed.

Tree estimation: We estimated phylogenetic trees using two methods: Neighbor Joining (NJ)

and Maximum Likelihood (ML). The NJ trees were estimated using d_N or d_S distance matrices

which estimated using codeml (Yang 2007) with parameters " runmode= -2, CodonFreq = 2,

cleandata = 1". To obtain bootstrap values, we bootstrapped the multiple sequence alignments

1,000 times, repeating the inference procedure for each bootstrap sample. The NJ tree was

estimated using the 'neighbor' software from the PHYLIP package (Felsenstein 2009). For ML

trees, we used IQ-TREE (Nguyen, et al. 2015) (v1.5.2) with parameter "-TEST -alrt 1000" which

did substitution model selection for the alignments and performed maximum-likelihood tree

estimation with the selected substitution model for 1,000 bootstrap replicates. For this analysis,

we masked all regions (Supplementary Table 5) that show recombination signals in any of the

five studied viral genome. We masked regions from all sequences when at least one sequence

showed evidence for recombination in that region. All masked regions are listed in

Supplementary Table 5. The coordinates (based on the Wuhan-Hu-1 genome) of the three

recombination regions (merged set of all the regions in Supplementary Table 5) were: 14611-

15225, 21225-24252 and 25965-28297. We also estimate genome-wide divergence between

RaTG13 and Wuhan-Hu-1 only excluding the region (position 22853-23092) where potential

recombination was detected for the Wuhan-Hu-1 strain (Supplementary Table 5).

Simulations: We simulated divergence with realistic parameters for SARS-CoV-2 using a

continuous time Markov chain under the F3x4 codon-based model (Goldman and Yang 1994;

Muse and Gaut 1994) (Yang, et al. 2000), which predicts codon frequencies from the empirical

nucleotide frequencies in all 3 codon positions and using the global genomic maximum

likelihood estimates of the transition/transversion bias κ(=2.9024) and the d_N/d_S ratio ω

(=0.0392) estimated from the human SARS-CoV-2 comparison to the nearest outgroup

sequence, RaTG13 (see Results). For the simulations of short 300 bp sequences we kept ω

constant but varied time such that the number of synynoymous substitutions per synonymous

sites, d_S, varied between 0.25 and 3.00. Estimates of d_S > 3 are truncated to 3. For simulations

of genome-wide divergence between RaTG13 and human strains, we fix d_S at 0.1609 (the

maximum likelihood estimate outside the RBD region reported in the Results section). In all

cases, we use 10,000 independent replicate simulations for each parameter setting.

Estimation of sequence divergence in 300-bp windows: d_N and d_S were estimated using two

different methods implemented in the PAML package (Yang 2007) (version 4.9d): a count-

based method, YN00 (Yang and Nielsen 2000) as implemented in the program 'yn00' with

parameters "icode = 0, weighting = 0, commonf3x4 = 0", and a maximum-likelihood method

(Goldman and Yang 1994; Muse and Gaut 1994) implemented in codeml applied with

arguments "runmode= -2, CodonFreq = 2". The estimates in 300-bp windows were further bias-

corrected as described below.

Bias correction for d_S estimates in 300-bp window: To correct for the biases observed in the

estimation of d_S (see results section) we identifed a quartic function which maps from $\hat{d_S}$, the

estimates of d_S, into $\widehat{d_S}^{*}$, the bias corrected estimate such that to a close approximation, $E[\widehat{d_S}^{*}]$

= d_S. To identify the coefficients of this function we used 10,000 simulations as previously

described, on a grid of d_S values (0.25, 0.5, 0.75, ..., 3.0). We then identified coefficients such

that sum of $(E[\widehat{d_S}^{*}] - d_S)^2$ is minimized over all simulation values.

Results

Database searches

The genome of human coronavirus can effectively recombine with other viruses to form a

chimeric new strain when they co-infect the same host (Forni, et al. 2017; Boni, et al. 2020).

Complicated recombination histories have been observed in the receptor binding motif region of

the spike protein (Lam, et al. 2020; Xiao, et al. 2020; Zhang, et al. 2020) and several other

regions (Boni, et al. 2020) of the SARS-CoV-2, it is thus important to exhaustively search along

the viral genome for other regions potentially of recombination origin and identify possible

donors associated with them. To identify possible viral strains that may have contributed, by

recombination, to the formation of human SARS-CoV-2, we searched NCBI and EMBL virus

entries along with GISAID Epiflu and EpiCov databases for similar sequences using BLAST in

100bp windows stepping every 10bp (Fig. 1b). The majority of the genome (78.1%, 2330/2982

of the windows) has one unique best hit, likely reflecting the high genetic diversity of the

coronavirus. 21.9% of the genomic regions has multiple best hits, which suggests that these

regions might be more conserved. Among the windows with unique best hits, 97.0%

(2260/2330) of them were the RaTG13 or RmYN02 bat strains and 1.9% of them, including the

ACE2 contact residues region of the S protein, were the pangolin SARS-CoV-2 virus. These

observations are consistent with previous results that RaTG13 and RmYN02 are the most

closely related viral strains, while the region containing the *ACE2* contact residues is more

closely related to the pangolin virus strain (Lam, et al. 2020; Li, Giorgi, et al. 2020; Xiao, et al.

2020; Zhang, et al. 2020). A considerable amount of genomic regions (20 windows with unique

hits) show highest sequence identity with other coronaviruses of the SARS-CoV-2 related

lineage (Lam, et al. 2020) (bat-SL-CoVZC45 and bat-SL-CoVZXC21 (Hu, et al. 2018)). In

addition, there were 6 windows whose unique top hits are coronavirus of a SARS-CoV related

lineage (Lam, et al. 2020) (Supplementary Table 4). The mosaic pattern that different regions of

the genome show highest identity to different virus strains is likely to have been caused by the

rich recombination history of the SARS-CoV-2 lineage (Boni, et al. 2020; Li, Giorgi, et al. 2020;

Patiño-Galindo, et al. 2020). Moreover, its unique connection with SARS-CoV related lineages

in some genomic regions may suggest recombination between the ancestral lineage of SARS-

CoV-2 and distantly related virus lineages, although more formal analyses are needed to

determine the recombination history (see also Boni, et al. 2020 for further discussion).

Searching databases with BLAST using the most closely related viral strains, RaTG13 and

RmYN02, we observe a very similar pattern, as that observed for SARS-CoV-2, in terms of top

hits across the genome (Fig. 1b), suggesting that these possible recombination events with

distantly related lineages are not unique to the SARS-CoV-2 lineage, but happened on the

ancestral lineage of SARS-CoV-2, RaTG13, and RmYN02. A notable exception is a large region

around the *S* gene, where RmYN02 show little similarity to both SARS-CoV-2 and RaTG13.

Sequence similarity and recombination

We focus further on studying the synonymous evolution of SARS-CoV-2, and analyzing Wuhan-

Hu-1 as the human nCoV19 reference strain (Wu, et al. 2020) along with the four viral strains

with highest overall identity: the bat strains RmYN02 and RaTG13 (Zhou, Chen, et al. 2020;

Zhou, Yang, et al. 2020), and the Malayan pangolin strains, GD410721 and GX_P1E, which

were isolated from Malayan pangolin samples seized by Guangdong and Guangxi Customs of

China, respectively. These four strains have previously been identified as the strains most

closely related to SARS-CoV-2 (Lam, et al. 2020; Xiao, et al. 2020). Other available

phylogenetically related, but less similar viral strains, such as bat-SL-CoVZXC21 and bat-SL-

CoVZC45 (Hu, et al. 2018), are not included due to nearly saturated synonymous mutations

when compared with SARS-CoV-2 (maximum likelihood estimates of d_S = 3.2067 and 2.8445,

respectively).

We performed recombination analyses across the five viral genomes based on the

concensus of the seven recombination-detection methods implemented in RDP5 (see Methods).

We identified nine recombination regions affecting at least one of the sequences

(Supplementary Table 5). Phylogenetic analyses of these regions confirm phylogenetic

incongruence when compared with genome-wide trees (Fig. 2 and Supplementary Figure 1-3).

Particularly, a recombination signal is found in a region encompassing the RBD of the S protein,

suggesting that the human SARS-CoV-2 (Wuhan-Hu-1) sequence is a recombinant with the

Pangolin-CoV (GD410721) as the donor (Supplementary Table 5). Phylogenetic analyses also

support that Wuhan-Hu-1 and GD410721 form a clade relative to RaTG13 (Supplementary

Figure 1c, 1d). Phylogenetic analyses (Fig. 2) in genomic regions with all recombination tracts

(Supplementary Table 5) masked using Maximum-likelihood (Fig. 2a) and Neighbor-joining

based on synonymous (Fig. 2b) or non-synoymous (Fig. 2c) mutation distance metrics,

consistently support RmYN02 as the nearest outgroup to human SARS-CoV-2, in contrast to

previous analyses before the discovery of RmYN02, which instead found RaTG13 to be the

nearest outgroup (Lam, et al. 2020; Wu, et al. 2020). This observation is also consistent with the

genome-wide phylogeny constructed in previous study (Zhou, Chen, et al. 2020).

We plot the overall sequence similarity (% nucleotides identical) between SARS-CoV-2

and the four other strains analyzed in windows of 300 bp (Fig. 1). Notice that the divergences

between human SARS-CoV-2 and the bat viral sequences, RaTG13 and RmYN02, in most

regions of the genome, are quite low compared to the other comparisons. A notable exception is

the suspected recombination region in RmYN02 that has an unusual high level of divergence

with all other viruses (Fig. 1e). However, there is also another exception: a narrow window in the

RBD of the S gene where the divergence between SARS-CoV-2 and GD410721 is moderate

and the divergences between GD410721 and both SARS-CoV-2 and RaTG13 are quite high

and show very similar pattern. This, as also found in the recombination analyses based on

methdos implemented in RDP5, would suggest a recombination event from a strain related to

GD410721 into an ancestor of the human strain (Lam, et al. 2020; Xiao, et al. 2020; Zhang, et

al. 2020), or alternatively, from some other species into RaTG13, as previously hypothesized

(Boni, et al. 2020). We note that RmYN02 is not informative about the nature of this event as it

harbors a long and divergent haplotype in this region, possibly associated with another

independent recombination event with more distantly related viral strains (Fig. 1e). The other

four sequences are all highly, and approximately equally, divergent from RmYN02 in this large

region (Fig. 1e), suggesting that the RmYN02 strain obtained a divergent haplotype from the

recombination event. When BLAST searching using 100-bp windows along the RmYN02

genome, we find no single viral genome as the top hit, instead the top hits are found

sporadically in different viral strains of the SARS-CoV lineage (Fig. 1f), suggesting that the

sequence of the most proximal donor is not represented in the database.

Estimating synonymous divergence and bias correction

While the overall divergence in the *S* gene encoding the *spike* protein could suggest the

presence of recombination in the region, previous study (Lam, et al. 2020) reported that the tree

based on synonymous substitutions supported RaTG13 as the sister taxon to the human SARS-

CoV-2 also in this region. That would suggest the similarity between GD410721 and human

SARS-CoV-2 might be a consequence of convergent evolution, possibly because both strains

adapted to the use of the same receptor. An objective of the current study is to examine if there

are more narrow regions of the spike protein that might show evidence of recombination. We

investigate this issue using estimates of synonymous divergence per synonymous site (d_S) in

sliding windows of 300 bp. However, estimation of d_S is complicated by the high levels of

divergence and extremely skewed nucleotide content in the 3rd position of the sequences

(Table 1) which will cause a high degree of homoplasy. We, therefore, entertain methods for

estimation that explicitly account for unequal nucleotide content and multiple hits in the same

site such as maximum likelihood methods and the YN00 method (Yang and Nielsen 2000). It is

shown that for short sequences, some counting methods, such as the YN00 method, can

perform better in terms of Mean Squared Error (MSE) for estimating d_N and d_S(Yang and

Nielsen 2000). However, it is unclear in the current case how best to estimate d_S. For this

reason, we performed a small simulations study (see Methods) for evaluating the performance

of the maximum likelihood (ML) estimator of d_N and d_S (as implemented in codeml (Yang 2007))

under the F3x4 model and the YN00 method implemented in PAML. In general, we find that

estimates under the YN00 are more biased with slightly higher MSE than the ML estimate for

values in the most relevant regime of d_S < 1.5 (Fig. 3). However, we also notice that both

estimators are biased under these conditions. For this reason, we perform a bias correction

calibrated using simulations specific to the nucleotide frequencies and d_N/d_S ratio observed for

SARS-CoV-2 (see Methods). The bias corrections we obtain are $\hat{d_S}^* = \hat{d_S} + 0.455\hat{d_S}^2 - 0.824\hat{d_S}^3$

$+ 0.264\hat{d_S}^4$, for the ML estimator and $\hat{d_S}^* = \hat{d_S}+1.492\hat{d_S}^2 - 3.166\hat{d_S}^3 + 1.241\hat{d_S}^4$ for yn00. Notice

that there is a trade-off between mean and variance (Fig. 3) so that the MSE becomes very

large, particularly for the for yn00 method, after bias correction. For d_S >2 the estimates are

generally not reliable, however, we note that for d_S <1.5 the bias-corrected ML estimator tends

overall to have slightly lower MSE, and we, therefore, use this estimator for analyses of 300 bp

regions.

Synonymous divergence

We estimate d_N and d_S under the F3x4 model in codeml (Goldman and Yang 1994; Muse and

Gaut 1994) and find genome-wide estimates of d_S = 0.1604, d_N = 0.0065 (d_N/d_S = 0.0405)

between SARS-CoV-2 and RaTG13 and d_S = 0.2043, d_N = 0.0220 (d_N/d_S = 0.1077) between

SARS-CoV-2 and RmYN02. However, a substantial amount of this divergence might be caused

by recombination with more divergent strains. We, therefore, also estimate d_N and d_S for the

regions with inferred recombination tracts (Supplementary Table 5) removed from all

sequences (Table 3). We then find values of d_S = 0.1462 (95% C.I., 0.1340-0.1584) and d_S =

0.1117 (95% C.I., 0.1019-0.1215) between SARS-CoV-2 and RaTG13 and RmYN02,

respectively. This confirms that RmYN02 is the virus most closely related to SARS-CoV-2. The

relative high synonymous divergence also shows that the apparent high nucleotide similarity

between SARS-CoV-2 and the bat strains (96.2% (Zhou, Yang, et al. 2020) and 97.2%(Zhou,

Chen, et al. 2020)) is caused by conservation at the amino acid level (d_N/d_S = 0.0410 and

0.0555) exacerbated by a high degree of synonymous homoplasy facilitated by a highly skewed

nucleotide composition at the third position of codons (with an AT content >72%, Table 1).

The synonymous divergence to the pangolin sequences GD410721 and GX_P1E in

genomic regions with inferred recombination tracts removed is 0.5095 (95% C.I., 0.4794-

0.5396) and 1.0304 (95% C.I., 0.9669-1.0939), respectively. Values for other comparisons are

shown in Tables 2 and 3. In comparisons between SARS-CoV-2 and more distantly related

strains, d_S will be larger than 1, and with this level of saturation, estimation of divergence is

associated with high variance and may be highly dependent on the accuracy of the model

assumptions. This makes phylogenetic analyses based on synonymous mutations unreliable

when applied to these more divergent sequences. Nonetheless, the synonymous divergence

levels seem generally quite compatible with a molecular clock with a d_S of 0.9974 (95% C.I.,

0.9381-1.0567, GD410721), 1.0366 (95% C.I., 0.9737-1.0995, RaTG13), 1.0333 (95% C.I.,

0.9699-1.0967, RmYN02) and 1.0304 (95% C.I., 0.9669-1.0939, Wuhan-Hu-1) between the

outgroup, GX_P1E , and the three ingroup strains. The largest value is observed for RaTG13

(d_S = 1.0366), despite this sequence being the most early sampled sequence, perhaps caused

by additional undetected recombination into RaTG13.

Sliding windows of synonymous divergence

To address the issue of possible recombination we plot d_S between SARS-CoV-2, GD410721,

and RaTG13 and the ratio of d_S(SARS-CoV-2, GD410721) to d_S(SARS-CoV-2, RaTG13) in 300

bp sliding windows along the genome. Notice that we truncate the estimate of d_S at 3.0.

Differences between estimates larger than 2.0 should not be interpreted strongly, as these

estimates have high variance and likely will be quite sensitive to the specifics of the model

assumptions.

We find that d_S(SARS-CoV-2, GD410721) approximately equals d_S(GD410721,

RaTG13) and is larger than d_S(SARS-CoV-2, RaTG13) in almost the entire genome showing

than in these parts of the genome GD410721 is a proper outgroup to (SARS-CoV-2, RaTG13)

assuming a constant molecular clock. One noticeable exception from this is the RBD region of

the *S* gene. In this region the divergence between SARS-CoV-2 and GD410721 is substantially

lower than between GD410721 and RaTG13 (Fig. 4a,4c). The same region also has much

smaller divergence between SARS-CoV-2 and GD410721 than between SARS-CoV-2 and

RaTG13 (Fig. 4a,4c). The pattern is quite different than that observed in the rest of the genome,

most easily seen by considering the ratio of d_S(SARS-CoV-2, GD410721) to d_S(SARS-CoV-2,

RaTG13) (Fig. 2b, 2d). In fact, the estimates of d_S(SARS-CoV-2, RaTG13) are saturated in this

region, even though they are substantially lower than 1 in the rest of the genome. This strongly

suggests a recombination event in the region and provides independent evidence of that

previously reported based on amino acid divergence (e.g.,(Zhang, et al. 2020)).

The combined evidences from synonymous divergence and the topological

recombination inference, provide strong support for the recombination hypothesis. However,

these analyses alone do not distinguish between recombination into RaTG13 from an unknown

source as previously hypothesized (Boni, et al. 2020) and recombination between SARS-CoV-2

and GD410721 as proposed as one possible explanation by Lam et al. (Lam, et al. 2020). To

distinguish between these hypotheses we searched for sequences that might be more closely

related, in the RBD region, to RaTG13 than SARS-CoV-2 and we plotted sliding window

similarities across the genome for RaTG13 (Fig. 1c). We observe relatively low sequence

identity between RaTG13 and all three other strains in the *ACE2* contact residue region of the

spike protein, which is more consistent with the hypothesis of recombination into RaTG13, as

proposed in (Boni, et al. 2020). Moreover, our BLAST search analyses of RaTG13 in this region

show highest local sequence similarity with GX pangolin virus strains which is the genome-wide

outgroup for the three other sequences (Lam, et al. 2020). This observation is more compatible

with the hypothesis of recombination from a virus related to GX pangolin strains, than with

recombination between SARS-CoV-2 and GD410721.

Unfortunately, because of the high level of synonymous divergence to the nearest

outgroup, tree estimation in small windows is extremely labile in this region. In fact, synonymous

divergence appears fully saturated in the comparison with GX_P1E, eliminating the possibility to

infer meaningful trees based on synonymous divergence. However, we can use the overall

maximum likelihood tree using both synonymous and nonsynonymous mutations (Fig. 2d). The

ML tree using sequence from the *ACE2* contact residue region supports the clustering of SARS-

CoV-2 and GD410721, but with unusual long external branches for all strains except SARS-

CoV-2, possibly reflecting smaller recombination regions within the *ACE2* contact residue

region.

Weakly deleterious mutations and clock calibrations

The use of synonymous mutations provides an opportunity to calibrate the molecular clock

without relying on amino acid changing mutations that are more likely to be affected by

selection. The rate of substitution of weakly and slightly deleterious mutations is highly

dependent on ecological factors and the effective population size. Weakly deleterious mutations

are more likely to be observed over small time scales than over long time scales, as they are

unlikely to persist in the population for a long time and go to fixation. This will lead to a

decreasing d_N/d_S ratio for longer evolutionary lineages. Furthermore, changes in effective

population size will translate into changes in the rate of substitution of slightly deleterious

mutations. Finally, changes in ecology (such as host shifts, host immune changes, changes in

cell surface receptor, etc.) can lead to changes in the rate of amino acid substitution. For all of

these reasons, the use of synonymous mutations, which are less likely to be the subject of

selection than nonsynonymous mutations, are preferred in molecular clock calculations. For

many viruses, the use of synonymous mutations to calibrate divergence times is not possible,

as synonymous sites are fully saturated even at short divergence times. However, for the

comparisons between SARS-CoV-2 and RaTG13, and SARS-CoV-2 and RmYN02,

synonymous sites are not saturated and can be used for calibration. We find an estimate of ω =

0.0391 between SARS-CoV-2 and RaTG13, excluding just the small RDB region showing a

recombination signal in SARS-CoV-2 (Supplementary Table 5, coordinates: 22851-23094).

Using 1000 parametric simulations under the estimated values and the F3x4 codon model, we

find that the estimate is approximately unbiased ($\hat{\omega} = 0.0398$, S.E.M.= 0.0001) and with

standard deviation 0.0033, providing an approximate 95% confidence interval of (0.0332,

0.0464). Also, using 59 human strains of SARS-CoV-2 from Genbank and National Microbiology

Data Center (See Methods) we obtain an estimate of ω = 0.5604 using the F3x4 model in

codeml. Notice that there is a 14-fold difference in d_N/d_S ratio between these estimates.

Assuming very little of this difference is caused by positive selection, this suggests that the vast

majority of mutations currently segregating in the SARS-CoV-2 are slightly or weakly deleterious

for the virus.

Dating of divergence between Bat viruses and SARS-CoV-2

To calibrate the clock we use the estimate provided by (http://virological.org/t/phylodynamic-

analysis-of-sars-cov-2-update-2020-03-06/420) of $\mu =1.04\times10^{-3}$ substitutions/site/year (95% CI:

0.71x10-3, 1.40x10-3). The synonymous specific mutation rate can be found from this as

d_S/year = μ_S = $\mu/(pS +\omega pN)$, where ω is the d_N/d_S ratio, and *pN* and *pS* are the proportions of

nonsynonymous and synonymous sites, respectively. The estimate of the total divergence on

the two lineages is then $\hat{t} = dS\,(pS + \omega pN)/\mu$. Inserting the numbers from Table 3 for the

divergence between SARS-CoV-2 and RaTG13 and RmYN02 ,respectively, we find a total

divergence of 96.92 years and 74.05 years respectively. Taking into account that RaTG13 was

isolated July 2013, we find an estimated tMRCA between that strain and SARS-CoV-2 of

$\hat{t} =$(96.92 +6.5)/2 = 51.71 years. Similarly, we find an estimate of divergence between SARS-

CoV-2 and RmYN02 of $\hat{t} =$74.05/2 = 37.02 years, assuming approximately equal sampling

times. The estimate for SARS-CoV-2 and RaTG13 is compatible with the values obtained using

different methods for dating (Boni, et al. 2020). The variance in the estimate in d_S is small and

the uncertainty is mostly dominated by the uncertainty in the estimate of the mutation rate. We

estimate the S.D. in $\hat{t}$ using 1000 parametric simulations, using the ML estimates of all

parameters, for both RaTG13 vs. SARS-CoV-2 and for RmYN02 vs. SARS-CoV-2, and for each

simulated data also simulating values of μ and ω from normal distributions with mean 1.04×10^{-3}

and S.D. 0.18×10^{-3}, and mean 0.5604 and S.D. 0.1122, respectively. We subject each

simulated data set to the same inference procedure as done on the real data. Our estimate of

the S.D. in the estimate is 11.8 for RaTG13 vs. SARS-CoV-2 and 9.41 for RmYN02 vs. SARS-

CoV-2, providing an approximate 95% confidence interval of (28.11, 75.31) and (18.19, 55.85),

respectively. For RaTG13, if including all sites, except the 244-bp in the RBD of the *S* gene

(Supplementary Table 5), the estimate is 55.02 years with an approx. 95% C.I. of (29.4, 80.7).

As more SARS-CoV-2 sequences are being obtained, providing more precise estimates of the

mutation rate, this confidence interval will become narrower. However, we warn that the

estimate is based on a molecular clock assumption and that violations of this assumption

eventually will become a more likely source of error than the statistical uncertainty quantified in

the calculation of the confidence intervals. We also note that, so far, we have assumed no

variation in the mutation rate among synonymous sites. However, just from the analysis of the

300 bp windows, it is clear that is not true. The variance in the estimate of d_S among 300 bp

windows from the RaTG13-SARS-CoV-2 comparison is approximately 0.0113. In contrast, in

the simulated data assuming constant mutation rate, the variance is approximately 0.0034,

suggesting substantial variation in the synonymous mutation rate along the length of the

genome. Alternatively, this might be explained by undetected recombination in the evolutionary

history since the divergence of the strains.

Discussion

The highly skewed distribution of nucleotide frequencies in synonymous sites in SARS-CoV-2

(Kandeel, et al. 2020), along with high divergence, complicates the estimation of synonymous

divergence in SARS-CoV-2 and related viruses. In particular, in the third codon position the

nucleotide frequency of T is 43.5% while it is just 15.7% for C. This resulting codon usage is not

optimized for mammalian cells (e.g, (Chamary, et al. 2006)). A possible explanation is a strong

mutational bias caused by Apolipoprotein B mRNA-editing enzymes (APOBECs) which can

cause Cytosine-to-Uracil changes (Giorgio, et al. 2020).

A consequence of the skewed nucleotide frequencies is a high degree of homoplasy in

synonymous sites that challenges estimates of d_S. We here evaluated estimators of d_S in 300 bp

sliding windows and found that a bias-corrected version of the maximum likelihood estimator

tended to perform best for values of $d_S < 2$. We used this estimator to investigate the

relationship between SARS-CoV-2 and related viruses in sliding windows. We show that

synonymous mutations show shorter divergence to pangolin viruses, than the otherwise most

closely related bat virus, RaTG13, in part of the receptor-binding domain of the *spike* protein.

This strongly suggests that the previously reported amino acid similarity between pangolin

viruses and SARS-CoV-2 is not due to convergent evolution, but more likely is due to

recombination. In the recombination analysis, we identified recombination from pangolin strains

into SARS-CoV-2, which provides further support for the recombination hypothesis. However,

we also find that the synonymous divergence between SARS-CoV-2 and pangolin viruses in this

region is relatively high, which is not consistent with a recent recombination between the two. It

instead suggests that the recombination was into RaTG13 from an unknown strain, rather than

between pangolin viruses and SARS-CoV-2, as proposed in (Boni, et al. 2020). The alternative

explanation of recombination into SARS-CoV-2 from the pangolin virus, would require the

additional assumption of a mutational hotspot to account for the high level of divergence in the

region between SARS-CoV-2 and the donor pangolin viral genome. To fully distinguish between

these hypotheses, additional strains would have to be discovered that either are candidates for

introgression into RaTG13 or can break up the lineage in the phylogenetic tree between

pangolin viruses and RaTG13.

The fact that synonymous divergence to the outgroups, RaTG13 and RmYN02, is not

fully saturated, provides an opportunity for a number of different analyses. First, we can date the

time of the divergence between the bat viruses and SARS-CoV-2 using synonymous mutations

alone. In doing so, we find estimates of 51.71 years (95% C.I., 28.11-75.31) and 37.02 years

(95% C.I., 18.19-55.85), respectively. Most of the uncertainty in these estimates comes from

uncertainty in the estimate of the mutation rate reported for SARS-CoV-2. As more data is being

produced for SARS-CoV-2, the estimate should become more precise and the confidence

interval significantly narrowed. We note that the mutation rate we use here are estimated based

on the entire genome, which may differ from that in non-recombination regions. To address this

problem, we downloaded all the SARS-CoV-2 sequences that are available until 2020-08-17

from GISAID, and obtained an estimate of 1:0.81 for the ratio of mutation rates in the

recombination and non-recombination regions, using the "GTRGAMMA" model implemented in

the RAxML (Stamatakis 2014). Given the length ratio between the two partitions is 1:4, the

difference between the partitions will cause a slight overestimate of the mutation rate by ~5%,

which is relatively small compared to the confidence intervals and the potential for other

unknown sources of uncertainty. However, we warn that a residual cause of unmodeled

statistical uncertainty is deviations from the molecular clock. Variation in the molecular clock

could be modeled statistically (see e.g., (Drummond, et al. 2006) and (Lartillot, et al. 2016)), but

the fact that synonymous mutations are mostly saturated for more divergent viruses that would

be needed to train such models, is a challenge to such efforts. On the positive side, we note that

the estimates of d_S given in Table 3 in general are highly compatible with a constant molecular

clock. Boni et al. (Boni et al. 2020) obtained divergence time estimates similar to ours using a

very different approach based on including more divergent sequences and applying a relaxed

molecular clock. We see the two approaches as being complimentary. In the traditional relaxed

molecular clock approach more divergent sequences are needed that may introduce more

uncertainty due to various idiosyncrasies such as alignment errors. Furthermore, the relaxed

molecular clock uses both synonymous and non-synonymous mutations and is, therefore, more

susceptible to the effects of selection. Our approach allows us to focus on just the relevant in-

group species and to use only synonymous mutations. The disadvantage is that we cannot

accommodate a relaxed molecular clock. However, the fact that both approaches provide

similar estimates is reassuring and suggests that neither idiosyncrasies of divergent sequences,

natural selection, or deviations from a molecular clock has led to grossly misleading conclusions

Another advantage of estimation of synonymous and nonsynonymous rates in the

outgroup lineage, is that it can provide estimates of the mutational load of the current pandemic.

The d_N/d_S ratio is almost 14 times larger in the circulating SARS-CoV-2 strains than in the

outgroup lineage. While some of this difference could possibly be explained by positive

selection acting at a higher rate after zoonotic transfer, it is perhaps more likely that a

substantial proportion of segregating nonsynonymous mutations are deleterious, suggesting a

very high and increasing mutation load in circulating SARS-CoV-2 strains.

Acknowledgements

We are grateful to Dr. E.C Holmes for providing the genome sequence of RmYN02. We thank

Dr. Sergei L Kosakovsky Pond for providing aligned sequences of SARS-CoV-2. We also thank

Dr. Adi Stern for discussion. The research was funded by Koret-UC Berkeley-Tel Aviv University

Initiative in Computational Biology and Bioinformatics to RN.

Data Availability

The pangolin virus sequences, GD410721 and GX_P1E, were downloaded from GISAID with

accession numbers EPI_ISL_410721 and EPI_ISL_410539, respectively, and RmYN02

sequence was provided by E. C. Holmes. All other sequences analyzed in this study were

downloaded from either NCBI Genbank or National Microbiology Data Cente (NMDC). The

accession codes for non-human sequences can be found in Supplementary Table 2 and the

accession codes for human sequences can be found in Supplementary Table 3.

Reference

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J Mol Biol 215:403-410.

Boni MF, Lemey P, Jiang X, Lam TT, Perry BW, Castoe TA, Rambaut A, Robertson DL. 2020. Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. Nat Microbiol. https://doi.org/10.1038/s41564-020-0771-4

Boni MF, Posada D, Feldman MW. 2007. An exact nonparametric method for inferring mosaic structure in sequence triplets. Genetics 176:1035-1047.

Chamary JV, Parmley JL, Hurst LD. 2006. Hearing silence: non-neutral evolution at synonymous sites in mammals. Nat Rev Genet 7:98-108.

Drummond AJ, Ho SY, Phillips MJ, Rambaut A. 2006. Relaxed phylogenetics and dating with confidence. PLoS Biol 4:e88.

Felsenstein J. 2009. PHYLIP (Phylogeny Inference Package) version 3.7a. Distributed by the author. Department of Genome Sciences, University of Washington, Seattle.

Forni D, Cagliani R, Clerici M, Sironi M. 2017. Molecular Evolution of Human Coronavirus Genomes. Trends Microbiol 25:35-48.

Gibbs MJ, Armstrong JS, Gibbs AJ. 2000. Sister-scanning: a Monte Carlo procedure for assessing signals in recombinant sequences. Bioinformatics 16:573-582.

Di Giorgio S, Martignano F, Torcia MG, Mattiuz G, Conticello SG. 2020. Evidence for host-dependent RNA editing in the transcriptome of SARS-CoV-2. Sci Adv 6:eabb5813.

Goldman N, Yang Z. 1994. A codon-based model of nucleotide substitution for protein-coding
DNA sequences. Mol Biol Evol 11:725-736.

Hu D, Zhu C, Ai L, He T, Wang Y, Ye F, Yang L, Ding C, Zhu X, Lv R, et al. 2018. Genomic
characterization and infectivity of a novel SARS-like coronavirus in Chinese bats. Emerg
Microbes Infect 7:154.

Kandeel M, Ibrahim A, Fayez M, Al-Nazawi M. 2020. From SARS and MERS CoVs to SARS-
CoV-2: Moving toward more biased codon usage in viral structural and nonstructural genes. J
Med Virol.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30:772-780.

Lam TT, Shum MH, Zhu HC, Tong YG, Ni XB, Liao YS, Wei W, Cheung WY, Li WJ, Li LF, et al.
2020. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. Nature.

Lartillot N, Phillips MJ, Ronquist F. 2016. A mixed relaxed clock model. Philos Trans R Soc
Lond B Biol Sci 371.

Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KSM, Lau EHY, Wong JY, et al.
2020. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected
Pneumonia. N Engl J Med 382:1199-1207.

Li X, Giorgi EE, Marichannegowda MH, Foley B, Xiao C, Kong X-P, Chen Y, Gnanakaran S,
Korber B, Gao F. 2020. Emergence of SARS-CoV-2 through recombination and strong purifying
selection. Sci. Adv. 6:eabb9153.

Loytynoja A. 2014. Phylogeny-aware alignment with PRANK. Methods Mol Biol 1079:155-170.

Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, Wang W, Song H, Huang B, Zhu N, et al. 2020.
Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus
origins and receptor binding. Lancet 395:565-574.

Martin D, Rybicki E. 2000. RDP: detection of recombination amongst aligned sequences.
Bioinformatics 16:562-563.

Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. 2015. RDP4: Detection and analysis of
recombination patterns in virus genomes. Virus Evol 1:vev003.

Muse SV, Gaut BS. 1994. A likelihood approach for comparing synonymous and

nonsynonymous nucleotide substitution rates, with application to the chloroplast genome. Mol

Biol Evol 11:715-724.

Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32:268-274.

Padidam M, Sawyer S, Fauquet CM. 1999. Possible emergence of new geminiviruses by

frequent recombination. Virology 265:218-225.

Patiño-Galindo JÁ, Filip I, AlQuraishi M, Rabadan R. 2020. Recombination and lineage-specific

mutations led to the emergence of SARS-CoV-2. bioRxiv DOI: 10.1101/2020.02.10.942748.

Posada D, Crandall KA. 2001. Evaluation of methods for detecting recombination from DNA

sequences: computer simulations. Proc Natl Acad Sci U S A 98:13757-13762.

Salminen MO, Carr JK, Burke DS, McCutchan FE. 1995. Identification of breakpoints in

intergenotypic recombinants of HIV type 1 by bootscanning. AIDS Res Hum Retroviruses

11:1423-1425.

Smith JM. 1992. Analyzing the mosaic structure of genes. J Mol Evol 34:126-129.

Valitutto MT, Aung O, Tun KYN, Vodzak ME, Zimmerman D, Yu JH, Win YT, Maw MT, Thein

WZ, Win HH, et al. 2020. Detection of novel coronaviruses in bats in Myanmar. PLoS One

15:e0230802.

Wan Y, Shang J, Graham R, Baric RS, Li F. 2020. Receptor Recognition by the Novel

Coronavirus from Wuhan: an Analysis Based on Decade-Long Structural Studies of SARS

Coronavirus. J Virol 94.

Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, et al.

2020. A new coronavirus associated with human respiratory disease in China. Nature 579:265-

269.

Xiao K, Zhai J, Feng Y, Zhou N, Zhang X, Zou JJ, Li N, Guo Y, Li X, Shen X, et al. 2020.

Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins. Nature 583:286-289.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-

1591.

Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution rates under

realistic evolutionary models. Mol Biol Evol 17:32-43.

Yang Z, Nielsen R, Goldman N, Pedersen AM. 2000. Codon-substitution models for

heterogeneous selection pressure at amino acid sites. Genetics 155:431-449.

Zhang T, Wu Q, Zhang Z. 2020. Probable Pangolin Origin of SARS-CoV-2 Associated with the

COVID-19 Outbreak. Curr Biol 30:1346-1351 e1342.

Zhang YZ, Holmes EC. 2020. A Genomic Perspective on the Origin and Emergence of SARS-

CoV-2. Cell 181:223-227.

Zhou H, Chen X, Hu T, Li J, Song H, Liu Y, Wang P, Liu D, Yang J, Holmes EC, et al. 2020. A

Novel Bat Coronavirus Closely Related to SARS-CoV-2 Contains Natural Insertions at the

S1/S2 Cleavage Site of the Spike Protein. Curr Biol 30:2196-2203 e2193.

Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, et al.

2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature

579:270-273.

Figure 1. Genome-wide identity plot and top blast hits for SARS-CoV-2, RaTG13 and RmYN02. (a) 300 bp sliding-windows of nucleotide identity between SARS-CoV-2 and the four most closely related viral strains, RmYN02, RaTG13, GD410721 and GX_P1E. Orange shading

marks the recombinant region in SARS-CoV-2 inferred by 3SEQ (details in Supplementary Table 5). (b) the plot lists all the viral strains that are the unique best BLAST hit in at least three 100-bp windows, when blasting with SARS-CoV-2, with the regions where each strain is the top blast hit marked. (b) and (c). Figures for RaTG13 (c, d) and RmYN02 (e, f) generated in the same way as for SARS-CoV-2 in (a) and (b). The *ACE2* contact residues of RBD region (left) and the furin sites (right) of the *S* protein are marked in both plots with grey lines.

Figure 2. Unrooted phylogenies of the virus strains. (a) Maximum-likelihood tree in genomic regions with recombination tracts removed; (b) Neighbor-joining tree using synonymous mutation (d_S) distance in genomic regions with recombination tracts removed; (c) Neighbor-joining tree using non-synonymous mutation (d_N) distances in genomic regions with recombination tracts removed; (d) The maximum-likelihoods tree at the receptor-binding domain *ACE2* contact residues (51 amino acids) region. The bootstrap values are based on 1,000 replicates. The associated distance matrix for (b) and (c) can be found in Table 3.

Figure 3. Bias correction for d_S estimate in 300-bp windows. (a) The mean of d_S estimates using different methods; ML.corr and yn00.corr are the bias corrected versions of the ML and yn00 methods, respectively. (b) Errors in d_S estimates as measured using the ratio of square root of mean squared error (MSE) to true d_S. All the estimates are based on 10,000 simulations. ML: maximum-likelihood estimates using the f3x4 model in codeml; ML.corr, maximum-likelihood estimates with bias correction; yn00, count-based estimates in (Yang and Nielsen 2000); yn00.corr, yn00 estimates with bias correction. All d_S estimates are truncated at 3, explaining the reduction in MSE with increasing values of d_S as d_S approaches 3.

Figure 4. d_S and d_N estimates across the virus genome. (a) Pairwise d_S estimates in 300-bp sliding windows for RaTG13, GD410721 and Wuhan-Hu-1, the estimates are truncated at 4. (b) d_S ratio of d_S (Wuhan-Hu-1,RaTG13) to d_S(Wuhan-Hu-1,GD410721). (c) and (d) are the zoom-in plot for d_S and d_S-ratio at the *spike* (S) protein region. The receptor-binding domain contact residues (left) and furin site regions (right) are marked with grey lines. (e) the pairwise d_N estimates in 300-bp sliding windows in the *S* protein for these strains. The d_S values are truncated at 4 in the plots.

Table 1. Genome-wide nucleotide composition at the third position of the codons in the viral strains. The nucletodie compositions at the first and second positions can be found in Supplementary table 1.

Accession	T	C	A	G
GD410721	42.71%	16.17%	28.55%	12.57%
GX_P1E	42.52%	16.40%	28.27%	12.81%
RaTG13	43.57%	15.74%	27.98%	12.71%
RmYN02	43.31%	15.90%	27.98%	12.81%
Wuhan-Hu-1	43.49%	15.73%	28.16%	12.62%

Table 2. Whole genome d_N and d_S estimates among the viral strains. The d_S estimates are shaded in green, and the d_N estimates are in orange shade. The 95% confidence intervals, calculated based on 1,000 bootstrap replicates, are included in the brackets for each estimates.

	GD410721	GX_P1E	RaTG13	RmYN02	Wuhan-Hu-1
GD410721		0.0372 (0.0341-0.0403)	0.0171 (0.0152-0.0190)	0.0293 (0.0266-0.0320)	0.0160 (0.0142-0.0178)
GX_P1E	0.9883 (0.9338-1.0428)		0.0347 (0.0318-0.0376)	0.0485 (0.0450-0.0520)	0.0342 (0.0314-0.0370)
RaTG13	0.5392 (0.5105-0.5679)	1.0156 (0.9608-1.0704)		0.0235 (0.0210-0.0260)	0.0065 (0.0053-0.0077)
RmYN02	0.6001 (0.5681-0.6321)	1.0757 (1.0166-1.1348)	0.2438 (0.2285-0.2591)		0.0220 (0.0195-0.0245)
Wuhan-Hu-1	0.5425 (0.5131-0.5719)	0.9973 (0.9434-1.0512)	0.1604 (0.1491-0.1717)	0.2043 (0.1901-0.2185)	

Table 3. Genome-wide d_N and d_S estimates after removing recombination regions inferred by 3SEQ . The d_S estimates are shaded in green, and the d_N estimates are in orange shade. The coordinates relative to the Wuhan-Hu-1 genome of the masked region can be found in the method section. The 95% confidence intervals, calculated based on 1,000 bootstrap replicates, are included in the brackets for each estimates.

	GD410721	GX_P1E	RaTG13	RmYN02	Wuhan-Hu-1
GD410721		0.0348 (0.0317-0.0379)	0.0138 (0.0120-0.0156)	0.0152 (0.0133-0.0171)	0.0135 (0.0117-0.0153)
GX_P1E	0.9974 (0.9381-1.0567)		0.0357 (0.0325-0.0389)	0.0361 (0.0329-0.0393)	0.0349 (0.0318-0.0380)
RaTG13	0.4962 (0.4669-0.5255)	1.0366 (0.9737-1.0995)		0.0079 (0.0066-0.0092)	0.0060 (0.0048-0.0071)
RmYN02	0.5070 (0.4773-0.5366)	1.0333 (0.9699-1.0967)	0.1522 (0.1395-0.1649)		0.0062 (0.0050-0.0074)
Wuhan-Hu-1	0.5095 (0.4794-0.5396)	1.0304 (0.9669-1.0939)	0.1462 (0.1340-0.1584)	0.1117 (0.1019-0.1215)	