Antibiotics may influence gut microbiome signaling to the brain in preterm neonates

Jordan T. Russell[1]†, J. Lauren Ruoss[2]†, Diomel de la Cruz[2], Nan Li[2], Catalina Bazacliu[2], Laura

Patton[2], Kelley Lobean McKinley[1], Timothy J. Garrett[3], Richard A. Polin[4], Eric W. Triplett[1]*,

and Josef Neu[2]

[1]Department of Microbiology and Cell Science, Institute of Food and Agricultural Sciences,

University of Florida, Gainesville, FL, USA.

[2]Division of Neonatology, Department of Pediatrics, University of Florida, Gainesville, FL,

USA.

[3]Department of Pathology, Immunology and Laboratory Medicine, College of Medicine,

University of Florida, Gainesville, FL, USA.

[4]Department of Pediatrics, College of Physicians and Surgeons, Columbia University, New

York, NY, USA.

*Corresponding author: Eric W. Triplett, PhD (email: ewt@ufl.edu)

†These authors contributed equally to this work

Abstract

Antibiotic use in neonates can have detrimental effects on the developing gut microbiome,

increasing the risk of morbidity. A majority of preterm neonates receive antibiotics after birth

without clear evidence to guide this practice. Here we present microbiome and metabolomic

results from the Routine Early Antibiotic use in SymptOmatic preterm Neonates (REASON)

study, which is the first trial to randomize symptomatic preterm neonates to receive or not

receive antibiotics in the first 48 hours after birth. Using 16S rRNA sequencing of stool samples

collected longitudinally for 91 neonates, we show that antibiotics alter trends in microbiome

diversity and development between randomized neonates and illustrate that type of nutrition

helps shape the early infant gut microbiome. Integrating multi-omic data for the gut microbiome,

stool immune markers, stool metabolites, and inferred metabolic pathways, we identified an

association between *Veillonella* and the neurotransmitter gamma-aminobutyric acid (GABA).

Our results suggest early antibiotic use may impact the gut-brain axis with the potential for

consequences in early life development.

Main

Premature infants are particularly susceptible to infections secondary to increased need for

invasive procedures and immaturity of the immune system, skin, and gastrointestinal tract[1–3].

Increasingly, there is growing concern that risk factors for mortality may originate from

underlying pathologies that could also be responsible for premature birth[4]. Symptoms of

prematurity are difficult to discern from symptoms of infection which, compounded by the

increased risk of infection, have led to most premature infants being exposed to antibiotics early

in life[5–7]. Despite high mortality rates, the incidence of culture positive early onset sepsis (EOS)

is relatively low, between 0.2-0.6%[8]. In the absence of a positive culture, a majority of preterm

infants receive antibiotics immediately after birth based on maternal risk factors (e.g. intra-

amniotic infection) or laboratory abnormalities (e.g. elevated serum C-reactive protein (CRP)

because of the risk of mortality[8]. Given the low incidence of culture positive EOS in this

population, it is possible that such high rates of antibiotic use are unnecessary and may increase

morbidity in these infants[9]. Other morbidities in the neonatal intensive care unit (NICU) such as

necrotizing enterocolitis (NEC) also have high mortality rates and have been associated with

prolonged antibiotic exposure.[10,11]. Nevertheless, antibiotics remain the most commonly

prescribed medication in the NICU[12,13].

The gut microbiome comprises a highly volatile community structure early in life[14]. Microbial

colonization is influenced as early as birth by mode of delivery, and perhaps even in the uterine

environment by maternal factors[15,16]. Not surprisingly, antibiotic use has been shown to also

change the composition of the preterm gut community[17–20]. Furthermore, antibiotic use early in

life has increasingly been associated with adverse outcomes both short- and long-term[21,22]. One

possible consequence is the disruption of the gut-brain axis (GBA), which involves bi-directional

transmission of bio-molecular signals between the gut microbiota and the nervous system[23].

Aberrations in the GBA have been associated with altered immune homeostasis, as well as

psychiatric, behavioral and metabolic conditions in adulthood[24]. It is therefore imperative to

determine if such high rates of antibiotic use in preterm infants is necessary, as it could have

lifelong consequences on future health.

Randomized clinical trials have the advantage of controlling for many of the numerous

covariates that could interfere with answering whether preemptive antibiotic use in preterm

infants affects outcomes. The Routine Early Antibiotic use in SymptOmatic preterm Neonates

(REASON) study is the first to randomize symptomatic premature infants to either receive or not

receive antibiotics soon after birth. Previously reported results from this study demonstrate the

feasibility of such a trial and that withholding antibiotics did not lead to a significant increase in

neonatal mortality or morbidity[25]. By employing a multi-omic approach, this randomized trial

also provides the unique opportunity to understand how antibiotic intervention perturbs the early

life gut microbiome, metabolome, and inflammatory environment in ways that may be

consequential to health and development.

Results

Cohort and study description

Ninety-eight infants were enrolled in the study over the two-year period: group A – antibiotics

indicated (n=32), group B – antibiotics not indicated (n=11), and group C – eligible for

randomization (n=55). Twenty-eight infants were randomized to C1 (antibiotics x 48 hours) and

27 infants were randomized to group C2 (no antibiotics). For 13 infants (48%) in group C2,

antibiotics were prescribed in the first 48 hours after birth. Neither sex nor mode of delivery

were significantly different between all groups and were not significantly different between the

two randomized subgroups using the chi-square test. Both weight (p-value=0.0001) and

gestational age (p-value=0.0002) were significantly different between groups overall, with group

A infants on average with lower gestational age (GA) and at lower birth weights. Neither birth

weight nor GA were significantly different between the randomized subgroups by ANOVA. A

summary of the infants in this analysis is provided in Table 1. A full description of enrollment

has been described previously[25].

Six hundred ninety-three stool samples were collected longitudinally for 91 of the total 98 infants

enrolled. Stool data was not available for 7 infants due to early mortality. Sequencing data for

16S rRNA was able to be obtained for 656 of those samples. After rarefying to an even

sequencing depth of 10,000 reads per sample, 642 samples remained for further analysis. The

breakdown of samples by group is provided in Table 1. The aim for this analysis is to test the

effects of randomization to antibiotics vs. no antibiotics has on the developing gut microbiome,

metabolome and inflammatory environment using high-throughput 16S rRNA sequencing,

quantitative PCR (qPCR), metabolomics, pathway inference, immune marker analysis and open-

source statistical tools.

Antibiotic use affects trends in early gut microbiome development over time

The within-sample diversity differed by enrollment groups A, B, C1, and C2. Using amplicon

sequencing variants (ASVs), the richness, and the alpha diversity were not significantly different

between enrollment groups overall (Kruskal-Wallis, Richness: $p = 0.31$, Shannon: $p = 0.90$) and

did not differ across time using a linear mixed effects model (LME) through Qiime2[26–28]

(Shannon p-values: group A = 0.98, group B = 0.70, group C1 = 0.83, group C2 = 0.926).

Though not statistically significant, scatter plots of each sample's alpha diversity, including

regression lines for each group, indicate trends in diversity development over time between the

groups (Fig. 1a, Shannon diversity). These trends are expected given the variation in antibiotic

levels used by each group. For example, group B (antibiotics not indicated) tended to have the

highest trend in diversity over time and is the group with the lowest antibiotic use. In contrast,

group A (antibiotics indicated) infants saw very little diversity development over time while

receiving the most antibiotics. Certain considerations need to be made when comparing infants

by enrollment group. For example, group A infants had significantly lower GA and birth

weights, particularly compared to group B infants who on average had the highest GAs and birth

weights, and the shortest stays in the NICU. Furthermore, infants in randomized group C2 who

were changed to receive antibiotics in the first 48 hours after birth can no longer be considered as

receiving no antibiotics and would be more similar to group C1. To account for this in testing the

effect of antibiotic use, we focus only on the randomization group, taking into consideration

those infants who were changed to receive antibiotics by instead separating infants based on any

antibiotic use within 48 hours after birth (any_abx).

Samples from group C infants that were collected during antibiotic use (abx_during_sample =

TRUE) were compared with those during periods of no treatment (abx_during_sample =

FALSE). As anticipated, samples taken during periods of antibiotic use had lower richness and

lower bacterial load, as determined by quantitative PCR (qPCR) of 16S rRNA copies per gram of

stool (Figs. 1b,c). As this result may be driven by samples collected early in life, just those

samples collected in the first week of life were also compared and were not significantly

different in richness (Wilcoxon, $p = 0.70$), alpha diversity (Wilcoxon, $p = 0.43$) and bacterial

load (Wilcoxon, $p = 0.50$).

Taking into consideration those infants in group C2 who were changed to antibiotic use, bacterial

richness was plotted as a function of time (Collection_week) considering whether the infant

received antibiotics (any_abx) (Fig. 1d). Pairwise comparisons of the antibiotic and no antibiotic

infants at each collection week were not statistically significant (Wilcoxon test, $p > 0.05$).

Initially within the first few weeks of life, bacterial richness is similar regardless of antibiotic

use. However, beginning at the fifth collection week (30 days after birth on average) infants who

never received antibiotics trended towards higher bacterial richness compared to infants that

were treated. Average bacterial richness remained higher among the no antibiotic infants through

collection week 8 (53 days after birth on average). Again using linear mixed effects modelling,

we found that both bacterial richness and Shannon diversity trended higher in infants who never

received antibiotics compared with those that did, though neither diversity metric was significant

as a function of time (Richness: $p = 0.35$, Shannon: $p = 0.26$) (Figs. 1e,f). As with the infants in

group C, infants who were randomized to not receive antibiotics and were not changed to receive

antibiotics (any_abx = FALSE) had shorter stays in the NICU. Furthermore, although the

randomized groups are almost evenly powered without considering change in antibiotic use

status, removing C2 infants who received antibiotics underpowers the group of infants who never

received antibiotics, limiting the strength of comparisons.

Changes in overall bacterial community structure were apparent between groups when assessing

beta-diversity. Both the Bray-Curtis and Jaccard distance indices were used to assess community

structure taking into account quantitative ASV abundance and qualitative presence/absence

information, respectively. Beta dispersion was not significantly different between groups using

the Bray-Curtis metric (ANOVA; Df = 3, Sum of Squares = 0.01589, Mean Squares =

0.0052974, F = 2.547, p = 0.055) but was significant using Jaccard (ANOVA: Df = 3, Sum of

Squares = 0.0315, Mean Squares = 0.01050, F = 8.4013, p = 1.75E-05), specifically group A

versus all other groups (TukeyHSD, B-A p = 0.0153, C1-A p-value = 0.0001, C2-A p = 0.0028).

With the test assumption met for the Bray-Curtis index, PERMANOVA also showed distance

dissimilarity to be significant between groups (PERMANOVA; Df = 3, Sum of Squares = 4.642,

R_2 = 0.01632, F = 3.544, p = 0.001). Principle coordinates analysis (PCoA) suggests an effect of

antibiotic use by enrollment group. The infants exposed to the least amount of antibiotics

through the NICU course (group B and group C2 that were not changed to receive antibiotics)

had tighter clustering using both Bray-Curtis and Jaccard, whereas groups A and C1 had the

highest variability, suggesting less antibiotic use led to greater microbiome stability (Figs. 2a,b).

Similarly, group C1 had greater spread by PCoA using Bray-Curtis but not Jaccard (Figs. 2c,d).

Infants in group C2 who were changed to receive antibiotics in the first 48 hours but did not

receive antibiotics again had much tighter clustering, indicating the outcome of early antibiotics

in the randomized group and the effect antibiotics use has on microbiome stability (Figs. 2e,f).

Feeding patterns drive changes in gut diversity and bacterial load

For preterm infants, diet generally consists of mother's breast milk, pasteurized donor breast

milk, and/or formula. To control for any effect of antibiotics, only samples from group C2

infants that were not changed to receive antibiotics in the first 48 hours were used. Using the

calculated alpha diversity metrics used previously, feeding type was found to significantly

associated with bacterial community richness (Kruskal-Wallis, p = 0.0016) and Shannon

diversity (Kruskal-Wallis, p = 6.2E-05) (Figs. 3a,b). In addition, feeding type had an effect on

total bacterial load measured by the number of 16S rRNA copies per gram of stool for each

sample (Kruskal-Wallis, p = 0.003) (Fig. 3c). As expected, stool collected during times of no

enteral feeding (NPO) had the lowest richness and total bacterial load. Samples collected during

formula feeding had just as high of richness and diversity as samples during feeding with

mother's milk. However, samples collected during feeding with mother's milk tended to have the

higher bacterial load (Wilcoxon, $p = 0.0057$). Donor breast milk (DBM), which is pasteurized

prior to use, tended to have lower richness, diversity, and bacterial load compared to other

feeding types except NPO. Interestingly however, combining formula with DBM trended

towards increased diversity and bacterial load, though not significantly, most likely due to the

limited number of samples collected during feeding with DBM + formula from group C (n= 2).

Clustering analysis by PCoA also showed trends in gut composition based on feeding type (Figs.

3d,e). Stool collected while feeding with mother's milk had the widest variability in overall

composition by both Bray-Curtis and Jaccard, whereas donor milk had the tightest clustering.

Notably, samples taken during formula or formula mixed with mother's milk had similar

clustering by Bray-Curtis, but formula feeding alone led to more variability with Jaccard,

suggesting addition of mother's milk had little effect on the abundance of bacteria but formula

feeding did contribute more variability in the presence or absence of certain bacteria. Not

considering samples from infants in group C2 who were changed to antibiotics, feeding type had

a significant effect on community structure by both Bray-Curtis (PERMANOVA, Df = 7, Sum of

Squares = 5.527, R_2 = 0.133, F = 1.98, $p = 0.001$) and Jaccard (PERMANOVA, Df = 7, Sum of

Squares = 4.692, R_2 = 0.118, F = 1.720, $p = 0.001$). Along the same lines, pairwise comparisons

identified significant differences in beta diversity between mother's milk vs. formula, donor milk

vs. formula, mother's milk vs. donor milk, and mother's milk vs. donor milk mixed with formula

(PERMANOVA, $p < 0.05$). These results indicate feeding type as a significant source of

variation in the composition of the preterm infant gut in addition to antibiotic use.

Gut microbial community development is highly variable and unique to each infant

Although the infants in this study were enrolled into 4 groups, each infant's stay in the NICU is

highly personalized, by type, frequency, number or length of antibiotic use, type and length of

feeding patterns, and adverse clinical events over time. To aid in visual identification of patterns

throughout the NICU course, detailed charts were created for each infant that depict both clinical

and laboratory data over time (Days Post Birth) integrated into a single graphic per infant (Fig. 4

and Supplementary Figure 1). This includes results from the 16S rRNA analysis as pie charts for

each stool sample, color coded by bacterial taxonomy and sized based on the $\log_{10}$-transformed

number of 16S rRNA copies per gram of stool, as well as adverse clinical events coded by a

single letter code (Fig. 5). With these visualizations, patterns are more easily observed between

antibiotic treatments, feeding types, and the gut microbiome. For example, anti-fungal

fluconazole administration resulted in lower bacterial loads and diversity. Also, the effects of

antibiotic use and feeding patterns become visually apparent. To illustrate, infant 5 was

exclusively fed mother's milk from day 14 through day 71 post birth. At day 31, antibiotics

vancomycin and piperacillin were administered (Supplementary Figure 1). During treatment,

Veillonella was entirely removed from the stool microbiome, falling from 3.23E+06 cells per

gram (73%) from the pre-treatment time point to undetected while antibiotics were used. At day

47, 9 days after antibiotic treatment ceased, *Veillonella* again dominated the stool at 3.83E+06

cells per gram (80%), almost a complete replacement of levels before treatment. Also, the

proportions of the other 2 genera found in the stool, *Escherichia* and an unclassified

Enterobacteriaceae spp., were nearly identical after treatment and continuation of mother's milk

as before treatment (pre-treatment: *Escherichia* – 21%, Enterobacteriaceae spp. – 4.7%; post-

treatment: *Escherichia* – 15%, Enterobacteriaceae spp. – 3.3%), demonstrating that mother's

milk effectively restored the stool microbiome to its pre-treatment state.

Bacterial genera correlate with stool metabolites and inferred metabolic pathways

In addition to 16S rRNA profiling, 90 stool samples from 10 infants were sent for metabolomic

profiling. Each of the four enrollment groups were included in these samples for comparison.

Peak height responses were recorded for 454 identifiable metabolites. To determine if gut

bacteria were associated with relative concentrations of metabolites in stool, the top 10 most

abundant bacterial genera associated with identified metabolites were determined. Spearman

correlation values were plotted using a heatmap, which indicated numerous significant, positive

and negative, associations between bacteria and metabolites (Fig. 6a). Next, 6 of the top 10

bacterial genera were correlated between infant groups (Fig. 6b). Interestingly, *Veillonella* was

positively associated with the neurotransmitter 4-aminobutanoate (GABA) (Spearman R = 0.48,

p = 0.003) and *Veillonella* counts were significantly different between groups A and C2

(Wilcoxon, p = 0.0475). *Veillonella* counts were plotted using a boxplot, along with the

normalized concentrations of GABA between groups (Figs. 6c,d). Furthermore, using

PICRUSt2, functional pathway abundances were inferred based on the rarefied 16S rRNA data[29].

The *Veillonella* counts of predicted pathways were strongly correlated with biosynthesis of the

GABA precursor L-glutamate and L-glutamine (Spearman R = 0.88, p = 3.02E-27) (Fig. 7).

Thus, *Veillonella* may be at least partially responsible for GABA neurotransmitter production

and that this function is negatively impacted by antibiotic use early in life. Alternatively, a

negative correlation between *Bifidobacterium* counts and glycocholic acid was observed

(Spearman R = -0.39, p = 0.0098), which was impacted by antibiotic use between groups

(Supplementary Figure 2). Furthermore, bifidobacteria were negatively associated with other

conjugated bile acids including taurocholic (Spearman R = -0.45, p = 0.0021) and glycocholic

acids (Spearman R = -0.39, p = 0.0098), but positively associated with deconjugated cholic

(Spearman R = 0.36, p = 0.0203). Thus, gut microbiota affected by antibiotic use may be

responsible for modification of neuroactive metabolites (i.e. deconjugated bile salts) in addition

to production of neurotransmitters.

Immune markers in stool correlate with bacterial abundance

Antibiotic use was examined for its correlation with inflammatory marker level. These levels

were also correlated with gut bacterial abundances. Twelve immune markers were measured in

110 samples across 18 of the first enrolled infants. Twelve bacterial genera had at least one

significant correlation with an immune marker (p < 0.05) (Fig. 8a). Significant correlations

between the bacterial genera and stool immune markers were classified as either inflammatory or

anti-inflammatory based on the known function of the marker (Fig. 8b). Interestingly,

Enterococcus was consistently negatively correlated with most of the immune markers

measured. *Enterococcus* counts were lower in group A compared to the randomized groups

(Wilcoxon, A vs. C1 p = 1.6E-06, A vs. C2 p = 9.4E-07) (Fig. 8c). *Enterococcus* counts did not

differ significantly between infants who received or did not receive antibiotics (Wilcoxon, p =

0.2) (Fig. 8d). *Enterococcus* had a significant negative correlation with IL8:IL10 ratios, whereas

this genus was positively correlated with IL1a:IL1ra ratios, suggesting that mechanisms of

controlling inflammation can be unique depending on the microbe. Notably, MIP1a was the only

marker that differed significantly depending on antibiotic use (Wilcoxon, p = 0.039). *Citrobacter*

was positively correlated with MIP1a (Spearman R = 0.47, p = 3.74E-05), and was significantly

higher in group C1 compared to group C2 (Wilcoxon, p = 7.2E-08). *Citrobacter* was higher in

those infants that received antibiotics compared to those that did not (Wilcoxon, p = 0.037).

Discussion

There is an urgent need for evidence supporting or refuting the widespread practice of routine

antibiotic use after birth in symptomatic preterm neonates. The REASON study represents a

significant step as it is the first randomized controlled trial to test the feasibility of randomizing

symptomatic preterm infants to antibiotics versus no antibiotics, evaluating the effect of

antibiotic treatment on the developing gut microbiome, metabolome, and inflammatory

environment. Our results expand upon previous reports with new evidence that early routine

antibiotic use leads to greater beta-diversity dissimilarity and lowered trends in richness and

Shannon diversity development over time, even after discontinuation of antibiotics[17,30,31]. The

power to detect significant associations was decreased mainly because many of the infants

randomized to not receive antibiotics were changed to antibiotic administration. A larger multi-

center randomized study is needed to validate and expand upon the extended effect of antibiotics

on the developing gut microbiome.

Our results confirm that feeding type also have a significant influence on gut microbiome

richness, diversity and bacterial load[32-34]. Exclusive or partial feeding with mother's own milk

tended to associate with high bacterial load, which is supported by evidence that breast milk

harbors maternal-originating bacteria, as well as nutritional components (prebiotics) that support

bacterial proliferation in the intestinal tract[35,36]. Interestingly, formula-fed infants had

comparable levels of richness, diversity and bacterial load as mother's milk. This supports the

notion that mother's milk drives early colonization of a limited set of dominant microbes through

nutrient and antimicrobial-mediated selection[37–39].

Integrating detailed and personalized records of clinical and laboratory data led us to identify

overlooked patterns in the data. One such peculiar pattern was that stool samples taken during

administration of the anti-fungal fluconazole had lowered copies of 16S rRNA, suggesting lower

bacterial load. A previous study reported that fluconazole, though not inherently bactericidal,

increased the bactericidal activity of neutrophils[40]. Immune marker data were collected for 18 of

the first enrolled infants, and one or more of those markers, such as calprotectin which is

secreted by neutrophils, may help explain this pattern[41]. However, unfortunately only 3 of the 18

infants that had immune marker data received fluconazole, and therefore more data is needed to

test this hypothesis. Interestingly, counts of *Enterococcus* were often negatively correlated with

levels of many of the immune markers tested, both pro- and anti-inflammatory, an odd finding

considering Enterococci have been associated with risk for infection in preterm neonates[42,43]. On

the other hand, *Citrobacter* counts were associated with increased levels of the macrophage

chemokine MIP1α and counts were significantly higher in infants who received antibiotics.

Increased levels of MIP1α are likely related to recruitment of intestinal macrophages leading to a

heightened inflammatory environment, suggesting that antibiotic use may select for bacteria

which lead to intestinal inflammation[44]. Further work including a larger sample size will be

needed to understand how changes in the preterm infant gut caused by routine antibiotics impacts

the gut inflammatory environment.

Neurological development can be impaired in infants born very prematurely compared to their

full-term counterparts; a trend that extends into delayed cognitive and behavioral development

through childhood[45–47]. Could routine antibiotic use in preterm neonates play a role in this

association? Intestinal microbes produce a plethora of metabolites and bio-active compounds that

can be absorbed by the host[48]. Some of these compounds have direct neurologic implications

including neurotransmitters such as GABA, which is reduced in preterm infants, is critical for

early brain development, and possesses immunomodulatory properties[49,50]. Our results indicate

that antibiotic use negatively affected the abundance of *Veillonella* and that *Veillonella* were

positively correlated with GABA concentrations in the gut. Furthermore, *Veillonella* correlated

strongly with the L-glutamine biosynthesis pathway, the precursor to GABA production. Aside

from production of neurotransmitters, we identified negative correlations between

Bifidobacterium abundance and concentrations of conjugated bile acids, particularly glyco- and

taurocholic acid, which were significantly different based on antibiotic use. Bifidobacteria,

which were more abundant in infants that did not receive antibiotics, are known to deconjugate

bile acids to primary forms including cholic acid, which was positively correlated with

bifidobacteria abundance[51,52]. Cholic acid can passively diffuse into the brain where it blocks

signaling in the GABA$_A$ receptor[53]. Bifidobacteria may therefore be essential in regulating

GABA signaling in the developing brain. These are significant findings, for they suggest routine

antibiotic use could be disrupting processes involved in the gut-brain axis and

immunomodulatory pathways critical for neonatal and future childhood development.

Evidence-based antibiotic use to prevent infection in preterm neonates is critical in preventing

unnecessary treatment that may be doing more harm than good. Routine overuse of antibiotics

can change the developmental trajectory of the infant gut microbiome during a time of critical

establishment and interaction. Given the potential for extensive crosstalk between gut microbiota

and the host, changes in microbiome composition could have both short- and long-term effects

on outcomes and overall health and development. Future randomized studies with greater infant

enrollment will be crucial in our understanding of the effects current neonatal practice has on

health which will allow for the reevaluation of practices. Such trials will need to expand on the

findings from this pilot study from a multi-omic standpoint to identify direct links between

antibiotic-induced dysbiosis and health outcomes.

Materials and Methods

Experimental design, enrollment, and clinical sample and data collection

The REASON study was conducted from January 2017 - January 2019 at the University of

Florida and was approved by the institutional review board (IRB201501045). This study is

funded by the NIH (R21HD088005). A detailed description of the study design including

enrollment, group selection, randomization, and collection of clinical samples and data including

outcomes has been previously described[25]. Briefly, 98 premature infants were enrolled in the

study and placed into one of three groups according to previously described criteria: group A

with indication for antibiotic use, group B without indication for antibiotic use, and group C

eligible for randomization to antibiotics (C1) or no antibiotics (C2) in the first 48 hours after

birth. Infants not receiving antibiotics in the first 48 hours after birth (group B, C2) could be

changed to receive antibiotics at any time at the medical team's discretion. Clinical samples

relevant to this analysis include weekly fecal collection starting with meconium when possible

(all stored at -80°C) and results of bacterial and fungal cultures (blood, urine, sputum, and

cerebrospinal fluid - when available) and laboratory measurements of CRP, white blood cell

count and immature to neutrophil ratio. Clinical metadata from the mothers such as antepartum

antibiotic use, type, duration, and proximity to delivery were recorded. Pertinent clinical

metadata from the infants include group placement, antibiotic use status, antibiotics and

antifungal use including type and duration throughout NICU course, feeding type and duration,

GA at birth, sex, mode of delivery and any serious adverse events (SAEs) including NEC, late

onset sepsis, spontaneous intestinal perforations, bronchopulmonary dysplasia, and death.

Stool DNA extraction, 16S rRNA PCR and Sequencing Analysis

DNA extraction and 16S rRNA barcoded PCR was carried out exactly as described previously[54].

Approximately 60 gigabases of nucleotide sequencing data was generated across 5 Illumina

Miseq flowcells for stool samples collected from 91 (of the 98 total) study participants where

samples were collected (ICBR, Gainesville, FL, USA). The resulting sequencing reads were

merged, demultiplexed, trimmed, filtered for quality and processed into amplicon sequencing

variants (ASVs) as previously described with no alterations in method[54]. Briefly, sequences were

processed to ASVs using the DADA2 package in R (https://www.R-project.org) and assigned

taxonomy using the SILVA_v132 training dataset[55–59]. Samples were rarefied to 10,000 reads per

sample, leaving 642 of the total 656 individual longitudinal stool samples for analysis.

Total bacterial quantification by universal 16S qPCR

Total bacterial load per gram of stool was determined by universal 16S rRNA qPCR using the

same primer set used for amplicon sequencing (341F and 806R). QPCR assays were performed

on a QuantStudio 3 system (Applied Biosystems, Life Technologies, USA). The reaction mixture

contained 12.5 µl PowerUp SYBR Green 2X Master Mix (Applied Biosystems), 1 µl each of

forward (341F) and reverse (806R) primer (10 µM), 1 µl of DNA template, 0.1 µg/µl BSA and

brought to a final volume of 25 µl with nuclease free water. Cycling conditions were identical to

those of the endpoint PCR used for sequencing, however with a total of 40 cycles and replacing

the final elongation step with a melt curve. Each sample reaction was performed in triplicate and

these values were averaged for each sample copy calculation. A standard curve was generated

for copy quantification using known concentrations of the expected PCR product amplified from

a similar stool sample. Copies of 16S rRNA per gram of stool was calculated by multiplying the

average copy number per replicate reaction (i.e. 1 μl DNA template) by the total DNA extraction

volume (75 μl) and dividing this value by the mass of stool extracted in grams.

Absolute bacterial abundance by copy number correction

Absolute bacterial abundance was calculated on a per gram of stool basis by correcting the

relative sequencing abundance by the variable number of copies of the 16S rRNA gene in each

observed organism. This correction was done using the "Estimate" tool provided as part of the

rrnDB copy number database[60]. Briefly, after rarefying each sample to an even sequencing depth,

the ASV sequences were submitted through the rrnDB online portal where they were classified

down to the genus level using the RDP classifier version 2.12 and copy number adjusted using

rrnDB copy number data version 5.6[60,61]. The copy number adjusted relative abundance for each

observed taxon was multiplied by the total number of 16S rRNA copies obtained by qPCR,

resulting in the absolute abundance of each taxon per gram of stool.

Fecal inflammatory markers

Inflammatory markers were analyzed using a combination of multiplex technologies using the

Bio-Rad Bio-Plex platform (Bio-Rad, California, USA). The markers evaluated include common

markers of intestinal inflammation including calprotectin and S100A12, in addition to other

markers such as IL-6, TNF, IL-10 and other cytokines and chemokines that may play a role in

inflammatory or anti-inflammatory processes. The data were analyzed using direct comparisons

of all infant groups using analysis of variance (ANOVA) and subsequent individual

comparisons. Fecal calprotectin and S100A12 levels were measured using the fCal ELISA kit

from BUHLMANN Laboratories AG (Schonenbuch, Switzerland) and the Inflamark S100A12

kit from Cisbio Bioassays (Codolet, France), respectively, according to the manufacturer's

instructions. Samples were then analyzed for the presence of both pro-inflammatory and anti-

inflammatory cytokines/chemokines using Multiplex Human Cytokine Magnetic kit, Milliplex

MAP Kit (Millipore, Billerica, MA, USA). Twelve cytokines/chemokines, including EGF, IL-

10, IL-1RA, IL-B, IL-4, IL-6, IL-8, IP-10, MCP-1, MIP-1a, TNFα, and VEGF were analyzed

according to the manufacturer's instructions. Plates were read using the MILLIPLEX Analyzer

3.1 xPONENT™ System (Luminex 200). Cytokine concentrations were determined using

BeadView software (Millipore, Billerica, MA, USA).

Metabolomics

The infant stool samples were suspended in 400 µl 5 mM ammonium acetate. Homogenization

was done 3 times for 30 seconds each time using a cell disruptor. Protein concentrations of the

homogenates were measured using Qubit Protein Assay. Samples were normalized to 500 µg/ml

protein at 25 µl for extraction. Each normalized sample was spiked with 5 µl of internal

standards solution. Extraction of metabolites was performed by protein precipitation by adding

200 µl of extraction solution consisting of 8:1:1 acetonitrile: methanol: acetone to each sample.

Samples were mixed thoroughly, incubated at 4°C to allow protein precipitation, and centrifuged

at 20,000 x g to pellet the protein. 190 µl supernatant was transferred into clean tube and dried

using nitrogen. Samples were reconstituted with 25 µl of reconstitution solution consisting of

injection standards, mixed, and incubated at 4° C for 10-15 min. Samples were centrifuged at

20000 x g. Supernatants were transferred into LC-vials.

Global metabolomics profiling was performed as previously described using a Thermo Q-

Exactive Orbitrap mass spectrometer with Dionex UHPLC and autosampler[62]. Briefly, samples

were analyzed in positive and negative heated electrospray ionization with a mass resolution of

35,000 at m/z 200 as separate injections. Separation was achieved on an ACE 18-pfp 100 x 2.1

mm, 2 µm column with mobile phase A as 0.1% formic acid in water and mobile phase B as

acetonitrile. The flow rate was 350 µl/min with a column temperature of 25°C. 4 µl was injected

for negative ions and 2 µl for positive ions.

Data from positive and negative ions modes were processed separately. LC-MS files were first

converted to open-format files (i.e. mzXML) using MSConvert from Proteowizard[63]. Mzmine

was used to identify features, deisotope, align features and perform gap filling to fill in any

features that may have been missed in the first alignment algorithm[64]. Features were matched

with SECIM internal compound database to identify metabolites. All adducts and complexes

were identified and removed from the data set prior to statistical analysis.

Statistical Analysis

The ASV and taxonomy tables resulting from DADA2 were manipulated using the phyloseq R

package v1.30.0[65]. Inferred metabolic pathway abundances were determined from the rarefied

16S rRNA data using PICRUSt2[29]. Alpha diversity measures, including the observed number of

ASVs and the Shannon diversity index, as well as Spearman correlations and associated

heatmaps, were calculated and drawn using the microbiome R package v1.8.0

(https://bioconductor.org/packages/devel/bioc/html/microbiome.html). Violin and box plots

(including statistical testing where applicable) were generated using the ggpubr R package v0.2.4

(https://github.com/kassambara/ggpubr), which serves as a wrapper for ggplot2[66]. The linear

mixed effects modeling, volatility analysis and associated plots were done using the longitudinal

plugin "q2-longitudinal" offered in Qiime2 v2019.4[26–28]. The biomformat R package

(https://biom-format.org) was used to convert data in phyloseq format to BIOM format for

import into Qiime2[67]. Bray-Curtis and Jaccard distance dissimilarities were calculated using the

vegan R package v2.5.6 (https://github.com/vegandevs/vegan) and PCoA plots were made using

ggplot2 v3.3.0[66]. Individual infant charts were also generated using ggplot2. Non-parametric

statistical tests including the Wilcoxon and Kruskal-Wallis tests were used for pairwise and

overall comparisons of 3 or more factors, respectively[68,69]. The permutational analysis of

variance (PERMANOVA) test was used in the vegan package to compare overall microbiome

dissimilarities between antibiotic use, feeding type, and enrollment groups. P-values were

adjusted for false discovery rate (FDR) via the Benjamin-Hochberg method[70].

Data availability

The demultiplexed 16S rRNA sequencing data generated in this study is deposited in the NCBI

Sequence Read Archive (SRA) under BioProject PRJNA515272.

Trial Registration

This project is registered at clinicaltrials.gov under the name "Antibiotic 'Dysbiosis' in Preterm

Infants" with trial number NCT02784821.

References

1. Israel, E. J. Neonatal necrotizing enterocolitis, a disease of the immature intestinal mucosal

barrier. *Acta. Paediatr. Suppl.* **396**, 27–32 (1994).

2. Sadeghi, K. *et al.* Immaturity of Infection Control in Preterm and Term Newborns Is

Associated with Impaired Toll-Like Receptor Signaling. *J. Infect. Dis.* **195**, 296–302 (2007).

3. Hunter, C. J., Upperman, J. S., Ford, H. R. & Camerini, V. Understanding the Susceptibility of

the Premature Infant to Necrotizing Enterocolitis (NEC). *Pediatr. Res.* **63**, 117–123 (2008).

4. Basso, O. & Wilcox, A. Mortality risk among preterm babies: Immaturity vs. underlying

pathology. *Epidemiology* **21**, 521–527 (2010).

5. Tripathi, N., Cotten, C. M. & Smith, P. B. Antibiotic use and misuse in the neonatal intensive

care unit. *Clin. Perinatol.* **39**, 61–68 (2012).

6. Schulman, J. *et al.* Neonatal intensive care unit antibiotic use. *Pediatrics* **135**, 826–833

(2015).

7. Greenberg, R. G. *et al.* Prolonged duration of early antibiotic therapy in extremely

premature infants. *Pediatr. Res.* **85**, 994–1000 (2019).

8. Puopolo, K. M., Benitz, W. E., Zaoutis, T. E., Newborn, C. on F. A. & Diseases, C. on I.

Management of Neonates Born at ≤34 6/7 Weeks' Gestation With Suspected or Proven

Early-Onset Bacterial Sepsis. *Pediatrics* **142**, 1006 (2018).

9. Fajardo, C., Alshaikh, B. & Harabor, A. Prolonged use of antibiotics after birth is associated

with increased morbidity in preterm infants with negative cultures. *J. Matern.-Fetal*

Neonatal Med. **32**, 4060–4066 (2019).

10. Samuels, N., van de Graaf, R. A., de Jonge, R. C. J., Reiss, I. K. M. & Vermeulen, M. J. Risk

factors for necrotizing enterocolitis in neonates: a systematic review of prognostic studies.

BMC Pediatr. **17**, 105 (2017).

11. Alganabi, M., Lee, C., Bindi, E., Li, B. & Pierro, A. Recent advances in understanding

necrotizing enterocolitis. *F1000Res.* **8**, 107 (2019).

12. Clark, R. H., Bloom, B. T., Spitzer, A. R. & Gerstmann, D. R. Reported Medication Use in the

Neonatal Intensive Care Unit: Data From a Large National Data Set. *Pediatrics* **117**, 1979–

1987 (2006).

13. Hsieh, E. M. *et al.* Medication use in the neonatal intensive care unit. *Am. J. Perinatol.* **31**,

811–821 (2014).

14. Derrien, M., Alvarez, A.-S. & Vos, W. M. de. The Gut Microbiota in the First Decade of Life.

Trends Microbiol. **27**, 997–1010 (2019).

15. Shao, Y. *et al.* Stunted microbiota and opportunistic pathogen colonization in caesarean-

section birth. *Nature* **574**, 117–121 (2019).

16. Stinson, L. F., Boyce, M. C., Payne, M. S. & Keelan, J. A. The Not-so-Sterile Womb: Evidence

That the Human Fetus Is Exposed to Bacteria Prior to Birth. *Front. Microbiol.* **10**, 1124

(2019).

17. Dardas, M. *et al.* The impact of postnatal antibiotics on the preterm intestinal microbiome.

Pediatr. Res. **76**, 150–158 (2014).

18. Gasparrini, A. J. *et al.* Antibiotic perturbation of the preterm infant gut microbiome and

resistome. *Gut Microbes* **7**, 443–449 (2016).

19. Sun, X., Zhuan, C., Xiao, J., Yao, E. & Chen, L. Impact of postnatal exposure to antibiotics on

intestinal microbiome in preterm infants. *Chinese Journal of Perinatal Medicine* **21**, 458–464

(2018).

20. Zou, Z.-H. *et al.* Prenatal and postnatal antibiotic exposure influences the gut microbiota of

preterm infants in neonatal intensive care units. *Ann. Clin. Microbiol. Antimicrob.* **17**, 9

(2018).

21. Cantey, J. B., Pyle, A. K., Wozniak, P. S., Hynan, L. S. & Sánchez, P. J. Early Antibiotic

Exposure and Adverse Outcomes in Preterm, Very Low Birth Weight Infants. *J. Pediatr.* **203**,

62–67 (2018).

22. Langdon, A., Crook, N. & Dantas, G. The effects of antibiotics on the microbiome

throughout development and alternative approaches for therapeutic modulation. *Genome*

Med. **8**, 39 (2016).

23. Sherman, M. P., Zaghouani, H. & Niklas, V. Gut microbiota, the immune system, and diet

influence the neonatal gut–brain axis. *Pediatr. Res.* **77**, 127–135 (2015).

24. Collins, S. M., Surette, M. & Bercik, P. The interplay between the intestinal microbiota and

the brain. *Nat. Rev. Microbiol.* **10**, 735–742 (2012).

25. Ruoss, J. L., *et al.* Routine Early Antibiotic use in SymptOmatic preterm Neonates (REASON):

a prospective randomized controlled trial. *medRxiv* 2020.04.17.20069617 (2020)

doi:10.1101/2020.04.17.20069617

26. Lindstrom, M. J. & Bates, D. M. Newton—Raphson and EM Algorithms for Linear Mixed-

Effects Models for Repeated-Measures Data. *J. Am. Stat. Assoc.* **83**, 1014–1022 (1988).

27. Bokulich, N. A. *et al.* q2-longitudinal: Longitudinal and Paired-Sample Analyses of

Microbiome Data. *mSystems* **3**, e00219-18 (2018).

28. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data science

using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

29. Douglas, G. M. *et al.* PICRUSt2: An improved and customizable approach for metagenome

inference. *bioRxiv* 672295 (2020) doi:10.1101/672295.

30. Tanaka, S. *et al.* Influence of antibiotic exposure in the early postnatal period on the

development of intestinal microbiota. *FEMS Immunol. Med. Mic.* **56**, 80–87 (2009).

31. Greenwood, C. *et al.* Early Empiric Antibiotic Use in Preterm Infants Is Associated with

Lower Bacterial Diversity and Higher Relative Abundance of Enterobacter. *J. Pediatr.* **165**,

23–29 (2014).

32. Cong, X. *et al.* Gut Microbiome Developmental Patterns in Early Life of Preterm Infants:

Impacts of Feeding and Gender. *PLoS One* **11**, e0152751 (2016).

33. Cong, X. *et al.* Influence of Infant Feeding Type on Gut Microbiome Development in

Hospitalized Preterm Infants. *Nurs. Res.* **66**, 123–133 (2017).

34. Parra-Llorca, A. *et al.* Preterm Gut Microbiome Depending on Feeding Type: Significance of

Donor Human Milk. *Front. Microbiol.* **9**, 1376 (2018).

35. Gregory, K. E. *et al.* Influence of maternal breast milk ingestion on acquisition of the

intestinal microbiome in preterm infants. *Microbiome* **4**, 68 (2016).

36. Kordy, K. *et al.* Contributions to human breast milk microbiome and enteromammary

transfer of Bifidobacterium breve. *PLoS ONE* **15**, e0219633 (2020).

37. Pannaraj, P. S. *et al.* Association Between Breast Milk Bacterial Communities and

Establishment and Development of the Infant Gut Microbiome. *JAMA Pediatr.* **171**, 647–

654 (2017).

38. Forbes, J. D. *et al.* Association of Exposure to Formula in the Hospital and Subsequent Infant

Feeding Practices With Gut Microbiota and Risk of Overweight in the First Year of Life.

JAMA Pediatr. **172**, e181161 (2018).

39. Mueller, E. & Blaser, M. Breast milk, formula, the microbiome and overweight. *Nat. Rev.*

Endocrinol. **14**, 510–511 (2018).

40. Zervos, E. E., Bass, S. S., Robson, M. C. & Rosemurgy, A. S. Fluconazole increases bactericidal

activity of neutrophils. *J. Trauma* **41**, 10–14 (1996).

41. Rougé, C. *et al.* Fecal Calprotectin Excretion in Preterm Infants during the Neonatal Period.

PLoS One **5**, e11083 (2010).

42. Coggins, S. A., Wynn, J. L. & Weitkamp, J.-H. Infectious Causes of Necrotizing Enterocolitis.

Clin. Perinatol. **42**, 133–154 (2015).

43. Dobson, S. R. M. & Baker, C. J. Enterococcal Sepsis in Neonates: Features by Age at Onset

and Occurrence of Focal Infection. *Pediatrics* **85**, 165–171 (1990).

44. O'Grady, N. P. *et al.* Detection of Macrophage Inflammatory Protein (MIP)-1α and MIP-β

during Experimental Endotoxemia and Human Sepsis. *J. Infect.* **179**, 136-141 (1999).

45. Bhutta, A. T., Cleves, M. A., Casey, P. H., Cradock, M. M. & Anand, K. J. S. Cognitive and

Behavioral Outcomes of School-Aged Children Who Were Born Preterm: A Meta-analysis.

JAMA **288**, 728–737 (2002).

46. Counsell, S. J. & Boardman, J. P. Differential brain growth in the infant born preterm:

Current knowledge and future developments from brain imaging. *Semin. Fetal. Neonat. M.*

10, 403–410 (2005).

47. Marlow, N. & Samara, M. Neurologic and Developmental Disability at Six Years of Age after

Extremely Preterm Birth. *N. Engl. J. Med.* **352**, 9-19 (2005).

48. Sharon, G. *et al.* Specialized Metabolites from the Microbiome in Health and Disease. *Cell*

Metab. **20**, 719–730 (2014).

49. Kwon, S. H. *et al.* GABA, Resting State Connectivity and the Developing Brain. *Neonatology*

106, 149–155 (2014).

50. Jin, Z., Mendu, S. K. & Birnir, B. GABA is an effective immunomodulatory molecule. *Amino*

Acids **45**, 87–94 (2013).

51. Grill, J. P., Manginot-Dürr, C., Schneider, F. & Ballongue, J. Bifidobacteria and probiotic

effects: Action of Bifidobacterium species on conjugated bile salts. *Curr. Microbiol.* **31**, 23–

27 (1995).

52. Jia, W., Xie, G. & Jia, W. Bile acid–microbiota crosstalk in gastrointestinal inflammation and

carcinogenesis. *Nat. Rev. Gastro. Hepat.* **15**, 111–128 (2018).

53. Kiriyama, Y. & Nochi, H. The Biosynthesis, Signaling, and Neurological Functions of Bile

Acids. *Biomolecules* **9**, 232 (2019).

54. Russell, J. T. *et al.* Genetic risk for autoimmunity is associated with distinct changes in the

human gut microbiome. *Nat. Commun.* **10**, 1–12 (2019).

55. Callahan, B. J. *et al.* DADA2: High resolution sample inference from Illumina amplicon data.

Nat. Methods. **13**, 581–583 (2016).

56. Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and aligned

ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* **35**, 7188–7196

(2007).

57. Glöckner, F. O. *et al.* 25 years of serving the community with ribosomal RNA gene reference

databases and tools. *J. Biotechnol.* **261**, 169–176 (2017).

58. Yilmaz, P. *et al.* The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks.

Nucleic Acids Res. **42**, D643–D648 (2014).

59. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing

and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2013).

60. Stoddard, S. F., Smith, B. J., Hein, R., Roller, B. R. K. & Schmidt, T. M. rrnDB: improved tools

for interpreting rRNA gene abundance in bacteria and archaea and a new foundation for

future development. *Nucleic Acids Res.* **43**, D593–D598 (2015).

61. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*

73, 5261–5267 (2007).

62. Chamberlain, C. A., Hatch, M. & Garrett, T. J. Metabolomic and lipidomic characterization of

Oxalobacter formigenes strains HC1 and OxWR by UHPLC-HRMS. *Anal. Bioanal. Chem.* **411**,

4807–4818 (2019).

63. Chambers, M. C. *et al.* A cross-platform toolkit for mass spectrometry and proteomics. *Nat.*

Biotechnol. **30**, 918–920 (2012).

64. Pluskal, T., Castillo, S., Villar-Briones, A. & Orešič, M. MZmine 2: Modular framework for

processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC*

Bioinform. **11**, 395 (2010).

65. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. *PLOS ONE* **8**, e61217 (2013).

66. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York,

2009).

67. McDonald, D. *et al.* The Biological Observation Matrix (BIOM) Format Or: How I Learned to

Stop Worrying and Love the Ome-Ome. *Gigascience* **1**, 1-7 (2012).

68. Wilcoxon, F. Individual Comparisons by Ranking Methods. *Biometrics Bulletin* **1**, 80–83

(1945).

69. Kruskal, W. H. & Wallis, W. A. Use of Ranks in One-Criterion Variance Analysis. *J. Am. Stat.*

Assoc. **47**, 583–621 (1952).

70. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *J. R. Statist. Soc.* **57**, 289–300 (1995).

Acknowledgements

This study was funded by the NIH under grant R21HD088005. The funder did not have a role in

the design of the study nor how the study was carried out including sample collection, data

analysis and interpretation or manuscript submission. We gratefully acknowledge the University

of Florida Data Safety Monitoring Board; Susmita Datta, PhD - Biostatistician and

Epidemiologist at University of Florida; Michael Cotton, M.D. - Neonatologist at Duke

University School of Medicine; William Benitz, M.D. - Neonatologist at Lucile Packard

Children's Hospital, Stanford; and Robert Lawrence, M.D. - Pediatric Infectious Disease

specialist at University of Florida

Author Information

These authors contributed equally to this work: Jordan T. Russell & J. Lauren Ruoss

Affiliations

Department of Microbiology and Cell Science, Institute of Food and Agricultural Sciences,

University of Florida, Gainesville, FL, USA

Jordan T. Russell, Kelley Lobean McKinley & Eric W. Triplett

Division of Neonatology, Department of Pediatrics, University of Florida, Gainesville, FL, USA

J. Lauren Ruoss, Diomel de la Cruz, Nan Li, Catalina Bazacliu & Josef Neu

Department of Pathology, Immunology and Laboratory Medicine, College of Medicine,

University of Florida, Gainesville, FL, USA

Timothy J. Garrett

Department of Pediatrics, College of Physicians and Surgeons, Columbia University, New York,

NY, USA

Richard A. Polin

Author contributions

J.N. designed and oversaw the study. J.L.R. aided in implementation in the NICU, review of

clinical events for the cohort, preparation for the DSMB and IRB reviews, and analyzed clinical

data. J.N, J.L.R., D.C., and C.B. carried out patient enrollment/consent/group allocation,

provided care for the infants in the NICU during the study. J.L.R, C.B. L.P. recorded clinical

data. N.L. maintained/distributed the samples and performed the stool immune marker assays.

T.J.G performed the stool metabolomics assays and analysis. K.L.M. assisted with stool DNA

extraction, 16S PCR/qPCR, and figure generation. J.T.R. performed the stool microbiome

analysis, data integration, figure generation and wrote the manuscript. R.A.P. and E.W.T.

assisted with data interpretation. All authors reviewed the manuscript before submission.

Corresponding author

Correspondence to Eric W. Triplett

Ethics Declarations

Competing Interests

Dr. Josef Neu is the principal investigator of a study with Infant Bacterial Therapeutics and

on the Scientific Advisory Boards of Medela and Astarte. No other authors have conflicts of

interest to disclose.

Tables and Figures

	Group A	Group B	Group C1	Group C2	Changed from C2
Total	32	11	28	27	13
Changed to antibiotics (bailed)	NA	1	NA	13	NA
Sex (M::F)	14::14	4::7	13::15	17::10	10::3
Delivery mode (V::C)	13::19	8::3	11::17	8::19	11::2
Gestational age range (median)	23 – 32 (28)	29 – 32 (32)	23 – 32 (29)	23 – 32 (29)	24 – 32 (28)
Maternal antibiotic exposure (Y::N)	23::9	6::5	21::7	19::8	12::1
Number of samples post-rarefying	232	42	171	197	97

Table 1 – Summary of infant enrollment and covariates

Summary of the number of enrolled infants per group used in this analysis and the number of
infants changed from groups C2 and B (no antibiotics). Enrollment groups are also summarized
by infant sex (male::female), mode of delivery (vaginal::caesarean), gestational ages and
maternal antibiotic exposure (yes::no). The number of individual stool samples used in the
analysis are also summarized by group.

Figure 1 – Antibiotic use affects early gut richness, bacterial load and trends in diversity

a, Regression scatter plot applying linear mixed effects (LME) modeling to test the relationship

between enrollment group and Shannon diversity longitudinally. (group A n=232, group B n=42,

group C1 n=171, group C2 n=197). Time scale on the x-axis is day after birth (DOL). Greyed

area represents a 95% confidence interval. **b**, **c**, Violin boxplots showing the **(b)** observed

number of ASVs (richness) and **(c)** the bacterial load (log_{10} 16S rRNA copies) of samples which

were taken when antibiotics either were (TRUE, n=83) or were not (FALSE, n=562)

administered (abx_during_sample). Comparisons performed using the Wilcoxon test. **d**,

Volatility plot of the observed number of ASVs by whether the infants in the randomized group

received antibiotics (any_abx, TRUE, n=323 vs. FALSE, n=45). Pairwise comparisons at each

timepoint done using the Wilcoxon test. **e**, **f**, Regression scatter plots applying LME to the **(e)**

observed number of ASVs and **(f)** Shannon diversity longitudinally for infants in group C that

either received (TRUE, n=323) or did not receive (FALSE, n=45) antibiotics (any_abx). Greyed

area represents a 95% confidence interval.

Figure 2 – Beta-diversity shows effects of antibiotics on the infant gut communities

a, **b**, **c**, **d**, **e**, **f**, PCoA ordination of stool samples by **(a, b)** enrollment group, **(c, d)** group C1 or

C2, and **(e, f)** whether they received antibiotics (C1 plus C2 changed: any_abx = TRUE, n=323)

or not (C2 not changed: any_abx = FALSE, n=45) using the quantitative Bray-Curtis distance

index **(a, c, e)** and the qualitative Jaccard distance index **(b, d, f)**. (group A n=232, group B

n=42, group C1 n=171, group C2 n=197). Ellipses are calculated based on a 95% confidence

interval.

Figure 3 – Feeding patterns influence bacterial community richness, diversity, size and structure

a, **b**, **c**, Violin boxplots comparing the **(a)** observed number of ASVs (richness), **(b)** Shannon diversity, and **(c)** log10-transformed number of 16S copies (bacterial load) of samples across different feeding types. Overall comparisons between feeding types was performed using the Kruskal-Wallis test and pairwise comparisons by the Wilcoxon test. Only significant pairwise comparisons are shown for clarity (p < 0.05). **d, e**, PCoA ordination of stool samples collected during different feeding types using the **(d)** Bray-Curtis and **(e)** Jaccard distance indices. Ellipses are calculated based on a 95% confidence interval. DBM: donor breast milk, MBM: mother's breast milk, NPO: no enteral nutrition, ABX: antibiotics. (MBM n=49, DBM n=10, Formula n=27, MBM_Formula n=6, DBM_Formula n=2, NPO n=2).

Figure 4 – Integration of clinical and laboratory data gives detailed view of infant stay in NICU

Extensive clinical and laboratory data, when combined, provide a detailed summary of each infant's stay in the NICU. Data included in each chart from top to bottom include: the infant ID, group assignment, antibiotic change status (bail), gestational age, any adverse clinical events (which are further described in Supplementary Figure 1), the type and duration of antibiotic use (if any), the copy-number corrected absolute composition of each weekly stool sample and it's $\log_{10}$-scale number of bacterial 16S rRNA copies, the type and duration of each feeding including administration of human milk fortifier, the relative levels of C-reactive protein measured from blood, and relative concentrations of measured stool immune markers (for infants where these measurements were performed). DBM: donor breast milk, MBM: mother's breast milk, NPO: no enteral nutrition, CRP: C-reactive protein, EGF: epidermal growth factor.

Figure 5 – Bacterial taxa and adverse clinical event key for infant charts

(a) Key for the color-coded bacterial taxa used in the stool 16S rRNA copy-number corrected

composition pie chart for each infant chart. **(b)** Key for the adverse clinical events (including

infections by body site) and the administration of human milk fortifier for each infant chart.

Figure 6 – Metabolites in stool correlate with abundance of bacterial genera

a, Heatmap of Spearman correlation coefficients between peak response heights of identified

metabolites in stool and the top 10 bacterial genera from the same samples (n=90 stool samples).

Only genera with at least one significant correlation with an identified metabolite are displayed

(9 of 10). Significant correlations are indicated by a '+' with FDR-corrected p-values < 0.05. **b**,

Bacterial genera which had significantly different counts when comparing at least 2 enrollment

groups (at least 1 pairwise comparison). P-values are FDR-adjusted. **c**, Boxplot comparing the

number of rarefied *Veillonella* counts between the enrollment groups. **d**, Original and normalized

concentrations of the neurotransmitter 4-aminobutanoate (GABA) by enrollment group and

significant comparisons of normalized concentrations between groups by ANOVA.

Figure 7 – Abundance of top bacterial genera correlate with abundances of inferred metabolic pathways

730 Heatmap of Spearman correlation values between counts of the most highly abundant bacterial

731 genera and abundance of inferred metabolic pathways using PICRUSt2. Only bacterial genera

732 with at least one significant correlation are depicted (p-values < 0.05). n=90 stool samples.

Figure 8 – Stool immune marker levels show modest correlation with gut microbiota

a, Heatmap of Spearman correlation coefficients between immune markers measured from stool

and the most abundant bacterial genera from the same samples (n=110 stool samples). Only the

bacterial genera with at least one significant correlation with an immune marker are displayed

(n=12). Significant correlations are marked with an '*' by the coefficient, with FDR-adjusted p-

values < 0.05. **b**, Table listing the immune markers used for correlation analysis and their

commonly known general functions. **c**, Comparison of $\log_{10}$-transformed number of

Enterococcus counts by enrollment group and their significance by Kruskal-Wallis. **d**,

Comparison of $\log_{10}$-transformed number of *Enterococcus* counts by whether the infant in

received any antibiotics (any_abx) and their significance by the Wilcoxon test. VEGF: vascular

endothelial growth factor, TNFalpha: tumor necrosis factor alpha, MIP1a: macrophage

inflammatory protein 1a, MCP1: monocyte chemoattractant protein 1, IP10: interferon inducible

protein 10, EGF: epidermal growth factor.