Downregulation of a Type III Secretion System under Reducing Conditions in

Bordetella pertussis

Masataka Goto[a], Tomoko Hanawa[b], Akio Abe[a] and Asaomi Kuwae[a#]

[a]Laboratory of Bacterial Infection, Graduate School of Infection Control Sciences, Kitasato

University, Tokyo, Japan

[b]Department of Infectious Diseases, Kyorin University School of Medicine, Tokyo, Japan

Running Head: Reducing Conditions Downregulate *Bordetella* T3SS.

[#]**Address Correspondence to:** Asaomi Kuwae, kuwae@lisci.kitasato-u.ac.jp

ABSTRACT

Bordetella pertussis uses a type III secretion system (T3SS) to inject virulence proteins into host cells. Although the *B. pertussis* T3SS would play a role in host colonization, the efficient secretion of type III secreted proteins from *B. pertussis* has not been observed. To investigate the roles of type III secreted proteins during infection, we attempted to optimize culture conditions for the production and secretion of a type III secreted protein, BteA, in *B. pertussis*. We observed that *B. pertussis* efficiently secretes BteA in ascorbic acid-depleted (AsA⁻) medium. When L2 cells, a rat lung epithelial cell line, were infected with *B. pertussis* cultured in the AsA⁻ medium, BteA-dependent cytotoxicity was observed. We also performed an immunofluorescence assay of L2 cells infected with *B. pertussis*. The clear fluorescence signals of Bsp22, a needle structure of T3SS, were detected on the bacterial surface of *B. pertussis* cultured in the AsA⁻ medium. Moreover, when *B. pertussis* was cultured in liquid medium containing other reducing agents such as 2-mercaptoethanol and dithioerythritol, the production of type III secreted proteins was repressed. These results suggest that in *B. pertussis*, the production and secretion of type III secreted proteins are downregulated under reducing conditions.

IMPORTANCE

The type III secretion system (T3SS) of *Bordetella pertussis* forms a needle-like structure that protrudes from the bacterial cell surface. *B. pertussis* uses T3SS to translocate virulence proteins called effectors into host cells. The culture conditions for effector production in *B. pertussis* have not been investigated. We attempted to optimize culture medium compositions for producing and secreting type III secreted proteins. We found that *B. pertussis* secretes type III secreted proteins in reducing agent-deprived liquid medium, and that BteA-secreting *B. pertussis* provokes cytotoxicity against cultured mammalian cells. These results suggest that redox signaling is involved in the regulation of *B. pertussis* T3SS.

Introduction

The genus *Bordetella* is made up of Gram-negative bacteria and has been subdivided into

16 subspecies to date (1). *Bordetella pertussis* is a causative agent of whooping cough, also

known as pertussis. *B. pertussis* strictly adapts to humans, and it lacks an environmental

reservoir (2). A variety of virulence factors have been identified in *Bordetella*, e.g.,

filamentous hemagglutinin (3), adenyl cyclase toxin (4) and pertactin (5). These virulence

factors are regulated by a *Bordetella* two-component regulatory system, BvgAS. In addition

to these virulence factors, the type III secretion system (T3SS) of *B. pertussis* is also

regulated by BvgAS (6, 7).

A T3SS that forms a needle-like structure is conserved among many Gram-negative bacteria.

In order to subvert hosts, bacteria use a T3SS to inject type III secreted proteins into host

cells (8). Such type III secreted proteins are called effectors. Three types of effectors have

been identified in *Bordetella*: BteA ((9) [also referred to as BopC (10)], BopN (11, 12), and

BspR (13) [also referred to as BtrA (14, 15)]. BteA is 658 amino acids in length and is

localized to lipid rafts in host cells via its N-terminal region (16). BteA induces necrotic cell

death in several types of cultured mammalian cells (9, 10), and BteA translocation into host

cells is promoted by the functioning of BopN (11). BspR is a negative regulator for type III

secreted proteins (13) and is translocated into nuclei of cultured mammalian cells (17). BspR

has been suggested to be an anti-sigma factor against BtrS that functions as a positive

regulator for type III secreted proteins (15). Some type III secreted proteins are called

translocators rather than effectors. Three types of translocators have been identified in

Bordetella, i.e., BopB (18), BopD (19) and Bsp22 (20), and they function as the path for

effectors delivered into host cells. The complex of BopB and BopD forms the pores on the

host membrane (19), and Bsp22 forms a filamentous-like structure and is associated with

the T3SS needle and with a pore-forming component, BopD (21).

The optimal culture conditions for the production and secretion of BteA have not been

established. To investigate the roles of BteA and other type III secreted proteins during

B. pertussis infection, we investigated culture conditions for the maximal production and

secretion of type III secreted proteins in *B. pertussis*.

MATERIALS AND METHODS

Bacterial strains

The strains used in this study are listed in Table 1. Tohama I (22) was used as the wild-type

strain. BP344, BP348, BP350, and BP351 were used as clinical isolates and were a kind gift

from Dr. K. Kamachi (National Institue of Infectious Diseases) (23). *Escherichia coli* DH10B

and Sm10λ*pir* were used for DNA cloning and conjugation, respectively (Table 1).

B. pertussis was grown on Bordet-Gengou agar plates at 37ºC for 5 days. Fresh colonies of

B. pertussis on each Bordet-Gengou agar plate were suspended in Stainer-Scholt (SS)

medium [Table 2, (24)]. In this study, we used a liquid medium described as a cyclodextrin-

containing medium as the SS medium (25). Low casamino acids (LCA) medium was

prepared by removing the agar from cyclodextrin solid medium (CSM) contents [Table 2,

(26)]. *B. pertussis* was cultured in SS medium with a starting A_{600} of 0.23 under static

conditions at 37ºC. The liquid cultivation period was 22 hr for the protein preparation, mRNA

preparation, and infection.

Generation of a Tohama I nalidixic acid-resistant mutant

For the generation of a Tohama I nalidixic acid-resistant mutant, Tohama I cultured in SS

medium for 24 hr as described above was spread on 20 sets of Bordet-Gengou agar plates

containing nalidixic acid at the final concentration of 30 μg/ml. After 5 days, surviving

colonies were designated as MGKU1. We checked whether the MGKU1 cells showed

hemolysis on a Bordet-Gengou agar plate, and we observed that the amounts of the

production and secretion of type III secreted proteins in MGKU1 were not significantly

different from those of Tohama I wild-type (data not shown).

Cell culture

L2 cells, a rat lung epithelial cell line (ACTT CCL-149), were maintained in F-12K (Invitrogen,

Tokyo). The cell culture medium was supplemented with 10% fetal calf serum (FCS). L2

cells were grown at 37ºC under a 5% CO_2 atmosphere.

Construction of gene-disrupted mutants

The primers used in this study are listed in Table 3. To construct the *bspR* mutant, the 4.0 kb

DNA fragment encoding *bspR* and its flanking regions was amplified by polymerase chain

reaction (PCR) with primers B1-bspR and B2-bspR using *B. pertussis* Tohama I genomic

DNA as the template. The resulting PCR product was cloned into pDONR201 to obtain

pMGKU401, by means of adapter PCR and site-specific recombination techniques using a

Gateway cloning system (Invitrogen). An inverse PCR was carried out with the set of primers

R1-bspR and R2-bspR using circular pMGKU401 as the template.

The resulting PCR product was digested with BamHI and self-ligated to obtain pMGKU402.

This plasmid contained the BamHI site in addition to a 474-bp in-frame deletion from 175 bp

downstream of the 5' end of the *bspR* gene to 52 bp upstream of the 3' end of the gene. This

plasmid, pMGKU402, was mixed with a positive suicide vector, pABB-CRS2 (27), to obtain

pMGKU403 using the Gateway cloning system. This plasmid, pMGKU403, was introduced

into *E. coli* Sm10λ*pir* (18) and transconjugated into MGKU1 as described previously (28).

The resulting mutant strain was designated as MGKU4 (BspR-deficient strain). Similarly,

pABB-Δ*bscN* (18) and pABB-Δ*bopC* (10) were separately introduced into *E. coli* Sm10λ*pir*

and were transconjugated into MGKU1. The resulting mutant strains were designated as

MGKU2 (a T3SS-inactive strain) and MGKU3 (a BteA-deficient strain), respectively.

Preparation of proteins from culture supernatants and whole bacterial cell lysates

Secreted proteins released into bacterial culture supernatants and whole bacterial cell

lysates were prepared by trichloroacetic acid precipitation. The culture supernatants were

filtered, and bacterial pellets were resuspended in distilled water. Trichloroacetic acid was

then added to each sample at the final concentration of 10%. The samples were incubated

on ice for 15 min, then centrifuged for 5 min. The resulting precipitated proteins were

neutralized by 2 M Tris-base and dissolved in the sodium dodecyl sulfate-polyacrylamide gel

electrophoresis (SDS-PAGE) sample buffer. The protein samples were separated by SDS-

PAGE and analyzed by western blotting.

Antibodies

Anti-BteA, anti-BopB, and anti-Bsp22 antibodies were purified from rabbit serum in our

previous study (10, 18, 29). To detect filamentous hemagglutinin (FHA) signals, we used

mouse anti-FHA serum (30). Mouse anti-CyaA monoclonal antibody was purchased from

Santa Cruz Biotechnology (Santa Cruz, CA).

Quantitative reverse transcription-PCR (qRT-PCR)

The amounts of mRNA were measured by qRT-PCR. Bacterial total RNA was prepared

using a Trizol Max Bacterial RNA Isolation Kit (Invitrogen), an RNeasy Mini Kit (Qiagen,

Hilden, Germany) and an RNase-free DNase-free Kit (Qiagen). The RNA sample was

reverse transcribed with Transcriptor Universal cDNA Master (Roche Diagnostics,

Indianapolis, IN) and a T100 Thermal Cycler (Bio-Rad, Hercules, CA). The resulting cDNA

was amplified by FastStart Essential DNA Probes Master (Roche) using the following primer

pairs: 5-recA and 3-recA for *recA*; 5-bteA and 3-bteA for *bteA*; 5-bopB and 3-bopB for *bopB*

(Table 3). The results were calculated as described in the Roche manual. The amount of

recA mRNA was used as an internal control. The mRNA amounts are presented herein as

relative to the amounts in the Tohama I cultured in SS medium, which was defined as 1.

LDH Assays

To examine the release of lactate dehydrogenase (LDH) from *B. pertussis*-infected cells,

5.0×10^4 cells/well of L2 cells seeded in 24-well plates were infected with bacteria at the

multiplicity of infection (MOI) of 125, 250, or 500. The plates were centrifuged at 900 *g* for 5

min and incubated for 3 hr at 37ºC under a 5% CO_2 atmosphere. The amounts of LDH were

measured spectrophotometrically using a Cyto-Tox 96 Non-radioactive Cytotoxicity Assay

Kit (Promega, Madison, WI). The LDH (%) level is shown as a ratio, with the value obtained

from the well treated with 0.1% TritonX-100 set as 100%.

Immunofluorescent staining

For the immunofluorescent staining assay, 2.5×10^5 cells/well of L2 cells were seeded on

coverslips in six-well plates and incubated overnight, and then infected with *B. pertussis* at

an MOI of 125. The plates were centrifuged at 900 *g* for 5 min and incubated for 3 hr at 37ºC

under a 5% CO_2 atmosphere. The infected L2 cells were then immunostained as described

previously (12). Briefly, the cells were treated with 4% paraformaldehyde, 50 mM NH_4Cl,

0.2% Triton-X, and 4% bovine serum albumin (BSA). After blocking, the cells were stained

by anti-Bsp22 antibody (29). As a secondary antibody, Alexa Fluor 488 goat anti-rabbit IgG

(Invitrogen) was used. F-actin was stained with Rhodamine Phalloidin (Molecular Probes,

Eugene, OR). Bsp22 signals per one cell were counted under a fluorescence microscope.

Statistical analyses

The statistical analyses were performed using the nonparametric unpaired *t*-test with a one-

tailed p-value with Prism ver. 5.0f software (Graphpad, La Jolla, CA). Values of $p<0.05$ were

considered significant.

RESULTS

Type III secreted proteins are secreted from *B. pertussis* cultured in LCA medium

Cyclodextrin solid medium (CSM) is reported as a synthetic medium for clinical isolation of

B. pertussis (26). We sought to determine whether *B. pertussis* grown on this agar secretes

type III effectors, and we prepared a liquid medium, designated "low casamino acids (LCA)"

medium, by removing the agar from CSM contents (Table 2). *B. pertussis* was cultured in SS

or LCA medium, and then the whole cell lysates (WCL) and supernatant fractions (Sup)

samples were subjected to western blotting with antibodies against BteA (an effector, a type

III secreted protein) and BopB (a translocator, a type III secreted protein).

BteA and BopB signals were detected in WCL and Sup samples of LCA medium (Fig.

1). As reported previously (10), BteA forms SDS-resistant multimers, and the signals were

also detected around 200 kDa. In contrast, the BteA signal was faintly detected in the WCL

samples of SS medium, and no BteA signal was detected in the Sup samples of SS medium

(Fig. 1). BopB signals were not detected in WCL or Sup samples of SS medium (Fig. 1).

These results suggest that *B. pertussis* secretes type III secreted proteins in LCA medium.

Ascorbic acid downregulates the production and secretion of type III secreted

proteins in *B. pertussis*

Although the ingredients of the LCA and SS media are the same, the amounts of ascorbic

acid and casamino acids in LCA medium are lower than those in SS medium (Table 2). To

determine which substance is involved in the production and secretion of type III secreted

protein, we cultured *B. pertussis* in SS medium, ascorbic acid-deprived SS medium

(SS_AsA⁻), casamino acid-deprived SS medium (SS_CaA⁻), or both ascorbic acid- and

casamino acid-deprived medium (SS_AsA⁻_CaA⁻). WCL and Sup samples were subjected

to western blotting with antibodies against BteA and BopB (Fig. 2).

The BteA multimer signal intensities in the WCL samples of the SS_AsA⁻, SS_CaA⁻ and

SS_AsA⁻_CaA⁻ media were stronger than that in the SS medium (Fig. 2). A faint BteA signal

intensity in the Sup samples of SS was detected (Fig. 2). The BteA signal intensities in the

Sup samples were evident in the SS_AsA⁻ and SS_AsA⁻_CaA⁻ media (Fig. 2). Again, BopB

signal intensities in both WCL and Sup samples were evident in the SS_AsA⁻ and

SS_AsA⁻_CaA⁻ media (Fig. 2).

We also investigated the potential involvement of ascorbic acid or casamino acids in

Bvg-regulated virulence factors, e.g., FHA and CyaA. WCL and Sup samples were subjected

to western blotting with antibodies against FHA and CyaA (Fig. 2). The results revealed that

there were no significant differences in the FHA or CyaA signal intensities in the WCL and

Sup samples among the media used here (Fig. 2). Collectively, these results suggest that

ascorbic acid has a specific influence on the production and secretion of BteA and BopB in

B. pertussis.

Standard SS medium contains ascorbic acid at a final concentration of 2270 μM. To further

explore whether ascorbic acid affects the production and secretion of type III secreted

proteins, we cultured *B. pertussis* in SS medium in the presence of ascorbic acid at the final

concentrations of 91, 454, or 2270 μM. WCL and Sup samples were prepared from each

bacterial culture and analyzed by western blotting (Fig. 3). The BteA signal was detected in

both WCL and Sup samples of SS medium at 0–454 μM ascorbic acid, but the BteA signal of

the SS medium at 2270 μM ascorbic acid was faint or absent (Fig. 3). The BopB signals

showed a similar pattern (Fig. 3). These results indicate that *B. pertussis* produces and

secretes type III secreted proteins under low ascorbic acid concentrations.

BteA and BopB are secreted from other *B. pertussis* clinical isolates cultured in

SS_AsA⁻ medium

To determine whether *B. pertussis* clinical isolates secrete type III secreted proteins under low ascorbic acid concentrations, we cultured four strains of *B. pertussis* (Table 1) in SS medium or ascorbic acid-deprived medium (SS_AsA⁻). The prepared WCL and Sup samples were analyzed by western blotting (Fig. 4). BteA signals in the Sup samples of BP350 and BP351 were evident in the SS_AsA⁻ medium (Fig. 4, AsA⁻). Again, BopB signals in the Sup samples of BP350 and BP351 were evident in SS_AsA⁻ medium. These results suggest that some clinical strains, e.g., BP350 and BP351, also secrete type III secreted proteins under low ascorbic acid concentrations.

Translocator BopB is upregulated at the transcriptional level in *B. pertussis* cultured

in SS_AsA⁻ medium

To investigate the gene expressions of type III secreted proteins under ascorbic acid-starved conditions, we prepared total RNA from *B. pertussis* cultured in SS or SS_AsA⁻ medium. The cDNA samples reverse-transcribed from the total RNA samples were subjected to a quantitative RT-PCR analysis to quantify the relative amounts of *bteA* and *bopB* mRNA. The results demonstrated that the relative amount of *bteA* mRNA of Tohama I in SS_AsA⁻ medium was increased compared to that in SS medium (Fig. 5A). However, the relative amount of *bteA* mRNA of BP350 in the SS_AsA⁻ medium was not significantly different from that in the SS medium (Fig. 5A). The relative amounts of *bopB* mRNA of Tohama I and BP350 in SS_AsA⁻ medium were increased compared to those in SS medium (Fig. 5B).

These results suggest that BopB is upregulated at the transcriptional level in Tohama I and BP350 under low ascorbic acid concentrations. BteA is also upregulated at the transcriptional level in Tohama I under low ascorbic acid concentrations but not in BP350.

B. pertussis **induces cytotoxicity against cultured mammalian cells**

A *B. pertussis* BspR-deficient strain was reported to induce cytotoxicity against cultured

mammalian cells (15). In the present study, we identified better culture conditions for the

production and secretion of BteA in *B. pertussis* than those provided with standard SS

medium. We thus investigated whether wild-type *B. pertussis* provokes cytotoxicity against

cultured mammalian cells under these conditions. L2 cells, a rat lung epithelial cell line, were

infected in SS medium comprised of one-twentieth of ascorbic acid (SS_1/20AsA), SS

medium comprised of one-twentieth of ascorbic acid and one-half of casamino acids

(SS_1/20AsA_1/2CaA), or SS medium comprised of one-twentieth of ascorbic acid and no

casamino acids (SS_1/20AsA_CaA$^-$) at an MOI of 500 for 3 hr. The amounts of lactate

dehydrogenase (LDH) released into the extracellular medium were then measured.

The results revealed that released LDH was detected from L2 cells infected with Tohama I in

the SS_1/20AsA or SS_1/20AsA_1/2CaA medium (Fig. 6). The L2 cells incubated in the

SS_AsA$^-$ or SS_1/20AsA_CaA$^-$ medium were injured by the medium itself. Therefore, these

media were not useful for this assay. Taken together, these results demonstrate that we

successfully observed cytotoxicity of mammalian cells induced by wild-type *B. pertussis*.

The T3SS of *B. pertussis* cultured in SS_AsA$^-$ medium is active during infection

against cultured mammalian cells

As described above, we observed cytotoxicity of *B. pertussis* against cultured cells. In order

to confirm that *B. pertussis* indeed secretes type III secreted proteins during infection, we

infected L2 cells with *B. pertussis* in SS or SS_1/20AsA_1/2CaA medium at an MOI of 125

for 3 hr. After infection, Bsp22 (a translocator, and a component of the filamentous-like

structure) and F-actin were stained with anti-Bsp22 and rhodamine phalloidin, respectively.

The results showed that the amounts of Bsp22 signals (Fig. 7A, green) on L2 cells infected

with Tohama I and BP350 cultured in SS_1/20AsA_1/2CaA medium were greater than those

in SS medium (Fig. 7). Thus, we demonstrated that *B. pertussis* T3SS is activated during the

infection of cultured mammalian cells under appropriate conditions.

The cytotoxicity induced by wild-type *B. pertussis* depends on the T3SS

It has been reported that *B. bronchiseptica* induces BteA-dependent necrotic cell death

against cultured mammalian cells (10). To investigate whether the cytotoxicity provoked by

wild-type *B. pertussis* also depends on BteA function, we infected L2 cells with MGKU1 (a

Tohama I nalidixic acid-resistant strain), MGKU2 (a T3SS-inactive strain), MGKU3 (a BteA-

deficient strain), or MGKU4 (a BspR-deficient strain that allows the constitutive activation of

T3SS) in SS_1/20AsA_1/2CaA medium at an MOI of 125 or 250 for 3 hr. The amounts of

LDH released into extracellular medium were measured.

As a result, LDH was not detected in the medium of L2 cells infected with MGKU2 (Fig.

8A,B). The amount of LDH detected in the medium of L2 cells infected with MGKU3 was

significantly lower than that in the medium of L2 cells infected with MGKU1 (Fig. 8A,B). The

amount of LDH detected in the medium of L2 cells infected with MGKU4 was significantly

greater than that in the medium of L2 cells infected with MGKU1 at an MOI of 125 (Fig. 8A),

but not at an MOI of 250 (Fig. 8B).

WCL and Sup samples were prepared from each strain and separated by SDS-PAGE, then

analyzed by western blotting with anti-BteA antibody (Fig. 8C). BteA signals of the WCL

samples were detected in MGKU1, MGKU2, and MGKU4, but not in MGKU3 (Fig. 8C). BteA

signals of the Sup samples were detected in MGKU1 and MGKU4, but not in MGKU2 or

MGKU3 (Fig. 8C). These results suggest that the cytotoxicity induced by wild-type

B. pertussis (Fig. 6) is dependent on T3SS activity and is partially dependent on BteA.

The secretion of type III secreted proteins in *B. pertussis* is downregulated under

reducing conditions

Ascorbic acid is known as a reducing agent. To determine whether the secretion of type III

secreted proteins is downregulated under reducing conditions, we cultured Tohama I in

ascorbic acid-deprived SS medium (SS_AsA⁻, Fig. 9, "none") and in SS_AsA⁻ medium

containing the reducing agents ascorbic acid (AsA), 2-mercaptoethanol (2-ME) or

dithioerythritol (DTE), respectively. The prepared WCL and Sup samples were analyzed by

western blotting with anti-BteA, anti-BopB, or anti-FHA antibodies (Fig. 9). The BteA and

BopB signal intensities of both WCL and Sup samples were decreased in the presence of

each of the reducing agents (Fig. 9). In contrast, the FHA signal intensities of both WCL and

Sup samples were not affected by the reducing agents. The growth of bacteria cultured in

SS medium with reducing agents was not significantly different from that of the bacteria

cultured in SS medium (data not shown). These results suggest that the production and

secretion of type III secreted proteins in *B. pertussis* are downregulated under reducing

conditions.

DISCUSSION

In this study, *B. pertussis* increased the production and secretion of type III secreted proteins,

BteA and BopB, when cultured in ascorbic acid-deprived SS medium (SS_AsA⁻). We

successfully produced cytotoxicity against mammalian cells by the infection of wild-type

Tohama I cultured in SS 1/20AsA_1/2CaA medium. *B. pertussis* decreased the production

and secretion of type III secreted proteins when cultured in SS_AsA⁻ medium containing

reducing agents such as 2-ME or DTT. These results suggest that redox signaling is

involved in regulation of the T3SS in *B. pertussis*.

It has been reported that no secretion of the type III secreted proteins BteA and Bsp22

was detected from Tohama I (31), and that the secretion of the type III secreted proteins

BteA and BopD was detected from a non-vaccine-type strain, BP157. However, BP157

showed no cytotoxicity against L2 cells, J774 mouse macrophage-like cells, and HeLa cells

(32). In another report, glutamate limitation upregulates production of the type III secreted

proteins (33). However, the requirement of a type III secretion system (T3SS) for virulence in

B. pertussis thus remains unclear. On the other hand, a BscN-deficient mutant in which the

T3SS was inactive colonized mouse lungs to a significantly lower degree compared to the

wild-type (31). In addition, cytotoxicity against HeLa cells was observed by infection with a

BspR (a negative regulator for type III secreted proteins)-deficient *B. pertussis* strain (15).

These results suggest that the T3SS has a significant role in the virulence of *B. pertussis*.

The production and the relative amounts of *bopB* mRNA of Tohama I cultured in SS

AsA⁻ medium were significantly increased compared to those of Tohama cultured in SS

medium, but this was not the case for BteA mRNA (Figs. 2, 5). It was reported that *bteA* and

bopB genes are upregulated by BtrS (a sigma factor for type III secreted proteins) and

downregulated by BspR (13, 15). It has also been reported that the amount of *bteA* mRNA

was increased 2-fold in a BspR-deficient strain compared to the wild-type, whereas the

amount of *bopB* mRNA was increased by 18-fold compared to the wild-type by RNA-seq

(15). The *bopB* gene is localized in the T3SS apparatus locus (*bsc* locus) adjacent to the

*btr*S gene. Since the *bteA* gene is separated from the *bsc* locus by >2.5 Mb (9), BtrS would

preferentially regulate transcription of the *bopB* gene over that of the *bteA* gene.

The cytotoxicity against L2 cells was observed by the infection of Tohama I cultured in

SS_1/20AsA _1/2CaA medium (Fig. 6). Before infection, we exchanged Ham's F-12K

medium for the SS_1/20AsA _1/2CaA medium. When L2 cells were infected with Tohama I

in Ham's F-12K medium, the cytotoxicity against L2 cells was not observed (data not shown).

We therefore speculate that a substance which inhibits the secretion of type III secreted

proteins from Tohama I was present in the Ham's F-12K medium, or a substance that is

necessary for the secretion of type III secreted proteins from Tohama I was absent in the

Ham's F-12K medium. We found that infection of the human alveolar epithelium cell line

A549 with wild-type *B. pertussis* cultured in SS_1/20AsA _1/2CaA medium did not result in

cytotoxicity (data not shown). For the observation of cytotoxicity against A549 cells, it is

necessary to optimize the conditions for the production and secretion of type III secreted

proteins in *B. pertussis*.

A previous study showed that cytotoxicity against the cell line J774A.1 was achieved

by infection with wild-type *B. pertussis* 18323, but not by infection with a CyaA-deficient

strain (34). This finding suggested that CyaA has the potential to provoke the cytotoxicity.

However, we confirmed herein that the production and secretion of CyaA in a BteA-deficient

strain (MGKU3) were not significantly different from those of the wild-type (data not shown).

In our results, the amounts of LDH released from L2 cells infected with MGKU3 were greater

than those of the T3SS-inactive strain (MGKU2), and no LDH release was detected from L2

cells infected with MGKU2 (Fig. 8). These results raised the possibility that an unknown type

III effector other than BteA is involved in the cytotoxicity of *B. pertussis*.

Although we observed no significant difference in the growth rate of Tohama I

between SS medium and SS_AsA⁻ medium (data not shown), the production and secretion

of type III secreted proteins of Tohama I cultured in the SS_AsA⁻ medium were increased

compared to those in the SS medium (Fig. 2). It had not been determined whether the

expression of type III secreted proteins is regulated by reducing agents at the molecular

level, and in this study, the production and secretion of CyaA and FHA in Tohama I cultured

in SS medium were not significantly different from those in SS_AsA⁻ medium (Fig. 2). It was

reported that there is no difference in Prn production by Tohama I cultured in modified SS

medium compared to culture in modified SS medium containing dithiothreitol (DTT), a

reducing agent (35). These results suggest that BvgAS-regulated virulence factors other

than type III secreted proteins are not affected by reducing agents.

Ascorbic acid, urea, and glutathione (GSH) function as reducing agents in the

epithelial lining fluid (ELF) of the human airway. The concentrations of ascorbic acid and

urea in ELF in the upper respiratory tract are not significantly different from those in the

lower respiratory tract. However, the concentration of GSH in the ELF in the upper

respiratory tract is 100-fold lower than that in the lower respiratory tract. One of the

possibilities is that GSH forms GS-SG (an oxidized from of GSH) in the ELF in the upper

respiratory tract (36). It may thus be considered that the upper respiratory tract is more

oxidized than the lower respiratory tract. It was reported that the transcription of genes

encoding type III secreted proteins of *B. pertussis* recovered from mouse nasal lavage was

higher than that from bronchoalveolar lavage (37). We thus speculate that the T3SS is

important for *B. pertussis* to colonize the upper respiratory tract of the host.

In this study, the production and secretion of BopB by Tohama I cultured in the

SS_AsA⁻ medium were significantly increased compared to those in SS medium (Fig. 2).

Bacteria use redox-sensing proteins to resist oxidative stress (38). For example, OxyR (a

member of the LysR family of transcriptional regulators) of *Escherichia coli* is converted to

an active mode by the oxidation of intramolecular cysteine residues, which then increases

the expression of antioxidants genes, e.g., hydroperoxidase I [*katG*, (39)]. Vfr (a member of

the CRP/FNR superfamily) of *Pseudomonas* forms an active mode by retaining

intramolecular cysteine residues as reducing states, and then increases the expression of

type III secreted proteins (40). Btr is a regulator protein produced by *B. pertussis* and is

homologous to Vfr (41). A cysteine cluster is located in the N-terminal domain of Btr, and a

helix-turn-helix DNA binding domain is located in the C-terminal domain of Btr. Therefore,

when *B. pertussis* is cultured in reducing agent-deprived SS medium, Btr might be oxidized

and change its conformation, thereby increasing the expression of type III secreted proteins.

Although the signal pathway that *B. pertussis* uses to increase the production of type III

secreted proteins is unclear, it is possible that Btr regulates gene expressions for type III

secreted proteins.

REFERENCES

1.   Ivanov YV, Linz B, Register KB, Newman JD, Taylor DL, Boschert KR, Le Guyon S,

Wilson EF, Brinkac LM, Sanka R, Greco SC, Klender PM, Losada L, Harvill ET. 2016.

Identification and taxonomic characterization of *Bordetella pseudohinzii* sp. nov.

isolated from laboratory-raised mice. Int J Syst Evol Microbiol 66:5452-5459.

2.   Mattoo S, Cherry JD. 2005. Molecular pathogenesis, epidemiology, and clinical

manifestations of respiratory infections due to *Bordetella pertussis* and other

Bordetella subspecies. Clin Microbiol Rev 18:326-82.

3.   Relman DA, Domenighini M, Tuomanen E, Rappuoli R, Falkow S. 1989. Filamentous

hemagglutinin of *Bordetella pertussis*: nucleotide sequence and crucial role in

adherence. Proc Natl Acad Sci U S A 86:2637-41.

4.   Hewlett EL, Gordon VM, McCaffery JD, Sutherland WM, Gray MC. 1989. Adenylate

cyclase toxin from *Bordetella pertussis*. Identification and purification of the holotoxin

molecule. J Biol Chem 264:19379-84.

5.   Roberts M, Fairweather NF, Leininger E, Pickard D, Hewlett EL, Robinson A,

Hayward C, Dougan G, Charles IG. 1991. Construction and characterization of

Bordetella pertussis mutants lacking the vir-regulated P.69 outer membrane protein.

Mol Microbiol 5:1393-404.

6.   Smith AM, Guzman CA, Walker MJ. 2001. The virulence factors of *Bordetella*

pertussis: a matter of control. FEMS Microbiol Rev 25:309-33.

7.   Yuk MH, Harvill ET, Miller JF. 1998. The BvgAS virulence control system regulates

type III secretion in *Bordetella bronchiseptica*. Mol Microbiol 28:945-59.

8.   Cornelis GR, Van Gijsegem F. 2000. Assembly and function of type III secretory

systems. Annu Rev Microbiol 54:735-74.

9.   Panina EM, Mattoo S, Griffith N, Kozak NA, Yuk MH, Miller JF. 2005. A genome-wide

screen identifies a *Bordetella* type III secretion effector and candidate effectors in

other species. Mol Microbiol 58:267-79.

10.   Kuwae A, Matsuzawa T, Ishikawa N, Abe H, Nonaka T, Fukuda H, Imajoh-Ohmi S,

Abe A. 2006. BopC is a novel type III effector secreted by *Bordetella bronchiseptica*

and has a critical role in type III-dependent necrotic cell death. J Biol Chem

281:6589-600.

11.   Abe A, Nishimura R, Kuwae A. 2017. *Bordetella* effector BopN is translocated into

host cells via its N-terminal residues. Microbiol Immunol 61:206-214.

12.   Nagamatsu K, Kuwae A, Konaka T, Nagai S, Yoshida S, Eguchi M, Watanabe M,

Mimuro H, Koyasu S, Abe A. 2009. *Bordetella* evades the host immune system by

inducing IL-10 through a type III effector, BopN. J Exp Med 206:3073-88.

13.   Kurushima J, Kuwae A, Abe A. 2012. The type III secreted protein BspR regulates

the virulence genes in *Bordetella bronchiseptica*. PLoS One 7:e38925.

14.   Ahuja U, Liu M, Tomida S, Park J, Souda P, Whitelegge J, Li H, Harvill ET, Parkhill J,

Miller JF. 2012. Phenotypic and genomic analysis of hypervirulent human-associated

Bordetella bronchiseptica. BMC Microbiol 12:167.

15.   Ahuja U, Shokeen B, Cheng N, Cho Y, Blum C, Coppola G, Miller JF. 2016.

Differential regulation of type III secretion and virulence genes in *Bordetella pertussis*

and *Bordetella bronchiseptica* by a secreted anti-sigma factor. Proc Natl Acad Sci U

S A 113:2341-8.

16.   French CT, Panina EM, Yeh SH, Griffith N, Arambula DG, Miller JF. 2009. The

Bordetella type III secretion system effector BteA contains a conserved N-terminal

motif that guides bacterial virulence factors to lipid rafts. Cell Microbiol 11:1735-49.

17.   Abe A, Nishimura R, Tanaka N, Kurushima J, Kuwae A. 2015. The *Bordetella*

Secreted Regulator BspR Is Translocated into the Nucleus of Host Cells via Its N-

459   Terminal Moiety: Evaluation of Bacterial Effector Translocation by the *Escherichia*

460   *coli* Type III Secretion System. PLoS One 10:e0135140.

461 18. Kuwae A, Ohishi M, Watanabe M, Nagai M, Abe A. 2003. BopB is a type III secreted

462   protein in *Bordetella bronchiseptica* and is required for cytotoxicity against cultured

463   mammalian cells. Cell Microbiol 5:973-83.

464 19. Nogawa H, Kuwae A, Matsuzawa T, Abe A. 2004. The type III secreted protein BopD

465   in *Bordetella bronchiseptica* is complexed with BopB for pore formation on the host

466   plasma membrane. J Bacteriol 186:3806-13.

467 20. Yuk MH, Harvill ET, Cotter PA, Miller JF. 2000. Modulation of host immune

468   responses, induction of apoptosis and inhibition of NF-kappaB activation by the

469   *Bordetella* type III secretion system. Mol Microbiol 35:991-1004.

470 21. Medhekar B, Shrivastava R, Mattoo S, Gingery M, Miller JF. 2009. *Bordetella* Bsp22

471   forms a filamentous type III secretion system tip complex and is immunoprotective in

472   vitro and in vivo. Mol Microbiol 71:492-504.

473 22. Kasuga T, Nakase Y, Ukishima K, Takatsu K. 1954. Studies on *Haemophilus*

474   *pertussis*. V. Relation between the phase of bacilli and the progress of the whooping-

475   cough. Kitasato Arch Exp Med 27:57-62.

476 23. Otsuka N, Han HJ, Toyoizumi-Ajisaka H, Nakamura Y, Arakawa Y, Shibayama K,

477   Kamachi K. 2012. Prevalence and genetic characterization of pertactin-deficient

478   *Bordetella pertussis* in Japan. PLoS One 7:e31985.

479 24. Stainer DW, Scholte MJ. 1970. A simple chemically defined medium for the

480   production of phase I *Bordetella pertussis*. J Gen Microbiol 63:211-20.

481 25. Imaizumi A, Suzuki Y, Ono S, Sato H, Sato Y. 1983. Effect of heptakis (2,6-O-

482   dimethyl) beta-cyclodextrin on the production of pertussis toxin by *Bordetella*

483   *pertussis*. Infect Immun 41:1138-43.

26. Aoyama T, Murase Y, Iwata T, Imaizumi A, Suzuki Y, Sato Y. 1986. Comparison of

blood-free medium (cyclodextrin solid medium) with Bordet-Gengou medium for

clinical isolation of *Bordetella pertussis*. J Clin Microbiol 23:1046-8.

27. Sekiya K, Ohishi M, Ogino T, Tamano K, Sasakawa C, Abe A. 2001. Supermolecular

structure of the enteropathogenic *Escherichia coli* type III secretion system and its

direct interaction with the EspA-sheath-like structure. Proc Natl Acad Sci U S A

98:11638-43.

28. Donnenberg MS, Kaper JB. 1991. Construction of an eae deletion mutant of

enteropathogenic *Escherichia coli* by using a positive-selection suicide vector. Infect

Immun 59:4310-7.

29. Kurushima J, Kuwae A, Abe A. 2012. Btc22 chaperone is required for secretion and

stability of the type III secreted protein Bsp22 in *Bordetella bronchiseptica*. FEMS

Microbiol Lett 331:144-51.

30. Watanabe M, Nagai M, Funaishi K, Endoh M. 2000. Efficacy of chemically cross-

linked antigens for acellular pertussis vaccine. Vaccine 19:1199-203.

31. Fennelly NK, Sisti F, Higgins SC, Ross PJ, van der Heide H, Mooi FR, Boyd A, Mills

KHG. 2008. *Bordetella pertussis* expresses a functional type III secretion system that

subverts protective innate and adaptive immune responses. Infection and Immunity

76:1257-1266.

32. Han HJ, Kuwae A, Abe A, Arakawa Y, Kamachi K. 2011. Differential expression of

type III effector BteA protein due to IS481 insertion in *Bordetella pertussis*. PLoS One

6:e17797.

33. Hanawa T, Kamachi K, Yonezawa H, Fukutomi T, Kawakami H, Kamiya S. 2016.

Glutamate Limitation, BvgAS Activation, and (p)ppGpp Regulate the Expression of

the *Bordetella pertussis* Type 3 Secretion System. J Bacteriol 198:343-51.

34. Khelef N, Zychlinsky A, Guiso N. 1993. *Bordetella pertussis* induces apoptosis in

macrophages: role of adenylate cyclase-hemolysin. Infect Immun 61:4064-71.

35. Stenson TH, Patton AK, Weiss AA. 2003. Reduced glutathione is required for

pertussis toxin secretion by *Bordetella pertussis*. Infect Immun 71:1316-20.

36. van der Vliet A, O'Neill CA, Cross CE, Koostra JM, Volz WG, Halliwell B, Louie S.

1999. Determination of low-molecular-mass antioxidant concentrations in human

respiratory tract lining fluids. Am J Physiol 276:L289-96.

37. van Beek LF, de Gouw D, Eleveld MJ, Bootsma HJ, de Jonge MI, Mooi FR, Zomer A,

Diavatopoulos DA. 2018. Adaptation of *Bordetella pertussis* to the Respiratory Tract.

J Infect Dis 217:1987-1996.

38. Lee SJ, Kim DG, Lee KY, Koo JS, Lee BJ. 2018. Regulatory mechanisms of thiol-

based redox sensors: lessons learned from structural studies on prokaryotic redox

sensors. Arch Pharm Res 41:583-593.

39. Aslund F, Zheng M, Beckwith J, Storz G. 1999. Regulation of the OxyR transcription

factor by hydrogen peroxide and the cellular thiol-disulfide status. Proc Natl Acad Sci

U S A 96:6161-5.

40. Zhang Y, Zhang C, Du X, Zhou Y, Kong W, Lau GW, Chen G, Kohli GS, Yang L,

Wang T, Liang H. 2019. Glutathione Activates Type III Secretion System Through Vfr

in *Pseudomonas aeruginosa*. Front Cell Infect Microbiol 9:164.

41. Bannan JD, Moran MJ, MacInnes JI, Soltes GA, Friedman RL. 1993. Cloning and

characterization of btr, a *Bordetella pertussis* gene encoding an FNR-like

transcriptional regulator. J Bacteriol 175:7228-35.

ACKNOWLEDGMENTS

This work was supported in part by grants from the Ministry of Education, Culture, Sports,

Sciences and Technology and the Japan Society for the Promotion of Science (KAKENHI),

nos. 19K07542 (to A.A.), 17K08838 (A.K.), and 19K07561 (T.H.), and by a research grant

from the Takeda Science Foundation in 2016 (A.K.). The funders had no role in the study

design, data collection or analysis, the decision to publish, or the preparation of the

manuscript.

Figure legends

Fig. 1. BteA and BopB production by *B. pertussis* cultured in LCA medium. Tohama I was

cultured in SS or LCA medium. Whole cell lysates (WCL) and supernatant fractions (Sup)

samples were separated by SDS-PAGE and analyzed by western blotting with antibodies

against BteA (*upper panel*) or BopB (*lower panel*). BteA forms SDS-resistant multimers (10).

NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal

volume of bacterial culture. Experiments were performed at least three times, and

representative data are shown.

Fig. 2. Effects of ascorbic acid and casamino acids on Bvg-regulated virulence factors.

Tohama I was cultured in SS medium or ascorbic acid-deprived SS medium (SS_AsA$^-$),

casamino acid-deprived SS medium (SS_CaA$^-$) or both ascorbic acid- and casamino acid-

deprived SS medium (SS AsA$^-$_CaA$^-$). WCL and Sup samples were separated by SDS-

PAGE and analyzed by western blotting with antibodies against BteA, BopB, FHA, or CyaA.

NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal

volume of bacterial culture. Experiments were performed at least three times, and

representative data are shown.

Fig. 3. Effects of ascorbic acid on type III secreted proteins. Whole cell lysates and secreted

proteins prepared from Tohama I cultured in SS medium in the presence of ascorbic acid at

the final concentration of 91, 454 or 2270 μM were separated by SDS-PAGE and then

analyzed by western blotting with antibodies against BteA or BopB. NS: nonspecific signals.

Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture.

Experiments were performed at least three times, and representative data are shown.

Fig. 4. Effects of ascorbic acid on the secretion of BteA and BopB in clinical isolates cultured

in SS_AsA⁻ medium. The *B. pertussis* clinical isolates BP344, BP348, BP350, and BP351

were cultured in SS (+) or SS_AsA⁻ (–) medium. WCL and Sup samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA (*upper panel*) or

BopB (*lower panel*). NS: nonspecific signals. Loaded WCL and Sup samples were prepared

from an equal volume of bacterial culture. Experiments were performed at least three times,

and representative data are shown.

Fig. 5. Results of the RT-PCR analysis for mRNA levels of *bteA* and *bopB* from *B. pertussis*

cultured under ascorbic acid-starved conditions. Total RNA was prepared from Tohama I or

BP350 cultured in SS or SS_AsA⁻ medium and subjected to a quantitative RT-PCR analysis.

The histograms show the relative amounts of *bteA* (**A**) and *bopB* (**B**) mRNA normalized by

the housekeeping gene, *recA* mRNA. Error bars: the standard error of the mean (SEM) from

triplicate experiments. *$p<0.05$; N.S.: no significant difference. Experiments were performed

at least three times, and representative data are shown.

Fig. 6. Cytotoxic activity of *B. pertussis* cultured in SS_AsA⁻ medium. L2 cells were infected

with Tohama I cultured in SS medium, SS_AsA⁻ medium, SS medium containing one-

twentieth of ascorbic acid (SS_1/20AsA), SS medium containing one-twentieth of ascorbic

acid and one-half of casamino acids (SS_1/20AsA_1/2CaA), or SS medium containing one-

twentieth of ascorbic acid and no casamino acids (SS_1/20AsA_CaA⁻) at an MOI of 500 for

3 hr. +, 1/2 and 1/20 show the amounts of indicated substances that are equal to, one-half of,

and one-twentieth of those in SS medium, respectively. The amounts of LDH released into

the extracellular medium from infected cells are shown, and the relative cytotoxicity (%) was

determined as described in the Materials and Methods section. Error bars: the SEM from

triplicate experiments. n.a.: No available result. Experiments were performed at least three

times, and representative data are shown.

Fig. 7. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. pertussis.* **A:** L2

cells were infected with Tohama I or BP350 in the indicated media at an MOI of 125 for 3 hr.

+, 1/2 and 1/20 indicate the amounts of indicated substances that are equal to, one-half of,

and one-twentieth of those in SS medium, respectively. After fixation, cells were stained with

anti-Bsp22 antibody (*green*) and rhodamine phalloidin (*red*). *Arrowheads* indicate the Bsp22

(*green*) signal. **B:** Bsp22 signals per one cell were counted under a fluorescent microscope.

At least 120 cells were randomly chosen. Error bars: the SEM from triplicate experiments.

*p<0.05; N.S.: no significant difference. Experiments were performed at least three times,

and representative data are shown.

Fig. 8. Cytotoxic activities of *B. pertussis* mutants. The results of LDH assays at an MOI of

125 (**A**) or 250 (**B**) are shown as a histogram. Error bars: the SEM from triplicate

experiments. *p<0.05. Experiments were performed at least three times, and representative

data are shown. **C:** MGKU1 (Tohama I nalidixic acid-resistant strain, 1: WT Nalʳ), MGKU2

(T3SS-inactive strain, 2: ΔT3SS), MGKU3 (BteA-deficient strain, 3: ΔBteA), or MGKU4

(BspR-deficient strain, 4: ΔBspR) was cultured in SS_1/20AsA_1/2CaA medium. WCL and

Sup samples were separated by SDS-PAGE and analyzed by western blotting with anti-BteA

antibody. We loaded a 3-fold smaller amount of MGKU4 supernatant fraction on the SDS-

PAGE gel to avoid obtaining an excess signal intensity. NS: nonspecific signals. Loaded

WCL and Sup samples were prepared from an equal volume of bacterial culture.

Experiments were performed at least three times, and representative data are shown.

Fig. 9. Effects of reducing agents on the production and secretion of type III secreted

proteins in *B. pertussis*. Tohama I was cultured in SS_AsA⁻ medium or SS_AsA⁻ medium

containing ascorbic acid (AsA), 2-mercaptoethanol (2-ME), or dithioerythritol (DTE) at final

concentrations of 2.3, 2.1, or 3.0 mM, respectively. WCL and Sup samples were separated

by SDS-PAGE, then analyzed by western-blotting with antibodies against BteA, BopB, or

FHA. Loaded WCL and Sup samples were prepared from an equal volume of bacterial

culture. NS: nonspecific signals. Experiments were performed at least three times, and

representative data are shown.

Table 1. Bacterial strains and plasmids used in study

Strains or plasmids	Relevant genotype/description	Reference
Strains		
B. pertussis		
Tohama I	Clinical isolate	22
BP344	Clinical isolate	23
BP348	Clinical isolate	23
BP350	Clinical isolate	23
BP351	Clinical isolate	23
MGKU1	Tohama I nalidixic-acid resistant strain	This study
MGKU2	T3SS-inactive strain	This study
MGKU3	BteA-deficient strain	This study
MGKU4	BspR-deficient strain	This study
E. coli		
DH10B	Host strain for pDONR201	invitrogen
Sm10λ*pir*	Host strain for pABB-CRS2	18
Plasmids		
pDONR201	DNA cloning vector, Km^r	invitrogen
pABB-CRS2	Suicide vector for conjugation, Amp^r	27
pABB-Δ*bscN*	pABB-CRS2, Δ*bscN* gene	18
pABB-Δ*bopC*	pABB-CRS2, Δ*bopC* (Δ*bteA*) gene	10
pMGKU401	pDONR201, *bspR* gene	this study
pMGKU402	pDONR201, Δ*bspR* gene	this study
pMGKU403	pABB-CRS2, Δ*bspR* gene	this study

Table 2. Medium compositions

Basal medium (g/L)	SS	LCA
Sodium Gultamate	10.7	10.7
L-proline	0.24	0.24
NaCl	2.5	2.5
KCl	0.2	0.2
KH_2PO_4	0.5	0.5
$MgCl_2 \cdot 6H_2O$	0.1	0.1
$CaCl_2$	0.02	0.02
Tris	6.1	6.1
Casamino acids	10	0.5
Dimethyl-β-cyclodextrin	1	1

Supplement (g/L)	SS	LCA
L-cysteine·HCl	4	4
$FeSO_4 \cdot 7H_2O$	1	1
L-ascorbic acid	40	2
Niacin	0.4	0.4
L-glutathione reduced	15	15

Table 3. Primers used in this study.

Name	Sequence
B1-bspR	5'-AAAAAGCAGGCTCGCCGGTTTCCACCACGTAA-3'
B2-bspR	5'-AGAAAGCTGGGTGGCCAATATCCTGTCTCAGC-3'
R1-bspR	5'-CGCGGATCCCATACGGCGCAAGAATGGCT-3'
R2-bspR	5'-CGCGGATCCAAAGACATCAAGCTCCAAAG-3'
5-recA	5'-ATGAAGATCGGCGTGATGT-3'
3-recA	5'-TAGAACTTGAGCGCGTTGC-3'
5-bteA	5'-AATGGCCTTGGTGGGAAC-3'
3-bteA	5'-ATTTCAGCGCCGTGATCTT-3'
5-bopB	5'-GTCTGCTGACAGCGTTGG-3'
3-bopB	5'-GGGGTTCGATCAGCAAGAC-3'

Restriction enzyme recognition sites were underlined.

Fig. 1. BteA and BopB production by *B. pertussis* cultured in LCA medium. Tohama I was cultured in SS or LCA medium. Whole cell lysates (WCL) and supernatant fractions (Sup) samples were separated by SDS-PAGE and analyzed by western blotting with antibodies against BteA (*upper panel*) or BopB (*lower panel*). BteA forms SDS-resistant multimers (10). NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 2. Effects of ascorbic acid and casamino acids on Bvg-regulated virulence factors.

Tohama I was cultured in SS medium or ascorbic acid-deprived SS medium (SS_AsA⁻),

casamino acid-deprived SS medium (SS_CaA⁻) or both ascorbic acid- and casamino

acid-deprived SS medium (SS AsA⁻_CaA⁻). WCL and Sup samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA, BopB, FHA, or

CyaA. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal

volume of bacterial culture. Experiments were performed at least three times, and

representative data are shown.

Fig. 3. Effects of ascorbic acid on type III secreted proteins. Whole cell lysates and secreted proteins prepared from Tohama I cultured in SS medium in the presence of ascorbic acid at the final concentration of 91, 454 or 2270 μM were separated by SDS-PAGE and then analyzed by western blotting with antibodies against BteA or BopB. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 4. Effects of ascorbic acid on the secretion of BteA and BopB in clinical isolates cultured

in SS_AsA⁻ medium. The *B. pertussis* clinical isolates BP344, BP348, BP350, and BP351

were cultured in SS (+) or SS_AsA⁻ (−) medium. WCL and Sup samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA (*upper panel*) or

BopB (*lower panel*). NS: nonspecific signals. Loaded WCL and Sup samples were prepared

from an equal volume of bacterial culture. Experiments were performed at least three times,

and representative data are shown.

Fig. 5. Results of the RT-PCR analysis for mRNA levels of *bteA* and *bopB* from *B. pertussis* cultured under ascorbic acid-starved conditions. Total RNA was prepared from Tohama I or BP350 cultured in SS or SS_AsA⁻ medium and subjected to a quantitative RT-PCR analysis. The histograms show the relative amounts of *bteA* (**A**) and *bopB* (**B**) mRNA normalized by the housekeeping gene, *recA* mRNA. Error bars: the standard error of the mean (SEM) from triplicate experiments. *p<0.05; N.S.: no significant difference. Experiments were performed at least three times, and representative data are shown.

Fig. 6. Cytotoxic activity of *B. pertussis* cultured in SS_AsA⁻ medium. L2 cells were infected

with Tohama I cultured in SS medium, SS_AsA⁻ medium, SS medium containing

one-twentieth of ascorbic acid (SS_1/20AsA), SS medium containing one-twentieth of

ascorbic acid and one-half of casamino acids (SS_1/20AsA_1/2CaA), or SS medium

containing one-twentieth of ascorbic acid and no casamino acids (SS_1/20AsA_CaA⁻) at an

MOI of 500 for 3 hr. +, 1/2 and 1/20 show the amounts of indicated substances that are equal

to, one-half of, and one-twentieth of those in SS medium, respectively. The amounts of LDH

released into the extracellular medium from infected cells are shown, and the relative

cytotoxicity (%) was determined as described in the Materials and Methods section. Error

bars: the SEM from triplicate experiments. n.a.: No available result. Experiments were

performed at least three times, and representative data are shown.

Fig. 7. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. pertussis*. **A:** L2 cells

were infected with Tohama I or BP350 in the indicated media at an MOI of 125 for 3 hr. +, 1/2

and 1/20 indicate the amounts of indicated substances that are equal to, one-half of, and

one-twentieth of those in SS medium, respectively. After fixation, cells were stained with

anti-Bsp22 antibody (*green*) and rhodamine phalloidin (*red*). *Arrowheads* indicate the Bsp22

(*green*) signal. **B:** Bsp22 signals per one cell were counted under a fluorescent microscope.

At least 120 cells were randomly chosen. Error bars: the SEM from triplicate experiments.

*p<0.05; N.S.: no significant difference. Experiments were performed at least three times, and

representative data are shown.

Fig. 8. Cytotoxic activities of *B. pertussis* mutants. The results of LDH assays at an MOI of 125 (**A**) or 250 (**B**) are shown as a histogram. Error bars: the SEM from triplicate experiments *p<0.05. Experiments were performed at least three times, and representative data are shown. **C:** MGKU1 (Tohama I nalidixic acid-resistant strain, 1: WT Nal^r), MGKU2 (T3SS-inactive strain, 2: ΔT3SS), MGKU3 (BteA-deficient strain, 3: ΔBteA), or MGKU4 (BspR-deficient strain, 4: ΔBspR) was cultured in SS_1/20AsA_1/2CaA medium. WCL and Sup samples were separated by SDS-PAGE and analyzed by western blotting with anti-BteA antibody. We loaded a 3-fold smaller amount of MGKU4 supernatant fraction on the SDS-PAGE gel to avoid obtaining an excess signal intensity. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 9. Effects of reducing agents on the production and secretion of type III secreted proteins

in *B. pertussis*. Tohama I was cultured in SS_AsA⁻ medium or SS_AsA⁻ medium containing

ascorbic acid (AsA), 2-mercaptoethanol (2-ME), or dithioerythritol (DTE) at final

concentrations of 2.3, 2.1, or 3.0 mM, respectively. WCL and Sup samples were separated by

SDS-PAGE, then analyzed by western-blotting with antibodies against BteA, BopB, or FHA.

Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. NS:

nonspecific signals. Experiments were performed at least three times, and representative

data are shown.