Transcriptional Downregulation of a Type III Secretion System under Reducing

Conditions in *Bordetella pertussis*

Masataka Goto[a], Tomoko Hanawa[b], Akio Abe[a] and Asaomi Kuwae[a#]

[a]Laboratory of Bacterial Infection, Graduate School of Infection Control Sciences,

Kitasato University, Tokyo, Japan

[b]Department of Infectious Diseases, Kyorin University School of Medicine, Tokyo,

Japan

Running Head: Reducing Conditions Downregulate *Bordetella* T3SS.

[#]**Address Correspondence to:** Asaomi Kuwae, kuwae@lisci.kitasato-u.ac.jp

ABSTRACT

Bordetella pertussis uses a type III secretion system (T3SS) to inject virulence

proteins into host cells. Although the *B. pertussis* T3SS was presumed to be involved

in host colonization, the efficient secretion of type III secreted proteins from

B. pertussis has not been observed. To investigate the roles of type III secreted

proteins during infection, we attempted to optimize culture conditions for the

production and secretion of a type III secreted protein, BteA, in *B. pertussis*. We

observed that *B. pertussis* efficiently secretes BteA in ascorbic acid-depleted (AsA⁻)

medium. When L2 cells, a rat lung epithelial cell line, were infected with *B. pertussis*

cultured in the AsA⁻ medium, BteA-dependent cytotoxicity was observed. We also

performed an immunofluorescence assay of L2 cells infected with *B. pertussis*. The

clear fluorescence signals of Bsp22, a needle structure of T3SS, were detected on

the bacterial surface of *B. pertussis* cultured in the AsA⁻ medium. Since ascorbic

acid is known as a reducing agent, we cultured *B. pertussis* in liquid medium

containing other reducing agents such as 2-mercaptoethanol and dithioerythritol.

Under these reducing conditions, the production of type III secreted proteins was

repressed. These results suggest that in *B. pertussis*, the production and secretion of

type III secreted proteins are downregulated under reducing conditions.

IMPORTANCE

The type III secretion system (T3SS) of *Bordetella pertussis* forms a needle-like

structure that protrudes from the bacterial cell surface. *B. pertussis* uses T3SS to

translocate virulence proteins called effectors into host cells. The culture conditions

for effector production in *B. pertussis* have not been investigated. We attempted to

optimize culture medium compositions for producing and secreting type III secreted

proteins. We found that *B. pertussis* secretes type III secreted proteins in reducing

agent-deprived liquid medium, and that BteA-secreting *B. pertussis* provokes

cytotoxicity against cultured mammalian cells. These results suggest that redox

signaling is involved in the regulation of *B. pertussis* T3SS.

Introduction

The genus *Bordetella* is made up of Gram-negative bacteria and has been

subdivided into 16 subspecies to date (1). *Bordetella pertussis* is a causative agent

of whooping cough, also known as pertussis. *B. pertussis* strictly adapts to humans,

and it lacks an environmental reservoir (2). A variety of virulence factors have been

identified in *Bordetella*, e.g., filamentous hemagglutinin (3), adenyl cyclase toxin (4)

and pertactin (5). These virulence factors are regulated by a *Bordetella* two-

component regulatory system, BvgAS. In addition to these virulence factors, the type

III secretion system (T3SS) of *B. pertussis* is also regulated by BvgAS (6, 7).

A T3SS that forms a needle-like structure is conserved among many Gram-negative

bacteria. In order to subvert hosts, bacteria use a T3SS to inject type III secreted

proteins into host cells (8). Such type III secreted proteins are called effectors. Three

types of effectors have been identified in *Bordetella*: BteA ((9) [also referred to as

BopC (10)], BopN (11, 12), and BspR (13) [also referred to as BtrA (14, 15)]. BteA is

658 amino acids in length and is localized to lipid rafts in host cells via its N-terminal

region (16). BteA induces necrotic cell death in several types of cultured mammalian

cells (9, 10), and BteA translocation into host cells is promoted by the functioning of

BopN (11). BspR is a negative regulator for type III secreted proteins (13) and is

translocated into nuclei of cultured mammalian cells (17). BspR has been suggested

to be an anti-sigma factor against BtrS that functions as a positive regulator for type

III secreted proteins (15). Some type III secreted proteins are called translocators

rather than effectors. Three types of translocators have been identified in *Bordetella*,

i.e., BopB (18), BopD (19) and Bsp22 (20), and they function as the path for effectors

delivered into host cells. The complex of BopB and BopD forms the pores on the

host membrane (19), and Bsp22 forms a filamentous-like structure and is associated

with the T3SS needle and with a pore-forming component, BopD (21).

The optimal culture conditions for the production and secretion of BteA have

not been established. To investigate the roles of BteA and other type III secreted

proteins during *B. pertussis* infection, we investigated culture conditions for the

maximal production and secretion of type III secreted proteins in *B. pertussis* strain

Tohama I.

MATERIALS AND METHODS

Bacterial strains

The strains used in this study are listed in Table 1. *B. pertussis* Tohama I (22) was

used as the wild-type strain. BP344, BP348, BP350, and BP351 were used as

clinical isolates and were a kind gift from Dr. K. Kamachi (National Institute of

Infectious Diseases) (23). *Escherichia coli* DH10B and Sm10λ*pir* were used for DNA

cloning and conjugation, respectively (Table 1). *B. pertussis* was grown on Bordet-

Gengou agar plates at 37°C for 5 days. Bordet-Gengou agar plates were prepared

as follows. We mixed 15 g of Bordet-Gengou agar base (BD #248200), 5 mL of

glycerol, 100 mL of defibrinated horse blood (Nihon Bio-Supp Center), and 400 mL

of distilled water. The mixture was autoclaved and poured onto plastic plates. Fresh

colonies of *B. pertussis* on each Bordet-Gengou agar plate were suspended in

Stainer-Scholt (SS) medium [Table 2, (24)]. In this study, we used a liquid medium

described as a cyclodextrin-containing medium as the SS medium (25). Low

casamino acids (LCA) medium was prepared by removing the agar from cyclodextrin

solid medium (CSM) contents [Table 2, (26)]. *B. pertussis* was cultured in SS

medium with a starting A_{600} of 0.23 under static conditions at 37ºC. To measure the

bacterial density, a spectrophotometer (Ultrospec 2100 *pro*, Amersham) was used.

The final A_{600} of *B. pertussis* Tohama I culture after 22 hr was ca 0.67, and the

culture contained ca 1.0×10^9 cfu/mL. The liquid cultivation period was 22 hr for the

protein preparation, mRNA preparation, and infection.

Generation of a Tohama I nalidixic acid-resistant mutant

For the generation of a Tohama I nalidixic acid-resistant mutant, Tohama I cultured

in SS medium for 24 hr as described above was spread on 20 sets of Bordet-

Gengou agar plates containing nalidixic acid at the final concentration of 30 μg/ml.

After 5 days, surviving colonies were designated as MGKU1. The growth rate, colony

size and hemolysis on the BG plates of the resulting nalidixic acid-resistant strain

were almost equivalent to those of the wild-type strain, and we observed that the

amounts of the production and secretion of type III secreted proteins in MGKU1 were

not significantly different from those of Tohama I wild-type (data not shown).

Cell culture

L2 cells, a rat lung epithelial cell line (ACTT CCL-149), were maintained in F-12K

(Invitrogen, Tokyo). The cell culture medium was supplemented with 10% fetal calf

serum (FCS). L2 cells were grown at 37°C under a 5% CO_2 atmosphere.

Construction of gene-disrupted mutants

The primers used in this study are listed in Table 3. To construct the *bspR* mutant,

the 4.0 kb DNA fragment encoding *bspR* and its flanking regions was amplified by

polymerase chain reaction (PCR) with primers B1-bspR and B2-bspR using

B. pertussis Tohama I genomic DNA as the template. The resulting PCR product

was cloned into pDONR201 to obtain pMGKU401, by means of adapter PCR and

site-specific recombination techniques using a Gateway cloning system (Invitrogen).

An inverse PCR was carried out with the set of primers R1-bspR and R2-bspR using

circular pMGKU401 as the template.

The resulting PCR product was digested with BamHI and self-ligated to obtain

pMGKU402. This plasmid contained the BamHI site in addition to a 474-bp in-frame

deletion from 175 bp downstream of the 5' end of the *bspR* gene to 52 bp upstream

of the 3' end of the gene. This plasmid, pMGKU402, was mixed with a positive

suicide vector, pABB-CRS2 (27), to obtain pMGKU403 using the Gateway cloning

system. This plasmid, pMGKU403, was introduced into *E. coli* Sm10λ*pir* (18) and

transconjugated into MGKU1 as described previously (28). The resulting mutant

strain was designated as MGKU4 (BspR-deficient strain). Similarly, pABB-Δ*bscN*

(18) and pABB-Δ*bopC* (10) were separately introduced into *E. coli* Sm10λ*pir* and

were transconjugated into MGKU1. The resulting mutant strains were designated as

MGKU2 (a T3SS-inactive strain) and MGKU3 (a BteA-deficient strain), respectively.

Preparation of proteins from culture supernatants and whole bacterial cell

lysates

Secreted proteins released into bacterial culture supernatants and whole bacterial

cell lysates were prepared by trichloroacetic acid precipitation. In the case of the

supernatant fraction (Sup) samples, the culture supernatants were filtered with a

0.22 μm filter (Millipore #SLGVR33RB). Next, 2 μL of 5% deoxycholic acid and 100

μL of 100% trichloroacetic acid were added to the 1 mL of filtered supernatants. After

being incubated on ice for 15 min, the samples were centrifuged at 21130 ×g for 5

min. The resulting precipitated proteins were neutralized with 2 μL of 2 M Tris-base

and dissolved in 13 μL of 2×SDS-PAGE sample buffer. In the case of the whole cell

lysate (WCL) samples, bacterial pellets collected from 1 mL of bacterial cultures

were resuspended in distilled water. 100 μL of 100% trichloroacetic acid was added

to 1 mL of the bacterial suspension. After being incubated on ice for 15 min, the

samples were centrifuged at 21130 ×g for 5 min. The resulting precipitated proteins

were neutralized with 5 μL of 2 M Tris-base and dissolved in 95 μL of 2×SDS-PAGE

sample buffer. When the A_{600} of bacterial culture was 0.67, 0.5 μL of WCL and 5 μL

of Sup samples were loaded on the SDS-PAGE gels. The loaded sample amounts

were adjusted by the A_{600} of each bacterial culture in order to load samples prepared

from the same number of bacteria. The protein samples were separated by SDS-

PAGE and analyzed by western blotting.

Antibodies

Anti-BteA, anti-BopB, and anti-Bsp22 antibodies were purified from rabbit serum in

our previous study (10, 18, 29). To detect filamentous hemagglutinin (FHA) signals,

we used mouse anti-FHA serum (30). Mouse anti-CyaA monoclonal antibody was

purchased from Santa Cruz Biotechnology (Santa Cruz, CA). To prepare the anti-

BtrS antibody, the peptide corresponding to the C-terminus region of BtrS

(CALREALRERGYDSVP) were conjugated with hemocyanin from keyhole limpets

(Sigma) by using 3-maleimidobenzoic acid N-hydroxysuccinimide ester (Sigma).

These cross-linked peptides were used to immunize rabbits, and resulting anti-sera

were incubated with peptide immobilized on epoxy-activated sepharose 6B

(Amersham) to obtain specific Ig-fractions.

Quantitative reverse transcription-PCR (qRT-PCR)

The amounts of mRNA were measured by qRT-PCR. Bacterial total RNA was

prepared using a Trizol Max Bacterial RNA Isolation Kit (Invitrogen), an RNeasy Mini

Kit (Qiagen, Hilden, Germany) and an RNase-free DNase-free Kit (Qiagen). The

RNA sample was reverse transcribed with Transcriptor Universal cDNA Master

(Roche Diagnostics, Indianapolis, IN) and a T100 Thermal Cycler (Bio-Rad, Hercules,

CA). The resulting cDNA was amplified by FastStart Essential DNA Probes Master

(Roche) using the following primer pairs: 5-recA and 3-recA for *recA*; 5-bteA and 3-

bteA for *bteA*; 5-bopB and 3-bopB for *bopB* (Table 3). The results were calculated as

described in the Roche manual. The amount of *recA* mRNA was used as an internal

control. The mRNA amounts are presented herein as relative to the amounts in the

Tohama I cultured in SS medium, which was defined as 1.

LDH Assays

To examine the release of lactate dehydrogenase (LDH) from *B. pertussis*-infected

cells, 5.0×10^4 cells/well of L2 cells seeded in 24-well plates were infected with

bacteria at the multiplicity of infection (MOI) of 125, 250, or 500. The plates were

centrifuged at 900 g for 5 min and incubated for 3 hr at 37°C under a 5% CO_2

atmosphere. The amounts of LDH were measured spectrophotometrically using a

Cyto-Tox 96 Non-radioactive Cytotoxicity Assay Kit (Promega, Madison, WI). We

added 50 μL of 10 %Triton X-100 to the 950 μL of extracellular medium of L2 cells in

24-well plate. We then mixed the extracellular medium by pipetting. We subtracted

the LDH value obtained from extracellular medium of uninfected cells from the value

obtained from the Triton X-100-treated cells and used the resulting value as 100%.

Immunofluorescent staining

For the immunofluorescent staining assay, 2.5×10^5 cells/well of L2 cells were

seeded on coverslips in six-well plates and incubated overnight, and then infected

with *B. pertussis* at an MOI of 125. The plates were centrifuged at 900 g for 5 min

and incubated for 3 hr at 37°C under a 5% CO_2 atmosphere. The infected L2 cells

were then immunostained as described previously (12). Briefly, the cells were

treated with 4% paraformaldehyde, 50 mM NH_4Cl, 0.2% Triton-X, and 4% bovine

serum albumin (BSA). After blocking, the cells were stained by anti-Bsp22 antibody

(29). As a secondary antibody, Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen) was

used. F-actin was stained with Rhodamine Phalloidin (Molecular Probes, Eugene,

OR). Bsp22 signals per one cell were counted under a fluorescence microscope.

Statistical analyses

The statistical analyses were performed using the nonparametric unpaired *t*-test with a one-tailed p-value with Prism ver. 5.0f software (Graphpad, La Jolla, CA). Values of $p < 0.05$ were considered significant.

RESULTS

Type III secreted proteins are secreted from *B. pertussis* cultured in LCA medium

Cyclodextrin solid medium (CSM) is reported as a synthetic medium for clinical isolation of *B. pertussis* (26). We sought to determine whether *B. pertussis* grown on this agar secretes type III effectors, and we prepared a liquid medium, designated "low casamino acids (LCA)" medium, by removing the agar from CSM contents (Table 2). *B. pertussis* was cultured in SS or LCA medium, and then the whole cell lysates (WCL) and supernatant fractions (Sup) samples were subjected to western blotting with antibodies against BteA (an effector, a type III secreted protein) and BopB (a translocator, a type III secreted protein).

   BteA and BopB signals were detected in WCL and Sup of the wild-type Tohama I cultured in LCA medium (Fig. 1). As reported previously, BteA forms SDS-resistant multimers, and in our present study we also detected the signals at around 200 kDa. In contrast, the BteA and BopB signals were faintly detected in the WCL samples of SS medium, and no BteA and BopB signals were detected in the Sup samples of MGKU2 (a T3SS-inactive strain) or the wild-type cultured in SS medium (Fig. 1). These results suggest that *B. pertussis* secretes type III secreted proteins in LCA medium.

Ascorbic acid downregulates the production and secretion of type III secreted proteins in *B. pertussis*

Although the ingredients of the LCA and SS media are the same, the amounts of ascorbic acid and casamino acids in LCA medium are lower than those in SS medium (Table 2). To determine which substance is involved in the production and secretion of type III secreted protein, we cultured *B. pertussis* in SS medium, ascorbic acid-deprived SS medium (SS_AsA$^-$), casamino acid-deprived SS medium (SS_CaA$^-$), or both ascorbic acid- and casamino acid-deprived medium (SS_AsA$^-$_CaA$^-$). WCL and Sup samples were subjected to western blotting with antibodies against BteA and BopB (Fig. 2).

The BteA multimer signal intensities in the WCL samples of the SS_AsA$^-$, SS_CaA$^-$ and SS_AsA$^-$_CaA$^-$ media were stronger than that in the SS medium (Fig. 2). A faint BteA signal intensity in the Sup samples of SS was detected (Fig. 2). The BteA signal intensities in the Sup samples were evident in the SS_AsA$^-$ and SS_AsA$^-$_CaA$^-$ media (Fig. 2). Again, BopB signal intensities in both WCL and Sup samples were evident in the SS_AsA$^-$ and SS_AsA$^-$_CaA$^-$ media (Fig. 2).

We also investigated the potential involvement of ascorbic acid or casamino acids in Bvg-regulated virulence factors, e.g., FHA and CyaA. WCL and Sup samples were subjected to western blotting with antibodies against FHA and CyaA (Fig. 2). The results revealed that there were no significant differences in the FHA or CyaA signal intensities in the WCL and Sup samples among the media used here (Fig. 2). Collectively, these results suggest that ascorbic acid has a specific influence on the production and secretion of BteA and BopB in *B. pertussis*.

Standard SS medium contains ascorbic acid at a final concentration of 2270 μM. To

further explore whether ascorbic acid affects the production and secretion of type III

secreted proteins, we cultured *B. pertussis* in SS medium in the presence of ascorbic

acid at the final concentrations of 91, 454, or 2270 μM. WCL and Sup samples were

prepared from each bacterial culture and analyzed by western blotting (Fig. 3). The

BteA signal was detected in both WCL and Sup samples of SS medium at 0–454 μM

ascorbic acid, but the BteA signal of the SS medium at 2270 μM ascorbic acid was

faint or absent (Fig. 3). The BopB signals showed a similar pattern (Fig. 3). These

results indicate that *B. pertussis* produces and secretes type III secreted proteins

under low ascorbic acid concentrations.

BteA and BopB are secreted from other *B. pertussis* clinical isolates cultured

in SS_AsA$^-$ medium

To determine whether *B. pertussis* clinical isolates secrete type III secreted proteins

under low ascorbic acid concentrations, we cultured four strains of *B. pertussis*

(Table 1) in SS medium or ascorbic acid-deprived medium (SS_AsA$^-$). The prepared

WCL and Sup samples were analyzed by western blotting (Fig. 4). BteA signals in

the Sup samples of BP350 and BP351 were evident in the SS_AsA$^-$ medium (Fig. 4,

AsA$^-$). Again, BopB signals in the Sup samples of BP350 and BP351 were evident in

SS_AsA$^-$ medium. These results suggest that some clinical strains, e.g., BP350 and

BP351, also secrete type III secreted proteins under low ascorbic acid

concentrations.

Translocator BopB is upregulated at the transcriptional level in *B. pertussis*

cultured in SS_AsA⁻ medium

To investigate the gene expressions of type III secreted proteins under ascorbic

acid-starved conditions, we prepared total RNA from *B. pertussis* cultured in SS or

SS_AsA⁻ medium. The cDNA samples reverse-transcribed from the total RNA

samples were subjected to a quantitative RT-PCR analysis to quantify the relative

amounts of *bteA* and *bopB* mRNA. The results demonstrated that the relative

amount of *bteA* mRNA of Tohama I in SS_AsA⁻ medium was increased compared to

that in SS medium (Fig. 5A). However, the relative amount of *bteA* mRNA of BP350

in the SS_AsA⁻ medium was not significantly different from that in the SS medium

(Fig. 5A). The relative amounts of *bopB* mRNA of Tohama I and BP350 in SS_AsA⁻

medium were increased compared to those in SS medium (Fig. 5B).

These results suggest that BopB is upregulated at the transcriptional level in

Tohama I and BP350 under low ascorbic acid concentrations. BteA is also

upregulated at the transcriptional level in Tohama I under low ascorbic acid

concentrations but not in BP350.

***B. pertussis* induces cytotoxicity against cultured mammalian cells**

A *B. pertussis* BspR-deficient strain was reported to induce cytotoxicity against

cultured mammalian cells (15). In the present study, we identified better culture

conditions for the production and secretion of BteA in *B. pertussis* than those

provided with standard SS medium. We thus investigated whether wild-type

B. pertussis provokes cytotoxicity against cultured mammalian cells under these

conditions. L2 cells, a rat lung epithelial cell line, were infected at an MOI of 500 for 3

hr in several modified SS media shown in Fig. 6. The amounts of ascorbic acid

and/or casamino acids was reduced to one-twentieth of and/or to one-half of those in

SS medium. We also used the SS media without ascorbic acid and/or casamino

acids.

The amounts of lactate dehydrogenase (LDH) released into the extracellular

medium were then measured.

The results revealed that released LDH was detected from L2 cells infected with

Tohama I in the SS_1/20AsA or SS_1/20AsA_1/2CaA medium (Fig. 6). The

SS_1/20AsA medium and SS_1/20AsA_1/2CaA medium showed no cytotoxicity. On

the other hand, the L2 cells incubated in the SS_AsA$^-$ or SS_1/20AsA_CaA$^-$

medium were injured by the medium itself. Therefore, these media were not useful

for this assay. Taken together, these results demonstrate that we successfully

observed cytotoxicity of mammalian cells induced by wild-type *B. pertussis*.

The T3SS of *B. pertussis* cultured in SS_AsA$^-$ medium is active during

infection against cultured mammalian cells

As described above, we observed cytotoxicity of *B. pertussis* against cultured cells.

In order to confirm that *B. pertussis* indeed secretes type III secreted proteins during

infection, we infected L2 cells with *B. pertussis* in SS or SS_1/20AsA_1/2CaA

medium at an MOI of 125 for 3 hr. After infection, Bsp22 (a translocator, and a

component of the filamentous-like structure) and F-actin were stained with anti-

Bsp22 and rhodamine phalloidin, respectively.

The results showed that the amounts of Bsp22 signals (Fig. 7A, green) on L2 cells

infected with Tohama I and BP350 cultured in SS_1/20AsA_1/2CaA medium were

greater than those in SS medium (Fig. 7). Thus, we demonstrated that *B. pertussis*

T3SS is activated during the infection of cultured mammalian cells under appropriate

conditions.

The cytotoxicity induced by wild-type *B. pertussis* depends on the T3SS

It has been reported that *B. bronchiseptica* induces BteA-dependent necrotic cell

death against cultured mammalian cells (10). To investigate whether the cytotoxicity

provoked by wild-type *B. pertussis* also depends on BteA function, we infected L2

cells with MGKU1 (a Tohama I nalidixic acid-resistant strain), MGKU2 (a T3SS-

inactive strain), MGKU3 (a BteA-deficient strain), or MGKU4 (a BspR-deficient strain

that allows the constitutive activation of T3SS) in SS_1/20AsA_1/2CaA medium at

an MOI of 125 or 250 for 3 hr. The amounts of LDH released into extracellular

medium were measured.

As a result, LDH was not detected in the medium of L2 cells infected with MGKU2

(Fig. 8A,B). The amount of LDH detected in the medium of L2 cells infected with

MGKU3 was significantly lower than that in the medium of L2 cells infected with

MGKU1 (Fig. 8A,B). The amount of LDH detected in the medium of L2 cells infected

with MGKU4 was significantly greater than that in the medium of L2 cells infected

with MGKU1 at an MOI of 125 (Fig. 8A), but not at an MOI of 250 (Fig. 8B).

WCL and Sup samples were prepared from each strain and separated by SDS-

PAGE, then analyzed by western blotting with anti-BteA antibody (Fig. 8C). BteA

signals of the WCL samples were detected in MGKU1, MGKU2, and MGKU4, but

not in MGKU3 (Fig. 8C). BteA signals of the Sup samples were detected in MGKU1

and MGKU4, but not in MGKU2 or MGKU3 (Fig. 8C). These results suggest that the

cytotoxicity induced by wild-type *B. pertussis* (Fig. 6) is dependent on T3SS activity

and is partially dependent on BteA.

The secretion of type III secreted proteins in *B. pertussis* is downregulated

under reducing conditions

Ascorbic acid is known as a reducing agent. To determine whether the secretion of

type III secreted proteins is downregulated under reducing conditions, we cultured

Tohama I in ascorbic acid-deprived SS medium (SS_AsA$^-$, Fig. 9, "none") and in

SS_AsA$^-$ medium containing the reducing agents ascorbic acid (AsA), 2-

mercaptoethanol (2-ME) or dithioerythritol (DTE), respectively. The prepared WCL

and Sup samples were analyzed by western blotting with anti-BteA, anti-BopB, or

anti-FHA antibodies (Fig. 9). The BteA and BopB signal intensities of both WCL and

Sup samples were decreased in the presence of each of the reducing agents (Fig. 9).

In contrast, the FHA signal intensities of both WCL and Sup samples were not

affected by the reducing agents. The growth of bacteria cultured in SS medium with

reducing agents was not significantly different from that of the bacteria cultured in SS

medium (data not shown). These results suggest that the production and secretion of

type III secreted proteins in *B. pertussis* are downregulated under reducing

conditions.

DISCUSSION

In this study, *B. pertussis* increased the production and secretion of type III secreted

proteins, BteA and BopB, when cultured in ascorbic acid-deprived SS medium

(SS_AsA$^-$). We successfully produced cytotoxicity against mammalian cells by the

infection of wild-type Tohama I cultured in SS 1/20AsA_1/2CaA medium.

B. pertussis decreased the production and secretion of type III secreted proteins

when cultured in SS_AsA⁻ medium containing reducing agents such as 2-ME or DTT.

These results suggest that redox signaling is involved in regulation of the T3SS in

B. pertussis.

It has been reported that no secretion of the type III secreted proteins BteA

and Bsp22 was detected from Tohama I (31), and that the secretion of the type III

secreted proteins BteA and BopD was detected from a non-vaccine-type strain,

BP157. However, BP157 showed no cytotoxicity against L2 cells, J774 mouse

macrophage-like cells, and HeLa cells (32). In another report, glutamate limitation

upregulates production of the type III secreted proteins (33). However, the

requirement of a type III secretion system (T3SS) for virulence in *B. pertussis* thus

remains unclear. On the other hand, a BscN-deficient mutant in which the T3SS was

inactive colonized mouse lungs to a significantly lower degree compared to the wild-

type (31). In addition, cytotoxicity against HeLa cells was observed by infection with

a BspR (a negative regulator for type III secreted proteins)-deficient *B. pertussis*

strain (15). These results suggest that the T3SS has a significant role in the

virulence of *B. pertussis*.

The production and the relative amounts of *bopB* mRNA of Tohama I cultured

in SS AsA⁻ medium were significantly increased compared to those of Tohama

cultured in SS medium, but this was not the case for BteA mRNA (Figs. 2, 5). It was

reported that *bteA* and *bopB* genes are upregulated by BtrS (a sigma factor for type

III secreted proteins) and downregulated by BspR (13, 15). It has also been reported

that the amount of *bteA* mRNA was increased 2-fold in a BspR-deficient strain

compared to the wild-type, whereas the amount of *bopB* mRNA was increased by

18-fold compared to the wild-type by RNA-seq (15). The *bopB* gene is localized in

the T3SS apparatus locus (*bsc* locus) adjacent to the *btrS* gene. Since the *bteA*

gene is separated from the *bsc* locus by >2.5 Mb (9), BtrS would preferentially

regulate transcription of the *bopB* gene over that of the *bteA* gene.

The cytotoxicity against L2 cells was observed by the infection of Tohama I

cultured in SS_1/20AsA _1/2CaA medium (Fig. 6). Before infection, we exchanged

Ham's F-12K medium for the SS_1/20AsA _1/2CaA medium. When L2 cells were

infected with Tohama I in Ham's F-12K medium, the cytotoxicity against L2 cells was

not observed (Fig. S1). In order to establish oxidizing conditions in the tissue culture

media, we used several oxidizing agents, i.e., hydrogen peroxide (H_2O_2), potassium

ferricyanide(III) ($K_3[Fe(CN)_6]$), methyl 3-nitro-2-pyridinesulfenate (Npys-OMe), and

5,5'-dithiobis(2-nitrobenzoic acid) (DTNB). We also used another tissue culture

medium, i.e., Minimum Essential Medium Eagle (MEM), because MEM contains no

reducing agents, while F-12K, which is the medium for the L2 cells culture, contains

a reducing agent, i.e., cysteine. Even when L2 cells were infected in F-12K medium

containing the oxidizing agent or MEM, *B. pertussis* showed no cytotoxicity (Fig. S1).

We therefore speculate that a substance which inhibits the secretion of type III

secreted proteins from Tohama I was present in the Ham's F-12K medium, or a

substance that is necessary for the secretion of type III secreted proteins from

Tohama I was absent in the Ham's F-12K medium. We found that infection of the

human alveolar epithelium cell line A549 with wild-type *B. pertussis* cultured in

SS_1/20AsA _1/2CaA medium did not result in cytotoxicity (data not shown). For the

observation of cytotoxicity against A549 cells, it is necessary to optimize the

conditions for the production and secretion of type III secreted proteins in

B. pertussis.

A previous study showed that cytotoxicity against the cell line J774A.1 was

achieved by infection with wild-type *B. pertussis* 18323, but not by infection with a

CyaA-deficient strain (34). This finding suggested that CyaA has the potential to

provoke the cytotoxicity. However, we confirmed herein that the production and

secretion of CyaA in a BteA-deficient strain (MGKU3) were not significantly different

from those of the wild-type (data not shown). In our results, the amounts of LDH

released from L2 cells infected with MGKU3 were greater than those of the T3SS-

inactive strain (MGKU2), and no LDH release was detected from L2 cells infected

with MGKU2 (Fig. 8). These results raised the possibility that an unknown type III

effector other than BteA is involved in the cytotoxicity of *B. pertussis*.

Although we observed no significant difference in the growth rate of Tohama I

between SS medium and SS_AsA⁻ medium (data not shown), the production and

secretion of type III secreted proteins of Tohama I cultured in the SS_AsA⁻ medium

were increased compared to those in the SS medium (Fig. 2). It had not been

determined whether the expression of type III secreted proteins is regulated by

reducing agents at the molecular level, and in this study, the production and

secretion of CyaA and FHA in Tohama I cultured in SS medium were not significantly

different from those in SS_AsA⁻ medium (Fig. 2). It was reported that there is no

difference in Prn production by Tohama I cultured in modified SS medium compared

to culture in modified SS medium containing dithiothreitol (DTT), a reducing agent

(35). These results suggest that BvgAS-regulated virulence factors other than type III

secreted proteins are not affected by reducing agents.

Ascorbic acid, urea, and glutathione (GSH) function as reducing agents in the

epithelial lining fluid (ELF) of the human airway. The concentrations of ascorbic acid

and urea in ELF in the upper respiratory tract are not significantly different from

those in the lower respiratory tract. However, the concentration of GSH in the ELF in

the upper respiratory tract is 100-fold lower than that in the lower respiratory tract.

One of the possibilities is that GSH forms GS-SG (an oxidized from of GSH) in the

ELF in the upper respiratory tract (36). It may thus be considered that the upper

respiratory tract is more oxidized than the lower respiratory tract. It was reported that

the transcription of genes encoding type III secreted proteins of *B. pertussis*

recovered from mouse nasal lavage was higher than that from bronchoalveolar

lavage (37). We thus speculate that the T3SS is important for *B. pertussis* to

colonize the upper respiratory tract of the host.

In this study, the production and secretion of BopB by Tohama I cultured in the

SS_AsA$^-$ medium were significantly increased compared to those in SS medium (Fig.

2). Bacteria use redox-sensing proteins to resist oxidative stress (38). For example,

OxyR (a member of the LysR family of transcriptional regulators) of *Escherichia coli*

is converted to an active mode by the oxidation of intramolecular cysteine residues,

which then increases the expression of antioxidants genes, e.g., hydroperoxidase I

[*katG*, (39)]. Vfr (a member of the CRP/FNR superfamily) of *Pseudomonas* forms an

active mode by retaining intramolecular cysteine residues as reducing states, and

then increases the expression of type III secreted proteins (40). Btr is a regulator

protein produced by *B. pertussis* and is homologous to Vfr (41). A cysteine cluster is

located in the N-terminal domain of Btr, and a helix-turn-helix DNA binding domain is

located in the C-terminal domain of Btr. Therefore, when *B. pertussis* is cultured in

reducing agent-deprived SS medium, Btr might be oxidized and change its

conformation, thereby increasing the expression of type III secreted proteins.

Although the signal pathway that *B. pertussis* uses to increase the production of type

III secreted proteins is unclear, it is possible that Btr regulates gene expressions for

type III secreted proteins.

REFERENCES

1.   Ivanov YV, Linz B, Register KB, Newman JD, Taylor DL, Boschert KR, Le

Guyon S, Wilson EF, Brinkac LM, Sanka R, Greco SC, Klender PM, Losada L,

Harvill ET. 2016. Identification and taxonomic characterization of *Bordetella*

pseudohinzii sp. nov. isolated from laboratory-raised mice. Int J Syst Evol

Microbiol 66:5452-5459.

2.   Mattoo S, Cherry JD. 2005. Molecular pathogenesis, epidemiology, and

clinical manifestations of respiratory infections due to *Bordetella pertussis* and

other *Bordetella* subspecies. Clin Microbiol Rev 18:326-82.

3.   Relman DA, Domenighini M, Tuomanen E, Rappuoli R, Falkow S. 1989.

Filamentous hemagglutinin of *Bordetella pertussis*: nucleotide sequence and

crucial role in adherence. Proc Natl Acad Sci U S A 86:2637-41.

4.   Hewlett EL, Gordon VM, McCaffery JD, Sutherland WM, Gray MC. 1989.

Adenylate cyclase toxin from *Bordetella pertussis*. Identification and

purification of the holotoxin molecule. J Biol Chem 264:19379-84.

5.   Roberts M, Fairweather NF, Leininger E, Pickard D, Hewlett EL, Robinson A,

Hayward C, Dougan G, Charles IG. 1991. Construction and characterization

of *Bordetella pertussis* mutants lacking the vir-regulated P.69 outer membrane

protein. Mol Microbiol 5:1393-404.

6.   Melvin JA, Scheller EV, Miller JF, Cotter PA. 2014. *Bordetella pertussis*

pathogenesis: current and future challenges. Nat Rev Microbiol 12:274-88.

7.   Cummings CA, Bootsma HJ, Relman DA, Miller JF. 2006. Species- and

strain-specific control of a complex, flexible regulon by *Bordetella* BvgAS. J

Bacteriol 188:1775-85.

8. Cornelis GR, Van Gijsegem F. 2000. Assembly and function of type III

secretory systems. Annu Rev Microbiol 54:735-74.

9. Panina EM, Mattoo S, Griffith N, Kozak NA, Yuk MH, Miller JF. 2005. A

genome-wide screen identifies a *Bordetella* type III secretion effector and

candidate effectors in other species. Mol Microbiol 58:267-79.

10. Kuwae A, Matsuzawa T, Ishikawa N, Abe H, Nonaka T, Fukuda H, Imajoh-

Ohmi S, Abe A. 2006. BopC is a novel type III effector secreted by *Bordetella*

bronchiseptica and has a critical role in type III-dependent necrotic cell death.

J Biol Chem 281:6589-600.

11. Abe A, Nishimura R, Kuwae A. 2017. *Bordetella* effector BopN is translocated

into host cells via its N-terminal residues. Microbiol Immunol 61:206-214.

12. Nagamatsu K, Kuwae A, Konaka T, Nagai S, Yoshida S, Eguchi M, Watanabe

M, Mimuro H, Koyasu S, Abe A. 2009. *Bordetella* evades the host immune

system by inducing IL-10 through a type III effector, BopN. J Exp Med

206:3073-88.

13. Kurushima J, Kuwae A, Abe A. 2012. The type III secreted protein BspR

regulates the virulence genes in *Bordetella bronchiseptica*. PLoS One

7:e38925.

14. Ahuja U, Liu M, Tomida S, Park J, Souda P, Whitelegge J, Li H, Harvill ET,

Parkhill J, Miller JF. 2012. Phenotypic and genomic analysis of hypervirulent

human-associated *Bordetella bronchiseptica*. BMC Microbiol 12:167.

15. Ahuja U, Shokeen B, Cheng N, Cho Y, Blum C, Coppola G, Miller JF. 2016.

Differential regulation of type III secretion and virulence genes in *Bordetella*

pertussis and *Bordetella bronchiseptica* by a secreted anti-sigma factor. Proc

Natl Acad Sci U S A 113:2341-8.

16.   French CT, Panina EM, Yeh SH, Griffith N, Arambula DG, Miller JF. 2009.

The *Bordetella* type III secretion system effector BteA contains a conserved

N-terminal motif that guides bacterial virulence factors to lipid rafts. Cell

Microbiol 11:1735-49.

17.   Abe A, Nishimura R, Tanaka N, Kurushima J, Kuwae A. 2015. The *Bordetella*

Secreted Regulator BspR Is Translocated into the Nucleus of Host Cells via

Its N-Terminal Moiety: Evaluation of Bacterial Effector Translocation by the

Escherichia coli Type III Secretion System. PLoS One 10:e0135140.

18.   Kuwae A, Ohishi M, Watanabe M, Nagai M, Abe A. 2003. BopB is a type III

secreted protein in *Bordetella bronchiseptica* and is required for cytotoxicity

against cultured mammalian cells. Cell Microbiol 5:973-83.

19.   Nogawa H, Kuwae A, Matsuzawa T, Abe A. 2004. The type III secreted

protein BopD in *Bordetella bronchiseptica* is complexed with BopB for pore

formation on the host plasma membrane. J Bacteriol 186:3806-13.

20.   Yuk MH, Harvill ET, Cotter PA, Miller JF. 2000. Modulation of host immune

responses, induction of apoptosis and inhibition of NF-kappaB activation by

the *Bordetella* type III secretion system. Mol Microbiol 35:991-1004.

21.   Medhekar B, Shrivastava R, Mattoo S, Gingery M, Miller JF. 2009. *Bordetella*

Bsp22 forms a filamentous type III secretion system tip complex and is

immunoprotective in vitro and in vivo. Mol Microbiol 71:492-504.

22.   Kasuga T, Nakase Y, Ukishima K, Takatsu K. 1954. Studies on *Haemophilus*

pertussis. V. Relation between the phase of bacilli and the progress of the

whooping-cough. Kitasato Arch Exp Med 27:57-62.

23.   Otsuka N, Han HJ, Toyoizumi-Ajisaka H, Nakamura Y, Arakawa Y,

Shibayama K, Kamachi K. 2012. Prevalence and genetic characterization of

pertactin-deficient *Bordetella pertussis* in Japan. PLoS One 7:e31985.

24.   Stainer DW, Scholte MJ. 1970. A simple chemically defined medium for the

production of phase I *Bordetella pertussis*. J Gen Microbiol 63:211-20.

25.   Imaizumi A, Suzuki Y, Ono S, Sato H, Sato Y. 1983. Effect of heptakis (2,6-O-

dimethyl) beta-cyclodextrin on the production of pertussis toxin by *Bordetella*

pertussis. Infect Immun 41:1138-43.

26.   Aoyama T, Murase Y, Iwata T, Imaizumi A, Suzuki Y, Sato Y. 1986.

Comparison of blood-free medium (cyclodextrin solid medium) with Bordet-

Gengou medium for clinical isolation of *Bordetella pertussis*. J Clin Microbiol

23:1046-8.

27.   Sekiya K, Ohishi M, Ogino T, Tamano K, Sasakawa C, Abe A. 2001.

Supermolecular structure of the enteropathogenic *Escherichia coli* type III

secretion system and its direct interaction with the EspA-sheath-like structure.

Proc Natl Acad Sci U S A 98:11638-43.

28.   Donnenberg MS, Kaper JB. 1991. Construction of an eae deletion mutant of

enteropathogenic *Escherichia coli* by using a positive-selection suicide vector.

Infect Immun 59:4310-7.

29.   Kurushima J, Kuwae A, Abe A. 2012. Btc22 chaperone is required for

573   secretion and stability of the type III secreted protein Bsp22 in *Bordetella*

574   *bronchiseptica*. FEMS Microbiol Lett 331:144-51.

575   30.   Watanabe M, Nagai M, Funaishi K, Endoh M. 2000. Efficacy of chemically

576   cross-linked antigens for acellular pertussis vaccine. Vaccine 19:1199-203.

577   31.   Fennelly NK, Sisti F, Higgins SC, Ross PJ, van der Heide H, Mooi FR, Boyd A,

578   Mills KHG. 2008. *Bordetella pertussis* expresses a functional type III secretion

579   system that subverts protective innate and adaptive immune responses.

580   Infection and Immunity 76:1257-1266.

581   32.   Han HJ, Kuwae A, Abe A, Arakawa Y, Kamachi K. 2011. Differential

582   expression of type III effector BteA protein due to IS481 insertion in *Bordetella*

583   *pertussis*. PLoS One 6:e17797.

584   33.   Hanawa T, Kamachi K, Yonezawa H, Fukutomi T, Kawakami H, Kamiya S.

585   2016. Glutamate Limitation, BvgAS Activation, and (p)ppGpp Regulate the

586   Expression of the *Bordetella pertussis* Type 3 Secretion System. J Bacteriol

587   198:343-51.

588   34.   Khelef N, Zychlinsky A, Guiso N. 1993. *Bordetella pertussis* induces apoptosis

589   in macrophages: role of adenylate cyclase-hemolysin. Infect Immun 61:4064-

590   71.

591   35.   Stenson TH, Patton AK, Weiss AA. 2003. Reduced glutathione is required for

592   pertussis toxin secretion by *Bordetella pertussis*. Infect Immun 71:1316-20.

593   36.   van der Vliet A, O'Neill CA, Cross CE, Koostra JM, Volz WG, Halliwell B,

594   Louie S. 1999. Determination of low-molecular-mass antioxidant

concentrations in human respiratory tract lining fluids. Am J Physiol 276:L289-

96.

37. van Beek LF, de Gouw D, Eleveld MJ, Bootsma HJ, de Jonge MI, Mooi FR,

Zomer A, Diavatopoulos DA. 2018. Adaptation of *Bordetella pertussis* to the

Respiratory Tract. J Infect Dis 217:1987-1996.

38. Lee SJ, Kim DG, Lee KY, Koo JS, Lee BJ. 2018. Regulatory mechanisms of

thiol-based redox sensors: lessons learned from structural studies on

prokaryotic redox sensors. Arch Pharm Res 41:583-593.

39. Aslund F, Zheng M, Beckwith J, Storz G. 1999. Regulation of the OxyR

transcription factor by hydrogen peroxide and the cellular thiol-disulfide status.

Proc Natl Acad Sci U S A 96:6161-5.

40. Zhang Y, Zhang C, Du X, Zhou Y, Kong W, Lau GW, Chen G, Kohli GS, Yang

L, Wang T, Liang H. 2019. Glutathione Activates Type III Secretion System

Through Vfr in *Pseudomonas aeruginosa*. Front Cell Infect Microbiol 9:164.

41. Bannan JD, Moran MJ, MacInnes JI, Soltes GA, Friedman RL. 1993. Cloning

and characterization of btr, a *Bordetella pertussis* gene encoding an FNR-like

transcriptional regulator. J Bacteriol 175:7228-35.

ACKNOWLEDGMENTS

This work was supported in part by grants from the Ministry of Education, Culture,

Sports, Sciences and Technology and the Japan Society for the Promotion of

Science (KAKENHI), nos. 19K07542 (to A.A.), 17K08838 (A.K.), and 19K07561

(T.H.), and by a research grant from the Takeda Science Foundation in 2016 (A.K.).

The funders had no role in the study design, data collection or analysis, the decision

to publish, or the preparation of the manuscript.

Figure legends

Fig. 1. BteA and BopB production by *B. pertussis* Tohama I cultured in LCA medium.

Tohama I and MGKU2 (a T3SS-inactive strain) were cultured in SS or LCA medium.

Whole cell lysates (WCL) and supernatant fraction (Sup) samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA (*upper*

panel), BopB (*middle panel*) or BtrS (*lower panel*). BtrS, an RNA polymerase sigma

factor (15), was used as the unsecreted protein control. BteA forms SDS-resistant

multimers (10). NS: nonspecific signals. Loaded WCL and Sup samples were

prepared from an equal volume of bacterial culture. Experiments were performed at

least three times, and representative data are shown.

Fig. 2. Effects of ascorbic acid and casamino acids on Bvg-regulated virulence

factors. Tohama I was cultured in SS medium or ascorbic acid-deprived SS medium

(SS_AsA$^-$), casamino acid-deprived SS medium (SS_CaA$^-$) or both ascorbic acid-

and casamino acid-deprived SS medium (SS AsA$^-$_CaA$^-$). WCL and Sup samples

were separated by SDS-PAGE and analyzed by western blotting with antibodies

against BteA, BopB, FHA, or CyaA. NS: nonspecific signals. Loaded WCL and Sup

samples were prepared from an equal volume of bacterial culture. Experiments were

performed at least three times, and representative data are shown.

Fig. 3. Effects of ascorbic acid on type III secreted proteins. Whole cell lysates and

secreted proteins prepared from Tohama I cultured in SS medium in the presence of

ascorbic acid at the final concentration of 91, 454 or 2270 μM were separated by

SDS-PAGE and then analyzed by western blotting with antibodies against BteA or

BopB. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from

an equal volume of bacterial culture. Experiments were performed at least three

times, and representative data are shown.

Fig. 4. Effects of ascorbic acid on the secretion of BteA and BopB in clinical isolates

cultured in SS_AsA$^-$ medium. The *B. pertussis* clinical isolates BP344, BP348,

BP350, and BP351 were cultured in SS (+) or SS_AsA$^-$ (–) medium. WCL and Sup

samples were separated by SDS-PAGE and analyzed by western blotting with

antibodies against BteA (*upper panel*) or BopB (*lower panel*). NS: nonspecific signals.

Loaded WCL and Sup samples were prepared from an equal volume of bacterial

culture. Experiments were performed at least three times, and representative data

are shown.

Fig. 5. Results of the RT-PCR analysis for mRNA levels of *bteA* and *bopB* from

B. pertussis cultured under ascorbic acid-starved conditions. Total RNA was

prepared from Tohama I or BP350 cultured in SS or SS_AsA$^-$ medium and

subjected to a quantitative RT-PCR analysis. The histograms show the relative

amounts of *bteA* (**A**) and *bopB* (**B**) mRNA normalized by the housekeeping gene,

recA mRNA. Error bars: the standard error of the mean (SEM) from triplicate

experiments. *$p<0.05$; N.S.: no significant difference. Experiments were performed at

least three times, and representative data are shown.

Fig. 6. Cytotoxic activity of *B. pertussis* cultured in SS_AsA⁻ medium. L2 cells were

infected with Tohama I cultured in SS medium, SS_AsA⁻ medium, SS medium

containing one-twentieth of ascorbic acid (SS_1/20AsA), SS medium containing one-

twentieth of ascorbic acid and one-half of casamino acids (SS_1/20AsA_1/2CaA), or

SS medium containing one-twentieth of ascorbic acid and no casamino acids

(SS_1/20AsA_CaA⁻) at an MOI of 500 for 3 hr. +, 1/2 and 1/20 show the amounts of

indicated substances that are equal to, one-half of, and one-twentieth of those in SS

medium, respectively. The amounts of LDH released into the extracellular medium

from infected cells are shown, and the relative cytotoxicity (%) was determined as

described in the Materials and Methods section. Error bars: the SEM from triplicate

experiments. n.a.: No available result. Experiments were performed at least three

times, and representative data are shown.

Fig. 7. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. pertussis*.

A: L2 cells were infected with Tohama I or BP350 in the indicated media at an MOI

of 125 for 3 hr. +, 1/2 and 1/20 indicate the amounts of indicated substances that are

equal to, one-half of, and one-twentieth of those in SS medium, respectively. After

fixation, cells were stained with anti-Bsp22 antibody (*green*) and rhodamine

phalloidin (*red*). *Arrowheads* indicate the Bsp22 (*green*) signal. **B:** Bsp22 signals per

one cell were counted under a fluorescent microscope. At least 120 cells were

randomly chosen. Error bars: the SEM from triplicate experiments. *$p<0.05$; N.S.: no

significant difference. Experiments were performed at least three times, and

representative data are shown.

Fig. 8. Cytotoxic activities of *B. pertussis* mutants. The results of LDH assays at an

MOI of 125 (**A**) or 250 (**B**) are shown as a histogram. Error bars: the SEM from

triplicate experiments. *$p<0.05$. Experiments were performed at least three times,

and representative data are shown. **C:** MGKU1 (Tohama I nalidixic acid-resistant

strain, 1: WT Nal^r), MGKU2 (T3SS-inactive strain, 2: ΔT3SS), MGKU3 (BteA-

deficient strain, 3: ΔBteA), or MGKU4 (BspR-deficient strain, 4: ΔBspR) was cultured

in SS_1/20AsA_1/2CaA medium. WCL and Sup samples were separated by SDS-

PAGE and analyzed by western blotting with anti-BteA antibody. We loaded a 3-fold

smaller amount of MGKU4 supernatant fraction on the SDS-PAGE gel to avoid

obtaining an excess signal intensity. NS: nonspecific signals. Loaded WCL and Sup

samples were prepared from an equal volume of bacterial culture. Experiments were

performed at least three times, and representative data are shown.

Fig. 9. Effects of reducing agents on the production and secretion of type III secreted

proteins in *B. pertussis*. Tohama I was cultured in SS_AsA$^-$ medium or SS_AsA$^-$

medium containing ascorbic acid (AsA), 2-mercaptoethanol (2-ME), or dithioerythritol

(DTE) at final concentrations of 2.3, 2.1, or 3.0 mM, respectively. WCL and Sup

samples were separated by SDS-PAGE, then analyzed by western blotting with

antibodies against BteA, BopB, or FHA. Loaded WCL and Sup samples were

prepared from an equal volume of bacterial culture. NS: nonspecific signals.

Experiments were performed at least three times, and representative data are shown.

Table 1. Bacterial strains and plasmids used in study

Strains or plasmids	Relevant genotype/description	Reference
Strains		
B. pertussis		
Tohama I	Clinical isolate	22
BP344	Clinical isolate	23
BP348	Clinical isolate	23
BP350	Clinical isolate	23
BP351	Clinical isolate	23
MGKU1	Tohama I nalidixic-acid resistant strain	This study
MGKU2	T3SS-inactive strain	This study
MGKU3	BteA-deficient strain	This study
MGKU4	BspR-deficient strain	This study
E. coli		
DH10B	Host strain for pDONR201	invitrogen
Sm10λ*pir*	Host strain for pABB-CRS2	18
Plasmids		
pDONR201	DNA cloning vector, Km^r	invitrogen
pABB-CRS2	Suicide vector for conjugation, Amp^r	27
pABB-Δ*bscN*	pABB-CRS2, Δ*bscN* gene	18
pABB-Δ*bopC*	pABB-CRS2, Δ*bopC* (Δ*bteA*) gene	10
pMGKU401	pDONR201, *bspR* gene	this study
pMGKU402	pDONR201, Δ*bspR* gene	this study
pMGKU403	pABB-CRS2, Δ*bspR* gene	this study

Table 2. Medium compositions

Basal medium (g/L)	SS	LCA
Sodium Gultamate	10.7	10.7
L-proline	0.24	0.24
NaCl	2.5	2.5
KCl	0.2	0.2
KH_2PO_4	0.5	0.5
$MgCl_2 \cdot 6H_2O$	0.1	0.1
$CaCl_2$	0.02	0.02
Tris	6.1	6.1
Casamino acids	10	0.5
Dimethyl-β-cyclodextrin	1	1

Supplement (g/L)	SS	LCA
L-cysteine•HCl	4	4
$FeSO_4 \cdot 7H_2O$	1	1
L-ascorbic acid	40	2
Niacin	0.4	0.4
L-glutathione reduced	15	15

Table 3. Primers used in this study.

Name	Sequence
B1-bspR	5'-AAAAAGCAGGCTCGCCGGTTTCCACCACGTAA-3'
B2-bspR	5'-AGAAAGCTGGGTGGCCAATATCCTGTCTCAGC-3'
R1-bspR	5'-CGC<u>GGATCC</u>CATACGGCGCAAGAATGGCT-3'
R2-bspR	5'-CGC<u>GGATCC</u>AAAGACATCAAGCTCCAAAG-3'
5-recA	5'-ATGAAGATCGGCGTGATGT-3'
3-recA	5'-TAGAACTTGAGCGCGTTGC-3'
5-bteA	5'-AATGGCCTTGGTGGGAAC-3'
3-bteA	5'-ATTTCAGCGCCGTGATCTT-3'
5-bopB	5'-GTCTGCTGACAGCGTTGG-3'
3-bopB	5'-GGGGTTCGATCAGCAAGAC-3'

Restriction enzyme recognition sites were underlined.

Fig. 1. BteA and BopB production by *B. pertussis* Tohama I cultured in LCA medium.

Tohama I and MGKU2 (a T3SS-inactive strain) were cultured in SS or LCA medium. Whole

cell lysates (WCL) and supernatant fraction (Sup) samples were separated by SDS-PAGE

and analyzed by western blotting with antibodies against BteA (*upper panel*), BopB (*middle*

panel) or BtrS (*lower panel*). BtrS, an RNA polymerase sigma factor (15), was used as the

unsecreted protein control. BteA forms SDS-resistant multimers (10). NS: nonspecific signals

Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture.

Experiments were performed at least three times, and representative data are shown.

Fig. 2. Effects of ascorbic acid and casamino acids on Bvg-regulated virulence factors.

Tohama I was cultured in SS medium or ascorbic acid-deprived SS medium (SS_AsA⁻),

casamino acid-deprived SS medium (SS_CaA⁻) or both ascorbic acid- and casamino

acid-deprived SS medium (SS AsA⁻_CaA⁻). WCL and Sup samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA, BopB, FHA, or

CyaA. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal

volume of bacterial culture. Experiments were performed at least three times, and

representative data are shown.

Fig. 3. Effects of ascorbic acid on type III secreted proteins. Whole cell lysates and secreted proteins prepared from Tohama I cultured in SS medium in the presence of ascorbic acid at the final concentration of 91, 454 or 2270 μM were separated by SDS-PAGE and then analyzed by western blotting with antibodies against BteA or BopB. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 4. Effects of ascorbic acid on the secretion of BteA and BopB in clinical isolates cultured

in SS_AsA⁻ medium. The *B. pertussis* clinical isolates BP344, BP348, BP350, and BP351

were cultured in SS (+) or SS_AsA⁻ (−) medium. WCL and Sup samples were separated by

SDS-PAGE and analyzed by western blotting with antibodies against BteA (*upper panel*) or

BopB (*lower panel*). NS: nonspecific signals. Loaded WCL and Sup samples were prepared

from an equal volume of bacterial culture. Experiments were performed at least three times,

and representative data are shown.

Fig. 5. Results of the RT-PCR analysis for mRNA levels of *bteA* and *bopB* from *B. pertussis*

cultured under ascorbic acid-starved conditions. Total RNA was prepared from Tohama I or

BP350 cultured in SS or SS_AsA⁻ medium and subjected to a quantitative RT-PCR analysis.

The histograms show the relative amounts of *bteA* (**A**) and *bopB* (**B**) mRNA normalized by

the housekeeping gene, *recA* mRNA. Error bars: the standard error of the mean (SEM) from

triplicate experiments. *p<0.05; N.S.: no significant difference. Experiments were performed

at least three times, and representative data are shown.

Fig. 6. Cytotoxic activity of *B. pertussis* cultured in SS_AsA⁻ medium. L2 cells were infected

with Tohama I cultured in SS medium, SS_AsA⁻ medium, SS medium containing

one-twentieth of ascorbic acid (SS_1/20AsA), SS medium containing one-twentieth of

ascorbic acid and one-half of casamino acids (SS_1/20AsA_1/2CaA), or SS medium

containing one-twentieth of ascorbic acid and no casamino acids (SS_1/20AsA_CaA⁻) at an

MOI of 500 for 3 hr. +, 1/2 and 1/20 show the amounts of indicated substances that are equal

to, one-half of, and one-twentieth of those in SS medium, respectively. The amounts of LDH

released into the extracellular medium from infected cells are shown, and the relative

cytotoxicity (%) was determined as described in the Materials and Methods section. Error

bars: the SEM from triplicate experiments. n.a.: No available result. Experiments were

performed at least three times, and representative data are shown.

Fig. 7. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. pertussis*. **A:** L2 cells

were infected with Tohama I or BP350 in the indicated media at an MOI of 125 for 3 hr. +, 1/2

and 1/20 indicate the amounts of indicated substances that are equal to, one-half of, and

one-twentieth of those in SS medium, respectively. After fixation, cells were stained with

anti-Bsp22 antibody (*green*) and rhodamine phalloidin (*red*). *Arrowheads* indicate the Bsp22

(*green*) signal. **B:** Bsp22 signals per one cell were counted under a fluorescent microscope.

At least 120 cells were randomly chosen. Error bars: the SEM from triplicate experiments.

*p<0.05; N.S.: no significant difference. Experiments were performed at least three times, and

representative data are shown.

A

MOI of 125

B

MOI of 250

C

Fig. 8. Cytotoxic activities of *B. pertussis* mutants. The results of LDH assays at an MOI of 125 (**A**) or 250 (**B**) are shown as a histogram. Error bars: the SEM from triplicate experiments *p<0.05. Experiments were performed at least three times, and representative data are shown. **C:** MGKU1 (Tohama I nalidixic acid-resistant strain, 1: WT Nal^r), MGKU2 (T3SS-inactive strain, 2: ΔT3SS), MGKU3 (BteA-deficient strain, 3: ΔBteA), or MGKU4 (BspR-deficient strain, 4: ΔBspR) was cultured in SS_1/20AsA_1/2CaA medium. WCL and Sup samples were separated by SDS-PAGE and analyzed by western blotting with anti-BteA antibody. We loaded a 3-fold smaller amount of MGKU4 supernatant fraction on the SDS-PAGE gel to avoid obtaining an excess signal intensity. NS: nonspecific signals. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 9. Effects of reducing agents on the production and secretion of type III secreted proteins in *B. pertussis*. Tohama I was cultured in SS_AsA⁻ medium or SS_AsA⁻ medium containing ascorbic acid (AsA), 2-mercaptoethanol (2-ME), or dithioerythritol (DTE) at final concentrations of 2.3, 2.1, or 3.0 mM, respectively. WCL and Sup samples were separated by SDS-PAGE, then analyzed by western-blotting with antibodies against BteA, BopB, or FHA. Loaded WCL and Sup samples were prepared from an equal volume of bacterial culture. NS: nonspecific signals. Experiments were performed at least three times, and representative data are shown.

Fig. S1. Cytotoxic activity of *B. pertussis* Tohama I in F-12K medium containing oxidizing agents. L2 cells were infected with *B. pertussis* Tohama I in the modified SS medium (SS_1/20AsA_1/2CaA, MSS). The amounts of ascorbic acid and casamino acids in MSS are one-twentieth of, and one-half of those in SS medium, respectively. L2 cells were also infected in F-12K medium (none), F-12K medium containing hydrogen peroxide (H_2O_2), potassium ferricyanide(III) ($K_3[Fe(CN)_6]$), methyl 3-nitro-2-pyridinesulfenate (Npys-OMe) or 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB). The final concentrations of each oxidizing agent were 1 mM, 1 mM, 10 μM, or 2 mM, respectively. Minimum Essential Medium Eagle (MEM) was also used. The cells were infected at an MOI of 500 for 3 hr. The amounts of LDH released into the extracellular medium from infected cells are shown, and the relative cytotoxicity (%) was determined as described in the Materials and Methods section. Error bars are the SEM from triplicate experiments. Experiments were performed at least three times, and representative data are shown.