Tendon and Motor Phenotypes in the *Crtap[-/-]* Mouse Model of Recessive Osteogenesis

Imperfecta

Matthew W. Grol[1†], Nele A. Haelterman[1], Joohyun Lim[1], Elda M. Munivez[1], Marilyn Archer[2],

David M. Hudson[2], Sara F. Tufa[3], Douglas R. Keene[3], Kevin Lei[1], Dongsu Park[1], David R. Eyre[2,]

and Brendan H. Lee[1†]

[1] Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX

[2] Department of Orthopaedics and Sports Medicine, University of Washington, Seattle, WA

[3] Shriners Hospital for Children, Portland, OR

[†] Correspondence and reprint requests:

Brendan H. Lee, Department of Human and Molecular Genetics, One Baylor Plaza, R814, Baylor

College of Medicine, Houston, TX, USA, 77030, Email: blee@bcm.edu, Tel: +1 713-798-7246,

Fax: +1 713-798-4523; Matthew W. Grol, Department of Physiology and Pharmacology, Dental

Sciences Building, Room 0079, University of Western Ontario, London, ON, Canada, N6A 5C1,

Email: mgrol2@uwo.ca, Tel: +1 519-661-2111 ext. 81060

Running Title: Tendon and motor deficits in Osteogenesis Imperfecta

Keywords: osteogenesis imperfecta, tendon, ligament, cross-link, motor behavior

ABSTRACT

Osteogenesis imperfecta (OI) is characterized by short stature, skeletal deformities, low bone

mass with bone fragility, and motor deficits. A subset of OI patients also present with joint

hypermobility; however, the role of tendon/ligament dysfunction in OI pathogenesis is largely

unknown. Using the $Crtap^{-/-}$ mouse model of severe, recessive OI, we found that mutant Achilles

tendons and patellar ligaments were thinner with increased collagen cross-links and reduced

collagen fibril size at 1- and 4-months compared to wildtype. Patellar ligaments from $Crtap^{-/-}$

mice also had fewer progenitors with a concomitant increase in immature cells. RNA-seq

analysis of Achilles tendons and patellar ligaments from 1-month $Crtap^{-/-}$ mice revealed

dysregulation in matrix gene expression concomitant with predicted alterations in TGF-β,

inflammatory, and metabolic signaling. Finally, a series of behavioral tests revealed severe motor

impairments and reduced grip strength in 4-month $Crtap^{-/-}$ mice – a phenotype that correlates

with the tendon/ligament pathology.

INTRODUCTION

Tendon is a fibrous tissue that connects skeletal muscle to bone to facilitate motion, whereas ligaments connect articulating bones to support joint alignment and function[1, 2]. The extracellular matrix of tendons/ligaments is primarily composed of type I collagen as well as smaller quantities of other collagens and proteoglycans[3]. During development, the collagen fibrils in tendons and ligaments develop through addition and lengthening before transitioning to the appositional fusion of existing fibers with continued lengthening in postnatal life[4]. The synthesis and assembly of this collagen-rich matrix are influenced by other small collagens and proteoglycans as well as by the cross-linking chemistry of type I procollagen fibrils, which in turn regulates fibril size and strength[5]. Like tendon and ligament, the organic matrix of bone consists largely of type I collagen[6], and disruptions in collagen synthesis and folding are known to negatively impact its biochemical and mechanical properties in connective tissue diseases such as Osteogenesis Imperfecta (OI)[7]. However, despite evidence of joint mobility phenotypes and motor deficits in OI patients[8, 9], tendon and ligament phenotypes in this disease are relatively understudied.

OI is a heterogeneous group of disorders characterized by variable short stature, skeletal deformities, low bone mass, and increased bone fragility. Approximately 80% of OI cases are caused by dominantly inherited mutations in the genes encoding the $\alpha 1(I)$ or $\alpha 2(I)$ chains of type I collagen. Mutations in genes responsible for the synthesis, post-translational modification, and processing of collagen, such as cartilage-associated protein (CRTAP), lead to severe, recessive forms of this disease[7]. In addition to skeletal defects, other connective tissue manifestations including joint hypermobility and skin hyperlaxity are observed in a subset of OI patients[8, 9]. Our and others' studies have shown that CRTAP forms a complex with Prolyl 3-hydroxylase 1

(P3h1, encoded by *Lepre1*) and Cyclophilin B (CypB, encoded by *Ppib*), and is required for prolyl 3-hydroxylation of type I procollagen at Pro986 of chain α1(I) and Pro707 of chain α2(I)[10-12]. In this regard, loss of either CRTAP or P3H1 leads to loss of this complex and its activity, causing a severe recessive form of OI characterized by short stature and brittle bones[11, 13-15]. Collagen isolated from *Crtap*[-/-] and *Lepre1*[-/-] mice is characterized by lysine over-modifications and abnormal fibril diameter[11, 14]. While a comprehensive analysis of *Crtap*[-/-] mice has revealed multiple connective tissue abnormalities, including in lungs, kidneys, and skin[16], the impact of the loss of Crtap on tendons and ligaments remains unknown.

Alterations in collagen fibril size and cross-linking have been noted in a limited number of studies using dominant or recessive mouse models of OI[17-19]; however, whether loss of Crtap impacts tendon and ligament development and structure remains unknown. In this study, we show that *Crtap*[-/-] mice have thinner Achilles tendons and patellar ligaments at 1 and 4 months-of-age that are hypercellular with a reduction in tendon progenitors and total ligament volume. Moreover, the patellar and cruciate ligaments from *Crtap*[-/-] show increased cell size and variable ectopic chondrogenesis by 4 months-of-age. Examining the collagen matrix, we found an increase in stable (irreversible) collagen cross-links at both timepoints, accompanied by alterations in fibril diameter at 4-months compared to wildtype controls. RNA-seq analyses revealed both common and distinct changes in the transcriptome of the Achilles tendon and patellar ligaments of *Crtap*[-/-] mice compared to wildtype with a predicted activation of transforming growth factor-β (TGF-β) and inflammatory signaling in both tissues. These changes in *Crtap*[-/-] tendons and ligaments were accompanied by motor deficits and reduced strength at 4 months-of-age. In conclusion, loss of CRTAP in mice causes tendon/ligament abnormalities and significant behavioral impairments.

MATERIALS AND METHODS

Animals

Crtap⁻/⁻ mice were generated as previously described[11] and maintained on a mixed C57BL/6J

and 129Sv genetic background. All studies were performed with approval from the Institutional

Animal Care and Use Committee (IACUC) at Baylor College of Medicine. Mice were housed 3

to 4 mice to a cage in a pathogen-free environment with *ad libitum* access to food and water and

under a 14h light/10h dark cycle.

Histological analysis

Mice were euthanized and ankle and knee joints were dissected and fixed for 48 h on a shaker at

4ºC or room temperature in freshly prepared 4% paraformaldehyde (PFA) in 1×phosphate-

buffered saline (PBS). Samples were decalcified at 4°C using 10% ethylenediaminetetraacetic

acid (EDTA) in 1×PBS for 10 days (with one change out at 5 days) before paraffin embedding

using a standard protocol. Samples were sectioned at 6-μm and stained with hematoxylin and

eosin (H&E) to visualize tendon structures. Cell number per tissue area was determined using the

Fiji release of ImageJ[20].

Fluorescence-activated cell sorting analysis of tendon progenitors

After dissection of patellar ligaments from 5-month-old wildtype and *Crtap⁻/⁻* mice, tissues were

cut into small pieces in 1×PBS with 10% fetal bovine serum (FBS) and incubated with 500 μl of

PBS + 10% FBS and 0.1% collagenase at 37°C for 3 hours. After digestion, cells were filtered

with a 40-μm strainer, washed, resuspended in 1×PBS at a concentration of 10^6 cells/mL, and

stained with CD45-pacific blue (clone: 30-F11), CD31-eFlour 450 (clone: 390), CD146-PE-Cy7

(clone ME-9F1) and CD200-APC (clone OX-90) (eBioscience). Propidium iodide was used for

selecting viable cells. Cell analysis was performed using a LSRII Fortessa, and FACS

experiments were done using an AriaII cytometer (BD Biosciences, San Jose, CA). Data were

analyzed with FlowJo software (TreeStar, OR).

Phase-contrast µCT imaging and analysis

To quantify tendon/ligament volume, knee joints were dissected from mice, stained with contrast

agents, scanned by phase-contrast µCT, and analyzed using TriBON software (RATOC, Tokyo,

Japan) as previously described for articular cartilage[21-24]. In addition to the articular surfaces

(data not shown), we performed contrast-enhanced visualization of the patellar ligament using

this protocol. To quantify volume, samples were examined in transverse where the patellar

ligament boundary was easily distinguished from the joint capsule. Ligament volume was

assessed from its origin within the patella to its insertion at the tibia.

Transmission electron microscopy analysis of collagen fibril size

Mouse ankle and knee joints were dissected and fixed in fresh 1.5% glutaraldehyde/1.5% PFA

(Tousimis) with 0.05% tannic acid (Sigma) in 1×PBS at 4°C overnight to preserve the native

tension on relevant tendons/ligaments. The next day, flexor digitorum longus (FDL) and Achilles

tendons as well as patellar ligament were dissected out in 1×PBS, and placed back into fixative.

Samples were then post-fixed in 1% osmium tetroxide (OsO_4), rinsed in Dulbecco's Modified

Eagle Medium (DMEM), and dehydrated in a graded series of ethanol to 100%. Samples were

rinsed in propylene oxide, infiltrated in Spurrs epoxy, and polymerized at 70°C overnight. TEM

images were acquired using an FEI G20 TEM at multiple magnifications to visualize transverse

sections of collagen fibrils. Collagen fibril diameter was measured using the Fiji release of

ImageJ[20].

Tendon collagen cross-linking analysis

Collagen hydroxylysyl-pyridinoline (HP) cross-links were quantified as previously described[25,

26]. In brief, tendons and ligaments isolated from hindlimbs were hydrolyzed by 6M HCl for 24h

at 108°C. Dried samples were then dissolved in 1% (v/v) n-heptafluorobutyric acid for

quantitation of HP by reverse-phase HPLC with fluorescence monitoring.

RNA-seq analysis

At 1 month-of-age, Achilles tendons were excised using scissors proximal to the calcaneal

insertion and distal to the tendon-muscular-junction, whereas patellar ligaments were removed

via scalpel just proximal to the tibial insertion and distal to the patella. Tendons and ligaments

were not cleaned of their paratenon layers in either case. RNA extraction for both tissues was

performed with the Fibrous Connective Tissue kit (Qiagen), with columns from the RNeasy

Micro Kit (Qiagen) employed for the patellar ligament. Quality control, library preparation,

sequencing, and differential gene expression analysis including gene ontology analysis was

performed by GENEWIZ® (South Plainfield, NJ). For the examination of predicted Upstream

Regulators, we utilized the Ingenuity Pathway Analysis (IPA) platform (Qiagen) with an

adjusted *p*-value of 0.05. The expression is shown as Base Means and read counts of genes were

normalized per million transcripts (Transcripts Per Million; TPM).

Open field assessment of spontaneous motor activity

Open field activity was measured using the VersaMax Animal Activity Monitoring System (AccuScan Instruments, Columbus, OH). On the day of assessment, mice were transferred to the test room and allowed to acclimate in their home cage for 30 min at 50 Lux of illumination with 60 dB of white noise. Mice were then placed individually into clear 40 cm × 40 cm × 30 cm chambers and allowed to move freely for 30 min. Locomotion parameters and zones were recorded using the VersaMax activity monitoring software. Chambers were cleaned with 30-50% ethanol to remove the scent of previously tested mice between each run.

Rotarod analysis of motor coordination and endurance

On the day of assessment, mice were transferred to the test room and allowed to acclimate within their home cage for 30 min at 50 Lux of illumination with 60 dB of white noise. Mice were then placed on a rotarod (UGO Basile, Varese, Italy) set to accelerate from 5-to-40 RPM over 5 min. Five trials were performed per day for 2 consecutive days (trials 1-10) with a rest time of 5 min between trials. Latency to fall was recorded when the mouse fell from the rotating rod or went for two revolutions without regaining control. The rotarod was cleaned with 30-50% ethanol between mice to remove the scent of previously tested animals.

Grid foot slip analysis of motor coordination

The grid foot slip assay consisted of a wire grid set atop a stand where the movement was recorded by a suspended digital camera. Mice were transferred to the test room on the day of assessment and allowed to acclimate within their home cage for 30 min at 50 Lux of illumination with 60 dB of white noise. Mice were then placed one at a time on the grid and allowed to move freely for 5 min. The observer sat 6-8 feet away at eye-level to the mouse and recorded forelimb

and hindlimb foot slips using the ANY-maze video tracking software (Stoelting Co., Wood Dale,

IL). After the test, mice were removed to their original home cage. Forelimb and hindlimb foot

slips were normalized to the total distance traveled during the test.

Inverted grid analysis of strength and endurance

On the day of assessment, mice were transferred to the test room and allowed to acclimate within

their home cage for 30 min at 50 Lux of illumination with 60 dB of white noise. Mice were then

placed in the middle of a wire grid, held approximately 18-in above a cushioned pad, and

inverted. The latency to fall for each mouse was recorded. At the completion, mice were returned

to their home cage.

Grip strength analysis

Mice were transferred to the test room on the day of assessment and allowed to acclimate within

their home cage for 30 min at 50 Lux of illumination with 60 dB of white noise. Each mouse was

then lifted by its tail onto the bar of a digital grip strength meter (Columbus Instruments,

Columbus, OH). Once both forepaws had gripped the bar, the mouse was gently pulled away

from the meter by its tail at a constant speed until the forepaws were released. The grip (in N of

force) was recorded and the procedure repeated twice for a total of three measurements, which

were averaged for the final result.

Statistical analysis

Determination of sample size was based on previous publications. Biological replicates were

defined as an individual mouse for each experiment. Respective tendons and ligaments from the

left and right hindlimbs were combined for TEM, collagen cross-linking, tendon progenitor, and

RNA-seq analyses. Data are presented as means ± S.D. or min-to-max box and whisker plots

with individual data points. For data whose residuals passed the Shapiro-Wilk test for normality,

groups of two were compared using unpaired t-tests, and groups of three or more were compared

using one-way ANOVA followed by Tukey's post-hoc tests. For data whose residuals did not

have a normal distribution, groups of two were compared using a Mann-Whitney test, and groups

of three or more were compared using Kruskal-Wallis followed by Dunn's post-hoc tests. For the

Rotarod assay, where time was a variable, groups were compared using repeated measures two-

way ANOVA followed by Tukey's post-hoc tests. For all tests reported above, statistical analysis

was performed using Prism 8.3.1 (GraphPad Software, La Jolla, CA). For all tests, the exact p-

value is reported, and a p-value of < 0.05 was considered statistically significant.

RESULTS

Crtap[-/-] mice exhibit abnormal tendon development

Mice lacking Crtap present with growth delay, rhizomelia, and severe osteoporosis together with

disruption of other connective tissues including lung and skin[11, 16]. To assess whether *Crtap[-/-]*

mice exhibit disruptions in the development of tendons and ligaments, we harvested ankle and

knee joints at 1 and 4 months-of-age to histologically examine the Achilles tendon and patellar

ligament. At 1-month, *Crtap[-/-]* mice presented with thinner Achilles tendons and patellar

ligaments (**Figure 1C,F**) compared to wildtype (**Figure 1A,D**) and heterozygous mice (**Figure**

1B,E) with increased cell density in both structures (**Figure 1M**). There was also prominent

hyperplasia of the synovial membrane in *Crtap* mutant knees at this age (**Figure 1F**). By 4

months-of-age, *Crtap[-/-]* Achilles tendons and patellar ligaments remained thinner and

hypercellular compared to wildtype and heterozygous mice (**Figure 1G-L,N**). In addition,

synovial hyperplasia persisted within *Crtap*[-/-] knees (**Figure 1L**). Interestingly, ectopic

chondrogenesis was present towards either end of the patellar ligament and/or cruciate ligaments

in some (but not all) 4-month-old *Crtap*[-/-] mice (**Figure 1L**) – a phenomenon that can occur in

tendinopathy[27].

To more accurately quantify the differences in tendon/ligament size between genotypes,

we performed phase-contrast μCT to examine differences in patellar ligament volume between

genotypes (**Figure 1O-Q**) – a technique we previously used to examine soft tissues including

articular cartilage in intact murine knee joints[21-24]. In 4-month-old mice, we observed a thinning

of the patellar ligament compared to wildtype mice, with no notable changes in articular cartilage

surface (**Figure 1O,P**). Confirming our histological findings, quantification revealed a decrease

in patellar ligament volume in *Crtap*[-/-], but not in heterozygous mice, compared to wildtype

controls (**Figure 1Q**).

Postnatal progenitor populations are thought to be critical for the maintenance and

regeneration of tendon/ligament tissue. Previous studies have shown that a subset of fibroblasts

in the tendon midsubstance (endotenon) has the potential to contribute to tendon homeostasis and

regeneration at various developmental stages[28-30]. In addition, platelet-derived growth factor

receptor α (PDGFRa)[+] cells in the tendon sheath (paratenon) can contribute to tendon

regeneration after acute injury[31]. These progenitor cells can be labeled by *Tppp3*-CreER[31] or

aSMA-CreER[28] with the expression of tendon progenitor markers such as CD146[32], CD140a

(PDGFRa)[31], and CD200[33]. Given the significant hypercellularity seen in the Achilles tendons

and patellar ligaments from *Crtap*[-/-] mice, we examined whether the postnatal progenitor

population is altered compared to wildtype controls. Using fluorescence-activated cell sorting

(FACS) analysis of 5-month-old patellar ligaments, we observed a significant decrease in the

percentage of CD45$^-$CD31$^-$CD146$^+$CD200$^+$ stem/progenitor cells (~2%) compared to wild-type

mice (~4%) (**Figure 2A-C**, red box), while CD45$^-$CD31$^-$CD146$^-$CD200$^+$ immature cells were

concomitantly increased in *Crtap$^{-/-}$* mice (**Figure 2D**).

Taken together, these data demonstrate that *Crtap$^{-/-}$* mice display disruptions in the

development and postnatal maturation of tendons and ligaments. Moreover, the matrix

disruptions caused by loss of CRTAP lead to the dysregulation of the postnatal tendon progenitor

pool as well as tenocyte maturation.

Collagen fibril formation is altered in heterozygous and Crtap$^{-/-}$ mice

Tendons develop embryonically by increasing fibril length and number, whereas postnatal

growth arises from an increase in fibril length and diameter – the latter of which is driven by the

lateral fusion of smaller fibrils[4]. To investigate the role of Crtap in collagen fibril maturation, we

utilized transmission electron microscopy (TEM) to examine changes in fibril diameter in flexor

digitorum longus (FDL) and Achilles tendons as well as patellar ligaments (**Figure 3**). In the

FDL tendon, there was a marked increase in small collagen fibrils (20-60 nm in size), a reduction

in 80-320 nm fibrils, and a slight increase in larger fibrils (>340 nm in diameter) in *Crtap$^{-/-}$* mice

compared to wildtype (**Figure 3A,C,D**). Despite similarities seen in histology, heterozygous

mutant FDL tendons also exhibited a slight increase in 20-40 nm fibrils in mice compared to

wildtype controls (**Figure 3A-B,D**). Similar trends were observed for the Achilles tendon,

namely an increase in small fibrils (20-60 nm), a reduction in 80-240 nm fibrils, and an increase

in large fibrils (>280 nm) upon loss of *Crtap* (**Figure 3E,G-H**). In contrast to what we observed

for the FDL, heterozygous Achilles tendons did not have increased numbers of smaller fibers

(**Figure 3F,H**). Instead, a greater number of fibrils ranging from 140-200 nm in size were noted

compared to wildtype controls.

Compared to the FDL and Achilles tendons, the greatest differences were seen within the

patellar ligament, though the pattern of changes remained consistent (**Figure 3I-L**). Specifically,

we observed a dramatic increase in 20 nm collagen fibrils compared to both heterozygous and

wildtype mice (**Figure 3I-L**). Fibrils ranging from 100-180 nm in diameter were reduced in

heterozygous and *Crtap*[-/-] mice compared to wildtype. Interestingly, the greatest difference from

wildtype was an increase in large collagen fibrils (>200 nm) in both heterozygous and *Crtap*[-/-]

mice (**Figure 3I-L**). Taken together, these data indicate that heterozygous and complete loss of

CRTAP alter collagen fibril assembly in tendons and ligaments. In addition, the degree to which

collagen assembly is affected is site-dependent.

Collagen cross-linking is increased in heterozygous and Crtap[-/-] *mice*

Along with P3H1 and CyPB, CRTAP is an integral part of the collagen prolyl 3-hydroxylation

complex that is responsible for the 3-hydroxylation of Pro986 of the type I procollagen α1

chain[7]. Loss of this complex blocks 3-hydroxyproline formation and affects lysine hydroxylation

and cross-linking in bone collagen[11, 12]; however, whether *Crtap*[-/-] tendons and ligaments display

altered collagen cross-linking is unknown. To investigate this, we harvested tendons and

ligaments at 1- and 4-months and assessed collagen cross-linking by quantifying the levels of

hydroxylysyl-pyridinoline (HP) (**Figure 4**). Overall, we observed an increase in these stable,

mature collagen cross-links from 1 to 4 months-of-age in all genotypes for the FDL and Achilles

tendons (**Figure 4A-B**). In contrast, for the patellar ligament, age-dependent increases in

collagen cross-links were only observed in *Crtap*[-/-] mice (**Figure 4C**). For FDL tendons, *Crtap*[-/-]

mice had more of these collagen cross-links at 1- and 4-months compared to heterozygous and

wildtype mice; however, the content of HP residues per collagen decreased with age in this tissue

(**Figure 4A**). Interestingly, in Achilles tendons, an increase in collagen cross-linking was

observed in both heterozygous and *Crtap*[-/-] mice at 1-month compared to wildtype. In contrast, at

4-months, only *Crtap*[-/-] mice had elevated collagen cross-links, and these levels were greater than

those observed at the earlier time point (**Figure 4B**).

Of the tissues examined, the patellar ligament showed the greatest increase in collagen

cross-links both with time and across genotypes. Specifically, collagen cross-links were elevated

by 5- to 10-fold in *Crtap*[-/-] patellar ligaments compared to heterozygous and wildtype at 1- and

4-months, respectively (**Figure 4C**). Taken together, these data suggest that *Crtap* is required for

proper hydroxylation and cross-linking of collagen fibrils in tendons and ligaments in a semi-

dominant fashion, as heterozygous mutant tendons/ligaments display a phenotype that is milder

than the phenotype observed for homozygous mutant mice. Importantly, the chemical quality of

collagen cross-linking appears to be spatiotemporally regulated and this regulation is

differentially affected by the loss of a single or both copies of *Crtap*.

Signaling and metabolic dysregulation in Crtap[-/-] *tendons and ligaments*

To investigate the molecular changes underlying tendon/ligament phenotypes, we performed

bulk RNA-seq with RNA isolated from Achilles tendons and patellar ligaments of 1-month-old

wildtype and *Crtap*[-/-] mice. To determine global changes in differentially expressed genes as well

as predicted upstream regulators, we performed Ingenuity Pathway Analysis (IPA, Qiagen,

Germany). For the Achilles tendon, a total of 178 genes (consisting of 99 upregulated genes and

79 downregulated genes) were significantly differentially expressed between wildtype and *Crtap*[-]

$^{-/-}$ samples (**Figure 5A**). Of the top 30 differentially expressed genes, several extracellular matrix

proteins, including matrilin-3 (*Matn3*), matrilin-4 (*Matn4*), and fibronectin 1 (*Fn1*), and

proteolytic enzymes such as matrix metallopeptidase 2 (*Mmp2*) were dysregulated. Gene

ontology analysis revealed "GO:000715 – Cell Adhesion", "GO:0045778 – Positive Regulation

of Ossification", and "GO:0051928 – Positive Regulation of Calcium Ion Transport" to be

enriched (**Figure 5B**). Examination of upstream regulators based on the differential gene

expression data also revealed a predicted activation of TGF-β1 in *Crtap*$^{-/-}$ mice and predicted

inhibition of dystrophin (DMD) along with several for which activation state was unclear,

including platelet-derived growth factor-BB (PDGF-BB), β-catenin (CTNNB1), and tumor

necrosis factor (TNF) (**Figure 5C**).

333   In keeping with the increased severity seen in ligaments from *Crtap*$^{-/-}$ mice compared to

tendons, a greater number of total genes were differentially expressed between wildtype and

Crtap$^{-/-}$ patellar ligament samples (**Figure 5D**). Specifically, we saw significant differential

expression of 519 genes, with 224 being upregulated and 295 downregulated in *Crtap*$^{-/-}$

compared to wildtype. Several of the top 30 differentially expressed genes were small collagens

such as type IX collagen α3 chain (*Col9a3*), type IX collagen α1 chain (*Col9a1*), and type XXII

collagen α1 chain (*Col22a1*) as well as other extracellular matrix proteins, including lumican

(Lum) and fibronectin 1 (*Fn1*). Unlike the Achilles tendon, gene ontology analysis revealed

significant enrichment for metabolic processes, including "GO:0055114 – Oxidation-Reduction

Process", "GO:0050873 – Brown Fat Cell Differentiation", and "GO:0050872 – White Fat Cell

Differentiation" as well as for "GO:0008284 – Positive Regulation of Proliferation" and

"GO:0001525 – Angiogenesis" (**Figure 5E**). Examination of upstream regulators also suggested

a highly significant activation of TNF and interleukin 6 (IL-6) as well as fibroblast growth factor

2 (FGF2) and TGF-β3 in *Crtap⁻ᐟ⁻* patellar ligaments (**Figure 5F**). Indeed, the representation of

TNF, TGF-β, and PDGF-BB in the ligament dataset is consistent with but more severe than that

observed for those same regulators in the Achilles tendon results (**Figure 5C**).

Loss of CRTAP leads to deficiencies in motor activity, coordination, and strength

To investigate whether the tendon phenotypes observed in *Crtap⁻ᐟ⁻* mice have functional

consequences, we performed a series of behavioral assays at 4 months-of-age. Using the open

field assay to quantify changes in spontaneous motor activity, we observed that *Crtap⁻ᐟ⁻* mice

displayed significant reductions in both horizontal and vertical activity compared to

heterozygous and wildtype mice (**Figure 6A-B**). We next examined changes in motor

coordination and endurance using the rotarod assay. While no genotype-dependent differences

were observed during the learning phase of the assessment (Trials 1-5), *Crtap⁻ᐟ⁻* displayed a

reduction in latency to fall for Trials 6, 9-10 compared to wildtype and Trials 6, 8-10 compared

to heterozygous mice (**Figure 6C**). To confirm this observation, we evaluated the mice using the

grid foot slip assay – an alternative metric for motor coordination. In this regard, we found that

Crtap⁻ᐟ⁻ mice exhibited a modest increase in forelimb and hindlimb foot slips compared to

heterozygous and wildtype mice (**Figure 6D-E**). Taken together, these findings indicate that

Crtap⁻ᐟ⁻ mice have deficiencies in motor activity and coordination compared to controls.

We next evaluated strength in the *Crtap⁻ᐟ⁻* mice using the inverted grid and grip strength

assays. Interestingly, we observed a decrease in the latency to fall during the inverted grid assay

for *Crtap⁻ᐟ⁻* mice compared to heterozygous and wildtype controls (**Figure 6F**). Using a more

quantitative metric, we examined these mice using the grip strength test and found that while

wildtype and heterozygous mice could generate approximately 1.3 N of force, mice lacking

CRTAP were weaker with a mean grip strength of 0.62 N (**Figure 6G**). Thus, *Crtap*$^{-/-}$ mice

display significant reductions in strength together with perturbations in motor activity and

coordination – behavioral changes that could be related in part to their tendon phenotype.

DISCUSSION

In this study, we examined the histological, ultrastructural, biochemical, and transcriptional

characteristics of tendons and ligaments in the *Crtap*$^{-/-}$ mouse model of severe, recessive OI. We

demonstrate that at 1- and 4 months-of-age, *Crtap*$^{-/-}$ have thinner, more cellular Achilles tendons

and patellar ligaments with a reduction in tendon progenitors compared to heterozygous and

wildtype mice. Using phase-contrast µCT imaging, we confirmed that tissue volume is reduced

in *Crtap*$^{-/-}$ patellar ligaments at 4-months than those from heterozygous and wildtype mice.

Examining collagen fibril organization, we found a marked alteration in small and large fibrils in

both heterozygous and *Crtap*$^{-/-}$ mice compared to wildtype that varied in severity depending on

the tendon or ligament being examined. Stable, HP cross-links were also elevated at 1- and 4-

months in *Crtap*$^{-/-}$ mice compared to wildtype, which indicates an increase in telopeptide lysine

hydroxylation and irreversible intermolecular cross-links (see below). RNA-seq analysis

revealed alterations in extracellular matrix proteins, growth factor signaling, and metabolic

pathways in *Crtap*$^{-/-}$ tendons and ligaments compared to wildtype controls. Finally, we

demonstrate that *Crtap*$^{-/-}$ mice exhibit motor impairments concomitant with reductions in grip

strength – a phenomenon that may be related to the tendon pathology observed in these mice.

Collagen ultrastructure and cross-linking has been well-documented in the bones of

mouse models for both dominant and recessive OI; however, quantitative differences in the

volume of tendons and ligaments in these models has not been examined and tendon/ligament

histological characterization has been modest. In this regard, we demonstrate that *Crtap*[-/-] mice

have thinner Achilles tendons and patellar ligaments at 1- and 4-months. Despite the reduction in

size and extracellular matrix, tendons and ligaments from these mice are hypercellular compared

to wildtype and heterozygous animals. At the same time, there is a significant reduction in

postnatal CD146+CD200+ tendon progenitors with a concomitant increase of immature cells.

While the alterations in tendon and ligament size and cellularity may be a result of alterations in

the collagen extracellular matrix, it could also be associated with alterations in cellular signaling.

In this regard, TGF-β signaling is upregulated in bones from *Crtap*[-/-] mice[34], and elevated TGF-β

signaling has been noted in mouse models with increased tendon cellularity and alterations in

collagen fibril distribution[35]. Indeed, our RNA-seq analysis identified TGF-β signaling as an

activated upstream regulator predicted to be driving differential gene expression observed in the

tendons and ligaments from *Crtap*[-/-] mice. We also observed significant metabolic dysregulation

in *Crtap*[-/-] ligaments. Interestingly, this a phenotype that has been observed in mammalian target

of rapamycin complex 1 (mTORC1) knockout mouse tendons that have a similar histological

phenotype to those of *Crtap*[-/-] mice[35]. These data are consistent with the principle that an altered

extracellular matrix can lead to altered cellular signaling as previously shown for vascular tissue

in Marfan syndrome and OI bone. Here, we extend this theme to tendon and ligamentous defects

in OI.

Collagen fibril assembly is a dynamic process that begins with the formation of many

small fibrils that grow longitudinally during development, followed by appositional fusion and

continued longitudinal growth as tendons mature[4]. In the present study, we found an increased

proportion of small and large collagen fibrils in the FDL and Achilles tendons as well as patellar

ligaments of *Crtap*[-/-] mice compared to wildtype. These distributions varied in severity across the

three tissues, with the greatest differences being observed in the patellar ligament. *Leprel*[-/-] mice

have been reported to display an increased proportion of small collagen fibrils in tail tendons[19],

indicating that loss of the 3-prolyl hydroxylase complex alters collagen fibrillogenesis. At the

same time, the increased number of small fibers was more pronounced in *Leprel*[-/-] mice, and

there was no evidence of an increased proportion of large fibers[19] as observed for *Crtap*[-/-] mice in

this study. Similar to the *Leprel*[-/-] mouse model, mice lacking CypB also exhibit a pronounced

increase in the number of small collagen fibrils within tail tendons[18]. Together, this suggests that

despite forming a complex with P3h1 and CypB, loss of Crtap has distinct consequences on

collagen fibrillogenesis. In this regard, it is important to note that tail tendons were used for

collagen fibril distribution assessments in both *Leprel*[-/-] and *Ppib*[-/-] mice[18, 19], whereas

appendicular tendons and ligaments were examined in our study. The difference in tissue type

examined might also explain why we observed unique alterations in collagen fibril distribution in

heterozygous *Crtap* mice, whereas *CypB* heterozygous tendons were indistinguishable from

wildtype[18].

Type I procollagen molecules undergo post-translational modifications within the

endoplasmic reticulum, including lysyl-hydroxylation and prolyl-hydroxylation, that are critical

for proper collagen synthesis, transport, and stability. Specifically, telopeptide lysine

hydroxylation results in mature lysyl-pyridinoline (LP) or hydroxylysyl-pyridinoline (HP)

residues after lysyl oxidase oxidation, which as permanent, irreversible crosslinks play a role in

regulating fibril growth and strength[25, 36]. Previous literature has demonstrated that loss of the 3-

prolyl hydroxylase complex caused by deletion of P3h1 (*Leprel*[-/-]) or Crtap (*Crtap*[-/-]) prevents

prolyl 3-hydroxylation of clade A (type I, II and III) collagens and can lead to changes in lysine

post-translational modifications due to loss of its chaperone function[10, 11]. In this study, we found

that mature collagen cross-links (HP residues per collagen) are markedly increased in the FDL

and Achilles tendons as well as the patellar ligament of *Crtap*[-/-] mice relative to wildtype.

Interestingly, we observed increased cross-links in the Achilles, but not FDL or patellar tendons

of 1-month old heterozygous mice, indicating a mild haploinsufficient effect of Crtap on this

tendon biochemical property thereby suggesting a rate-limited contribution of this complex in

this function. Outside of the genotype-specific effects, we saw an age-dependent increase in HP

residues per collagen in all genotypes. This observation is consistent with a study by Taga and

colleagues that reported an increase in 3-hydroxyproline residues in rat tendon collagen (but not

bone or skin) that plateaued at 3 months-of-age[37]. Together with the TEM analyses, these data

suggest that altered collagen cross-linking in tendons and ligaments from *Crtap*[-/-] mice may

adversely affect collagen fibril assembly.

In addition to skeletal deformities and frequent fractures, severe OI is associated with

motor impairments including gait abnormalities, chronic pain, and reduced muscle strength[8, 9, 38].

In this study, we showed that *Crtap*[-/-] mice exhibit reduced motor activity and coordination using

the open field, rotarod, and grid foot slip assays. We also observed a reduction in latency to fall

on the inverted grid assay that was mirrored by a dramatic loss of grip strength compared to

heterozygous and wildtype mice. These results are consistent with findings reported for the

Col1a1[Jrt/+] mouse model of severe OI and Ehlers-Danlos syndrome[17]. Specifically, Abdelaziz

and colleagues found that *Col1a1*[Jrt/+] mice displayed reduced motor activity using the open field

and running wheel assays – a phenotype they attributed to thermal hyperalgesia and mechanical

allodynia in these mice[39]. In this regard, the reduced vertical activity we observed in *Crtap*[-/-]

mice may be indicative of a pain or spinal phenotype, suggesting that the characterization of pain

in this model is a useful avenue for future research. Overall, this study represents one of the first

extensive characterizations of behavioral deficits in a mouse model of severe, recessive OI that

also correlates these deficits to tendon phenotypes.

Taken together, this study provides the first evidence for tendon and ligament phenotypes

in the *Crtap*[-/-] mouse model of severe recessive OI. We also provide compelling evidence for a

strong motor activity and coordination phenotype in these mice. As the quality of life is so

impacted in patients with OI, a more comprehensive evaluation of behavioral outcomes in future

preclinical studies may provide important insights into the efficacy of therapeutic interventions.

AUTHOR CONTRIBUTIONS

M. W. Grol: Conception and design of the study, acquisition, analysis and interpretation of data,

drafting and editing of the manuscript

N. A. Haelterman: Acquisition and analysis of data, editing of the manuscript

J. Lim: Acquisition and analysis of data, editing of the manuscript

E. Munivez: Acquisition of data

M. Archer: Acquisition and analysis of data

D. M. Hudson: Acquisition, analysis, and interpretation of data, editing of the manuscript

S. F. Tufa: Acquisition and analysis of data

D. R. Keene: Acquisition, analysis, and interpretation of data, editing of the manuscript

K. Lei: Acquisition and analysis of data

D. Park: Analysis and interpretation of data

D. R. Eyre: Analysis and interpretation of data, editing of the manuscript

B H. Lee: Conception and design of the study, interpretation of data, editing of the manuscript

Competing Interests

No competing interests to report.

Funding

BCM Intellectual and Developmental Disabilities Research Center (HD024064) from the Eunice

Kennedy Shriver National Institute Of Child Health & Human Development, the Rolanette and

Berdon Lawrence Bone Disease Program of Texas, the BCM Center for Skeletal Medicine and

Biology, and the Pamela and David Ott Center for Heritable Disorders of Connective Tissue. D.

Eyre was supported by a NIAMS R37 (AR37318).

REFERENCES

1. Nourissat G, Berenbaum F, Duprez D. Tendon injury: from biology to tendon repair. Nat Rev
    Rheumatol 2015; 11: 223-233.

2. Screen HR, Berk DE, Kadler KE, Ramirez F, Young MF. Tendon functional extracellular
    matrix. J Orthop Res 2015; 33: 793-799.

3. Kannus P. Structure of the tendon connective tissue. Scand J Med Sci Sports 2000; 10: 312-
    320.

4. Kalson NS, Lu Y, Taylor SH, Starborg T, Holmes DF, Kadler KE. A structure-based
    extracellular matrix expansion mechanism of fibrous tissue growth. Elife 2015; 4.

5. Saito M, Marumo K. Collagen cross-links as a determinant of bone quality: a possible
    explanation for bone fragility in aging, osteoporosis, and diabetes mellitus. Osteoporos Int
    2010; 21: 195-214.

6. Alford AI, Kozloff KM, Hankenson KD. Extracellular matrix networks in bone remodeling.

Int J Biochem Cell Biol 2015; 65: 20-31.

7. Lim J, Grafe I, Alexander S, Lee B. Genetic causes and mechanisms of osteogenesis

imperfecta. Bone 2017; 102: 40-49.

8. Arponen H, Makitie O, Waltimo-Siren J. Association between joint hypermobility, scoliosis,

and cranial base anomalies in paediatric osteogenesis imperfecta patients: a retrospective

cross-sectional study. BMC Musculoskelet Disord 2014; 15: 428.

9. Primorac D, Anticevic D, Barisic I, Hudetz D, Ivkovic A. Osteogenesis imperfecta--multi-

systemic and life-long disease that affects whole family. Coll Antropol 2014; 38: 767-772.

10. Hudson DM, Eyre DR. Collagen prolyl 3-hydroxylation: a major role for a minor post-

translational modification? Connect Tissue Res 2013; 54: 245-251.

11. Morello R, Bertin TK, Chen Y, Hicks J, Tonachini L, Monticone M, et al. CRTAP is

required for prolyl 3- hydroxylation and mutations cause recessive osteogenesis imperfecta.

Cell 2006; 127: 291-304.

12. Baldridge D, Schwarze U, Morello R, Lennington J, Bertin TK, Pace JM, et al. CRTAP and

LEPRE1 mutations in recessive osteogenesis imperfecta. Hum Mutat 2008; 29: 1435-1442.

13. Barnes AM, Chang W, Morello R, Cabral WA, Weis M, Eyre DR, et al. Deficiency of

cartilage-associated protein in recessive lethal osteogenesis imperfecta. N Engl J Med 2006;

355: 2757-2764.

14. Cabral WA, Chang W, Barnes AM, Weis M, Scott MA, Leikin S, et al. Prolyl 3-hydroxylase

1 deficiency causes a recessive metabolic bone disorder resembling lethal/severe

osteogenesis imperfecta. Nat Genet 2007; 39: 359-365.

15. van Dijk FS, Nesbitt IM, Zwikstra EH, Nikkels PG, Piersma SR, Fratantoni SA, et al. PPIB

mutations cause severe osteogenesis imperfecta. Am J Hum Genet 2009; 85: 521-527.

16. Baldridge D, Lennington J, Weis M, Homan EP, Jiang MM, Munivez E, et al. Generalized

connective tissue disease in *Crtap-/-* mouse. PLoS One 2010; 5: e10560.

17. Chen F, Guo R, Itoh S, Moreno L, Rosenthal E, Zappitelli T, et al. First mouse model for

combined osteogenesis imperfecta and Ehlers-Danlos syndrome. J Bone Miner Res 2014; 29:

1412-1423.

18. Terajima M, Taga Y, Chen Y, Cabral WA, Hou-Fu G, Srisawasdi S, et al. Cyclophilin-B

modulates collagen cross-linking by differentially affecting lysine hydroxylation in the

helical and telopeptidyl domains of tendon type I collagen. J Biol Chem 2016; 291: 9501-

9512.

19. Vranka JA, Pokidysheva E, Hayashi L, Zientek K, Mizuno K, Ishikawa Y, et al. Prolyl 3-

hydroxylase 1 null mice display abnormalities in fibrillar collagen-rich tissues such as

tendons, skin, and bones. J Biol Chem 2010; 285: 17253-17262.

20. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods 2012; 9: 676-682.

21. Nixon AJ, Grol MW, Lang HM, Ruan MZC, Stone A, Begum L, et al. Disease-modifying

osteoarthritis treatment with interleukin-1 receptor antagonist gene therapy in small and large

animal models. Arthritis Rheumatol 2018; 70: 1757-1768.

22. Ruan MZ, Erez A, Guse K, Dawson B, Bertin T, Chen Y, et al. Proteoglycan 4 expression

protects against the development of osteoarthritis. Sci Transl Med 2013; 5: 176ra134.

23. Ruan MZ, Patel RM, Dawson BC, Jiang MM, Lee BH. Pain, motor and gait assessment of

murine osteoarthritis in a cruciate ligament transection model. Osteoarthritis Cartilage 2013;

21: 1355-1364.

24. Stone A, Grol MW, Ruan MZC, Dawson B, Chen Y, Jiang MM, et al. Combinatorial *Prg4*

and *Il-1ra* gene therapy protects against hyperalgesia and cartilage degeneration in post-

traumatic osteoarthritis. Hum Gene Ther 2019; 30: 225-235.

25. Eyre D. Collagen cross-linking amino acids. Methods Enzymol 1987; 144: 115-139.

26. Hudson DM, Archer M, King KB, Eyre DR. Glycation of type I collagen selectively targets

the same helical domain lysine sites as lysyl oxidase-mediated cross-linking. J Biol Chem

2018; 293: 15620-15627.

27. Steinmann S, Pfeifer CG, Brochhausen C, Docheva D. Spectrum of tendon pathologies:

triggers, trails and end-state. Int J Mol Sci 2020; 21.

28. Dyment NA, Hagiwara Y, Matthews BG, Li Y, Kalajzic I, Rowe DW. Lineage tracing of

resident tendon progenitor cells during growth and natural healing. PLoS One 2014; 9:

e96113.

29. Dyment NA, Liu CF, Kazemi N, Aschbacher-Smith LE, Kenter K, Breidenbach AP, et al.

The paratenon contributes to scleraxis-expressing cells during patellar tendon healing. PLoS

One 2013; 8: e59944.

30. Howell K, Chien C, Bell R, Laudier D, Tufa SF, Keene DR, et al. Novel model of tendon

regeneration reveals distinct cell mechanisms underlying regenerative and fibrotic tendon

healing. Sci Rep 2017; 7: 45238.

31. Harvey T, Flamenco S, Fan CM. A Tppp3(+)Pdgfra(+) tendon stem cell population

contributes to regeneration and reveals a shared role for PDGF signalling in regeneration and

fibrosis. Nat Cell Biol 2019; 21: 1490-1503.

32. Lee CH, Lee FY, Tarafder S, Kao K, Jun Y, Yang G, et al. Harnessing endogenous

stem/progenitor cells for tendon regeneration. J Clin Invest 2015; 125: 2690-2701.

33. Feng H, Xing W, Han Y, Sun J, Kong M, Gao B, et al. Tendon-derived cathepsin K-

expressing progenitor cells activate Hedgehog signaling to drive heterotopic ossification. J

Clin Invest 2020.

34. Grafe I, Yang T, Alexander S, Homan EP, Lietman C, Jiang MM, et al. Excessive

transforming growth factor-beta signaling is a common mechanism in osteogenesis

imperfecta. Nat Med 2014; 20: 670-675.

35. Lim J, Munivez E, Jiang MM, Song IW, Gannon F, Keene DR, et al. mTORC1 Signaling is a

critical regulator of postnatal tendon development. Sci Rep 2017; 7: 17175.

36. Bateman JF, Boot-Handford RP, Lamande SR. Genetic diseases of connective tissues:

cellular and extracellular effects of ECM mutations. Nat Rev Genet 2009; 10: 173-183.

37. Taga Y, Kusubata M, Ogawa-Goto K, Hattori S. Developmental stage-dependent regulation

of prolyl 3-hydroxylation in tendon type I collagen. J Biol Chem 2016; 291: 837-847.

38. Garman CR, Graf A, Krzak J, Caudill A, Smith P, Harris G. Gait deviations in children with

osteogenesis imperfecta type I. J Pediatr Orthop 2019; 39: e641-e646.

39. Abdelaziz DM, Abdullah S, Magnussen C, Ribeiro-da-Silva A, Komarova SV, Rauch F, et

al. Behavioral signs of pain and functional impairment in a mouse model of osteogenesis

imperfecta. Bone 2015; 81: 400-406.

    FIGURE LEGENDS

Figure 1. **Loss of Crtap causes thinning and hypercellularity of tendons and ligaments in**

young and mature mice. (A-C) representative H&E images of 1-month ankle joints. **(D-F)**,

representative H&E images of 1-month knee joints. **(G-I)** representative H&E images of 4-

month ankle joints. **(J-L)** representative H&E images of 4-month knee joints. For all

micrographs, higher magnification images of the mid-tendon/ligament are illustrated. n = 3-4

mice per group. Scale bar is 1-mm. **(M-N)** quantification of cell density for Achilles tendons and

patellar ligaments of wildtype and *Crtap^{-/-}* mice at 1-month **(M)** and 4-months **(N)** of age. Data

are min-to-max box and whisker plots with individual points indicated. n = 3 mice per group.

Data passed the Shapiro-Wilk test for normality, and groups were compared using two-tailed

unpaired t-tests. Exact *p*-values are reported. **(O-P)** representative phase-contrast μCT images of

4-month wildtype **(O)** and *Crtap^{-/-}* **(P)** knee joints. Blue indicates the patellar ligament, green

indicates the femoral articular cartilage, and red indicates the tibial articular cartilage. Scale bar

is 1-mm. **(Q)** volumetric quantification of the patellar ligament volume in wildtype,

heterozygous, and *Crtap^{-/-}* mice. Data are min-to-max box and whisker plots with individual

points indicated. n = 3 mice per group. Data passed the Shapiro-Wilk test for normality, and

groups were compared using one-way ANOVA with Tukey's post-hoc tests. Exact *p*-values are

reported.

Figure 2. **Loss of Crtap in patellar ligaments leads to a decrease in progenitor cells and an**

accumulation of immature resident tissue cells. (A-B) patellar ligament cells isolated from 5-

month-old wildtype **(A)** or *Crtap^{-/-}* **(B)** mice were analyzed for the expression of CD200 and

CD146 tendon/ligament progenitor markers (*top histogram*) within the CD45^-CD31^- population

(*bottom histogram*). The plots are representative from a single wildtype or *Crtap*[-/-] mouse. **(C-D)**

graphs show the percentage of CD45[-]CD31[-]CD146[+]CD200[+] progenitor cells **(C)** and CD45[-]

CD31[-]CD146[-]CD200[+] immature ligament cells **(D)** from 5-month wildtype and *Crtap*[-/-] patellar

ligaments. Data are min-to-max box and whisker plots with individual points indicated. n = 5

mice per group. Data passed the Shapiro-Wilk test for normality, and groups were compared

using two-tailed unpaired t-tests. Exact *p*-values are reported.

Figure 3. **Collagen fibril diameter is altered in tendons and ligaments from heterozygous**

and *Crtap*[-/-] mice. (A-C) representative TEM images of 4-month FDL tendon collagen fibrils.

Scale bar is 500-nm. **(D)** representative histogram of the size distribution for collagen fibrils in

FDL tendons. Data are representative of n = 3 mice. **(E-G)** representative TEM images of 4-

month Achilles tendon collagen fibrils. Scale bar is 500-nm. **(H)** representative histogram of the

size distribution for collagen fibrils in Achilles tendons. Data are representative of n = 3 mice. **(I-**

K) representative TEM images of 4-month patellar ligament collagen fibrils. Scale bar is 500-

nm. **(L)** representative histogram of the size distribution for collagen fibrils in patellar ligaments.

Data are representative of n = 3 mice per group.

Figure 4. **Collagen cross-linking is increased in tendons and ligaments from young and**

mature *Crtap*[-/-] mice. Quantification of collagen cross-links as hydroxylysyl-pyridinoline (HP)

residues per collagen molecule for **(A)** FDL tendons; **(B)** Achilles tendons; and **(C)** patellar

ligaments. Data are shown as means ± S.D. n = 3-4 mice per group. Data passed the Shapiro-

Wilk test for normality, and groups were compared using one-way ANOVA with Tukey's post-

hoc tests. Exact *p*-values are reported.

Figure 5. **Transcriptome analysis of tendons and ligaments from 1-month-old *Crtap*$^{-/-}$ mice.**

DESeq2 was used to compare gene expression between wildtype and *Crtap*$^{-/-}$ Achilles tendon

and patellar ligament RNA samples, and genes with an adjusted *p*-value < 0.05 and absolute log$_2$

fold change > 1 were considered as differentially expressed. **(A)** a bi-clustering heatmap of the

top 30 differentially expressed genes between wildtype and *Crtap*$^{-/-}$ Achilles tendons sorted by

adjusted *p*-value and plotted according to log$_2$ transformed expression values. The Wald test was

used to generate p-values and log$_2$ fold changes. **(B)** significantly differentially expressed genes

between Achilles tendons from wildtype and *Crtap*$^{-/-}$ mice were clustered by their gene ontology,

and enrichment for gene ontology terms was tested using Fisher exact test. All gene ontology

terms with an adjusted *p*-value < 0.05 are plotted according to their -log$_{10}$ adjusted *p*-value. **(C)**

select upstream regulators predicted as being activated (shown in green), inhibited (shown in

red), or of unclear state (shown in grey) in *Crtap*$^{-/-}$ compared to wildtype Achilles tendons plotted

according to their -log$_{10}$ adjusted *p*-value. n = 3 mice per genotype for **(A-C)**. **(D)** a bi-clustering

heatmap of the top 30 differentially expressed genes between wildtype and *Crtap*$^{-/-}$ patellar

ligaments sorted by adjusted *p*-value and plotted according to log$_2$ transformed expression

values. The Wald test was used to generate p-values and log$_2$ fold changes. **(E)** significantly

differentially expressed genes between Achilles tendons from wildtype and *Crtap*$^{-/-}$ mice were

clustered by their gene ontology, and enrichment for gene ontology terms was tested using Fisher

exact test. All gene ontology terms with an adjusted *p*-value < 0.05 are plotted according to their

-log$_{10}$ adjusted *p*-value. **(F)** select upstream regulators predicted as being activated (shown in

green), inhibited (shown in red), or of unclear state (shown in grey) in *Crtap*$^{-/-}$ compared to

wildtype patellar ligaments plotted according to -log$_{10}$ adjusted *p*-value. n = 2 mice per genotype

for **(D-F)**.

Figure 6. **Motor activity and coordination are impaired in 4-months-old *Crtap*$^{-/-}$ mice. (A-B)**

quantification of spontaneous motor activity including horizontal **(A)** and vertical **(B)** activity

over a 30-min period using the open field assay. Data are min-to-max box and whisker plots with

individual points indicated. n = 9-12 mice per group. For **(A)**, data passed the Shapiro-Wilk test

for normality, and groups were compared using one-way ANOVA with Tukey's post-hoc tests.

For **(B)**, data failed the Shapiro-Wilk test for normality, and groups were compared using

Kruskal-Wallis with Dunn's post-hoc tests. Exact p-values are reported for **(A-B)**. **(C)**

quantification of motor activity, coordination, and endurance across 10 trials conducted over 2

days using an accelerating rotarod assay. Data are means ± S.D. n = 9-12 mice per group. Groups

were compared using repeated-measures two-way ANOVA with Tukey's post-hoc tests. Exact *p*-

values are reported, with black *p*-values being compared to wildtype, and blue *p*-values being

compared to heterozygous mice. **(D-E)** quantification of forelimb **(D)** and hindlimb **(E)** motor

coordination using the grid foot slip assay. Data are min-to-max box and whisker plots with

individual points indicated. n = 9-11 mice per group. Data failed the Shapiro-Wilk test for

normality, and groups were compared using Kruskal-Wallis with Dunn's post-hoc tests. Exact p-

values are reported. **(F)** quantification of forelimb and hindlimb grip strength using the inverted

grid assay conducted for 120 s. A reduction in latency to fall indicates reduced strength. Data are

min-to-max box and whisker plots with individual points indicated. n = 8-9 mice per group. Data

failed the Shapiro-Wilk test for normality, and groups were compared using Kruskal-Wallis with

Dunn's post-hoc tests. Exact p-values are reported. **(G)** quantification of forelimb grip strength

in N of force measured over 3 trials and then averaged. Data are min-to-max box and whisker

plots with individual points indicated. n = 9-12 mice per group. Data passed the Shapiro-Wilk

test for normality, and groups were compared using one-way ANOVA with Tukey's post-hoc

tests. Exact p-values are reported.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 5

Achilles Tendon (1-month)

A

Crtap⁻/⁻ Wildtype

Log² Expression Value:

Upregulated: 99   Downregulated: 79

B

GO Terms

0051928 — Positive Regulation of Calcium Ion Transport

0045778 — Positive Regulation of Ossification

0007155 — Cell Adhesion

Adjusted *p*-value (-log₁₀)

C Upstream Regulators

TP53
TNF
CTNNB1
DMD
PDGF-BB
β-Estradiol
TGF-β1

Adjusted *p*-value (-log₁₀)

Patellar Ligament (1-month)

D

Crtap⁻/⁻ Wildtype

Log² Expression Value:

Upregulated: 224   Downregulated: 295

E

GO Terms

0045906 — Negative Reg. of Vasoconstriction
0045444 — Fat Cell Differentiation
0008284 — Positive Reg. of Cell Proliferation
0055114 — Oxidation-Reduction Process
0050872 — White Fat Cell Differentiation
0006629 — Lipid Metabolic Process
0007155 — Cell Adhesion
0001525 — Angiogenesis
0050873 — Browin Fat Cell Differentiation

Adjusted *p*-value (-log₁₀)

F Upstream Regulators

ATF4
TGF-β3
IL-6
FGF2
PDGF-BB
CTNNB1
TNF

Adjusted *p*-value (-log₁₀)

FIGURE 6