<u>Anti-biofilm efficacy of a medieval treatment for bacterial infection requires</u>

<u>the combination of multiple ingredients</u>.

<u>Jessica Furner-Pardoe*[1,2], Blessing O Anonye*[1], Ricky Cain[1†], John Moat[3], Catherine A. Ortori[4],</u>

<u>Christina Lee[5], David A. Barrett [4], Christophe Corre[1,6] and Freya Harrison[1]</u>

[1] School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry, CV4 7AL

[2] Warwick Medical School, Gibbet Hill Campus, University of Warwick, Coventry, CV4 7AL

[3] Warwick Antimicrobial Screening Facility, School of Life Sciences, Gibbet Hill Campus, University of

Warwick, Coventry, CV4 7AL

[4] Centre for Analytical Bioscience, Advanced Materials and Healthcare Technologies Division, School

of Pharmacy, University of Nottingham, Nottingham NG7 2RD, UK

[5] School of English, University of Nottingham, Nottingham NG7 2RD, UK

[6] Department of Chemistry, University of Warwick, University of Warwick, Coventry, CV4 7AL

[†]Present address: Evotec (U.K.) Ltd., 114 Innovation Drive, Milton Park, Abingdon, Oxfordshire OX14

4RZ, United Kingdom

* These authors contributed equally

<u>Author contributions</u>

Conceived the study: CL, FH

Formulated hypotheses and designed experiments: JFP, BOA, RC, JM, CAO, DAB, CC, FH

Conducted experimental work: JFP, BOA, RC, FH

Analysed data: JFP, BOA, FH

Drafted the manuscript: JFP, BOA

Contributed to manuscript preparation: RC, JM, CAO, CL, DAB, CC, FH

Data supplement will be supplied with accepted/published article

<u>Abstract</u>

Novel antimicrobials are urgently needed to combat the increasing occurrence of drug-resistant

bacteria and to overcome the inherent difficulties in treating biofilm-associated infections. Research

into natural antimicrobials could provide candidates to fill the antibiotic discovery gap, and the study

of plants and other natural materials used in historical infection remedies may enable further

discoveries of natural products with useful antimicrobial activity. We previously reconstructed a

1,000-year-old remedy containing onion, garlic, wine, and bile salts, which is known as 'Bald's

eyesalve', and showed it to have promising antibacterial activity. In this paper, we have found this

remedy has bactericidal activity against a range of Gram-negative and Gram-positive wound

pathogens in planktonic culture and, crucially, that this activity is maintained against *Acinetobacter*

baumannii, Stenotrophomonas maltophilia, Staphylococcus aureus, *Staphylococcus epidermidis* and

Streptococcus pyogenes in a model of soft-tissue wound biofilm. While the presence of garlic in the

mixture is sufficient to explain activity against planktonic cultures, garlic alone has no activity against

biofilms. We have found the potent anti-biofilm activity of Bald's eyesalve cannot be attributed to a

single ingredient and requires the combination of all ingredients to achieve full activity. Our work

highlights the need to explore not only single compounds but also mixtures of natural products for

treating biofilm infections. These results also underline the importance of working with biofilm models

when exploring natural products for the anti-biofilm pipeline.

<u>Importance</u>

Bacteria can live in two ways, as individual planktonic cells or as a multicellular biofilm. Biofilm helps

protect bacteria from antibiotics and makes them much harder to treat. Both the biofilm lifestyle and

the evolution of antibiotic resistance mean we urgently need new drugs to treat infections. Here, we

show that a medieval remedy made from onion, garlic, wine, and bile can kill a range of problematic

bacteria grown both planktonically and as biofilms. A single component of the remedy – allicin, derived

from garlic – is sufficient to kill planktonic bacteria. However, garlic or allicin alone do nothing against

bacteria when they form a biofilm. All four ingredients are needed to fully kill bacterial biofilm

communities, hinting that these ingredients work together to kill the bacteria. This suggests that

future discovery of antibiotics from natural products could be enhanced by studying combinations of

ingredients, rather than single plants or compounds.

Introduction

Widespread multidrug resistance of once-susceptible pathogens, combined with a lack of

success in developing novel antimicrobials, has resulted in a looming crisis. Antimicrobial

resistance leads to problematic infections, threatens the success of routine surgery and

cancer treatments (1), and is estimated to kill 10 million people per year by 2050 (2,3). One

particularly troublesome area is biofilm-associated infection. Biofilm infections are estimated

to cost the UK's National Health Service over a billion pounds every year, with this cost only

set to increase (4,5). Biofilms are communities of bacteria that produce a protective

extracellular matrix and are especially persistent (6). Biofilm eradication often requires 100-

1,000 times higher antibiotic concentrations to achieve clearance than the same bacteria

growing planktonically (as individual free-floating cells)(7). *In vivo*, biofilms may essentially be

completely impervious to antibiotic treatment (8).

The path to overcoming biofilm infections requires a multifaceted response, including the

discovery and clinical deployment of novel antimicrobials. This search for novel candidates

must be directed towards those pathogenic bacteria with the highest health and economic

impact (e.g. ESKAPE group; (9–11)) and will be expedited if early discovery work on leads takes

into account the inherent difficulties of treating biofilm-associated infections (6,12).

The heightened urgency for novel antibiotics has resulted in natural products being revisited.

Just 200 years ago, our pharmacopoeia was dominated by herbal medicines. These medicines

have had varying levels of success for the treatment of infections in modern research, with a

number of potent compounds against planktonic cultures being identified (13,14); however,

these have been far less successful against biofilm cultures. We do not fully understand the

reasons for this lack of translational success of plant extracts.

It could be that the conventional process for developing drugs may miss key aspects of those

herbal remedies which could be effective against biofilms. Conventional drug development

calls for the isolation of single active compounds, whereas historical medicine usually calls for

combinations of whole plants (and other natural materials). There is some evidence that

whole plant extracts can have stronger biological effects than individual isolated compounds

(15). Examples of such synergy include observations that the anti-malarial activity of

artemisinin is enhanced by the presence of other compounds from the same plant (16,17) or

the combination of flavonoids used in Citrox® (15). Synergy may result from the presence of

molecules which potentiate the activity of antimicrobial compounds, or from multiple active

compounds with different mechanisms of action. It is also possible that in purifying individual

compounds to achieve readily quantifiable and characterised treatments, we may lose vital

interactions of natural products within the original mixture which prevent irritation or

toxicity.

Additionally, research into natural antimicrobial products is often limited by a methodological

focus on planktonic activity testing. The current gold standard for antibacterial testing is broth

microdilution, usually conducted in cation-adjusted Mueller-Hinton Broth (18), and this is

used extensively for the discovery of novel antibacterial compounds. However, whilst

providing a great high-throughput screening method, this does not provide information on

how the test substance interacts with bacteria in biofilms, or in settings which better mimic

the chemical environment experienced by pathogens *in vivo* (18,19). This focus may in part

explain the lack of translational success of isolated plant compounds, as their activity may be

sufficient to kill planktonic cells, but not to penetrate or kill biofilms – i.e. by not using biofilm

assays, we create a high false positive rate in the early stages of activity testing.

In this paper, we investigate the importance of the combination of ingredients for the

anti-biofilm activity of a 10th century remedy used for eye infection from the manuscript

known as Bald's Leechbook (London, British Library, Royal 12, D xvii). This remedy has

previously been shown to kill *S. aureus* biofilms by Harrison *et al.* (2015) (20), *Pseudomonas*

aeruginosa planktonic cultures (21) and recently *Neisseria gonorrhoeae* in a disk diffusion

assay (22). The recipe, known as 'Bald's eyesalve', requires equal volumes of garlic (*Allium*

sativum) and another *Allium* species (referred to as *cropleac* in the original Old English) to be

crushed together and mixed with equal volumes of wine and ox gall (bovine bile). We now

report two key new findings. First, we show the bactericidal activity of Bald's eyesalve against

a panel of clinically relevant strains of ESKAPE pathogens in planktonic culture, and against

biofilms of a subset of these strains. Second, we conducted a detailed assessment of whether

(a) any individual ingredient or (b) the sulphur-containing compound allicin (from garlic) can

explain this activity. No single ingredient recaptures the anti-biofilm activity of Bald's

eyesalve, and the presence of allicin cannot fully explain the ability of the whole mixture to

kill either planktonic cultures grown in host-mimicking medium (synthetic wound fluid), or

biofilms grown in an *in vivo*-like model of a soft tissue wound. The combination of ingredients

in the full mixture is the key to the remedy's apparent efficacy, and this is only apparent when

it is tested in host-mimicking models as opposed to planktonic cultures in standard laboratory

medium.

<u>Materials & Methods</u>

Bacterial strains and culture conditions

Strains used are listed in Table 1. All cultures were grown aerobically at 37°C and Miller's LB

agar (Melford) was used for plating.

Table 1. Bacterial strains used in the study

Species	Strain	Notes
Gram-positive		
Staphylococcus aureus	Newman	Standard laboratory strain, used in initial activity testing of all fresh eyesalve batches
Staphylococcus aureus	USA300 Los Angeles County clone	Well-studied example of community-acquired MRSA
Staphylococcus aureus	6 human carriage isolates	1 MRSA + 5 MSSA, gifted by Ruth Massey
Staphylococcus aureus	6 clones isolated from a chronic post-surgical wound	MSSA, with thanks to Tim Sloan
Staphylococcus epidermidis	1457	Standard laboratory strain, with thanks to Kim Hardie
Streptococcus pyogenes	Clinical isolate	Nottingham University Hospitals biobank, strain ref. 15W905066
Gram-negative		
Acinetobacter baumannii	Clinical isolate	Nottingham University Hospitals biobank, strain ref. 12W125768

Enterobacter cloacae	Clinical isolate	Nottingham University Hospitals biobank, training isolate
Escherichia coli	K12	Standard laboratory strain
Pseudomonas aeruginosa	PA14	Standard laboratory strain
Stenotrophomonas maltophilia	Clinical isolate	Nottingham University Hospitals biobank, training isolate

Media

Synthetic wound fluid (SWF) comprised equal volumes of foetal bovine serum (Gibco) and

peptone water (Sigma-Aldrich), (23). Mueller-Hinton Broth (MHB) was obtained from VWR.

HyClone water (GE Healthcare) was used throughout.

Bald's Eyesalve Reconstruction

Bald's eyesalve was prepared as previously reported (20). The outer skin of the garlic and

onion (sourced from local greengrocers) was removed. The garlic and onion were finely

chopped, and equal volumes of garlic and onion were crushed together using a mortar and

pestle for 2 minutes. Various sized batches of Bald's eyesalve were used throughout this

paper, ranging from final volumes of 30 ml – 400 ml, the average weight used was 14.1 ± 1.5

g of onion and 15.0 ± 1.3 g of garlic per 100 ml of Bald's eyesalve.

The crushed onion and garlic were then combined with equal volumes of wine (Pennard's

organic dry white, 11% ABV, sourced from Avalon Vineyard, Shepton Mallet) and bovine bile

salts (Sigma Aldrich) made up to 89 mg·ml^{-1} in water and sterilised by exposing to UV radiation

for 10 minutes (Carlton Germicidal Cabinet fitted with a 2537Å, 8-Watt UV tube). The mixture

was stored in sterilised glass bottles in the dark at 4°C for 9 days, after which it was strained

and centrifuged for 5 minutes at 1,811 g. The supernatant was then filtered using Whatman™

1001-110 Grade 1 Qualitative Filter Paper, Diameter: 11 cm, Pore Size: 11 μm. Filtered Bald's

eyesalve was stored in sterilised glass vials in the dark at 4°C.

For ease of reading, batches are numbered as they appear within the paper. The key for their

batch name is provided in the Data Supplement.

Preparation of Individual Ingredients and Dropouts

Individual ingredients were prepared such that their concentrations were equal to

concentrations present in the full remedy, a schematic is provided in Figure S1. To achieve

comparable concentrations, each ingredient was prepared as it would be for the full remedy

(onion and garlic crushed separately from each other), with all other ingredients substituted

with the same volume of water. A similar process was followed for "dropout" batches, where

one ingredient at a time was systematically excluded from the remedy and replaced with an

equal volume of water to maintain comparable concentrations to the original remedy.

Planktonic killing assay

Bacteria were cultured aerobically at 37°C on LB agar for 18-24 hours. Several colonies were

then inoculated into 5 ml SWF and incubated for 6 hours at 37°C on an orbital shaker. Aliquots

of each bacterial culture (100 μl) were added to wells of Corning® Costar® TC-Treated 96-Well

Plate and 50 μL of the eyesalve batch to be tested was added to 5 wells per strain. 50 μL of

water was used as a negative control. Plates were incubated in Tecan SPARK 10M at 37°C for

18 hours with periodic orbital shaking (10 seconds at 20 min intervals). Serial dilutions were

then performed and plated on LB agar plates.

Minimum Inhibitory Concentration Assay by Broth Microdilution

The minimum inhibitory concentrations (MIC) of eyesalve, single ingredients, dropout batches

and allicin were determined as described by Wiegand, Hilpert and Hancock (2008) in MHB

and in SWF. Treatments, or water as a negative control, were serially diluted with media in

Corning® Costar® TC-Treated 96-Well Plates, following the scheme in Table 2. Bacterial

isolates were streaked onto LB agar to obtain single colonies. After 18 - 24 hours incubation

at 37°C, three to five morphologically similar colonies were transferred to phosphate-

buffered saline (PBS) and diluted to 0.5 McFarland standard (OD_{600} 0.08-0.1). This suspension

was diluted 1 in 100 in media, resulting in 5 x 10^5 CFU/ml. The bacterial suspension was then

added to the treatment dilutions, resulting in another twofold dilution. The highest final

concentration of eyesalve, single ingredient or dropout batch tested was 50%. Plates were

incubated for 18 hours at 37°C. Results were visually checked for turbidity, and MIC values

are the lowest concentration where growth was not visible.

Table 2. Final volumes and calculated percentages of Bald's eyesalve in wells of 96-well plates

used for MIC.

Well	Bald's eyesalve (µl)	Media (µl)	Bacterial suspension (µl)	Final volume (µl)	% ESO in well
A	50	0	50	100	50
B	33	17	50	100	33
C	25	25	50	100	25
D	12.5	37.5	50	100	12.5
E	6.25	43.75	50	100	6.25
F	3.13	46.87	50	100	3.13
G	1.56	48.44	50	100	1.56
H	0.78	49.22	50	100	0.78

Biofilm Killing Assay in Synthetic Wounds

Biofilms were created in a synthetic soft-tissue wound model as described by Werthén *et al.*

(2010; (23)) and as used in our previous work (20). Briefly, synthetic wounds were created on

ice and comprised 2 mg·ml^{-1} collagen, 0.01% acetic acid, 60% [vol/vol] SWF, 10 mM sodium

hydroxide. Synthetic wounds were incubated at 37°C for 1 hour to allow collagen to

polymerise and placed under UV light for 10 minutes to ensure no contamination. Wounds

were either 400 μl in 24-well culture plates or 200 μl in 48-well plates.

Bacteria were incubated aerobically in SWF on an orbital shaker for 6 hours at 37°C, after

which cultures were diluted to OD$_{600}$ of 0.1 - 0.2 with fresh SWF. Bacteria suspension was

added to each synthetic wound (100 μl in 400 μl wounds or 50 μl in 200 μl wounds) and

incubated at 37°C for 24 hours to allow biofilm formation.

Wounds containing mature biofilms were then exposed to Bald's eyesalve or water (200 μl in

400 μl wounds or 100 μl in 200 μl wounds). Bacteria were recovered by treating with 300 μl

(400 μl wounds) or 150 μl (200 μl wounds) of 0.5 mg·ml^{-1} collagenase type 1 (EMD Millipore

Corp, USA) for 1 hour at 37°C to break down the matrix; the resulting liquid was serially diluted

and plated on LB plates. Plates were incubated at 37°C overnight, colonies were counted, and

CFU/wound calculated.

High-Performance Liquid Chromatography Quantification of Allicin

Allicin is an unstable and reactive compound under gas chromatography settings (24),

therefore, high-performance liquid chromatography (HPLC) was used to determine allicin

concentrations (24).

Various preparations of Bald's eyesalve were analysed by reversed-phase HPLC on an Agilent

1200 series system fitted with an Agilent ZORBAX Eclipse XDB-C18, 150 x 4.6 mm, 5 μm

particle size column and diode array detection at 210 nm. A gradient of methanol (5- 95%) in

water was used at a flow rate of 1 ml·min^{-1} over 30 minutes. Injection volume was 10 μl, and

the column temperature was 25°C.

To identify the concentration of allicin in eyesalve preparations, a calibration curve was

prepared by serially diluting an external allicin standard (Abcam, Cambridge) in water

(twofold dilutions, from 750 μg·ml^{-1} to 0.5 μg·ml^{-1}). External standards were run through HPLC

as described above (n = 2) and allicin peak area, in the 210 nm reading, was plotted against

concentration (μg·ml^{-1}) (Figure S2A and Data Supplement). Allicin external standards had a

retention time of approximately 15 minutes (Figure S2B). This peak was confirmed in Bald's

eyesalve by comparing fresh eyesalve to the same batch spiked with additional allicin

standard. The suspected peak increased as expected (Figure S2C-D).

Statistics

Data were analysed in R v3.5.1 (R Core Team, 2018) using the *car* (25), *lsmeans* (26), *multcomp*

(27) and *FSA* (28) packages. Raw data and R code for all experiments are provided in the Data

Supplement.

Results

*Consistent anti-biofilm activity of Bald's eyesalve and decision to use onion (*Allium cepa*) as*

"cropleac"

The meaning of Old English *cropleac,* as used in the original remedy, is ambiguous and may

refer to a variety of *Allium* species (29,30). Two likely translations are onion (*Allium cepa*) or

leek (*Allium porrum*). For this reason, multiple batches of the remedy using either *Allium*

species were made in our laboratory between 2014 and 2019. After the nine-day brewing

period, the activity of the remedies was tested against mature biofilms of *S. aureus* Newman

in an *in vivo*-like model of a soft tissue wound. In this assay, 24-hour-old biofilms, that have

been grown at 37°C in collagen-based synthetic wounds, are exposed to Bald's eyesalve or

water for 24 hours, and surviving bacteria are counted (20,23).

In total, 75 batches of Bald's eyesalve were made, including 15 pairs of batches where onion

(ESO) and leek (ESL) variants were made at the same time, using the same garlic, wine and

bile. As shown in Figure 1A, biofilm killing was achieved by 14/15 paired batches, with 22/30

preparations causing a >3-log drop in viable bacteria, when compared with water-treated

control biofilms. The mean log drop was not significantly different between paired ESO and

ESL batches (paired *t*-test t_{14}= 0.025, *p* = 0.981). We tested one example pair of ESO and ESL

variants (batch 6) against a panel of twelve clinical isolates of *S. aureus* grown in the synthetic

wound biofilm model. As shown in Figure 1B*,* ESO consistently showed more bactericidal

activity than ESL. This result may be due to the presence of antimicrobial compounds unique

to onion, or it may simply be due to onions being easier to crush in a mortar and pestle,

potentially making the extraction of natural compounds more efficient. To simplify further

analysis, follow on work was focussed on ESO.

As shown in Figure 2, biofilm killing was consistently achieved by multiple batches of ESO. Of

these batches, 62 of the 75 caused a >3-log drop in viable CFU, with 45/75 causing >5 log

drop, and 28 of these causing > 6-log drop, all relative to control biofilms treated with sterile

water.

Figure 1. **The activity of Bald's eyesalve, ESO and ESL variants, against *S. aureus*.** (A) 15 pairs of Bald's

eyesalve translating *cropleac* as either onion (ESO, grey) and leek (ESL, white), were prepared at the

same time (batches 1-15). Their activity was assessed against *S. aureus* Newman biofilms. Mature

biofilms of *S. aureus* Newman were grown in a model of a soft tissue wound, then treated with either sterile water or Bald's eyesalve (n = 3-5 replicates per treatment) for 24 hours before recovering bacteria for CFU counts. The mean log change in viable bacteria in treated vs control wounds was calculated for each preparation and no significant difference was seen (paired *t*-test t_{14}= 0.025, p = 0.981). Raw data are supplied in the Data Supplement. (B) Six carriage isolates of *S. aureus*, and six isolates from a chronic post-surgical wound (detailed in Table 1), were grown in the synthetic wound biofilm model and treated with ESO, ESL or water (n=3 per treatment). Each strain grew to different densities in untreated wounds (range approx. 10^6–10^8). The $\log_{10}$ drop in bacteria associated with treatment was calculated for each treated wound relative to the mean CFU in the three untreated wounds. Log drop data was analysed by ANOVA which revealed significant differences in log drop between eyesalve variants ($F_{1,48}$ = 732, p < 0.001) and strains ($F_{11,48}$ = 14.4, p < 0.001) and a significant strain-dependent difference in the magnitude of the effect of eyesalve variant (strain*variant interaction $F_{11,48}$ = 8.66, p < 0.001). The family-wise error rate was calculated from the ANOVA table and used to conduct planned contrasts of ESO vs ESL treatment effects for each strain using *t*-tests. ESO caused a larger reduction in viable cells than ESL for all strains (all p < 0.001). Raw data and R scripts are supplied in the Data Supplement.

Figure 2. The activity of 75 Bald's eyesalve (ESO) batches made by our group against *S. aureus* Newman

biofilms. Mature *S. aureus* Newman biofilms were grown in a model of a soft tissue wound, then treated with

either sterile water (control) or eyesalve (n = 3-5 replicates per treatment) for 24h before recovering bacteria

for CFU counts. All batches showed killing, and the log drop in treated vs control wounds was calculated. Raw

data are provided in the Data Supplement.

Broad-spectrum antibacterial activity of Bald's eyesalve against common wound pathogens

The antibacterial activity of three batches of ESO was tested in both planktonic and biofilm

cultures of eight strains of bacteria that commonly cause chronic wound infections. As shown

in Figure 3, ESO had potent activity against planktonic cultures of Gram-negative

(*P. aeruginosa* PA14, *A. baumannii* clinical isolate, *E. cloacae, S. maltophilia*) and Gram-

positive (*S. aureus* Newman, *S. aureus* USA300, *S. epidermidis* and *S. pyogenes*) wound

pathogens. ESO eradicated all planktonic cultures in all strains tested with the exception of

S. aureus USA300 and *S. maltophilia*, where a 3-4 log drop in viable bacteria was seen.

Figure 3. **Bald's eyesalve (ESO) activity against eight example strains of bacteria that commonly cause chronic**

wound infections. Planktonic cultures were grown for 6 hours in synthetic wound fluid. Cultures were treated

with ESO (batches 16-18) or sterile water (CT) to a final concentration of 33% v/v (n = 3-5 replicates per

treatment). After 18 hours bacteria were recovered for CFU counts. The dashed line represents the limit of

detection by plating. For *S. maltophila* and *S. aureus* USA300, where we did not observe complete killing, we

used ANOVA to determine that the CFU recovered from ESO-treated wounds was significantly different from

control wounds (*S. aureus* USA300: $F_{2,14}$ = 3458, $p < 0.001$; *S. maltophilia*: $F_{3,8}$ = 87.41, $p < 0.001$). Data were log

transformed to meet the assumptions of linear modelling. A Dunnett's test was conducted to compare the CFU

of the three batches with the CFU of the controls. All 3 batches of ESO were significantly different from their

respective controls for both isolates (all $p < 0.001$). Raw data and R scripts are supplied in the Data Supplement.

Mature biofilms of the above isolates were grown in the synthetic wound model and treated

with the same batches of eyesalve used for planktonic killing experiments. A 2-6-log drop in

viable cells was observed for the Gram-positives *S. aureus* Newman, *S. aureus* USA300,

S. epidermidis, S. pyogenes and the Gram-negative *A. baumannii* (Figure 4). No or inconsistent

killing was observed for *P. aeruginosa*, *E. cloacae* and *S. maltophilia* biofilms (Figure S3).

Figure 4. Bald's eyesalve (ESO) anti-biofilm activity. Mature biofilms of various isolates were grown in a model

of a soft tissue wound, then treated with either sterile water (control, CT) or eyesalve (batches 16-18), to a final

concentration of 33% (v/v), for 24h before recovering bacteria for CFU counts (n = 3-5 replicates per treatment).

The dashed line represents the limit of detection by plating. For *S. aureus* Newman, *S. aureus* USA300, and *S.*

epidermidis, where we did not observe complete killing, we used ANOVA to determine that the CFU recovered

from ESO-treated wounds was significantly different from control wounds (*S. aureus* Newman: $F_{3,8}$ = 107, *p* <

0.001; *S. aureus* USA300: $F_{3,14}$ = 103.5, *p* < 0.001; *S. epidermidis*: $F_{3,14}$ = 63.86, *p* < 0.001). Data were log

transformed to meet the assumptions of linear modelling. A Dunnett's test was conducted to compare the CFU

of the three batches with the CFU of the controls. All 3 batches of ESO were significantly different from their

respective controls for all three isolates (all *p* < 0.001). Raw data and R scripts are supplied in the Data

Supplement.

Garlic is responsible for the majority of planktonic killing by ESO

Our initial work indicated that all four ingredients in ESO were required to kill biofilms of *S.*

aureus Newman in synthetic wound biofilms (20). A recent publication by Fuchs *et al.* (2018;

(21)) concluded that the bactericidal activity of ESO was due to the presence of allicin from

garlic, however, the Fuchs *et al.* study only investigated planktonic killing in standard Mueller-

Hinton Broth (MHB). It is well known that planktonic cultures of bacteria can be up to 1,000

times more sensitive to antibiotics than the same isolates grown as biofilms (7), and that

antibiotic sensitivity is highly dependent on growth medium (31). This means conducting

assays only on planktonic MHB cultures risks underestimating the number or concentration

of bioactive agents in ESO required for killing in *in vivo*-like conditions (18,19).

To more robustly determine if the activity of ESO stems from one ingredient or several, we

prepared individual ingredients and preparations omitting one ingredient, such that the

concentrations of each ingredient in the whole remedy and the single ingredient or dropout

variants were equal. The MICs of these preparations, along with MICs of the full recipe, were

assessed in MHB and in SWF, in standard broth microdilution assays using four of the isolates

previously tested (two Gram-negatives, *A. baumannii* clinical isolate strain and *P. aeruginosa*

PA14, and two Gram-positives, *S. aureus* Newman and *S. aureus* USA300).

As shown in Table 3, the bacterial isolates varied in their sensitivity to the full ESO. Crucially,

there was an effect of the growth medium, such that MICs at least doubled in SWF compared

with MHB for *S. aureus* Newman and *A. baumannii*. Wine, bile or onion alone were much less

effective than the full remedy. In MHB, the MIC of garlic alone was the same as the ESO MIC

for all strains except *P. aeruginosa*, and in SWF the garlic and ESO MICs were either equal or

exhibited a max. 2-fold difference. This suggests that the planktonic activity of the remedy is

due to the presence of garlic. The results of MIC testing with ingredients omitted gave similar

results (Table 4). Preparations omitting garlic lost all or most of their antibacterial activity.

Removal of any other ingredient left activity largely unaffected, although the omission of wine

or bile doubled the MIC against *P. aeruginosa* PA14 in SWF. The discordance between MICs

in different media is more prominent here, with the removal of an ingredient having a minimal

effect in MHB but a large loss in activity when tested in SWF. This stresses the common

discordance between testing in standard rich lab medium and host-mimicking medium.

Table 3. **Minimum inhibitory concentration (MIC) of ESO or individual ingredients.** MICs were tested in both

Mueller-Hinton Broth (MHB) or synthetic wound fluid (SWF). MICs are presented as modal values of 3 different

batches (ESO 19-21; with 3 replicate MIC tests per batch) and are the percentage of treatment present at MIC

(v/v).

Treatment	*S. aureus* Newman		*S. aureus* USA300		*A. baumannii* clinical isolate		*P. aeruginosa* PA14	
	MHB	SWF	MHB	SWF	MHB	SWF	MHB	SWF
ESO	1.56	6.25	3.13	3.13	1.56	3.13	25	25
Onion	>50	>50	>50	>50	>50	>50	>50	>50

Garlic	1.56	12.5	3.13	3.13	1.56	6.25	33	33
Wine	>50	>50	>50	>50	>50	>50	>50	>50
Bile	>50	>50	>50	>50	>50	>50	>50	>50

357 **Table 4**. **Minimum inhibitory concentration (MIC) of ESO or preparations omitting a single ingredient.** MICs

358 were tested in both Mueller-Hinton Broth (MHB) and synthetic wound fluid (SWF). MICs are presented as modal

359 values of 3 different batches (ESO 22-24; with 3 replicate MIC tests per batch) and are the percentage of

360 treatment present at MIC (v/v).

Treatment		*S. aureus* Newman		*S. aureus* USA300		*A. baumannii* clinical isolate		*P. aeruginosa* PA14	
		MHB	SWF	MHB	SWF	MHB	SWF	MHB	SWF
	ESO	<0.78	12.5	3.13	12.5	1.56	6.25	25	25
Omitting	Onion	<0.78	12.5	3.13	12.5	3.13	6.25	25	25
	Garlic	50	50	50	50	50	>50	>50	>50
	Wine	<0.78	6.25	3.13	12.5	1.56	6.25	25	50
	Bile	<0.78	6.25	3.13	12.5	1.56	6.25	25	50

All four ingredients are necessary for activity against mature* S. aureus *biofilms in synthetic wounds

Consistent with previous results from our group, no single ingredient alone had any

bactericidal activity against mature *S. aureus* Newman biofilms in the synthetic wound model,

and removing any single ingredient resulted in a reduction in the anti-biofilm activity (Figure

5). Surprisingly, removal of wine caused a decrease in activity on par with that seen with the

removal of garlic, despite wine possessing very limited antimicrobial activity on its own in

either biofilm or planktonic assays (Table 3). For comparison with planktonic MIC data, the

final concentration of Bald's eyesalve in synthetic wound biofilm killing assays (Figures 1-6),

is 33% vol/vol. Therefore, in more *in vivo*-like biofilm conditions, combining all the ingredients

is necessary for full activity against *S. aureus* Newman.

Figure 5. **Contribution of each ingredient to Bald's eyesalve (ESO) anti-biofilm activity.** Mature biofilms of *S.*

aureus Newman were grown in a model of a soft tissue wound, then treated with either water (CT) or treatment

for 24 hours, before recovering bacteria for CFU counts. The dashed lines represent the limit of detection by

plating. A) ESO and individual ingredients prepared such that the concentrations of each ingredient in the whole

remedy and individual preparations are equal (n = 3 replicates per treatment). Three independent batches of

preparations are shown, corresponding to batches ESO 19, 20 and 21, shown in white, light grey and dark grey,

respectively. Analysis of the data by ANOVA showed a significant effects of treatment ($F_{4,30}$ = 389.215, $p < 0.001$)

but no effect of batch ($F_{2,30}$ = 1.24, $p = 0.305$) and no interaction between treatment and batch (i.e. the effect of

treatment did not depend on batch: $F_{8,30} = 0.930$, $p = 0.507$). Data were square root transformed to meet the

assumptions of linear modelling. To compare each treatment group to the untreated controls, we then fitted a

new ANOVA to fit least-squares means and variances for viable CFU recovered after each treatment (excluding

controls), averaged over the three batches. These fitted means were then compared to the mean observed in

the untreated controls using a two-tailed, unpaired Welch's t-test, which takes into account the unequal sample

sizes and variances in control and treatment groups. The decrease in CFU on treatment with full ESO was

significant ($t_{2.93} = 9.18$, $p = 0.003$). Garlic, onion and wine caused small but significant increases in viable CFU ($t_{2.93}$

$= 9.05$, $p = 0.003$, $t_{2.93} = 9.61$, $p = 0.003$, $t_{2.93} = 11.1$, $p = 0.002$), while bile had no significant effect on CFU ($t_{2.93} = $

2.44, $p = 0.094$). B) ESO and batches with one ingredient omitted and replaced with water, to keep the

concentrations of remaining ingredients equal to those in the full recipe. Three independent batches of

preparations are shown, corresponding to ESO batches 22, 23 and 24, shown in white, light grey and dark grey,

respectively. ANOVA showed significant differences in CFU between treatments ($F_{4,26} = 162$, $p < 0.001$) and

between batches ($F_{2,26} = 18.7$, $p < 0.001$), and a significant interaction between treatment and batch ($F_{8,26} = 5.85$,

$p < 0.001$); the graph shows that this interaction is due to different batches showing larger or smaller differences

between the killing effects of the preparations, but that the rank order of preparations by CFU is the same in

each batch. Data were log-transformed to meet the assumptions of linear modelling. As only two control data

points were obtained for this experiment, one-sample t-tests were performed using least-squares mean CFU in

the treated wounds to the mean CFU in control wounds; the decrease in CFU vs control was statistically

significant for ESO and all dropouts ($p \leq 0.002$) except the wine dropout ($p = 0.202$). A Dunnett's test was

conducted to compare each dropout with the full ESO. All dropouts resulted in higher CFU than the full ESO ($p < $

0.001). Raw data and full statistical results are supplied in the Data Supplement.

The antibacterial activity of ESO is not simply due to its allicin content

The study which attributed the planktonic activity of ESO to garlic further concluded that this

was specifically due to the presence of allicin (21), an organosulfur compound with well

known, potent antimicrobial activity *in vitro* (32–34). We used HPLC, calibrated against allicin

standards of known concentrations, to determine the allicin concentration in 12 batches of

fresh ESO. These batches were then assayed for MIC against *S. aureus* Newman in SWF. We

were then able to calculate the concentration of allicin present in the MIC of each batch and

compare this with the MIC of purified allicin.

As shown in Table 5, the allicin concentration in the 12 batches of ESO was found to be in the

range of 391 ± 19 µg·ml^{-1}. Therefore, the estimated concentration of allicin in their respective

MICs would be in the range 11 - 39 µg·ml^{-1}. This is much lower than the MIC obtained for

purified allicin, which was 62.5 µg·ml^{-1}. Due to experimental error, two MIC test (ESO batches

28, 29) were lost; we did not repeat this due to the time delay between the initial and

repeated test potentially affecting concentrations of compounds present, and thus activity.

Table 5. Minimum inhibitory concentrations (MIC) were calculated for fresh Bald's eyesalve (n = 12; batches

19-30), garlic only preparations (n = 3) and external allicin standards (n = 3). Allicin concentrations for each batch

were quantified using HPLC and calculated using an allicin calibration curve. MICs were measured in synthetic

wound fluid. The allicin concentration in the MIC of each batch, and the allicin concentration present in the

biofilm treatment experiments, is calculated and the results of biofilm treatment experiments are provided for

reference. Raw data is supplied in the Data Supplement.

Treatment	Allicin concentration range (mean ± s.e.)	Allicin concentration in MIC	Biofilm activity (log drop from control)	Average predicted allicin concentration in biofilm treatment
ESO	275 - 486 (391 ± 19) µg·ml^{-1}	11 - 39 µg·ml^{-1}	Eradicated biofilm (>5)	142 µg·ml^{-1}
Garlic only	358 - 434 (389 ± 19) µg·ml^{-1}	45 - 54 µg·ml^{-1}	No activity (0)	160 µg·ml^{-1}
Allicin Standard		62.5 µg·ml^{-1}	Partial activity (2.6)	167 µg·ml^{-1}

All of these 12 batches eradicated mature *S. aureus* Newman biofilms in the synthetic wound

assay (see Data Supplement). The results of biofilm killing assays for five representative

batches are depicted in Figure 6A, alongside a biofilm killing assay conducted for preparations

of 100, 300 or 500 µg·ml⁻¹ of purified allicin. Purified allicin at concentrations similar to those

found in ESO, had much less anti-biofilm activity (Figure 6B), and garlic preparations with the

same allicin concentration as ESO had no activity (data supplied in the supplementary data

file). Therefore, allicin alone does not explain either the activity of ESO against biofilm-grown

bacteria or its activity against planktonic cultures.

Figure 6. **Antimicrobial activity of Bald's eyesalve (ESO) and allicin standards.** Mature *S. aureus* Newman

biofilms were grown in a model of a soft tissue wound, then treated with either water (CT) or treatment for 24

hours, before recovering bacteria for CFU counts. A) Biofilms treated with 5 ESO batches, with their respective

allicin concentrations indicated. B) External allicin standards diluted in water. The dashed line represents the

limit of detection by plating. To statistically analyse, data were log transformed to meet the assumptions of

linear modelling. A one-way ANOVA was performed and found a significant difference between allicin

treatments and control cells ($F_{3,9} = 141$, $p < 0.001$). A Dunnett's test was conducted to compare the CFU of the

three allicin concentrations with the CFU of the controls. Of the concentrations, 300 µg·ml^{-1} and 500 µg·ml^{-1}

were significantly different ($p < 0.001$), 100 µg·ml^{-1} was not significantly different ($p = 0.247$). Raw data and full

statistical results are supplied in the Data Supplement.

Discussion

The activity of a range of natural products against microbes in simple *in vitro* assays (agar

diffusion, broth microdilution or simple surface-attached biofilm assays) demonstrates their

potential as a source of novel antibiotics (35–37). However, the standard approach of

purifying individual compounds from natural products rarely produces clinically useful

products and potent activity against planktonic bacteria in standard lab media rarely

translates to *in vivo* efficacy. This is an especially pressing problem in the case of biofilm

infections. Biofilms are much harder to treat due to reduced penetration of antibiotics

through the extracellular matrix and the enhanced tolerance of biofilm-grown cells to many

in-use antibiotics (6,12). Biofilm infections of wounds (e.g. burns, diabetic foot ulcers),

medical implants (e.g. artificial joints, catheters), the lungs (e.g. in cystic fibrosis) and other

body sites impose a major health and economic burden (38) and can be effectively

untreatable. Non-healing, infected foot ulcers, which can be a complication of diabetes,

provide an especially sobering example. Even if the infection is apparently successfully

treated, there is a high chance of recurrence and an estimated 50% of those affected die

within five years of ulcer development. Management of diabetic foot ulcers costs the UK's

NHS £650M per year (39).

Historical medical manuscripts often prescribe complex preparations of several ingredients

to treat infections. Thus, when considering natural products as a potential source of

anti-biofilm agents, we must consider the possibility that any efficacy they may possess could

rely on creating a cocktail of different products. Understanding the relationship between

combinations of natural products and antimicrobial activity may generate a novel way to

create new antibiotics from botanicals. Here, we confirm Bald's eyesalve as an example of an

"ancientbiotic" that requires the combination of all ingredients for potent activity against a

panel of clinically important bacterial strains.

Our research builds on previous work (20–22) to show that Bald's eyesalve can eradicate

planktonic cultures of a range of problematic Gram-positive and Gram-negative bacteria

including *P. aeruginosa*, *A. baumannii*, *E. cloacae*, *S. maltophilia*, *S. aureus*, *S. epidermidis* and

S. pyogenes. It is also able to cause a 4-log reduction in viable cell counts in planktonic cultures

of the especially problematic MRSA. Despite the widely understood problems of treating

biofilms (6,12), Bald's eyesalve was also able to significantly reduce viable cell counts in

biofilms of *S. epidermidis* and MRSA and was able to completely eradicate biofilms of *S. aureus*

Newman, *A. baumannii* and *S. pyogenes,* in an established soft-tissue wound model (23).

However, although there was promising planktonic activity against *P. aeruginosa*, *E. cloacae*

and *S. maltophilia,* variable or no activity was seen against biofilm cultures of these isolates

in the wound model. This highlights the importance of investigating the anti-biofilm activity

of candidate antibacterial agents, rather than extrapolating from the results of planktonic

assays. Bald's eyesalve shows great promise as an effective antimicrobial candidate, although

further development and combination with biofilm-degrading adjuvants may be necessary to

achieve activity against species such as *P. aeruginosa*.

Each of Bald's eyesalve ingredients has known antimicrobial properties or compounds (onion

and garlic: (34,37,40,41), bile: (42–44), wine: (45–47)). We explored the contribution of all

four ingredients to both planktonic and biofilm activity of Bald's eyesalve to build a picture of

their relative contributions. Planktonic activity appeared almost entirely attributable to garlic.

However, tests against *S. aureus* Newman biofilms, grown in a synthetic wound model,

showed garlic exhibited no antibacterial activity in this more clinically-relevant setting. In fact,

no preparation which omitted any one ingredient possessed full activity in the biofilm assay.

This confirms our previously published finding that Bald's eyesalve anti-biofilm activity is

contingent on the presence of all four ingredients (20).

Our results against planktonic cultures of *S. aureus* and *P. aeruginosa* align with a report by

Fuchs *et al.* 2018, that garlic alone, and specifically allicin, accounts for the majority of the

planktonic activity of Bald's remedy. Allicin is a defensive compound that is converted from

alliin by the enzyme alliinase, upon damage to the plant tissue. Allicin can kill a wide range of

both Gram-negative and Gram-positive bacteria *in vitro* (32) due to the thiosulfinate group (–

S(O)–S–) reacting with many cellular proteins in the pathogen (48). We therefore explored

the role of allicin in Bald's eyesalve in detail.

Crucially, we found that 12 batches of Bald's eyesalve were able to elicit much greater anti-

biofilm activity than purified allicin at comparable concentrations. Pure allicin at a

concentration of 500 $\mu g \cdot ml^{-1}$ was able to reduce viable cell numbers by 2-3 logs, whereas

Bald's eyesalve batches containing 275 - 486 $\mu g \cdot ml^{-1}$ allicin were able to cause a 6-7 log drop

and eradicate the biofilms. Even in planktonic tests, we found that the concentration of allicin

in the MIC of these batches of Bald's eyesalve was lower than the MIC of purified allicin.

Together, these results clearly illustrate that in addition to allicin, other ingredients in Bald's

eyesalve contribute to its activity. This highlights the importance of the combination of

ingredients.

The differences between our findings and those of Fuchs *et al.* are partly explained by the

contrasts between bacterial tolerance to killing (i) in MHB *versus* synthetic wound fluid; and

(ii) in planktonic culture *versus* established biofilms. This highlights the differences in

antibiotic susceptibility often seen in host-mimicking media versus standard MHB (18,19,49).

Further, the allicin concentrations found in our batches of Bald's eyesalve were lower than

that found in Fuchs *et al.* The highest allicin concentration we measured was 486 $\mu g \cdot ml^{-1}$,

compared with Fuchs *et al.*'s 836 $\mu g \cdot ml^{-1}$.

We think it unlikely that this is due to the differences in quantification methods used as both

are appropriate. We used HPLC, a method that has been widely used previously (33,50–53)

and is accepted to produce consistent and accurate results. Previous papers using HPLC have

detected allicin concentrations ranging from 1 $\mu g \cdot ml^{-1}$ to 2 $mg \cdot ml^{-1}$ (33,52), a range that the

results in this paper lie within. Fuchs *et al.* 2018 used quantitative NMR (qNMR), which too

has advantages and is being increasingly used in quantification of concentrations (54). It is

more likely that our Bald's eyesalve preparations really do have different allicin

concentrations. Alliin is released from storage vesicles and converted to allicin upon damage

to tissue: Fuchs *et al.*, 2018 used a food processor, in comparison to our pestle and mortar,

which may have resulted in greater damage to the tissue and therefore greater release of

alliin and more allicin created. Further, different garlic varieties are thought to have different

activities and compositions (55–59) and our research labs are based in different continents,

which presumably results in different garlic varieties and growth conditions. The higher allicin

concentration may explain the slightly increased planktonic activity seen in Fuchs *et al*. This

may also be due to their use of agitation in MIC assays, as agitation is known to lower the MIC

(60).

The step in the early Medieval remedy which specifies that the onion and garlic be ground

together would have been conducted manually – most likely with a pestle and mortar. We

have found that this process generates sub-bactericidal concentrations of allicin, which is

complemented or synergised by the presence of other ingredients to form a biofilm-killing

preparation. This demonstrates how contingent explorations of natural product

antimicrobials are on material preparation and testing conditions. This result is also

interesting because allicin can be toxic in high concentrations (61). Reducing the amount of

allicin present but maintaining full activity against bacterial biofilms might have been key to

producing a safe topical treatment. Further work is needed to elucidate the exact

combination of natural products responsible for the anti-biofilm activity.

Contrary to many papers in the literature (45,62,63), we found that wine possessed very

limited planktonic antimicrobial activity, and could not kill *S. aureus* in biofilms. Despite this,

its absence from the full remedy causes a large drop in activity against *S. aureus* biofilms.

Taken together, these observations suggest that the role of wine in the full recipe may be

more to do with its physical properties, perhaps its ethanol content and/or low pH. Ethanol is

a well-known extraction solvent and may allow better extraction of compounds from the

plant matter (64), or allow better diffusion through biofilms, while the lower pH may activate

pH-dependent compounds. The answer may lie in the combination of these properties, as the

activity of wine reported in the literature cannot be attributed exclusively to one aspect of it:

the physicochemical environment plays a large role in the activity of wine (65). Although pH

and ethanol concentrations corresponding to those found in various wines have minimal

effects on pathogens, without them the compounds within the wine have reduced activity

(65).

Similarly, individual preparations of onion and bile, prepared as they would be for Bald's

eyesalve, possessed no planktonic antibacterial activity. This is contrary to the literature that

has shown these ingredients to have activity against various strains (42,66). The most likely

cause for this disparity is the difference in the preparation or the dilution effect when

combined with other ingredients. However, the removal of onion or bile significantly reduced

the anti-biofilm activity against *S. aureus*, indicating the importance of their presence within

the whole remedy. They may be providing additional compounds necessary for the full killing

effect in biofilms or aiding penetration/activity of antibacterial compounds from other

ingredients.

Here, we have shown for the first time that Bald's eyesalve has anti-biofilm activity against

several clinically relevant bacteria, and we confirm our earlier work which concluded that this

activity is dependent on combining all the ingredients. Interestingly, statistical analysis of the

surviving portion of Bald's Leechbook (London, British Library, Royal MS 12 D XVII) showed

that garlic is combined with a second *Allium* species significantly more often than would be

expected given the frequency of use of either garlic or other *Allium* species across the book if

ingredients were combined randomly (67). Perhaps the patterns of ingredient combinations

used by pre-modern physicians do, in at least some cases, reflect a requirement for

combinatorial activity of several natural products to produce an efficacious antimicrobial

preparation.

Research into natural products is often focussed on isolating single compounds, however,

here we provide evidence that in doing so potent anti-biofilm mixtures can be overlooked.

Viewing natural products in this way has the potential to open a vast new source of

antimicrobials that can overcome the inherent difficulties of treating biofilm infections. In a

companion manuscript (22), we describe the low potential of Bald's eyesalve for producing

irritation or impeding wound healing. Future work will determine whether our results

translate into a candidate natural product cocktail for incorporation into wound ointments or

dressings. At present, we conclude by re-stating the exciting potential for pre-modern

European medical texts to contain antibacterial preparations of clinical interest.

Acknowledgements

We thank Mathew Diggle, Kim Hardie, Ruth Massey and Tim Sloan for bacterial strains; Meera

Unnikrishnan for helpful comments on an early draft of the manuscript; Callum Parsons,

Colman Ó Cathail, Jason Millington and Thorulf Vargsen for pilot work; Callum Parsons, Jenny

Littler, Shanjini Subhaskaran, Jason Millington, Thorulf Vargsen, Esther Sweeney and Colman

O'Cathail for preparation of the eyesalve batches summarised in Figure 1; and our colleagues

who helped initiate work on this project – Steve Diggle, Aled Roberts, Kendra Rumbaugh and

Rebecca Gabrilska. We also thank Cerith Harries, Caroline Stewart and the University of

Warwick Media Preparation team for preparing the media used for this work. This work was

supported by a Diabetes UK Project Grant to FH (ref. 17/0005690) and Jessica Furner-Pardoe

is funded by the MRC Doctoral Training Partnership [grant number MR/N014294/1].

Supplementary Figures

Figure S1. Schematic of the process to generate Bald's eyesalve and various batch variations.

Figure S2. Confirmation of allicin in Bald's eyesalve. All chromatograms were run against a 0-95% methanol

gradient, with a flow rate of 1 ml·min⁻¹ on HPLC Aligent 1200 series, at 210 nm UV wavelength. A) Allicin

calibration curve. External allicin standards, serially diluted in water were run and peak area was measured. A

linear relationship was found between HPLC peak area and allicin concentrations between 8 - 750 μg·ml⁻¹ with

a correlation coefficient R^2 value > 0.99. The mid-range of the curve (8-512 μg·ml⁻¹) has 2 repeats. B) Exemplary

allicin external standard (128 μg·ml⁻¹) chromatogram, suspected allicin peak has a retention time of approx. 15

minutes. C-D) Bald's eyesalve fresh batch, ESO 31 without (C) and with (D) an additional 100 μg·ml⁻¹ of allicin

standard; the differences in peak area are indicated. The highlighted region is the allicin peak.

Figure S3. Bald's eyesalve (ESO) anti-biofilm activity. Mature biofilms of various isolates were grown in a model

of a soft tissue wound, then treated with either 0.5 vols sterile water (control, CT) or 0.5 vols of ESO for 24h

before recovering bacteria for CFU counts (n = 3-5 replicates per treatment). The dashed line represents the

limit of detection by plating. Data could not be transformed to fit assumptions of parametric tests, therefore,

Kruskall-Wallis tests were used to determined that the CFU recovered from ESO-treated wounds was not

significantly different from control wounds for *P. aeruginosa* PA14 (X^2 = 6.88, df = 3, *p* value = 0.076), *S.*

maltophilia (X^2 = 7.63, df = 3, *p* value = 0.54) or *E. cloacae* (X^2 = 7.74, df = 3, *p* value = 0.051). Raw data and R

scripts are supplied in the Data Supplement.

References

1. Teillant A, Gandra S, Barter D, Morgan DJ, Laxminarayan R. Potential burden of

antibiotic resistance on surgery and cancer chemotherapy antibiotic prophylaxis in

the USA: A literature review and modelling study. Lancet Infect Dis.

2015;15((12)):1429–37.

2. O'Neill J. Tackling drug-resistant infections globally : final report and

recommendations the review on antimicrobial resistance. 2016.

3. Wright GD. Opportunities for natural products in 21st century antibiotic discovery.

Nat Prod Rep [Internet]. 2017 Jul 6 [cited 2019 May 21];34(7):694–701. Available

from: http://xlink.rsc.org/?DOI=C7NP00019G

4. Guest JF, Ayoub N, Mcilwraith T, Uchegbu I, Gerrish A, Weidlich D, et al. Health

economic burden that wounds impose on the National Health Service in the UK. Natl

Heal Serv UK BMJ Open [Internet]. 2015 [cited 2019 Nov 4];5:9283. Available from:

http://dx.doi.org/10.1136/bmjopen-2015-009283

5. Guest JF, Ayoub N, McIlwraith T, Uchegbu I, Gerrish A, Weidlich D, et al. Health

economic burden that different wound types impose on the UK's National Health

Service. Int Wound J [Internet]. 2017 Apr 1 [cited 2019 Mar 18];14(2):322–30.

Available from: http://doi.wiley.com/10.1111/iwj.12603

6. Penesyan A, Gillings M, Paulsen IT. Antibiotic Discovery: Combatting Bacterial

Resistance in Cells and in Biofilm Communities. Molecules [Internet]. 2015 [cited

2019 Nov 4];20:5286–98. Available from: www.mdpi.com/journal/molecules

7. Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A. The Calgary Biofilm Device:

New Technology for Rapid Determination of Antibiotic Susceptibilities of Bacterial

Biofilms. Vol. 37, Journal of Clinical Microbiology. 1999.

8.   Høiby N, Bjarnsholt T, Moser C, Bassi GL, Coenye T, Donelli G, et al. ESCMID guideline

for the diagnosis and treatment of biofilm infections 2014. Clin Microbiol Infect

[Internet]. 2015 [cited 2020 Feb 26];21((S1)):1–25. Available from:

http://dx.doi.org/10.1016/j.cmi.2014.10.024

9.   Tacconelli E, Carrara E, Savoldi A, Kattula D, Burkert F. Global priority list of antibiotic-

resistant bacteria to guide research, discovery, and development of new antibiotics

[Internet]. 2019 [cited 2019 Nov 24]. Available from:

http://www.cdc.gov/drugresistance/threat-report-2013/

10.   Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens.

Expert Rev Anti Infect Ther. 2013;11(3):297–308.

11.   WHO. Who Publishes List of Bacteria for Which New Antibiotics Are Urgently Needed

[Internet]. Saudi Medical Journal. 2017 [cited 2020 Feb 18]. Available from:

https://www.who.int/news-room/detail/27-02-2017-who-publishes-list-of-bacteria-

for-which-new-antibiotics-are-urgently-needed

12.   Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: A common cause of

persistent infections. Science (80-). 1999;284(5418):1318–22.

13.   Benkeblia N. Antimicrobial activity of essential oil extracts of various onions (Allium

cepa) and garlic (Allium sativum). LWT - Food Sci Technol. 2004;37(2):263–8.

14.   Chen C, Liu CH, Cai J, Zhang W, Qi WL, Wang Z, et al. Broad-spectrum antimicrobial

activity, chemical composition and mechanism of action of garlic (Allium sativum)

extracts. Food Control. 2018;86:117–25.

15.   Hooper SJ, Lewis MAO, Wilson MJ, Williams DW. Antimicrobial activity of Citrox[TM]

bioflavonoid preparations against oral microorganisms. Br Dent J. 2011 Jan

8;210(1):1–5.

16.   Elfawal MA, Towler MJ, Reich NG, Golenbock D, Weathers PJ, Rich SM. Dried Whole

Plant Artemisia annua as an Antimalarial Therapy. PLoS One. 2012 Dec 20;7(12).

17.   Elfawal MA, Towler MJ, Reich NG, Weathers PJ, Rich SM. Dried whole-plant artemisia

annua slows evolution of malaria drug resistance and overcomes resistance to

artemisinin. Proc Natl Acad Sci U S A. 2015 Jan 20;112(3):821–6.

18.   Kubicek-Sutherland JZ, Heithoff DM, Ersoy SC, Shimp WR, House JK, Marth JD, et al.

Host-dependent Induction of Transient Antibiotic Resistance: A Prelude to Treatment

Failure. EBioMedicine [Internet]. 2015 [cited 2019 Nov 4];2:1169–78. Available from:

http://dx.doi.org/10.1016/j.ebiom.2015.08.012

19.   Ersoy SC, Heithoff DM, Barnes L, Tripp GK, House JK, Marth JD, et al. Correcting a

Fundamental Flaw in the Paradigm for Antimicrobial Susceptibility Testing.

EBioMedicine [Internet]. 2017 Jun [cited 2019 Nov 4];20:173–81. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28579300

20.   Harrison F, Roberts AEL, Gabrilska R, Rumbaugh KP, Lee C, Diggle SP. A 1,000-year-old

antimicrobial remedy with antistaphylococcal activity. MBio. 2015;6(4):1–7.

21.   Fuchs AL, Weaver AJ, Tripet BP, Ammons MCB, Teintze M, Copié V. Characterization

of the antibacterial activity of Bald's eyesalve against drug resistant Staphylococcus

aureus and Pseudomonas aeruginosa. Becker K, editor. PLoS One [Internet]. 2018 Nov

28 [cited 2019 May 28];13(11):e0208108. Available from:

http://dx.plos.org/10.1371/journal.pone.0208108

22.   Anonye BO, Nweke V, Furner-Pardoe J, Gabrilska R, Rafiq A, Ukachukwu F, et al. The

safety profile of Bald's eyesalve for the treatment of bacterial infections. BioRXiv.

2020;

23.   Werthén M, Henriksson L, Jensen PØ, Sternberg C, Givskov M, Bjarnsholt T. An in vitro

692   model of bacterial infections in wounds and other soft tissues. Apmis.

693   2010;118(2):156–64.

694   24.   Mondy N, Naudin A, Christides JP, Mandon N, Auger J. Comparison of GC-MS and

695   HPLC for the analysis of Allium volatiles. Chromatographia. 2001;53(SPEC. ISS):356–

696   60.

697   25.   Fox J, Weisberg S. An R Companion to Applied Regression. 2011.

698   26.   Lenth R V. Lsmeans: R Package. 2017.

699   27.   Hothorn T, Bretz F, Westfall P. Simultaneous Inference in General Parametric Models.

700   Biometrical J. 2008;50:346–63.

701   28.   Dinno DHO and PW and A. FSA: Fisheries Stock Analysis [Internet]. 2020. Available

702   from: https://github.com/droglenc/FSA

703   29.   Bierbaumer P, Sauer H, Klug HW, Krischke U. Dictionary of Old English Plant Names.

704   [Internet]. Toronto: Dictionary of Old English Project 2009. 2009 [cited 2019 Dec 18].

705   Available from: http://oldenglish-plantnames.org

706   30.   diPaolo Healey A, Price Wilkin J, Xiang X. Dictionary of Old English Web Corpus

707   [Internet]. Toronto: Dictionary of Old English Project 2009. 2009 [cited 2019 Dec 18].

708   Available from: https://tapor.library.utoronto.ca/doecorpus/

709   31.   Brenner VC, Sherris JC. Influence of different media and bloods on results of diffusion

710   antibiotic susceptibility tests. Antimicrob Agents Chemother. 1972;1(2):116–22.

711   32.   Ankri S, Mirelman D. Antimicrobial properties of allicin from garlic. Microbes Infect.

712   1999;1(2):125–9.

713   33.   Fujisawa H, Watanabe K, Suma K, Origuchi K, Matsufuji H, Seki T, et al. Antibacterial

714   Potential of Garlic-Derived Allicin and Its Cancellation by Sulfhydryl Compounds.

Biosci Biotechnol Biochem [Internet]. 2009 [cited 2019 Sep 13];73(9):1948–55.

Available from:

https://www.tandfonline.com/action/journalInformation?journalCode=tbbb20

34.   Marchese A, Barbieri R, Sanches-Silva A, Daglia M, Nabavi SF, Jafari NJ, et al.

Antifungal and antibacterial activities of allicin: A review. Trends Food Sci Technol

[Internet]. 2016 Jun 1 [cited 2019 Sep 17];52:49–56. Available from:

https://www.sciencedirect.com/science/article/pii/S0924224416300073

35.   Salam AM, Quave CL. Opportunities for plant natural products in infection control.

Vol. 45, Current Opinion in Microbiology. Elsevier Ltd; 2018. p. 189–94.

36.   Ta CAK, Arnason JT. Mini review of phytochemicals and plant taxa with activity as

microbial biofilm and quorum sensing inhibitors. Vol. 21, Molecules. MDPI AG; 2016.

37.   Lanzotti V, Bonanomi G, Scala F. What makes Allium species effective against

pathogenic microbes? Phytochem Rev. 2013;12(4):751–72.

38.   Wolcott RD, Rhoads DD, Bennett ME, Wolcott BM, Gogokhia L, Costerton JW, et al.

Chronic wounds and the medical biofilm paradigm. J Wound Care. 2010 Sep

29;19(2):45–53.

39.   National Institute for Health and Care Excellence. NICE guideline NG19: Diabetic foot

problems: prevention and management. [Internet]. NICE; 2015 [cited 2020 Feb 26].

Available from: https://www.nice.org.uk/guidance/ng19

40.   Lanzotti V, Bonanomi G, Scala F. What makes Allium species effective against

pathogenic microbes? Vol. 12, Phytochemistry Reviews. Kluwer Academic Publishers;

2013. p. 751–72.

41.   Azu N, Onyeagba R. Antimicrobial Properties Of Extracts Of Allium cepa (Onions) And

Zingiber officinale (Ginger) On Escherichia coli, Salmonella typhi And Bacillus subtilis.

Internet J Trop Med. 2007;3(2):1–10.

42.   Begley M, Gahan CGM, Hill C. The interaction between bacteria and bile. FEMS

Microbiol Rev. 2005;29(4):625–51.

43.   Sung JY, Costerton JW, Shaffer EA. Defense system in the biliary tract against bacterial

infection. Dig Dis Sci. 1992;37(5):689–96.

44.   Sannasiddappa TH, Lund PA, Clarke SR. In vitro antibacterial activity of unconjugated

and conjugated bile salts on Staphylococcus aureus. Front Microbiol. 2017;8(AUG):1–

11.

45.   Rodríguez Vaquero MJ, Alberto MR, Manca de Nadra MC. Influence of phenolic

compounds from wines on the growth of Listeria monocytogenes. Food Control.

2007;18(5):587–93.

46.   Rodríguez Vaquero MJ, Alberto MR, de Nadra MCM. Antibacterial effect of phenolic

compounds from different wines. Food Control. 2007;18(2):93–101.

47.   Carneiro A, Couto JA, Mena C, Queiroz J, Hogg T. Activity of wine against

Campylobacter jejuni. Food Control [Internet]. 2008 Aug 1 [cited 2019 Aug

22];19(8):800–5. Available from:

https://www.sciencedirect.com/science/article/pii/S0956713507001661

48.   Müller A, Eller J, Albrecht F, Prochnow P, Kuhlmann K, Bandow JE, et al. Allicin induces

thiol stress in bacteria through S-allylmercapto modification of protein cysteines. J

Biol Chem [Internet]. 2016 [cited 2019 Aug 29];291(22):11477–90. Available from:

http://www.jbc.org/

49.   Imani Rad H, Arzanlou M, Ranjbar Omid M, Ravaji S, Peeri Doghaheh H. Effect of

culture media on chemical stability and antibacterial activity of allicin. J Funct Foods

[Internet]. 2017 Jan 1 [cited 2019 Mar 19];28:321–5. Available from:

https://www.sciencedirect.com/science/article/pii/S1756464616303450?via%3Dihub

50.   Bose S, Laha B, Banerjee S. Quantification of allicin by high performance liquid

chromatography-ultraviolet analysis with effect of post-ultrasonic sound and

microwave radiation on fresh garlic cloves. Pharmacogn Mag [Internet]. 2014 Apr

[cited 2019 May 30];10(Suppl 2):S288-93. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24991105

51.   Fujisawa H, Suma K, Origuchi K, Kumagai H, Seki T, Ariga T. Biological and Chemical

Stability of Garlic-Derived Allicin. J Agric Food Chem [Internet]. 2008 [cited 2019 Sep

13];56:4229–35. Available from: https://pubs.acs.org/sharingguidelines

52.   Fujisawa H, Suma K, Origuchi K, Seki T, Ariga T. Thermostability of Allicin Determined

by Chemical and Biological Assays. Biosci Biotechnol Biosci [Internet]. 2008 [cited

2019 Sep 13];72(11):2877–83. Available from:

https://www.tandfonline.com/action/journalInformation?journalCode=tbbb20

53.   Bocchini P, Andalò C, Pozzi R, Galletti GC, Antonelli A. Determination of diallyl

thiosulfinate (allicin) in garlic (Allium sativum L.) by high-performance liquid

chromatography with a post-column photochemical reactor. Anal Chim Acta.

2001;441(1):37–43.

54.   Li X, Hu K. Quantitative NMR Studies of Multiple Compound Mixtures. Annu Reports

NMR Spectrosc. 2017;90:85–143.

55.   Shobana S, Vidhya VG, Ramya M. Antibacterial Activity of Garlic Varieties

(Ophioscordon and Sativum) on Enteric Pathogens. Curr Res J Biol Sci. 2009;1(3):123–

6.

56.   Eagling D, Sterling S. A cholesterol-lowering extract from GARLIC A report for the

Rural Industries Research and Development Corporation [Internet]. 2000 [cited 2020

Feb 20]. Available from: http://www.rirdc.gov.au

57.   Bhandari SR, Yoon MK, Kwak JH. Contents of phytochemical constituents and

antioxidant activity of 19 garlic (Allium sativum L.) parental lines and cultivars. Hortic

Environ Biotechnol. 2014 May 10;55(2):138–47.

58.   Beato VM, Orgaz F, Mansilla F, Montaño A. Changes in Phenolic Compounds in Garlic

(Allium sativum L.) Owing to the Cultivar and Location of Growth. Plant Foods Hum

Nutr [Internet]. 2011 Sep [cited 2020 Feb 20];66(3):218–23. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21667145

59.   Fotopoulos V, Goulas V, Alexandros Petropoulos S, Cheng Z, Hayat S, Ahmad H, et al.

Garlic, from Remedy to Stimulant: Evaluation of Antifungal Potential Reveals Diversity

in Phytoalexin Allicin Content among Garlic Cultivars; Allicin Containing Aqueous

Garlic Extracts Trigger Antioxidants in Cucumber. 2016 [cited 2020 Feb 19]; Available

from: www.frontiersin.org

60.   Sawer IK, Berry MI, Ford JL. Effect of medium composition, agitation and the presence

of EDTA on the antimicrobial activity of cryptolepine. Lett Appl Microbiol.

1997;25(3):207–11.

61.   Gruhlke MCH, Nicco C, Batteux F, Slusarenko AJ. The effects of allicin, a reactive sulfur

species from garlic, on a selection of mammalian cell lines. Antioxidants. 2017 Mar

1;6(1).

62.   Friedman M. Antibacterial, Antiviral, and Antifungal Properties of Wines and Winery

Byproducts in Relation to Their Flavonoid Content. J Agric Food Chem [Internet]. 2014

[cited 2019 Dec 17];62(26):6025–42. Available from:

https://pubs.acs.org/sharingguidelines

63.   Møretrø T, Daeschel MA. Wine is Bactericidal to Foodborne Pathogens. J Food Sci

[Internet]. 2006 May 31 [cited 2019 Aug 22];69(9):M251–7. Available from:

http://doi.wiley.com/10.1111/j.1365-2621.2004.tb09938.x

64. Zhang QW, Lin LG, Ye WC. Techniques for extraction and isolation of natural products:

A comprehensive review. Vol. 13, Chinese Medicine. BioMed Central Ltd.; 2018.

65. Boban N, Tonkic M, Budimir D, Modun D, Sutlovic D, Punda-Polic V, et al.

Antimicrobial effects of wine: Separating the role of polyphenols, pH, Ethanol, and

other wine components. J Food Sci. 2010;75(5):322–6.

66. Ahiabor C, Gordon A, Ayittey K, Agyare R. In vitro assessment of antibacterial activity

of crude extracts of onion (Allium cepa L .) and shallot (Allium aescalonicum L .) on

isolates of Escherichia coli (ATCC 25922) , Staphylococcus aureus (ATCC 25923) ,

and Salmonella typhi (ATCC 19430). Int J Appl Res. 2016;2(5):1029–32.

67. Harrison F, Connelly E. Could Medieval Medicine Help the Fight Against Antimicrobial

Resistance? In: Jones C, Kostick C, Oschema K, editors. Making the Medieval Relevant

[Internet]. Berlin, Boston: De Gruyter; 2019. p. 113–34. Available from:

http://www.degruyter.com/view/books/9783110546316/9783110546316-

005/9783110546316-005.xml

827