**Strain background, species frequency and environmental conditions are**

**important in determining population dynamics and species co-existence between**

***Pseudomonas aeruginosa* and *Staphylococcus aureus***

Selina Niggli,[a]# Rolf Kümmerli[a]

[a] Department of Quantitative Biomedicine, University of Zürich, Winterthurerstrasse 190,

8057 Zürich, Switzerland

# Address correspondence to Selina Niggli, selina.niggli@uzh.ch

**Running title:** How ecology affects competition between two pathogens

**Keywords:** opportunistic human pathogens, interspecies interactions, polymicrobial

infections, ecological factors, community dynamics

**Abstract**

Bacterial communities in the environment and in infections are typically diverse, yet we know little about the factors that determine interspecies interactions. Here, we apply concepts from ecological theory to understand how biotic and abiotic factors affect interaction patterns between the two opportunistic human pathogens *Pseudomonas aeruginosa* and *Staphyloccocus aureus*, which often co-occur in polymicrobial infections. Specifically, we conducted a series of short- and long-term competition experiments between *P. aeruginosa* PAO1 (as our reference strain) and three different *S. aureus* strains (Cowan I, 6850, JE2) at three starting frequencies and under three environmental (culturing) conditions. We found that the competitive ability of *P. aeruginosa* strongly depended on the strain background of *S. aureus*, whereby *P. aeruginosa* dominated against Cowan I and 6850, but not against JE2. In the latter case, both species could end up as winners depending on conditions. Specifically, we observed strong frequency-dependent fitness patterns, including positive frequency dependence, where *P. aeruginosa* could dominate JE2 only when common, but not when rare. Finally, changes in environmental (culturing) conditions fundamentally altered the competitive balance between the two species, in a way that *P. aeruginosa* dominance increased when moving from shaken to static environments. Altogether, our results highlight that ecological details can have profound effects on the competitive dynamics between co-infecting pathogens, and determine whether two species can co-exist or invade each others' populations from rare. Moreover, our findings might parallel certain dynamics observed in chronic polymicrobial infections.

**Importance**

Bacterial infections are frequently caused by more than one species and such polymicrobial infections are often considered more virulent and more difficult to treat than the respective monospecies infections. *Pseudomonas aeruginosa* and *Staphyloccocus aureus* are among the most important pathogens in polymicrobial infections and their co-occurrence is linked to worse disease outcome. There is great interest in understanding how these two species interact with each other and what the consequences for the host are. While previous studies have mainly looked at molecular mechanisms implicated in interactions between *P. aeruginosa* and *S. aureus*, here we show that ecological factors such as strain background, species frequency and environmental conditions are important elements determining population dynamics and species co-existence patterns. We propose that the uncovered principles may also play major roles in infections, and therefore proclaim that an integrative approach combining molecular and ecological aspects is required to fully understand polymicrobial infections.

**Introduction**

Bacteria typically live in complex multi-species communities in the environment and associated with host organisms (1–3). The same holds true in the case of disease, as it is increasingly recognized that a majority of bacterial infections are polymicrobial, meaning that they are caused by more than one bacterial species (4, 5). There is great interest in understanding how bacteria interact and how interactions affect a community and the associated hosts (6–8). At the mechanistic level, a multitude of ways have been unraveled through which bacterial species can interact, with mechanisms including cross-feeding, quorum sensing-based signaling, toxin-mediated interference and physical interactions via contact-dependent systems (e.g. type VI secretion system) (9–11). In the context of disease, a key question is how interactions affect species successions in chronic infections and whether multispecies infections are more virulent and more difficult to treat than the respective monospecies infections, as it is commonly assumed (5, 12–14).

Studying interactions between the two opportunistic human pathogens *Pseudomonas aeruginosa* (PA) and *Staphylococcus aureus* (SA) has emerged as a popular and relevant model system (15–17). The reason for this is that the two species often co-occur in infections, including cystic fibrosis (CF) lung and wound infections (18–20). Results from laboratory experiments suggest that PA is the superior species, suppressing growth of SA (21–23) and indeed, PA seems to be a well-equipped competitor. For example, it has been shown that 4-hydroxy-2-heptylquinoline N-oxide (HQNO) released by PA inhibits the electron transport chain of SA and induces the

formation of small colony variants (SCVs) (22, 24). Furthermore, the PA endopeptidase

LasA is capable of lysing SA cells, a process that releases iron into the environment,

potentially providing a direct benefit to PA (21, 25). While it was observed that

experimental co-infections of PA and SA seem to be more virulent than the respective

monospecies infections (26, 27), evolutionary studies revealed that PA can adapt to the

presence of SA (28) and become more benign in the context of chronic co-infections

(29, 30). Moreover, of clinical relevance is the observation that PA and SA exhibited

increased antibiotic resistance or tolerance when co-cultured compared to being

cultured alone (31–33).

In this study, we follow a complementary approach to examine how biotic and abiotic

ecological factors influence interactions between PA and SA. Previous work has

primarily focused on the molecular mechanisms driving interactions between specific

strain pairs under defined laboratory conditions. Here we hypothesize that not only

molecular mechanisms, but also ecological factors will have a major impact on species

interactions, particularly on community composition and temporal dynamics between

species. To test our predictions, we used PA strain PAO1 as our focal strain and asked

how (competitive) interactions with SA vary when manipulating: (1) the genetic

background of SA; (2) the frequency of SA in competition with PA; and

(3) environmental (culturing) conditions.

To vary the genetic background of SA, we competed PA against the three different SA

strains Cowan I, 6850 and JE2. These strains fundamentally differ in several

characteristics (Table 1). Cowan I is a methicillin-sensitive SA strain (MSSA), which is

highly invasive towards host cells, non-cytotoxic and defective in the accessory gene

regulator (agr) quorum-sensing system (34). 6850 is another MSSA strain, which is

highly invasive, cytotoxic and haemolytic (35–37). Finally, JE2 is a methicillin-resistant

(MRSA) USA300 strain, which is highly virulent, cytotoxic and hemolytic (38, 39). Given

the tremendous differences between these SA strains, we expect PA performance in

competition with SA to vary substantially.

To manipulate strain frequency, we competed PA against SA at three different starting

frequencies (1:9 ; 1:1 ; 9:1). Frequency-dependent fitness effects occur in many

microbiological systems (40–43). A common pattern is that species have a relative

fitness advantage when rare in a population, but not when common (so-called negative

frequency dependence). This phenomenon can lead to stable co-existence of

competitors. On the other hand, the fitness of a species can also be positive frequency

dependent, which means that a species is dominant when common in the community,

but not when rare. An important consequence of this pattern is that initially rare species

cannot invade an established population.

To manipulate environmental conditions, we changed simple parameters of our

culturing conditions. First, we compared the performance of PA against SA strains in

shaken liquid vs. viscous medium. Increased environmental viscosity has been shown

to increase spatial structure, thereby decreasing strain interaction rates (44–46).

Second, we compared the performance of PA against SA in shaken vs. static

environments. While static conditions also reduce strain mixing, it further leads to a

more heterogeneous environment characterized by gradients from the aerated air-liquid

interface down to the microoxic bottom of a culture (47, 48).

In a first set of experiments, we assessed the growth performance of all strains in

monoculture under the three different environmental culturing conditions used. Basic

growth differences between strains could induce frequency shifts in co-cultures even in

the absence of direct interactions. We then performed high-throughput 24 hours batch

culture competition experiments between PA and SA using a full-factorial design. All

three species combinations were competed at all three starting frequencies under all

three environmental conditions (see Figure 1 for an illustration of the workflow). Finally,

we followed the temporal dynamics between PA and SA over five days to assess

whether results from 24 hours competitions are predictive for more long-term dynamics

between species and whether species coexistence is possible. (A copy of this

manuscript has been posted on bioRxiv doi: https://doi.org/10.1101/2020.04.21.052670)

## Results

### PA grows better than SA in monoculture

We used tryptic soy broth (TSB) as the standard medium for all our assays. In this medium, we found that the number of doublings varied significantly among strains during a 24 hours growth cycle under all conditions tested (ANOVA, shaken: $F_{3,20} = 10.71$, $P = 0.0002$; viscous: $F_{3,20} = 4.12$, $P = 0.0199$; static: $F_{3,20} = 20.75$, $P < 0.0001$, Figure 2 and see Table S1 for the full statistical analysis). Under shaken conditions, PA had the highest number of doublings (20.6 ± 0.63, mean ± SD), followed by SA strains 6850 (18.9 ± 1.16), Cowan I (18.1 ± 0.41) and JE2 (18.0 ± 1.18). While PA grew significantly better than all SA strains, the number of doublings did not differ between the three SA strains (TukeyHSD pairwise comparisons: Cowan I vs. 6850, $P_{adj} = 0.3673$; 6850 vs. JE2, $P_{adj} = 0.3096$; Cowan I vs. JE2, $P_{adj} = 0.9994$). Due to its moderate growth advantage, PA is expected to slightly increase in frequency in competition with SA strains under shaken conditions, even in the absence of any direct species interactions.

Under viscous conditions, PA had the highest number of doublings as well (17.2 ± 1.02), followed by 6850 (17.0 ± 0.76), JE2 (16.3 ± 0.52) and Cowan I (15.6 ± 1.06). However, differences in the number of doublings were only significant between PA and Cowan I and between Cowan I and 6850 (TukeyHSD pairwise comparisons: PA vs. Cowan I, $P_{adj} = 0.0265$; Cowan I vs. 6850, $P_{adj} = 0.0496$). Thus, based on growth rate differences alone, one would expect PA to increase in frequency in competition with Cowan I but not in competition with the other two SA strains.

Under static conditions, PA again showed the highest number of doublings (18.6 ± 0.64)

followed by 6850 (16.5 ± 0.45), JE2 (15.9 ± 0.96) and Cowan I (15.1 ± 1.05). PA grew

significantly faster than all the three SA strains (TukeyHSD pairwise comparisons: PA

vs. Cowan I, $P_{adj} < 0.0001$; PA vs. 6850, $P_{adj} = 0.0009$; PA vs. JE2, $P_{adj} < 0.0001$), and

one would therefore expect PA to substantially increase in frequency against all three

SA strains under static conditions.

**Genetic background, strain frequency and environmental factors all influence**

**competition outcomes**

The full-factorial design allowed us to simultaneously analyze the impact of SA strain

genetic background, starting frequency, and culturing condition on the competitive

outcomes between PA and SA strains. Our linear statistical model yielded a significant

triple interaction between the three manipulated factors (strain genetic background,

starting frequency and culturing condition; ANCOVA: $F_{4,509} = 3.41$, $P = 0.0091$). While

this shows that all three manipulated factors influence the competitive outcomes

between PA and SA in complex ways, the triple interaction makes it difficult to tease

apart the various effects. The statistical procedure for such cases is to split the model

into sub-models. We followed this approach by first analyzing separate models for each

of the three environmental conditions (shaken, viscous, static), and then split models

according to SA strain background to test for differences between environmental

conditions.

**The competitive ability of PA depends on the SA strain genetic background**

Under all three environmental conditions, we found that the relative fitness of PA

significantly depended on the SA strain background (ANCOVA, shaken: $F_{2,170} = 90.87$,

$P < 0.0001$; viscous: $F_{2,168} = 116.76$, $P < 0.0001$; static: $F_{2,170} = 56.52$, $P < 0.0001$;

Figure 3). Against Cowan I (Figure 3, column 1), we noted that PA consistently won the

competitions across all starting frequencies and culturing conditions. SA strain 6850

(Figure 3, column 2) turned out to be more competitive than Cowan I under shaken

conditions ($t_{176} = -6.74$, $P < 0.0001$), while it lost similarly against PA under viscous and

static conditions (viscous: $t_{174} = 0.78$, $P = 0.4350$; static: $t_{176} = -1.99$, $P = 0.0482$). In

contrast, JE2 was the most competitive SA strain in our panel (Figure 3, column 3),

performing significantly better than the other two SA strains under all conditions (see

Table S2 for the full statistical analysis), and outcompeted PA under shaken and

viscous conditions.

**The competitive ability of PA depends on its starting frequency in the population**

We found that the starting frequency of the two competitors had varying but always

significant effects on the competitive ability of PA (ANCOVA, shaken: $F_{1,170} = 52.81$,

$P < 0.0001$; viscous, interaction with strain background: $F_{2,168} = 10.05$, $P < 0.0001$;

static: $F_{1,170} = 162.32$, $P < 0.0001$). Under shaken conditions (Figure 3, row 1), we

observed that the relative fitness of PA increased when initially more common, thus

following a positive frequency-dependent pattern. Under viscous conditions (Figure 3,

row 2), the same positive frequency-dependent effect was only observed when PA

competed with JE2. In competition with Cowan I or 6850, we noted that the relative

211    fitness of PA peaked at intermediate starting frequencies. Under static conditions

212    (Figure 3, row 3), we observed a pattern opposite to the one seen under shaken

213    conditions for all strain pair combinations. The relative fitness of PA decreased when

214    initially more common, thus following a negative frequency-dependent pattern (see

215    Table S2 for the full statistical analysis).

**The competitive ability of PA is highest under static conditions**

Next, we compared the competitive outcomes among the different culturing conditions

(shaken, viscous and static) for each strain combination separately. For all strain

combinations, the culturing condition significantly affected competition outcomes

(ANCOVA, Cowan I: $F_{2,168} = 461.73$, $P < 0.0001$; 6850: $F_{2,167} = 428.16$, $P < 0.0001$; JE2:

$F_{2,168} = 199.95$, $P < 0.0001$). In competition with all three SA strains, we found that the

relative fitness of PA was significantly higher under static compared to shaken

conditions (Cowan I: $t_{174} = 19.99$, $P < 0.0001$; 6850: $t_{174} = 17.99$, $P < 0.0001$; JE2: $t_{174} =$

15.39, $P < 0.0001$). In contrast, there were no significant differences in the relative

fitness of PA between shaken and viscous conditions for Cowan I ($t_{174} = 0.91$, $P =$

0.3644) and JE2 ($t_{174} = 0.82$, $P = 0.4117$), while against 6850, PA was more competitive

under viscous than shaken conditions ($t_{174} = 3.53$, $P = 0.0005$) (see Table S2 for the full

statistical analysis).

**Temporal dynamics between PA and SA**

In a next experiment, we competed PA and SA strains over five days under shaken

conditions using the same three starting frequencies and by transferring cultures to

fresh medium every 24 hours. The aim of this experiment was to follow the more long-

term species dynamics and to assess whether stable coexistence between PA and SA

can arise.

In competition with Cowan I, we found PA to be the dominant species (Figure 4a). It

strongly increased in frequency already at day 1 under all starting frequencies and

almost completely outcompeted Cowan I by day 3 (i.e., Cowan I remained below

detection limit). Thus, we could not observe coexistence between PA and Cowan I. In

competition with 6850, we observed similar population dynamics (Figure 4b). PA

strongly increased in frequency from day 1 onwards at all starting frequencies and after

three days, the bacterial populations almost entirely consisted of PA. Only in 10 out of

30 populations, 6850 managed to persist at very low frequencies by day 5 (< 3% in nine

cases, and 13% in one case). In competition with JE2, we found community trajectories

that were strikingly different from the other two strain combinations (Figure 4c). First, we

observed that JE2 was a strong competitor, keeping PA at bay in many populations

during the first 24 hours of the experiment. Following day 1, community dynamics

followed positive frequency-dependent patterns. In all populations with intermediate or

high PA starting frequencies, PA became the dominant species, and SA was recovered

at low frequency in only a minority of populations by day 5 (3 out of 20 at < 10% of the

population). In stark contrast, in populations where PA was initially rare, it did not

increase in frequency, could not invade the SA populations and remained at a low

frequency (< 10%) throughout the 5 days.

**Discussion**

*Pseudomonas aeruginosa* (PA) and *Staphylococcus aureus* (SA) frequently occur

together in polymicrobial infections, where they cause severe host damage and lead to

increased morbidity and mortality in patients (14, 49, 50). Consequently, there is high

interest in understanding how PA and SA interact and how their interactions may

influence disease outcome (12, 15). While most previous studies have focused on

molecular aspects (51, 52), we here examined how a set of ecological factors affect

competitive interactions between the two species. Our study, carried out in an *in vitro*

batch culture system, revealed that: (i) the competitive ability of PA varied extensively

as a function of the genetic background of SA; (ii) there were strong frequency-

dependent fitness patterns, including positive-frequency dependent relationships where

PA could only dominate a particular SA strain (JE2) when common, but not when rare;

and (iii) changes in environmental (culturing) conditions fundamentally affected the

competitive balance between the two species. The key conclusion from our results is

that ecology matters, and that variation in biotic and abiotic factors affect interactions

between pathogenic bacterial species. This is most likely not only the case in *in vitro*

systems, but also in the context of polymicrobial infections.

PA has often been described as the dominant pathogen possibly displacing SA in

infections (21, 53–55). Our results support this view, as PA dominated over SA under

many conditions in 24 hours and 5-day competition experiments (Figures 3 and 4).

However, PA did not always emerge as the winner and its success significantly varied in

response to the genetic background of SA. Specifically, JE2 was the strongest

competitor, followed by 6850 and Cowan I. While differences in monoculture growth

performance can explain why PA dominates in many cases, they cannot explain the

variation in competitive abilities among SA strains, because the three SA strains grew

similarly under all conditions (Figure 2). PA vs. JE2 makes the strongest case for a

mismatch between growth performance in mono- vs. mixed cultures, as PA grew

significantly better than JE2 in monoculture under shaken conditions, but typically lost

the competition in mixed cultures in this environment (Figures 2 and 3). This suggests

that apart from resource competition via growth rate differences, other factors must

contribute to the competitive ability of SA strains towards PA. Such factors could include

interference mechanisms. For example, it is known that genetically different SA strains

widely differ in virulence among each other (56–58). Interestingly, we found that the

competitive ability of our three SA strains against PA correlated with their reported

virulence level in infections (34, 36, 39). This could indicate that factors important for SA

virulence (e.g. toxins or secreted enzymes) might also be involved in interactions with

competitor bacteria. For JE2 and related USA300 isolates, there are many genetic

determinants known to be important for their success as opportunistic human

pathogens (59). Among them are the cytotoxin Panton-Valentine leukocidin (PVL), the

arginine catabolic mobile element (ACME) and the phenol soluble modulins (PSMs)

(60). Derivatives of PSMs have previously been shown to exhibit inhibitory activity

against *Streptococcus pyogenes* (61). The authors of this work suggested that high

production of PSMs might not only benefit SA in host colonization, but also in

competition against co-infecting pathogens. Thus, it seems plausible that the USA300

derivative JE2 deploys a similar mechanism against PA in our competition experiments.

Strain 6850 showed intermediate competitiveness against PA. As Cowan I, 6850 is a

MSSA strain, but it is known to be more virulent than Cowan I and therefore likely

produces certain substances that could also be important in competition with PA (34,

36). Conversely, Cowan I is known to have a nonfunctional accessory gene regulator

(agr) quorum sensing system (34). The agr controls most virulence determinants in SA

(62). If virulence determinants also played a role in interspecies competition, then this

could explain why Cowan I turned out to be the least competitive SA strain against PA.

Important to note is that we only manipulated the SA but not the PA strain background,

so that we cannot draw conclusions on genotype-by-genotype interactions. Such

interactions are likely to play a role as evidenced by previous studies showing that

genetically diverse PA clinical isolates widely differ in their ability to inhibit SA (29, 63,

64). More recently, it has also been shown that SA clinical isolates vary in their

interactions with PA, from being highly sensitive to completely tolerant against PA-

mediated effects (65). One aim of our future work is to follow up these proposed

mechanistic leads and test some of the outlined hypotheses above in order to explain

differences in the competitive abilities between SA strains towards PA.

Another insight from our experiments is that the competitive ability of PA often

depended on its starting frequency in the population, and that the type of frequency-

dependent interactions (positive or negative) varied across environmental conditions

(Figure 3). Our purpose was to compare three experimentally defined starting

frequencies to mimic what is happening when a species is either rare (1:9), at parity

with its competitor (1:1) or dominant (9:1). Under natural conditions, including infections,

species frequencies could of course be more extreme, and invasion from rare could

start at species frequencies that would be below the detection limit of our methods.

In our experiments, we observed that under static conditions, the relative fitness of PA

declined when more common in the population, but PA still won at all frequencies. This

pattern is common for a highly dominant species that drives a competitor to extinction

(66). Its decline in relative fitness simply reflects the fact that the room for further

absolute frequency gains is reduced when a high frequency is already reached. In stark

contrast, under shaken conditions, we found that the relative fitness of PA increased

when it was more common in the population. Against Cowan I and 6850, this positive

frequency-dependent fitness pattern did not affect the long-term community dynamics

and PA won at all frequencies (Figure 4a+b). Against JE2, however, the 24 hours

competition data suggest that, in most cases, PA cannot invade populations when

initially rare and this is exactly what we observed in the long-term experiments: when its

initial frequency was below 10%, PA did not increase in frequency, while it always fixed

in the population or reached very high frequencies when initially occurring above 10%.

There were two additional interesting observations with regards to PA-JE2 long-term

dynamics. First, there were no major changes in PA frequency relative to JE2 during the

first 24 hours (compatible with the competition assay data in Figure 3), and clear

positive-frequency dependent patterns only emerged from 48 hours onwards. One

possible explanation for this pattern is that PA is initially naïve, but then senses and

mounts a more competitive response over time (67). Similarly, SA might also respond,

for example through increased formation of small colony variants (SCVs), which are

known to be induced by inhibitory exoproducts released by PA (22, 24). Second, one

replicate (starting frequency 1:1) did not follow the above rules: PA continuously

dropped in frequency until day 4 (11%) and then sharply increased to 93% on day 5.

This frequency "zigzag" pattern is an indicator of antagonistic co-evolution (68), where

the spread of a beneficial mutation in one species (SA) is followed by a counter-

adaptation in the competing species (PA). It therefore seems that such evolutionary

dynamics can already occur within relatively short periods of time. This finding supports

evidence from studies on clinical PA isolates, which showed patterns of adaptations

towards SA favoring co-existence between the two species over time (29, 30).

Our results further show that the competitive ability of PA is profoundly influenced by

environmental (culturing) conditions (Figure 3). The largest differences arose between

shaken and static culturing conditions with PA being most competitive in the latter

environment. PA is known to be metabolically versatile, it is motile and grows well under

microoxic conditions (69, 70). Static conditions introduce strong oxygen and nutrient

gradients, and our results from monoculture growth show that PA grows better under

these conditions than SA (Figure 2). This is certainly part of the reason why PA ends up

as the competition winner against all SA strains under static conditions. With regard to

medium viscosity, we initially hypothesized that increased spatial structure could temper

competitive interactions and favor species co-existence, as competitors are spatially

more segregated from each other (44, 71, 72). However, we found no support for this

hypothesis as the competitive ability of PA did not much differ between shaken and

viscous environments. While the spatial structure, introduced through the addition of

agar to the liquid growth medium, had significant effects on within-species social

interactions in other study systems (66, 73), it did not appreciably affect the between-

species interactions in our setup. One reason might be that the degree of spatial

structure introduced (0.2% agar in TSB) was simply not high enough to see an effect.

This could especially be true if toxins were involved in mediating interactions – small

molecules that can freely diffuse and target competitors that are not physically close-by.

We argue that our results, even though they stem from an *in vitro* system, could have at

least three important implications for our understanding of polymicrobial infections. First,

we show that the biological details of the strain background matter and determine who

is dominant in a co-infection and whether co-existence between species is possible.

Thus, we need to be careful not to overinterpret interaction data from a single PA-SA

strain pair and conclude that the specific details found apply to PA-SA interactions in

general. Second, there might be strong order effects, such that the species that infects

a host first cannot be invaded by a later arriving species. This scenario applied to the

interactions between PA and SA strain JE2, which were both unable to invade

populations of the other species from rare. Finally, local physiological conditions at the

infection site, like the degree of spatial structure or oxygen supply, can shift the

competitive balance between species. This suggests that infections at certain sites

might be more prone than others to polymicrobial infections, or to experience ecological

shifts from one pathogen to another. To sum up, we wish to reiterate our take home

message that the ecology of interactions between pathogens should receive more

attention and may explain so far unresolved aspects of polymicrobial infections.

**Materials and Methods**

**Bacterial strains, media and growth conditions**

We used the *Pseudomonas aeruginosa* (PA) strain PAO1 (74) as our PA reference

strain and the *Staphylococcus aureus* (SA) strains Cowan I, 6850 and JE2 for all

experiments (Table 1). To distinguish PA from SA strains, we used a variant of our PA

strain PAO1, which constitutively expresses the green fluorescent protein, from a single-

copy gene (attTn7::ptac-gfp), stably integrated in the chromosome (75, 76). We chose

the rich laboratory medium tryptic soy broth (TSB, Becton Dickinson) for all our

experiments, because it supports growth of all the strains used. Bacterial stocks were

prepared by mixing 50% of culture with 50% of a 85% glycerol solution and were stored

at -80 °C. For all experiments, overnight cultures were grown in 10 ml TSB in 50 ml

falcon tubes for ± 16 hours at 37 °C and 220 rpm with aeration. After centrifugation and

removal of the supernatant, we washed bacterial cells using 10 ml 0.8% NaCl solution

and adjusted the $OD_{600}$ (optical density at 600 nm) to obtain similar cell numbers per ml

for each strain. All media, buffer and washing solutions were sterilized by autoclaving at

121°C for 20 minutes and subsequently stored at room temperature in the dark. For all

experiments, blanks were used to ensure the sterility of the media during

experimentation.

**Calculating number of doublings for each strain in monoculture**

To assess the number of doublings of each strain in monoculture, we grew our strains in

TSB (or TSB + 0.2% agar, respectively) under the same conditions and using the same

starting $OD_{600}$ as for the competition experiments (see below). We serially diluted cells

at the start ($t_0$) and after 24 hours ($t_{24}$), and plated aliquots on TSB + 1.2% agar. The

plates were incubated overnight at 37 °C and colony- forming units (CFUs) counted for

both timepoints on the following day. We estimated the number of doublings (D) for

each strain as $D = (\ln(x_{24}/x_0))/\ln(2)$, where $x_0$ and $x_{24}$ are the initial and the final CFU/ml,

respectively (23). We performed this experiment two times with three replicates per

strain per experiment.

**Competition experiments**

To initiate competitions, we mixed PA and SA strain pairs at three different starting

frequencies (1:9, 1:1, 9:1) from washed and $OD_{600}$-adjusted overnight cultures (see

above). Competitions occurred in 24-well plates filled with 1.5 ml TSB per well. The

starting $OD_{600}$ of both mixed and monocultures was $10^{-5}$. Monocultures of each strain

served as controls in each experiment. We incubated plates for 24 hours at 37 °C under

three different culturing conditions: shaken (170 rpm), viscous (170 rpm with 0.2% agar

in TSB) and static. Prior and after the 24 hours competition period, we estimated the

actual strain frequencies for each replicate using flow cytometry. We performed four

independent experiments each featuring five replicates for each strain/starting

frequency/condition combination. A graphical representation of the competition workflow

is provided in Figure 1.

To follow community dynamics over time, we set up competitions in the same way as

described above. After the first 24 hours of competition, we diluted cultures 1:10,000

into fresh TSB medium. This process was repeated for five consecutive days. Strain

frequencies were assessed using flow cytometry prior and after each 24 hours

competition cycle. We carried out two independent experiments for each strain pair and

starting frequency combination with 5 replicates per strain pair and frequency.

**Flow cytometry to estimate relative species frequency**

We assessed the relative strain frequencies at the beginning and at the end of each

competition using a BD LSR II Fortessa flow cytometer (flow cytometry facility,

University of Zürich) and the FlowJo™ software (BD, Bioscience) for data analysis. As

our PA strain expresses a constitutive gfp tag, PA cells could unambiguously be

distinguished from the gfp-negative SA cells with a blue laser line (excitation at 488 nm)

and the FITC channel (emission: mirror 505 longpass, filter 530/30) (see supplementary

Figure 1). Cytometer Setup and Tracking settings of the instrument were used for each

experiment and the threshold of particle detection was set to 200 V (lowest possible

value). We diluted cultures appropriately in sterile-filtered 1x phosphate buffered saline

(PBS, Gibco, Thermo Fisher) and recorded 100,000 events with a low flow rate. The

following controls were used for data acquisition in every experiment: 1) PBS blank

samples (to estimate number of background counts of the flow cytometer), 2) untagged

monocultures (negative fluorescence control, used to set a fluorescence threshold in

FlowJo™) and 3) constitutive gfp-expressing monocultures (positive fluorescence

control, set to 100% gfp-positive cells). Using our fluorescence threshold, we extracted

the percentage of gfp-positive cells for each sample and scaled these values to the

positive fluorescence control. The resulting percentage corresponds to the frequency of

PA present in the respective replicate. Initial and final strain frequencies were used to

calculate the relative fitness (v) of the focal strain PA as $v = [a_t \times (1-a_0)] / [a_0 \times (1-a_t)]$,

where $a_0$ and $a_t$ are the initial and final frequencies of PA, respectively (43). We ln-

transformed all relative fitness values to obtain normally distributed residuals. Values of

$\ln(v) > 0$ or $\ln(v) < 0$ indicate whether the frequency of the focal strain PA increased (i.e.

PA won the competition) or decreased (i.e. PA lost competition) relative to its SA

competitor.

We know from previous experiments in our laboratory that due to the gfp tag, our PA

strain does have a slight fitness defect in competition with its untagged parental strain

($\ln(v) = -0.358 \pm 0.13$, mean ± 95% CI, see (66)). As we consistently used the same

gfp-tagged PA strain for all experiments in this study, results are fully comparable

among treatments.

To test whether flow cytometry counts (measuring all viable and non-viable cells)

correlate with CFUs (measuring only viable cells), we serially diluted and plated initial

and final strain frequencies from competitions performed under shaken conditions for all

three strain combinations on TSB + 1.2% agar. We compared the obtained CFUs with

the flow cytometry counts from the same samples and found strong positive correlations

for the strain frequency estimates between the two methods (see supplementary

figure 2). This means that flow cytometry adequately measures strain frequencies and

that the two methods (flow cytometry and CFU counts) yield similar results.

**Statistical analysis**

All statistical analyses were performed with R Studio version 3.6.1. We used analysis of

variance (ANOVA) and Tukey's HSD to compare number of doublings in monocultures

of PA and SA. To test whether the relative fitness of PA varies in response to the SA

strain genetic background, starting frequency and culturing conditions, we first built a

factorial analysis of co-variance (ANCOVA), with SA strain genetic background and

culturing conditions as factors and the starting frequency as covariate. We further

included 'experimental block' as an additional factor to account for variation between

experiments. This full model yielded a significant triple interaction between SA strain

genetic background, starting frequency and culturing condition. We therefore split the

full model into a set of ANCOVA sub-models, separated either by culturing condition

(shaken, viscous, static) or by SA strain genetic background (Cowan I, 6850, JE2). For

post-hoc pairwise comparisons between culturing conditions or SA strains in the sub-

models, we removed 'experimental block' as additional factor from the model. To test

whether PA relative fitness is significantly different from zero under a given

strain/starting frequency/condition combination, we performed one sample t-tests and

used the false discovery rate method to correct p-values for multiple comparisons (77).

To compare strain frequencies obtained by flow cytometry with those obtained by

plating (CFUs), we used Pearson correlation analysis. For all data sets, we consulted

Q-Q plots and results from the Shapiro-Wilk test to ensure that our residuals were

normally distributed. Summary tables for linear models and t-tests used to analyze

Figures 2 and 3 can be found in the supplemental material (Tables 1-3).

**Data availability**

All raw data sets have been deposited in the figshare repository (DOI will be provided

upon the acceptance of the manuscript).

**Conflict of Interest**

The authors declare no conflict of interest.

**Acknowledgements**

We thank Markus Huemer (University Hospital of Zürich) for providing *S. aureus* strains

and the flow cytometry facility (University of Zürich) for technical support and

maintenance of resources. This project has received funding from the European

Research Council (ERC) under the European Union's Horizon 2020 research and

innovation programme (grant agreement no. 681295) to RK. Illustration for Figure 1 was

created using BioRender (www.biorender.com).

**Author contributions**

S.N. and R.K. designed research, S.N. performed research, S.N. and R.K. analysed

data and wrote the paper.

**References**

1.  Stubbendieck RM, Vargas-Bautista C, Straight PD. 2016. Bacterial communities: Interactions to scale. Front Microbiol 7:1–19.

2.  Donaldson GP, Lee SM, Mazmanian SK. 2016. Gut biogeography of the bacterial microbiota. Nat Rev Microbiol 14:20–32.

3.  Human Microbiome Project Consortium. 2012. Structure, function and diversity of the healthy human microbiome. Nature 486:207–214.

4.  Brogden KA, Brogden KA, Guthmiller JM, Guthmiller JM, Taylor CE, Taylor CE, Guthmiller JM, Guthmiller JM. 2005. Human polymicrobial infections. Lancet 365:253–255.

5.  Peters BM, Jabra-Rizk MA, O'May GA, William Costerton J, Shirtliff ME. 2012. Polymicrobial interactions: Impact on pathogenesis and human disease. Clin Microbiol Rev 25:193–213.

6.  Rogers GB, Hoffman LR, Carroll MP, Bruce KD. 2013. Interpreting infective microbiota: The importance of an ecological perspective. Trends Microbiol 21:271–276.

7.  De Vos MGJ, Zagorski M, McNally A, Bollenbach T. 2017. Interaction networks, ecological stability, and collective antibiotic tolerance in polymicrobial infections. Proc Natl Acad Sci U S A 114:10666–10671.

8.  Rezzoagli C, Granato ET, Kümmerli R. 2020. Harnessing bacterial interactions to manage infections: A review on the opportunistic pathogen Pseudomonas aeruginosa as a case example. J Med Microbiol 69:147–161.

9.  Short FL, Murdoch SL, Ryan RP. 2014. Polybacterial human disease: The ills of

social networking. Trends Microbiol 22:508–516.

10.   Tashiro Y, Yawata Y, Toyofuku M, Uchiyama H, Nomura N. 2013. Interspecies

interaction between Pseudomonas aeruginosa and other microorganisms.

Microbes Environ 28:13–24.

11.   Ryan RP, Dow JM. 2008. Diffusible signals and interspecies communication in

bacteria. Microbiology 154:1845–1858.

12.   Murray JL, Connell JL, Stacy A, Turner KH, Whiteley M. 2014. Mechanisms of

synergy in polymicrobial infections. J Microbiol 52:188–199.

13.   Hubert D, Réglier-Poupet H, Sermet-Gaudelus I, Ferroni A, Le Bourgeois M,

Burgel PR, Serreau R, Dusser D, Poyart C, Coste J. 2013. Association between

Staphylococcus aureus alone or combined with Pseudomonas aeruginosa and

the clinical condition of patients with cystic fibrosis. J Cyst Fibros 12:497–503.

14.   Maliniak ML, Stecenko AA, McCarty NA. 2016. A longitudinal analysis of chronic

MRSA and Pseudomonas aeruginosa co-infection in cystic fibrosis: A single-

center study. J Cyst Fibros 15:350–356.

15.   Limoli DH, Hoffman LR. 2019. Help, hinder, hide and harm: What can we learn

from the interactions between Pseudomonas aeruginosa and Staphylococcus

aureus during respiratory infections. Thorax 74:684–692.

16.   Orazi G, Ruoff KL, O'Toole GA. 2019.  Pseudomonas aeruginosa Increases the

Sensitivity of Biofilm-Grown Staphylococcus aureus to Membrane-Targeting

Antiseptics and Antibiotics . MBio 10:1–15.

17.   Tognon M, Köhler T, Luscher A, Van Delden C. 2019. Transcriptional profiling of

Pseudomonas aeruginosa and Staphylococcus aureus during in vitro co-culture.

BMC Genomics 20:1–15.

18. Cystic Fibrosis Trust. 2014. UK Cystic Fibrosis Registry Annual data report 2018.

Registry.

19. Gjødsbøl K, Christensen JJ, Karlsmark T, Jørgensen B, Klein BM, Krogfelt KA.

2006. Multiple bacterial species reside in chronic wounds: A longitudinal study. Int

Wound J 3.

20. Malic S, Hill KE, Hayes A, Percival SL, Thomas DW, Williams DW. 2009.

Detection and identification of specific bacteria in wound biofilms using peptide

nucleic acid fluorescent in situ hybridization (PNA FISH). Microbiology 155:2603–

2611.

21. Mashburn LM, Jett AM, Akins DR, Whiteley M. 2005. Staphylococcus aureus

serves as an iron source for Pseudomonas aeruginosa during in vivo coculture. J

Bacteriol 187:554–566.

22. Hoffman LR, Déziel E, D'Argenio DA, Lépine F, Emerson J, McNamara S, Gibson

RL, Ramsey BW, Miller SI. 2006. Selection for Staphylococcus aureus small-

colony variants due to growth in the presence of Pseudomonas aeruginosa. Proc

Natl Acad Sci U S A 103:19890–19895.

23. Harrison F, Paul J, Massey RC, Buckling A. 2008. Interspecific competition and

siderophore-mediated cooperation in Pseudomonas aeruginosa. ISME J 2:49–55.

24. Biswas L, Biswas R, Schlag M, Bertram R, Götz F. 2009. Small-colony variant

selection as a survival strategy for Staphylococcus aureus in the presence of

Pseudomonas aeruginosa. Appl Environ Microbiol 75:6910–6912.

25. Kessler E, Safrin M, Olson JC, Ohman DE. 1993. Secreted LasA of

*Pseudomonas aeruginosa* is a Staphylolytic Protease. J Biol Chem 268:7503–7508.

26. Dalton T, Dowd SE, Wolcott RD, Sun Y, Watters C, Griswold JA, Rumbaugh KP. 2011. An in vivo polymicrobial biofilm wound infection model to study interspecies interactions. PLoS One 6.

27. Pastar I, Nusbaum AG, Gil J, Patel SB, Chen J, Valdes J, Stojadinovic O, Plano LR, Tomic-Canic M, Davis SC. 2013. Interactions of Methicillin Resistant Staphylococcus aureus USA300 and Pseudomonas aeruginosa in Polymicrobial Wound Infection. PLoS One 8:1–11.

28. Tognon M, Köhler T, Gdaniec BG, Hao Y, Lam JS, Beaume M, Luscher A, Buckling A, Van Delden C. 2017. Co-evolution with Staphylococcus aureus leads to lipopolysaccharide alterations in Pseudomonas aeruginosa. ISME J 11:2233–2243.

29. Michelsen CF, Christensen AMJ, Bojer MS, Høiby N, Ingmer H, Jelsbak L. 2014. Staphylococcus aureus alters growth activity, autolysis, and antibiotic tolerance in a human host-adapted Pseudomonas aeruginosa lineage. J Bacteriol 196:3903–3911.

30. Michelsen CF, Khademi SMH, Johansen HK, Ingmer H, Dorrestein PC, Jelsbak L. 2016. Evolution of metabolic divergence in Pseudomonas aeruginosa during long-term infection facilitates a proto-cooperative interspecies interaction. ISME J 10:1330–1336.

31. DeLeon S, Clinton A, Fowler H, Everett J, Horswill AR, Rumbaugh KP. 2014. Synergistic interactions of Pseudomonas aeruginosa and Staphylococcus aureus

in an In vitro wound model. Infect Immun 82:4718–4728.

32.  Beaudoin T, Yau YCW, Stapleton PJ, Gong Y, Wang PW, Guttman DS, Waters V.

2017. Staphylococcus aureus interaction with Pseudomonas aeruginosa biofilm

enhances tobramycin resistance. npj Biofilms Microbiomes 3:25.

33.  Orazi G, O'Toole GA. 2017. Pseudomonas aeruginosa alters Staphylococcus

aureus sensitivity to vancomycin in a biofilm model of cystic fibrosis infection.

MBio 8:1–17.

34.  Grundmeier M, Tuchscherr L, Brück M, Viemann D, Roth J, Willscher E, Becker

K, Peters G, Löffler B. 2010. Staphylococcal Strains Vary Greatly in Their Ability

to Induce an Inflammatory Response in Endothelial Cells. J Infect Dis 201:871–

880.

35.  Vann JM, Proctor RA. 1987. Ingestion of Staphylococcus aureus by bovine

endothelial cells results in time- and inoculum-dependent damage to endothelial

cell monolayers. Infect Immun 55:2155–2163.

36.  Haslinger-Löffler B, Kahl BC, Grundmeier M, Strangfeld K, Wagner B, Fischer U,

Cheung AL, Peters G, Schulze-Osthoff K, Sinha B. 2005. Multiple virulence

factors are required for Staphylococcus aureus-induced apoptosis in endothelial

cells. Cell Microbiol 7:1087–1097.

37.  Balwit JM, Langevelde P Van, Vann JM, Proctorn RA. 1994. Gentamicin-resistant

menadione and hemin auxotrophic staphylococcus aureus persist within cultured

endothelial cells. J Infect Dis 170:1033–1037.

38.  Nimmo GR. 2012. USA300 abroad: Global spread of a virulent strain of

community-associated methicillin-resistant Staphylococcus aureus. Clin Microbiol

Infect 18:725–734.

39.    Thurlow LR, Joshi GS, Richardson AR. 2012. Virulence strategies of the dominant

USA300 lineage of community-associated methicillin-resistant Staphylococcus

aureus (CA-MRSA). FEMS Immunol Med Microbiol 65:5–22.

40.    Rendueles O, Amherd M, Velicer GJ. 2015. Positively Frequency-Dependent

Interference Competition Maintains Diversity and Pervades a Natural Population

of Cooperative Microbes. Curr Biol 25:1673–1681.

41.    Healey D, Axelrod K, Gore J. 2016. Negative frequency-dependent interactions

can underlie phenotypic heterogeneity in a clonal microbial population. Mol Syst

Biol 12:877.

42.    Levin BR. 1988. Frequency-dependent selection in bacterial populations. Philos

Trans R Soc Lond B Biol Sci 319:459–472.

43.    Ross-Gillespie A, Gardner A, West SA, Griffin AS. 2007. Frequency dependence

and cooperation: Theory and a test with bacteria. Am Nat 170:331–342.

44.    Kerr B, Riley MA, Feldman MW, Bohannan BJM. 2002. Local dispersal promotes

biodiversity in a real-life game of rock-paper-scissors. Nature 418:171–174.

45.    Van Gestel J, Weissing FJ, Kuipers OP, Kovács ÁT. 2014. Density of founder

cells affects spatial pattern formation and cooperation in Bacillus subtilis biofilms.

ISME J 8:2069–2079.

46.    Weigert M, Ross-Gillespie A, Leinweber A, Pessi G, Brown SP, Kümmerli R.

2017. Manipulating virulence factor availability can have complex consequences

for infections. Evol Appl 10:91–101.

47.    Spiers AJ, Bohannon J, Gehrig SM, Rainey PB. 2003. Biofilm formation at the air-

liquid interface by the Pseudomonas fluorescens SBW25 wrinkly spreader

requires an acetylated form of cellulose. Mol Microbiol 50:15–27.

48.    Borer B, Tecon R, Or D. 2018. Spatial organization of bacterial populations in

response to oxygen and carbon counter-gradients in pore networks. Nat Commun

9.

49.    Limoli DH, Yang J, Khansaheb MK, Helfman B, Peng L, Stecenko AA, Goldberg

JB. 2016. Staphylococcus aureus and Pseudomonas aeruginosa co-infection is

associated with cystic fibrosis-related diabetes and poor clinical outcomes. Eur J

Clin Microbiol Infect Dis 35:947–953.

50.    Serra R, Grande R, Butrico L, Rossi A, Settimio UF, Caroleo B, Amato B, Gallelli

L, De Franciscis S. 2015. Chronic wound infections: The role of Pseudomonas

aeruginosa and Staphylococcus aureus. Expert Rev Anti Infect Ther 13:605–613.

51.    Hotterbeekx A, Kumar-Singh S, Goossens H, Malhotra-Kumar S. 2017. In vivo

and In vitro interactions between Pseudomonas aeruginosa and Staphylococcus

spp. Front Cell Infect Microbiol 7:1–13.

52.    Nguyen AT, Oglesby-Sherrouse AG. 2016. Interactions between Pseudomonas

aeruginosa and Staphylococcus aureus during co-cultivations and polymicrobial

infections. Appl Microbiol Biotechnol 100:6141–6148.

53.    Machan ZA, Pitt TL, White W, Watson D, Taylor GW, Cole PJ, Wilson R. 1991.

Interaction between Pseudomonas aeruginosa and Staphylococcus aureus:

Description of an anti-staphylococcal substance. J Med Microbiol 34:213–217.

54.    Pernet E, Guillemot L, Burgel PR, Martin C, Lambeau G, Sermet-Gaudelus I,

Sands D, Leduc D, Morand PC, Jeammet L, Chignard M, Wu Y, Touqui L. 2014.

688   Pseudomonas aeruginosa eradicates Staphylococcus aureus by manipulating the

689   host immunity. Nat Commun 5:5105.

690 55. Filkins LM, O'Toole GA. 2015. Cystic Fibrosis Lung Infections: Polymicrobial,

691   Complex, and Hard to Treat. PLoS Pathog 11:1–8.

692 56. King JM, Kulhankova K, Stach CS, Vu BG, Salgado-Pabón W. 2016. Phenotypes

693   and Virulence among Staphylococcus aureus USA100, USA200, USA300,

694   USA400, and USA600 Clonal Lineages. mSphere 1:1–15.

695 57. Tuchscherr L, Pöllath C, Siegmund A, Deinhardt-Emmer S, Hoerr V, Svensson

696   CM, Figge MT, Monecke S, Löffler B. 2019. Clinical S. aureus isolates vary in

697   their virulence to promote adaptation to the host. Toxins (Basel) 11.

698 58. Montgomery CP, Boyle-Vavra S, Adem PV, Lee JC, Husain AN, Clasen J, Daum

699   RS. 2008.  Comparison of Virulence in Community-Associated Methicillin-

700   Resistant Staphylococcus aureus Pulsotypes USA300 and USA400 in a Rat

701   Model of Pneumonia . J Infect Dis 198:561–570.

702 59. Diep BA, Gill SR, Chang RF, Phan TH, Chen JH, Davidson MG, Lin F, Lin J,

703   Carleton HA, Mongodin EF, Sensabaugh GF, Perdreau-Remington F. 2006.

704   Complete genome sequence of USA300, an epidemic clone of community-

705   acquired meticillin-resistant Staphylococcus aureus. Lancet 367:731–739.

706 60. Watkins RR, David MZ, Salata RA. 2012. Current concepts on the virulence

707   mechanisms of meticillin-resistant Staphylococcus aureus. J Med Microbiol

708   61:1179–1193.

709 61. Joo HS, Cheung GYC, Otto M. 2011. Antimicrobial activity of community-

710   associated methicillin-resistant Staphylococcus aureus is caused by phenol-

711      soluble modulin derivatives. J Biol Chem 286:8933–8940.

62. Cheung AL, Bayer AS, Zhang G, Gresham H, Xiong YQ. 2004. Regulation of virulence determinants in vitro and in vivo in Staphylococcus aureus. FEMS Immunol Med Microbiol 40:1–9.

63. Baldan R, Cigana C, Testa F, Bianconi I, De Simone M, Pellin D, Di Serio C, Bragonzi A, Cirillo DM. 2014. Adaptation of Pseudomonas aeruginosa in cystic fibrosis airways influences virulence of Staphylococcus aureus in vitro and murine models of co-infection. PLoS One 9.

64. Radlinski L, Rowe SE, Kartchner LB, Maile R, Cairns BA, Vitko NP, Gode CJ, Lachiewicz AM, Wolfgang MC, Conlon BP. 2017. Pseudomonas aeruginosa exoproducts determine antibiotic efficacy against Staphylococcus aureus. PLoS Biol 15:1–25.

65. Bernardy EE, Petit III RA, Raghuram V, Alexander AM, Read TD, Goldberg JB. 2020. Genotypic and Phenotypic Diversity of Staphylococcus aureus Isolates from Cystic Fibrosis Patient Lung Infections and Their Interactions with Pseudomonas aeruginosa. MBio 11:1–18.

66. Leinweber A, Inglis FR, Kümmerli R. 2017. Cheating fosters species co-existence in well-mixed bacterial communities. ISME J 11:1179–1188.

67. Cornforth DM, Foster KR. 2013. Competition sensing: The social side of bacterial stress responses. Nat Rev Microbiol.

68. Zhang QG, Buckling A, Ellis RJ, Godfray HCJ. 2009. Coevolution between cooperators and cheats in a microbial system. Evolution (N Y) 63:2248–2256.

69. Valot B, Guyeux C, Rolland JY, Mazouzi K, Bertrand X, Hocquet D. 2015. What it

takes to be a Pseudomonas aeruginosa? The core genome of the opportunistic

pathogen updated. PLoS One 10:1–15.

70. Wolfgang MC, Kulasekara BR, Liang X, Boyd D, Wu K, Yang Q, Miyada CG, Lory

S. 2003. Conservation of genome content and virulence determinants among

clinical and environmental isolates of Pseudomonas aeruginosa. Proc Natl Acad

Sci U S A 100:8484–8489.

71. Hyun JK, Boedicker JQ, Jang WC, Ismagilov RF. 2008. Defined spatial structure

stabilizes a synthetic multispecies bacterial community. Proc Natl Acad Sci U S A

105:18188–18193.

72. Lowery NV, Ursell T. 2019. Structured environments fundamentally alter

dynamics and stability of ecological communities. Proc Natl Acad Sci U S A

116:379–388.

73. Refardt D, Bergmiller T, Kümmerli R. 2013. Altruism can evolve when relatedness

is low: Evidence from bacteria committing suicide upon phage infection. Proc R

Soc B Biol Sci 280.

74. Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ,

Brinkman FSL, Hufnagle WO, Kowallk DJ, Lagrou M, Garber RL, Goltry L,

Tolentino E, Westbrock-Wadman S, Yuan Y, Brody LL, Coulter SN, Folger KR,

Kas A, Larbig K, Lim R, Smith K, Spencer D, Wong GKS, Wu Z, Paulsen IT,

Relzer J, Saler MH, Hancock REW, Lory S, Olson M V. 2000. Complete genome

sequence of Pseudomonas aeruginosa PAO1, an opportunistic pathogen. Nature

406:959–964.

75. Choi KH, DeShazer D, Schweizer HP. 2006. mini-Tn7 insertion in bacteria with

multiple glmS-linked attTn7 sites: Example Burkholderia mallei ATCC 23344. Nat

Protoc 1:162–169.

76.  Rezzoagli C, Granato ET, Kümmerli R. 2019. In-vivo microscopy reveals the

impact of Pseudomonas aeruginosa social interactions on host colonization. ISME

J 488262.

77.  Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a pratical

and powerful approach to multiple testing. J R Stat Soc Ser B 57:289–300.

## Table and figures

**Table 1.** PA and SA strains used for this study.

| Strain name | Origin | Description | Reference |
|---|---|---|---|
| ***Pseudomonas aeruginosa* (PA)** | | | |
| PAO1::gfp | Wound | Constitutive GFP expression from the chromosome (attTn7::Ptac-GFP). | Our laboratory |
| ***Staphylococcus aureus* (SA)** | | | |
| Cowan I | Septic arthritis | MSSA isolate. Highly invasive, but not cytotoxic. Agr-defective. | ATCC 12598 |
| 6850 | Osteomyelitis | MSSA isolate. Highly invasive, cytotoxic and hemolytic. | ATCC 53657 |
| JE2 | Skin and soft tissue infection | USA300 CA-MRSA isolate. Highly virulent, cytotoxic and hemolytic. | NARSA |

CA-MRSA: Community-acquired methicillin-resistant *S. aureus*

MSSA: Methicillin-sensitive *S. aureus*

Agr: Accessory gene regulator

**Figure 1.** Workflow for the competition experiments. Bacterial overnight cultures were

grown in 10 ml TSB in 50 ml falcon tubes for ~ 16 hours at 37 °C and 220 rpm with

aeration. After washing and adjustment of $OD_{600}$ to obtain similar cell numbers for all

strains, strain pairs (PA-Cowan I ; PA-6850 ; PA-JE2) were mixed at three different

volumetric starting frequencies (1:9 ; 1:1 ; 9:1). Flow cytometry was used to measure

the actual starting frequencies. Competitions were started with diluted cultures ($OD_{600}$ =

$10^{-5}$) in 24-well plates filled with 1.5 ml TSB per well. Plates were incubated for 24 hours

at 37 °C under three different culturing conditions: shaken (170 rpm), viscous (170 rpm

+ 0.2% agar in TSB) and static. After the 24 hours competition period, final strain

frequencies were measured for each replicate by flow cytometry. Using the initial and

final strain frequencies, the relative fitness (v) of the focal strain PA was calculated as

$v = [a_t \times (1-a_0)]/ [a_0 \times (1-a_t)]$, where $a_0$ and $a_t$ are the initial and final frequencies of PA,

respectively.

**Figure 2.** The number of doublings in monoculture is higher for *P. aeruginosa* PAO1

(PA) than for the three *S. aureus* strains (Cowan I, 6850, JE2) under most conditions.

Strains were grown as monocultures in TSB for 24 hours at 37 °C under the same

conditions and using the same starting $OD_{600}$ as for the competition experiments. The

box plots show the median (bold line) with the first and the third quartiles. The whiskers

cover the 1.5* inter-quartile range (IQR) or extend from the lowest to the highest value if

they fall within the 1.5* IQR. Data are shown from two independent experiments with

three replicates each. * $p < 0.05$, *** $p < 0.001$, pairwise comparisons without bars are

all not significant.

**Figure 3.** Relative fitness ln(v) of *P. aeruginosa* PAO1 (PA) after 24-hours competitions

against three different *S. aureus* (SA) strains (Cowan I, 6850, JE2) at three different

starting frequencies (1:9 ; 1:1 ; 9:1) and across three different environmental conditions

(shaken, viscous, static). Values of ln(v) < 0, ln(v) > 0, or ln(v) = 0 (dotted line), indicate

whether PA lost, won, or performed equally well in competition against the respective

SA strain. The box plots show the median (bold line) with the first and third quartiles.

The whiskers cover the 1.5* inter-quartile range (IQR) or extend from the lowest to the

highest value if they fall within the 1.5* IQR. Each strain pair/culturing condition/starting

frequency combination was repeated 20 times (four experiments featuring five

replicates each). Asterisks indicate whether the relative fitness of PA is significantly

different from zero in a specific treatment (one-sample t-tests with p-values corrected for

multiple comparisons using the false discovery rate method:  n.s. = not significant, * p <

0.05, ** p < 0.01, *** p < 0.001).  Detailed information on all statistical comparisons are

provided in Table S3.

**Figure 4.** Multi-day competitive dynamics between *P. aeruginosa* PAO1 (PA) and the

three *S. aureus* strains (a) Cowan I, (b) 6850 and (c) JE2 under shaken conditions.

Competitions started at three volumetric starting frequencies of PA:SA (red 1:9, blue

1:1, green 9:1). Community composition was followed over five days with daily transfer

of diluted cultures to fresh TSB medium. Strain frequencies were assessed using flow

cytometry. The experiment was carried out two times with five replicates per treatment

combination and experiment.