Naïve orangutans (*Pongo abelii & Pongo pygmaeus*) individually acquire nut-cracking

using hammer tools

Elisa Bandini[1], Johannes Grossmann[2]; Martina Funk[3], Anna Albiach Serrano[4*] & Claudio

Tennie[1*]

[1]*Department for Early Prehistory and Quaternary Ecology, The University of Tübingen,*

Tübingen, 72070, Germany

[2]*Max Plank Institute for Evolutionary Anthropology, Deutscher Platz 6, Leipzig, 04103,*

Germany

[3]*Independent researcher*

[4]*Ethology and Animal Welfare Section, Universidad Cardenal Herrera-CEU, CEU*

Universities, Valencia, Spain

[*]Shared senior authors

Corresponding author: Elisa Bandini, elisa-bandini@hotmail.it

Abstract

Nut-cracking using hammer tools has been argued to be one of the most complex tool-use
behaviours observed in non-human animals (henceforth: animals). Recently, even the United
Nations Convention on the Conservation of Migratory Species (CMS) recognised the unique
nature of chimpanzee nut-cracking by making it the first animal behaviour to be awarded UN-
protected status (Picheta, 2020). So far, only chimpanzees, capuchins and macaques have
been observed using tools to crack nuts in the wild (Boesch & Boesch, 1990; Gumert, Kluck,
& Malaivijitnond, 2009; Ottoni & Mannu, 2001). However, the learning mechanisms behind
this behaviour, and the extent of nut-cracking in other primate species are still unknown. The
aim of this study was two-fold. First, we aimed to examine whether other great ape species
would develop nut-cracking when provided with all the tools and motivation to do so. Second,
we wanted to examine the mechanisms behind the emergence of nut-cracking in a naïve
sample. Orangutans (*Pongo abelii; pygmaeus*) have not been observed cracking nuts in the
wild, despite having the second most extensive tool-use repertoire of the great apes (after
chimpanzees), having the materials for the behaviour in the wild (albeit rarely) and possessing
flexible problem-solving capacities. Therefore, orangutans are a valid candidate species for
the investigation of the development of nut-cracking. Four nut-cracking-naïve orangutans at
Leipzig zoo (*Pongo abelii*; M_{age}=16; age range=10-19; 4F; at time of testing) were provided
with nuts and hammers but were not demonstrated the nut-cracking behavioural form, in order
to control for the role of copying social learning in the acquisition of this behaviour.
Additionally, we report data from a previously unpublished study by one of the authors (MF)
with eight orangutans housed at Zürich zoo (10 *Pongo abelii* and two *Pongo pygmaeus;*
M_{age}=14; age range =2-30; 5F; at time of testing) that followed a similar testing paradigm.
Out of the twelve orangutans across both testing institutions, at least four individuals, one
from Leipzig (*Pongo abelii*) and three from Zürich (*Pongo abelii*; *pygmaeus*), spontaneously
expressed nut-cracking with a wooden hammer. These results suggest that the behavioural
form of nut-cracking using hammer tools can emerge in orangutans when required through
individual learning combined, in some cases, with non-copying social learning mechanisms.

Keywords: Nut-cracking, Orangutan tool-use, Individual learning, Social learning, Animal
tool-use

Introduction

Once heralded as the main distinguishing feature of humans in the animal kingdom, it is now known that several other species also possess the ability to use tools (Shumaker, Walkup, Beck, & Burghardt, 2011). Of these species, non-human great apes (henceforth apes), alongside New Caledonian crows (Kenward et al., 2011), demonstrate the most extensive and potentially most 'complex' tool-use repertoires (van Schaik, Deaner, & Merrill, 1999), with complex tool-use defined here as: *"tool-use variants that include at least two tool elements (for example, hammer and anvil), flexibility in manufacture or use (that is, tool properties are adjusted to the task at hand), and that skills are acquired in part by social learning"* (Meulman, Sanz, Visalberghi, & van Schaik, 2012, 58). Of the tool-use behaviours observed in wild and captive great apes, nut-cracking using hammer tools (henceforth: nut-cracking) in chimpanzees is often cited as one of the most complex behaviours within their tool-use repertoires (Boesch, 1991; Lonsdorf, 2013). The claims for the complexity of nut-cracking in chimpanzees are based on several assumptions on the behaviour, including the learning mechanisms underlying the acquisition of nut-cracking in wild chimpanzees. Whilst we will discuss these assumptions in further detail in the following section, below we provide definitions for the main social learning mechanisms discussed here.

The animal social learning field is rife with terminology and mechanisms (for an overview of these terms, see Whiten, Horner, Litchfield, & Marshall-Pescini, 2004). However, throughout this paper we will mainly be referring to two types of social learning: copying and non-copying social learning mechanisms. Copying social learning refers to social learning mechanisms that can transmit the actual form of a behaviour and/or artefact (where form is defined as "*the specific action [and/or artefact] component(s) and organisation of a behaviour";* Bandini & Tennie, 2020, 2). Copying social learning includes mechanisms such as imitation and some types of emulation (Bandini & Tennie, 2020). On the other hand, non-copying social learning mechanisms are those that cannot transmit the form of a behaviour and/or artefact. Instead, these mechanisms can regulate the frequency of the acquisition of a behaviour, for example by increasing an individual's motivation to interact or manipulate a certain object or location. Non-copying social learning mechanisms include local and/or stimulus enhancement (Bandini & Tennie, 2020).

Nut-cracking in primates

Nuts represent an important source of calories and fat in primates' diets (Biro et al., 2003).

The encased condition of these nutrients, however, makes it often beneficial and sometimes

neccessary to use stone and/or wooden hammers to crack open the nuts against hard surfaces.

Nut-cracking has been observed in chimpanzees, long-tailed macaques, and capuchins

(Boesch & Boesch, 1990; Gumert, Kluck, & Malaivijitnond, 2009; Ottoni & Mannu, 2001).

Of all nut-cracking behaviours, however, the best-studied example is that of chimpanzees

(Biro et al., 2003; Boesch & Boesch, 1990; Luncz & Boesch, 2014; Luncz, Mundry, &

Boesch, 2012). Indeed, this behaviour in chimpanzees has recently been selected for

conservation by the United Nations Convention on the Conservation of Migratory Species

(CMS) body, showing how important – and how cultural - this behaviour is considered to be,

even by organisations outside of academia (Picheta, 2020).

Wild chimpanzees in the Taï Forest (Ivory Coast) and in Bossou (Guinea) use hammer tools

to access the kernels of several nut species - *Panda oleosa, Parinari excelsa, Saccoglottis*

gabonensis, Coula edulis, and *Detarium senegalensis* (Proffitt, Haslam, Mercader, Boesch, &

Luncz, 2018). . The crux of the nut-cracking behavioural form in chimpanzees (see also

Foucart et al., 2005) involves three steps: (1) Retrieving a nut from the surrounding area and

placing it on an anvil (e.g., a tree root or a stone), (2) Picking up a stone- or wooden *hammer*

(with one hand or both hands) and (3) Hitting the nut with the *hammer* (holding it with one or

both hands) until it is open and the inside kernel can be retrieved and consumed (Boesch &

Boesch, 1983; Carvalho, Biro, McGrew, & Matsuzawa, 2009). Sometimes more steps are

described, such as the transportation of the materials to the nut-cracking site (Carvalho et al.,

2009) and the stabilisation of the anvil on the ground (although this is a rare behaviour;

Carvalho et al., 2009). This multi-step approach has been regarded as a complex tool use

behaviour (Meulman et al., 2012), because it is improbable that such a compound behaviour

is acquired in its entirety by chance, especially considering that it is only rewarded at the end

of the chain of actions (note that most of the other behavioural forms within chimpanzee tool-

use repertoire only involve the manipulation of a single object (usually a stick) and include

only one action (e.g., marrow picking; see Whiten et al., 2001 for an overview of chimpanzee

behaviours and their description). Moreover, the precision needed to crack open nuts is said to

contribute to the complexity of the behaviour since (at least at the beginning) many attempts

will go unrewarded. Perhaps for these reasons, chimpanzee nut-cracking is often assumed to

be driven and maintained by action copying (Boesch, 1991; Boesch, Marchesi, Marchesi,
Fruth, & Joulian, 1994), although this assumption is heavily debated (e.g., see Neadle et al.,
2020).

Acquisition of nut-cracking in chimpanzees

Juvenile chimpanzees take a long time to acquire nut-cracking (Biro et al., 2003; Boesch &
Boesch, 1990) and observations of wild juvenile chimpanzees suggest that the acquisition of
nut-cracking may occur within a sensitive learning period, most likely when chimpanzees are
between the ages of three and five years old (Inoue-Nakamura & Matsuzawa, 1997). If the
behaviour is not acquired within this sensitive learning period, chimpanzees will seemingly
never develop the behaviour (Biro et al., 2003). This seems to also be the case for nut-
cracking in other primates, such as long-tailed macaques (Tan, 2017).

Furthermore, it has been proposed that during this learning period, juveniles acquire nut-
cracking by observing and then copying their mother's actions (e.g., see Biro et al., 2003) and
that a repeated cycle of observation and practice sessions is required before nut-cracking can
be expressed (e.g., what Whiten, 2017, 7795, describes as a "helical process of learning"). In
a similar interpretation, de Waal (2008) suggests that juvenile chimpanzees copy their
mothers' nut-cracking via 'Bonding and Identification-based Observational Learning'
(BIOL), where a juvenile copies actions in order "to be like others" (de Waal, 2008, 231).
Yet, a lengthy learning period alone is not necessarily indicative of copying. Instead, it can be
also be explained by mere maturation processes, alongside an extended period of individual
learning (likely encouraged by non-copying variants of social learning, such as stimulus and
local enhancement; Whiten et al., 2004 and "peering"; Corp & Byrne, 2002; Schuppli et al.,
2016). Furthermore, conclusive evidence for (unenculturated) apes possessing the ability to
copy behaviours is still lacking. Whilst enculturated and heavily trained primates have
demonstrated an ability to copy actions (indeed, this type of training seems to change the
individuals' brain structures to allow action copying; Hecht et al., 2013; Pope, Taglialatela,
Skiba, & Hopkins, 2017), experimental paradigms aimed at identifying this ability in
unenculturated chimpanzees have so far been unsuccessful (Clay & Tennie, 2017; Tennie,
Call, & Tomasello, 2012; Tomasello et al., 1997). Therefore, whilst further studies should be
carried out on this ability in different ape species, the current state of knowledge suggests that
alternative explanations to those based solely on copying social learning should be explored.

One alternative approach towards explaining the acquisition of nut-cracking in primates is

provided by the zone of latent solutions hypothesis (ZLS; Tennie, Call, & Tomasello, 2009).

The ZLS hypothesis argues that behavioural forms are acquired in many species via an

interplay of individual learning and non-copying variants of social learning. According to this

hypothesis, the behavioural form of primate nut-cracking would not be copied, but

individually derived. There are many ways in which this individual learning may work. To

give just one example, the difficulty of learning individually such a complex behaviour may

be overcome by individuals having a general predisposition to explore and manipulate objects

plus some cognitive capacities like good spatial memory (that allows to locate the needed

materials), inhibitory control (that allows to delay a reward), planning abilities and working

memory (that allow to chain steps towards a goal) and some understanding of the physical

affordances of objects and of object relations (that can aid in the selection of appropriate

materials and actions to process the materials). As a result, such individuals are able to solve

problems in a flexible way. This would explain why, for example, wild nut-cracking

chimpanzees use different types of anvils (stationary and non-stationary) depending on their

needs and material availability. The ZLS hypothesis also suggests that the observed

differences in nut-cracking activity across chimpanzee populations are fostered by non-

copying social learning mechanisms (such as local and/or stimulus enhancement), which

increase the likelihood of reinnovation once a population already contains individuals who

have innovated the behavioural form. This can then lead to a frequency increase and

maintenance of the behavioural forms in question in some populations but not in others. The

ZLS hypothesis predicts successful reinnovation of behavioural forms by naïve ape subjects

provided the right conditions and in the absence of copying opportunities. This prediction has

been met several times, as demonstrated by a fast-growing experimental literature detailing

successful individual acquisitions of various wild-type behavioural forms (including tool use)

in various species of naïve, captive great apes (Allritz, Tennie, & Call, 2013; Bandini &

Tennie, 2017; 2019; 2020; Bandini & Harrison, 2020; Menzel, Fowler, Tennie, & Call, 2013;

Neadle, Allritz, & Tennie, 2017; Tennie, Hedwig, Call, & Tomasello, 2008). Therefore, the

ape ZLS hypothesis has growing support, but whether it can also explain the behavioural form

of nut-cracking is still an open question.

Latent solutions testing methodology

In order to examine whether a subject can acquire a behavioural form and the mechanisms behind this acquisition (whether it is driven by copying social learning or by individual learning and non-copying social learning), Tennie & Hedwig (2009) described the 'latent solutions' (LS) testing methodology. This methodology allows for the role of individual learning in the acquisition of a target behavioural form. All the ecological materials of the target behavioural form, but no demonstrations, are provided to naïve subjects, who have never seen, or been trained in, the target behaviour before. Subjects should be so-called 'ecologically-representative' individuals, i.e. unenculturated captive animals who live in social groups (Henrich & Tennie, 2017). If the target behavioural form emerges under these conditions, then, logically, it can be concluded that copying is not *required* for the form of behaviour to emerge. If the behaviour does not emerge in this baseline condition, then it could be that some variant of copying is necessary for the behaviour to be acquired (for these cases, Bandini & Tennie, 2018 provide an extended LS testing methodology that allows for the level and variant of social learning required (if any) to be identified), *or* that other factors, such as sensitive periods, opportunities to practice or motivation levels, play a role (Bandini & Tennie, 2018; Neadle et al., 2020). Past LS studies have demonstrated that multiple behavioural forms – including tool use behavioural forms – can be individually acquired by primates (see above). For example, naïve, captive, capuchins were tested following the LS testing methodology and two individuals spontaneously started cracking nuts, without copying the behaviour (Visalberghi, 1987). Furthermore, this methodology has also shown that different species may sometimes overlap in their latent solution repertoires. Indeed, human children were recently found to spontaneously acquire various tool-use behaviours observed in wild chimpanzees and orangutans (Reindl, Beck, Apperly, & Tennie, 2016, Nelder et al., 2020; and see also: Allritz et al., 2013; Bandini & Tennie, 2019, 2017; Menzel et al., 2013; Neadle et al., 2017; Tennie et al., 2008 for examples of latent solutions across primate species).

With regard to great apes, apart from the observations of chimpanzee nut-cracking, anecdotal reports exist of gorillas and bonobos cracking nuts in sanctuaries (Wrangham, 2006) – though the exact circumstances of these innovations remain unclear, whilst orangutans have not (perhaps not yet) been reported to crack nuts (Fox et al., 2004; Parker & Gibson, 1977). Yet, despite being the most arboreal great apes, orangutans have the second most extensive tool-

use repertoire in the wild after chimpanzees (Fox, van Schaik, Sitompul, & Wright, 2004;
Meulman & van Schaik, 2013). Orangutans' tool-use repertoires include extractive foraging
tool-use and, similarly to that of chimpanzees, demonstrate intra and inter-population
variation (van Schaik & Knott, 2001). Moreover, there are some reports of orangutans using
sticks to 'hammer' open termite or bee nests (Fox, Sitompul, & van Schaik, 1999). These
behaviours are not fully comparable to chimpanzee nut-cracking, as they generally involve
thin sticks (whilst chimpanzee wooden hammers are usually thicker and/or heavier than the
sticks they use for other tool-use behaviours) and the hammering action used by orangutans is
unlikely to carry enough force to break an object as sturdy as an encased nut (Fox et al.,
1999). Therefore, orangutans present a promising candidate to acquire nut-cracking, whilst
still remaining naïve to this form of hammering behaviour..

Four naïve captive orangutans housed at Leipzig zoo (M_{age}=16; age range=10-19; 4F; at time
of testing) were provided with all the raw materials necessary for nut-cracking (nuts, wooden
hammers, cracking locations), but with no information or demonstrations on *how* to crack
nuts. This was to test whether orangutans could individually acquire this behavioural form of
nut-cracking, without having access to a model to copy. After conducting this study, it was
brought to the authors' attention that an unpublished thesis contained a similar study with
orangutans, carried out by Dr. Funk (one of the co-authors of the current manuscript) between
1983-1984. In this study, eight naïve captive orangutans housed at Zürich zoo (10 *Pongo*
abelii and two *Pongo pygmaeus;* M_{age}=14; age range =2-30; 5F; at time of testing) were
tested, following a similar (baseline) paradigm as our study. Therefore, the methods and
findings from this previous, unpublished, study are also described here. For the sake of
clarity, we will refer to Leipzig study and Zürich study in each case.

Leipzig study

Methods

Ethical approval

In accordance with ethical recommendations, all subjects were housed in indoor and outdoor
enclosures containing climbing structures and natural features. Subjects received their
regularly scheduled feedings and had access to enrichment devices and water *ad lib.* Subjects
were never food or water deprived for the purposes of this study. All research was conducted
in the subjects sleeping rooms. An internal committee of the Max Planck Institute for
Evolutionary Anthropology (director, research coordinator) and the Leipzig zoo (head keeper,
curator, vet) granted ethical approval for this project. No medical, toxicological or
neurobiological research of any kind is conducted at the WKPRC. This research was non-
invasive and strictly adhered to the legal requirements of Germany. Animal husbandry and
research comply with the "EAZA Minimum Standards for the Accommodation and Care of
Animals in Zoos and Aquaria", the "WAZA Ethical Guidelines for the Conduct of Research
on Animals by Zoos and Aquariums" and the "Guidelines for the Treatment of Animals in
Behavioral Research and Teaching" of the Association for the Study of Animal Behavior
(ASAB).

Subjects

Research was carried out at the Wolfgang Köhler Primate Research Center (WKPRC),
Leipzig, Germany in 2007. Four orangutans (M_{age}=16; age range=10-19; 4F; at time of
testing) participated in the study (see the demographic information in Table 1; all subjects
were born (except for DK) and raised at the testing institution). The keepers confirmed that
none of the individuals in this study had prior experience with macadamia nuts. Hazelnuts and
walnuts, however, had occasionally been provided by the keepers. Yet, the orangutans either
opened these with their teeth or, occasionally, by hitting them with their hand against a hard
surface. Crucially, none of the orangutans at the WKPRC had ever been observed using a *tool*
for nut-cracking before this study. Indeed, heavy objects that could potentially be used as
hammers (such as stones or wooden stumps) are not allowed inside the enclosures of the
WKPRC, for health and safety reasons, and therefore the subjects can confidently be assumed
to have been naïve to the target behaviour prior to this study.

 Procedure

We implemented three conditions sequentially (see Table 2): The first condition was a
baseline, in which subjects could only acquire the nut-cracking behaviour individually, as no
information on the actions required for the behaviour was provided. The second condition was
another baseline, which we called locked-anvil condition, that guaranteed that the object
provided as an anvil could *only* be used as an anvil and not as a hammer (see below). The
third condition was a demonstration condition, in which subjects could potentially learn nut-
cracking behaviour through social learning (of any variant) after observing a conspecific
model (PD; age 10 at the time of testing). Subjects were tested separately with no visual or
acoustic access to each other. While the sub-adult (PD) was tested alone, the adult females
were tested together with their dependent offspring (however no data were analysed from the
behaviour of the offspring as they were too young at the time of testing to attempt performing
the task).

Baseline condition

During each of five baseline sessions, subjects had access to one large wooden block (the
anvil; height 30 cm, diameter 50 cm, approximate weight 50 kg) with 5 depressions (diameter
2.5 cm) carved into the top side to facilitate the placement of the nuts, mirroring similar
depressions of anvils in the wild (e.g., Carvalho et al., 2009; Luncz et al., 2012), two smaller
wooden blocks (the wooden hammers; height 30 cm, diameter 50 cm, approximate weight 2.4
kg each) and five macadamia nuts (see figure 1 below). The materials were scattered evenly
on the floor in the testing room, which was emptied of any other objects prior to the test to
avoid distractions, within approx. one square meter. The subjects were not allowed to enter
the room until all the materials were in place. Sessions lasted a maximum of twenty minutes
but were discontinued earlier if the subjects had successfully opened all the nuts. The shells of
the opened nuts and any nuts that the subjects did not open were retrieved after each session
and discarded. A video camera were used to record the subjects' behaviour. For each subject,
the between-session interval was at least 24 hours.

Locked-anvil condition

After the baseline condition, the single successful subject (PD, see the results section)
participated in four additional sessions that were similar to the initial baseline sessions but
with the anvil fixed on the ground (by being pressed down with a sliding door). This way, we

encouraged the subject to explore options other than using the anvil as a hammer to crack open the nuts (as in the baseline the subject used an anvil-dropping and rolling technique to crack the nuts).

Demonstration condition

After the baseline and locked-anvil conditions, the three orangutans that did not demonstrate the nut-cracking behaviour in the baseline condition participated in five subsequent demonstration condition sessions (15 sessions in total). Before each session, PD, who had reliably started the nut-cracking behaviour in the previous phases, served as a demonstrator, cracking (and eating) five macadamia nuts. The subject, who had access to two hammers and a fixed anvil, could observe PD's performance from an adjacent cage. As soon as the subject had observed at least one successful nut-cracking bout (coded when the subject had its head oriented towards the demonstrator and its eyes were open during a successful nut-cracking bout by the demonstrator), five macadamia nuts were placed into the subject's enclosure and the session started. The demonstrations continued even after the nuts were placed in the enclosure. The rest of the testing procedure remained the same as in the baseline condition (see above).

Data collection and reliability

We live and video coded the behaviour performed by subjects to try to open the macadamia nuts (see Tables 3 & 4). Two second coders, who were not familiar with the aims and results of the study, watched the videos and coded the same categories as the original coder to assess inter-rater reliability. One coded the ethogram of behaviours, and how often each individual demonstrated the methods, whilst the other coded the number of successes in opening the nuts and the time spent with a nut in the subject's mouth. A Cohens kappa was calculated to assess the inter-rater reliability of both sets of data. All data are available in OSF (please see: https://osf.io/43fbr/?view_only=fd9290ce18b542c7a43a102f600ab22d).

Reliability testing results

A Cohen's kappa was run to assess the reliability of the coded data. We did not expect to find a very high inter-observer reliability for some of the behaviours, as the data were collected in

the orangutans' management areas (due to the testing facilities requirements), which were
dark and often did not allow for a clear view from the filming platform (also as subjects
moved dynamically and often blocked camera angles with their bodies). Despite this, in terms
of the general coding of the ethogram a moderate agreement (Cohen, 1968) was found
(k=0.60). However, note that a substantial interobserver agreement (k=0.80) was found for our
main experimental outcome, namely for coding the target nut-cracking behavioural forms
(anvil on floor and hammer on floor behaviours shown by PD). Therefore, whilst we
acknowledge that the reliability scores are lower than ideal for some additional behaviours
coded in the ethogram, the main behavioural outcome (nut-cracking) was reliably coded. The
overarching aim of this study was to examine whether nut-cracking with a tool would develop
in naïve orangutans, and these observations received a substantial agreement across coders
(k=0.80).

Results

Table 3 presents the behaviours coded, their descriptions, how many individuals attempted
each behaviour, the first session in which behaviours were observed, in which experimental
conditions they were observed, whether or not they allowed opening nuts and the percentage
of times each behaviour resulted in successfully cracking open a nut (see supplementary
material for video clips of the most common behaviours observed). In the baseline condition,
the juvenile individual, PD (F, 10 years old at time of testing, mother-reared and born at the
testing institution; see Table 1), successfully cracked nuts by using the large wooden anvil-
block as a hammer-tool (see also Table 3). When, in the locked-anvil condition, the large
anvil-block was fixed to the ground and therefore could no longer be used as a hammer, this
same subject cracked nuts by using the wooden hammers (see supplementary videos),
showing a similar behaviour to chimpanzee nut-cracking. No other individual in the study
performed the nut-cracking behaviour with a tool. Instead, the other (all adult) subjects
opened the nuts with their teeth (*bite,* see Tables 3 & 4). This *bite* behaviour of adults
continued even after the demonstration condition, in which the adults had the opportunity to
observe PD cracking nuts using the target behaviour. Indeed, the adults used primarily the *bite*
method, followed by the only other method they used: *hit with hand* (see more below).

Baseline condition

The *bite* method was the first method attempted in the baseline by all individuals, and the one
used in a higher number of sessions in this phase (*bite* was attempted in 20/20 (100%) of the
sessions, followed by *hit with hand* (6/20, 30%;), *anvil on floor* (4/20, 20%) and *step* (3/20,
15%;). All subjects attempted to open at least some nuts with their mouth, feet or hands in
most sessions, whereas only PD used the *anvil on floor* method, in 4/5, 80% of her sessions
(from the 2nd session of the baseline). Of these methods, only the *bite* and *anvil on floor* led to
successful kernel access. The adult females accessed an average of 4.4 out of 5 nut kernels per
session using the *bite* method (and were successful from the first session). PD also tried to
open nuts first with the mouth in her first session but failed to open them. However, in the
second and third sessions, PD tilted the large wooden block, placed a nut under the block, and
then dropped it on the nut. By using this method (*anvil on floor*), PD successfully opened six
nuts overall (the remaining nuts stayed unopened, as PD then reverted to attempting the *bite*
methodology unsuccessfully). In the fourth session, PD successfully cracked one nut with her
mouth but failed to open more nuts with either the *bite* or *anvil on floor* techniques. These
data may suggest that PD was relatively incapable of cracking open the nuts with her teeth. In
the last session, PD opened all five nuts using the *anvil on floor* method.

Anvil-locked condition (note: only PD was tested)

This condition (4 sessions) was carried out to examine whether PD would be able to change
from her technique of using the large wooden block (which had been devised as an anvil) to
using the smaller wooden pieces provided (which were designed to resemble the hammers
used by wild chimpanzees). From the first session, PD used the wooden hammers to perform
the target nut-cracking behaviour, albeit ignoring the large block as an anvil. Instead, PD
placed nuts on the floor (which was sufficiently hard), and then used the wooden hammer to
forcibly hit the nut until it opened (i.e., *hammer on floor* occurred in all sessions). Other than
hammer on floor, only *bite* and *hit with hand* were recorded in this condition. PD cracked 19
of 20 nuts using the *hammer on floor* method and no nuts using the *bite* method.

Demonstration condition

Despite being provided with live demonstrations from PD of the target nut-cracking
behaviour in the demonstration condition (15 sessions in total), none of the adult females

subsequently used any of the provided tools to open nuts. All adults continued to crack the
nuts using their teeth or by trying to open the nuts (unsuccessfully) using the *hit with hand*
method (*bite* 100%, 15/15 of the sessions; *hit with hand* 13%, 2/15 of the sessions). All the
nuts that were opened in the demonstration condition were opened with the *bite* behaviour.

Zürich study

Methods

Ethical approval

No specific ethical protocols were required when this study was carried out (in the 1980s).
However, the experiments were non-invasive, and were designed alongside the animals'
keepers to ensure the well-being of the animals. The study strictly adhered to the legal
requirements of Switzerland. All subjects were housed in indoor and outdoor enclosures
containing climbing structures and natural features. Subjects received their regularly
scheduled feedings and had access to enrichment devices and water *ad lib*. Subjects were
never food or water deprived for the purposes of this study. The research was conducted in
the subjects' familiar environment and any kind of stress to the animals was avoided.

Subjects

Research was carried out at Zürich zoo, Switzerland, between 1983-1984 by MF. Twelve
orangutans were tested (10 *Pongo abelii* and two *Pongo pygmaeus;* M_{age}=14; age range= 2-
30; 5F; see demographic information in Table 6). The keepers confirmed that none of the
subjects had any previous experience with coula nuts or Brazil nuts. However, the orangutans
had occasionally been provided with walnuts and coconuts. Whilst the walnuts could be easily
opened by biting, the coconuts were opened by hitting them against the hard floor or walls of
the enclosure. Despite the potential utility of a tool to open the coconuts, the keepers had
never observed the orangutans using hammer tools to open the coconuts and had never
demonstrated the hammer technique to them – for these or any other nuts. The two *Pongo*
pygmaeus were tested together, and the remaining subjects (all the *Pongo abelii*) were tested
in two groups (see Table 6 for a list of individuals in each group). These two groups did not
have visual access to each other.

Procedure

A baseline condition was implemented, in which a wooden hammer (25cm long, 8-10cm in diameter) was provided alongside shelled coula nuts, Brazil nuts and coconuts. No demonstrations on how to use the hammers to crack the nuts were provided before or during testing. The two *Pongo pygmaeus* received one coula nut per session for both individuals (one nut between both individuals was provided), five Brazil nuts each and one coconut each per session whilst the *Pongo abelii* received one coula nut and one coconut each and five Brazil nuts each per session. Each subject received one hammer. Initially no anvils were provided, as there were already objects inside the enclosure that could be used as anvils, such as logs with indentations that were in the orangutans' outside enclosure. However, once it became clear that the *Pongo pygmaeus* preferred to manipulate the testing equipment in the indoor enclosure, an extra anvil location was created by chiselling a hole into the concrete of the *Pongo pygmaeus'* indoor enclosure flooring. The *Pongo abelii* were not provided with additional anvils (due to zoo management requirements, as their enclosure had just been renovated and they had more anvil-type objects that could be used). The nuts were scattered on the floor of the enclosure one hour before feeding. Following the testing protocols adopted at the time, once the testing session started, focal animals (defined as the ones that were currently holding a nut at the start of the session) were followed and observations reported at 10second intervals. In the cases when more that one individual was manipulating a nut, both subjects were followed. However, this happened rarely, and only a maximum of two individuals at a time manipulated a nut, making the combined focal follows manageable. Each session lasted one hour, and 15 sessions were carried out in total. Once the session ended, the keepers removed the hammers and the nuts from the enclosure. No video recording was made of the tests, as the video equipment was found to be too distracting to the subjects by the experimenter (MF). Observations were recorded by voice instead using a handheld tape recorder (unfortunately these tapes no longer exist).

Results

The orangutans were observed practicing several different techniques to open the nuts (see Table 7 for an ethogram of these behaviours). Three individuals, one in each group (PG, AD & RJ), used the wooden hammer to crack open the coula nuts in the first testing session. In the subsequent second to seventh testing sessions, seven out of the total eight tested orangutans also demonstrated the same target hammering action (see Table 8 for the number

of interactions with nuts with the hammering method observed across subjects). Therefore, at
least three individuals spontaneously acquired nut-cracking with a hammer without having
any demonstrators available. The only individual who did not demonstrate nut-cracking with
a hammer was the youngest subject (HU), who rolled or threw away the nuts at the start of
each session, and did not seem motivated to manipulate the nuts or the hammer. Although
seven individuals attempted to crack nuts with the provided hammer, only three individuals
(RS, RJ, TR) were successful in opening the nuts with this method, potentially due to the fact
that not all individuals used an anvil to stabilise the nut. Most individuals only used the
hammer to crack the coula nuts, as they were the only type of nut that required the hammer
(both the Brazil nuts and the coconuts could be opened by biting or by hitting them on the
floor or walls). However, two individuals (PG and RJ) also used the hammer on the coconut,
of which only RJ was successful. PG was the only individual who was able to open the coula
nuts with his teeth from the first session, and continued with this method for the duration of
the study.

Discussion

Four naïve, unenculturated, captive orangutans spontaneously -without any copying or
training opportunities- cracked nuts using a wooden hammer as a tool. This finding suggests
that orangutans possess the individual ability to express nut-cracking in a similar way to wild
chimpanzees, and that copying is not required for this behaviour to be expressed in this
species. As orangutans do not crack nuts with tools in the wild, it is highly unlikely that any
of the orangutans in Leipzig and Zürich studies had previously observed this behaviour from
wild, or related, conspecifics. Furthermore, the subjects' experiences in previous experiments,
training and enrichment were discussed in detail at both institutions with their keepers, who
assured us that the subjects had never been trained or demonstrated how to crack nuts, or even
the hammering action, before testing. Although it is impossible to account for every minute of
a subject's life, keepers at both zoological institutions are heavily involved in the research
studies and enrichment programs with their animals, making them the most reliable source on
the animals' knowledge of behavioural forms. Therefore, based on all the information
available from the wild and the testing facilities, we are confident in assuming that all the
orangutans tested here were naïve to nut cracking (and the fact that nut-cracking appeared
across both institutions renders the naivety of the subjects even more likely).

This is not to say that the orangutans did not benefit from any previous experiences. Indeed,
all of the subjects had prior experience with nuts, and therefore knew that force could be
applied to the shells of the nuts to access the kernel inside. However, they only had
experience with walnuts and hazelnuts, which can be opened relatively easily by using the
teeth - even by juvenile orangutans -, and with coconuts, which can be opened by hitting them
against the floor or walls. In contrast, the coula nuts provided in the Zürich study and the
macadamia nuts provided in the Leipzig study require 2.8kn and 2.2kn to be opened,
respectively (Visalberghi et al., 2008), which makes these alternative methods unviable. This,
and the lack of suitable tool materials prior to our studies, may explain, at least in part, why
none of the subjects in these studies had ever been observed using tools to crack nuts before.

Candidate mechanisms behind nut-cracking in orangutans

The findings of both the Leipzig and the Zürich zoo studies suggest that nut-cracking with
tools can be acquired individually by naïve orangutans without the need of copying, provided
the availability of appropriate materials and the subjects' motivation to do so, as well as,
perhaps, a basic knowledge of the problem at hand (i.e., that shells often encase kernels and
that force can be applied to break them open). We are not suggesting that nut-cracking is a
hard-wired behaviour in orangutans. Although the ZLS hypothesis can also include such
cases, 'latent solutions' is an umbrella term that subsumes behaviours spanning from highly
genetically-determined behaviours to more learning-dependent behaviours, with the exception
of copying-dependent behaviours (Tennie et al., in press). In the case of orangutan nut-
cracking, there are several reasons to believe that more than instinct is at play. Firstly, despite
the fact that several orangutans in our studies performed nut-cracking, long-term field studies
with wild orangutans have not (yet) observed such behaviour (e.g., Krützen, Willems, &
van Schaik, 2011, although note that this might also be due to a lack of need to crack open
nuts). This suggests that nut-cracking is not (and is unlikely to have been) a critical behaviour
for the species' survival that may have been fixed in their genes throughout evolution
(although note that some of the abilities required for nut-cracking, such as the manipulative
tendency of the species, their hand affordance and some of their cognitive capacities may be
more genetically determined). Secondly, there was variability in the performance of subjects
in our studies. Despite experiencing the same conditions, not all the orangutans acquired the
behaviour within the time frame given (although we acknowledge that motivation plays a role
as well). Lastly, the orangutans in our studies demonstrated flexibility in their approach to the

problem at hand – as an example, PD attempted several different methods to access the
kernels before arriving at the target behavioural form of nut-cracking and, even after
discovering the target behaviour, she did not then use it in every session. Perhaps most
importantly, PD proved able to crack open nuts with a variety of tool use techniques.

If strong genetic predispositions and reliance on copying forms of social learning are
excluded as explanations for the acquisition of this behaviour, a plausible alternative
candidate mechanism is individual learning, supported by non-copying social learning. All
apes demonstrate impressive abilities for such type of learning (see Tomasello & Call 1997;
Whiten & Mesoudi, 2008 for an overview of these studies). Whilst individual learning
abilities probably involve some genetic predispositions (see above), they also rely on
cognitive skills that allow for considerable behavioural flexibility, including the capacity of
finding different solutions to a given problem. One example of this flexibility is PD's
performance in this study. In the baseline, before the locked-anvil condition, PD used the
provided large wooden block to crack open nuts, already demonstrating a similar tool use to
wild chimpanzee nut-cracking but using a different tool and action. PD might have initially
preferred to use the large block instead of the small wooden hammers as, although the former
required more effort when being lifted due to its large weight (approx. 50kg vs. 2.4kg), it did
not require the application of hitting force and speed to crack the nut, but could simply be
part-lifted and/or rolled, and then dropped on top of the nuts. Moreover, the large block may
have been easier to manipulate since its larger width required less precision when aiming to
hit the nut than a hammer does. Once the large block was rendered inaccessible in the locked-
anvil condition, however, PD flexibly switched her approach and used a hammer,
demonstrating the behavioural form of nut-cracking similar to that observed in some wild
chimpanzee populations (Biro et al., 2003; Boesch et al., 1994; Luncz & Boesch, 2014; Luncz
et al., 2012). In brief, individual learning, alongside some genetic predispositions, non-
copying social learning, and enhanced cognitive capacities that allow flexibility in the search
for solutions to problems, may drive the acquisition of nut-cracking in orangutans.

Potential explanations for the lack of reinnovation of the target behaviour by the remaining
orangutans

None of the adult orangutans in the Leipzig zoo study used tools to crack macadamia nuts,
and one juvenile individual in the Zürich zoo never attempted to use a hammer on coula nuts

either. In the case of the Leipzig study, the adults were immediately and consistently

successful in cracking open the nuts with their teeth, and continued doing so even after they

were exposed to five sessions of live demonstrations of nut-cracking with a tool by PD. One

explanation for the absence of nut-cracking with tools behaviour in these subjects could be

precisely the fact that, as we observed, the adults were strong enough to bite through the

shells of the nuts (note that, although macadamia nuts are hard, orangutans have a remarkable

bite strength; Daegling, 2007), which might have rendered the use of a tool superfluous for

them. Similarly, PG in the Zürich study was successful in opening the coula nuts with the bite

method and he persisted with this strategy (he only used the hammer on coconuts). In

contrast, the sub-adult PD attempted to bite nuts in the first session but failed, most likely

because she had not yet developed the same jaw strength as the adults in the group. Therefore,

PD may have been the only test subject motivated to find alternative methods to biting in

order to access the kernels, including the use of tools to open the nuts. According to this

explanation, if even harder nuts had been provided, rendering the bite methodology

impossible, the adults in the group might have also acquired the tool-use behaviour. This

hypothesis is supported by the findings of the Zürich study in which the harder coula nuts

were provided and adults were reported to acquire the behaviour. An alternative to the jaw

force explanation would be that age differences in inhibitory control (defined as the ability to

stop a planned or on-going thought or action; Carlson & Wang, 2007; see also Albiach-

Serrano, Guillén-Salazar, & Call, 2007; Amici, Aureli, & Call, 2008; Parrish et al., 2014) and

functional fixedness (defined as "*the disinclination to use familiar objects or methods in novel*

ways" Brosnan & Hopper, 2014, 2) encouraged PD to explore new solutions to the problem at

hand while preventing the adults in the Leipzig study from finding the same solution. The fact

that the only individual who did not attempt to use a hammer in the earlier Zürich study (HU)

was the youngest member of the group does not support this explanation, although it may

have been that this subject simply lacked the motivation to manipulate the nuts and the

hammers or even to consume the kernels. Indeed, HU immediately rolled the nuts away when

they were in his possession and did not attempt to retrieve them when other members of the

group took them.

Conclusion

Collectively, the results of our two studies demonstrate that individual learning (probably

aided by several factors, such as genetic predispositions and cognitive capacities that allow to

find solutions to problems in a flexible way) is sufficient for the acquisition of the behavioural

form of nut-cracking in orangutans. Although this study did not find direct evidence for non-

copying social learning increasing the frequency of the nut-cracking behaviour (as the older

orangutans may have been fixed in their alternative, successful method of cracking open the

nuts with their teeth in the Leipzig study), it is likely that, similar to other ape behaviours,

non-copying variants of social learning can increase and stabilise the frequency of nut-

cracking within populations – at least when these mechanisms apply across generations (see

also the discussion in Moore, 2013). Therefore, the behavioural form of nut-cracking could, in

principle, become another example of a SMR (Bandini & Tennie, 2017; 2019), for

orangutans. Indeed, it is possible that orangutans may one day be found to express (or have

expressed) nut-cracking behaviour in the wild - as a latent solution.

Acknowledgements

The authors are very grateful to the Wolfgang Köhler Primate Research Centre (WKPRC), in

Leipzig, Germany and to Zürich Zoo, in Switzerland, for providing the testing facilities. The

authors also thank William Daniel Snyder, Li Li and David Boysen for coding and second

coding, and Christian Nawroth, Yasmin Möbius and Daniela Hedwig for helpful discussions.

We also thank Josep Call, Heinz Gretscher, Vincent Müller, Franziska Zemke, Josefine

Kalbitz for support and feedback in the past, either directly or indirectly related to the current

studies and Christophe Boesch for allowing access to the Zürich zoo thesis. MF is grateful to

Hans Kummer for helping design the study, Christophe Boesch and Dr. G. Anzeberger for

sourcing the coula nuts from the Ivory Coast and Bruno Schnyder for help during testing. EB

and CT are supported by the Institutional Strategy of the University of Tübingen (Deutsche

Forschungsgemeinschaft, ZUK 63). At the time of writing, CT was also supported by the

ERC: This project has received funding from the European Research Council (ERC) under

the European Union's Horizon 2020 research and innovation programme (grant agreement n°

714658; STONECULT project).

Conflict of Interest

The authors have no conflict of interest to declare.

References cited

Albiach-Serrano, A., Guillén-Salazar, F., & Call, J. (2007). Mangabeys (*Cercocebus torquatus lunulatus*) solve the reverse contingency task without a modified procedure. *Animal Cognition*, *10*(4), 387–396.

Allritz, M., Tennie, C., & Call, J. (2013). Food washing and placer mining in captive great apes. *Primates*, *54*(4), 361–370. https://doi.org/10.1007/s10329-013-0355-5

Amici, F., Aureli, F., & Call, J. (2008). Fission-fusion dynamics, behavioral flexibility, and inhibitory control in primates. *Current Biology*, *18*(18), 1415–1419. https://doi.org/10.1016/J.CUB.2008.08.020

Bandini, E., & Tennie, C. (2019). Individual acquisition of 'stick pounding' behaviour by naïve chimpanzees. *American Journal of Primatology, 81*(6), e22987.

Bandini, E., & Tennie, C. (2017). Spontaneous reoccurrence of "scooping", a wild tool-use behaviour, in naïve chimpanzees. *PeerJ*, *5*, e3814. https://doi.org/10.7717/peerj.3814

Bandini, E., & Tennie, C. (2018). Naive, captive long-tailed macaques (*Macaca fascicularis fascicularis*) fail to individually and socially a tool-use behaviour. *Royal Society open science*, *5*(5), 171826.

Bandini, E., & Tennie, C. (2020). Exploring the role of individual learning in animal tool-use. *PeerJ*, *8*, e9877.

Bandini, E., & Harrison, R. A. (2020). Innovation in chimpanzees. *Biological Reviews Early view*.

Biro, D., Inoue-Nakamura, N., Tonooka, R., Yamakoshi, G., Sousa, C., & Matsuzawa, T. (2003). Cultural innovation and transmission of tool use in wild chimpanzees: evidence from field experiments. *Animal Cognition*, *6*(4), 213–223. https://doi.org/10.1007/s10071-003-0183-x

Boesch, C. (1991). Teaching among wild chimpanzees. *Animal Behaviour*, *41*(3), 530–532. https://doi.org/10.1016/S0003-3472(05)80857-7

Boesch, C., Marchesi, P., Marchesi, N., Fruth, B., & Joulian, F. (1994). Is nut cracking in wild chimpanzees a cultural behavior? *Journal of Human Evolution*, *26*(4), 325–338. https://doi.org/10.1006/jhev.1994.1020

Boesch, C., & Boesch, H. (1983). Optimisation of nut-cracking with natural hammers by wild chimpanzees. *Behaviour*, *83*(3–4), 265–286.

Boesch, C., & Boesch, H. (1990). Tool Use and Tool Making in Wild Chimpanzees. *Folia Primatologica*, *54*(1–2), 86–99. https://doi.org/10.1159/000156428

Brosnan, S. F., & Hopper, L. M. (2014). Psychological limits on animal innovation. *Animal*

Behaviour, *92*, 325–332. https://doi.org/10.1016/j.anbehav.2014.02.026

Carlson, S. M., & Wang, T. S. (2007). Inhibitory control and emotion regulation in preschool
children. *Cognitive Development*, *22*(4), 489–510.

Carvalho, S., Biro, D., McGrew, W. C., & Matsuzawa, T. (2009). Tool-composite reuse in
wild chimpanzees (*Pan troglodytes*): archaeologically invisible steps in the technological
evolution of early hominins? *Animal Cognition*, *12*(S1), 103–114.
https://doi.org/10.1007/s10071-009-0271-7

Clay, Z., & Tennie, C. (2017). Is overimitation a uniquely human phenomenon? Insights from
human children as compared to bonobos. *Child Development, 89(5), 1535-1544.*
https://doi.org/10.1111/cdev.12857

Cohen, J. (1968). Weighted kappa: Nominal scale agreement provision for scaled
disagreement or partial credit. *Psychological Bulletin*, *70*(4), 213–220.
https://doi.org/10.1037/h0026256

Corp, N., & Byrne, R. W. (2002). The ontogeny of manual skill in wild chimpanzees:
evidence from feeding on the fruit of *Saba florida*. *Behaviour, 139,* 137–168.

Daegling, D. J. (2007). Morphometric estimation of torsional stiffness and strength in primate
mandibles. *American Journal of Physical Anthropology*, *132*(2), 261–266.
https://doi.org/10.1002/ajpa.20508

De Waal, F. (2008). *The ape and the sushi master: Cultural reflections of a primatologist*.
Basic Books: New York.

Foucart, J., Bril, B., Hirata, S., Monimura, N., Houki, C., Ueno, Y., & Matsuzawa, T. (2005).
A preliminary analysys of nut-cracking movements in a captive chimpanzee: adaptation
to the properties of tools and nuts. *Stone Knapping: The Necessary Conditions for a*
Uniquely Hominin Behaviour, 147–157.

Funk, M. (1985).

Fox, A., van Schaik, C. P., Sitompul, A., & Wright, D. N. (2004). Intra-and interpopulational
differences in orangutan (*Pongo pygmaeus*) activity and diet: Implications for the
invention of tool use. *American Journal of Physical Anthropology*, *125*(2), 162–174.
https://doi.org/10.1002/ajpa.10386

Fox, A, Sitompul, A. F., & van Schaik, C. P. (1999). Intelligent tool use in wild Sumatran
orangutans. *The Mentality of Gorillas and Orangutans*, *480*, 99–116.

Gumert, M. D., Kluck, M., & Malaivijitnond, S. (2009). The physical characteristics and
usage patterns of stone axe and pounding hammers used by long☐tailed macaques in the
Andaman Sea region of Thailand. *American Journal of Primatology, 71*(7), 594-608.

Hecht, E. E., Gutman, D. A., Preuss, T. M., Sanchez, M. M., Parr, L. A., & Rilling, J. K.
(2013). Process versus product in social learning: Comparative diffusion tensor imaging
of neural systems for action execution–observation matching in macaques, chimpanzees,
and humans. *Cerebral Cortex*, *23*(5), 1014–1024. https://doi.org/10.1093/cercor/bhs097

Henrich, J., & Tennie, C. (2017). Cultural Evolution in Chimpanzees and Humans.
Chimpanzees and Human Evolution. Harvard University Press.

Inoue-Nakamura, N., & Matsuzawa, T. (1997). Development of stone tool use by wild
chimpanzees (*Pan troglodytes*). *Journal of Comparative Psychology*, *111*(2), 159–173.

Kenward, B., Schloegl, C., Rutz, C., Weir, A., Bugnyar, T., & Kacelnik, A. (2011). On the
evolutionary and ontogenetic origins of tool-oriented behaviour in New Caledonian
crows (*Corvus moneduloides*). *Biological Journal of the Linnean Society*, *102*(4), 870–
877. https://doi.org/10.1111/j.1095-8312.2011.01613.x

Krützen, M., Willems, E. P., & van Schaik, C. P. (2011). Culture and geographic variation in
orangutan behavior. *Current Biology*, *21*(21), 1808–1812.
https://doi.org/10.1016/J.CUB.2011.09.017

Lonsdorf, E. V. (2013). The role of mothers in the development of complex skills in
chimpanzees. In *Building Babies*. Springer, New York, NY, 2013.

Luncz, L. V., & Boesch, C. (2014). Tradition over trend: Neighboring chimpanzee
communities maintain differences in cultural behavior despite frequent immigration of
adult females. *American Journal of Primatology*, *76*(7), 649–657.
https://doi.org/10.1002/ajp.22259

Luncz, L. V., Mundry, R., & Boesch, C. (2012). Evidence for cultural differences between
neighboring chimpanzee communities. *Current Biology*, *22*(10), 922–926.
https://doi.org/10.1016/j.cub.2012.03.031

Menzel, C., Fowler, A., Tennie, C., & Call, J. (2013). Leaf surface roughness elicits leaf
swallowing behavior in captive chimpanzees (*Pan troglodytes*) and bonobos (*Pan
paniscus*), but not in gorillas (*Gorilla gorilla*) or orangutans (*Pongo abelii*).
International Journal of Primatology, *34*(3), 533–553. https://doi.org/10.1007/s10764-
013-9679-7

Mesoudi, A., & Whiten, A. (2008). The multiple roles of cultural transmission experiments in
understanding human cultural evolution. *Philosophical Transactions of the Royal Society
B: Biological Sciences*, *363*(1509), 3489-3501

Meulman, E. J. M., Sanz, C. M., Visalberghi, E., & van Schaik, C. P. (2012). The role of
terrestriality in promoting primate technology. *Evolutionary Anthropology*, *21*(2), 58–68.
https://doi.org/10.1002/evan.21304

Meulman, E. J. M., & van Schaik, C. P. (2013). Orangutan tool use and the evolution of
technology. *Tool Use in Animals: Cognition and Ecology*, 176–202.

Moore, R. (2013). Social learning and teaching in chimpanzees. *Biology & Philosophy*, *28*(6),
879-901. https://doi.org/10.1007/s10539-013-9394-y

Neadle, D., Allritz, M., & Tennie, C. (2017). Food cleaning in gorillas: Social learning is a
possibility but not a necessity. *PLOS ONE*, *12*(12), e0188866.
https://doi.org/10.1371/journal.pone.0188866

Neadle, D., Bandini, E., & Tennie, C. (2020). Testing the individual and social learning abilities of task-naïve captive chimpanzees (*Pan troglodytes sp.)* in a nut-cracking task. *PeerJ*, *8*, e8734.

Neldner, K., Reindl, E., Tennie, C., Grant, J., Tomaselli, K., & Nielsen, M. (2020). A cross-cultural investigation of young children's spontaneous invention of tool use behaviours. *Royal Society Open Science*, *7*(5), 192240.

Ottoni, E. B., & Mannu, M. (2001). Semi-free ranging tufted capuchin monkeys (*Cebus apella*) spontaneously use tools to crack open nuts. *International Journal of Primatology*, *22*(3), 347–358.

Parker, S. T., & Gibson, K. R. (1977). Object manipulation, tool use and sensorimotor intelligence as feeding adaptations in cebus monkeys and great apes. *Journal of Human Evolution*, *6*(7), 623–641. https://doi.org/10.1016/S0047-2484(77)80135-8

Parrish, A. E., Perdue, B. M., Stromberg, E. E., Bania, A. E., Evans, T. A., & Beran, M. J. (2014). Delay of gratification by orangutans (*Pongo pygmaeus*) in the accumulation task. *Journal of Comparative Psychology*, *128*(2), 209.

Picheta, R. (2020). The UN wants to protect these chimps' unique culture. CNN. Retrieved from: https://edition.cnn.com/2020/03/09/africa/chimpanzees-west-africa-un-conservation-scli-scn-intl/index.html

Pope, S., Taglialatela, J., Skiba, S., & Hopkins, W. D. (2017). Changes in fronto-parieto-temporal connectivity following Do-As-I-Do imitation training in chimpanzees (*Pan troglodytes*). *Journal of cognitive neuroscience*, *30*(3), 421-431.

Proffitt, T., Haslam, M., Mercader, J. F., Boesch, C., & Luncz, L. V. (2018). Revisiting Panda 100, the first archaeological chimpanzee nut-cracking site. *Journal of Human Evolution*. https://doi.org/10.1016/J.JHEVOL.2018.04.016

Reindl, E., Beck, S. R., Apperly, I. A., & Tennie, C. (2016). Young children spontaneously invent wild great apes' tool-use behaviours. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1825), 20152402. https://doi.org/10.1098/rspb.2015.2402

Schuppli, C., Meulman, E. J. M., Forss, S. I. F., Aprilinayati, F., van Noordwijk, M. A., & van Schaik, C. P. (2016). Observational social learning and socially induced practice of routine skills in immature wild orang-utans. *Animal Behaviour*, *119*, 87–98. https://doi.org/10.1016/J.ANBEHAV.2016.06.014

Shumaker, R. W., Walkup, K. R., Beck, B. B., & Burghardt, G. M. (2011). *Animal Tool Behavior: The Use and Manufacture of Tools by Animals*. Johns Hopkins University Press.

Tan, A. W. Y. (2017). From play to proficiency: The ontogeny of stone-tool use in coastal-foraging long-tailed macaques (*Macaca fascicularis*) from a comparative perception-action perspective. *Journal of Comparative Psychology*, *131*(2), 89–114. https://doi.org/10.1037/com0000068

Tennie, C., Call, J., & Tomasello, M. (2009). Ratcheting up the ratchet: on the evolution of
cumulative culture. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, *364*(1528), 2405–2415. https://doi.org/10.1098/rstb.2009.0052

Tennie, C., Call, J., & Tomasello, M. (2012). Untrained chimpanzees (*Pan troglodytes*
schweinfurthii) fail to imitate novel actions. *PLoS ONE*, *7*(8), e41548.
https://doi.org/10.1371/journal.pone.0041548

Tennie, C., & Hedwig, D. (2009). How latent solution experiments can help to study
differences between human culture and primate traditions. *Primatology: Theories,*
Methods and Research. Hauppauge, NY: Nova Science Publishers, 95–112.

Tennie, C., Hedwig, D., Call, J., & Tomasello, M. (2008). An experimental study of nettle
feeding in captive gorillas. *American Journal of Primatology*, *70*(6), 584–593.
https://doi.org/10.1002/ajp.20532

Tomasello, M., & Call, J. (1997). *Primate cognition*. Oxford University Press.

Tomasello, M., Call, J., Warren, J., Frost, G. T., Carpenter, M., & Nagell, K. (1997). The
ontogeny of chimpanzee gestural signals: A comparison across groups and generations.
Evolution of communication, *1*(2), 223-259.

van Schaik, C. P., Deaner, R. O., & Merrill, M. Y. (1999). The conditions for tool use in
primates: implications for the evolution of material culture. *Journal of Human Evolution*,
36(6), 719–741. https://doi.org/10.1006/jhev.1999.0304

van Schaik, C. P., & Knott, C. D. (2001). Geographic variation in tool use on Neesia fruits in
orangutans. *American Journal of Physical Anthropology, 114*(4), 331–342.

Visalberghi, E., Sabbatini, G., Spagnoletti, N., Andrade, F. R. D., Ottoni, E., Izar, P., &
Fragaszy, D. (2008). Physical properties of palm fruits processed with tools by wild
bearded capuchins (*Cebus libidinosus*). *American Journal of Primatology*, *70*(8), 884–
891. https://doi.org/10.1002/ajp.20578

Visalberghi, E. (1987). Acquisition of nut-cracking behaviour by two capuchin monkeys
(*Cebus apella*). *Folia Primatologica*, *49*(3–4), 168–181.
https://doi.org/10.1159/000156320

Whiten, A., Tutin, C. E. G., McGrew, W. C., & Wrangham, R. W. (2001). Charting cultural
variation in chimpanzees. *Behaviour*, *138*(11-12), 1481-1516.
https://doi.org/10.1163/156853901317367717

Whiten, A. (2017). Culture extends the scope of evolutionary biology in the great apes.
Proceedings of the National Academy of Sciences, *114*(30), 7790–7797.
https://doi.org/10.1073/pnas.1620733114

833   *Table 1: Demographic information on the subjects included in the Leipzig zoo study*

Name	Species	Sex	Date of birth	Place of birth	Breeding
Dokana (DK)	*Pongo abelii*	F	31/01/1989	Dresden, DE	Parent
Padana (PD)	*Pongo abelii*	F	18/11/1997	Leipzig, DE	Parent
Pini (PI)	*Pongo abelii*	F	30/06/1988	Leipzig, DE	Parent
Dunja (DJ)	*Pongo abelii*	F	19/04/1990	Leipzig, DE	Hand

835   *Table 2: Table showing the number of conditions, sessions and role of each subject from the*
836   *Leipzig zoo study*

Subject	Conditions participated in	Number of sessions per condition	Role
DK	Baseline, Demonstration	Baseline: 5 Demonstration: 5	Subject Subject
PD	Baseline, Locked-anvil, Demonstration	Baseline: 5 Locked-anvil: 4 Demonstration: 5	Subject Subject Conspecific model in the demonstration condition
PI	Baseline, Demonstration	Baseline: 5 Demonstration: 5	Subject Subject
DJ	Baseline, Demonstration	Baseline: 5 Demonstration: 5	Subject Subject

838   *Table 3: Ethogram of behaviours directed towards the nuts by subjects in the Leipzig zoo*
839   *study across all conditions*

Behaviour	Description	Number of subjects & name	First session in which the behaviour was observed	Condition in which the behaviour was observed	Successful for opening nuts	Percentage of success per individual
Bite	The subject inserts the nut in its mouth	All subjects	1[st] session baseline	All conditions	Yes	DK: 100% DJ: 100% PD: 11% PI: 100%

Hit with Hand	The subject hits the nut with its hand against a hard surface	2 (PD & DK)	1st session baseline	All conditions	No	N/A
Step	The subject hits the nut with its foot against a hard surface	2 (PD & PI)	1st session baseline	Only baseline condition	No	N/A
Anvil on Floor	The subject tilts the anvil and either drops it or rolls it on the nut that is on the floor	1 (PD)	2nd session baseline	Only baseline condition	Yes	PD: 75%
Hammer on Floor	The subject lifts the hammer and drops it on the nut which is on the floor	1 (PD)	1st session locked anvil	Only locked anvil condition	Yes	PD: 100%

 Table 4: Count and percentage of each behaviour directed towards the nuts per individual per condition and sessions in the Leipzig zoo study

Condition	Subject	Number of sessions	Number of sessions in which Bite was observed	%	Number of sessions in which Hit with Hand was observed	%	Number of sessions in which Step was observed	%	Number of tsessions in which Anvil on Floor was observed	%	Number of sessions in which Hammer on Floor was observed	%
	DK	5	5	100%	3	60%	0	0%	0	0%	0	0%
	DJ	5	5	100%	0	0%	0	0%	0	0%	0	0%
Baseline	PD	5	5	100%	3	60%	2	40%	4	80%	0	0%
	PI	5	5	100%	0	0%	1	20%	0	0%	0	0%
Baseline Total		20	20	100%	6	30%	3	15%	4	20%	0	0%
Anvil Locked	PD	4	4	100%	2	50%	0	0%	0	0%	4	100%
Anvil Locked Total		4	4	100%	2	50%	0	0%	0	0%	4	100%
	DK	5	5	100%	0	0%	0	0%	0	0%	0	0%
Demonstration	DJ	5	5	100%	2	40%	0	0%	0	0%	0	0%
Social	PI	5	5	100%	0	0%	0	0%	0	0%	0	0%
Demo Total		15	15	100%	2	13%	0	0%	0	0%	0	0%

Table 6: Demographic information on the subjects included in the Zürich zoo study

Name	Species	Testing Group	Sex	Date of birth	Place of birth	Breeding
Rosa (RS)	*Pongo pygmaeus*	1	F	1958	Wild Caught	Parent
Adam (AD)	*Pongo pygmaeus*	1	M	1953	Wild Caught	Parent
Pongo (PG)	*Pongo abelii*	2	M	1961	Wild Caught	Parent
Lea (LA)	*Pongo abelii*	2	F	22/08/1967	Zürich, CH	Hand
Farida (FA)	*Pongo abelii*	3	F	3/5/1979	Zürich, CH	Hand/Parent
Hantu (HU)	*Pongo abelii*	2	M	3/5/1981	Zürich, CH	Parent
Radja (RJ)	*Pongo abelii*	3	F	19/1/1973	Zürich, CH	Parent
Timor (TR)	*Pongo abelii*	2	F	23/11/1973	Basel, CH	Parent

Table 7: Ethogram of behaviours directed towards the nuts observed in the Zürich study

Behaviour	Description	Successful for opening nut?
Bite	The subject inserts the nut in its mouth and presses it between the teeth	Yes
Shake	The subject holds the nut in one or both hands and moves it back and forth quickly	No
Hit with hand/foot	The subject strikes the nut with either hands or feet, without using any tools	No
Press nut against teeth	After positioning the nut on the outer surface of the teeth, the subject presses it against the teeth either with the hands directly or by using a hard object such as the iron bars of the cage or wooden sticks	No
Scratch nut	The subject uses the nails to scratch the nut by damaging the surface of the shell	No
Rub nut	The subject moves the nut back and forth on a surface such as the ground or the walls	No
Beat against substrate	The subject holds the nut in one or both hands or feet and hits it against the floor or the walls or any other element in the enclosure	No
Press nut	The subject uses its weight or force on the nut without causing a sudden impact on the nut	No
Bounce nut	The subject throws or drops the nut onto the floor	No
Poke with branch	The subject pokes the shell of the nut with a branch	No
Wrapping nut	The subject wraps the nut with objects such as cardboard, wood wool or cotton bags	No
Hammering	The subject uses a hard object such as the wooden hammer to pound against the nut	Yes

Subject	Number of interactions with nuts	Number of hammering bouts	Number of hammering bouts before first success or at all if no success	Number of successes with hammering	Total number of successes	Proportion of hammering bouts in %
RS	41	390	3	30	37	57.5
AD	9	18	18	0	0	2.6
PG	24	1	1	0	11	0.1
LA	9	3	3	0	8	0.4

RJ	17	72	22	2	8	10.6
TR	16	192	13	4	7	28.3
FA	11	2	2	0	0	0.3
HU	8	0	0	0	0	0.0
All subjects	120	678	62	36	74	100

Table 8: Total number of nut manipulation and hammering bouts in the Zürich zoo study

Figure 1: Photograph of the testing apparatus with the anvil, wooden hammers and macadamia nuts in the Leipzig zoo study.