Bioinformatics Study on Structural Protein of Severe Acute Respiratory Syndrome Coronavirus 2 (Sars-Cov-2) For Better Understanding the Vaccine Development

[1] *Sumaira Gulzar*,*

Department of Biotechnology and Bioinformatics International Islamic University Islamabad Pakistan

sumaira.msbt317@iiu.edu.pk

[2]*Saqib Husssain*,*

International Center for Chemical and Biological Sciences University of Karachi Pakistan

Saqib670@gmail.com

Abstract

Novel coronavirus 2019 (2019-nCoV), also known as SARS-CoV-2), leads high morbidity and mortality in global epidemics. Four structural protein (surface glycoprotein (QIQ22760.1), envelop glycoprotein (QIQ22762.1), nucleocapsid phosphoprotein (QIQ22768.1) and membrane glycoprotein (QIQ22763.1)) of SARS-CoV-2 are extracted from the NCBI database and further analyzed with ExPASy ProtParam tool. Phenyl alanine and serine are absolutely missing in all proteins. Histidine and glutamic acid is absent in envelope protein and cystein lacks in nucleocapsid phosphoprotein. Lucien is the highest in envelope, surface and membrane glycoprotein that is an optimal environment for rapid virus fixation on host cell's surface to the receptor molecule. Transmembrane region prediction was performed by SOSUI server. For all structural proteins, except nucleocapsid Phosphoprotein, the trans-membrane prediction indicates that the virus can enter the host easily. Domain analysis was done by SMART tool. Domain information helps in the function of the viral protein. Lastly, the 3D structure prediction was carried out by Swiss Model and the result validation was achieved by PROCHECK. This fairly simple method may help us understand how antivirals and vaccines could be produced against it Such models are the starting point of the community for structural drug and vaccine designs as well as virtual computational screening.

Keywords – **SARS-CoV-2,** *Structural protein, Transmembrane, SMART tool, Swiss Model*

Introduction

 The latest 2019–nCoV, now officially known as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), was established as the responsible pandemic(Gorbalenya et al., 2020). Infections of (SARS-CoV-2) have been reported in more than 200 countries including Pakistan. The first genome sequence of SARS-CoV-2 to be published from Pakistan and is now available on NCBI https://www.ncbi.nlm.nih.gov/nuccore/MT240479, GISAID and NEXTRAIN.

Coronaviruses, a genus belonging to the Coronaviridae family, contain the largest genome (26.4 kb to 31.7 kb) of RNA viruses with a diameter of 120–160 nm, consisting of a single-stranded positive-sense RNA molecule (de Groot et al., 2012)

 The broad genome has given this family of viruses extra plasticity in their adaptation and alteration of genes. The G + C content of coronavirus genomes ranges from 32% TO 43% The genome consists of seven genes arranged in sequence [5′- replicase ORF1ab, spike (S), envelope (E), membrane (M), nucleocapside (N)-3′] with small untranslated regions in both termini and additional ORFs in each subgroup of coronavirus. (Woo et al., 2009)

5 'Non-structural protein coding regions consisting of two-thirds genome replicase genes 1 and structural and non-essential gene. 2-7 regions consisting of structural and non-essential protein coding regions [5]. The genes are translated from genomic mRNA 2 -7. Subgenomic RNAs encode the main Surface Proteins (S), Envelope Protein (E), Membranes Protein (M), and Nucleocapside Protein (Ng et al., 2003)

Surface protein of SARS CoV-2 join ACE2 (angiotensin- converting enzyme2) and to infect cells. After this initial process, Surface Protein must be produced with an enzyme known as protease to complete entry into the cell. SARS CoV-2 uses TMPRSS 2 in the same way as SARS-CoV is used to complete this process (Graham et al., 2010)

Surface Glycoproteins are outside of the virion and give the typical shape to the virion. The S proteins form homotrimers that allow sun-like morphologies to be developed that give the name Coronaviruses [10]. Via the C-terminal transmembrane regions, S proteins bind to the virion membrane and interact with M proteins. Virions can be attached through the N-terminus of the S proteins to different surface receptors in the host cell's plasma membrane. The S protein is the receptor-binder and viral input in host cells and is therefore a major therapeutic objective (Du et al., 2017)

Membrane Glycoproteins proteins inside the Golgi system are glycosylated. The modification of the M protein is essential for the virion to attach into the cell and make antigenic protein (de Haan et al., 2003)

The protein M plays a key role in the cell's regeneration of virions. N protein forms a complex by binding to genomic RNA and M protein activates the development of interacting virions in this intermediate endoplasmic reticulum-Golgi interface (ERGIC) compartment with this complex (de Haan et al.,1998 ; Escorset al., 2001)

Envelop Glycoproteins are tiny proteins consisting of around 76 to 109 amino acids. Approximately 30 amino acids in the E protein N-terminus allow attachment to the virus membrane (Raamsman et al., 2000)

Additionally, coronavirus E proteins play a critical role in virion assembly and morphogenesis within the cell. For one research coronavirus E and M proteins were expressed for conjunction with vectors of mammalian expression to form virus-like structures inside the cell (Baudoux et al., 1998; Vennema et al., 1996)

In another review, the ability of recombinant mouse hepatitis virus (MHV) and SARS viruses to generate E protein expression in the genome to sustain this status has been significantly reduced (Kuo et al., 2003; DeDiego. et al 2007)

N Proteins are helix-capable phosphoproteins with a versatile viral genomic RNA structure. It plays an important role in coronavirus virion assembly, replication and transcription, as the N protein locates both the coronavirus replication / transcription region and the ERGIC region where the virus is collected (Stertzet al., 2007)

In this paper, we tried to describe the four structural proteins of SARS-CoV-2 by the use of bioinformatics tools

Materials and Methods

Four structural proteins of SARS COV-2 included in this research named as surface glycoprotein (QIQ22760.1), envelop glycoprotein (QIQ22762.1), nucleocapsid phosphoprotein (QIQ22768.1), membrane glycoprotein (QIQ22763.1) available in the NCBI was retrieved. For further study of bioinformatics, the FASTA sequence was selected and used.

Protein statistics

Using the Protparam method, protein statistics were reviewed. ProtParam method enables the measurement of various physicochemical protein sequence parameters. Parameters included

molecular weight, Theoretical Pi (Isoelectric point), Half-life calculation, instability index, aliphatic index and Grand range of hydropathicity (GRAVY)

Trans-membrane sequence prediction

SOSUI tools from EXPASY repository are used for predicting transmembrane. Results normally return within 1 min. It provides the following production number of TMHs with sequence and protein type

Functional Site Prediction

One of the main goals of molecular biology is functional assignment to protein. Protein functional sites that are responsible for or perform all of the essential protein functions are more conserved than other regions over the time of evolution. These accessible websites are called domains. SMART Internet server was forecasting domains

3D Model predictions and validation

The SARS COV-2 structural protein amino acid sequences are used as targets for homology modeling using the SWISS-MODEL server. Depending on the Global Model Quality Estimation (GMQE) and QMEAN, the top-ranked models are further analyzed and sorted. GMQE is a quality estimate that combines the target – template alignment properties with the template search method. The corresponding GMQE value is given as 0 to 1. The QMEAN Z-score gives a global estimation of the "degree of nativeness" of the structural characteristics observed in the model (Benkert et al., 2011) QMEAN Z-scores about zero indicate strong compatibility between the structure of the model and similar-size experimental structures. Scores of -4.0 or below are indicative of poor quality models. The overall stereochemical output was checked by PROCHECK (Colovos et al., 1993) including torsional angles to the backbone through the Ramachandran plot

Results

Protein Statistics

Phenyl alanine and serine are absolutely missing in all proteins tested as displayed in chemical parameters (Table 1). Histidine and glutamic acid is absent in envelope protein and cystein lacks in nucleocapsid phosphoprotein. lucien quantity is more in envelop protein, surface and membrane glycoprotein while glycine is more in nucleocapsid phosphoprotein

The physical parameters (Table 2) show that Surface protein contain highest amino acids (1273), negatively charged residues (110) positively charged residues (103), EC (148960) and molecular weight (141178.47). Higher molecular weight of the surface glycoprotein suggests that its tertiary structure may contain strong amino acid side chains. All three Proteins are basic in nature, excluding surface glycoprotein, as the isoelectric point value is more than 9.The

instability index value greater than 40 is considered unstable (Lagerstrom et al.,) This is derived from the Instability index surface, membrane and envelop glycoprotein is stable and nucleocapsid phosphoprotein is highly unstable .the range of aliphatic index from 33.01 -55.09 that suggests a tendency to be sensitive to a wide range of temperatures (Ikai ,1980) and GRAVY value shows protein's hydropathicity and whether the nature protein side chains are hydrophilic or hydrophobic (Kyte et al., 1982) Leaving the surface

Glycoprotein and nucleocapsid phosphoprotein all other protein are hydrophobic in nature

Trans-membrane sequence prediction

Analysis of the transmembrane protein reveals (Table 3) that it is absent in nucleocapsid phosphoprotein. The other three proteins show transmembrane sequence. All three other proteins show different sequence from each other. Membrane glycoprotein have three transmembrane region yet there is a sequence of length 22 and 23 amino acids in all the proteins that show trans-membrane. All the proteins have different C and N terminals

Functional Site Prediction

SMART results (Table 4) clearly indicates the Surface Glycoprotein contains two domain .One is Pfam:Spike_rec_bind starts from 330 to 583 residues predicted with the e-value (expected value) second is Pfam:Corona_S2 domain starts from 671 to 1279 residues predicted with the e-value Envelop Glycoprotein contains one domain Pfam:NS3_envE starts from 1 to 74 residues predicted with the e-value 7.7e-8.Membrane Glycoprotein contains one domain Pfam:Corona_Mstarts from 4 to 221 residues predicted with the e-value 1.3e-95.Nucleocapsid Phosphoprotein contains one domain Pfam:Corona_nucleoca starts from 14 to 337 residues predicted with the e-value 1.2e-164

3D Model predictions and validation

(Table 5 and Figure. 2) shows the evaluation of the 3D model that all four proteins have satisfactory model with QMEAN > 0.Model validity research Shows that the models are at the optimum point, just like in the Ramchandran series. (Figure 3).

The SWISS-MODEL system was used for protein structure homology modelling and alignments for all structural proteins of SARS COV 2. First, the right protein template structures in PDB were selected using the given measures: should high reporting of the template (i.e. > 60% target associated with the template) and sequence uniqueness > 35%. Instead, as an initial criterion, we used the GMQE and QMEAN4 scoring method to distinguish well from poor ones

We also carried out PROCHECK analysis to measures protein structure stereochemical consistency by comparing geometry with residue-by-residue and particularly structural geometry.

There should be more than 90 % amino acid for good quality model according to the PROCHECK standard acid resided in the most favored regions. Table 5 and Figure 4 indicate that percentage of modeled proteins residues in most favoured regions (red) is 83%─91.4%, percentage of residues of the modeled proteins in additional allowed regions (yellow) is 8.6─15.6% , percentage of residues of the modeled proteins in generously allowed regions (beige) is 0.0─1.5%, and percentage of residues in disallowed regions (white) is 0.0%. These findings show that the models are generated overall stereochemical properties were highly stable, and that future molecular modelling studies can benefit from the models. Of those four proteins, the Ramachandran plots provide more proof of their acceptability (Figure 3).

Discussion

The physical parameters shows that Surface protein contain highest amino acids, negatively charged residues, positively charged residues, EC and molecular weight than other proteins. The chemical factors show in proteins phenyl alanine and serine is totally absent in all structural proteins, there is a lack of histidine and glutamic acid in the protein envelope and a lack of cysteine in the nucleocapside phosphoprotein. The envelope, surface and membrane glycoprotein is rich in Leucine so have more affinity to the host cell receptor surface as stated in Luo et al, 1999. Although glycine is more in nucleocapside phosphoprotein, researchers should be able to isolate the protein with little effort along with these data and iso-electric point value. Higher molecular weight of the surface glycoprotein suggests that its tertiary structure may contain strong amino acid side chains.Instability index surface, membrance and envelop glycoprotein is stable and nucleocapsid phosphoprotein is highly unstable Leaving the surface Glycoprotaein and
 nucleocapsid phosphoprotein all other protein are hydrophobic in nature. The prediction of trans-membrane in all structural proteins except Nucleocapsid Phosphoprotein shows the virus ability to reach the host with ease. This fairly simple method may help us understand how antivirals and vaccines could be produced against it Furthermore, as a beginning for docking studies (small and large scale) we are now offering homology models, Diverse knowledge of the molecular biology of SARS COV_2 is required to

learn more.

Developing technologies will gain

valuable insight into the structure of the protein in order to determine how protein disease induces, and understanding the relationship between protein-protein and protein RNA would greatly enhance our ability to develop vaccines. Meanwhile, methods of molecular simulation provide important solutions to the struggle

Acknowledgment

All authors acknowledge and thank their respective Institutes and Universities

References

Baudoux P, Carrat C, Besnardeau L, Charley B, Laude H. (1998) . Coronavirus pseudo particles formed with recombinant M and E proteins induce alpha interferon synthesis by leukocytesJ Virol; 72: 8636-8643. https://goo.gl/Fzrwdb

Benkert, P., Biasini, M., Schwede, T. (2011). Toward the estimation of the absolute quality of individual protein structure models. Bioinformatics 27, 343-350

Colovos, C. and T.O. Yeates. (1993). Verification of protein structures: Patterns of nonbonded atomic interactions. Prot. Sci., 2: 1511-1519.

 de Groot RJ, Baker SC, Baric R, Enjuanes L, Gorbalenya AE, Holmes KV, Perlman S, Poon LL, Rottier PJM, Talbot PJ, Woo PCY, Ziebuhr J. Family Coronaviridae. In: King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ, (2012.) Virus taxonomy. Ninth report of the international committee on taxonomy of viruses Amsterdam: Elsevier Academic Press; pp. 806–828

De Haan CA, de Wit M, Kuo L, Montalto-Morrison C, Haagmans BL, Weiss SR, et al. (2003) The glycosylation status of the murine hepatitis coronavirus M protein aff ects the interferogenic capacity of the virus in vitro and its ability to replicate in the liver but not the brainVirology; 312: 395-406. https://goo.gl/AYsmyB

De Haan CA, Masters PS, Lili Kuo, Harry Vennema, Peter JM, Rottier. (1998) Coronavirus particle assembly: primary structure requirements of the membrane protein J Virol.; 72: 6838-6850.

DeDiego ML, Alvarez E, Almazan F, Rejas MT, Lamirande E, Roberts A. (2007). A severe acute respiratory syndrome coronavirus that lacks the E gene is attenuated in vitro and in vivo J Virol.; 81: 1701-13. https://goo.gl/zv5cyu

Du, L.; Yang, Y.; Zhou, Y.; Lu, L.; Li, F.; Jiang, S. (2017). MERS-CoV spike protein: A key target for antivirals Expert Opin. Ther.Targets, 21, 131–143.

Graham RL, Baric RS. (2010). Recombination, reservoirs, and the modular spike: mechanisms of coronavirus cross-species transmission J Virol.; 84: 3134-3146. https://goo.gl/89BVoZ

Escors D, Ortego J, Enjuanes L. (2001) The membrane M protein of the transmissible gastroenteritis coronavirus binds to the internal core through the carboxyterminus Adv Exp Med Biol; 494: 589-593. https://goo.gl/Erg1Yi

Gorbalenya A.E, Baker S.C, RS Baric RS, Groot RJd, Drosten. C, AA Gulyaeva AA, Haagmans B.L, Lauber C, Leontovich AM, Neuman BW. (2020) Severe acute respiratory syndrome-related coronavirus-The species and its viruses, a statement of the Coronavirus Study Group

Ikai, A., (1980). Thermostability and aliphatic index of globular proteins. J. Biochem., 88: 1895-1898

Kuo L, Masters PS. (2003) .The small envelope protein E is not essential for murine coronavirus replication J Virol.; 77: 4597-4608. https://goo.gl/RT1og2

Kyte, J. and R.E. Doolittle, (1982). A simple method for displaying the hydropathic character of a protein. J. Mol. Biol., 157: 105-132

Lagerstrom, M.C. and H.B. Schioth, (2008). Structural diversity of G protein-coupled receptors and significance for drug discovery. Nat. Rev. Drug Discov., 7: 339-357

M. Hoffmann, H. Kleine-Weber, S. Schroeder, N. Krüger, T. Herrler, S. Erichsen. (2020). SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor Cell

Masters PS. The molecular biology of coronaviruses (2006) Adv Virus Res; 66: 193-292.
Ng ML, Tan SH, See EE, Ooi EE, Ling AE. (2003) .Proliferative growth of SARS coronavirus in Vero E6 cells J Gen Virol.; 84: 3291-3303.

Raamsman MJB, Locker JK, de Hooge A, de Vries AA, Griffi ths G, Vennema H, (2000) Characterization of the coronavirus mouse hepatitis virus strain A59 small membrane protein E. J Virol; 74: 2333-2342. https://goo.gl/GwygyF

Stertz S, Reichelt M, Spiegel M, Kuri T, Martinez Sobrido L, Garcia Sastre A. (2007). The intracellular sites of early replication and budding of SARS coronavirus Virology; 361: 304-15

Vennema H, Godeke GJ, Rossen JW, Voorhout WF, Horzinek MC, Opstelten DJ. (1996). Nucleo capsid-independent assembly of coronavirus-like particles by co-expression of viral envelope protein genesEMBO J; 15: 20202028. https://goo.gl/tFRLYY

Woo PC, Lau SK, Lam CS, Lai KK, Huang Y, Lee P, Luk GS, Dyrting KC, Chan KH, Yuen KY. (2009). Comparative analysis of complete genome sequences of three avian coronaviruses reveals a novel group 3c coronavirus J Virol. ; 83:908–917

Woo PC, Wang M, Lau SK, Xu H, Poon RW, Guo R, Wong BH, Gao K, Tsoi HW, Huang Y, Li KS, Lam CS, Chan KH, Zheng BJ, Yuen KY. (2007). Comparative analysis of twelve genomes of three novel group 2c and group 2d coronaviruses reveals unique group and subgroup features, J Virol, 81: 1574–1585.

Table 1 Amino Acid Composition the Structural Protein Extracted from SARS-Cov-2

Protein	envelope protein	Surface protein	membrane glycoprotein	nucleocapsid phosphoprotein
Number	75	1273	222	419
Molecular	8365.04	141178.47	25146.62	45625.70
Theoretical pI	8.57	6.24	9.51	10.07
negatively	3	110	13	36
positively charged	5	103	21	60
Formula	$C_{390}H_{62}...N...O_{...}$	$C_{6336}H_{9770}N_{1...}O_{...}S_{...}$	$C_{1165}H_{1823}N_{...}O_{...}S_{...}$	$C_{1971}H_{3137}N_{607}O_{62}S_{...}$
Total number of atoms	1213	19710	3600	6351
Extinction coefficient	6085	148960	52160	43890
half-life	30 hours	30 hours	30 hours	
Instability index	38.68	33.01	39.14	55.09
(GRAVY)	1.128	-0.079	0.446	-0.971
Aliphatic index	144.00	84.67	120.86	52.53

Table 2-Physical Parameters of the Structural Protein from SARS-Cov-2

Proteins	Envelope Protein	Surface Glycoprotein	Membrane Glycoprotein	Nucleocapsid Phosphoprotein
Ala (A)	4	79	19	37
Arg (R)	3	42	14	29
Asn (N)	5	88	11	22
Asp (D)	1	62	6	24
Cys (C)	3	40	4	0
Gln (Q)	0	62	4	35
Glu (E)	2	48	7	12
Gly (G)	1	82	14	43
His (H)	0	17	5	4
Ile (I)	3	76	20	14
Leu (L)	14	108	35	27
Lys (K)	2	61	7	31
Met (M)	1	14	4	7
Phe (F)	5	77	11	13
Pro (P)	2	58	5	28
Ser (S)	10	99	15	37
Thr (T)	8	97	13	32
Trp (W)	4	12	7	5
Tyr		54	9	11
Val (V)	13	97	12	8
Pyl (O)	0	0	0	0
Sec (U)	0	0	0	0

Table 3 Transmembrane of the Protein Extracted from SARS-Cov-2

	No	Transmembrane Region	N Terminal	C Terminal	Type	Length	Predicted Location
Surface glycoprotein	1	MFVFLVLLPLVSSQCVNLTTRT	1	22	SECONDARY	22	outside
	2	GLIAIVMVTIMLCCMTSCCSCLK	12	124	PRIMARY	23	outside
Envelop glycoprotein	1	TLIVNSVLLFLAFVVFLLVTLAI	11	33	PRIMARY	23	inside
	2	LRLCAYCCNIVNVSLVKPSFYVY	37	59	PRIMARY	23	inside
Nucleocapsid phosphoprotein		NO TRANSMEMBRANE PROTEIN					
Membrane glycoprotein	1	LLEQWNLVIGFLFLTWICLLQF	16	37	SECONDARY	22	inside
	2	YIIKLIFLWLLWPVTLACFVLAA	47	69	PRIMARY	23	inside
	3	GIAIAMACLVGLMWLSYFIASFR	79	101	PRIMARY	23	inside

Table 4-Predicted domains of Structural Proteins of SARS-CoV-2 by SMART

Proteins	Name	Start	End	E-value
Surface Glycoprotein	Low complexity	2	11	N/A
	Pfam:Spike_rec_bind	330	583	6.1e-75
	Pfam:Corona_S2	671	1279	1.3e-266
Envelop Glycoprotein	Pfam:NS3_envE	1	74	7.7e-8
Membrane Glycoprotein	Pfam:Corona_M	4	221	1.3e-95
Nucleocapsid Phosphoprotein	Pfam:Corona_nucleoca	14	337	1.2e-164

Table 5-Summary of Structural Proteins of SARS-CoV-2 Homology Models and Statistical Validations

Figure 1-The COVID-19 region of 5′ UTR and 3′ UTR and coding, the number of base pairs shown. [7]

					favoured regions	allowed regions	y allowed regions	lowed regions
Surface Glycoprotein	6vsb.1.B	0.77	0.87	99.26	86.4%	13.0%	0.6%	0 0.0%
Envelop Glycoprotein	5x29.1.A	0.78	0.46	91.38	84.4%	14.1%	1.5%	0.0%
Nucleocapsid Phosphoprotein	1ssk.1.A	0.25	Nil	92.37	83.3%	15.6%	1.1%	0.0%
Membrane Glycoprotein	5ctg.1.A	0.08	00	14.29	91.4%	8.6%	0.0%	0.0%

Figure -2 3D Model generated from Swiss Model of SARS COV-2 Structural proteins; Surface glycoprotein (A), Envelop glycoprotein (B), Membrane glycoprotein (C), Nucleocapsid phosphoprotein(D)

Figure-3 Ramachandran plots for structural proteins of SARS COV-2 Surface glycoprotein (A), Envelop glycoprotein (B), Membrane glycoprotein (C), Nucleocapsid phosphoprotein obtained by PROCHECK, most favored regions are display in red ; additional allowed regions are display in yellow; generously allowed regions are display in beige; and disallowed regions are display in white