Human iPSC-derived cerebral organoids model features of Leigh Syndrome and reveal abnormal corticogenesis

Alejandra I. Romero-Morales[1], Anuj Rastogi[1], Gabriella L. Robertson[1],Megan L. Rasmussen[1], Hoor Temuri[1], Ram Prosad Chakrabarty[3], Gregory Scott McElroy[3], Lawrence Hsu[2], Paula M. Almonacid[4], Bryan A. Millis[1,5], Navdeep S. Chandel[3,6], Jean-Philippe Cartailler[2] and Vivian Gama[1,2,7]

[1]Vanderbilt University, Cell and Developmental Biology, Nashville, TN.
[2]Vanderbilt University, Creative Data Solutions, Vanderbilt Center for Stem Cell Biology, Nashville, TN
[3]Northwestern University, Feinberg School of Medicine Department of Medicine Division of Pulmonary and Critical Care Medicine, Chicago, IL
[4]Universidad EAFIT, School of Economics and Finances, Colombia
[5]Vanderbilt University, Vanderbilt Biophotonics Center, Nashville, TN
[6]Northwestern University, Feinberg School of Medicine Department of Biochemistry and Molecular Genetics, Chicago, IL
[7]Vanderbilt University, Vanderbilt Brain Institute, Nashville, TN

Corresponding author

Correspondence should be addressed to:

Vivian Gama
PMB407935
465 21st Avenue South, 4150A
Vanderbilt University
Nashville, TN 37240-7935
Tel: (615) 875-9490
vivian.gama@vanderbilt.edu

Running title: Modeling the effect of Leigh syndrome-associated mutations using human cerebral organoids

50 **Keywords:** Leigh syndrome, stem cells, glycolysis, oxidative phosphorylation, mitochondria,
51 neural precursor cells, neural rosettes, brain organoids
52

Summary

Leigh syndrome (LS) is a rare, inherited neurometabolic disorder that presents with bilateral brain lesions, caused by defects in the mitochondrial respiratory chain and associated nuclear-encoded proteins. We generated iPSCs from three available LS fibroblast lines and identified, through whole exome and mitochondrial sequencing, unreported mutations in pyruvate dehydrogenase (GM0372, PDH; GM13411, MT-ATP6/PDH) and dihydrolipoyl dehydrogenase (GM01503, DLD). LS derived cell lines were viable and able to differentiate into key progenitor populations, but we identified several abnormalities in three-dimensional differentiation models of brain development. LS-derived cerebral organoids showed defects in neural epithelial bud generation, reduced size, and loss of cortical architecture at 100 days. The MT-ATP6/PDH line produced organoid neural progenitor cells with an abnormal mitochondrial morphology characterized by fragmentation and disorganization and demonstrated an increased generation of astrocytes. These studies aim to provide a comprehensive phenotypic characterization of available patient-derived cell lines that could be used as LS model systems.

Introduction

Leigh syndrome (LS), or sub-acute necrotizing encephalomyelopathy, is an inherited neurometabolic disorder that affects the central nervous system (CNS) (Baertling et al., 2014; Gerards et al., 2016; Leigh, 1951; Sorbi and Blass, 1982). LS is a rare, progressive, early-onset disease with a prevalence of 1 in 40,000 live births (Lake et al., 2016). The pathologic features of LS are focal, bilateral lesions in one or more areas of the CNS, including the brainstem, thalamus, basal ganglia, cerebellum, and spinal cord. The most common underlying cause is defective oxidative phosphorylation (OXPHOS), due to mutations in genes encoding complexes of the mitochondrial respiratory chain (Baertling et al., 2014; Lake et al., 2015, 2016). The LS phenotype caused by mitochondrial-encoded gene mutations can be rescued by mitochondrial replacement, highlighting the importance of this organelle in LS disease progression (Ma et al., 2015).

The availability of animal models and brain tissue from biopsies has provided critical insight into this disease. However, our understanding of the etiology and pathology of complex neurological diseases like LS would benefit from human-derived platforms such as an induced pluripotent stem cell-derived model (Quadrato et al., 2016). The ability to reprogram somatic cells into induced pluripotent stem cells (iPSCs), followed by differentiation into specific lineages has become a useful tool for complex disease modeling (Kelava and Lancaster, 2016; Di Lullo and Kriegstein, 2017). In the context of LS, iPSCs have been successfully generated from patients with mutations in Mitochondrially Encoded ATP Synthase Membrane Subunit 6 (MT-ATP6) (Galera-Monge et al., 2016; Grace et al., 2019; Lorenz et al., 2017; Ma et al., 2015), Mitochondrially Encoded NADH:Ubiquinone Oxidoreductase Core Subunit 3 (MT-ND3) subunit (Hattori et al., 2016) and the nuclear-encoded gene Surfeit locus protein 1 (SURF1) (Inak et al., 2019). These iPSC-model systems have been proposed for drug discovery (Inak et al., 2017; Lorenz et al., 2017) as well as testing platforms for potential metabolic rescue treatments (Ma et al., 2015).

Many studies have used LS patient fibroblasts commercially available at the Coriell

Institute. This repository provides scientists around the world with resources for cell and genetic

research. Here we report our findings on the genomic and phenotypic characterization of iPSCs

derived from three LS fibroblast lines available at Coriell. Whole exome and mitochondrial

sequencing revealed previously unidentified mutations in these patient-derived cell lines. While

these LS mutations had no effects on pluripotency, trilineage differentiation, or apoptotic

sensitivity, three-dimensional differentiation into neural rosettes and cerebral organoids resulted

in severe abnormalities. LS-iPSC-derived 100-day cerebral organoids showed decreased size

as well as defects in the generation of neural epithelial buds. Corticogenesis was impaired in all

LS mutant cell lines. MT-ATP6/PDH showed a decrease in NPC and cortical plate markers at

day 30, while DLD and PDH mutants showed a reduction of the upper layer markers SATB2,

BRN2, and CUX1 at day 100. These results point to aberrant corticogenesis as a driver of LS

pathogenesis and demonstrate the utility of iPSC-derived systems to recapitulate CNS

phenotypes and to test potential strategies to restore neurogenesis in LS.

Results

Genomic characterization of Leigh syndrome fibroblasts.

Due to the limited genomic information available for the three cell lines, we performed

whole-exome sequencing (WES) and mitochondrial sequencing of the fibroblasts prior to

reprogramming (Figure 1A-C). Targeted analysis of the genes associated with Leigh syndrome

(Lake et al., 2016) revealed a loss of function insertion/deletion (indel) frameshift in pyruvate

dehydrogenase complex (PDHc) E1 alpha 1 subunit or pyruvate dehydrogenase (PDHA1,

c.79delC, p.Arg27fs) in the cell lines GM03672 and GM13411. A single-nucleotide

polymorphism (SNP) in the PDHc E3 subunit or dihydrolipoyl dehydrogenase (DLD, c.100A>G,

p.Thr34Ala) was identified in GM01503 (Table S1). In addition to being part of the PDHc, DLD is

also a component of the α-ketoglutarate and branched-chain α-ketoacid dehydrogenase

complexes (Craigen, 1996). Despite the lack of genomic data, dysfunction of the PDH complex

had been previously suggested in GM03672 and GM01503 (Hinman et al., 1989; Huh et al.,

1990; Sorbi and Blass, 1982) as the main driver of the disease in these patients. To our

knowledge, mutations in the nuclear genome of GM13411 have not been reported to date.

Mitochondrial sequencing identified several SNPs in all of the cell lines (Figure 1D). A loss

of function SNP in the MT-ATP6 gene was identified in the GM13411 line. These mutations

were reported in the original clinical case (Pastores et al., 1994). The authors described the T to

G transition at the 8993 position that results in the substitution of a highly conserved leucine

residue for an arginine (L156R). MT-ATP6 is part of the F0 domain of ATP synthase that

functions as a proton channel. The L156R substitution prevents the induction of c-ring rotation,

resulting in decreased ATP synthesis (Uittenbogaard et al., 2018). Heteroplasmic analysis of

fibroblasts showed a 92% frequency of this mutation in the cell population, which is consistent

with previous reports (Galera-Monge et al., 2016; Iyer et al., 2012; Pastores et al., 1994).

Characterization of iPSCs derived from commercially available Leigh syndrome

fibroblasts.

Metabolic remodeling is a crucial step toward reprogramming somatic cells into iPSCs

(Mathieu and Ruohola-Baker, 2017; Panopoulos et al., 2012; Rasmussen and Gama, 2020;

Rastogi et al., 2019; Wu et al., 2016). Reprogramming of fibroblasts was performed as

previously described (Takahashi et al., 2007) (Figure S1A); we generated iPSCs from a healthy

age-matched control and the three LS cell lines. Pluripotency was evaluated using the

microarray-based analysis PluriTest (Müller et al., 2011). All three LS cell lines showed a high

pluripotency score and a low novelty score (Figure S1B-C), congruent with the transcriptional

profile of pluripotent stem cells. Moreover, all the reprogrammed cells expressed the pluripotent

markers *NANOG* and *OCT4* (Figure S1D) and had a normal karyotype (Figure S1E)

To assess the ability of the LS and control cell lines to differentiate into the three germ

layers, we performed trilineage differentiation. Commitment into ectodermal fate was evaluated

by immunofluorescence staining of the marker PAX6 (Figure 2A), as well as the mRNA

expression of the genes *GATA3* and *PAX6* (Figure 2B). The endoderm lineage was assessed

by protein expression of the marker SOX17 (Figure 2C) and by qPCR with the expression of the

genes *CDX2* and *SOX17* (Figure 2D). The PDH mutant cell line showed increased expression

of the endodermal gene SOX17 (Figure 2D) compared to control and the other LS cell lines.

Clinical data available from the patient with the PDH mutation showed elevated blood pyruvate

levels. High concentrations of pyruvate have been shown to potentiate the differentiation of

human embryonic stem cells (hESCs) into endodermal and mesodermal lineages while

suppressing the expression of ectodermal markers in a lineage-specific fashion (Song et al.,

2019). Finally, mesodermal lineage was confirmed by the staining of the protein markers

Brachyury and CXCR4 (Figure 2E) and the expression of the genes *TBXT* and *NCAM* (Figure

2F).

Two-dimensional neural differentiation is not affected by Leigh syndrome-associated

mutations.

To determine if the LS mutations impact the commitment and development of the neural

lineage, neural progenitor cells (NPCs) were generated by a dual SMAD inhibition protocol

(Chambers et al., 2009) (Figure 3A). NPCs showed the expected expression of the neural

markers PAX6, NESTIN, and SOX2 (Figure 3B), with no differences between the protein

expression of the LS cell lines and control (Figure S2A). The multipotent capacity of NPCs to

generate the three basic neural lineages (neurons, astrocytes, and oligodendrocytes) was

evaluated using standard protocols (Chambers et al., 2009; TCW et al., 2017; Vescovi and

Snyder, 1999). Immunostaining showed β3 TUBULIN-positive neurons, OLIG2-positive

oligodendrocytes, and S100β-positive astrocytes (Figure 3C). Thus, LS NPCs constitute a

multipotent population capable of differentiation into the main cellular populations of the nervous

system.

Considering that neural cell death is a hallmark of LS, we performed a cell viability assay to

investigate the sensitivity of the LS derived NPCs to different apoptotic stimuli (Figure S2B).

Treatment with DNA damaging agents etoposide and neocarzinostatin, as well as the

microtubule depolymerizing agent nocodazole, did not show increased sensitivity to cell death in

LS NPCs compared to control. To evaluate the susceptibility of the NPCs to mitochondrial

damage, we treated these cells with the mitochondrial oxidative phosphorylation uncoupler,

CCCP. No difference was observed in the viability of the LS NPCs compared to control-treated

with this mitochondrial toxicant.

Mitochondria in murine NPCs form an elongated network (Khacho et al., 2016) which

fragments as cells differentiate. Characterization of the mitochondrial network using structured

illumination microscopy (SIM) showed no major morphological changes in LS NPCs (Figure 3D

and Figure S3). Thus, LS mutations do not affect the capacity of the NPCs to further

differentiate into the three neural lineages, do not affect the viability of the NPCs when exposed

to different exogenous stressors in a monolayer culture, and do not perturb the morphology of

the mitochondrial network in these cells.

LS mutations cause differential accumulation of metabolites in neuralized embryoid

bodies.

Previous studies using cells from LS patients carrying homozygous *SURF1* (c.769G>A and

c.530T>G) (Inak et al., 2019) and MT-ATP6 (m.9185T>C) (Lorenz et al., 2017) mutations

showed an abnormal generation of neural lineages *in vitro* using two-dimensional culture

systems. SURF1 mutations caused impaired neurogenesis in cerebral organoids (Inak et al.,

2019). Therefore, we investigated the effects of the PDH, DLD, and MT-ATP6/PDH LS

mutations on neurogenesis using three-dimensional (3D) models of neural development as

described previously (Lancaster and Knoblich, 2014; Romero-Morales et al., 2019). Most of

these 3D neural systems start with the formation of neuralized embryoid bodies (EBs). EBs are

generated from a single cell suspension (Itskovitz-Eldor et al., 2000) plated in ultra-low

attachment microwells with neural induction media (Figure 4A). EB diameter was measured on

day 5 before plating on Matrigel-coated imaging plates. All three LS mutant lines showed an

increase in their diameter compared to control (Figure 4B, PDH vs Control & DLD vs Control:

$p<0.0001$, MT-ATP6/PDH vs Control: $p=0.0209$).

We then measured metabolite levels in EBs derived from LS cell lines using LC-MS-based

metabolomics (Table S2). Several pathways were found to be potentially altered in the LS EBs

compared to controls (Figure 4C and Table S3). These include central carbon metabolism (for

example glycolytic intermediates like Fructose 2,6-bisphosphate (f2,6p) and tricarboxylic acid

cycle (TCA) intermediates like citrate seem to be consistently enriched), glutathione

metabolism, serine metabolism, cysteine metabolism, and nucleotide metabolism.

Dihydroorotate, lactate, and aspartate are metabolites that were altered in the setting of electron

transport chain (ETC) inhibition, and there is evidence of a trend towards the enrichment of

dihydroorotate, lactate, and aspartate compared to control. There also appears to be an

enrichment of the NAD pool. These results suggest that LS mutations result in differential

changes in metabolite concentrations, which may cause downstream changes in gene

expression or cellular function.

Neural rosettes derived from Leigh syndrome iPSCs show morphological differences.

To investigate the effects of LS-associated mutations in the early stages of CNS

development, we generated neural rosettes using EBs grown in the presence of SMAD inhibitor

media (Figure 5A) (Elkabetz et al., 2008; Zhang et al., 2001). These structures have previously

been established to recapitulate the early neural tube formation stage of development (Elkabetz

et al., 2008; Wilson and Stice, 2006).

Neural rosettes were stained with the tight junction marker ZO-1 (Elkabetz et al., 2008;

Hříbková et al., 2018) and the centrosomal marker CDK5RAP2 (Figure 5B). Quantification of

the number of neural rosettes per field of view showed a reduction in the counts of these

structures in the DLD mutant compared to controls (Figure 5C, p<0.001). Lumen area

quantification showed an increase in the PDH and MT-ATP6/PDH mutants, while the DLD

mutant line showed a decrease in their area relative to controls (Figure 5D). The neural rosettes

obtained from all cell lines followed the expected formation sequence described previously

(Hříbková et al., 2018). The polymerization of α-tubulin and the generation of the ZO-1 ring at

the apical region of the rosettes are conserved in the LS mutants. Increased neural rosette

lumen size has previously been associated with TGFβ pathway activation (Medelnik et al.,

2018), Notch and SHH pathway activation, and WNT inhibition (Elkabetz et al., 2008). Large

rosette formation has been previously shown to be a consequence of apical domain opening

and expansion rather than a process dependent on cell proliferation (Medelnik et al., 2018).

Leigh syndrome-associated mutations disrupt corticogenesis in cerebral organoids.

To investigate the effects of these mutations during corticogenesis, we generated cerebral

organoids from LS iPSCs (Figure 6A). Differences between the cell lines became apparent as

early as the neuroepithelial bud expansion phase. After Matrigel embedding, the MT-ATP6/PDH

mutant cell line showed poor budding with large areas of non-neuroepithelial cells (Figure S4A).

Defective organoid formation in this cell line was significantly higher than control and the other

two LS cell lines (Figure S4B). A previous report showed that when iPSCs generated from

fibroblasts harboring the same T8993G mitochondrial mutation were differentiated into EBs,

there was rapid regression and death after 7 days in suspension, while the monolayer culture

did not show obvious deficits in cell growth (Grace et al., 2019). Given the fact the

neuroectoderm expansion phase happens during days 7-10, the degeneration of the MT-

ATP6/PDH organoids after embedding may recapitulate these previously reported observations.

Higher metabolic requirements have been shown to be associated with NPC proliferation

and migration in three-dimensional scaffolds and development (Fang et al., 2020; Homem et al.,

2015). As the PDH mutant line did not show this particular phenotype during the expansion of

the neural epithelium, the presence of the additional mitochondrial mutation in the MT-

ATP6/PDH line and the metabolic burden generated by the double mutation may be the reason

for the reduction of organoid formation efficiency.

Cerebral organoid growth was tracked for 100 days (Figure 6B). Growth differences in the

average diameter of the organoids become significant at day 80 between control and the PDH

mutant line. At days 90 and 100, organoids derived from all LS cell lines were significantly

smaller than the controls. Phenotypic information available for the patient from which the PDH

mutant cell line was derived indicates that this particular patient presented with microcephaly.

Hence, we were able to mimic an aspect of the patient phenotype *in vitro* using the cerebral

organoid model system.

To assess the effect of the LS mutations during the first stages of corticogenesis, day 30

organoids were sectioned and stained for the ventricular zone (VZ), subventricular zone (sVZ),

and cortical plate (CP) markers (Figure 6C). As expected, based on the defective

neuroepithelial expansion, MT-ATP6/PDH organoids showed a significant reduction in the

number of cells that stained positive for the NPC markers PAX6 and NESTIN, as well as the

intermediate progenitor marker TBR2 (Figure 6D). In addition to these findings, the overall

architecture in MT-ATP6/PDH organoids was compromised. There were few to no ventricle-like

structures present and the foci of PAX6+ cells were not organized in the expected radial pattern.

Migration of early-born neurons *in vivo* depends on pioneer Cajal-Retzius neurons that are

positive for the glycoprotein REELIN (Lancaster et al., 2017). Cells positive for this marker were

identified in superficial regions of control, PDH, and DLD organoids, while none were observed

in the MT-ATP6/PDH organoids (Figure 6E). Early born neurons at this time point are expected

to migrate into the CP that will later give rise to the deep cortical layers (Camp et al., 2015).

CTIP2+ and TBR1+ cells were observed in control, PDH, and DLD organoids but absent in MT-
ATP6/PDH organoids, demonstrating once again a profound impairment in the early stages of
corticogenesis when both MT-ATP6 and PDH mutations are present.

Mitochondrial morphology was evaluated in the VZ NPCs of the cerebral organoids. Cells
positive for NPC marker SOX2+ demonstrated elongated mitochondrial networks that extend
radially from the ventricle-like lumen (Figure 6F). This observation is consistent with what has
been previously reported in SOX2+ cells in the developing mouse cortex (Khacho et al., 2016).
DLD mutant organoid NPCs showed a fragmented mitochondrial network in comparison with
control and PDH. As mentioned earlier, the stereotypical arrangement of the VZ was
compromised in most MT-ATP6/PDH organoids. In the few areas where ventricle-like structures
were identified with a conserved SOX2+ VZ, the mitochondrial network seemed more
fragmented and aggregated. This morphology was also observed in the clusters of SOX2+ cells
that were scattered throughout the organoid.

In order to assess the proper cortical layer fate specification during normal cortical
development, we grew the cerebral organoids until day 100 and probed for upper cortical layer
markers (Figure 7A) (Florio and Huttner, 2014; Lui et al., 2011; Saito et al., 2011). Late-born
superficial layer markers CUX1, BRN2, and SATB2 (layers II, III, and IV, respectively) were
reduced in PDH and DLD organoids and were absent in MT-ATP6/PDH organoids (Figure 7B-
D). CTIP2+ cells were also reduced in PDH and DLD organoids (Figure 7B-C). Interestingly, at
day 100, few PAX6+ cells were observed in PDH and DLD organoids, whereas in MT-
ATP6/PDH organoids they were present. The reduction in the number and variety of cell types
found in PDH and DLD may explain the decrease in the diameter of the organoids derived from
these cell lines (Figure 6B).

Compared to controls, LS cerebral organoids showed increased staining for the astrocyte
marker S100β at day 100 (Figure 7C). The decrease in the diversity of neuronal cell types and
the increase in the presence of S100β+ cells may suggest a switch to astrocyte fate during

cortical development in LS. Clinical data from LS patients include marked gliosis as part of the

characteristic findings (Lake et al., 2015). While this marked gliosis is potentially associated with

a reactive process secondary to neuronal damage, an intriguing alternate possibility is that

progenitor cells may have an increased propensity to differentiate down the astrocyte lineage

due to LS-causative mutations. Previous studies have shown that reactive astrocytes acquire

molecular hallmarks of radial glial cells. It was also shown through genetic fate mapping that

mature astroglial cells can dedifferentiate and resume proliferation (Robel et al., 2009, 2011).

Thus, the phenotypes uncovered here could either reflect that chronic metabolic stress induced

by Leigh syndrome mutations activates a brain injury response, or that the inhibition of

mitochondrial metabolism in neural progenitor cells could cause defects in lineage selection.

We then explored the changes in metabolites in Day 40 organoids from control and LS

samples (Figure S5). From the top 10 metabolites identified, the organic osmolyte betaine was

increased in the control compared to the DLD and PDH samples, and the quaternary amine

carnitine was increased in the control compared with all LS samples. L-Carnitine is essential for

the transport of long-chain fatty acids into mitochondria for degradation by β-oxidation (Jernberg

et al., 2017). Carnitine concentration in the brain increases during gestation (Nakano et al.,

1989) and has been reported as a therapeutic option for children affected by metabolic

diseases. Moreover, this compound has been shown to decrease oxidative stress, improve

brain energy status, and prevent neuronal cell death in animal models of brain injury and

ischemia (Ferreira and McKenna, 2017).

Pseudouridine, an isomer of uridine and the most abundant RNA modification, was

decreased in DLD organoids. Pseudouridine and pseudouridine synthases (Pus) have been

associated with regulating neuronal functions (Angelova et al., 2018). Specifically, Pus3 has

been implicated in neural development (Diez-Roux et al., 2011). A truncated form of Pus3

accompanied with reduced levels of pseudouridine in tRNA has been identified in patients with

intellectual disabilities (Shaheen et al., 2016).

High lactate was identified in both PDH and MT-ATP6/PDH organoids, correlating with the

available clinical information that both patients presented elevated lactic acid in the blood (Table

1). Lactic acid accumulation in blood and cerebrospinal fluid is the key clinical indicator for LS

diagnosis and is particularly expected in the setting of PDH dysfunction (Miyabayashi et al.,

1985). Also, in these two lines, the metabolite proline was elevated. High levels of proline have

been associated with negative effects in brain function by interference in glutamatergic

neurotransmission (Gogos et al., 1999; Vorstman et al., 2009). In addition, hydroxyphenyllactic

acid was elevated in PDH mutant organoids. High levels of this metabolite have been reported

in association with high lactate and pyruvate in pediatric lactic acidosis in patients with PDHc

deficiency (Stern, 1994). Not surprisingly, in the MT-ATP6/PDH mutant organoids, pyruvate was

also increased, correlating with the lactic acidosis expected in the organoids based on the

patient phenotypes and the presence of the PDH mutation that hinders flux from pyruvate into

the TCA cycle through acetyl-CoA.

Besides the previously mentioned metabolites, MT-ATP6/PDH mutant organoids presented

increased levels of choline, cytidine, and leucine. Choline is a crucial metabolite for normal CNS

development. Neural tube defects have been associated with a lack of choline during early

pregnancy (Zeisel, 2006). It has also been shown to increase cell proliferation and decrease

apoptosis in fetal rat hippocampal progenitor cells (Albright et al., 1999b, 1999a; Zeisel and

Niculescu, 2006). Choline is also crucial for the production of the neurotransmitter acetylcholine,

the sphingolipid sphingomyelin, and myelin (Oshida et al., 2003). Concomitantly, cytidine is

used with choline for the generation of cytidine-5-diphosphocholine, a crucial intermediate in the

biosynthesis of the cell membrane phospholipids phosphatidylcholine and

phosphatidylethanolamine (Cansev, 2006; Rema et al., 2008). Increased abundance of the

branched-chain amino acid leucine has been associated with the metabolic illness maple syrup

urine disease and can be extremely neurotoxic (Bridi et al., 2005; García-Cazorla et al., 2014).

This amino acid is considered ketogenic as its end products can enter the TCA cycle for energy

generation or act as precursors for lipogenesis and ketone body production (Manoli and

Venditti, 2016).

Considering all the aforementioned findings, the cerebral organoid system emerges as a

promising model for the study of the effects of LS associated mutations in brain development

and the effects of metabolic impairment during corticogenesis.

Discussion

Inborn errors of metabolism are rare genetic disorders resulting from defects in metabolic

pathways (Agana et al., 2018; Das et al., 2010). Mitochondrial diseases are the most common

group of inherited metabolic disorders and are among the most common forms of inherited

neurological disorders (Gorman et al., 2016). These illnesses are challenging not only at the

time of diagnosis but also during their medical evolution, as they involve multiple organ systems

and there are limited therapeutic options (Grier et al., 2018; Parikh et al., 2017; Schaefer et al.,

2019).

Leigh syndrome is one of these rare inherited neurometabolic diseases, with more than 75

causal genes identified in both nuclear and mitochondrial DNA. It has an early onset, affecting

most patients within their first year of life, although cases during teenage years and adulthood

have been reported (Finsterer, 2008; Lake et al., 2016). As it is a highly heterogeneous disease,

the establishment of animal and *in vitro* models has been challenging and limited to only select

mutations. The few animal models available have been utilized for the development of

therapeutic approaches with mixed results. Gene editing using adeno-associated virus in

Ndufs4-/- mice has shown partial rescue of the phenotype (Di Meo et al., 2017).

Supplementation of nicotinamide riboside to Sco2-/- mice showed improvement of the

respiratory chain defect and increased exercise tolerance due to improved mitochondrial

biogenesis (Cerutti et al., 2014). Hypoxia and low oxygen availability in the brain have also been

shown to increase the life span and improve neurological findings in Ndufs4-/- mice (Ferrari et

al., 2017; Jain et al., 2016, 2019).

Molecular testing and prenatal diagnosis of respiratory chain disorders using skin

fibroblasts and muscle biopsies for diagnostic and research purposes are becoming mainstream

procedures (Baertling et al., 2014; Calvo et al., 2006, 2012; Schubert and Vilarinho, 2020).

Nevertheless, heteroplasmy in mitochondrial DNA among tissues can cause less pronounced or

absent phenotypes in cultured cells (Baertling et al., 2014). Here we report the characterization

and the subsequent generation of brain organoids from three commercially available Leigh

syndrome fibroblast cell lines and an age-matched control.

Three-dimensional differentiation generates higher numbers of NPCs and more mature

neurons than two-dimensional differentiation (Chandrasekaran et al., 2017; Di Lullo and

Kriegstein, 2017; Muratore et al., 2014; Paşca et al., 2015) in part due to an improved spatial

cellular environment that influences cell fate specification. Tissue architecture, mechanical cues,

cell-to-cell communication (Pampaloni et al., 2007), nutrient accessibility, oxygen tension, as

well as morphogen gradients characteristic of 3D systems (Tibbitt and Anseth, 2012) aid to

recapitulate the development of the central nervous system (CNS) up to approximately 16

weeks post-fertilization (Camp et al., 2015; Paşca et al., 2015). We observed that all the LS

cerebral organoids failed to thrive at different time points. Although organoid development

initially appeared normal in cell lines with nuclear-encoded LS mutations, at later time points,

the overall diameter decreased, presumably due to failure to generate upper-layer neurons. This

reduction of late-born neurons may be due to the overall reduction in the PAX6+ NPCs in both

PDH and DLD compared to controls. Interestingly, PDH cerebral organoids displayed arrested

growth very early on, correlating with the microcephaly observed in the source patient.

The increase in the glial-specific marker S100β in DLD and MT-ATP6/PDH organoids could

also correlate with the observation that gliosis is a common clinical finding in LS patients

(Baertling et al., 2014, 2016; Schubert and Vilarinho, 2020). The formation of lesions in LS has

been described as the result of OXPHOS dysfunction and subsequent ATP depletion. Neuronal

dysfunction is suspected to trigger chronic gliosis (Baertling et al., 2016). In patients, the gliosis

phenotype can be accompanied by vascular hypertrophy and production of excess ROS, which

increases neuronal damage (Lake et al., 2015). However, due to the lack of vascularization in

the organoid model, replicating the vascular abnormalities associated with LS is not feasible in

this system.

The profound dysregulation of corticogenesis in the double mutant MT-ATP6/PDH may

suggest that some pregnancies harboring LS-causing mutations may not be viable, which could

lead to an underestimation of the prevalence of the disease in the population (Feeney et al.,

2019). Prenatal genetic evaluation is now being performed for cases where there is a known

risk for mitochondrial mutations (Craven et al., 2017; White et al., 1999). This testing can be

performed as early as 10-12 weeks post conception and is usually requested if there is a history

of a previously affected child or first-degree relative (Nesbitt et al., 2014). Interpretation of these

tests is challenging in the case of mitochondrial mutations due to heteroplasmy; the biopsied

tissues may exhibit a different mutational burden compared with other fetal tissues (Ferlin et al.,

1997; Harding et al., 1992; Nesbitt et al., 2014; Steffann et al., 2007).

In a previous study (Hattori et al., 2016), the metabolic signature analysis of iPSCs derived

from a mitochondrial encoded LS mutation (m.10191T>C) showed differences in the abundance

of pyruvate and lactate, among others. Interestingly, the metabolic difference between control

and LS cells was reported to revert to normal after EB differentiation (Hattori et al., 2016). In our

study, metabolomic analysis from neuralized EBs and organoids show that the observed

changes in the metabolites are in line with the clinical observations of LS patients. Changes in

blood and cerebral spinal fluid concentration of lactate and pyruvate are common diagnostic

tools for LS (Hattori et al., 2016) and other mitochondrial diseases (Barshop, 2004; Buzkova et

al., 2018; Esterhuizen et al., 2017; Rahman and Rahman, 2018). We also identified trends

toward changes in other metabolites, like dihydroorotate and aspartate, that highlight the overall

impact of LS mutations at different steps of the metabolic pathway (ETC complex III, TCA cycle,

and *de novo* nucleotide synthesis, and NADH/NAD+ ratio).

The metabolic dysregulation of the affected tissues in LS may have a direct effect on

mitochondrial morphology and function. Mitochondrial fragmentation is a hallmark of glycolytic

cell types such as stem cells and cancer cells (Chen and Chan, 2017; Rastogi et al., 2019).

Moreover, neurogenesis defects have been observed in the context of mitochondrial

morphology dysregulation and are considered to be upstream regulators of self-renewal and cell

fate decisions in stem cells (Khacho et al., 2016). In addition, energetic requirements have been

shown to directly impact the capacity of progenitor cells to migrate and thrive in 3D

environments (Zanotelli et al., 2018, 2019). Hence, mitochondrial morphology disruption

observed in the double mutant MT-ATP6/PDH organoids is expected when analyzed in

conjunction with the metabolic and developmental profile of these mutant organoids. To our

knowledge, this is the first time that mitochondrial morphology in the cortex has been analyzed

in a human model system of LS brain development and highlights the potential importance of

mitochondria network plasticity for the proper specification of cell fate and survival.

Here, we presented the characterization of three commercially available LS cell lines from

fibroblasts to cerebral organoids. We were able to identify new genetic alterations in these

samples by using whole-exome sequencing and mitochondrial DNA sequencing. We described

the effects of these mutations in a 3D cerebral organoid system. We aim to provide a

comprehensive phenotypic characterization of available patient samples to encourage their

utilization as model systems for uncovering the mechanisms underlying neuronal cell death in

the context of LS and for drug discovery.

Figure Legends

Figure 1. Whole exome sequencing identifies novel mutations in Leigh syndrome fibroblasts. A. Schematic of the WES and mitochondrial sequencing workflow. B. Representation of whole-genome sequencing data, highlighting the top 20 genes containing high impact indels. C. Representation of whole-genome sequencing data, highlighting the top 20 genes containing high impact SNPs (increased likelihood of disrupting protein function). D. Mitochondrial sequencing identifies novel mutations in LS fibroblasts. Representation of mitochondrial sequencing data, highlighting mitochondrial genes containing mutations (transitions, deletions, or transversions). Red dots: represent DLD line. Green dots: represent the PDH line. Orange dots: represent the MT-ATP6/PDH line. PDH: Pyruvate dehydrogenase. DLD: Dihydrolipoyl dehydrogenase. MT-ATP6/PDH: Mitochondrially Encoded ATP Synthase Membrane Subunit 6/ Pyruvate dehydrogenase.

Figure 2. Induced pluripotent stem cells derived from Leigh syndrome patient fibroblasts are capable of differentiation into specific lineages. A-B. Representative images of the ectoderm marker Pax6 (A) and qPCR for the ectodermal genes *GATA3* and *PAX6* (B). C-D. Representative images of the endoderm marker SOX17 (C) and qPCR for the ectodermal genes *CDX2* and *SOX17* (D). E-F. Representative images of the mesoderm markers Brachyury and CXCR4 (E) and qPCR for the mesodermal genes *TBXT* and *NCAM* (F). Scale bar: 100µm.

Figure 3. Leigh syndrome derived NPCs are multipotent and do not show aberrant mitochondrial morphology compared to control. A. C. Immunofluorescence for multipotency markers. Neural progenitor cells stained by Pax6 and Nestin, neurons marked with βIII-tubulin, oligodendrocytes stained with Olig2, and astrocytes are marked with S100β. Scale bar: 100µm D. Representative super-resolution images of mitochondrial morphology (Mitotracker) in LS and control NPCs. Scale bar: 5µm.

Figure 4. Generation of neuralized EBs from Leigh syndrome derived iPSCs lead to changes in EB diameter and the abundance of metabolites involved in central carbon metabolism. A. Schematic of EB generation protocol. B. Embryoid body diameter is increased in the three LS cell lines. C. Metabolite abundance pathway analysis. Top metabolic pathways affected in day 10 LS neuralized EBs. Statistical p values from enrichment analysis are adjusted for multiple testing. Total: number of total compounds in the pathway. Hits: matched number from the uploaded data. Raw p: original p-value calculated from the enrichment analysis. Holm p: p-value adjusted by Holm-Bonferroni method. FDR p: adjusted p-value using False Discovery Rate. Impact: pathway impact value calculated from pathway topology analysis. *$p < 0.05$; **$p < 0.001$, ***$p < 0.0001$.

Figure 5. Three-dimensional differentiation reveals abnormalities during induction of neural rosettes in LS cell lines. A. Schematic of neural rosette generation protocol. B-D. Representative confocal images of neural rosettes (C) show decreased numbers of neural rosettes per field in the DLD mutant line (D). Quantification of the lumen area (μm^2) indicates increased lumen area in the PDH and MT-ATP6/PDH mutant cell lines and a decreased lumen area in the DLD mutant line. Scale bar: 50μm. *$p < 0.05$; **$p < 0.001$, ***$p < 0.0001$.

Figure 6. Leigh syndrome derived brain organoids show a reduction in size and MT-ATP6/PDH mutant brain organoids show defects in SVZ/VZ and CP formation and aberrant mitochondrial morphology in SOX2+ cells. A. Schematic of brain organoid generation protocol. B. Brain organoid growth curves. LS derived brain organoids show reduction in size when grown for 100 days. C. Schematic representation of the expected organization of the brain organoids on day 30. D-E. Representative immunostaining confocal images of day 30 brain organoids. MT-ATP6/PDH mutant presents severe disorganization of the SVZ/VZ markers PAX6 and TBR2, as well as the neural progenitor marker NESTIN (D). MT-ATP6/PDH mutant does not express the

CP and deep layer markers CTIP2 and TBR1 (E). Scale bar: 100μm. F. Representative confocal

images of day 30 brain organoids showing mitochondrial morphology (TOM20). The red line

divides the Sox2+ neural progenitor cells surrounding the lumen (L) from newly committed

neurons. MT-ATP6/PDH mutant organoids show disorganization and fragmentation of the

mitochondrial network compared to control. Scale bar: 10μm. SVZ: subventricular zone, VZ:

ventricular zone, CP: cortical plate, MZ: marginal zone. * $p < 0.05$ for Control and PDH mutant;

□ $p < 0.05$ for Control and DLD mutant, φ $p <$ for Control and MT-ATP6/PDH mutant.

Figure 7. Leigh syndrome derived brain organoids show defects in cortical layer formation at

day 100. A. Schematic representation of the expected organization of the brain organoids at day

100. B-D. Representative immunostaining confocal images of day 100 brain organoids. LS

derived brain organoids present reduced expression of the upper layer markers SATB2 (B),

BRN2 (C) and CUX1 (D) and deep layer marker CTIP2 (C & D). Increased expression of the

astrocyte marker S100β was also noticed in the MT-ATP6/PDH mutant line. Scale bar: 100μm.

SVZ: subventricular zone, VZ: ventricular zone, DL: deep layers UL: upper layers, MZ: marginal

zone.

525 **STAR Methods**

526 **Key Resource Table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
ANTIBODIES		
Primary Antibodies (Immunocytochemistry)		
Rabbit anti-PAX6	Cell Signaling Technology	Cat # 60433, AB_2797599
Rabbit anti-SOX17	Cell Signaling Technology	Cat # 81778S, AB_2650582
Rabbit anti-BRACHYURY	Cell Signaling Technology	Cat # 81694S, AB_2799983
Mouse anti-CD184 (CXCR4)	STEMCELL Technologies	Cat # 60089
Mouse anti-β3 tubulin	Cell Signaling Technology	Cat # 4466, AB_10270973
Rabbit anti-S100	Abcam	Cat # ab868, AB_306716
Mouse anti-Olig2	Millipore Sigma	Cat # MABN50, AB_10807410
Rat anti-α tubulin	Thermo Fisher Scientific	Cat # MA180017, AB_2210201
Mouse anti-ZO-1	Thermo Fisher Scientific	Cat # 339100, AB_2533147
Rabbit anti-CDK5RAP2	Bethyl Laboratories	Cat # IHC00063, AB_2076863
Mouse anti-Nestin	STEMCELL Technologies	Cat # 60091, AB_2650581
Chicken anti-TBR2	Millipore Sigma	Cat # AB15894, AB_10615604
Mouse anti-Reelin	Millipore Sigma	Cat # MAB5366, AB_2285132
Rat anti-CTIP2	Abcam	Cat # ab18465, AB_2064130
Rabbit anti-TBR1	Abcam	Cat # ab31940, AB_2200219
Rabbit anti-TOM20	Cell Signaling Technology	Cat # 42406, AB_2687663
Mouse anti-SATB2	Abcam	Cat # ab51502, AB_882455
Mouse anti-BRN2 (POU3F2)	Millipore Sigma	Cat # MABD51, AB_11204531
Mouse anti-CASP (CUX1)	Abcam	Cat # ab54583, AB_941209
Secondary Antibodies (Immunocytochemistry)		
Goat anti Chicken Alexa Fluor 647	Thermo Fisher Scientific	Cat # A-21449, AB_2535866
Goat anti Rat Alexa Fluor 647	Thermo Fisher Scientific	Cat # A-21247, AB_2535864
Donkey anti Rabbit Alexa Fluor 647	Thermo Fisher Scientific	Cat # A-31573, AB_2536183
Donkey anti Mouse Alexa Fluor 647	Thermo Fisher Scientific	Cat # A-31571, AB_162542
Donkey anti Rabbit Alexa Fluor 546	Thermo Fisher Scientific	Cat # A-10040, AB_2534016

Donkey anti Mouse Alexa Fluor 546	Thermo Fisher Scientific	Cat # A-10036, AB_2534012
Donkey anti Rabbit Alexa Fluor 488	Thermo Fisher Scientific	Cat # A-21206, AB_2535792
Donkey anti Mouse Alexa Fluor 488	Thermo Fisher Scientific	Cat # A-21202, AB_141607
Primary Antibodies (Western Blotting)		
Rabbit anti-PAX6	Cell Signaling Technology	Cat # 60433, AB_2797599
Mouse anti-Nestin	STEMCELL Technologies	Cat # 60091, AB_2650581
Rabbit anti-Sox2	Cell Signaling Technology	Cat # 3579, AB_2195767
Mouse anti- α tubulin	Sigma-Aldrich	Cat # T9026, AB_477593
Secondary Antibodies -HRP conjugated (Western Blotting)		
Peroxidase AffiniPure Donkey Anti-Rabbit IgG (H+L)	Jackson ImmunoResearch Inc	Cat # 711-035-152, AB_10015282
Peroxidase AffiniPure Donkey Anti-Mouse IgG (H+L)	Jackson ImmunoResearch Inc	Cat # 715-035-151, AB_2340771
CHEMICALS, PEPTIDE, AND RECOMBINANT PROTEINS		
Y-27632 Rho/Rock pathway inhibitor	STEMCELL Technologies	Cat # 72307
Dorsomorphin	Millipore Sigma	Cat # P5499
SB431542	REPROCELL	Cat # 04-0010-10
Etoposide	Millipore Sigma	Cat # E1383
Carbonyl cyanide 3-chlorophenylhydrazone (CCCP)	Sigma Aldrich	Cat # C2759
Nocodazole	Sigma Aldrich	Cat # M1404
Neocarzinostatin	Sigma Aldrich	Cat # 9162
CRITICAL COMMERCIAL ASSAYS AND KITS		
PluriTest Assay	Thermo Fisher Scientific	Cat# A38154
KaryoStat Assay	Thermo Fisher Scientific	Cat# A38153
Mitochondrial DNA sequencing	Girihlet	
Whole Exome sequencing	Genewiz LLC	
CytoTune iPS 2.0 Sendai Reprogramming Kit	Thermo Fisher Scientific	Cat # A16517
STEMdiff Trilineage Differentiation Kit	STEMCELL Technologies	Cat# 05230
NeuroCult™ media	STEMCELL	Cat # 05752

	Technologies	
Astrocyte medium	ScienCell	Cat # 1801
CellTiter Blue Viability Assay	Promega	Cat # G8081

DEPOSITED DATA

Raw and analyzed sequencing data	Done by Creative Solutions (J.P.C, Vanderbilt University	https://www.ncbi.nlm.nih.gov/sra/PRJNA626388 https://vandydata.github.io/Romero-Morales-Gama-Leigh-Syndrome-WES/

EXPERIMENTAL MODELS: CELL LINES

AG16409 control fibroblasts	Coriell Institute	https://www.coriell.org/0/Sections/Search/Sample_Detail.aspx?Ref=AG16409&Product=CC
GM13411 (MT-ATP6/PDH) fibroblasts	Coriell Institute	https://www.coriell.org/0/Sections/Search/Sample_Detail.aspx?Ref=GM13411&Product=CC
GM03672 (PDH Mutant) fibroblasts	Coriell Institute	https://www.coriell.org/0/Sections/Search/Sample_Detail.aspx?Ref=GM03672&Product=CC
GM01503 (DLD Mutant) fibroblasts	Coriell Institute	https://www.coriell.org/0/Sections/Search/Sample_Detail.aspx?Ref=GM01503&Product=CC

SEQUENCE-BASED REAGENTS

Primers for Trilineage assay		
OCT4	Integrated DNA Technologies	Forward GGGCTCTCCCATGCATTCAAAC Reverse CACCTTCCCTCCAACCAGTTGC

NANOG	Integrated DNA Technologies	Forward
		    TGGGATTTACAGGCGTGAGCCAC
		Reverse
		    AAGCAAAGCCTCCCAATCCCAAAC

GAPDH	Integrated DNA Technologies	Harvard PrimerBank, ID: 378404907c2 Forward ACAACTTTGGTATCGTGGAAGG Reverse GCCATCACGCCACAGTTTC
GATA3	Integrated DNA Technologies	Forward TGGAGGAGGAATGCCAATGGG Reverse GCCGGGTTAAACGAGCTGTTCTTG
PAX6	Integrated DNA Technologies	Harvard Primer Bank, ID: 189083679c1 Forward TGGGCAGGTATTACGAGACTG Reverse ACTCCCGCTTATACTGGGCTA

CDX2	Integrated DNA Technologies	Forward
		    CTGGAGCTGGAGAAGGAGTTTCAC
		Reverse
		    GACACTTCTCAGAGGACCTGGCTG

SOX17	Integrated DNA Technologies	Harvard Primer Bank, ID:145275218c1 Forward GTGGACCGCACGGAATTTG Reverse GGAGATTCACACCGGAGTCA
TBXT	Integrated DNA Technologies	Forward ACAATGCCAGCCCACCTACCAG

		Reverse CGTACTGGCTGTCCACGATGTCTG
NCAM	Integrated DNA Technologies	Harvard Primer Bank, ID:316659209c1 Forward GGGGTTGCTTGTCAGTAGC Reverse TTCAGGTTCACCAATCGCTGT
SOFTWARE AND ALGORITHMS		
Image Studio™ Lite	LI-COR	https://www.licor.com/bio/image-studio-lite/download
Fiji	Schindelin et al., 2012	https://imagej.net/Fiji
GraphPad Prism v8.1.2	GraphPad	https://www.graphpad.com/scientific-software/prism/
NIS-Elements	Nikon Instruments	https://www.microscope.healthcare.nikon.com/products/software/nis-elements
MetaboAnalyst 4.0	(Chong and Xia, 2018; Chong et al., 2018, 2019; Xia and Wishart, 2010, 2011a, 2011b; Xia et al., 2009)	https://www.metaboanalyst.ca/home.xhtml
SnpSift	(Cingolani et al., 2012)	http://snpeff.sourceforge.net/SnpSift.html
R 3.5.3	R Foundation	https://www.r-project.org/
BioCircos	(Cui et al., 2016)	https://cran.r-project.org/web/packages/BioCircos/index.html
OTHER		
Mitotracker Red CMXRos	Fisher Scientific	Cat # M7512
Matrigel™	Corning	Cat # 354277
Gentle dissociation solution	STEMCELL Technologies	Cat # 07174
Aggrewell™ 800 24-well plate	STEMCELL Technologies	Cat # 34815
AggreWell™ Rinsing Solution	STEMCELL Technologies	Cat # 07010
STEMdiff™ Cerebral Organoid Kit	STEMCELL Technologies	Cat # 08570
STEMdiff™ Cerebral Organoid Maturation Kit	STEMCELL Technologies	Cat # 08571

Lead Contact and Materials Availability

529 Further information and requests for resources and reagents should be directed to the Lead

530 Contact, Vivian Gama (vivian.gama@vanderbilt.edu).

Experimental Model and Subject Details

The Coriell cell line IDs were as follows: GM01503, GM03672, GM1341. Information about the

Leigh syndrome cell lines used in this study can be found in Supplementary Table1. Control skin

fibroblast cell line AG16409 was also obtained from Coriell Institute, Philadelphia, PA. The

donor was a 12-year-old apparently healthy Caucasian male.

Fibroblasts were maintained in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12

DMEM/F-12 (Gibco cat # 11330032) supplemented with 10% Fetal bovine serum (Sigma cat#

F2442) in 100mm cell culture plates (Eppendorf, cat # 0030702115) in a 37°C 5% CO_2

incubator.

Method Details

Whole Exome sequencing

Fibroblast cell pellets from each cell line (>1 million cells) were shipped on dry ice for whole-

genome exome sequencing to Genewiz, Plainfield, NJ. The Illumina HiSeq-X was used to

perform 150nt paired-end sequencing.

Mitochondrial sequencing

Fibroblast cell pellets from each patient (>1 million cells) were shipped on dry ice for

mitochondrial sequencing to Girihlet Inc. Oakland, CA. The sequencing configuration used was

80bp single-end sequencing, 20 million reads per sample.

hiPSC Generation and Characterization

Human fibroblasts were purchased from healthy control and patients (Coriell Institute,

Philadelphia, PA, USA). Induced pluripotent stem cells were derived from human fibroblasts

using Sendai virus-based reprogramming kit (CytoTune-iPS Sendai Reprogramming Kit; cat #.

A13780-01; Thermo Fisher), according to manufacturer's instructions. After 3-4 weeks, 2-3

colonies per sample were transferred to fresh 6-well plates and were expanded and gardened

for 3 passages before freezing. All iPSC cell lines were maintained in E8 medium in plates

coated with Matrigel (Corning, cat # 354277) at 37°C with 5% CO_2. Culture medium was

changed daily. Cells were checked daily for differentiation and were passaged every 3-4 days

using Gentle dissociation solution (STEMCELL Technologies, cat # 07174). All experiments

were performed under the supervision of the Vanderbilt Institutional Human Pluripotent Cell

Research Oversight (VIHPCRO) Committee.

Analysis of Pluripotency

Pluripotency of each iPSC clone was determined using a microarray-based tool known as

PluriTest (Thermo Fisher; cat# A38154) as an alternative to the teratoma assay. Samples were

outsourced to Thermo Fisher for PluriTest and further analysis. Low passage iPSC cell pellets

(>1 million cells) were frozen and shipped on dry ice. Additionally, the expression of

pluripotency genes POU5F1 and NANOG was assessed by qPCR.

Analysis of Chromosomal abnormalities

The presence of any chromosomal abnormalities in the newly generated iPSCs was determined

using a microarray-based tool known as KaryoStat (Thermo Fisher; cat# A38153) as an

alternative to chromosomal G-banding. Samples were outsourced to Thermo Fisher for

KaryoStat and further analysis. Low passage iPSC cell pellets (>1 million cells) were frozen and

shipped on dry ice to Thermo Fisher.

Trilineage differentiation

The STEMdiff Trilineage differentiation kit (STEMCELL Technologies, cat# 05230) was used to

functionally validate the ability of newly established iPSCs to differentiate into three germ layers,

as per the manufacturer's instructions. Single-cell suspensions of $2x10^6$ cells/well, $5x10^5$

cells/well, $2x10^6$ cells/well were seeded for ectoderm, mesoderm, and endoderm, respectively,

in their corresponding medium at day 0 in 6 well plates. The cultures were maintained for 7

days, 5 days, and 5 days for ectoderm, mesoderm, and endoderm, respectively. The

differentiation was assessed by immunocytochemistry and qPCR.

NPC differentiation and multipotency characterization

For monolayer differentiation of the iPSCs into NPCs, cells were dissociated into single cells

using Gentle Cell Dissociation Reagent (STEMCELL Technologies, cat # 07174) for 8 minutes

at 37°C. Live cell counts were performed using Trypan blue (0.4%) staining (Invitrogen, cat #

T10282) using a Countess™ Automated Cell Counter. Cells were then seeded in a Matrigel-

coated 6-well plate (Eppendorf, cat # 0030720113) to a 2.5×10^6 cells/well with dual SMAD

inhibitor media supplemented with Dorsomorphin (1μM) and SB431542 (10μM) (Chambers et

al., 2009) supplemented with ROCK inhibitor. Daily media changes were performed and

passaging of the cells was done every 7-9 days. Cells for NPC marker analysis were collected

at the end of the first 9 days of differentiation.

Mitochondrial imaging was performed by adding 100nM MitoTracker Red CMXRos (Thermo

Fisher Scientific #M7512) to the media for 20 minutes before PFA fixation. Structured

Illumination Microscopy (SIM) was accomplished in 3D-SIM mode on a Nikon Instruments N-

SIM, equipped with an Apo TIRF 100x SR 1.49NA objective, DU-897 EMCCD camera (Andor),

405nm and 561nm lasers. Images presented herein are maximum intensity projections after

image stacks were first acquired (5 phase shifts and 3 rotations of diffraction grating,

120nm/axial step via piezo) and subsequent stack reconstruction in NIS-Elements software

(Nikon Instruments, Inc.). Other than linear intensity scaling, no further image processing was

performed post-reconstruction.

For multipotency analysis, culture media was changed to NeuroCult™ media and maintained for

4 weeks. Samples were then fixed and stained for neuron and oligodendrocyte markers.

Astrocyte differentiation was performed by seeding on a Matrigel-coated plate 1.5×10^6 cells/cm^2

(TCW et al., 2017). The following day, the media was changed to Astrocyte medium (ScienCell,

cat # 1801) and maintained for 20 days. Full media changes were done every 2 days. Samples

were then fixed and stained for an astrocyte marker. Images were acquired with a Nikon

Instruments Ti2 inverted fluorescence widefield microscope equipped with a Plan Apo Lambda

20X 0.75 NA objective, DS-Qi2 camera (Nikon Instruments), and X-Cite 120LED light source

(Excelitas).

Neural rosette differentiation

To generate neural rosettes, we dissociated the cells into a single-cell suspension and seeded

3.0×10^6 cells/well of an Aggrewell^{TM} 800 in dual SMAD inhibitor media. EBs were incubated at

37°C with 5% CO2, with minimal disruption during the first 48 hours. Media changes, 50-75% of

the total volume, were performed every 2 days. On Day 5, EBs were harvested according to the

manufacturer protocol and transferred to a 35mm imaging plate (Cellvis, cat # D35-14-1.5-N)

coated with Matrigel. Daily media changes were performed up to day 9 when cells were fixed

with 100% ice-cold Methanol (Fisher Scientific, cat # A454-4). Images were acquired on a Nikon

Instruments Ti2 inverted fluorescence microscope, equipped with a Yokogawa X1 spinning disk

head, Andor DU-897 EMCCD, Plan Apo Lambda 0.75 NA 20X objective for representative

figures, and Plan Fluor 0.45 NA 10X objective for neural rosette quantification, piezo Z-stage, as

well as 405-,488-,561-, and 647 -nm lasers. Acquisition and analysis were performed using NIS-

Elements. Neural rosette quantification was accomplished by scripting a segmentation-based

image analysis routine to detect, enumerate, and measure rosette lumen area based on the ZO-

1 signal. Briefly, max intensity projections of each field were generated, followed by GPU-based

denoising of the resulting image. Intensity-based thresholding was then applied based on

criteria established for ZO-1 signal segmentation using control images. Restrictions on resultant

binaries were implemented to throw out binaries intersecting image borders, morphometries

deviating severely from rosette associated geometries, as well as for those not meeting

minimum size requirements. This routine could be run in batch across many image stacks to

increase the sample size and robust nature of the data. Measured data was exported to Excel

for further analysis.

Cerebral Organoids

Cerebral organoids were generated as described in (Romero-Morales et al., 2019) with some

modifications. Briefly, organoids were generated using the STEMdiff™ Cerebral Organoid Kit

(STEMCELL Technologies; Cat# 08571, 08570). iPSCs were dissociated into single cells using

Gentle Cell Dissociation Reagent (STEMCELL Technologies, cat # 07174) for 8 minutes at

37°C. Homogeneous and reproducible EBs were generated by using a 24-well plate

AggreWell™ 800 (STEMCELL Technologies, cat # 34815). On Day 7, high-quality EBs were

embedded in Matrigel (Corning, cat # 354277). On Day 10, the Matrigel coat was broken by

vigorously pipetting up and down and the healthy organoids were transferred to a 60mm low

attachment culture plate (Eppendorf, cat # 003070119). The plates were then moved to a 37°C

incubator and to a Celltron benchtop shaker for CO_2 incubators (Infors USA, cat # I69222) set

at 85rpm. Full media changes were performed every 3–4 days. Transmitted-light images were

acquired using an EVOS® XL Core Imaging System. The software used for processing was

ImageJ.

Organoid tissue preparation and Immunohistochemistry

Tissue preparation was performed as described in (Romero-Morales et al., 2019). Briefly,

organoids were fixed in 4% Paraformaldehyde in Phosphate Buffered Saline (PBS), washed 3

times with PBS, and then incubated in 30% sucrose solution overnight at 4°C. Organoids were

embedded in 7.5% gelatin/10% sucrose solution (Sigma, catalog G1890-100G and S7903-

250G) and sectioned with a cryostat (Leica CM1950) at 15um thickness. For immunostaining,

slides were washed with PBS before permeabilization with 0.2% Triton-X in PBS for 1 hr.

Tissues were blocked with blocking medium consisting of 10% donkey serum in PBS with 0.1%

Tween-20 (PBST) for 30 min. Incubation with primary and secondary antibodies was done using

standard methods. Confocal images of the organoids were acquired using the aforementioned

spinning disk microscope with Plan Fluor 10X 0.45 NA and Plan Apo Lambda 0.75 NA 20X

objectives (macrostructures) and Apo TIRF 1.49 NA 100X objective (mitochondria imaging).

NIS-Elements software was used for image acquisition and rendering.

RNA Extraction and Synthesis of cDNA

Cells cultured in 6 well plate, were collected after a wash with PBS, using 600µl Trizol reagent.

The samples were spun down at 12,000 g after the addition of 130µl of chloroform and

incubated at room temperature for 3 minutes. The aqueous phase of the sample was collected

200µl at a time until reaching the edge of phase separation. RNA precipitation was done by

incubating with 300µl of isopropanol for 25 minutes, followed by centrifugation at 12,000 g for 10

min at 4°C. The RNA pellet was washed with ethanol, semi-dried, and resuspended in 30µl of

DEPC water. After quantification and adjusting the volume of all the samples to 1µg/µl, the

samples were treated with DNAse (New England Biolabs, cat # M0303). 10µl of this volume was

used to generate cDNA using the manufacturer's protocol (Thermofisher, cat#4368814)

Quantitative RT PCR

1ug of cDNA sample was used to run a q-PCR for the primers mentioned in the table.

QuantStudio 3 Real-Time PCR machine, SYBR green master mix (Thermo Fisher,

cat#4364346), and manufacturer instructions were used to set up the assay.

Immunocytochemistry

Cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences, cat # 15710-S) in

PBS for 20 min at 4ºC. Blocking and permeabilization were done in 5% donkey serum (Jackson

ImmunoResearch Inc, cat # 017-000-121) + 0.3% Triton X-100 (Sigma Aldrich, cat # T9284) in

TBS for 1 hr at room temperature. After this, cells were treated with primary and secondary

antibodies using standard methods. Cells were mounted in Vectashield (Vector Laboratories,

cat # H-1000) prior to imaging.

Western Blotting

Cultured cells were lysed in 1% Triton buffer containing PMSF (ThermoFisher Scientific, cat #

36978), PhosSTOP (Roche, cat # 4906837001), and protease inhibitor cocktail (Roche, cat #

4693132001). Protein concentrations were determined using the bicinchoninic acid (BCA)

method (Thermo Scientific, cat # 23227). Gel samples were prepared by mixing 30µg of protein

with LDS sample buffer (Life Technologies, cat # NP0007) and 2-Mercaptoethanol (BioRad, cat

1610710) and boiled at 95°C for 5 minutes. Samples were run on 4-20% Mini-PROTEAN TGX

precast gels (BioRad, cat # 4561096) and transferred onto polyvinylidene difluoride (PVDF)

membrane (BioRad, cat # 1620177) overnight at 4°C. Membranes were blocked in 5% milk in

TBST prior to primary antibody incubation. Antibodies used for Western blotting are described in

the Key Resource table.

Cell titer blue assay

After the 24-h exposure to individual treatments of 50µM etoposide, 80µM CCCP, 100ng/mL

nocodazole, and 5ng/mL neocarzinostatin, 20 µl of Cell Titer Blue reagent from Cell Titer Blue

assay (Promega, cat # G8081) was added to each well of 96 well plate. Background

fluorescence was calculated by adding 10% Triton in PBS to some wells. The fluorescence

generated by the reduction of resazurin to resorufin by live cells was measured using a

Beckman coulter DTX 880 multimode plate reader (Beckman Coulter, Brea, California) (570/600

nm).

Metabolomics analysis

Neuralized embryoid bodies were generated by seeding $3x10^6$ cells/well in an Aggewell800^{TM}.

Media change was performed on days 2 and 4. On day 5, EBs were transferred to a low

attachment 60mm plate and maintained in suspension with daily media changes. On day 10,

~150 EBs were collected in a 1.5mL centrifuge tube, rinsed with ice-cold sterile 0.9% NaCl and

flash-frozen in liquid nitrogen. For organoids, at least 4 individual organoids were collected at

day 40, rinsed with ice-cold sterile 0.9% NaCl and flash-freeze in liquid nitrogen.

For metabolite extraction, cells were resuspended in 225uL of cold 80% HPLC grade

methanol/20% HPLC grade water per $1x10^6$ cells. After resuspension, cells were flash-frozen in

liquid nitrogen and thawed rapidly in a 37°C water bath 3 times. Next debris was removed by

centrifugation at max speed in a tabletop microcentrifuge at 4°C for 15 min. Metabolite-

containing supernatant was transferred to a new tube, dried, and resuspended in 50%

acetonitrile while the pellet was used for protein quantification. Samples were analyzed by Ultra-

High-Performance Liquid Chromatography and High-Resolution Mass Spectrometry and

Tandem Mass Spectrometry (UHPLC-MS/MS). Specifically, the system consisted of a Thermo

Q-Exactive in line with an electrospray source and an Ultimate3000 (Thermo) series HPLC

consisting of a binary pump, degasser, and auto-sampler outfitted with an Xbridge Amide

column (Waters; dimensions of 4.6mm × 100mm and a 3.5µm particle size). Mobile phase A

contained 95% (vol/vol) water, 5% (vol/vol) acetonitrile, 10mM ammonium hydroxide, 10mM

ammonium acetate, pH = 9.0; and mobile phase B was 100% Acetonitrile. The gradient was as

follows: 0 min, 15% A; 2.5 min, 30% A; 7 min, 43% A; 16 min, 62% A; 16.1-18 min, 75% A; 18-

25 min, 15% A with a flow rate of 400µL/min. The capillary of the ESI source was set to 275°C,

with sheath gas at 45 arbitrary units, auxiliary gas at 5 arbitrary units, and the spray voltage at

4.0kV. In positive/negative polarity switching mode, an m/z scan range from 70 to 850 was

chosen, and MS1 data was collected at a resolution of 70,000. The automatic gain control

(AGC) target was set at $1x10^6$ and the maximum injection time was 200 ms. The top 5 precursor

ions were subsequently fragmented, in a data-dependent manner, using the higher energy

collisional dissociation (HCD) cell set to 30% normalized collision energy in MS2 at a resolution

power of 17,500. Data acquisition and analysis were carried out by Xcalibur 4.1 software and

Tracefinder 4.1 software, respectively (both from Thermo Fisher Scientific). The peak area for

each detected metabolite was normalized by the total ion current which was determined by

integration of all of the recorded peaks within the acquisition window.

Bioinformatic Analysis

Bioinformatic analysis began with Variant Call Format (VCF) files provided by GENEWIZ (see

Whole Exome sequencing section above), both for SNP and indels. SnpSift version 4.3t (PMID:

22435069) was used to process and filter these files for downstream analysis. Details extracted

included gene symbol, Entrez gene ID and name, UniProt ID, Ensembl ID, chromosome and

position, reference variant, alternative variant, quality of the call, allele name, type of SNP,

impact of the SNP, and the genotype of each sample. From these filtered outputs, we generated

SNP/indel reports that allowed us to look at sample-specific SNPs and indels, as well as

perform aggregate-level functions for grouping and statistical analysis.

To generate the SNP/indel circular chromosome plots, the top 20 genes that had variants in all

three samples were plotted, ranked by frequency of variants per gene. The outside track is used

to visualize the chromosomes and marked gene locations. For each sample, we used a single

track to show the variant frequency as a circular scatter plot, with the height of the scatter points

representative of the variant quality metric, which is a Phred-scaled probability that a REF/ALT

polymorphism exists at the variant site. Similarly, for SNPs in the mitochondrial chromosome,

we used the same approach for visualization.

Quantification and Statistical Analysis

No statistical methods were used to pre-determine sample sizes. All experiments were performed with a minimum of 3 biological replicates unless specified. Statistical significance was determined by unpaired Student's t-test or by one- or two-way ANOVA as appropriate for each experiment. GraphPad Prism v8.1.2 was used for all statistical analysis and data visualization. Error bars in all bar graphs represent standard error of the mean or standard deviation as described for each Figure, while scattered dot plots were represented with boxes (with median and SD) and whiskers (minimum and maximum values).

For neural rosette experiments, ROI were randomly selected using the nuclear (DAPI) staining channel. Images were processed with NIS Elements software with our Neural rosette lumen identification Macro. Outliers were removed from the neural rosette area analysis as post-processing quality control for the NIS Element macro using GraphPad Prism v8.1.2. ROUT (Robust regression and Outlier removal) method was used with a False Discovery Rate of 1%.

For cerebral organoid experiments, 4 independent batches were generated. At time points day 30 and day100, at least 5 organoids per cell line were collected. Immunofluorescence images of at least 3 independent organoids were acquired per condition slide. Image processing was done by NIS Elements and Fiji software. Organoid efficiency evaluation was performed on day 10 using 4X transmitted-light images acquired using an EVOS® XL microscope. Two observers were blinded to the cell line identifier and counted the number of normal and defective (no epithelial buds or more than 75% of the area is not developed) organoids. Criteria for normal and defective organoids was based on (Lancaster and Knoblich, 2014).

For the metabolomics experiments, two independent LC/MS runs of ~150-200 neuralized EBs/cell line/replica were utilized with enough material to account for 3×10^6 cells/run. For the cerebral organoid metabolomics, one replica of day 40 organoids was performed, with 4 samples per genotype.

Software and Data Availability

All raw data in FastQ format for whole-exome sequencing and mitochondrial sequencing have

been deposited to the Short Read Archive as BioProject PRJNA626388, available at

https://www.ncbi.nlm.nih.gov/sra/PRJNA626388. All source code and documents are available

via https://vandydata.github.io/Romero-Morales-Gama-Leigh-Syndrome-WES/.

Author contributions

A.R.M. and V.G. designed experiments, interpret the data, and wrote the manuscript. A.R.M.,

A.R., G.L.R., and M.L.R. performed experiments and analyzed data with technical support of

H.T. J.P.C. and L.H. analyzed the genomic data, generated the corresponding figures, and the

associated method section. P.M.A provided the pluripotency data analysis and technical support

for data analysis. B.M generated the neural rosette analysis macro for NIS Elements. N.S.C.

and G.S.M provided technical support, led the experimental design and data analysis for

metabolomics and its corresponding method section.

Conflict of interests

The authors declare no competing interests.

Acknowledgments

We thank Dr. Nicholas Mignemi (Vanderbilt Nikon Center for Excellence) for his technical

support with image acquisition and processing. We would like to thank members of the Gama

Laboratory for helpful discussions and comments on the manuscript. Funding was provided by

the following grants: 1R35GM128915-01 (to VG); 1R21 CA227483-01A1 (to VG); the Precision

Medicine and Mental Health Initiative sponsored by the Vanderbilt Brain Institute (to VG). Image

acquisition and analysis were performed in part through the use of the Nikon Center of

Excellence within the Vanderbilt Cell Imaging Shared Resource (supported by NIH grants

CA68485, DK20593, DK58404, DK59637, and EY08126), Vanderbilt University Medical

Center's Translational Pathology Shared Resource supported by NCI/NIH Cancer Center

Support Grant 2P30 CA068485-14, and the Vanderbilt Mouse Metabolic Phenotyping Center

Grant 5U24DK059637-13. Whole exome sequencing and mitochondrial sequencing results

were analyzed by Creative Solutions. Metabolite measurements were performed by the

Northwestern University RHLCCC Metabolomics Core (Peng Gao) and were supported by the

following NIH grants to N.S.C.: NIH2PO1HL071643-11A1, NIH1R35CA197532-01,

NIH1PO1AG049665-01 and NIH/NCI grant to G.S.M: T32CA09560.

References

Agana, M., Frueh, J., Kamboj, M., Patel, D.R., and Kanungo, S. (2018). Common metabolic disorder (inborn errors of metabolism) concerns in primary care practice. Ann. Transl. Med. *6*, 469–469.

Albright, C.D., Friedrich, C.B., Brown, E.C., Mar, M.H., and Zeisel, S.H. (1999a). Maternal dietary choline availability alters mitosis, apoptosis and the localization of TOAD-64 protein in the developing fetal rat septum. Dev. Brain Res. *115*, 123–129.

Albright, C.D., Tsai, A.Y., Friedrich, C.B., Mar, M.-H., and Zeisel, S.H. (1999b). Choline availability alters embryonic development of the hippocampus and septum in the rat. Dev. Brain Res. *113*, 13–20.

Angelova, M.T., Dimitrova, D.G., Dinges, N., Lence, T., Worpenberg, L., Carré, C., and Roignant, J.Y. (2018). The emerging field of epitranscriptomics in neurodevelopmental and neuronal disorders. Front. Bioeng. Biotechnol. *6*, 46.

Baertling, F., Rodenburg, R.J., Schaper, J., Smeitink, J.A., Koopman, W.J.H., Mayatepek, E., Morava, E., and Distelmaier, F. (2014). A guide to diagnosis and treatment of Leigh syndrome. J. Neurol. Neurosurg. Psychiatry *85*, 257–265.

Baertling, F., Klee, D., Haack, T.B., Prokisch, H., Meitinger, T., Mayatepek, E., Schaper, J., and Distelmaier, F. (2016). The many faces of paediatric mitochondrial disease on neuroimaging. Child's Nerv. Syst. *32*, 2077–2083.

Barshop, B.A. (2004). Metabolomic approaches to mitochondrial disease: Correlation of urine organic acids. Mitochondrion *4*, 521–527.

Bridi, R., Latini, A., Braum, C.A., Zorzi, G.K., Wajner, M., Lissi, E., and Dutra-Filho, C.S. (2005). Evaluation of the mechanisms involved in leucine-induced oxidative damage in cerebral cortex of young rats. Free Radic. Res. *39*, 71–79.

Buzkova, J., Nikkanen, J., Ahola, S., Hakonen, A.H., Sevastianova, K., Hovinen, T., Yki-Järvinen, H., Pietiläinen, K.H., Lönnqvist, T., Velagapudi, V., et al. (2018). Metabolomes of mitochondrial diseases and inclusion body myositis patients: treatment targets and biomarkers. EMBO Mol. Med. *10*.

Calvo, S., Jain, M., Xie, X., Sheth, S.A., Chang, B., Goldberger, O.A., Spinazzola, A., Zeviani, M., Carr, S.A., and Mootha, V.K. (2006). Systematic identification of human mitochondrial disease genes through integrative genomics. Nat. Genet. *38*, 576–582.

Calvo, S.E., Compton, A.G., Hershman, S.G., Lim, S.C., Lieber, D.S., Tucker, E.J., Laskowski, A., Garone, C., Liu, S., Jaffe, D.B., et al. (2012). Molecular diagnosis of infantile mitochondrial disease with targeted next-generation sequencing. Sci. Transl. Med. *4*, 118ra10-118ra10.

Camp, J.G., Badsha, F., Florio, M., Kanton, S., Gerber, T., Wilsch-Bräuninger, M., Lewitus, E., Sykes, A., Hevers, W., Lancaster, M., et al. (2015). Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. Proc. Natl. Acad. Sci. *112*, 201520760.

Cansev, M. (2006). Uridine and cytidine in the brain: Their transport and utilization. Brain Res. Rev. *52*, 389–397.

Cerutti, R., Pirinen, E., Lamperti, C., Marchet, S., Sauve, A.A., Li, W., Leoni, V., Schon, E.A., Dantzer, F., Auwerx, J., et al. (2014). NAD+-dependent activation of Sirt1 corrects the phenotype in a mouse model of mitochondrial disease. Cell Metab. *19*, 1042–1049.

Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. Nat. Biotechnol. *27*, 275–280.

Chandrasekaran, A., Avci, H.X., Ochalek, A., Rösingh, L.N., Molnár, K., László, L., Bellák, T., Téglási, A., Pesti, K., Mike, A., et al. (2017). Comparison of 2D and 3D neural induction

methods for the generation of neural progenitor cells from human induced pluripotent stem
cells. Stem Cell Res. *25*, 139–151.

Chen, H., and Chan, D.C. (2017). Mitochondrial Dynamics in Regulating the Unique Phenotypes
of Cancer and Stem Cells. Cell Metab. *26*, 39–48.

Chong, J., and Xia, J. (2018). MetaboAnalystR: an R package for flexible and reproducible
analysis of metabolomics data. Bioinformatics *34*, 4313–4314.

Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., Wishart, D.S., and Xia, J. (2018).
MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis.
Nucleic Acids Res. *46*, W486–W494.

Chong, J., Wishart, D.S., and Xia, J. (2019). Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. Curr. Protoc. Bioinforma. *68*.

Cingolani, P., Patel, V.M., Coon, M., Nguyen, T., Land, S.J., Ruden, D.M., and Lu, X. (2012).
Using Drosophila melanogaster as a model for genotoxic chemical mutational studies with
a new program, SnpSift. Front. Genet. *3*, 35.

Craigen, W.J. (1996). Leigh disease with deficiency of lipoamide dehydrogenase: Treatment
failure with dichloroacetate. Pediatr. Neurol. *14*, 69–71.

Craven, L., Alston, C.L., Taylor, R.W., and Turnbull, D.M. (2017). Recent Advances in
Mitochondrial Disease. Annu. Rev. Genomics Hum. Genet. *18*, 257–275.

Cui, Y., Chen, X., Luo, H., Fan, Z., Luo, J., He, S., Yue, H., Zhang, P., and Chen, R. (2016).
BioCircos.js: An interactive Circos JavaScript library for biological data visualization on
web applications. Bioinformatics *32*, 1740–1742.

Das, A.M., Steuerwald, U., and Illsinger, S. (2010). Inborn errors of energy metabolism
associated with myopathies. J. Biomed. Biotechnol. *2010*.

Diez-Roux, G., Banfi, S., Sultan, M., Geffers, L., Anand, S., Rozado, D., Magen, A., Canidio, E.,
Pagani, M., Peluso, I., et al. (2011). A high-resolution anatomical atlas of the transcriptome
in the mouse embryo. PLoS Biol. *9*, e1000582.

Elkabetz, Y., Panagiotakos, G., Al Shamy, G., Socci, N.D., Tabar, V., and Studer, L. (2008).
Human ES cell-derived neural rosettes reveal a functionally distinct early neural stem cell
stage. Genes Dev. *22*, 152–165.

Esterhuizen, K., van der Westhuizen, F.H., and Louw, R. (2017). Metabolomics of mitochondrial
disease. Mitochondrion *35*, 97–110.

Fang, Q., Zhang, Y., Chen, X., Li, H., Cheng, L., Zhu, W., Zhang, Z., Tang, M., Liu, W., Wang,
H., et al. (2020). Three-Dimensional Graphene Enhances Neural Stem Cell Proliferation
Through Metabolic Regulation. Front. Bioeng. Biotechnol. *7*, 436.

Feeney, C.L., Lim, A.Z., Fagan, E., Blain, A., Bright, A., Maddison, J., Devine, H., Stewart, J.,
Taylor, R.W., Gorman, G.S., et al. (2019). A case-comparison study of pregnant women
with mitochondrial disease – what to expect? BJOG An Int. J. Obstet. Gynaecol. *126*,
1380–1389.

Ferlin, T., Landrieu, P., Rambaud, C., Fernandez, H., Dumoulin, R., Rustin, P., and Mousson, B.
(1997). Segregation of the G8993 mutant mitochondrial DNA through generations and
embryonic tissues family at risk of Leigh syndrome. J. Pediatr. *131*, 447–449.

Ferrari, M., Jain, I.H., Goldberger, O., Rezoagli, E., Thoonen, R., Chen, K.-H., Sosnovik, D.E.,
Scherrer-Crosbie, M., Mootha, V.K., and Zapol, W.M. (2017). Hypoxia treatment reverses
neurodegenerative disease in a mouse model of Leigh syndrome. Proc. Natl. Acad. Sci.
114, E4241–E4250.

Ferreira, G.C., and McKenna, M.C. (2017). l-Carnitine and Acetyl-l-carnitine Roles and
Neuroprotection in Developing Brain. Neurochem. Res. *42*, 1661–1675.

Finsterer, J. (2008). Leigh and Leigh-Like Syndrome in Children and Adults. Pediatr. Neurol. *39*,
223–235.

Florio, M., and Huttner, W.B. (2014). Neural progenitors, neurogenesis and the evolution of the
neocortex. Development *141*, 2182–2194.

Galera-Monge, T., Zurita-Díaz, F., González-Páramos, C., Moreno-Izquierdo, A., Fraga, M.F., Fernández, A.F., Garesse, R., and Gallardo, M.E. (2016). Generation of a human iPSC line from a patient with Leigh syndrome caused by a mutation in the MT-ATP6 gene. Stem Cell Res. *16*, 766–769.

García-Cazorla, A., Oyarzabal, A., Fort, J., Robles, C., Castejón, E., Ruiz-Sala, P., Bodoy, S., Merinero, B., Lopez-Sala, A., Dopazo, J., et al. (2014). Two Novel Mutations in the BCKDK (Branched-Chain Keto-Acid Dehydrogenase Kinase) Gene Are Responsible for a Neurobehavioral Deficit in Two Pediatric Unrelated Patients. Hum. Mutat. *35*, 470–477.

Gerards, M., Sallevelt, S.C.E.H., and Smeets, H.J.M. (2016). Leigh syndrome: Resolving the clinical and genetic heterogeneity paves the way for treatment options. Mol. Genet. Metab. *117*, 300–312.

Gogos, J.A., Santha, M., Takacs, Z., Beck, K.D., Luine, V., Lucas, L.R., Nadler, J.V., and Karayiorgou, M. (1999). The gene encoding proline dehydrogenase modulates sensorimotor gating in mice. Nat. Genet. *21*, 434–439.

Gorman, G.S., Chinnery, P.F., DiMauro, S., Hirano, M., Koga, Y., McFarland, R., Suomalainen, A., Thorburn, D.R., Zeviani, M., and Turnbull, D.M. (2016). Mitochondrial diseases. Nat. Rev. Dis. Prim. *2*, 1–22.

Grace, H.E., Galdun, P., Lesnefsky, E.J., West, F.D., and Iyer, S. (2019). MRNA Reprogramming of T8993G Leigh's Syndrome Fibroblast Cells to Create Induced Pluripotent Stem Cell Models for Mitochondrial Disorders. Stem Cells Dev. *28*, 846–859.

Grier, J., Hirano, M., Karaa, A., Shepard, E., and Thompson, J.L.P. (2018). Diagnostic odyssey of patients with mitochondrial disease Results of a survey. Neurol. Genet. *4*.

Harding, A.E., Holt, I.J., Sweeney, M.G., Brockington, M., and Davis, M.B. (1992). Prenatal diagnosis of mitochondrial DNA8993 T→G disease. Am. J. Hum. Genet. *50*, 629–633.

Hattori, T., Hamazaki, T., Kudo, S., and Shintaku, H. (2016). Metabolic Signature of MELAS/Leigh Overlap Syndrome in Patient-specific Induced Pluripotent Stem Cells Model. Osaka City Med. J. *62*, 69–76.

Hinman, L.M., Sheu, K.-F.R., Baker, A.C., Kim, Y.T., and Blass, J.P. (1989). Deficiency of pyruvate dehydrogenase complex (PDHC) in Leigh's disease fibroblasts: An abnormality in lipoamide dehydrogenase affecting PDHC activation. Neurology *39*, 70–70.

Homem, C.C.F., Repic, M., and Knoblich, J.A. (2015). Proliferation control in neural stem and progenitor cells. Nat. Rev. Neurosci. *16*, 647–659.

Hříbková, H., Grabiec, M., Klemová, D., Slaninová, I., and Sun, Y.-M. (2018). Five steps to form neural rosettes: structure and function. J. Cell Sci. *131*, jcs206896.

Huh, T.L., Casazza, J.P., Huh, J.W., Chi, Y.T., and Song, B.J. (1990). Characterization of two cDNA clones for pyruvate dehydrogenase E1β subunit and its regulation in tricarboxylic acid cycle-deficient fibroblast. J. Biol. Chem. *265*, 13320–13326.

Inak, G., Lorenz, C., Lisowski, P., Zink, A., Mlody, B., and Prigione, A. (2017). Concise Review: Induced Pluripotent Stem Cell-Based Drug Discovery for Mitochondrial Disease. Stem Cells *35*, 1655–1662.

Inak, G., Rybak-Wolf, A., Lisowski, P., Juettner, R., Zink, A., Mlody, B., Glazar, P., Secker, C., Ciptasari, U.H., Stenzel, W., et al. (2019). SURF1 mutations causative of Leigh syndrome impair human neurogenesis. BioRxiv 551390.

Itskovitz-Eldor, J., Schuldiner, M., Karsenti, D., Eden, A., Yanuka, O., Amit, M., Soreq, H., and Benvenisty, N. (2000). Differentiation of Human Embryonic Stem Cells into Embryoid Bodies Comprising the Three Embryonic Germ Layers. Mol. Med. *6*, 88–95.

Iyer, S., Bergquist, K., Young, K., Gnaiger, E., Rao, R.R., and Bennett, J.P. (2012). Mitochondrial Gene Therapy Improves Respiration, Biogenesis, and Transcription in G11778A Leber's Hereditary Optic Neuropathy and T8993G Leigh's Syndrome Cells. Hum. Gene Ther. *23*, 647–657.

Jain, I.H., Zazzeron, L., Goli, R., Alexa, K., Schatzman-Bone, S., Dhillon, H., Goldberger, O.,

Peng, J., Shalem, O., Sanjana, N.E., et al. (2016). Hypoxia as a therapy for mitochondrial disease. Science (80-.). *352*, 54–61.

Jain, I.H., Zazzeron, L., Goldberger, O., Marutani, E., Wojtkiewicz, G.R., Ast, T., Wang, H., Schleifer, G., Stepanova, A., Brepoels, K., et al. (2019). Leigh Syndrome Mouse Model Can Be Rescued by Interventions that Normalize Brain Hyperoxia, but Not HIF Activation. Cell Metab. *0*, 824-832.e3.

Jernberg, J.N., Bowman, C.E., Wolfgang, M.J., and Scafidi, S. (2017). Developmental regulation and localization of carnitine palmitoyltransferases (CPTs) in rat brain. J. Neurochem. *142*, 407–419.

Kelava, I., and Lancaster, M.A. (2016). Stem Cell Models of Human Brain Development. Cell Stem Cell *18*, 736–748.

Khacho, M., Clark, A., Svoboda, D.S., Azzi, J., MacLaurin, J.G., Meghaizel, C., Sesaki, H., Lagace, D.C., Germain, M., Harper, M.E., et al. (2016). Mitochondrial Dynamics Impacts Stem Cell Identity and Fate Decisions by Regulating a Nuclear Transcriptional Program. Cell Stem Cell.

Lake, N.J., Bird, M.J., Isohanni, P., and Paetau, A. (2015). Leigh syndrome: neuropathology and pathogenesis. J. Neuropathol. Exp. Neurol. *74*, 482–492.

Lake, N.J., Compton, A.G., Rahman, S., and Thorburn, D.R. (2016). Leigh syndrome: One disorder, more than 75 monogenic causes. Ann. Neurol. *79*, 190–203.

Lancaster, M.A., and Knoblich, J.A. (2014). Generation of cerebral organoids from human pluripotent stem cells. Nat. Protoc. *9*, 2329–2340.

Lancaster, M.A., Corsini, N.S., Wolfinger, S., Gustafson, E.H., Phillips, A.W., Burkard, T.R., Otani, T., Livesey, F.J., and Knoblich, J.A. (2017). Guided self-organization and cortical plate formation in human brain organoids. Nat. Publ. Gr. *35*.

Leigh, D. (1951). Subacute necrotizing encephalomyelopathy in an infant. J. Neurol. Neurosurg. Psychiatry *14*, 216–221.

Lorenz, C., Lesimple, P., Bukowiecki, R., Zink, A., Inak, G., Mlody, B., Singh, M., Semtner, M., Mah, N., Auré, K., et al. (2017). Human iPSC-Derived Neural Progenitors Are an Effective Drug Discovery Model for Neurological mtDNA Disorders. Cell Stem Cell *20*, 659-674.e9.

Lui, J.H., Hansen, D. V., and Kriegstein, A.R. (2011). Development and evolution of the human neocortex. Cell *146*, 18–36.

Di Lullo, E., and Kriegstein, A.R. (2017). The use of brain organoids to investigate neural development and disease. Nat. Rev. Neurosci. *18*, 573–584.

Ma, H., Folmes, C.D.L., Wu, J., Morey, R., Mora-Castilla, S., Ocampo, A., Ma, L., Poulton, J., Wang, X., Ahmed, R., et al. (2015). Metabolic rescue in pluripotent cells from patients with mtDNA disease. Nature *524*, 234–238.

Manoli, I., and Venditti, C.P. (2016). Disorders of branched chain amino acid metabolism. Transl. Sci. Rare Dis. *1*, 91–110.

Mathieu, J., and Ruohola-Baker, H. (2017). Metabolic remodeling during the loss and acquisition of pluripotency. Dev. *144*, 541–551.

Medelnik, J.-P., Roensch, K., Okawa, S., del Sol, A., Chara, O., Mchedlishvili, L., and Tanaka, E.M. (2018). Signaling-Dependent Control of Apical Membrane Size and Self-Renewal in Rosette-Stage Human Neuroepithelial Stem Cells. Stem Cell Reports *10*, 1751–1765.

Di Meo, I., Marchet, S., Lamperti, C., Zeviani, M., and Viscomi, C. (2017). AAV9-based gene therapy partially ameliorates the clinical phenotype of a mouse model of Leigh syndrome. Gene Ther. *24*, 661–667.

Miyabayashi, S., Ito, T., Narisawa, K., Iinuma, K., and Tada, K. (1985). Biochemical study in 28 children with lactic acidosis, in relation to Leigh's encephalomyelopathy. Eur. J. Pediatr. *143*, 278–283.

Müller, F.J., Schuldt, B.M., Williams, R., Mason, D., Altun, G., Papapetrou, E.P., Danner, S., Goldmann, J.E., Herbst, A., Schmidt, N.O., et al. (2011). A bioinformatic assay for

1029   pluripotency in human cells. Nat. Methods *8*, 315–317.

1030 Muratore, C.R., Srikanth, P., Callahan, D.G., and Young-Pearse, T.L. (2014). Comparison and
1031   optimization of hiPSC forebrain cortical differentiation protocols. PLoS One *9*, e105807.

1032 Nakano, C., Takashima, S., and Takeshita, K. (1989). Carnitine concentration during the
1033   development of human tissues. Early Hum. Dev. *19*, 21–27.

1034 Nesbitt, V., Alston, C.L., Blakely, E.L., Fratter, C., Feeney, C.L., Poulton, J., Brown, G.K.,
1035   Turnbull, D.M., Taylor, R.W., and McFarland, R. (2014). A national perspective on prenatal
1036   testing for mitochondrial disease. Eur. J. Hum. Genet. *22*, 1255–1259.

1037 Oshida, K., Shimizu, T., Takase, M., Tamura, Y., Shimizu, T., and Yamashiro, Y. (2003). Effects
1038   of dietary sphingomyelin on central nervous system myelination in developing rats.
1039   Pediatr. Res. *53*, 589–593.

1040 Pampaloni, F., Reynaud, E.G., and Stelzer, E.H.K. (2007). The third dimension bridges the gap
1041   between cell culture and live tissue. Nat. Rev. Mol. Cell Biol. *8*, 839–845.

1042 Panopoulos, A.D., Yanes, O., Ruiz, S., Kida, Y.S., Diep, D., Tautenhahn, R., Herrerías, A.,
1043   Batchelder, E.M., Plongthongkum, N., Lutz, M., et al. (2012). The metabolome of induced
1044   pluripotent stem cells reveals metabolic changes occurring in somatic cell reprogramming.
1045   Cell Res. *22*, 168–177.

1046 Parikh, S., Goldstein, A., Karaa, A., Koenig, M.K., Anselm, I., Brunel-Guitton, C., Christodoulou,
1047   J., Cohen, B.H., Dimmock, D., Enns, G.M., et al. (2017). Patient care standards for
1048   primary mitochondrial disease: A consensus statement from the mitochondrial medicine
1049   society. Genet. Med. *19*, 1–18.

1050 Paşca, A.M., Sloan, S.A., Clarke, L.E., Tian, Y., Makinson, C.D., Huber, N., Kim, C.H., Park, J.-
1051   Y., O'Rourke, N.A., Nguyen, K.D., et al. (2015). Functional cortical neurons and astrocytes
1052   from human pluripotent stem cells in 3D culture. Nat Methods. *12*, 671–678.

1053 Pastores, G.M., Santorelli, F.M., Shanske, S., Gelb, B.D., Fyfe, B., Wolfe, D., and Willner, J.P.
1054   (1994). Leigh syndrome and hypertrophic cardiomyopathy in an infant with a mitochondrial
1055   DNA point mutation (T8993G). Am. J. Med. Genet. *50*, 265–271.

1056 Quadrato, G., Brown, J., and Arlotta, P. (2016). The promises and challenges of human brain
1057   organoids as models of neuropsychiatric disease. Nat. Med. *22*, 1220–1228.

1058 Rahman, J., and Rahman, S. (2018). Mitochondrial medicine in the omics era. Lancet *391*,
1059   2560–2574.

1060 Rasmussen, M.L., and Gama, V. (2020). A connection in life and death: The BCL-2 family
1061   coordinates mitochondrial network dynamics and stem cell fate. In International Review of
1062   Cell and Molecular Biology, (Elsevier Inc.), p.

1063 Rastogi, A., Joshi, P., Contreras, E., and Gama, V. (2019). Remodeling of mitochondrial
1064   morphology and function: an emerging hallmark of cellular reprogramming. Cell Stress *3*,
1065   181–194.

1066 Rema, V., Bali, K.K., Ramachandra, R., Chugh, M., Darokhan, Z., and Chaudhary, R. (2008).
1067   Cytidine-5-diphosphocholine supplement in early life induces stable increase in dendritic
1068   complexity of neurons in the somatosensory cortex of adult rats. Neuroscience *155*, 556–
1069   564.

1070 Robel, S., Mori, T., Zoubaa, S., Schlegel, J., Sirko, S., Faissner, A., Goebbels, S., Dimou, L.,
1071   and Götz, M. (2009). Conditional deletion of β1-integrin in astroglia causes partial reactive
1072   gliosis. Glia *57*, 1630–1647.

1073 Robel, S., Berninger, B., and Götz, M. (2011). The stem cell potential of glia: Lessons from
1074   reactive gliosis. Nat. Rev. Neurosci. *12*, 88–104.

1075 Romero-Morales, A.I., O'Grady, B.J., Balotin, K.M., Bellan, L.M., Lippmann, E.S., and Gama, V.
1076   (2019). Spin∞ an improved miniaturized spinning bioreactor for the generation of human
1077   cerebral organoids from pluripotent stem cells. BioRxiv 687095.

1078 Saito, T., Hanai, S., Takashima, S., Nakagawa, E., Okazaki, S., Inoue, T., Miyata, R., Hoshino,
1079   K., Akashi, T., Sasaki, M., et al. (2011). Neocortical layer formation of human developing

brains and lissencephalies: Consideration of layer-specific marker expression. Cereb.
Cortex *21*, 588–596.
Schaefer, A., Lim, A., and Gorman, G. (2019). Epidemiology of Mitochondrial Disease. In
Diagnosis and Management of Mitochondrial Disorders, (Cham: Springer International
Publishing), pp. 63–79.
Schubert, M.B., and Vilarinho, L. (2020). Molecular basis of Leigh syndrome: A current look.
Orphanet J. Rare Dis. *15*, 1–14.
Shaheen, R., Han, L., Faqeih, E., Ewida, N., Alobeid, E., Phizicky, E.M., and Alkuraya, F.S.
(2016). A homozygous truncating mutation in PUS3 expands the role of tRNA modification
in normal cognition. Hum. Genet. *135*, 707–713.
Song, C., Xu, F., Ren, Z., Zhang, Y., Meng, Y., Yang, Y., Lingadahalli, S., Cheung, E., Li, G.,
Liu, W., et al. (2019). Elevated Exogenous Pyruvate Potentiates Mesodermal
Differentiation through Metabolic Modulation and AMPK/mTOR Pathway in Human
Embryonic Stem Cells. Stem Cell Reports *13*, 338–351.
Sorbi, S., and Blass, J.P. (1982). Abnormal activation of pyruvate dehydrogenase in Leigh
disease fibroblasts. Neurology *32*, 555–558.
Steffann, J., Gigarel, N., Corcos, J., Bonnière, M., Encha-Razavi, F., Sinico, M., Prevot, S.,
Dumez, Y., Yamgnane, A., Frydman, R., et al. (2007). Stability of the m.8993T→G mtDNA
mutation load during human embryofetal development has implications for the feasibility of
prenatal diagnosis in NARP syndrome. J. Med. Genet. *44*, 664–669.
Stern, H.J. (1994). Lactic acidosis in paediatrics: Clinical and laboratory evaluation. Ann. Clin.
Biochem. *31*, 410–419.
Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., and Yamanaka, S.
(2007). Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined
Factors. Cell *131*, 861–872.
TCW, J., Wang, M., Pimenova, A.A., Bowles, K.R., Hartley, B.J., Lacin, E., Machlovi, S.I.,
Abdelaal, R., Karch, C.M., Phatnani, H., et al. (2017). An Efficient Platform for Astrocyte
Differentiation from Human Induced Pluripotent Stem Cells. Stem Cell Reports *9*, 600–
614.
Tibbitt, M.W., and Anseth, K.S. (2012). Dynamic microenvironments: The fourth dimension. Sci.
Transl. Med. *4*, 160ps24-160ps24.
Uittenbogaard, M., Brantner, C.A., Fang, Z.S., Wong, L.J.C., Gropman, A., and Chiaramello, A.
(2018). Novel insights into the functional metabolic impact of an apparent de novo
m.8993T>G variant in the MT-ATP6 gene associated with maternally inherited form of
Leigh Syndrome. Mol. Genet. Metab. *124*, 71–81.
Vescovi, A.L., and Snyder, E.Y. (1999). Establishment and properties of neural stem cell clones:
Plasticity in vitro and in vivo. In Brain Pathology, (International Society of Neuropathology),
pp. 569–598.
Vorstman, J.A.S., Turetsky, B.I., Sijmens-Morcus, M.E.J., De Sain, M.G., Dorland, B., Sprong,
M., Rappaport, E.F., Beemer, F.A., Emanuel, B.S., Kahn, R.S., et al. (2009). Proline
affects brain function in 22q11DS children with the low activity COMT158 allele.
Neuropsychopharmacology *34*, 739–746.
White, S.L., Collins, V.R., Wolfe, R., Cleary, M.A., Shanske, S., DiMauro, S., Dahl, H.H.M., and
Thorburn, D.R. (1999). Genetic counseling and prenatal diagnosis for the mitochondrial
DNA mutations at nucleotide 8993. Am. J. Hum. Genet. *65*, 474–482.
Wilson, P.G., and Stice, S.S. (2006). Development and differentiation of neural rosettes derived
from human embryonic stem cells. Stem Cell Rev. *2*, 67–77.
Wu, J., Ocampo, A., and Belmonte, J.C.I. (2016). Cellular Metabolism and Induced
Pluripotency. Cell *166*, 1371–1385.
Xia, J., and Wishart, D.S. (2010). MetPA: a web-based metabolomics tool for pathway analysis
and visualization. Bioinformatics *26*, 2342–2344.

Xia, J., and Wishart, D.S. (2011a). Metabolomic data processing, analysis, and interpretation
using MetaboAnalyst. Curr. Protoc. Bioinforma. *34*, 1–48.
Xia, J., and Wishart, D.S. (2011b). Web-based inference of biological patterns, functions and
pathways from metabolomic data using MetaboAnalyst. Nat. Protoc. *6*, 743–760.
Xia, J., Psychogios, N., Young, N., and Wishart, D.S. (2009). MetaboAnalyst: a web server for
metabolomic data analysis and interpretation. Nucleic Acids Res. *37*, W652–W660.
Zanotelli, M.R., Goldblatt, Z.E., Miller, J.P., Bordeleau, F., Li, J., VanderBurgh, J.A., Lampi,
M.C., King, M.R., and Reinhart-King, C.A. (2018). Regulation of ATP utilization during
metastatic cell migration by collagen architecture. Mol. Biol. Cell *29*, 1–9.
Zanotelli, M.R., Rahman-Zaman, A., VanderBurgh, J.A., Taufalele, P. V., Jain, A., Erickson, D.,
Bordeleau, F., and Reinhart-King, C.A. (2019). Energetic costs regulated by cell
mechanics and confinement are predictive of migration path during decision-making. Nat.
Commun. *10*, 1–12.
Zeisel, S.H. (2006). The fetal origins of memory: The role of dietary choline in optimal brain
development. J. Pediatr. *149*, S131–S136.
Zeisel, S.H., and Niculescu, M.D. (2006). Perinatal Choline Influences Brain Structure and
Function. Nutr. Rev. *64*, 197–203.
Zhang, S.-C., Wernig, M., Duncan, I.D., Brüstle, O., and Thomson, J.A. (2001). In vitro
differentiation of transplantable neural precursors from human embryonic stem cells. Nat.
Biotechnol. *19*, 1129–1133.
1151

Figure 2

A

B

C

D

A

B

C

Pathway	Hits	Total	Raw p	-log(p)	Holm adjust	FDR	Impact
Alanine, aspartate and glutamate metabolism	16	28	9.88E-13	27.643	8.30E-11	8.30E-11	0.76282
Synthesis and degradation of ketone bodies	1	5	0.29616	1.2169	1	0.85783	0.6
Arginine biosynthesis	9	14	3.24E-08	17.244	2.69E-06	1.36E-06	0.5736
Glycine, serine and threonine metabolism	8	33	0.001137	6.7797	0.088659	0.01364	0.55868
D-Glutamine and D-glutamate metabolism	3	6	0.005201	5.259	0.38485	0.036405	0.5
Cysteine and methionine metabolism	9	33	0.000209	8.4735	0.016923	0.003721	0.47436
Nicotinate and nicotinamide metabolism	4	15	0.015176	4.188	1	0.091058	0.42895
Beta-Alanine metabolism	2	21	0.42186	0.86309	1	1	0.39925
Pyrimidine metabolism	14	39	6.64E-08	16.527	5.45E-06	1.86E-06	0.32533
Arginine and proline metabolism	8	38	0.00301	5.806	0.22873	0.028089	0.28068
Citrate cycle (TCA cycle)	6	20	0.001488	6.5105	0.11456	0.015622	0.27512
Pentose phosphate pathway	3	22	0.18299	1.6984	1	0.61483	0.27226
Purine metabolism	12	65	0.000959	6.9499	0.075743	0.013423	0.27171
Tyrosine metabolism	7	42	0.02008	3.908	1	0.11245	0.2369
Pyruvate metabolism	3	22	0.18299	1.6984	1	0.61483	0.20684
Glycolysis / Gluconeogenesis	3	26	0.25592	1.3629	1	0.7962	0.20594
Glyoxylate and dicarboxylate metabolism	7	32	0.004395	5.4272	0.32966	0.036405	0.20371

A

B

Control

PDH

α-TUBULIN
CDK5RAP2
Nuclei
ZO-1

DLD

MT-ATP6/PDH

α-TUBULIN
CDK5RAP2
Nuclei
ZO-1

C

D