Experimental and computational study on motor control and recovery after stroke: towards a constructive loop between experimental and virtual embodied neuroscience

Anna Letizia Allegra Mascaro[1,2,⋆,*], **Egidio Falotico**[3,⋆], **Spase Petkoski**[4,⋆], **Maria Pasquini**[3], **Lorenzo Vannucci**[3], **Núria Tort-Colet**[5], **Emilia Conti**[2,6], **Francesco Resta**[2], **Cristina Spalletti**[1], **Shravan Tata Ramalingasetty**[7], **Axel von Arnim**[8], **Emanuele Formento**[9], **Emmanouil Angelidis**[8,10], **Camilla Hagen Blixhavn**[11], **Trygve Brauns Leergaard**[11], **Matteo Caleo**[1,12], **Alain Destexhe**[5], **Auke Ijspeert**[7], **Silvestro Micera**[3,9], **Cecilia Laschi**[3], **Viktor Jirsa**[4], **Marc-Oliver Gewaltig**[13,†] **& Francesco S. Pavone**[2,6,†]

[1]*Neuroscience Institute, National Research Council, Pisa*
[2]*European Laboratory for Non-Linear Spectroscopy, Sesto Fiorentino (Fi)*
[3]*The BioRobotics Institute, Scuola Superiore Sant'Anna, Pontedera (Pisa)*
[4]*Aix-Marseille Université, Inserm, INS UMR_S 1106, Marseille, France*
[5]*Paris-Saclay University, Institute of Neuroscience, CNRS, Gif Sur Yvette, France*
[6]*Department of Physics and Astronomy, University of Florence, Italy*
[7]*Biorobotics Laboratory, École polytechnique fédérale de Lausanne (EPFL), Lausanne, Switzerland*
[8]*Fortiss GmbH, Munich, Germany*
[9]*Bertarelli Foundation Chair in Translational NeuroEngineering, Institute of Bioengineering, Swiss Federal Institute of Technology (EPFL), Lausanne, Switzerland*
[10]*Chair of Robotics, Artificial Intelligence and Embedded Systems, Department of Informatics, Technical University of Munich, Germany*
[11]*Institute of Basic Medical Sciences, University of Oslo, Norway*
[12]*Department of Biomedical Sciences, University of Padua, Italy*
[13]*Blue Brain Project (BBP), École polytechnique fédérale de Lausanne (EPFL), Genève, Switzerland*
⋆, †*: These authors equally contributed to this work.*

Correspondence*:
A. L. Allegra Mascaro
allegra@lens.unifi.it

ABSTRACT

Being able to replicate real experiments with computational simulations is a unique opportunity
to refine and validate models with experimental data and redesign the experiments based on
simulations. However, since it is technically demanding to model all components of an experiment,
traditional approaches to modeling reduce the experimental setups as much as possible. In this

study, our goal is to replicate all the relevant features of an experiment on motor control and
motor rehabilitation after stroke. To this aim, we propose an approach that allows continuous
integration of new experimental data into a computational modeling framework. First, results show
that we could reproduce experimental object displacement with high accuracy via the simulated
embodiment in the virtual world by feeding a spinal cord model with experimental registration of the
cortical activity. Second, by using computational models of multiple granularities, our preliminary
results show the possibility of simulating several features of the brain after stroke, from the
local alteration in neuronal activity to long-range connectivity remodeling. Finally, strategies
are proposed to merge the two pipelines. We further suggest that additional models could be
integrated into the framework thanks to the versatility of the proposed approach, thus allowing
many researchers to achieve continuously improved experimental design.

1 INTRODUCTION

In nature, the activity of the brain of an individual interacting with the environment is conditioned by the
response of the environment itself, in that the output of the brain is relevant only if it has the ability to
impact the future and hence the input the brain receives. This "closed-loop" can be simulated in a virtual
world, where simulated experiments reproduce actions (output from the brain) that have consequences
(future input to the brain) (Zrenner et al., 2016). To the aim of reproducing *in silico* the complexity of real
experiments, different levels of modeling shall be integrated. However, since modeling all components of an
experiment is very difficult, traditional approaches of computational neuroscience reduce the experimental
   setups as much as possible. An "Embodied brain" (or "task dynamics", see Zrenner et al. (2016)) approach

Figure 1. Scheme of the proposed *Embodied brain* framework . The picture suggests a closed-loop
workflow linking real and simulated experiment. The different types of data obtained from the experiments,
from brain activity to dynamic and kinematics of goal-directed movement, are used to feed the whole brain
and spinal cord model, in addition to the virtual mouse and environment. The loop is closed by validation of
in silico results on real data. Eventually, the simulated experiment raises novel hypotheses, to be validated
on new real experiments.

could overcome these limits by associating the modelled brain activity with the generation of behavior
within a virtual or real environment, i.e. an entailment between an output of the brain and a feedback
signal into the brain (Tessadori et al., 2012; DeMarse et al., 2001; Reger et al., 2000). The experimenter
can interfere with the flow of information between the neural system and environment on the one hand
and the state and transition dynamics of the environment on the other. Closing the loop can be performed
effectively by (i) validating the models on experimental data, and (ii) designing new experiments based on
the hypotheses formulated by the simulations. On the example shown in Fig. 1, data on brain activity (be
it, for instance, from electrophysiological recordings or imaging) and on the environment (e.g. by means
of kinematic or dynamic measures) from the real experiment are used to feed the models of the *in silico*
representation of the experiment. From a comparison of the real and model-based data, the features that are
most important to replicate the real experiment are identified, and thus novel insights are generated (Fig. 1).
To realize such a complex virtual system, many choices can be made, for instance on the brain model or
spinal cord model that best represent the salient features of experimental measures to be replicated. The
ideal framework shall comprise a library of tools to choose from, to reproduce a variety of experimental
paradigms in the virtual environment. By briefly introducing the state of the art in brain and spinal cord
modeling, we will discuss few classes of models to pick from an ideal library.
41

1.1   State of the art

1.1.1   Local cortical network modeling

Biologically detailed models of a single neuron such as the Hodgkin and Huxley model (Hodgkin and
Huxley, 1952) take into account the activity of the ion channels in the cell membrane that lead to changes
in the membrane potential, eventually causing the neuron to spike. However, simpler but still biologically
realistic models of the single cell are preferable when interested in modeling the dynamics of a larger
number of cells. A good candidate is the adaptive exponential integrate and fire (adex) neuron model (Brette
and Gerstner, 2005), which has been shown to reproduce the intrinsic neuronal properties of a number of
cell types, including those with spike frequency adaptation (Destexhe, 2009). Interestingly, adex neurons
network models with different adaptation levels can reproduce the dynamical properties of distinctive brain
states such as wakefulness, sleep or anesthesia (Zerlaut and Destexhe, 2017; Nghiem et al., 2020). This
property makes adex neuron networks suitable to model the dynamics of the emerging activity in a local
network of neurons after injury, given that, after a stroke, the dynamics of the local network switches
to a slow oscillatory rhythm which resembles that of sleep or anesthesia Butz et al. (2004). Moreover,
alterations in low-frequency cortical activity in the peri-infact cortex after stroke are known to correlate
with motor recovery (e.g. Yilmaz et al. (2015); Ramanathan et al. (2018))
58

1.1.2   Brain Network Modeling

Efforts have been made to reconstruct single brain regions with as many details as possible, (Markram
et al., 2015), or to build detailed networks of multi-compartment oscillators (Izhikevich and Edelman,
2008). Contrary to the detailed models, top-down modelling seeks to elucidate whole-brain network
mechanisms, which may underpin a variety of apparently diverse neurophysiological phenomena. Neural
masses formalisms have been used over many years to develop macroscopic models that capture the
collective dynamics of large neural assemblies (Deco et al., 2008; Sanz-Leon et al., 2015). In this case
the activity of a macroscopic brain region is often directly derived from populations of spiking neurons
as a mean-field using concepts from statistical physics, e.g. (Stefanescu and Jirsa, 2008; Wong and
Wang, 2006; Zerlaut et al., 2017). In other cases the statistics of the macroscopic brain activity is derived

more phenomenologically while still conserving some basic physiological principles such as division on
excitatory and inhibitory neurons, e.g. the seminal Wilson Cowan model (Wilson and Cowan, 1972). The
third subclass of neural masses contains purely phenomenologically derived computational models that
aim to reproduce certain dynamical properties of the macroscopic neuronal activity, such as e.g. seizure
dynamics (Jirsa et al., 2014; Saggio et al., 2017), whilst different realizations of damped or self-sustained
oscillators are often used to model the coherent fluctuations of resting-state activity (Cabral et al., 2011;
Deco et al., 2016). Depending on the working point of the system, the macroscopic dynamics can be
described not only by physiologically derived mean-fields, but also by phenomenological models in their
canonical form (Deco et al., 2009, 2011; Izhikevich, 1998). Hence, phase oscillators (Kuramoto, 1984) are
often chosen to model and study the coactivation patterns in the brain, as some kind of a minimal model
explanation (Batterman and Rice, 2014) for the synchronized behaviour over a network (Pikovsky et al.,
2001; Breakspear et al., 2010).

Connecting the neural masses in large-scale brain network models (BNM) became possible with the
progress of non-invasive structural brain imaging (Johansen-Berg and Rushworth, 2009). This allowed
extraction of biologically realistic brain connectivity, the so-called connectome, which shapes the local
neuronal activity to the emergent network dynamics (Honey et al., 2007; Ghosh et al., 2008; Deco et al.,
2009; Sanz-Leon et al., 2015; Petkoski et al., 2018; Petkoski and Jirsa, 2019).

The large-scale BNM have been used to interpret healthy (Cabral et al., 2011; Deco et al., 2016) or
pathological (Nakagawa et al., 2013; Saenger et al., 2017; Zimmermann et al., 2016) brain activity. This
is often reflected in the coherence between brain rhythms (Lachaux et al., 1999) that also describes the
functional connectivity (FC) of the brain as an important marker of its spatio-temporal organization (Deco
et al., 2009, 2011; Ghosh et al., 2008; Deco and Jirsa, 2012; Petkoski et al., 2018).

The Virtual Brain (TVB) (Sanz Leon et al., 2013; Sanz-Leon et al., 2015) is a commonly used
neuroinformatics platform for full brain simulations. It supports a systematic exploration of the underlying
components of a large-scale BNM: the structural connectivity (SC) and the local dynamics that depend on
the neurophysiological mechanisms or phenomena being studied. In this way, the BNM allows to describe
structural changes (through connectivity variation including stroke, motor learning and recovery) and
subsequent functional consequences accessible to modeling and empirical data collection on the meso,
macro and behavioral level. The modelling with TVB thus represents a useful paradigm for multi-scale
integration. TVB has been already utilized in modeling functional mechanism of recovery after stroke in
humans (Falcon et al., 2015, 2016), identifying that the post-stroke brain favors excitation-over-inhibition
and local-over-global dynamics. For studying the changes in synchronization, as we intend to do, TVB
offers a range of oscillatory models for the neural activity. One of these is the Kuramoto model (KM),
which captures the emergent behaviour of a large class of oscillators that are near an Andronov-Hopf
bifurcation (Kuramoto, 1984), including some population rate models Ton et al. (2014). This makes
the KM well suited for assessing how the connectome governs the ynchronization between distant brain
regions (Breakspear et al., 2010; Cabral et al., 2011, 2012; Ponce-Alvarez et al., 2015; Petkoski et al., 2018).

1.1.3   Spinal cord modeling

The brain controls its body through neural signals originating from the brain and processed by the spinal
cord to control muscle activation in order to perform a large variety of behaviors. Several biologically
realistic functional models of the spinal cord have been developed and tested in closed loop simulations with
musculoskeletal embodiments. Stienen and colleagues (Stienen et al., 2007) developed a fairly complete

model that includes Ia, Ib and II sensory afferents, both monosynaptic and polysynaptic reflexes as well
as Renshaw cells, improving a previous work by Bashor (Bashor, 1998). The model was tested with a
musculoskeletal model consisting of a generic antagonistic couple of muscles, thus lacking a realistic
validation scenario. Cisi and Kohn developed a web-based framework for the simulation of generic spinal
cord circuits with associated muscles, that aims at replicating realistic experimental conditions (i.e. electrical
stimulation) (Cisi and Kohn, 2008). Sreenivasa and colleagues developed a specific neuro-musculoskeletal
system, upper limb with biceps and triceps, and validated it against human recordings (Sreenivasa et al.,
2016). In (Moraud et al., 2016), a simple spinal cord model of the rat, lacking any descending stimuli, was
developed in order to study how such circuitry can correct the gait after a spinal cord injury and embedded
in a closed loop simulation with biomechanical hindlimbs. All of the mentioned works were tested primary
for the generation of reflex motions, and not as intermediate levels of more complex controllers such as
ones capable of generating voluntary movements.

1.2 Aim of the work

We propose a framework ("*Embodied brain closed loop*") endowed with a library of modeling tools
that will eventually allow to realize entirely virtual experiments. We focused on an experiment on motor
control and motor recovery after stroke described in Spalletti et al. (2017) and Allegra Mascaro et al.
(2019), whose simulation requires two main tiles. The first is the realization of voluntary movements in a
virtual milieu. This piece requires monitoring and modeling of many components of movement control,
from brain activity to body kinematics and displacement of virtual objects. The second is the simulation
of brain injury. This includes modeling of acute consequences but also of neuronal plasticity after brain
damage, either spontaneous or supported by treatment. Both local and long-range modulation of neuronal
activity should be accounted to simulate the brain after stroke, since local alteration of neuronal activity
in the peri-infarct area is known to be associated to remodeling of long-range functional and structural
connectivity (several comprehensive reviews have summarized this research, e.g. Carmichael et al. (2017)).
To build those tiles, we developed two pipelines that target, on one side, the physiological execution of
movements and, on the other, pathological alterations and plasticity (Fig. 2). The first ("Movement-driven
models" pipeline) aims at reproducing in a virtual environment how a goal-directed movement is performed
and represented in the healthy brain. Data recorded on healthy mice are used as an input to the spinal
cord model, attached to the muscles of the simulated embodiment (see Fig. 2, red box). The goal of the
second pipeline ("Stroke models") is to reproduce both local and long-range consequences of stroke. We
developed a spiking neurons model that could simulate the local brain dynamics, and in particular the
abnormal oscillatory activity taking place in the peri-infarct cortex (see Fig. 2, lower line in the green box).
Also, we show how the simulation of brain activity by neural mass models allows replicating the evolution
of functional connectivity in mouse brain after a stroke and under rehabilitation (see Fig. 2, upper line in
the green box).

2 METHODS

149 Cortical recordings and behavioral data from the experiments described in this section are used to build
150 and validate the brain models and the output in the virtual environment.

2.1 *in vivo* experiments

152 On the experimental side, we performed electrophysiological recordings (Fig. 3, panel A) and wide-field
153 calcium imaging (Fig. 3, panel B) in awake mice performing active forelimb retraction on a robotic device
154 (M-Platform). These experiments allowed gathering simultaneous information on the neuronal activity,
155 force applied during active forelimb retraction and position of the forelimb, as displayed in the lower

Figure 2. Scheme of data and simulations. The scheme depicts the approach to build the *Embodied brain* framework from data to models and back. The workflow from data to models and simulation in the *Embodied brain closed loops* is shown. The upper, green box shows the *Stroke models* closed loop, the lower, red box shows the *Movement-driven models* closed loop. Colored images represent experiment data, brain and spinal cord models, and simulation of the environment (from left to right). Connections between the modeling components are presented as arrows: solid lines represent the output provided to other blocks; dashed lines indicate the output data of the models that are used for comparison with real data for validation.

panels of figure 3. The electrophysiological data and the recordings of limb position were used to feed the
spinal cord model, as described in section 4.1. The features of the wide-field calcium data recordings were
used to build the spiking neurons brain model and to validate the BNM, Section 3.3. All the procedures
were in accordance with the Italian Ministry of Health for care and maintenance of laboratory animals
(law 116/92) and in compliance with the European Communities Council Directive n. 2010/63/EU, under
authorizations n. 183/2016-PR (imaging experiments) and n. 753/2015-PR (electrophysiology experiments).
162
2.1.1   Robotic training on the M-Platform

The M-Platform is a robotic device designed to train mice to perform active forelimb retraction (Pasquini
et al., 2018). Briefly, the main component of the device is a linear actuator that moves a linear slide where
a custom handle is screwed. Moreover, the platform is provided with a system to control the friction on
the slide and a pump for the reward. During the experiments, while the mouse has its left paw connected
to the slide, first the linear actuator extends the forelimb then the animal has to perform an active pulling
movement to come back to the starting point and to receive a reward. Force signal and position of the
forelimb are recorded respectively by a load cell and a webcam.

In Section 2.1.3 and 2.1.4 we describe two different experiments with the robotic device. In the first one,
the M-Platform is embedded with Omniplex D System (Plexon, Usa) to obtain *in vivo* electrophysiological

Figure 3. The real robotic platform. (A) On the top a schematic representation of the experiment during electrophysiological recording in CFA. At the bottom the synchronized data: force peak (blue), movement of the slide (red), high frequency electrophysiological signal of a single channel (magenta) and the timestamp of a selected single unit. (B) On the top a scheme of the experiment with the setup to record calcium activity. At the bottom the recorded data after synchronization: force peak (blue), movement of the slide (red) and calcium response (green).

173   recording during the task. In the second one, the kinetic and kinematic parameters are synchronized with
174   wide-field calcium imaging recordings (Fig. 3).

175
176   ## 2.1.2 Photothrombotic Stroke

177   To induce focal stroke in the right hemisphere, mice were injected with Rose Bengal (0.2 ml, 10 mg/ml
178   solution in Phosphate Buffer Saline). Five minutes after intraperitoneal injection, a white light from an
179   LED lamp was focused with a 20X objective and used to illuminate the primary motor cortex (0.5 mm
180   anterior and 1.75 mm lateral from Bregma) for 15 min.

181
2.1.3 Electrophysiological recordings on the M-Platform

Two healthy mice were used for the experiments. Animals were housed on a 12h/12h light/dark cycle.
Mice were water deprived overnight before training on the platform; daily liquid supplement was given after
the test. Food was available ad libitum. To have access to the motor cortex, a craniotomy was performed 3
days before the training to expose the Caudal Forelimb Area (CFA) of the right hemisphere.The craniotomy
was filled with agarose and silicon (Kwik cast sealant, WPI) and could be opened and closed several times
for acute recordings."

Mice were gradually acclimated to the platform. Then they performed the task for two days, fifteen
trials each day. During the pulling experiment, mice were head fixed to the platform with their left wrist
constrained to the slide. The friction on the slide was set at 0.3 N. The force signal was acquired by a load

cell (Futek LSB200, CA, USA) along the direction of the movement at 100 Hz, at the same time a webcam
recorded the position of the slide at 25 Hz and the multi-unit activity was recorded by Omniplex D System
(Plexon, USA) with a frequency of 40 kHz thanks to a 16 channels linear probe (1 $M\Omega$, ATLAS, Belgium)
inserted into the CFA at 850 μm of depth. (Fig. 3A).

2.1.4　Wide-field calcium imaging of cortical activity during training on the M-Platform

The mouse was housed in clear plastic cage under a 12 h light/dark cycle and was given ad libitum access
to water and food. We used the following mouse line from Jackson Laboratories (Bar Harbor, Maine USA):
C57BL/6J-Tg(Thy1GCaMP6f)GP5.17Dkim/J (referred to as GCaMP6f mice). In this mouse model, the
fluorescence indicator GCaMP6f is mainly expressed in excitatory neurons H. Dana (2014). GCaMP6f
protein is ultra-sensitive to calcium ions concentration T.W. Chen and Kim (2013); H. Dana (2014) whose
increase is associated with neuronal firing activity Grienberger and Konnerth (2012); R. Yasuda and
Svoboda (2004).

For wide-field fluorescence imaging of GCaMP6f fluorescence, we used a custom made microscope
described in Conti et al. (2019). Briefly, the system is composed by a 505 nm LED (M505L3 Thorlabs, New
Jersey, United States) light deflected by a dichroic filter (DC FF 495-DI02 Semrock, Rochester, New York
USA) on the objective (2.5x EC Plan Neofluar, NA 0.085, Carl Zeiss Microscopy, Oberkochen, Germany).
The fluorescence signal is selected by a band pass filter (525/50 Semrock, Rochester, New York USA) and
collected on the sensor of a high-speed complementary metal-oxide semiconductor (CMOS) camera (Orca
Flash 4.0 Hamamatsu Photonics, NJ, USA).

The experiment starts with a mouse being trained and recorded for one week (five days) on the M-platform
("healthy" condition, see Fig. 3). The focal stroke is then induced at the beginning of the second week
by phototrombosis on the right primary motor cortex (rM1). Starting 26 days after stroke, the mouse
performance and spontaneous motor remapping was evaluated on the M-Platform for 5 days a week along
4 more weeks. The results from the first week one month after the injury is the so-called "stroke" condition,
while the results during the last week, when the animal recovers the motor function is referred to as "rehab".

Each day, the beginning of the wide-field imaging session was triggered by the start of the training
session on the M-Platform. To detect the movement of the wrist of the animal in the low-light condition of
the experiment, an infrared (IR) emitter was placed on the linear slide, and rigidly connected to the load
cell and thus to the animal's wrist. Slide displacement was recorded by an IR camera (EXIS WEBCAM
#17003, Trust) that was placed perpendicular to the antero-posterior axis of the movement. Position and
speed signals were subsequently extracted from the video recordings and synchronized with the force
signals recorded by the load cell (sampling frequency = 100 Hz) and with the fluorescence signal recorded
by the CMOS sensor (Fig. 3B).

2.2　Data analysis

2.2.1　Spikes and force analysis

Data were analyzed offline using custom routines in Matlab (MathWorks). First, the position signal was
extracted by the video using a white squared marker on the slide as reference. The recording frequency
of the video was 25 Hz. After applying an antialiasing FIR lowpass filter, a uniform linear resample of
the movement of the slide was performed, in order to synchronize the position signal with the force data,
recorded at 100 Hz. To identify the timing of the voluntary activity of the animal, a threshold method was
used to detect force peaks during the pulling phase of the task. For the following analysis, we picked out
peaks that produced a displacement of the slide, in addition to crossing the threshold; and we calculate the

onset of these peaks as the minimum of the force derivative just before the respective peak (Spalletti et al., 2014). The electrophysiological signal, recorded at 40 kHz as sampling rate, was analysed by Offline Sorter (Plexon, Dallas, TX). First, for each channel of the probe, we sorted waveform that crossed a detection threshold of the mean $\pm$ 3 standard deviations. Then, detected spikes were clustered using an automatic process based on principal component analysis. Starting from these clusters, a manual sorting was executed to isolate all single units which could be identified in the recorded multi-units signal. The time stamp of each unit was synchronized with the data of the robot. To evaluate the temporal behavior-related spike activity, the peri-stimulus time histograms (PSTHs, NeuroExplorer, Plexon) was generated with bins of 20 ms in an interval of 1 s around the onset of force peaks. In addition, the resting activity of each unit was evaluated selecting intervals of at least 0.6 s with no force peaks and calculating the average of the number of spikes in bins of 20 ms. Finally, the PSTHs was used to evaluated when a single neuron was active, that is when the number of spikes for bin cross the threshold, calculated as the mean $\pm$ 2 standard deviations of the number of spikes for bin during the respective resting activity.

2.2.2   Phase coherence and Functional connectivity

Functional connectivity (FC) among cortical regions was inferred from phase coherence of activity measurements, and used to determine changes in brain activity in "stroke" and "rehab" condition, as compared to the healthy mice. These inferred activity changes were used to parameterize simulations of the BNM built over the Allen Brain Atlas mouse connectivity data (http://connectivity.brain-map.org/; (Oh et al., 2014), below referred to as the Allen Mouse Brain Atlas - AMBA), incorporated in the extended virtual mouse brain (Melozzi et al., 2017)In each animal, the camera field-of-view used for activity measurements was placed in a standard position using the sagittal suture and its intersection with the coronal suture of the skull (bregma) as anatomical landmarks. To spatially correlate our activity measures with the structural connectivity data (Oh et al., 2014), the camera field-of-view (Fig. 3, panel B) was spatially translated to the Allen Common Coordinate Framework (CCF, v3, 2015; (Oh et al., 2014)). Since the CCF lacks stereotactic skull landmarks, these were introduced by spatially co-registering all diagrams from a standard stereotaxic mouse brain atlas (Franklin et al., 2008) to the CCF coordinate space with affine transformations defined using the QuickNii tool (Puchades et al., 2019). Using bregma and the sagittal suture as a reference, the four corners of the downsampled 128x128 pixels field-of-view of the recorded images were positioned in CCF, taking the 5 degree lateral tilt of the camera view into account. Delineations of layer IV cortical regions were then projected onto the camera field-of-view, and used as a custom atlas reference for all activity maps.

The spectral content of the signals is analyzed to identify the frequency band which captures the spontaneous brain activity that occurs simultaneously with the motor-evoked events. Even though the slowest dynamics $< 0.5 Hz$, which has the highest power, is often a marker of the resting state (Wright et al., 2017), in this experiment it also contains the propagation of waves generated during the limb movements on the platform. The mechanisms behind stimulation propagation (Spiegler et al., 2016) are different from the spontaneous oscillations at rest (Deco and Jirsa, 2012) that we try to study and model here, and hence the lowest frequencies are excluded from the analysis. In addition, the mice heart rate is between $6 - 8 Hz$, whilst the activity above $10 Hz$ is too close to the Nyquist frequency of $12.5 Hz$, defined as half of the sampling rate of the recordings. As a consequence these bands are generally avoided in the analysis of calcium signals, which is consequently often centered at the δ band between around 1 Hz and 5 Hz, (Vanni et al., 2017; Wright et al., 2017).

279   The FC is characterized with the phase coherence of the analytical phases of the band-passed time-series
280   obtained using the Hilbert transform (Pikovsky et al., 2001). For this we employ phase locking values
281   (PLV) (Lachaux et al., 1999) that are a statistical measure for similarity between the phases of two signals,
282   hence defined as

$$PLV_{ij} = |\frac{1}{M} \sum_{m=1}^{M} \mathrm{e}^{i(\theta_i(m) - \theta_j(m))}|, \tag{1}$$

283   where the phase difference $\theta_i(m) - \theta_j(m)$ between the regions i and j is calculated at times $m = 1 \ldots M$.
284   The same procedure is also applied to surrogate time-series to find the level of statistically significant phase
285   coherence (Lancaster et al., 2018).

3 MODELS

3.1 Spinal cord model

287   To develop the final model, an incremental approach was followed, starting from a circuit for a single
288   *muscle*, adding inhibitory connections between *antagonistic* pairs and finally interneurons to modulate
      descending stimuli (Fig. 4). For a single muscle, a network with muscle spindles providing Ia and II afferent

Figure 4. The spinal cord model for a pair of antagonistic muscles.

289
fibers activity, a pool of α-motoneurons and excitatory II-interneurons was considered (Stienen et al., 2007;
Moraud et al., 2016). Ia afferents directly provide excitatory inputs to the α-motoneurons (monosynaptic
stretch reflex mechanism), while the II afferents output is mediated by a set of interneurons before reaching
the α-motoneurons, creating a disynaptic reflex. The muscle spindles are implemented using the model
from (Vannucci et al., 2017). All other neurons are modeled as leaky integrate and fire neurons. The number

of neurons in the spinal cord populations, as well as parameters for the synaptic connections are taken from
(Moraud et al., 2016), with the exception of the synaptic weights of the monosynaptic connections, which
have been significantly lowered (see Supplementary Material). The parameters from the muscle spindle
models are taken from (Mileusnic et al., 2006), which are tuned on neurophisiological recordings of lower
mammals. Distribution of parameters for the α-motoneurons that influence the recruitment order and fiber
strength (membrane capacitance, membrane time constant, maximum twitch force, time to peak force) are
taken from (Sreenivasa et al., 2016):

$$
\begin{align}
D_i &= [d_{max} - d_{min} \cdot \log(N - i)] \cdot D_{SF} \tag{2}\\
C_i &= \pi D_i^2 \cdot c_{spf} \tag{3}\\
\tau_i &= \tau_{max} - (D_i - \tau_{adj}) \cdot \tau_{slp}\\
F_i &= [p_{max} - p_{min} \cdot \log(N - i)] \cdot F_{SF}\\
T_i &= \left[s_{min} - \frac{s_{sl}}{N} i \right] \cdot T_{SF} + s_{min}
\end{align}
$$

where i is the index of the α-motoneuron in the pool, N is the size of the pool and the others are free
parameters that can be adjusted for every muscle. In this work, the value of these parameters has not been
changed from (Sreenivasa et al., 2016).

In order to compute the actual muscle activation from the motoneurons activity, a special spike integration
unit that sums the fibers twitches was implemented. The spikes were integrated using the discrete time
equations of (Cisi and Kohn, 2008) with a non-linear scaling factor from (Fuglevand et al., 1993) that
prevents the activation to grow indefinitely:

$$
a_i(t) = 2e^{\frac{-\delta t}{T_i}} \cdot a_i(t - 1) - e^{\frac{-2\delta t}{T_i}} \cdot a_i(t - 2) + F_i \cdot g(t) \cdot \frac{\delta t^2}{T_i} e^{\frac{1 - \delta t}{T_i}} \cdot u(t) \tag{4}
$$

309 where δt is the integration time, and $u(t)$ and $g(t)$ are the spike function and the non-linear scaling, defined
310 as:

$$
u(t) = \begin{cases} 1 & \text{if a spike is received at } t \\ 0 & \text{if no spikes are received at } t \end{cases} \tag{5}
$$

$$
g(t) = \begin{cases} 1 & \text{if } T_i/ISI_i < 0.4 \\ \frac{1 - e^{-2(T_i/ISI_I)^3}}{T_i/ISI_I} & \text{otherwise} \end{cases} \tag{6}
$$

311 where ISI_i is the observed inter-spike interval of α-motoneuron i. Moreover, the activation can be scaled
312 between 0 and 1 by dividing by the maximum theoretical value:

$$
a_{i,max} = \lim_{\substack{t \to +\infty \\ ISI_i \to 0}} a_i(t) = F_i \frac{\frac{\delta t^3}{T_i^2} \left(1 - e^{-2\left(\frac{T_i}{\delta t}\right)^3} \right) \cdot e^{\left(1 - \frac{\delta t}{T_i}\right)}}{1 - 2e^{-\frac{\delta t}{T_i}} + e^{-2\frac{\delta t}{T_i}}} \tag{7}
$$

Therefore, the output of the twitch integration module is an activation value in $[0; 1]$ that is suitable for
the muscle model present on the mouse virtual embodiment. The effect of this integration is that at low
frequencies the individual twitches can still be seen, while at higher stimulation the twitches fuse into a

tetanic contraction. Moreover, thanks to the non-linear scaling, the activation reaches a maximum value
and higher stimulation frequencies do not produce any effect, in accordance with the contractile properties
of real muscle fibers.
In order to implement the polysynaptic inhibition reflex between antagonistic muscles, two populations
of Ia-interneurons were added to the network. Those receive inputs from all Ia afferents of a synergistic
muscle and provide inhibition to the α-motoneurons of the corresponding antagonistic muscle. Moreover,
as the activation of a muscle should provoke an inhibition of its antagonist (Pierrot-Deseilligny and
Burke, 2005), the Ia-interneurons also receive low-gain positive inputs from the corresponding descending
pathways. Again, the number of neurons in these population and their parameters have been taken from
Moraud et al. (2016). Finally, as there is lack of evidence for a direct connection between cortical neurons
and motoneurons in the spinal cord of rodents (Yang and Lemon, 2003), an intermediate population
of neurons mediating descending signals was added to the circuitry. This population aims at modelling
propriospinal neurons, which provide an inhibitory action on the signals coming from the corticospinal
tract (Alstermark, 1992). In general, the inhibition is generated from different peripheral afferents, but we
included only afferents from muscle spindles as these are the only present in the model. As there is no
definitive experimental evidence on the size of the population of propriospinal neurons and its parameters,
we set the values of these to those of the populations of Ia-interneurons. Conversely, the synaptic weights
and the number of connections between the descending inputs and the propriospinal interneurons were
empirically tuned starting from experimental data.
335

3.2   Simulation tools and physical models

337   This section describes simulation tools that have been used to synchronize neural and physical simulations
338   and the physical simulations models that have been developed and used. These tools and model were used
339   in the context of the *Movement-driven models* pipeline.
340

3.2.1   Embodied mouse in the Neurorobotics Platform

The full musculoskeletal model of the virtual rodent controlled by the spinal cord model was simulated in
the Neurorobotics Platform (NRP) developed in the Human Brain Project (Falotico et al., 2017). The main
components of the NRP are a world simulator, a brain simulator and the mechanism that enables the data
flow between the two in a closed-loop. The connection between the body and the brain is specified through
a domain specific language (Hinkel et al., 2015, 2017), via Python scripts called Transfer Functions. In
these scripts the output of devices that read neuronal output data can be processed and passed as input
for the virtual body actuators, and vice versa, the sensory information from the virtual body sensors, in
this case muscle length data, can be passed to devices that map sensory data to neural input. The brain
simulation, which currently is simulating point-neurons, follows closely the paradigm of NEST (Gewaltig
and Diesmann, 2007), interfaced through PyNN (Davison et al., 2009). On the other side of the closed
loop the world simulator of choice is Gazebo (Koenig and Howard, 2004), extended to support muscle
simulation through OpenSim (Millard et al., 2013a), which provides its' own muscle simulation engine.
354

3.2.2   Musculoskeletal embodiment

As described earlier the musculoskeletal system comprises of two elements, the skeletal and the muscle
system respectively. Here both systems are elaborated a bit more in the context of the experiment.
Developing animal skeletal systems is no trivial task. It involves many complex degrees of freedom
and physical properties such as mass, center of mass and inertias. To ease this process, NRP has developed

a toolkit for Blender (Open source modeling and animation tool) called RobotDesigner (HBPNeurorobotics,
2019). RobotDesigner allows to automate several steps needed to develop skeletal/robot models to be
simulated in the NRP. Using the same, currently NRP hosts state-of-the-art a full skeletal model of the
mouse consisting of 110 degrees of freedom. More details about the full model will be soon published
following the current article. For the current experiment, the mouse skeletal model is reduced in complexity
by constraining all the degrees of freedom except the left forelimb. The forelimb consists of four segments
and it is further constrained to only have flexion-extension movements, enough to reproduce the passive
extension-active retraction experiment on the M-Platform. The different segments and the joints of the
forelimb are shown in Fig. 5.

Figure 5. Mouse forearm musculoskeletal system (left panel) Forelimb skeletal system with three joints
(1) shoulder (Ball and socket) (2) elbow (hinge) (3) wrist (hinge). (right panel) Forelimb muscle system
with six muscles (1) humerus-extension (2) humerus-flexion (1) elbow-extension (2) elbow-flexion (1)
hand-extension (2) hand-flexion

The physical properties of the skeletal system such as mass, center of mass and inertia are automatically
estimated based on bounding objects generated for each link (segment) using the RobotDesigner. Once the
skeletal system is established, muscles-tendon system can be attached to the bones. As mentioned before,
NRP now supports OpenSim for integrating muscle models into physical animal bodies or even robots. In
the current experiment a pair of antagonist hill-type muscles were added to each of the joints in the mouse
forelimb. The muscle model in OpenSim is taken from Millard and colleagues (Millard et al., 2013b) (see
Supplementary Material). Again RobotDesigner offers a unique solution to visualize attachments and easily
add muscles to the body in blender. Using the same technique all the muscles for the mouse forelimb were
added. Muscle parameters used in the current experiment are hand tuned to produce flexion-extension
movements necessary for the experiment. Fig. 5 (right panel) shows the muscle attachments used in the
current model.

3.2.3   Robotic rehabilitation platform model

In the real experiment, the mouse forelimb is attached to the sliding mechanism, which is a prismatic
joint, driven by a DC motor whose rotational motion is converted into a linear one. The motor that is
controlled with a PID controller, whose reference can be set to a position of the joint between the minimum
and maximum positions. The controller is enabled when the operator decides to replace the sled in its
starting position and is disabled afterwards, so that the mouse can actually pull the sled. In simulation,
the same configuration has been implemented. The musculoskeletal mouse forelimb was attached to a

simulated M-platform, which has been modelled as a prismatic joint, actuated by a PID controller that can
directly apply a simulated force on the joint. Again, the reference to the PID controller is the position of
the prismatic joint in its range, this time normalized between 0 (minimum) and 1 (maximum). To simulate
the intervention of the operator that puts the slide back we employed a state machine that automatically
controls the slide, by making use of the PID controller setting 1 as a reference. Inputs to this state machine
are a list of times at which the slide should be put back. Conversely, to simulate the minimum amount of
force that is required to move the slide in the real setup, we deactivated the PID controller in simulation
only after a certain activation of the simulated muscles was reached (0.95).

3.3 Stroke Models

3.3.1 Brain Network Model with Kuramoto oscillators

To simulate the functional network reorganization during stroke and recovery given by the phase coherence
of the macroscopic brain activity reflected in calcium signals, we built our BNM based on Kuramoto
oscillators for the local oscillatory dynamics and the AMBA connectome that dictates the strength of the
couplings between brain regions (Melozzi et al., 2017; Choi and Mihalas, 2019), Fig. 6, and has been
validated with empirical functional data that justifies its use (Melozzi et al., 2019). The AMBA contains 86
cortical regions (43 per hemisphere), of which 18 were included in the field-of-view (Fig. 6, bottom left).
The average calcium signal of the pixels entirely located within a brain region was used to represent their
mean neuronal activity.

Besides their simplicity, phase models exhibit rich dynamics and a direct link to more complex biophysical
models, while admitting analytic approaches (Roy et al., 2011; Sheppard et al., 2013; Ton et al., 2014;
Stankovski et al., 2016). KM (Kuramoto, 1984), as a phenomenological model for emergent group dynamics
of weakly coupled oscillators (Pikovsky et al., 2001) is well suited for assessing how the connectome
governs the brain oscillatory dynamics that can be reflected in different neuroimaging modalities (Schmidt
et al., 2014; Váša et al., 2015; Cabral et al., 2017; Petkoski et al., 2018). The constructed BNM is thus used
to identify the structural alterations due to the stroke and the subsequent recovery, using their causal effects
on the functional changes captured by the calcium recordings of the cortical brain activity.

Even though delayed interactions due to axonal transmission can be of crucial importance for the observed
dynamics of the oscillatory systems (Ghosh et al., 2008; Petkoski et al., 2016, 2018), the impact of these
delays is much less pronounced for low frequencies compared with them, as it is the case here. Moreover,
the tracing used for obtaining the AMBA Connectome (Oh et al., 2014) does not allow tracking the length
of the white fibers. Hence, we assume instantaneous couplings and the utilized model gives the following
evolution of the phases for each of the N brain regions

$$\dot{\theta}_i = 2\pi f + \frac{1}{N}\sum_{j=1}^{N} K_{ij}\sin(\theta_j - \theta_i) + \eta_i(t), \ \ i = 1\ldots N. \tag{8}$$

Here the dynamics of each region i is driven by the natural frequencies f that are assumed to be identical
across the brain. A stochastic variability is introduced with additive Gaussian noise defined as $\langle\eta_i(t)\rangle = 0$
and $\langle\eta_i(t)\eta_j(t')\rangle = 2D\delta(t - t')\delta_{i,j}$, where D is the noise strength and $\langle\cdot\rangle$ denotes time-averaging. The
activity of the BNM is then constrained by the structural connectivity, which for every region is represented
by the inputs that they receive from the other regions j through the coupling strength $K_{ij} = Kw_{ij}$. This
contains the structural weight of the connectome between these areas, w_{ij}, scaled with the same global

Figure 6. Scheme of the mouse BNM. Brain Network Model consisting of neural masses superimposed over the AMBA connectome simulates the recorded calcium activity. The average oscillatory neuronal activity of the brain regions is described by Kuramoto oscillators, which are coupled due to the fiber tracts, giving rise to the simulated recordings. The brain network (right) is reconstructed from the AMBA, with the centers of subcortical regions being small black dots, while larger the circles are for the cortical regions, with the region of the stroke highlighted. On the left, the field of view during the recordings is overlayed on the reconstructed brain, and different colors represent the cortical regions according to the AMBA.

coupling K for every link.

3.3.2 Spiking network model for simulation of slow wave activity in peri-infarct cortex

Besides the phenomenological neural mass model for the oscillatory activity that we have used in the
BNM, we also show an alternative spiking neural model to reproduce local brain activity in the acute phase
after stroke. In future, this model should be integrated in the BNM and therefore in the *Embodied brain*
closed-loop simulation, either by deriving its mean-field representation, e.g. see (Zerlaut et al., 2017), or by
co-simulation. This model aims to reproduce the two-photon calcium signals of a population of spiking
neurons located at the peri-infarct area, since it is known that slow frequency patterns of synchronized
activity emerge from the damaged areas after an ischemic stroke (Butz et al., 2004; Carmichael and
Chesselet, 2002; Rijsdijk et al., 2008; Rabiller et al., 2015).

Network of adaptive exponential integrate and fire (adex) neurons:

The network consists of an excitatory (regular spiking, RS) and and inhibitory (fast spiking, FS) population
of neurons (Fig. 7A). All cells are modeled as adex neurons, which can be described by the following
equations:

$$
\begin{cases}
C_m \frac{dV(t)}{dt} = G_l(E_l - V(t)) + G_l \Delta_V e^{\left(\frac{V(t)-V_{thre}}{\Delta_V}\right)} + I_{syn}(t, V(t)) - w(t) + \sigma\xi(t) \\
\frac{dw(t)}{dt} = -\frac{G_l}{C_m} w(t) + b \sum_k \delta(t-t_k) + a(V(t) - E_l)
\end{cases}
\tag{9}
$$

443 where the synaptic input I_{syn} is defined as

$$
I_{syn}(t, V(t)) = \sum_i g_i^{syn}(t)(V(t) - E_i^{syn})
\tag{10}
$$

Figure 7. Two-photon calcium signal model from a spiking network model. (A) Schematic connectivity between the excitatory population of regular spiking (RS) neurons and the inhibitory population of fast spiking (FS) neurons of the modelled cortical network. (B) Activation curve of a high-voltage activated calcium channel that is used to compute the inward calcium current (I_{Ca}) from the changes in the V_m. (C) From top to bottom, simulated membrane potential of a neuron emitting three spikes which are represented by dashed lines, membrane potential with reconstructed spikes, inward calcium current associated with changes in the membrane potential, cytosolic calcium concentration and fluorescence emitted by the calcium indicator due to the intracellular concentration of calcium.

444   with

$$\frac{dg_i^{syn}(t)}{dt} = -g_i^{syn}(t)/\tau_{syn} \tag{11}$$

Here, $Gl = 10\ nS$ is the leak conductance and $Cm = 150\ pF$ is the membrane capacitance. The resting
potential, El, is $-60\ mV$ or $-65\ mV$, for excitatory or inhibitory cells, respectively. Similarly, the
steepness of the exponential approach to threshold, Δ_V is $2.0\ mV$ or $0.5\ mV$, for excitatory or inhibitory
cells, respectively. When the membrane potential V reaches the threshold, $V_{thre} = -50\ mV$, a spike is
emitted and V is instantaneously reset and clamped to $V_{reset} = -65\ mV$ during a refractory period of
$T_{refrac} = 5\ ms$. The membrane potential of excitatory neurons is also affected by the adaptation variable,
w, with time constant $\tau_w = 500\ ms$, and the dynamics of adaptation is given by parameter $a = 4\ nS$. At
each spike, w is incremented by a value b, which regulates the strength of adaptation. $b = 60 pA$ was used
to model deep anesthesia, and $b = 20\ pA$ for light anesthesia simulations.
454
455   ***From spikes to fluorescence of two photon signal***

456   In order to model the two-photon calcium signal the spikes and the values of the membrane potential
457   V_m (Fig. 7C, grey trace) of each neuron were recorded during the simulation, for each level of adaptation.
458   Increases in the V_m lead to an inward calcium current through voltage-dependent channels. We characterized
459   the L-type high voltage activated calcium current I_{Ca} (Fig. 7C, red trace) as in Rahmati et al. (2016):

$$\mathrm{I_{Ca} = g_{Ca}s(V_m - E_{Ca})} \tag{12}$$

460   where $E_{Ca} = 120\ mV$ and $g_{Ca} = 5\ mS/cm^2$ are the reversal potential and the maximal conductance of
461   this current, respectively. The steady-state voltage dependent activation of the channel (Fig. 7B), is defined

462   by the Boltzmann function:

$$s = \frac{1}{1 + \exp^{-\frac{(V_m - V_{1/2})}{\rho}}} \tag{13}$$

463   with a half-activation voltage $V_{1/2} = -25\ mV$ and a slope factor $\rho = 5mV$ (Ermentrout, 1998; Helton
464   et al., 2005). The intracellular concentration of calcium (cytosolic $[Ca^{2+}]$, Fig. 7C, dark red trace) increases
465   proportionally to the I_{Ca} current, and then it slowly decays back to a basal value $[Ca^{2+}]_i$ (Traub, 1982):

$$\frac{d[Ca^{2+}]}{dt} = -k_{Ca}I_{Ca} - \frac{[Ca^{2+}] - [Ca^{2+}]_i}{\tau_{Ca}} \tag{14}$$

466   with $K_{Ca} = 0.002\ (nM/ms)(\mu A/cm^2)^{-1}$, $\tau_{Ca} = 760\ ms$ and $[Ca^{2+}]_i = 0\ nM$.

467   Finally, the fluorescence $F(t)$ associated with the intracellular calcium concentration (Fig. 7C, green
468   trace) is then computed following the equation:

$$F(t) = dF + K_F \frac{[Ca^{2+}]^{n_H}}{[Ca^{2+}]^{n_H} + K_d} \tag{15}$$

469   where $dF = 0$ and $K_F = 10$ are the offset and the scaling of $F(t)$, $k_d = 375\ nM$ is the dissociation time
470   constant for GCaMP6f (Chen et al., 2013), a measure of the affinity of the fluorescent indicator to the
471   calcium ion, and $n_H = 2.3$ is the Hill coefficient (Chen et al., 2013).

4   RESULTS

472   Here we show first the results we obtained on the simulation of goal-directed movements ("Movement-
473   driven models" pipeline) and then on the modeling of brain alterations after stroke ("Stroke models"
474   pipeline).

475   4.1   Simulation of the experiment on goal-directed movements

As the first component of the proposed framework ("Movement-driven models"), we simulated the
experiment on goal-directed forelimb pulling in the virtual environment and validated the simulation on
experimental data.

In the in-vivo experiment, two healthy mice were trained on the M-Platform to perform active pulling of
the forelimb. As we expected, the contralateral motor cortex showed a highly coherent activation with the
kinetic data. The coherence between the force applied by the animal and the signal recorded in the CFA
was evident both in the low and in the high frequencies band (Fig. 8). For the data that were later used
in simulations we focused on the high band (300 to 40k Hz); in particular we found an high activation of
the motor cortex around the force peaks for both multi-unit activity and single units analysis. This result
proves that for each recording the SUs were successfully extracted by the multi-units. The PSTHs was
used to evaluate the temporal behaviour-related spike activity of every single unit. The behaviour around
the force peaks was different according to the single units selected, but all of them showed that the activity
began to increase before onset of force peaks and came back to the resting value after 0.4 s from the onset.
In order to simulate the descending signal from the motor cortex generating the movement of the forelimb,
we employed these neurophysiological recordings. In particular, the events resulting from the single unit
spike sorting were given as spike times for static spike generator in the neural simulation. As the number
of recorded neurons was low, the spike generators were copied 100 times, while also adding gaussian noise

(with mean = 0ms and standard deviation = 5ms) to the spike times of the copies to avoid synchronicity.
As the neural recording originates mainly from neurons that control the pulling, we decided to connect
the descending stimuli only to interneural populations associated with muscles that are active during the
pulling, i.e the flexors of the the two actuated joints. Therefore, the antagonistic muscles would only actuate
thanks to spinal reflexes. To tune the parameters of these connections, and to produce a muscular activation
that was similar in amplitude to the force recorded in the in-vivo experiments, we performed a preliminary
set of experiments, without the simulated embodiment, in which we empirically tuned the synaptic weights
and number of connections. Due to the absence of the embodiment, at this stage there is no muscle spindle
activity and thus no sensory feedback enabling reflexes. Then, the spinal cord model described in Section
3.1 was connected to the mouse forelimb. In principle, the musculoskeletal embodiment has three pairs of
muscles, but the one controlling the paw is not significantly involved in the pulling of the limb. We did not
consider those when building the neural network to decrease simulation times. Thus, we replicated the same
spinal cord circuitry two times and connected it to the four muscles controlling the elbow and shoulder
joints, named <u>humerus</u> and <u>radius</u> in the simulation. In the closed loop simulated by the Neurorobotics
Platform, the output of the spinal cord model (muscles activation between 0 and 1) could be directly given
to the simulated mouse actuators, while the muscle lengths and contraction speed had to be normalized
before sending them to the muscle spindles models.

Figure 8. Results from the simulated pulling experiment. Comparison between the simulated slide position and the one recorded in the in-vivo experiment (top) and comparison between simulated muscle activation levels and force applied to the physical slide (bottom). To increase the readability of the bottom figure, an upper peak envelope was applied to the signals.

In Fig. 8 we show the results for a simulation trial and a comparison with data recorded from a physical
experiment. We employed kinematic data recorded alongside neural activity in the same in-vivo experiment:
position of the slide and force applied to the slide through the trial. As expected, by comparing the activation
levels with the normalized force applied by the mouse to the slide we can observe that the flexor muscles
are active when there is also a force recorded, and conversely, there is low activation when the slide is still.
It is also worth mentioning that, although the two muscles receive the same inputs from the descending
stimuli, their activation levels are different due to the feedback circuitry of the spinal cord and the activity of
muscle spindles, which are different for the two muscles. Thus, the output of the spinal cord circuitry is not
a mere filtering of the input signals, but it also takes into account the feedback from the embodiment, which

can change during the experiment. This effect underlies the importance of embedding neural circuits in a
proper, realistic embodiment. The comparison between the simulated slide position and the recorded one
shows that, thanks to the recorded neural activity, the muscles are able to overcome the force threshold and
release the slide, and that an actual pulling is performed. Every pulling episode in the trial is reproduced,
even if with different degrees of accuracy. Overall, the mean absolute error between simulated and recorded
slide positions is 13%. The main discrepancies between the simulated and the recorded data come from the
fact the in the simulated mode, the muscular activity is mostly directly proportional to the neural activity,
while in the recorded data this is not always the case. While there is clearly a correlation between presence
of neural activity and motion, the intensity of such activity sometimes does not match the intensity of the
motion.
529

4.2   Local and global brain simulations after stroke and rehabilitation
4.2.1   BNM for brain connectivity changes after stroke and rehabilitation
Within the second pipeline of the framework ("Stroke models"), in this subsection we simulated different
extents of the brain injury and rehabilitation-induced plasticity after stroke. The results from the simulations
are compared with the experimental data, Fig. 2, allowing us to find the best fit with the empirical functional
reorganization in the parameters space of the structural changes in the white matter connectivity.

The averaged calcium activity has a local peak in the power spectrum at around 3.5 Hz (Fig. 9, (A)), which
is within the band relevant for the resting state activity. We hence focused the analysis of the experimental
data on the the upper delta band, $2.5 - 5Hz$, where we consequently band-pass filter the signals. From
these we calculated the pair-wise PLV in each condition, thus constructing the FC matrices for the cortical
regions of interest. Finally, to remove one condition, we calculate the changes of the FC during stroke
and recovery compared with the healthy state, and this is the data feature that is then compared with the
simulated data. For this we use the model described in 3.3.1, to identified which scenarios of structural
alterations cause the best agreement with the data in the modelled FC alterations, Fig. 9. To minimize the
effect of tissue displacement after stroke (Brown et al., 2007), the analysis includes only 12 ipsilesional
regions located outside the stroke core, Fig. 9 (B and D).

The stroke affects not only the inherent activity of the rM1, but all the connected regions. However,
the precise breadth and magnitude of the structural damage, namely which links and to what extent are
they disabled over time, is unknown. Similarly, it is not known which new links are created or reinforced
during the spontaneous recovery or what is modulated by the rehabilitation. On the other hand, the stroke
was shown to consistently change the alignment of dendrites and axons towards the core *in vivo* (Brown
et al., 2007), possibly meaning altered SC, confirming previous works on structural rewiring after stroke
(Dancause, 2005; Nudo, 2013). Hence a numerical exploration of the different possibilities of the stroke and
rewiring in the large-scale BNM is used to unveil the most probable structural alterations associated with
stroke and recovery. The calcium activity in the upper delta band that was chosen for the analysis shows
highly coherent co-activation of different parts of the cortex, compared with the surrogate time-series, Fig. 9
(A). We compared the functional reorganization associated with spontaneous recovery after stroke ("stroke"
group) to rehabilitation-supported recovery ("rehab" group). The changes in the functional connectivity in
"stroke" compared to "healthy" mice (Fig. 9 (B), left matrix) indicate an increased co-activation of all but
one somatosensory areas in the chronic phase after stroke, while visual areas have increased connectivity
with all the regions, and reduced with the retrosplenial cortex. In the rehabilitated mice (Fig. 9 (B), right
matrix), the increase in connectivity of the somatosensory is even higher across all the areas, and there is
also an increased FC of the visual areas between each other and with the somatosensory regions.

Figure 9. Simulated and empirical Functional connectivity and fitting the model. (A) Average power spectrum across the regions and the PLV values for each pair of regions (thin lines) during healthy state, as well as the significance levels from the surrogates (thick dotted lines). Vertical black lines show the boundaries of the lower and upper δ bands. (B) Relative changes of the FC at stroke and at rehabilitation compared to the healthy control for frequency band $f = 2.5 - 5Hz$. (C) Cross correlation of the model upper triangles of FC between the model and the data for fixed global coupling $K = 4.3$ and different levels of stroke (0 for complete damage and 0.9 for damage of 10% of the links) and rebound connectivity (0 for no rewiring and 5 for overall rewiring with strength of 5 times of the damaged links). Parameters: frequency $f = 2Hz$, noise strength $D = 1$. (D) Simulated relative changes of the FC at stroke and rehabilitation relative to the healthy control for the working points marked with red squares in the parameters space in the panel (C). The abbreviations for the areas in (B) and (D) are: VIS - visual, RSP - retrosplenial, SS - somatosensory, al - anterolateral, rl - rostrolateral, p - primary, pm - posteromedial, am - anteromedial, a - anterior, d - dorsal, agl - lateral agranular part, ptr - primary trunk, pll - primary lower limb, pun - primary unassigned, pul - primary upper limb.

A phenomenological neural mass is used to simulate how ipsilesional FC is changed by stroke and
rehabilitation based on the modifications of the SC. For this, we systematically modified the SC to account
for various impacts of stroke and subsequent recovery, in order to find the best match with the patterns
observed in the data. The damage due to the stroke is assumed to be homogeneous across the links
connecting rM1, but their magnitude is varied from 10% to 100%. Similarly, after the recovery it is
assumed that 0 to 500% of the lost connectivity due to stroke is restored homogeneously across the regions
with preexisting links towards rM1, proportionally to the initial strength of their link to rM1. We thus
explore the possibility of up to 5 times of weights of the damaged links to be redistributed along the rest of
the links of the nodes directly connected with the infarct area, in order to also allow for over-compensation
of the lost direct connectivity. The absence of time-delays and the focus on the phase locking, makes
the model insensitive on the chosen frequencies (Petkoski et al., 2016), which are therefore fixed in the
simulations. The natural time-variability of parameters Petkoski and Stefanovska (2012) is assumed to
be stochastic (Petkoski et al., 2018). We hence fix the level of the noise and we explore the impact of

Figure 10. Simulation of peri-stroke local network oscillations: experiments and models. (A) Fluorescence traces obtained from three example cells (two-photon signals) and from the entire field-of-view (wide field signals) recorded in mice under deep and light anesthesia. (B) Raster plot of the spikes (top) and averaged two-photon calcium signals (bottom) computed for the inhibitory (FS, in red) and excitatory (RS, in green) populations in a model of deep or light anesthesia.

the global coupling K and the described strategies of the stroke and recovery. For each combination
of parameters we obtain the same metric of FC as for the empirical data. The parameters space for the
agreement between the modelled and the experimental data about the changes in the FC for the two
parameters of the stroke-induced structural changes are shown in Fig. 9 (C).

Panel (D) of Fig. 9 illustrates the simulated FC for spontaneously recovered "stroke" and "rehab" mice
compared with pre-stroke conditions ("healthy" group), for fixed global coupling and for points in the
parameters space of the stroke damage and rebound connectivity that show the best fitting with the empirical
data. Comparing the simulated, Fig. 9 (B), with the empirical FC, Fig. 9 (D), we see that the best agreement
is achieved for the FC of the somatosensory areas, while that of the visual cortex areas could be improved
by testing different damage and rewiring strategies for those regions. From the model fitting for different
parameters, it is also visible that generally better fit is achieved if the extent of damaged links is decreased
due to rehabilitation-induced remapping. There is also a similar tendency for the rebound connectivity to
be decreased due to recovery training, although there are other possible recovery paths that keep roughly
the same level of rebound connectivity. In conclusion, the systematic exploration of the model parameters
to best fit the empirical data, allows us to obtain the sufficient structural changes that can reproduce the
modulation in FC after stroke and rehabilitation.
592

4.2.2   Simulation of the calcium activity of the peri-infarct network after stroke

Stroke profoundly alters the functionality at the local level in addition to long-range connections. The
local network next to the stroke core switches to slow wave activity (Butz et al., 2004), a type of brain
oscillation that is observed during deep sleep, but also during anesthesia and other pathological brain
states (Sanchez-Vives et al., 2017). Understanding the changes in the activity patterns at the level of the
peri-stroke region is necessary to get insight on the possible mechanisms that underlie functional recovery.
In order to explore the mechanisms that drive the neuronal networks of the peri-stroke areas to oscillate,
we developed a model that reproduces the spiking activity of a local network during slow oscillations and

extended the model to provide the two-photon calcium signal that one would record from that network. We
compared the simulated calcium data with that of the two-photon experiments conducted in anesthetized
mice (see the "Stroke models" box in Fig. 2).
604

We propose that a deficit in neuromodulation produced by the decreased cerebral blood flow in the
periphery of the region affected by the stroke could be responsible for the emergence of slow oscillations
and the general flattening of the EEG through an increase on the level of adaptation of the neurons (Nghiem
et al., 2020). To test this hypothesis, we developed a spiking network model capable of reproducing the
spontaneous activity of a cortical network during different depths of anesthesia (Fig. 7). The strength of
adaptation in the model can be regulated to produce different types of oscillations. In particular, we aimed
at reproducing the changes in frequency of the slow oscillation observed both in the two-photon and in the
wide field calcium imaging *in vivo* experiments when the anesthesia is reduced (Fig.10A). When adaptation
is strong, the model of deep anesthesia produces slow oscillations at 2.1 Hz, while decreasing the strength
of adaptation (model of light anesthesia) leads to slightly faster slow oscillations, at a frequency of 2.31 Hz
(Fig. 10B). Thus, we show that varying the strength of adaptation allows to reproduce the increase in
frequency of the slow oscillations observed in the calcium imaging experiments when decreasing the level
of anesthesia.
618

5   DISCUSSION

In this paper, we explored the steps and methods that are needed to develop a simulation model of a
complete experiment. We designed, validated and combined many components, even though they are
obviously not exhaustive to encounter for the complexity of the real world.

We first simulated execution of goal-directed movements with displacement of objects in the virtual reality
setting. Results show that we could replicate with high accuracy the displacement of an object by a virtual
(healthy) mouse in the simulated environment (Fig. 2). While this simulation is only an approximation of
brain-body interactions, it showcases the capability to simulate large scale neural networks as well as body
dynamics in a virtual environment.

The second pipeline of the framework involved modeling brain injury. We validated the potential of a
brain network model to predict the long-range stroke-induced connectivity changes measured in a real
experiment. We also tested an oscillatory spiking network model to simulate local peri-infarct activity
after stroke. In addition, this model could simulate the fluctuation in calcium concentration due to spiking
activity of homogeneous neuronal network, thus allowing modeling of calcium imaging data.
632

Towards the mechanistic understanding of behavior, recent studies took advantage of virtual reality (VR)
experiments conducted under controlled environment, where behavioral strategies could be isolated and
tested (Dombeck and Reiser, 2012). In a VR experiment, a simulated environment is updated based on
the animal's actions (Chronis et al., 2007; Ritter et al., 2001; Reiser and Dickinson, 2008; Dombeck et al.,
2010). The main drawback of this approach is that the activity of animals dictates not only the response
of the VR but also the properties of the neurons being measured. As a consequence, the closed-loop
VR system shall then be optimized on-line based on the animal's behaviour, which is very challenging.
The approach we propose here instead is based on an off-line simulation, that allows exploring multiple
dimensions in the parameter space of the dynamical model of mouse brain and the environment. Anyway,
both strategies are synergistic with the research of effective functional brain machine interfaces (Santhanam

Figure 11. Future perspective of data and simulations. The scheme depicts the approach to build the framework from data to models and back. The workflow from data to models and simulation in the *Embodied brain* closed loops is shown. The upper, green box shows the *Stroke models* closed loop, the lower, red box shows the *Movement-driven models* closed loop. Colored images represent experiment data, brain and spinal cord models, and simulation of the environment (from left to right). Connections between the components are presented as arrows: solid lines represent the output provided to other blocks; dashed lines indicate the output data of the models that are used for comparison with real data for validation. In gray, models and connections that are still under development. The overlapping green and red region pictures the future integration of the two pipelines, and in particular of the brain models within the NRP.

643 et al. (2006)).

5.1 Movement-driven models Closed-loop

The results showed in Section 4.1 demonstrate that is possible to achieve realistic simulations by
integrating some of the components described previously. Accuracy of the closed-loop simulation could
be increased by removing some simplifications that are currently in place. Some of them are related to
the physical models of the slide and of the musculoskeletal embodiment. Regarding the former, a more
accurate slide simulation will allow to introduce friction effects that are occurring in the real setup, thus we
could avoid putting a muscle activation level threshold for the release of the slide. Moreover, more detailed
spinal cord and musculoskeletal models will be essential to simulate finer movements.

Results shown in Fig. 8 demonstrate the the system is able to simulate the pulling task, albeit with some
inaccuracies on some pulling trials. A presumable cause of this inaccuracy can be identified in the low

number of neurons (less than 20) that is possible to record during an experiment on the platform with the
16 channels linear probe. For this reason, it is possible that the selected units do not encompass the entire
population of neurons involved in the movement. This issue could be mitigated by employing a multi-unit
analysis, however, this will add to the inputs a significant background activity which may not be useful to
generate the pulling movement.

Many parameters of the spinal cord circuitry can be adjusted, depending on the inputs, to accurately
reproduce the movements recorded in the in-vivo experiments. While in this work the tuning was
done manually, a more effective and generalized way would be to use different recordings, both
neurophysiological and kinematic, and employ an optimization similar to what has been done in (Sreenivasa
et al., 2016).

The level of detail of the spinal cord circuitry can clearly be improved. In this work we modelled a
minimal set of components that were capable of replicating experimental data with a certain degree of
realism. To achieve this, it was decided not to arbitrarily increase the complexity of the models by adding
subcircuits whose impact cannot be clearly measured from a comparison with experimental data. Among
these, it is worth mentioning the inclusion of proprioceptive feedback from Golgi tendon organs, which
could be potentially implemented with computational models such as (Mileusnic and Loeb, 2006). Perhaps
more interesting is the modulation of muscle spindle sensitivity from γ-motoneurons, as this is crucial in
the control of both voluntary and involuntary movements. While including a population of γ-motoneurons
could be done by replicating populations of α-motoneurons, measuring the impact of adding this component
is not trivial, especially considered that there is no experimental data measured, in the rehabilitation setup,
that can be used to validate the addition. As such, we decided not include γ-motoneurons in the spinal cord
circuitry.

5.2   Stroke models Closed-loop

AMBA was previously tested and demonstrated to have a predictive value for the resting state dynamics
in healthy conditions, compared with the gold standard individualized diffusion tensor imaging connectome
(Melozzi et al., 2019). One of the main aims of the stroke modeling pipeline in this study is to validate
the use of AMBA in the cases when there are significant changes in SC as compared to the healthy state
for which it was obtained (Oh et al., 2014). This requires finding the most probable structural alterations
corresponding to the stroke and recovery. From the perspective of the integrative neuroscience, this is
especially important as it will allow further application of these altered connectomes validated from the
resting state FC, to generate the particular brain dynamics associated with active forelimb pulling on the
M-platform by stroke and rehabilitated mice.

To this aim, the present study suggests that rehabilitative training could reinforce the connectivity between
motor and visual areas. The iterative loop between experiments and modeling goes towards the confirmation
of this hypothesis via stimulation experiments. New experiments shall verify the necessity of this feature in
promoting the recovery by stimulating the connections between motor and visual cortex, and modulation
of FC could be achieved via optogenetic stimulation, which recently showed to be a promising approach in
stroke recovery (e.g. Pendharkar et al. (2016); Conti et al. (2020); Cheng et al. (2014)).

The results from the model identify routes from the stroke to the recovery in the parameter space that
can be related to neurophysiological quantities, such as the white matter tracts. We could thus determine
links that need to be restored, or prevented from being established, for a successful recovery. One such a
recovery path proposed by the model is the rebound in SC after rehabilitative training, and this is especially

true for the links involving the visual-associated areas. The proposed rebound is due to newly established links from the regions afferent to the site of the stroke. This can lead to overall overcompensation for the SC, and some of these scenarios could be possible paths for recovery. However, it remains to be seen whether the structural changes of such magnitude can be achieved. Several studies previously showed that axonal growth is stimulated by neurorehabilitative activities after stroke, and that sprouting can extend to widespread brain systems (Carmichael et al., 2017). New experiments aimed at verifying the SC modifications shall verify the hypothesis on the importance of modified connectivity in the visual areas for recovery. In addition, stimulation experiments can also strengthen certain links, and with our modelling framework we can virtually compare the effects of each such modification to the observed dynamics.

For the best fitting of the data one would also need to allow different levels of the global coupling, which governs the global level of synchronization and that is already shown to be increased during stroke (Falcon et al., 2016; Corbetta et al., 2018), thus decreasing integration and information capacity (Adhikari et al., 2017) and modularity (Falcon et al., 2015). Thus, one could more precisely identify the path from stroke to recovery for a wider parameters range. This also includes numerically testing different scenarios for heterogeneous connectivity reinforcing (Nudo, 2013) such as reinforcing of contralateral links in general, or those to contralateral stroke region only, or towards the nodes (ipsi-, contra-lateral, or both) that were connected to the damaged region prior stroke.

Possible problems could arise from the alignment of the experimental data, especially after the stroke, due to the shrinkage and the movement of the tissue We have tried to avoid this by excluding from the analysis the regions adjacent to the stroke, but this reduces the predictive value of the model due to smaller number of analyzed regions.

Finally, these experiments provide a picture of the ipsilesional functionality after stroke and rehabilitation, but many other regions are involved, including the contralesional hemisphere (see, for instance, Dodd et al. (2017)). In the next experiments, we plan to enlarge the field-of-view of the wide-field imaging setup to provide longitudinal pictures of cortical functionality over both hemispheres. This should also refine the fitting across parameters, which now contains large areas or similar level of predictability, thus offering more precise recovery path. Individualized connectome data by Diffusion Tensor Imaging during the recovery process is another aspect of the future experiments that should test the predicted changes in the structure that we propose to be the cause of the observed functional alterations of different conditions. In addition, higher resolution SC performed with light-sheet microscopy on individual mice (Allegra Mascaro et al., 2015) could test the model prediction at the final time point of the experiment. As a final step, an individualized therapy could be proposed targeting specific parts of the brain (Spalletti et al. (2017), Allegra Mascaro et al. (2019) and Conti et al. (2020)), depending on the location and the size of the stroke.

5.3 Integration

We propose viable strategies to integrate the brain models described here and to embed them within the *Embodied brain* framework on the NRP (Falotico et al. (2017)) (pictured by the overlapping green and red boxes in Fig. 11).

Before applying it to the simulation of the whole-brain dynamics, the spiking neurons model shall be extended to include the heterogeneous long-range connections either via mean-field approximation or by means of co-simulation with other neural masses (see the spiking neurons model that receives calcium imaging data in Fig. 11). In addition, a model for embedding spiking model modules into the whole brain model is currently under development (displayed as a gray arrow from the spiking neurons to BNM in

Fig. 11). This work includes validating neuronal mass models against high-dimensional neuronal networks.
Once available, this tool will allow bridging the scales of brain models with different levels of description,
and they will be then implemented in the NRP and integrated into the *Embodied brain* framework (grey
arrows in upper box of Fig. 11).

To integrate the large-scale BNM with the proposed spinal cord model, we propose to modulate the
activity of the spiking neurons in the spinal cord by the output of the cortical regions, mainly those related
to the motor activity (displayed by grey arrows in the *Stroke models* closed loop, green upper box in
Fig. 11). In particular, the firing rate of the neurons in the spinal cord that triggers the movements on the
NRP can be driven by the mean activity of the cortical motor regions, or by some specific patterns of their
co-activation, such as a high-level activity propagation, similar as the one observed during the movements.
In this way, the mean neuronal activity of the brain regions at different conditions would trigger movements
at the NRP using the activity of the spinal cord. For the feedback link of the sensory activity, we envision
the information about muscle activity and limb displacement, which is encoded into the firing patterns
of the spinal cord spiking neurons, to directly modulate the mean activity of the sensory motor regions
(displayed as a dashed grey arrow in the *Stroke models* closed loop, upper box in Fig. 11). This on the
other hand would impact the overall brain network dynamics, including the activation patterns of the motor
regions.

To allow the flow of information from the brain to the virtual environment, we anticipate that the next step
will be the integration of a spiking network model of motor areas upstream to the spinal cord model. This
data-driven model of the motor cortex will include populations of pyramidal neurons and interneurons that
can be functionally attached to different lower circuits (displayed by grey arrows in the *Movement-driven*
closed loop, red lower box in Fig. 11). This integration in the proposed framework can be an effective
strategy to effectively close the *Embodied brain* loop.

6 CONCLUSIONS

To summarize, in this study we proposed a methodological framework (named *Embodied brain*) to
investigate a "brain in the loop" by a constructive refinement of experiments and simulation of an embodied
mouse.

Our findings suggest that simulation of real experiments within the proposed framework will help better
understand the complex mechanism that underlies the generation of behavior. Nevertheless, the actual
advantages of the "Embodiement" approach, still under construction, are largely unexplored. Even though
some aspects of complex animal behavior may be represented with good accuracy by modelling single
neural components, without embedding the neural simulations in a physical embodiment it is impossible to
show the effect of such neural systems on the body and the surrounding environment. In our study, it would
be impossible to assess whether or not the neural models are capable of performing the pulling task with
any degree of accuracy, computed on the kinematic data. Furthermore, we believe that new features (e.g.
activation of different brain regions for performing the same task due to degeneracy (Price and Friston,
2002) and its impact for stroke and recovery) will be disclosed by the simulation of the entire experiment.
In conclusion, the framework shown in this study will advance the field by formulating new hypothesis
on the mechanism underlying goal-directed voluntary movements, to be validated on *ad hoc* designed
experiments.In general, the framework could simulate new types of experiments that cannot be run in the
real word. Last but not least, the virtual environment will be an essential tool to reduce the number of
animals used in the experiments, thus making the "Reduction" rule on animal experimentation a feasible

goal.
784

CONFLICT OF INTEREST STATEMENT

785 The authors declare that the research was conducted in the absence of any commercial or financial
786 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

787 ALAM, EF, SP, LV, MOG conceived the study; ALAM, MP, NTC, EC, FR, CS performed experiments;
788 ALAM, EG, SP, LV, MP, NTC, EC, STR, EA, CHB, TBL wrote the paper; All authors agreed with the
789 manuscript; SP, VJ, NTC and AD developed the brain models; LV and EF developed the spinal cord model;
790 LV, AVA, EA, ST developed the simulation in the NRP.

FUNDING

791 This project was supported by the European Union's Horizon 2020 research and innovation programme
792 under grant agreement No. 720270 (SGA1) and No. 785907 (SGA2) Human Brain Project.

ACKNOWLEDGMENTS

793 We thank Krister Andersson, Oliver Schmidt, Martin Øvsthus, Ingrid Reiten and Jan G. Bjaalie of the
794 Human Brain Project curation team for expert assistance to share data via the Human Brain Project
795 Neuroinformatics Platform.
796

DATA AVAILABILITY STATEMENT

The datasets and data description on wide-field calcium imaging generated for this study can be found here:
Allegra Mascaro, A., Conti, E., Sacconi, L., & Pavone, F. (2019). Fluorescence cortical recording of mouse
activity after stroke [Data set]. Human Brain Project Neuroinformatics Platform. DOI: 10.25493/Z9J0-
ZZQ.
The raw datasets on neurophysiological recordings for this study can be found here: Pasquini, Maria,
Spalletti, Cristina, Caleo, Matteo, and Micera, Silvestro. (2019). Recordings of Caudal Forelimb Area in
healthy mice during a forelimb pulling task [Data set]. Zenodo. http://doi.org/10.5281/zenodo.3546068
Processed data and source code that can be used to reproduce the neurophysiological recordings experiment
can be found in the following repository: https://gitlab.com/lore.ucci/closed-loop-mouse-stroke-simulation
The BNM model and all the relevant information to reproduce the simulated BNM data can be found in the
following repository: . Data sharing license: This work is shared under a Creative Commons Attribution
CC BY 4.0 license (https://creativecommons.org)
809

REFERENCES

Adhikari, M. H., Hacker, C. D., Siegel, J. S., Griffa, A., Hagmann, P., Deco, G., et al. (2017). Decreased
integration and information capacity in stroke measured by whole brain models of resting state activity.
Brain 140, 1068–1085. doi:10.1093/brain/awx021
Allegra Mascaro, A. L., Conti, E., Lai, S., Di Giovanna, A. P., Spalletti, C., Alia, C., et al. (2019).
Combined rehabilitation promotes the recovery of structural and functional features of healthy neuronal
networks after stroke. Cell Reports 28, 3474 – 3485.e6. doi:https://doi.org/10.1016/j.celrep.2019.08.062

Allegra Mascaro, A. L., Silvestri, L., Sacconi, L., and Pavone, F. S. (2015). Towards a comprehensive
understanding of brain machinery by correlative microscopy. Journal of biomedical optics 20, 061105

Alstermark, B. (1992). The c3-c4 propriospinal system: target-reaching and food-taking. Muscle Afferents
and Spinal Control of Movement. , 327–354

Bashor, D. P. (1998). A large-scale model of some spinal reflex circuits. Biological cybernetics 78,
147–157

Batterman, R. W. and Rice, C. C. (2014). Minimal model explanations. Philosophy of Science 81,
349–376

Breakspear, M., Heitmann, S., and Daffertshofer, A. (2010). Generative models of cortical oscillations:
neurobiological implications of the Kuramoto model. Front. Hum. Neurosci. 4, 190

Brette, R. and Gerstner, W. (2005). Adaptive exponential integrate-and-fire model as an effective description
of neuronal activity. Journal of neurophysiology 94, 3637–3642

Brown, C. E., Li, P., Boyd, J. D., Delaney, K. R., and Murphy, T. H. (2007). Extensive turnover of dendritic
spines and vascular remodeling in cortical tissues recovering from stroke. Journal of Neuroscience 27,
4101–4109. doi:10.1523/JNEUROSCI.4295-06.2007

Butz, M., Gross, J., Timmermann, L., Moll, M., Freund, H.-J., Witte, O. W., et al. (2004). Perilesional
pathological oscillatory activity in the magnetoencephalogram of patients with cortical brain lesions.
Neuroscience letters 355, 93–96

Cabral, J., Hugues, E., Kringelbach, M. L., and Deco, G. (2012). Modeling the outcome of structural
disconnection on resting-state functional connectivity. NeuroImage 62, 1342–1353. doi:10.1016/j.
neuroimage.2012.06.007

Cabral, J., Hugues, E., Sporns, O., and Deco, G. (2011). Role of local network oscillations in resting-state
functional connectivity. NeuroImage 57, 130–139

Cabral, J., Vidaurre, D., Marques, P., Magalhães, R., Silva Moreira, P., Miguel Soares, J., et al. (2017).
Cognitive performance in healthy older adults relates to spontaneous switching between states of
functional connectivity during rest. Scientific Reports 7, 1–13. doi:10.1038/s41598-017-05425-7

Carmichael, S. T. and Chesselet, M.-F. (2002). Synchronous neuronal activity is a signal for axonal
sprouting after cortical lesions in the adult. Journal of Neuroscience 22, 6062–6070

Carmichael, S. T., Kathirvelu, B., Schweppe, C. A., and Nie, E. H. (2017). Molecular, cellular and
functional events in axonal sprouting after stroke. Experimental Neurology 287, 384 – 394. doi:https:
//doi.org/10.1016/j.expneurol.2016.02.007. Special Issue: Axon Regeneration Across Species

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., et al. (2013). Ultrasensitive
fluorescent proteins for imaging neuronal activity. Nature 499, 295

Cheng, M. Y., Wang, E. H., Woodson, W. J., Wang, S., Sun, G., Lee, A. G., et al. (2014). Optogenetic
neuronal stimulation promotes functional recovery after stroke. Proceedings of the National Academy
of Sciences 111, 12913–12918. doi:10.1073/pnas.1404109111

Choi, H. and Mihalas, S. (2019). Synchronization dependent on spatial structures of a mesoscopic
whole-brain network. PLoS computational biology 15, e1006978

Chronis, N., Zimmer, M., and Bargmann, C. I. (2007). Microfluidics for in vivo imaging of neuronal and
behavioral activity in caenorhabditis elegans. Nature methods 4, 727

Cisi, R. R. and Kohn, A. F. (2008). Simulation system of spinal cord motor nuclei and associated nerves
and muscles, in a web-based architecture. Journal of computational neuroscience 25, 520–542

Conti, E., Allegra Mascaro, A. L., and Pavone, F. S. (2019). Large scale double-path illumination system
with split field of view for the all-optical study of inter-and intra-hemispheric functional connectivity on
mice. Methods and Protocols 2. doi:10.3390/mps2010011

Conti, E., Allegra Mascaro, A. L., Scaglione, A., de Vito, G., Calugi, F., Pasquini, M., et al. (2020).
Synergic effect of optogenetic stimulation and motor training boosts recovery of motor functionality
after stroke supported by segregation of motor representation. bioRxiv doi:10.1101/2020.03.05.974972

Corbetta, M., Siegel, J. S., and Shulman, G. L. (2018). On the low dimensionality of behavioral deficits and
alterations of brain network connectivity after focal injury. Cortex , 1–9doi:10.1016/j.cortex.2017.12.017

Dancause, N. (2005). Extensive Cortical Rewiring after Brain Injury. Journal of Neuroscience 25,
10167–10179. doi:10.1523/JNEUROSCI.3256-05.2005

Davison, A. P., Brüderle, D., Eppler, J., Kremkow, J., Muller, E., Pecevski, D., et al. (2009). Pynn: A
common interface for neuronal network simulators. Front Neuroinform 2, 11–11. doi:10.3389/neuro.11.
011.2008

Deco, G. and Jirsa, V. K. (2012). Ongoing cortical activity at rest: criticality, multistability, and ghost
attractors. J. Neurosci. 32, 3366–3375

Deco, G., Jirsa, V. K., and McIntosh, A. R. (2011). Emerging concepts for the dynamical organization of
resting-state activity in the brain. Nat. Rev. Neurosci. 12, 43–56

Deco, G., Jirsa, V. K., McIntosh, A. R., Sporns, O., and Kötter, R. (2009). Key role of coupling, delay, and
noise in resting brain fluctuations. P. Natl. Acad. Sci. USA 106, 10302–10307

Deco, G., Jirsa, V. K., Robinson, P. A., Breakspear, M., and Friston, K. (2008). The Dynamic Brain: From
Spiking Neurons to Neural Masses and Cortical Fields. PLoS Comput. Biol. 4, e1000092

Deco, G., Kringelbach, M. L., Jirsa, V., and Ritter, P. (2016). The dynamics of resting fluctuations in the
brain: metastability and its dynamical cortical core. Scientific reports 7, 3095. doi:10.1101/065284

DeMarse, T. B., Wagenaar, D. A., Blau, A. W., and Potter, S. M. (2001). The neurally controlled animat:
biological brains acting with simulated bodies. Autonomous robots 11, 305–310

Destexhe, A. (2009). Self-sustained asynchronous irregular states and up–down states in thalamic,
cortical and thalamocortical networks of nonlinear integrate-and-fire neurons. Journal of computational
neuroscience 27, 493

Dodd, K. C., Nair, V. A., and Prabhakaran, V. (2017). Role of the contralesional vs. ipsilesional hemisphere
in stroke recovery. Frontiers in Human Neuroscience 11, 469. doi:10.3389/fnhum.2017.00469

Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L., and Tank, D. W. (2010). Functional imaging of
hippocampal place cells at cellular resolution during virtual navigation. Nature neuroscience 13, 1433

Dombeck, D. A. and Reiser, M. B. (2012). Real neuroscience in virtual worlds. Current Opinion in
Neurobiology 22, 3 – 10. doi:https://doi.org/10.1016/j.conb.2011.10.015. Neurotechnology

Ermentrout, B. (1998). Linearization of fi curves by adaptation. Neural computation 10, 1721–1729

Falcon, M. I., Riley, J. D., Jirsa, V., McIntosh, A. R., Elinor Chen, E., and Solodkin, A. (2016). Functional
Mechanisms of Recovery after Chronic Stroke: Modeling with the Virtual Brain. eNeuro 3. doi:10.1523/
ENEURO.0158-15.2016

Falcon, M. I., Riley, J. D., Jirsa, V., McIntosh, A. R., Shereen, A. D., Elinor Chen, E., et al. (2015). The
virtual brain: Modeling biological correlates of recovery after chronic stroke. Frontiers in Neurology 6,
1–13. doi:10.3389/fneur.2015.00228

Falotico, E., Vannucci, L., Ambrosano, A., Albanese, U., Ulbrich, S., Vasquez Tieck, J. C., et al. (2017).
Connecting artificial brains to robots in a comprehensive simulation framework: The neurorobotics
platform. Frontiers in Neurorobotics 11, 2. doi:10.3389/fnbot.2017.00002

Franklin, K. B., Paxinos, G., et al. (2008). The mouse brain in stereotaxic coordinates, vol. 3 (Academic
press New York:)

Fuglevand, A. J., Winter, D. A., and Patla, A. E. (1993). Models of recruitment and rate coding organization
in motor-unit pools. Journal of neurophysiology 70, 2470–2488

Gewaltig, M. and Diesmann, M. (2007). NEST (NEural Simulation Tool). Scholarpedia 2, 1430.
doi:10.4249/scholarpedia.1430. Revision #130182

Ghosh, A., Rho, Y., McIntosh, A. R., Kötter, R., and Jirsa, V. K. (2008). Noise during rest enables the
exploration of the brain's dynamic repertoire. PLoS Comput. Biol. 4, e1000196

Grienberger, C. and Konnerth, A. (2012). Imaging Calcium in Neurons. Neuron 73, 862–885. doi:10.
1016/j.neuron.2012.02.011

H. Dana, A. H. B. S. C. G. L. L. D. S. K. e. K. S., T.W. Chen (2014). Thy1-GCaMP6 Transgenic Mice for
Neuronal Population Imaging In Vivo. Plos One 9. doi:10.1371/journal.pone.0108697

[Dataset] HBPNeurorobotics (2019). Robot designer. `https://github.com/`
`HBPNeurorobotics/BlenderRobotDesigner`

Helton, T. D., Xu, W., and Lipscombe, D. (2005). Neuronal l-type calcium channels open quickly and are
inhibited slowly. Journal of Neuroscience 25, 10247–10251

Hinkel, G., Groenda, H., Krach, S., Vannucci, L., Denninger, O., Cauli, N., et al. (2017). A framework for
coupled simulations of robots and spiking neuronal networks. Journal of Intelligent & Robotic Systems
85, 71–91

Hinkel, G., Groenda, H., Vannucci, L., Denninger, O., Cauli, N., and Ulbrich, S. (2015). A domain-
specific language (dsl) for integrating neuronal networks in robot control. In Proceedings of the
2015 Joint MORSE/VAO Workshop on Model-Driven Robot Software Engineering and View-based
Software-Engineering (New York, NY, USA: ACM), MORSE/VAO '15, 9–15. doi:10.1145/2802059.
2802060

Hodgkin, A. L. and Huxley, A. F. (1952). A quantitative description of membrane current and its application
to conduction and excitation in nerve. The Journal of physiology 117, 500–544

Honey, C. J., Kötter, R., Breakspear, M., and Sporns, O. (2007). Network structure of cerebral cortex
shapes functional connectivity on multiple time scales. P. Natl. Acad. Sci. USA 104, 10240–10245

Izhikevich, E. M. (1998). Phase models with explicit time delays. Phys. Revi. E 58, 905–908. doi:10.
1103/PhysRevE.58.905

Izhikevich, E. M. and Edelman, G. M. (2008). Large-scale model of mammalian thalamocortical systems.
P. Natl. Acad. Sci. USA 105, 3593–3598

Jirsa, V. K., Stacey, W. C., Quilichini, P. P., Ivanov, A. I., and Bernard, C. (2014). On the nature of seizure
dynamics. Brain 137, 2210–2230. doi:10.1093/brain/awu133

Johansen-Berg, H. and Rushworth, M. F. S. (2009). Using diffusion imaging to study human connectional
anatomy. Annu. Rev. Neurosci. 32, 75–94. doi:10.1146/annurev.neuro.051508.135735

Koenig, N. and Howard, A. (2004). Design and use paradigms for gazebo, an open-source multi-robot
simulator. In 2004 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS) (IEEE
Cat. No.04CH37566). vol. 3, 2149–2154 vol.3. doi:10.1109/IROS.2004.1389727

Kuramoto, Y. (1984). Chemical Oscillations, Waves, and Turbulence (Berlin: Springer-Verlag)

Lachaux, J. P., Rodriguez, E., Martinerie, J., and Varela, F. J. (1999). Measuring phase synchrony in brain
signals. Hum. Brain Mapp. 8, 194–208

Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V., and Stefanovska, A. (2018). Surrogate data for
hypothesis testing of dynamical systems. Physics Reports doi:10.1016/j.physrep.2018.06.001

Markram, H., Muller, E., Ramaswamy, S., Reimann, M. W., Abdellah, M., Sanchez, C. A., et al. (2015).
Reconstruction and Simulation of Neocortical Microcircuitry. Cell 163, 456–492. doi:10.1016/j.cell.
2015.09.029

Melozzi, F., Bergmann, E., Harris, J. A., Kahn, I., Jirsa, V., and Bernard, C. (2019). Individual structural
features constrain the functional connectome. bioRxiv , 613307

Melozzi, F., Woodman, M. M., Jirsa, V. K., and Bernard, C. (2017). The Virtual Mouse Brain:
A Computational Neuroinformatics Platform to Study Whole Mouse Brain Dynamics. Eneuro 4,
ENEURO.0111–17.2017. doi:10.1523/ENEURO.0111-17.2017

Mileusnic, M. P., Brown, I. E., Lan, N., and Loeb, G. E. (2006). Mathematical models of proprioceptors. i.
control and transduction in the muscle spindle. Journal of neurophysiology 96, 1772–1788

Mileusnic, M. P. and Loeb, G. E. (2006). Mathematical models of proprioceptors. ii. structure and function
of the golgi tendon organ. Journal of Neurophysiology 96, 1789–1802

Millard, M., Uchida, T., Seth, A., and Delp, S. L. (2013a). Flexing computational muscle: Modeling and
simulation of musculotendon dynamics. Journal of Biomechanical Engineering 135, 021005–021005–
11. doi:10.1115/1.4023390

Millard, M., Uchida, T., Seth, A., and Delp, S. L. (2013b). Flexing computational muscle: modeling and
simulation of musculotendon dynamics. J Biomech Eng 135, 021005–021005. doi:10.1115/1.4023390

Moraud, E. M., Capogrosso, M., Formento, E., Wenger, N., DiGiovanna, J., Courtine, G., et al. (2016).
Mechanisms underlying the neuromodulation of spinal circuits for correcting gait and balance deficits
after spinal cord injury. Neuron 89, 814–828

Nakagawa, T. T., Jirsa, V. K., Spiegler, A., McIntosh, A. R., and Deco, G. (2013). Bottom up modeling
of the connectome: linking structure and function in the resting brain and their changes in aging.
Neuroimage 80, 318–329

Nghiem, T.-A. E., Tort-Colet, N., Górski, T., Ferrari, U., Moghimyfiroozabad, S., Goldman, J. S., et al.
(2020). Cholinergic Switch between Two Types of Slow Waves in Cerebral Cortex. Cerebral Cortex
doi:10.1093/cercor/bhz320. Bhz320

Nudo, R. J. (2013). Recovery after brain injury: mechanisms and principles. Frontiers in Human
Neuroscience 7, 1–14. doi:10.3389/fnhum.2013.00887

Oh, S. W., Harris, J. A., Ng, L., Winslow, B., Cain, N., Mihalas, S., et al. (2014). A mesoscale connectome
of the mouse brain. Nature 508, 207–214. doi:10.1038/nature13186

Pasquini, M., Lai, S., Spalletti, C., Cracchiolo, M., Conti, S., Panarese, A., et al. (2018). A robotic system
for adaptive training and function assessment of forelimb retraction in mice. IEEE Trans Neural Syst
Rehabil Eng. 26, 1803–1812

Pendharkar, A. V., Levy, S. L., Ho, A. L., Sussman, E. S., Cheng, M. Y., and Steinberg, G. K. (2016).
Optogenetic modulation in stroke recovery. Neurosurgical focus 40, E6

Petkoski, S. and Jirsa, V. K. (2019). Transmission time delays organize the brain network synchronization.
Philosophical Transactions of the Royal Society A 377, 20180132

Petkoski, S., Palva, J. M., and Jirsa, V. K. (2018). Phase-lags in large scale brain synchronization :
Methodological considerations and in-silico analysis. PLoS Comput Biol 14, 1–30. doi:10.1371/journal.
pcbi.1006160

Petkoski, S., Spiegler, A., Proix, T., Aram, P., Temprado, J.-J., and Jirsa, V. K. (2016). Heterogeneity
of time delays determines synchronization of coupled oscillators. Physical Review E 94, 012209.
doi:10.1103/PhysRevE.94.012209

Petkoski, S. and Stefanovska, A. (2012). Kuramoto model with time-varying parameters. Phys. Rev. E 86,
046212

Pierrot-Deseilligny, E. and Burke, D. (2005). The circuitry of the human spinal cord: its role in motor
control and movement disorders (Cambridge University Press)

Pikovsky, A., Rosenblum, M., and Kurths, J. (2001). Synchronization – A Universal Concept in Nonlinear
Sciences (Cambridge: Cambridge University Press)

Ponce-Alvarez, A., Deco, G., Hagmann, P., Romani, G. L., Mantini, D., and Corbetta, M. (2015). Resting-state temporal synchronization networks emerge from connectivity topology and heterogeneity. PLoS Comput. Biol. 11, e1004100

Price, C. J. and Friston, K. J. (2002). Degeneracy and cognitive anatomy. Trends in cognitive sciences 6, 416–421

Puchades, M. A., Csucs, G., Ledergerber, D., Leergaard, T. B., and Bjaalie, J. G. (2019). Spatial registration of serial microscopic brain images to three-dimensional reference atlases with the quicknii tool. PloS one 14, e0216796

R. Yasuda, V. S. T. P. B. S., E.A. Nimchinski and Svoboda, K. (2004). Imaging calcium concentration dynamics in small neuronal compartments. Sci STKE. 2004. doi:10.1126/stke.2192004pl5

Rabiller, G., He, J.-W., Nishijima, Y., Wong, A., and Liu, J. (2015). Perturbation of brain oscillations after ischemic stroke: a potential biomarker for post-stroke function and therapy. International journal of molecular sciences 16, 25605–25640

Rahmati, V., Kirmse, K., Marković, D., Holthoff, K., and Kiebel, S. J. (2016). Inferring neuronal dynamics from calcium imaging data using biophysical models and bayesian inference. PLoS computational biology 12, e1004736

Ramanathan, D. S., Guo, L., Gulati, T., Davidson, G., Hishinuma, A. K., Won, S.-J., et al. (2018). Low-frequency cortical activity is a neuromodulatory target that tracks recovery after stroke. Nature medicine 24, 1257

Reger, B. D., Fleming, K. M., Sanguineti, V., Alford, S., and Mussa-Ivaldi, F. A. (2000). Connecting brains to robots: an artificial body for studying the computational properties of neural tissues. Artificial life 6, 307–324

Reiser, M. B. and Dickinson, M. H. (2008). A modular display system for insect behavioral neuroscience. Journal of neuroscience methods 167, 127–139

Rijsdijk, M., Leijten, F., and Slooter, A. (2008). Continuous eeg monitoring in the intensive care unit. Neth J Crit Care 12, 157–62

Ritter, D. A., Bhatt, D. H., and Fetcho, J. R. (2001). In vivo imaging of zebrafish reveals differences in the spinal networks for escape and swimming movements. Journal of Neuroscience 21, 8956–8965

Roy, D., Ghosh, A., and Jirsa, V. K. (2011). Phase description of spiking neuron networks with global electric and synaptic coupling. Phys. Rev. E 83, 051909

Saenger, V. M., Ponce-Alvarez, A., Adhikari, M., Hagmann, P., Deco, G., and Corbetta, M. (2017). Linking Entropy at Rest with the Underlying Structural Connectivity in the Healthy and Lesioned Brain. Cerebral Cortex , 1–11doi:10.1093/cercor/bhx176

Saggio, M. L., Spiegler, A., Bernard, C., and Jirsa, V. K. (2017). Fast?Slow Bursters in the Unfolding of a High Codimension Singularity and the Ultra-slow Transitions of Classes. The Journal of Mathematical Neuroscience 7, 7. doi:10.1186/s13408-017-0050-8

Sanchez-Vives, M. V., Massimini, M., and Mattia, M. (2017). Shaping the default activity pattern of the cortical network. Neuron 94, 993–1001

Santhanam, G., Ryu, S. I., Byron, M. Y., Afshar, A., and Shenoy, K. V. (2006). A high-performance brain–computer interface. nature 442, 195

Sanz-Leon, P., Knock, S. A., Spiegler, A., and Jirsa, V. K. (2015). Mathematical framework for large-scale brain network modeling in the virtual brain. Neuroimage 111, 385–430

Sanz Leon, P., Knock, S. a., Woodman, M. M., Domide, L., Mersmann, J., McIntosh, A. R., et al. (2013). The Virtual Brain: a simulator of primate brain network dynamics. Frontiers in neuroinformatics 7, 10. doi:10.3389/fninf.2013.00010

Schmidt, H., Petkov, G., Richardson, M. P., and Terry, J. R. (2014). Dynamics on Networks: The Role of
Local Dynamics and Global Networks on the Emergence of Hypersynchronous Neural Activity. PLoS
Computational Biology 10. doi:10.1371/journal.pcbi.1003947

Sheppard, L. W., Hale, A. C., Petkoski, S., McClintock, P. V. E., and Stefanovska, A. (2013). Characterizing
an ensemble of interacting oscillators: The mean-field variability index. Phys. Rev. E 87, 012905

Spalletti, C., Alia, C., Lai, S., Panarese, A., Conti, S., Micera, S., et al. (2017). Combining robotic training
and inactivation of the healthy hemisphere restores pre-stroke motor patterns in mice. eLife 6, 1–31.
doi:10.7554/eLife.28662

Spalletti, C., Lai, S., Mainardi, M., Panarese, A., Ghionzoli, A., Alia, C., et al. (2014). A Robotic System
for Quantitative Assessment and Poststroke Training of Forelimb Retraction in Mice. Neurorehabilitation
and Neural Repair 28(2), 188–196. doi:10.1177/1545968313506520

Spiegler, A., Hansen, E. C. A., Bernard, C., McIntosh, A. R., and Jirsa, V. K. (2016). Selective activation of
resting state networks following focal stimulation in a connectome-based network model of the human
brain. eNeuro 3, ENEURO.0068–16.2016. doi:10.1523/ENEURO.0068-16.2016

Sreenivasa, M., Ayusawa, K., and Nakamura, Y. (2016). Modeling and identification of a realistic spiking
neural network and musculoskeletal model of the human arm, and an application to the stretch reflex.
IEEE Transactions on Neural Systems and Rehabilitation Engineering 24, 591–602

Stankovski, T., Petkoski, S., Smith, A. F., McClintock, P. V. E., Stefanovska, A., Raeder, J., et al.
(2016). Alterations in the coupling functions between cortical and cardio-respiratory oscillations
due to anaesthesia with propofol and sevoflurane. Philosophical Transactions of the Royal Society A:
Mathematical, Physical and Engineering Sciences 374, 20150186. doi:10.1098/rsta.2015.0186

Stefanescu, R. A. and Jirsa, V. K. (2008). A low dimensional description of globally coupled heterogeneous
neural networks of excitatory and inhibitory neurons. PLoS Computational Biology 4. doi:10.1371/
journal.pcbi.1000219

Stienen, A. H., Schouten, A. C., Schuurmans, J., and Van Der Helm, F. C. (2007). Analysis of reflex
modulation with a biologically realistic neural network. Journal of computational neuroscience 23, 333

Tessadori, J., Bisio, M., Martinoia, S., and Chiappalone, M. (2012). Modular neuronal assemblies
embodied in a closed-loop environment: toward future integration of brains and machines. Frontiers in
neural circuits 6, 99

Ton, R., Deco, G., and Daffertshofer, A. (2014). Structure-Function Discrepancy: Inhomogeneity and
Delays in Synchronized Neural Networks. PLoS Comput. Biol. 10. doi:10.1371/journal.pcbi.1003736

Traub, R. (1982). Simulation of intrinsic bursting in ca3 hippocampal neurons. Neuroscience 7, 1233–1242

T.W. Chen, Y. S. S. P. S. R. A. E. S. R. K. M. O. V. J. L. L. K. S., T.J. Wardill and Kim, D. (2013).
Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499, 295–300. doi:10.1038/
nature12354

Vanni, M. P., Chan, A. W., Balbi, M., Silasi, G., and Murphy, T. H. (2017). Mesoscale Mapping of Mouse
Cortex Reveals Frequency-Dependent Cycling between Distinct Macroscale Functional Modules. The
Journal of Neuroscience 37, 7513–7533. doi:10.1523/JNEUROSCI.3560-16.2017

Vannucci, L., Falotico, E., and Laschi, C. (2017). Proprioceptive feedback through a neuromorphic muscle
spindle model. Frontiers in Neuroscience 11, 341

Váša, F., Shanahan, M., Hellyer, P. J., Scott, G., Cabral, J., and Leech, R. (2015). Effects of lesions on
synchrony and metastability in cortical networks. NeuroImage 118, 456–467. doi:10.1016/j.neuroimage.
2015.05.042

Wilson, H. R. and Cowan, J. D. (1972). Excitatory and inhibitory interactions in localized populations of
model neurons. Biophysical journal 12, 1–24. doi:10.1016/S0006-3495(72)86068-5

Wong, K.-F. and Wang, X.-J. (2006). A Recurrent Network Mechanism of Time Integration in Perceptual Decisions. Journal of Neuroscience 26, 1314–1328. doi:10.1523/JNEUROSCI.3733-05.2006

Wright, P. W., Brier, L. M., Bauer, A. Q., Baxter, G. A., Kraft, A. W., Reisman, M. D., et al. (2017). Functional connectivity structure of cortical calcium dynamics in anesthetized and awake mice. PLoS ONE 12, 1–27. doi:10.1371/journal.pone.0185759

Yang, H.-W. and Lemon, R. (2003). An electron microscopic examination of the corticospinal projection to the cervical spinal cord in the rat: lack of evidence for cortico-motoneuronal synapses. Experimental brain research 149, 458–469

Yilmaz, O., Birbaumer, N., and Ramos-Murguialday, A. (2015). Movement related slow cortical potentials in severely paralyzed chronic stroke patients. Frontiers in Human Neuroscience 8, 1033. doi:10.3389/fnhum.2014.01033

Zerlaut, Y., Chemla, S., Chavane, F., and Destexhe, A. (2017). Modeling mesoscopic cortical dynamics using a mean-field model of conductance-based networks of adaptive exponential integrate-and-fire neurons. Journal of Computational Neuroscience , 1–17doi:10.1007/s10827-017-0668-2

Zerlaut, Y. and Destexhe, A. (2017). Enhanced responsiveness and low-level awareness in stochastic network states. Neuron 94, 1002–1009

Zimmermann, J., Ritter, P., Shen, K., Rothmeier, S., Schirner, M., and McIntosh, A. R. (2016). Structural architecture supports functional organization in the human aging brain at a regionwise and network level. Human Brain Mapping 37, 2645–2661. doi:10.1002/hbm.23200

Zrenner, C., Belardinelli, P., Müller-Dahlhaus, F., and Ziemann, U. (2016). Closed-loop neuroscience and non-invasive brain stimulation: A tale of two loops. Frontiers in Cellular Neuroscience 10, 92. doi:10.3389/fncel.2016.00092