Phenotypic innovation in one tooth induced concerted developmental evolution in another

Marie Sémon [1]*, Klara Steklikova[2,3], Marion Mouginot[1], Manon Peltier[1], Philippe Veber[4], Laurent Guéguen[4], Sophie Pantalacci[1]*

[1]Laboratoire de Biologie et Modelisation de la Cellule, Ecole Normale Superieure de Lyon, CNRS, UMR 5239, Inserm, U1293, Universite Claude Bernard Lyon 1, 46 allee d'Italie F-69364 Lyon, France.

[2]Institute of Histology and Embryology, First Faculty of Medicine, Charles University, Prague, Czech Republic.

[3]Department of Cell Biology, Faculty of Science, Charles University, Prague, Czech Republic

[4]Univ Lyon, Université Claude Bernard Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive, F-69100, Villeurbanne, France

* equal contribution & co-corresponding
correspondence: sophie.pantalacci@ens-lyon.fr, marie.semon@ens-lyon.fr

Short title:

Concerted developmental evolution underlaid independent phenotypic evolution

AUTHOR CONTRIBUTIONS

S.P. and M.S. conceived the study, designed and performed the experiments, analysed the data and wrote the manuscript. M.M., M. P. and S.P collected embryos and performed *in situ* hybridizations. KS performed immunostaining and 3D reconstructions. Models and code for cusp patterning were developed and refined by L.G, and models of temporal expression were initiated by P.V. All authors read and commented on the manuscript.

ABSTRACT

Serial appendages are similar organs found at different places in the body, such as

fore/hindlimbs or different teeth. They are bound to develop with the same pleiotropic

genes, apart from identity genes. These identity genes have logically been implicated in

cases where a single appendage evolved a drastically new shape while the other retained

an ancestral shape, by enabling developmental changes *specifically* in one organ. Here, we

showed that independent evolution involved developmental changes happening *in both*

organs, in two well characterized model systems.

Mouse upper molars evolved a new dental plan with two more cusps on the lingual side,

while the lower molar kept a much more ancestral morphology, as did the molars of

hamster, our control species. We obtained quantitative timelines of cusp formation and

corresponding transcriptomic timeseries in the 4 molars. We found that a molecular and

morphogenetic identity of lower and upper molars predated the mouse and hamster

divergence and likely facilitated the independent evolution of molar's lingual side in the

mouse lineage. We found 3 morphogenetic changes which could combine to cause the

supplementary cusps in the upper molar and a candidate gene, *Bmper.*

Unexpectedly given its milder morphological divergence, we observed extensive changes

in mouse lower molar development. Its transcriptomic profiles diverged as much as, and co-

evolved extensively with, those of the upper molar. Consistent with the transcriptomic

quantifications, two out of the three morphogenetic changes also impacted lower molar

development.

Moving to limbs, we show the drastic evolution of the bat wing also involved gene

expression co-evolution and a combination of specific and pleiotropic changes.

Independent morphological innovation in one organ therefore involves concerted

developmental evolution of the other organ. This is facilitated by evolutionary flexibility of

its development, a phenomenon known as Developmental System Drift.

KEYWORDS Correlated evolution, Pleiotropy, Cryptic developmental evolution,

Morphological innovation, Comparative transcriptomics, Developmental System Drift

AUTHOR SUMMARY

Serial organs, such as the different wings of an insect or the different limbs or teeth of a

vertebrate, can develop into drastically different shapes due to the position-specific

expression of so-called "identity" genes. Often during evolution, one organ evolves a new

shape while another retains a conserved shape. It was thought that identity genes were

responsible for these cases of independent evolution, by enabling developmental changes

specifically in one organ. Here, we showed that developmental changes evolved *in both*

organs to enable the independent evolution of the upper molar in mice and the wing in

bats. In the organ with the new shape, several developmental changes combine. In the

organ with the conserved shape, part of these developmental changes are seen as well.

This modifies the development but is not sufficient to drastically change the phenotype, a

phenomenon known as "Developmental System Drift", DSD. Thus, the independent

evolution of one organ relies on concerted molecular changes, which will contribute to

adaptation in one organ and be no more than DSD in another organ. This concerted

evolution could apply more generally to very different body parts and explain previous

observations on gene expression evolution.

77
78
79
80

INTRODUCTION

Serial appendages are repetitions of similar appendages in the body, such as different legs or wings in arthropods, vertebrate fore- and hindlimbs, or different teeth. According to their position and function, they can have very similar or very different shapes. Although a certain degree of individuation is often present deeply in evolution, its magnitude can evolve, with new shapes appearing specifically in one appendage. For example, fore- and hindlimb have had different shapes since early tetrapod evolution (Minelli, 2003; Siomava et al., 2020). In bats, the forelimb evolved into a wing, while the hindlimb conserved a more ancestral morphology (Cooper et al., 2012; Sadier et al., 2021). Similarly, in insects, one of the two pairs of wings was modified to form elytra and haltere in coleoptera and diptera, respectively (Tomoyasu, 2017). How selection could act to drastically change the shape of one serial appendage independently of another is an intriguing question.

The reason for that is the pleiotropy constraint. During development, serial organs develop with the same pleiotropic genes, except a handful of key transcriptional regulators, whose expression is appendage-specific. These "selector genes" or "identity genes", including the famous homeotic genes, are necessary to form the right appendage at the right place (Mann & Carroll, 2002; Tomoyasu, 2017; Weatherbee & Carroll, 1999). For example, *Ubx* is necessary to form a haltere instead of a wing by regulating hundreds of pleiotropic target genes specifically in the haltere (Hersh et al., 2007; Pavlopoulos & Akam, 2011)). The enhancers of these target genes are surprisingly pleiotropic as they are shared with the wing (McKay & Lieb, 2013). Intuitively, mutations in identity genes, or in the regulatory regions they target, might be easily selected because they will inherently have appendage-specific effects (Carroll, 2008; Morgalev et al., 2023). In contrast, mutations in other parts of pleiotropic genes might more often be counterselected, because they may have an effect on development of both appendages, which may be advantageous in one appendage but deleterious in the other. Therefore, there is an expectation that identity genes will play a central role in the independent evolution of serial appendages.

Many examples of independent appendage evolution have been studied, which confirm the

role of identity genes, but not only.

In the simplest cases, just a subpart of the appendage has changed, such as hairless parts

or specialized hair structures (eg. sexcombs) in fly legs. As expected, these new traits were

associated with new expression patterns of homeotic genes specific of one leg (eg. *Ubx*;

Scr) or their targets (eg. *Dsx*) and with the evolution of their cis-regulatory regions (G. K.

Davis et al., 2007; Eksi et al., 2018; Stern, 1998; Tanaka et al., 2011)   . This nicely explains

why these fly leg structures evolved at a specific position (or in one sex), but the

developmental and genetic changes through which selection has shaped these structures is

still under study (Atallah et al., 2014).

More complex cases concern the whole appendage, such as halteres and elytra in insects

(Tomoyasu, 2017), jump-adapted legs in insects (Mahfooz et al., 2007; Refki et al., 2014)

and rodents (Saxena et al., 2022), wings in bats (Cooper et al., 2012; Sadier et al., 2021), or

patterns of eyespots in butterflies fore- and hindwings (Matsuoka & Monteiro, 2021, 2022).

Here again studies have pointed to a role for homeotic and identity genes in general

(Booker et al., 2016; Matsuoka & Monteiro, 2022; Refki et al., 2014; Saxena et al., 2022).

Moreover, in species where wings are well differentiated (e.g. wing/haltere of flies, or to a

lesser extent, anterior/posterior wings of bees), the overall dose of hox genes appears more

different between the two appendages than in species with less differentiated appendages

(e.g. anterior/posterior wing of dragonflies), suggesting that the evolution of a differential

hox dose is an important determinant of appendage differentiation (Paul et al., 2021). Other

developmental genes have been implicated as well, because they evolved a new

expression pattern in the modified appendage that is consistent with its phenotype.

(Saxena et al., 2022; Z. Wang et al., 2014). More intriguingly, transcription factors which are

expressed in both appendages have appendage-specific functions revealed by knock-out

experiments or tests in heterologous species (Cretekos et al., 2008; Matsuoka & Monteiro,

2022; Ravisankar et al., 2016; Tomoyasu et al., 2009). It is assumed that this appendage-

specific function is provided by identity genes, either directly (through unknown differential

cis-regulation in the two appendages) or indirectly by providing context-dependency, but

this remains generally untested. Together this indicates that the independent evolution of

serial organs does cope with pleiotropic genes, but does not fully explain how.

Here we chose a new model and a different approach to address this question, focusing on

developmental dynamics. We studied the independent evolution of a drastically new shape

in the mouse upper molar, which is nicely described in the fossil record. Since it occurred

relatively recently, we could compare closely related species. Another strong advantage of

this model is that the development of mouse molars is very well understood

mechanistically, from years of developmental genetics and morphogenesis modeling.

Finally, we devised specific ways to quantify and decipher the evolution of development

based on the comparison of transcriptome time series.

Between 18-12 million years ago, the upper molars of mouse and rat ancestors gradually

acquired a new cusp row on the lingual side of the molar, and reduced cusps size on the

buccal side (Figure 1). This new dental plan and accompanying changes in mastication

movements are adaptive and associated with the success of murine rodent radiation (Lazzari

et al., 2008; Tiphaine et al., 2013). Changes in the lower molar were limited to the

connections between cusps, keeping cusp number and size constant (Figure 1). Because the

shape of upper and lower molars were different although less individuated in the basal

"cricetine" rodents, from which murine rodents emerged (Figure 1) this is not a case of *de*

novo individuation. In fact, lower and upper molars already had different morphologies in

the first mammals (B. M. Davis, 2011; Hillson, 2005). Hamsters are today's good

representative of the basal "cricetine" rodents. In the golden hamster lineage, both molars

kept the ancestral cusp number and organization. We can make the reasonable assumption

that the hamster presents ancestral developmental features, and in this study we use this

species as a phylogenetic control.

Morphogenesis

Cusp patterning

Expression profiling

PCA, Whole tooth samples

Molars develop from the physical and molecular interaction between an epithelium and a

mesenchyme (Jernvall & Thesleff, 2012), Figure 1A). The epithelium grows and folds to form

the crown and its cusps under the influence of two types of signalling centres, PEK and SEK

(Primary and Secondary Enamel Knots respectively) (Jernvall & Thesleff, 2012). First, the PEK

determines the field of the molar crown. As this field grows, the SEKs are patterned

sequentially and determine the cusps, starting with a buccal cusp (Cho et al., 2007;

Pantalacci et al., 2017). This spatio-temporal sequence depends on activation-inhibition

loops involving both epithelium and mesenchyme in a Turing-like mechanism (Salazar-

Ciudad, 2012). Tooth morphogenesis models and *in vivo* experiments have shown that

changes in the pathways controlling these loops can modify the number of cusps and

recapitulate evolutionary changes (Harjunmaa et al., 2012, 2014; Morita et al., 2020;

Salazar-Ciudad & Jernvall, 2010). More minimal modeling of Turing-like mechanisms in

teeth has shown how the interaction between activation-inhibition loops and growth of the

field dynamically determines the output pattern (Morita et al., 2022; Sadier et al., 2019) .

Applied to the case of supplementary lingual cusps, this theoretical framework predicts that

the new phenotype could be achieved by a change in the activation-inhibition loops (e.g.

allowing cusps forming closer from each other), a change in bucco-lingual (B/L) growth

(allowing more cusps to be fitted in a bigger field), or a combination of both.

Almost all what is known was established on the lower first molar of the mouse and few

studies have compared lower and upper molar development. The homeotic-code present

in the jaws at the early stages of development is still present when molars are initiated

(*Dlx1/2* and *Pou3f3* for upper jaw, *Dlx1/2/5/6* and *Nkx2.3* for lower jaw (Cobourne &

Sharpe, 2003; Hirschberger et al., 2021; Jeong et al., 2008). We have previously

demonstrated that later during morphogenesis, the expression of *Dlx5/6* genes is no longer

specific but remains biased (Pantalacci et al., 2017). Only three genes remain specific: in the

upper molar, *Pou3f3* and its non-coding regulator and in the lower molar, *Nkx2.3*. On top

of that, many genes are consistently biased throughout morphogenesis (Pantalacci et al.,

2017). The genetic architecture of lower and upper molars differs in mouse, since a few

mouse mutants have molar-specific phenotypes (including Dlx1/2 and Pitx1; reviewed in

(Hallikas et al., 2021; Kwon et al., 2017)), and molar specific loci are evidenced in the two

available quantitative genetic studies (Navarro & Murat Maga, 2018; Shimizu et al., 2004).

On top of working with this well characterized system, we took a comparative analysis of

RNA-seq timeseries, which proved successful to study development and its evolution. RNA-

seq temporal profiles describe the dynamic changes which orchestrate the development of

a complex structure, such as the many transcriptional and cell proportions changes

(Pantalacci & Sémon, 2015). In our published RNAseq timeseries comparing the

development of the mouse lower and upper first molar, we found that differences in their

transcriptomes were reflecting differences in their morphogenesis, such as different

proportions in the respective tissues of the tooth or different rates of cusp formation in the

tooth (Pantalacci et al., 2017). This showed that transcriptome timeseries contain valuable

information to compare the morphogenesis of two organs, even though this information is

not immediately noticeable. Comparing the transcriptomes of different species provides a

useful quantification of developmental similarity - once possible biases in estimating

expression levels have been controlled for (Cardoso-Moreira et al., 2019; Pantalacci &

Sémon, 2015). This has been successfully applied to answer questions on the periods of

maximal conservation of embryogenesis (so-called hourglass pattern, eg (Kalinka et al.,

2010; Levin et al., 2016)) or on the homology of organs (Fisher et al., 2020; Tschopp et al.,

2014; Z. Wang et al., 2011).

Here, to understand how lower and upper molars evolved independently from one another,

we compared the dynamics of their developmental systems in mouse and hamster with a

twofold strategy, analyzing 1) RNA-Seq time series and 2) the dynamics of cusp formation,

obtained by tracking a cusp marker in hundreds of molar samples. This revealed an

ancestral molecular identity for each molar type, associated with

morphogeneticspecificities. Consistent with the very peculiar mouse upper molar

morphology, the two mouse molars have more different temporal profiles than the two

hamster's. We found three morphogenetic changes which could combine to cause the

supplementary cusps in the upper molar, one of them building on the ancestral specificity

of upper molars. The biggest surprise came from the lower molar, which was first thought of

as an additional control. As many gene expression temporal profiles diverged in the lower

molar as in the upper molar, and a great part of them are co-evolving in the two molars.

This is associated with changes in B/L polarity and activation-inhibition mechanisms of cusp

formation seen in both molars. Based on these results, we propose that several mutations

combined to reach the new upper molar dental plan, some with specific developmental

effects, and others with effects in both molars. As a consequence, drift in lower molar

development went along with adaptation in upper molar development. We generalised our

findings by re-analysing transcriptome and literature data on bat foot and wing evolution.

We propose that mutations producing shared gene expression changes have a major

contribution to appendage-specific adaptation.

RESULTS

Mouse and hamster molars develop at different paces. We predicted developmental age

from embryonic weight in each species and aligned temporal series between species with

homologous start and end points of first molar morphogenesis (Figures 1 and S1). We then

devised a twofold strategy to compare the dynamics of cusp formation along with the

dynamics of gene expression. First, we established the sequence of PEK and SEKs

formation and modelled the relative stage durations with continuous Markov processes

(Figure 1B). Second, we obtained RNA-seq data from whole first molar germ at high time

resolution (Figure 1C) to model temporal profiles and added samples for dissociated

epithelium and mesenchyme as well as buccal and lingual half germs for specific purposes

(Figure 1C).

Conserved transcriptomic and morphogenetic features point to an

ancestral identity of lower and upper molars

Before focusing on the independent evolution of the first upper molar in mouse, we first

looked for molecular and developmental features which discriminate between lower and

upper molar development in the two species. This would form a molecular and

developmental identity of the molars common to both species, and likely present in their

common ancestor. It could have served as a basis for the independent evolution of the

upper molar in the mouse lineage.

Identity genes and many other key genes for tooth morphogenesis showed a conserved

expression bias that distinguish lower and upper molars in both species

Principal components in a PCA analysis separate samples according to the main axes of

variation in the data. In our dataset, the main effect was the species, followed by

development time (Figure 1C). Upper and lower molar samples were only separated from

each other on the sixth component, which carried 3% of the total variance. Hence only a

minor part of the variation distinguishes upper and lower samples while being common to

both species.

To detect genes carrying this conserved variation between upper and lower molars, we

modelled their temporal profiles with polynomials (Figure 2A). In each species, we fitted

two models: one with two distinct curves, one per tooth, and another with a single curve,

common for both teeth. By comparing the fit of these two models, we detected

differentially expressed genes between the two teeth (Table S2). In each species, we found

a similar proportion of differentially expressed genes (about 13%). There are 712 genes in

common between the two species (only a third of the genes in each species), including 550

genes with a consistent upper/lower bias.

A Set of genes with consistent upper/lower bias in both species

B Bucco-lingual difference in third cusp patterning

14532 genes

1 profiles up ≠ low

up ≠ low vs. up = low
Hamster
1917 genes

up ≠ low vs. up = low
Mouse
2088 genes

712 genes
+

2 consistent up/low bias

550 genes
+

3 profiles ham = mus

Upper
ham ≠ mus ham=mus

Lower
ham ≠ mus ham=mus

165 genes

C Difference in up/low expression peaks at stages 1-SEK and 2-SEK

D Transitory lingual phenotype

Fig 2 | Conserved differences of upper and lower molar development.

A. Pipeline to search for genes that distinguish lower and upper molars in the two species. 1/

Expression profiles were modelled separately in hamster (left) and mouse (right) to detect genes with

distinct profiles in upper and lower molars in each species. Colored curves are models allowing for

distinct profiles in upper and lower molars. Grey curves allow a single profile, common for both

teeth. The resulting number of differentially expressed genes is shown. 2/ Filtering for a consistent

The set of 550 genes with significant and consistent upper/lower bias in both species is

highly relevant: it contains the expected jaw-identity genes known for mouse (*Nkx2.3*,

Pou3f3, *Dlx1*, but also *Dlx5* and *6* which are no longer lower-jaw specific at this stage, but

show a lower molar bias as already described in mouse (Pantalacci et al., 2017)), a fifth of

the genes whose mutant shows a phenotype in the lower molar (21 out of 87 "keystone

genes" from (Hallikas et al., 2021)) and key transcriptional regulators of molar

morphogenesis (*Barx1*, *Msx1*, *Pitx1* Figure S2). Overall, these genes are strongly enriched

for transcriptional regulators. They are involved in epithelial and mesenchymal

development, cell adhesion, proliferation and differentiation and signalling, especially

WNT, BMP and NOTCH (enrichment for Gene Ontology terms, Figure S3). Among the 550

genes with a consistent upper/lower bias, only 165 display evolutionary conserved temporal

dynamics in both teeth (stars Figure S2, Table S2).

The earliest phase of cusp patterning gathers many features that distinguish lower and upper molars in both species

We next looked for criteria that would distinguish cusp patterning dynamics in lower and upper molar in both species, and may have been conserved from the common ancestor of mouse and hamster.

First, we noticed that in both species, SEK formation is initiated (transition to 1-SEK stage and 2-SEK stage) and completed (transition to 6-SEK stage) later in the upper molar as compared to the lower molar of the same embryo. This delay was also obvious in the transcriptome: the upper molar transcriptome looks "younger" than the lower molar transcriptome of the same embryo (Figure S4).

Second, we examined the details of the sequence of cusp acquisition. The first and second SEK are homologous in all teeth, the first SEK being buccal, and the second being its lingual neighbor. This bucco-lingual sequence is thus similar in lower and upper molars in both species, as previously shown in mouse (Cho et al., 2007; Pantalacci et al., 2017). The 3-SEK stage however distinguishes lower and upper molars in both species, with a buccal third SEK in upper molars, and a lingual third third SEK in lower molars (Figures 1B, 2B). It is striking that the three cups patterned at this 3-SEK stage recapitulate the bucco-lingual arrangement of their homologous cusps in the molars of early mammalian ancestors (Figures 2B, S5). These so-called "tribosphenic molars" were markedly asymmetric along the B/L axis (B. M. Davis, 2011; Hillson, 2005). On top of this geometric pattern, we found two other conserved distinctive features: the dynamics of the three first SEK stages (Figure 2B and figure S5), and a peculiar lingual epithelial bulge specific to the upper molar (Figure 2D). Both occur at this same early period of development. In fact this is the period where lower and upper molar transcriptomes differ most from one another, as seen on a multivariate analysis (BCA) performed on all samples to capture a lower/upper molar variation common to both species (Figure 2C).

Together, we found a clear conserved transcriptomic identity of each molar in the form of a

conserved expression bias for identity genes and many key regulators of tooth

development, and a conserved transcriptomic signature. A clear conserved morphogenetic

identity was obvious in the earliest phase of cusp formation, with different dynamics of cusp

formation along the bucco-lingual axis, that recapitulated the bucco-lingual specificities of

early mammals' molars. We next asked how the mouse upper molar evolved its new

morphology.

Increased upper-lower molar dissimilarity of mouse transcriptomes

Some transcription factors which made the ancestral identity showed increased levels and/or

biases in mouse molars

Because the dose of identity transcription factors was shown to correlate with the degree of

appendage differentiation, we first first asked whether the dose of transcription factors

forming the conserved transcriptomic identity has changed in mouse (Figure 3A). We

expected to observe an increased bias in mouse, especially in the upper molar.

A. Transcription factors with increased bias in mouse

B. Increased transcriptomic dissimilarity in mouse

Fig 3 | Increased upper-lower molar dissimilarity of mouse transcriptomes.

A. Temporal profiles for 6 key transcription factors, distinct curves were fit for upper (black) and lower

(grey) molars.

B Left: Model to detect genes which differ in their upper-lower expression profiles in one species.

Example of *Sfrp2* gene expression levels (grey dots), significantly better modelled by two curves in

mouse and one in hamster (purple curves), compared with one curve per species (grey curves). Right:

Number of genes detected by this tooth-specific model in mouse (purple) and in hamster (green, LRT

with adjusted p < 0.05). Barplots show their frequencies for the "total" gene set, genes from

developmental "pathways", genes from "bite-it" database, and genes with a mild or strong

phenotype in tooth mutants ("Dispensable" or "Keystone"). Size of each gene set into brackets.

Pou3f3 is the only TF specific of the upper molar in both species. Its expression showed a

twofold increase in mouse upper molar. *Dlx1/2* genes are expressed in both molars, but are

essential only for upper molar formation in mouse (Qiu et al., 1997). Their expression levels

were more than twice increased in mouse molars and the ratio, slightly in favor of the upper

molar in hamster, is increased in mouse. *Barx1* is a key molar-specific TF whose levels have

been correlated with cusp number in mammalian molars (Miletich et al., 2011). The bias in

favor of the upper molar was markedly increased in mouse. This effect is selective since

Msx1, another TF which cooperates with *Barx1 (Miletich et al., 2011)*, showed similar bias in

the two species (Figure S2).

Surprisingly, such changes were not restricted to the upper molar. The expression of

Nkx2.3, the specific TF of the lower molar, showed an almost twofold increase in mouse.

For *Pitx1*, a shared TF whose mutation impairs more specifically lower molar development

(Mitsiadis & Drouin, 2008), the ancestral bias in favor of the lower molar was increased. *Dlx1*

expression levels were also twice increased in the lower molar. This is not true for the *Dlx5-*

6 genes, which were more specifically associated with lower jaw identity (Depew et al.,

2005), but are expressed at this stage in the two molars (Pantalacci et al., 2017) (Figure S2).

Thus, the molecular identity of each molar was reinforced in mouse, partly in line with an

ancestral bias.

<u>The dissimilarity of upper/lower molar transcriptomes is increased in mouse</u>

Next, we asked whether the transcriptomes would capture a general increase of

differentiation of molar development in mouse as compared to hamster.

As exemplified above, a number of genes have a marginally or small significant bias in

hamster, which is increased in mouse. This is in agreement with the multivariate analysis

presented in Figure 2C. Along an axis that captures a lower/upper molar variation common

to both species, the variation is higher for mouse than for hamster. Thus, an ancestral

dissimilarity is exaggerated in mouse.

We detected genes whose expression profiles differ between upper and lower molars in

mouse but not in hamster, by a dedicated model based on the 4 molars altogether (Figure

3B). As a control, we built a reciprocal model to detect genes with tooth specific profiles in

the hamster. We found 2.5 times more genes with tooth-specific profiles in mouse as

compared to hamster. Even after removing the effect of baseline expression levels, which

may be impacted by differences in cell composition (Pantalacci et al., 2017), we still

observed 1.6 times more tooth-specific profiles in mouse. The function of the genes with

tooth specific profiles differs markedly between mouse and hamster (Figure S6). In mouse,

genes are linked to cell adhesion and migration, as well as cell cycle and mitosis. This is

consistent with our previous findings suggesting that morphogenetic movements are

enhanced in mouse upper molar during the period of lingual growth (Pantalacci et al.,

2017).

Thus, we quantify an increased dissimilarity of temporal profiles in mouse, consistent with

the increased morphological dissimilarity of the adult teeth. This involved the reinforcement

of an ancestral molecular identity as well as newly evolved gene expression differences.

Morphogenetic changes associated with mouse upper molar evolution

Next, we used the transcriptome as a starting point to investigate several possible

developmental mechanisms how the mouse upper molar forms additional lingual cusps. We

logically focused on specificities of mouse upper molar development as compared to any

other teeth.

The proportion of mesenchyme is increased in the mouse upper molar

We previously showed that the upper molar germ of the mouse contains more

mesenchyme relative to the epithelium than the lower germ since early cap stage. Tooth

engineering studies suggest that this higher proportion may help to form the

supplementary cusps. Indeed, in artificial teeth made by reassociating a varying amount of

mesenchymal cells to a single epithelium, the number of cusps formed increases with the

number of mesenchymal cells (Hu et al., 2006). To control whether this higher

mesenchyme:epithelium ratio is specific to the mouse, we extracted mesenchyme and

epithelium-specific marker genes from tissue-specific transcriptomes (Figure 1C), and used

in silico deconvolution to estimate the mesenchyme proportions from whole tooth germ

transcriptomes (Figure 4A). The proportion of mesenchyme was indeed significantly higher

in the upper molar in mouse, but not in hamster (Wilcoxon tests, $p < 2e-16$ and $p = 0.152$).

As shown in a previous publication (Pantalacci et al., 2017), and suggested above, this

change in tissue proportion should inflate the transcriptomic differences between mouse

and hamster.

Fig 4 | Morphogenetic changes specific to the mouse upper molar.

A. Percentage of mesenchymal tissue in tooth germs estimated by deconvolution of the RNAseq

time series with tissue-specific marker genes. Time scale in relative 0-10 scale (x axis) for each

species. Color code indicated for each molar.

B. Relative duration of the first morphological stages highlights a longer period of lingual growth in

mouse upper molar (1-SEK, purple). Stage durations from Figure 1B. PEK and SEK: primary and

secondary enamel knots respectively.

C. Levels of activation of BMP, SHH and WNT pathways in buccal and lingual sides of the mouse

molars at 15.0 dpc (1 SEK stage). Measurements made with an *in silico* method, ROMA, which

compares pathway activity in transcriptomic samples based on a list of targets for the pathway. Two

<u>A bucco-lingual polarity is maintained in mouse molars during the first steps of cusp</u>

<u>formation, and further enhanced in the upper molar</u>

The supplementary cusps of the upper molar form last, on the lingual side of the tooth

(Figure 1B), but the dynamics of SEK formation differs much earlier between mouse and

hamster upper molars. Indeed, the 1-SEK stage is longer in the mouse upper molar than in

any other tooth (likelihood ratio test, $p < 1e-16$, Figure 4B). At that stage, the tooth germ

grows rapidly on the lingual side. This finding prompted us to look into changes of the

bucco-lingual development in the mouse upper molar.

We do not know the molecular mechanisms whereby the first cusp forms on the buccal side

and a second one forms on its lingual side. However, we know the mechanisms which

decide if a first tooth is formed, and whether a second tooth can form lingually (Jia et al.,

2013, 2016; Lan et al., 2014; Zhang et al., 2009). To form a tooth, the WNT pathway must

be activated on the buccal side of the mouse jaw, while on the lingual side, WNT pathway

activation is prevented by OSR2, notably through the WNT inhibitor SFRP2 (Jia et al., 2016).

This bucco-lingual polarity of the jaw is set up by a mutual antagonism between BMP4

activity on the buccal side and OSR2 activity on the lingual side. In mice, displacing this

BMP4/OSR2 balance can suppress tooth formation (following loss of BMP4) or induce the

formation of a supplementary tooth on the lingual side (following the loss of OSR2 or of the

Wnt inhibitors) (Jia et al., 2013, 2016; Lan et al., 2014; Zhang et al., 2009)). Since the

mechanisms for tooth (PEK) formation are largely re-used for cusps (SEK) (Jernvall &

Thesleff, 2012), it would be unsurprising that this BMP4/OSR2 balance also controls the B/L

polarity of cusp formation.

516 To look for evidence of persistent molecular B/L polarity during cusp morphogenesis, we

517 collected and analyzed mouse transcriptomes of buccal and lingual halves at the early 1-

518 SEK stage (Figure 1C). We found that *Osr2* and *Sfrp2* are still expressed with a strong

519 lingual bias at 1-SEK stage, and in the timeseries, their expression is maintained at high

520 levels during the period of bucco-lingual development of the tooth germ (Figure S7). To

521 determine if the tooth germ is polarized at the 1-SEK stage, we estimated the levels of

522 activation of 3 pathways controlling cusp formation (BMP, WNT and SHH), in the buccal and

523 lingual halves separately. We used ROMA, a method that exploits the level of expression of

524 up and down-regulated transcriptional targets to quantify signaling pathway activity directly

525 from the transcriptomes (Martignetti et al., 2016). The list of WNT and BMP target genes

526 was specifically established in tooth epithelium and mesenchyme by others (O'Connell et al.

527 2012). WNT, BMP4 and SHH pathways are strongly activated on the buccal halves of both

528 molars, but very weakly on the lingual halves, which therefore still appears as a naive tissue

529 (Figure 4C). In the upper molar, the lingual half looked even more naive than in the lower

530 molar, with even lower levels of pathway activities, consistent with twice higher levels of

531 *Sfrp2* expression, and delayed downregulation (Figures 4C, S2, S3). In the buccal half,

532 activation of the BMP4 and WNT pathways in the mesenchyme is stronger in the upper

533 molar, which thus appears as more polarized than the lower molar. These findings support

534 the idea that the BMP4/OSR2 antagonism is still acting during early mouse molar

535 morphogenesis to set up the B/L polarity of the tooth. This polarity maintains naive tissue

536 on the lingual side of the germ at 15.0, which grows faster and shows delayed cusp

537 formation relative to the buccal side.

539 We reasoned that the larger the proportion of this naive lingual tissue, the stronger the

540 germ growth potential and its capacity to form cusps on its lingual side. We therefore

541 quantified this proportion in mouse and hamster tooth germs, by deconvoluting the

542 timeseries dataset with buccal and lingual tissue transcriptomes (Gong & Szustakowski,

543 2013). As expected due to progressive cusp formation, we found that the proportion of

544 naive lingual tissue decreases during morphogenesis in both species (Figure 4D). But in

545 mouse molars, and even more markedly in the mouse upper molar, the initial proportion of

naive tissue is larger, and diminishes more slowly. We noted that the naive tissue proportion

correlates with levels and temporal pattern of *Sfrp2* expression in mouse and hamster

molars (Figure S7).

In summary, a B/L polarity of the tooth germ is maintained during the first steps of cusp

formation. This is especially true in mouse molars whose proportion of lingual naive tissue is

increased, and correlates with higher levels of *Sfrp2* expression. This polarization is further

exaggerated in the mouse upper molar.

<u>A change in the BMP pathway may underlie the maintenance of the B/L polarity in the</u>

<u>mouse molars</u>

The findings above suggest a change in the BMP4/OSR2 balance in mouse molars, with an

exaggeration in the mouse upper molar. We came back to the list of genes with mouse

upper specific profiles (Figure 3B) and compared it with the list of genes with a marked

bucco-lingual bias. The *Bmper* gene is a good candidate since it is a regulator of the BMP4

pathway in many tissues, and it ranked well in both lists (respectively 211/14532 and

9/12008). In mouse early 1-SEK stage molars, *Bmper* is more expressed on the buccal side

in both molars and more expressed in the upper molars on both buccal and lingual sides. In

the time series, *Bmper* expression levels are higher in mouse than in hamster. It decreases

with time in all teeth, but this decrease is slower in the mouse upper molar (Figure 5A). This

is similar to the levels and dynamics observed for the proportion of naive lingual tissue, or

the expression of *Sfrp2* gene (Figures 2C, S2). This pattern suggested us that *Bmper*

participates in the regulation of the B/L polarity.

In situ hybridizations showed that in mouse molars, *Bmper* rapidly withdraws from the lingual side to remain strongly expressed in a small buccal domain only. This process is delayed in the upper molar. In the hamster, the withdrawal is symmetrical and similar in both teeth (Figures 5B, S6). Thus *Bmper* expression changed at two levels. First, it acquired a new bucco-lingual regulation leading to a strong buccal expression and an early withdrawal from the lingual side. This co-evolved in both teeth. Second, it acquired a new lower-upper molar difference, with delayed withdrawal in the upper molar.

To gain insight into the function of *Bmper*, we obtained a mouse null mutant and studied

the shape of its molars. Since the homozygous *Bmper* mutant are lethal at birth, this had to

be done by reconstructing the enamel-dentin junction at 19.5 days, carefully matching them

with controls of similar developmental age (see material and methods). The upper molar is

modified: one of the buccal cusp is absent or poorly grown (Figure 5C, arrowheads) and

one lingual supplementary cusp is more prominent (arrows). The lower molar is also

modified: the central lingual cusp, which is determined by formation of the 2-SEK following

lingual growth, is more prominent. Thus, the loss of *Bmper* modifies the bucco-lingual

equilibrium, favoring the lingual side of the molar at the expense of the buccal side. Since

Bmper modulates the BMP4 pathway, this mutant phenotype reinforces the idea that the

BMP4 pathway regulates B/L polarity during cusp formation, and that Bmper might have a

causative role in displacing the BMP4/OSR2 balance in mouse.

In summary, we show that the BMP4/OSR2 antagonism persists in mouse to regulate the

B/L polarity of the tooth during morphogenesis. As compared to hamster upper molar, the

mouse upper molar has an asymmetrical expression of *Bmper* (buccal side) and an

increased expression of *Sfrp2* (lingual side). This is associated with an increased and

persistent proportion of naïve lingual tissue. These differences seem very consistent with

the newly evolved lingual cusps of this tooth. We were very intrigued however that

qualitatively, these morphogenetic changes are also seen in the lower molar, although to a

lesser extent and without change in number and respective size of buccal/lingual cusps. We

next wanted to quantify this concerted developmental evolution of the lower molar with the

upper molar.

Concerted evolution of lower molar development with upper molar development

Developmental gene expression in the lower molar largely evolve in a concerted manner with the upper molar

We modelled the temporal profiles of the 4 molars altogether to quantify their concerted evolution. We fitted four models (Figure 6A): The most complex model has four curves (one distinct per tooth), intermediate models have two curves (distinguishing species: hamster/mouse or distinguishing tooth: upper/lower), and the most simple model has a single curve common to all teeth (1 curve). Models for different species account for different baseline expression levels, to make sure that we focus on species differences in temporal dynamics. We attributed the best model to each gene and from this we built an index of coevolution. We estimated that the expression profiles of 61% genes have coevolved. This is consistent with a cruder estimate from the PCA analysis, where the main axis of variation in the transcriptomes separates samples by species, but groups upper and lower molars (Figure 1D).

Fig 6 | Divergence and coevolution of expression profiles.

A. Nested models of temporal profiles taking the four teeth altogether. The percentage of

coevolution is computed as the proportion of "divergent" genes, among genes varying between

species and/or teeth.

B-C. *Bmp4* and *Wif1*'s expression profiles corroborated by *in situ* hybridization of dental

mesenchyme show that expression in upper and lower molar has coevolved. Dashed lines and

numbers map pictures to the timeseries. See Figure S9 for details.

D. Temporal profiles modelled by tooth type. *Dlx1* is shown with samples (grey dots), and models

(curves). Top: the "upper divergent" model, allowing different profiles in mouse and hamster (green),

is compared with the "upper non-divergent" fitting the same profile but different baseline

The profiles of many genes important for tooth development have coevolved, suggesting

that developmental processes have largely evolved in a concerted manner in the two teeth,

as seen earlier for the B/L axis. We decided to examine by *in situ* hybridization some of

those genes whose profile co-evolved in the transcriptomes. We wished to determine what

spatio-temporal profiles are behind this transcriptomic co-evolution, and which other

developmental processes besides B/L polarity may have co-evolved between the two teeth.

<u>The early dynamics of cusp formation show concerted evolution, with anticipated cusp</u>

<u>formation in mouse.</u>

We first examined *Bmp4*, since finding this essential gene for tooth development among

co-evolving genes was a surprise (Figure 6B). *Bmp4* is expressed in both the epithelium and

the mesenchyme, but the epithelial domain is so small relative to the mesenchymal domain

(see later Figure 7 and S9), that the latter will dictate the bulk transcriptomic profile. We

therefore first looked at the *Bmp4* expression in the mesenchyme. *Bmp4* reached a spatially

homogeneous expression earlier in the mouse mesenchyme (Figure 6B), consistent with an

earlier peak of expression in mouse transcriptomes.

We picked up two other mesenchymally expressed genes for *in situ* hybridization, *Wif1* and

Dkk1, because 1) these genes are likely involved in the gene regulatory network of tooth

formation: they are known modulators of the Wnt pathway which is critical for cusp

formation, and their expression is changed when the Bmp4 and/or Wnt pathways are

manipulated (data by O'Connell et al.) and 2) their temporal profiles markedly differ from

each other, as well as from *Bmp4.*

Wif1 rises earlier in mouse transcriptomes, and its mesenchymal expression is seen both

earlier and in a larger territory in mouse tooth germs (Figures 6C, S9). *Dkk1* expression

transiently decreases in mouse transcriptomes, which coincides with an earlier relocalisation

of its expression at future cusp tips beneath the SEK (Figure S9). This expression pattern

suggested to us that cusp formation might in fact be anticipated in mouse as compared to

hamster.

We thus turned back to our quantification of cusp formation dynamics, and realized that

both mouse molars quickly transition to 1-SEK after a rather short PEK stage (Figures 1, 7A).

By comparing epitheliums of the two species matched for growth advancement, we found

that both mouse molars already exhibit the rounded and focalized *Fgf4* expression typical

of a SEK when hamster's still exhibit the large and elongated *Fgf4* expression typical of the

PEK (Figure 7B stage 2). Thus, mouse PEK is rapidly turned into a precocious and focalized

1-SEK, marking an early beginning of cusp patterning in a very young cap stage, and this

happens very similarly for both mouse molars. The dynamics of activation-inhibition

mechanisms relative to the advancement of epithelial growth thus evolved in a concerted

manner in the two teeth. We have shown previously how the transcriptome carries

signatures of ongoing developmental processes, including cusp formation (Pantalacci et al.,

2017). We suspect that well beyond the few genes studied above, the transcriptome could

be deeply shaped by this anticipated formation of cusps in the young mouse molars, and

thereby be markedly different from hamster transcriptomes. Said differently, this

morphogenetic change could contribute to a substantial part of the transcriptomic

concerted evolution.

A.

B. small-range signalling centers in mouse

C. precocious transition to cusp morphogenesis in mouse

 Fig 7 | Activation-inhibition mechanisms co-evolved in mouse molars.

A. Duration of each stage estimated by Markov models from Figure 1B. Numbers in

triangles as in B. Timeline and cartoons on the right recapitulate changes in signalling

centres.

B. Mouse molars transition earlier to cusp patterning. Transition from the PEK to the 2-SEK

stage as seen on tooth germ epithelial parts hybridised against *Fgf4*. Pairs of

mouse/hamster embryos were selected to show four remarkable steps in this chronology (1-

4 in triangles). At stage 2, *Fgf4* expression is still elongated in hamster, as typical for a PEK,

while it is already roundish in mouse, as typical for a SEK.

C. Expression of *Bmp4* is more focalized in mouse than in hamster SEKs. Mouse and

hamster samples are paired for similar advancement of epithelial growth. Age of samples in

days is in the upper molar picture (for samples taken from the same embryo) or in both

pictures (samples taken from different embryos). See also Figure S10.

<u>Spatial aspects of the activation-inhibition mechanisms controlling cusp formation also show</u>

<u>concerted evolution, with more local inhibition in mouse.</u>

To get further insight into the evolution of activation-inhibition mechanisms in hamster and

mouse, we next studied the expression of diffusing signals, which are produced in the SEK

and inhibit the formation of other SEKs in the vicinity. Indeed, evolutionary changes in the

production and diffusion of these signals are thought to drive evolutionary changes in cusp

number (Salazar-Ciudad & Jernvall, 2010). We do not know how much each of these

molecules diffuses, but the spatial range of expression of each gene can be considered as a

minimal range for its inhibitory action on cusp formation. We studied two known inhibitors

of SEK formation, *Bmp4* (Meguro et al., 2019) and *Shh* (Kim et al., 2019). We also studied

Wif1, which we consider a likely inhibitor since it antagonizes the Wnt pathway, whose

activation in the epithelium promotes cusp formation (Järvinen et al., 2006; Liu et al., 2008).

Bmp4 and *Wif1* are expressed from the signalling centers : their expression pattern is much

more narrow and roundish in mouse than in hamster (both in PEK and SEKs) (Figures 7B and

S10). *Shh,* which more largely marks cells committed to cusp formation, also shows a more

restricted expression in mouse (Figure S10). Hence for the three pathways involved in the

activation-inhibition mechanisms, inhibition is more local in both mouse molars.

In conclusion, both dynamics and spatiality of activation-inhibition mechanisms have

evolved in a concerted manner in the molars of the two species. Both of them, rapid switch

to SEK formation relative to epithelial growth and a more local inhibition, are predicted to

favor the formation of more cusps. Features that make sense for the formation of the

supplementary cusps in the upper molar are thus observed also in the lower molar, as seen

above for the B/L polarity.

Developmental gene expression diverged as much in lower as in upper molar

These developmental phenotypes of the lower molar have evolved in concert with the

upper molar, but this evolution did not drive a major phenotypic change. Such discrepancy

between the divergence of development and the conservation of the final phenotype, is a

phenomenon known as Developmental System Drift (DSD). To measure the extent of this

phenomenon, we decided to compare levels of developmental evolution in both teeth.

Since the lower molar phenotype has been much more conserved during evolution, the

lower molar developmental phenotype captured by the temporal profiles should be more

conserved. Otherwise, this is an indication of DSD.

We scored the divergence between mouse and hamster upper and lower molars by

modelling temporal profiles with polynomials (LRT with adjusted $p < 0.05$). We found that

for 22.0% of genes, the profiles diverged in the lower molar, which is even more than in the

upper molar (17.3%, Figure 6D, Table S2). This is true as well for genes relevant for tooth

development and phenotype ("bite-it", "keystone", "pathways"; Figure 6E).

Put together, these observations suggest that the development of the lower molar has

drifted while co-evolving with the upper molar.

Bat limbs development show a similar pattern of concerted evolution

In order to generalize our results, we turned to another case of drastic independent

evolution: the bat limb. The evolution of the wing relied on drastic changes in the forelimb

development, including changes in digit patterning, growth, and webbing to form the wing

membrane. In comparison, the bat hindlimb kept a morphology more typical of

quadrupedal species, as did both mouse limbs. This provides a framework of independent

evolution allowing us to test the generality of our findings beyond mouse molars.

We collected raw sequencing data from a previous study comprising 3 stages of mouse and

bat fore/hindlimb development (Maier et al. 2017). We quantified expression levels and

classified temporal profiles with polynomial models dedicated to measure coevolution (as in

Figure 6A).

Just like in our molar dataset and consistent with the original analysis of this dataset (Maier

et al., 2017), the genes which have a limb-specific temporal profile and which have kept it in

mouse and bat are a minority (53 genes), but they are highly enriched for transcription

factors, including all the expected identity genes, eg. *Tbx4* and *Tbx5*, *Pitx1*, as well as 6

biased Hox genes.

The profiles of 714 genes differed both between species and limbs. The profiles of almost

four times more genes (2677) diverged between the two species, but co-evolved in the two

limbs, despite their drastic morphological differences. Such a large proportion of co-

evolving genes mirrors our finding in rodent molars. Importantly, genes with a well-

established role in controlling limb morphology co-evolved. It is the case of key genes

controlling limb patterning (*Shh*, *Fgf10*, *Fgf8*, *Grem1*...) and chondrogenesis (*Wnt3* and the

Activin pathway: *Inhba*, *Inhbb*, *Acvr2b*…). It is also the case of most of the genes known to

regulate webbing (*Fgf8*, *Grem1*, *Bmp7*, *Ihh,* Retinoic acid pathway: *Aldh1a2*, *Cyp26b1*).

We note that several of these genes have been pointed in the literature as key for bat wing

evolution. For three of them, we could compare the expression profile in the transcriptomic

dataset with published *in situ* hybridization in both limbs, and they were consistent with co-

evolution. The iconic *Shh* gene expression clearly peaks at the second stage in both bat

limbs, but not in mouse limbs (Figure S11), and peaking is exaggerated in the bat forelimb

(Figure S11). This is consistent with figure 2 in (Hockman et al., 2008). The new temporal

profiles of *Fgf8* and *Grem1* in both bat limbs are also consistent with a previous study,

which has shown the novel expression domain of these genes in both limbs (Weatherbee et

al., 2006).

As in mouse molars, co-evolution is pervasive in bats limbs and concerns genes whose

expression evolution was key for the independent phenotypic evolution of the forelimb.

DISCUSSION

Below we discuss how the independent phenotypic evolution of the mouse upper molar

involved reinforcing and building on ancestral specificities of the upper molar development,

in relation with identity genes. It was accompanied by extensive evolution of lower molar

development, including concerted evolution with upper molar development, which

contrasts with the limited phenotypic evolution in this tooth. These findings are best

understood in a model where developmental system evolution of the upper molar induced

developmental system drift in the lower molar.

<u>Conserved specificities of lower and upper molar morphogenesis may date back to early</u>

<u>mammals</u>

We found several conserved specificities which discriminate between lower and upper

molars. All mark the early period of cusp formation: the arrangement of cusps at 3-SEK

stage and the early dynamics of cusp formation, the morphology of lingual epithelium and a

transient reinforcement of transcriptomic identity (already seen in mouse in our previous

study (Pantalacci et al., 2017)). Since these findings were made in hamster and mouse,

these specificities of lower and upper molar were likely present in their common ancestor,

but we suspected they may even date from early mammals.

Early mammals evolved "tribosphenic molars", a major innovation of lower and

upper molar shape which enabled unprecedented occlusion (B. M. Davis, 2011; Hillson,

2005). For the first time in the reptilian evolution, lower and upper molars were developing

into drastically different shapes. In the Figure S5, we discuss in detail how the

developmental specificities of lower and upper molars of mouse and hamster strikingly

mirror the specificities of the lower and upper tribosphenic molars, taking into consideration

known homologies in mammalian molar cusps. In particular, we show how the spatio-

temporal dynamics of cusp formation in the lingual and posterior directions are combined

differently in the two molars. We propose that evolving a jaw-specific control of this

combination was the key developmental innovation underlying the invention of the

tribosphenic molars.

Most later mammals had less dissimilar teeth, such as the common ancestor of

mouse and hamster or the present golden hamster. Yet the heritage of the mammalian

innovation remains visible in the transient developmental dynamics of lower and upper

molars. This also constitutes a case of "recapitulation" since early ontogeny of cusp

formation recapitulates phylogeny (Gould, 1977).

These hidden developmental specificities could serve as a basis for the independent

evolution of upper molar in the mouse lineage. It is interesting that some fossil rodents

close to mouse/hamster common ancestor carried a crest along the lingual basis of the

upper, but not the lower molar, which may be seen as an ancestral predisposition to

enlarge lingually and form lingual cusps (Charles et al., 2009; Tiphaine et al., 2013).

<u>What does underlie the conserved morphogenetic identity of molars? Ancestral molecular</u>

<u>identity of molars</u>

We identified several conserved specificities of upper and lower molars at the

developmental system level, yet such conserved specificities remain rather discrete at the

transcriptomic level. We found relatively few genes consistently biased in the two species,

and their temporal profiles were most often not conserved, except some highly relevant

transcription factors. This includes the two jaw-specific genes *Nkx2-3* and *Pou3f3,* and

several *Dlx* genes. The *Dlx* genes are homeobox transcription factors which specify jaw

identity at early stages of craniofacial development in jawed vertebrates and might have

been implicated in the transition from a reptilian to a mammalian jaw (Depew et al., 2005;

Gillis et al., 2013). The dose as well as the complement of *Dlx* genes (*Dlx1/2* in upper jaw;

Dlx1/2/5/6 in the lower jaw) are important for normal jaw development in mouse (Depew et

al., 2005), and upper molars fail to develop without *Dlx1/2 (Qiu et al., 1997). Dlx1/2*

showed an upper-bias in both species, and Dlx5/6 a lower bias. Transcription factors known

to be essential to tooth development, such as *Msx1*, *Barx1*, *Pitx1* also showed a conserved

bias. It remains to be tested if this bias is directly controlled by identity genes.

Our results show that during evolution, the details of developmental interactions in serial

organs diverge extensively, but some developmental specificities of one organ with respect

to the other are conserved (e.g. the relative order and timing of appearance of the 3 first

cusps and the period of maximal transcriptomic divergence, the delayed development of

the upper molar). These specificities could be encoded in a conserved relative dose of the

key transcription factors specifying an organ (here a molar or a limb), and this conserved

relative dose could be controlled by identity genes. Altogether this forms an ancestral

molecular and developmental identity for the two teeth.

<u>Reinforcement of the ancestral molecular and morphogenetic identity in mouse molars</u>

We found that the molecular identity was reinforced in mouse, in the upper molar but also

more surprisingly in lower molar. Expression levels doubled for the upper-molar specific TF

Pou3f3 and the lower-molar specific TF *Nkx2-3*, and the ancestral bias of many genes was

exaggerated, whether in favor of upper or lower molar (e.g. *Barx1* and *Dlx1*; *Pitx1*,

respectively). Consistent with these changes in individual TFs, we found at genome-wide

scale that the temporal profiles in the mouse were exaggeratedly different from each other.

As for specific TF, divergence is seen both in upper and lower molar.

<u>Mouse upper molar displays three morphogenetic changes favoring supplementary cusp</u>

<u>formation, one is building on ancestral upper molar specificities</u>

The supplementary cusps form last, the most anterior one being the very last, just as

it happens in the fossil record (Lazzari et al., 2008; Tiphaine et al., 2013). This

"recapitulation" looks superficially like a case of "terminal addition", a mechanism for

evolutionary change whereby development is incremented with one more step (Gould,

1977). However, species divergence peaks early in development and we point to three

features that concern mouse early upper molar development, but could pave the way for

the supplementary cusps (Figure 8).

 Fig 8 | A summary of findings and working model from this study. For mouse molars and bat limbs,

 the "development" box shows key developmental stages with the time period covered by

 transcriptome data. The box "morphological evolution" represents the drastic morphological

 changes of the upper molar and forelimb as compared to the relative conservation of the lower

 molar and hindlimb. The "development evolution" box summarises the evolution of transcriptome

 and developmental mechanisms. Transcriptomes are dominated by co-evolution, with rare

 conservation of organ-specific expression besides identity genes. Organ-specific, shared, and shared

 but exaggerated developmental changes combine to achieve the organ-specific morphological

 change. For each change, the ancestral and derived state are represented, and candidate genes are

 indicated. In mouse molars, 3 changes (increased mesenchyme size, increased bucco-lingual

 asymmetry and smaller inhibitory signalling centres) combine to induce extra cusps on the lingual

 side of the upper molar only. In bat limbs, we took the example of early changes in AER size (apical

 ectodermal ridge, a signalling center) leading to altered digit patterning and late expression changes

 in *Fgf8* and *Grem1*, that efficiently combine to suppress interdigital apoptosis in the forelimb only. In

 our working model, shared changes are necessary but not sufficient. Combining them with more

 specific changes (e.g. mesenchyme size) and/or ancestral specific features (e.g. ancestral difference

 in molar Bucco/lingual axis) is necessary to achieve the morphological change. Related to Figure

 S12-13.

      The first feature is the larger mesenchyme compartment of the mouse upper molar.

 Increasing mesenchyme proportion increases cusp number in tooth engineering studies (Hu

 et al., 2006), probably because the mesenchyme promotes epithelial growth, which

enlarges the field where activation-inhibition mechanisms act and pattern SEKs. The

observed difference seems however too modest to drive the formation of supplementary

cusps on its own and does not explain why the supplementary cusps would form on the

lingual side only.

The second feature is the stronger polarisation of the upper molar field along the

bucco-lingual axis, associated with a precocious transition from PEK to 1-SEK and a very

long 1-SEK stage. As a consequence, a larger undifferentiated field is present on the lingual

side, where activation-inhibition loops can pattern SEKs. This feature seems highly relevant

because it could explain why the increase in cusp number is focused on the lingual side of

the tooth, while the size of buccal cusps is reduced. It seems to exploit an ancestral

specificity of the upper molar as compared to the lower molar, which produces a longer 1-

SEK stage. This specificity could combine together with a novelty in mouse responsible for

shortening the PEK stage in both teeth, and produce an upper molar with a very long 1-SEK

stage and a large undifferentiated field, while change remains more modest in the lower

molar.

The third feature is the narrower range of expression of signaling molecules in

mouse signalling centres, which is especially obvious for *Bmp4*. Reducing the range of

these known cusp formation inhibitors should allow to squeeze more SEKs in an equivalent

field. In agreement with this idea, a mouse mutant where *Bmp4* is overexpressed in all the

epithelium loses the supplementary cusps (Meguro et al., 2019), hence reverting to the

ancestral phenotype.

Molecular mechanisms and candidate genes for the observed morphogenetic changes

Mapping mutations corresponding to these developmental phenotypes is out of the

scope of this study, but transcriptomics provided us with molecular mechanisms and some

candidate genes. No expression change was observed in two obvious candidates from the

literature (*Fgf3* and *Activin A*, Figure S12, (Charles et al., 2009; Kwon et al., 2017).

In insects, the evolution of the dose of identity genes has been correlated with the

evolution of the size of serial organs (Paul et al., 2021). *Pou3f3* is expressed specifically in

the upper molar mesenchyme of both species and its dose is twice increased in mouse.

Hence *Pou3f3* is a good candidate to explain the larger proportion of mesenchyme

specifically in the mouse upper molar. *Pou3f3* loss-of-function mutants miss some skeletal

elements of the upper jaw, but their upper molars showed "no major defects" (data

unshown in (Jeong et al., 2008)). This may deserve re-examination, or study in sensitized

backgrounds. The causative mutation may also be upstream in the regulatory network, in

particular in *Dlx1/2*, since *Pou3f3* is regulated by *Dlx1*/2 in the early jaw (Jeong et al., 2008)

and the dose of *Dlx1/2* genes are also twofold increased. Both *Dlx1/2* and *Pou3f3* are also

involved in cranio-facial development. Since mastication has changed together with tooth

morphology in the mouse lineage (Lazzari et al., 2008; Tiphaine et al., 2013), changes in the

dose of these genes could have had pleiotropic effects beyond the molar.

The reinforcement of the B/L polarity in the mouse upper molar likely involved

changes in or upstream of the BMP4/OSR2 network. Interestingly, mutations in this network

have very different consequences on lower and upper molar development of mouse (eg. a

tooth is normally formed *versus* arrested at a very early stage) (Jia et al., 2013, 2016; Kwon

et al., 2017; Lan et al., 2014). We show that the BMP4/OSR2 antagonism, which earlier in

development regulates the B/L polarization of the molar-forming region, persists in mouse

during the first steps of cusp formation. We found expression changes in two genes which

should modify the output of this BMP4/Ors2 network: *Sfrp2*, whose role in the network was

known in mouse (Jia et al., 2013, 2016; Kwon et al., 2017; Lan et al., 2014) and *Bmper*,

whose role in tooth development was unknown.

BMPER is a known modulator of the BMP4 pathway, which seems to be pro- or anti-

BMP4 in different contexts (Correns et al., 2021; Ikeya et al., 2010; Kelley et al., 2009; Serpe

et al., 2008). Because the molars of the *Bmper* loss-of-function mutant are "hypo-

buccal/hyper-lingual", we deduce that *Bmper* is normally pro-BMP4 in the mouse

BMP4/OSR2 balance. In mouse, the strong buccal *Bmper* expression should favor BMP4

activity on the buccal side while the early lingual withdrawal should decrease BMP4 activity

on the lingual side. This sharper gradient of BMP4/OSR2 antagonism may link two

observations in mouse: on the buccal side, the earlier PEK/SEK transition; on the lingual

side, the larger undifferentiated field. Finally, we note that the teeth of the *Bmper* mutant are strikingly similar to the molars of a mouse relative, *Mastacomys fuscus brazenori*. Its first upper molars have very large supplementary cusps and have lost the same buccal cusp as the *Bmper* mutant, and its lower molars have larger lingual cusps as compared to buccal cusps (Museums Victoria Collections). Therefore, our data strongly suggest that evolutionary changes in the BMP4/OR2 network might be responsible for both the evolution of the murine dental plan and its further diversification.

Where could be the mutation which impacted this BMP4/Osr2 network? From the present data, we envision at least three possibilities. 1) A cis-regulatory change in the *Bmper* gene could have led to its new asymmetric profile, and feedback on *Sfrp2* expression through the network. In cichlids, a QTL containing *Bmper* is associated with variation in tooth number(Bloomquist et al., 2015). Species with more teeth have reduced *Bmper* expression, as mouse with more lingual cusps have reduced lingual *Bmper* expression (Bloomquist et al., 2015). 2) The mutation may lie in *Sfrp2*, and feedback on *Bmper* expression. 3) The mutation might also lie in *Dlx1/2* genes, because they control both *Sfrp2* and *Bmper* expression levels in the early mouse jaws (Jeong et al., 2008). It is striking that these 3 genes show only a minor expression difference in favor of the upper molar in hamster, but their expression is twice increased in mouse, together with a sharper upper molar bias. By acting on both a buccal (*Bmper*) and a lingual *(Sfrp2)* gene with antagonistic effects on the BMP4/Osr2 balance, the increased dose of *Dlx1/2* might have converted the mild ancestral polarization of the tooth into the sharp bucco-lingual polarisation seen in mouse. Future work focusing on the evolution of the cis-regulatory regions of these genes could test these hypotheses.

Finally, we noticed clear changes in the expression patterns of *Bmp4*, *Wif1* and *Dkk1*, three members of the BMP and Wnt pathways at the core of activation-inhibition networks (O'Connell et al., 2012; Salazar-Ciudad, 2012). Given the many regulatory feedback in these networks, a mutation may lie in one of these genes and feedback on the expression of the others, or lie in another gene to be identified.

1004   <u>Lower molar development evolved in a concerted manner with upper molar development</u>

1005

We show that the lower molar development has coevolved with the upper molar

development. Temporal expression profiles coevolved massively and several features of

cusp formation also evolved in a concerted manner: the precocious PEK to SEK transition,

the narrow expression of inhibitors in signaling centers, the marked bucco-lingual

asymmetry with persistence of some lingual naive tissue and the early *Bmper* withdrawal

from the lingual side. Concerted evolution in bucco-lingual development is especially

striking, since neither cusp number nor relative size along the B/L axis differs between

mouse and hamster lower molars (Figure 1). The only derived features of the mouse lower

molar are the connection between cusps (the crest connecting central and posterior cusps is

lost) and their slightly more parallel arrangement (Lazzari et al., 2008).

Why these concerted developmental changes translate into minor phenotypic change in the

lower molar but major ones in the upper molar is an open question. Molar development has

non-linear properties, characterized by threshold effects (Gjuvsland et al., 2013; Milocco &

Salazar-Ciudad, 2020; Morita et al., 2020; Urdy et al., 2016). Since the lower molar can form

supplementary lingual cusps when activation is boosted by adding ACTIVINβA to the

culture medium (Harjunmaa et al., 2012), it suggests that in wild type mouse, the lower

molar remains below a threshold, while the upper molar passes it. There could be two

different, non mutually exclusive reasons for that: 1) As mentioned above, there were

ancestral differences in the regulation of the bucco-lingual axis in the common ancestor.

When facing the same expression change (e.g. early *Bmper* withdrawal from the lingual

side), its upper and lower molar might then have reacted very differently, passing or not the

threshold. 2) The upper-specific increase in mesenchyme proportion may be just the small

effect needed to pass the threshold in the upper molar, while the lower molar remains

below it. Indeed, the mesenchyme is the endogenous source of ACTIVINβA, whose

supplementation produces lingual cusps in the lower molar (Harjunmaa et al., 2012).

1031

1032

1033

<u>A combinatory model to explain the independent phenotypic evolution of the upper molar</u>

<u>with concerted developmental evolution in the lower molar</u>

The three features that we observed hint at very complementary aspects of tooth development (Figure 8). The tooth literature shows it is difficult to increase cusp number in mouse molars: *in vitro* experiments have shown it can be necessary to play on multiple pathways, and mouse mutants show, at best, small accessory cusps, but no supplementary main cusps (Harjunmaa et al., 2012). None of these three changes should be sufficient on its own to induce the major changes in cusp size and proportions seen in mouse as compared to its ancestor. We therefore propose that the new phenotype involves combining mutations in at least two or three different genes, corresponding to these three features (Figure 8). Such a model with additive changes is also coherent with the stepwise addition of the supplementary cusps in the fossil record. Stem murine rodents had a single small extra cusp. Enlargement of this cusp, addition of a second extra-cusp, and size reduction of the buccal cusps came later (Tiphaine et al., 2013).

The 3 mutations which could lie behind the observed developmental phenotypes represent three different categories, with respect to their consequences for the lower molar (Figure 8).

i) mutation with organ-specific developmental effects - A mutation in the upper-molar specific *Pou3f3* gene which was part of the ancestral lower/upper code could have molecular effects specific to the upper molar. Such a mutation and effect are expected from the abundant literature on homeotic genes and serial appendage evolution.

ii) mutation with shared effects, but exaggerated in the upper molar - Another mutation could have the same molecular effects on the development of both molars (e.g. a mutation making *Bmper* expression asymmetric) but with a stronger expressivity in upper molar (e.g. larger lingual field) because the ancestral lower/upper code determines a different developmental context between the two teeth.

iii) mutation with fully shared effects on lower and upper molar development - A third mutation could have the same molecular effects and the same expressivity on the developmental phenotype (range of inhibitor expression in the signaling centers of the two

1064   teeth), but because it cannot combine with other effects in the lower molar as it does in the

1065   upper molar, it might have a very limited impact on the lower molar phenotype.

1066

1067   Such a genetic model is consistent with findings in butterflies' wings. Indeed,

1068   combinatory effects of mutations and context-dependency on the ancestral homeotic code

1069   have been proposed to explain the evolution of eyespot patterns in the fore and hind-wings

1070   (Monteiro 2007, 2021, 2022).

1071

1072   The patterns of transcriptome evolution seen in teeth resemble patterns observed in other

1073   serial organs

1074

Comparative transcriptomics in embryos may be confounded by methodological effects

that could inflate interspecies differences in expression levels. This includes estimating

expression levels with RNA-seq data from different species and sampling a few stages in a

continuous developmental window in species with different developmental rates. We

controlled for this by estimating expression levels on orthologous portions of the

transcripts, by matching the time window with homologous stages, by sampling many time

points, and working on temporal profiles instead of individual stages.

1082

To the best of our knowledge, there are only two other studies using interspecies

transcriptomics in serial organ evolution, in similar settings: at least two species, one

representing ancestral morphologies, and another one where a single organ strongly

diverged from the ancestral morphology. One study compares bones from fore/hind limbs

in mouse and jerboa at a single timepoint (Saxena et al., 2022). The other compares

limbs/wings in mouse and bat at three timepoints (Maier et al., 2017) which allowed us to

reanalyse the data with our methods.

Similar patterns are seen in all these datasets. 1) Only a small set of genes, enriched in

transcription factors, showed an organ-specific expression conserved between species (our

tooth data, reanalyzed limbs dataset, and (Maier et al., 2017; Saxena et al., 2022)). 2) The

expression of large numbers of genes co-evolved in the two organs (tooth and reanalyzed

limb data, (Saxena et al., 2022)). 3) Expression differences between serial organs are

increased in species with the morphological innovation (tooth and reanalyzed limbs data,

(Saxena et al., 2022)) 4) but the serial organ which kept the most ancestral morphology

does not show better expression conservation (tooth and reanalyzed limbs data, and

(Saxena et al., 2022).

1099

1100   <u>Co-evolution is also pervasive in bat limbs, where adaptation combines organ-specific with</u>

1101   <u>shared gene expression changes.</u>

1102

Our comparative analysis of early mouse and bat development revealed that

developmental dynamics of gene expression is largely shared by the two bat limbs, despite

their drastically different morphologies. This concerted evolution was largely overlooked so

far (e.g. *Shh* gene), because attention was mainly given to wing-specific developmental

features, which seem more logically susceptible to explain wing evolution. We however

realized that gene expression changes, previously pointed for their role in bat wing

evolution, are in fact accompanied by concerted expression changes in the hindlimb. The

bat wing membrane is achieved by suppressing the apoptosis which normally defines the

digits. Functional tests showed that this is achieved by simultaneously activating the anti-

apoptotic FGF pathway and downregulating the pro-apoptotic BMP pathway (Weatherbee

et al., 2006). *Fgf8* and the BMP inhibitor *Grem1* coevolved in our analysis (Figure S12), with

a new mesenchymal expression in both bat limbs (Figure 8, drawn from Figure 3 A-C, E-G in

(Weatherbee et al., 2006)). At later stages, *Fgf8* mesenchymal expression persists in the

interdigital area in the wing, but not in the foot (from Figure 3H in (Weatherbee et al.,

2006)). In contrast, the BMP inhibitor *Grem1* expression persists in both limbs, with higher

levels in the wing (from their Figure 3D, note the blue staining remaining around digits

whereas more proximal parts of the limb are not stained at all). Thus, at this stage, specific

and exaggerated shared gene expression changes seem to combine to pass the threshold

for apoptosis suppression in the wing, but not in the foot (Figure 8). This evolutionary

scenario of independent evolution is thus very similar to teeth, involving a combination of

specific, shared, and exaggerated shared expression changes and differential threshold

effects.

1125

1126 <u>Concerted evolution with Developmental System Drift is a mechanism facilitating</u>

1127 <u>independent evolution of serial organs</u>

1128

1129 We observed incongruent patterns of transcriptome and morphologies in molar evolution :

1130 transcriptomes diverged equally in the upper and lower molars, while the morphology of

1131 the lower molar remains largely conserved. This unexpected level of developmental

1132 divergence as compared to morphological conservation is called Developmental System

1133 Drift (DSD, (Cutter & Bundus, 2020; Félix, 2012; True & Haag, 2001)).

1134

There is now accumulating evidence that cryptic changes in developmental systems are

frequent in evolution (Félix, 2007; Guignard et al., 2020; Torres Cleuren et al., 2019; Wotton

et al., 2015). Because natural selection mainly acts on the final product of development,

divergent developmental paths may be taken to reach the same final phenotype and drift in

development is neutral with respect to natural selection. Further taking into account that

genomes are constantly mutating, DSD appears as a likely alternative to developmental

conservation (Félix & Wagner, 2008; Peter & Davidson, 2011).

1142

The situation here seems different from this classical definition of DSD since at least part of

lower molar and hindlimb DSD is not random: it is concerted with developmental

innovation in the other organ, and therefore likely induced by the adaptation of this other

organ. Because the lower molar and the hindlimb developmental systems could evolve

while robustly maintaining the final phenotype during evolutionary times, mutations with

shared effect could be used by adaptation. This is unexpected since it is commonly thought

that adaptive mutations need to be modular at the DNA level to have organ-specific effects

and thereby circumvent gene pleiotropy. The capacity of developmental systems to

undergo DSD is another way of circumventing gene pleiotropy, and thus appears as a

mechanism by which non-modular mutations can be selected in adaptation. We propose

this is the reason why independent evolution can be so frequently seen in nature despite

gene pleiotropy.

1155

<u>Pleiotropy, concerted evolution and DSD</u>

Serial organs such as molars and limbs have a heavy pleiotropy load and for this reason,

they are possibly especially prone to developmental co-evolution and DSD. We

nevertheless believe that our results in serial organs illustrate a much more general

correlation between pleiotropy and DSD at the organismal level, as suggested previously

(Félix, 2007; Pavlicev & Wagner, 2012).

The link between pleiotropy and DSD has been observed in experiments of *in silico*

evolution (Johnson & Porter, 2007; Tulchinsky et al., 2014). It has also been observed in

nematode genetics with a mutation increasing the fitness in laboratory conditions that has

induced DSD in the vulva (Duveau & Félix, 2012). Finally, a link between pleiotropy and

concerted transcriptomic evolution has already been suggested. In most multispecies

transcriptomic analyses, samples of different organs tend to group by species (like molar

samples in Figure 1C). This pattern, so-called "species signal", often dominates in samples

of adult tissues (e.g. kidney, brain, liver… (Brawand et al., 2011)) as well as in individual

embryonic timepoints (Liang et al., 2018; Tschopp et al., 2014). This has been reinterpreted

as a conspicuous concerted evolution, possibly driven by the pleiotropy of gene networks,

repeatedly used in different organs (Liang et al., 2018; Musser & Wagner, 2015).

1173

We further suggest that pleiotropy-induced DSD may explain another observation

concerning genes involved in human diseases and pleiotropic genes. It was expected that

the embryonic expression profiles of these important genes would evolve slowly, but they

evolve as fast as the rest of the genome (Cardoso-Moreira et al., 2019, 2020). Further work

may reveal which part of sequence and expression divergence which is usually attributed to

genetic drift (divergence by random chance) could in fact be attributed to "pleiotropy-

induced DSD".

1181

1182

MATERIAL AND METHODS

<u>Data analysis</u>

R scripts corresponding to the main methods and processed data are available on GitHub (https://github.com/msemon/DriftHamsterMouse).

<u>Rodent breeding and embryo sampling</u>

CD1 (CD1) adult mice and RjHan:AURA adult hamsters were purchased from Charles River (Italy) and Janvier (France) respectively. Females were mated overnight and the noon after morning detection of a vaginal plug or sperm, respectively, was indicated as ED0.5. Other breeding pairs were kept in a light-dark reversed cycle (12:00 midnight), so that the next day at 16:00 was considered as ED1.0.

The $Bmper^{tm1Emdr}$ strain (Zakin et al., 2008) was kept in a C57/BL6N background by crossing heterozygotes with wild types, as homozygotes die at birth. To avoid suffering at birth, we generated homozygotes and wild type samples for X-ray by crossing heterozygotes and sacrificing pregnant mice at 19.5 days (1 day before delivery). Pregnant mouse females were killed by cervical dislocation. Hamster females were deeply anesthetized with a ketamine-xylasine mix administered intraperitoneally before being killed with pentobarbital administered intracardially. All embryos were harvested and thereby anesthetized on cooled Hank's or DMEM advanced medium, weighted as described in (Peterka et al., 2002) and immediately decapitated.

This study was performed in strict accordance with the European guidelines 2010/63/UE and was approved by the Animal Experimentation Ethics Committee CECCAPP (Lyon, France, APAFIS#27308-2020092210045896 v1).

<u>Estimating embryonic age from embryo weight</u>

Embryo weight is well correlated with developmental age, allowing us to use it as a proxy in mouse and hamster, following (Pantalacci et al., 2009). We fitted age of development according to weight (in mg) for hamster and mouse data separately, based on 1047 mouse

embryos and 636 hamster embryos respectively, collected over more than 15 years of

research. We fitted generalised additive models (GAM) to the data after Box-Cox

transformation of weight (libraries mgv version 1.8-35 for GAM and MASS 7.3-53.1 for Box-

Cox). These models were prefered to log transformations and linear models, because they

allow to treat the data homogeneously between species, and because the relationship is

not perfectly linear between weight and age (Figure S1). These models were then used to

predict developmental age, based on weight, for all samples used in this study (RNA-seq

analysis, cusp patterning analysis, and *in situ* hybridizations for several genes).

1220

<u>Epithelium dissociations and in situ hybridizations</u>

Complete or hemi mandibles and maxillae were dissected in Hank's medium and treated

with Dispase (Roche) 10mg/ml in Hepes/KOH 50mM ph7.7; NaCl 150 mM at 37 °C for 30

min to 1h depending on embryonic stage. Epithelium and mesenchyme were carefully

separated and fixed overnight in PFA 4% at 4 °C. DIG RNA antisense mouse *Fgf4*, *Shh*,

Fgf10 (Bellusci et al., 1997), *Bmper/Cv2* probes were prepared from plasmids described

elsewhere (Coffinier et al., 2002). Mouse *Dkk1*, *Wif1*, hamster *Bmper* probes, Mouse and

hamster *Bmp4* probes were newly cloned following RT-PCR or DNA synthesis (Table S1). *In*

situ hybridizations were done according to a standard protocol (DIG mix, DIG antibody and

BM purple were purchased from ROCHE). Photographs were taken on a Leica M205FA

stereomicroscope with a Leica DFC450 digital camera (Wetzlar, Germany) or on a Zeiss

LUMAR stereomicroscope with a CCD CoolSNAP camera (PLATIM, Lyon, France).

1233

<u>Immunolocalisation and 3D reconstructions</u>

Tooth germs dissected from litter-mate embryos of RNA-seq samples were fixed overnight

in PFA4% and dehydrated through a methanol series. In toto immunolocalization protocol

was adapted from (Ahnfelt-Rønne et al., 2007). Following incubation in H202 5%, DMSO

10% in methanol for 4 hours, they were rehydrated, blocked and incubated successively

with a pan-cytokeratin antibody (overnight,1/200, Novus Biologicals) and a Dylight 549

conjugated Donkey Anti-rabbit antibody (overnight 1/200, Jackson immunoresearch) and

finally with Hoechst (overnight , 50µg/ml). Following methanol dehydration, they were

clarified and mounted in BABB as described in (Yokomizo & Dzierzak, 2010). They were

imaged with a Zeiss LSM710 confocal microscope at the PLATIM (Lyon, France). The outline

of the epithelium labelled by the pan-cytokeratin antibody and the outline of the tooth

germ labelled with hoechst were delineated manually and reconstructed in 3D in the

AMIRA software.

1247

<u>X-ray scanning and 3D reconstruction of *Bmper* and wild type tooth shape at 19.5 days</u>

We obtained 14 homozygote (Ho) and 19 wild type (Wt) samples from a total of 14 different

19.5 dpc litters, out of which we selected for reconstruction of the first molar morphology 7

Ho and 4 Wt with matching body weights (homozygotes: 1174-1329 mg; wt: 1227-1310

mg). At 19.5 dpc, female embryos are more developmentally advanced than male embryos

of a similar weight, therefore sex was also recorded. This was necessary to control for

differences in growth advancement, since we anticipated (and confirmed) that

supplementary cusps are still growing rapidly at this stage, due to their late formation.

Heads freed of skin were fixed in PFA, dehydrated in ethanol, stained with 0,3% PTA in 70%

ethanol for 2 weeks-1 month and scanned at 40kV on a Phoenix Nanotom S

microtomography for a voxel size of 4 μm. Semi-automatic reconstruction of the enamel-

dentin junction was performed with ITK-snap. Reconstructions were oriented for

comparison in MeshLab 2021.05. Due to variations in staining efficiency and advancement

of mineralization, only a total of 4 Ho and 4 Wt upper molars and 3 Ho and 4 Wt lower

molars were finally successfully reconstructed and considered to be directly comparable.

The aberrant upper molar morphology was obvious on microCT sections in 7/7 samples.

The loss of one cusp was observed in 4 Ho/4 3D reconstructions. The larger lingual cusps

were observed in all Ho 3D reconstructions when paired with a wt of corresponding age.

This (as well as cusp loss) was also confirmed by comparing epithelial dissociations of

Bmper Ho and Wt embryos at 19.5 dpc.

1268

<u>Modelling and comparing cusp patterning dynamics</u>

To compare the dynamic of crown morphogenesis in four teeth (lower and upper molars in

hamster and mouse) we need to establish the sequence of primary and secondary signalling

centres formation (respectively, PEK and SEK). In mouse, this could be done with time lapse

imaging of fluorescent lines (Harjunmaa et al., 2014). To integrate non-model species like

hamster, we had to set up a new method that infers the dynamic based on fixed embryos.

We hybridised developing molars against a *Fgf4* probe to reveal PEK and SEKs. The

patterns we observed among samples are consistent with a stereotypic and specific

sequence of SEK patterning in each tooth and species (Figure 1B, schemas on the sides).

We name each stage by the number of signalling centres (PEK stage then 1-SEK stage, 2-

SEK stage etc).

Cusp patterning can be seen as a succession of irreversible stages representing step-wise

cusp additions. Transition rates between these stages were modelled through continuous

time Markov modelling as in (Pantalacci et al., 2017). The rationale is that if sampling is

uniform over the time course of tooth morphogenesis, stages that are rarely sampled are

very transient (with high exit rate), while stages that are often sampled last for a longer

period of time. In continuous Markov models, the duration of each state follows an

exponential distribution, which is not realistic for the stage lengths. So, to have a more

realistic stage length distribution, each stage was modelled by several consecutive states,

so that its length followed an Erlang distribution, which has a mode different from zero. We

built independent models for each species and tooth types. Models are estimated on 121

embryos for the hamster lower molars, 113 for hamster upper, 217 for mouse lower, 187 for

mouse upper.

We estimated the duration of each stage in a complete model, with different transition rates

for all stages. We also fitted several simpler, nested models, with constraints on the number

of different transition rates, up to the most simple model with the same transition rate for all

stages. We retained models with three different rates in mouse, and two different rates in

hamster, by comparing the fit of the models by likelihood ratio tests in each tooth. Markov

models were built by custom scripts calling on R libraries maxLik and expm (maxLik_1.4-8

and expm_0.999-6).

1299

1300

1301

<u>RNA-seq sample preparation</u>

A total of 32 samples per species, coming from eight individuals, were prepared for the time serie RNA-seq analysis, representing consecutive stages in mouse (ED14.5, 15.0, 15.5, 16.0, 16.5, 17.0, 17.5, 18.0) and nine stages in hamster (ED11.8, 12.0, 12.2, 12.5, 13.0, 13.25, 13.5, 13.75, 14.0). Each sample contained two whole tooth germs, the left and right first molars (M1) of the same female individual, and for a given stage, the upper and lower samples came from the same individual. Harvesting and dissection were performed in a minimal amount of time in advanced DMEM medium. The M1 lower and upper germs were dissected under a stereomicroscope and stored in 200 uL of RNA later (SIGMA). Similarly dissected tooth germs from the same litter and same weight were fixed overnight in PFA 4% for immunolocalization and 3D reconstruction, to check for dissection leaving almost no non-tooth tissue. Examples of dissection are visible in (Pantalacci et al., 2017). Another embryo of the same litter and same weight was processed as indicated above for *Fgf4 in situ* hybridization to check the exact developmental stage. Total RNA was prepared using the RNeasy micro kit from QIAGEN following lysis with a Precellys homogenizer. RNA integrity was controlled on a Bioanalyzer (Agilent Technologies, a RIN of 10 was reached for all samples used in this study). PolyA+ libraries of the large-scale dataset were prepared with the Truseq V2 kit (Illumina, non stranded protocol), starting with 150 ng total RNA and reducing the amplification step to only 12 cycles and sequenced on an Illumina Hi-seq2000 sequencer (100 bp paired end reads) at the GENOSCOPE (Evry, France).

For the bucco-lingual dataset, we dissected the 4 first molars (left/right, lower/upper) from a unique mouse E15.0 embryo (weight: 359 mg) as above, except that tooth germs were cut in two halves to isolate buccal and lingual side. Replicates were thus obtained by comparing the right and left side of this same embryo. Total RNAs were extracted and libraries were prepared as above, starting with 50-70 ng total RNAs, where an equal amount of AmbionR ERCC RNA Spike-In Mix1 had been added according to the AmbionR protocol (e.g. 1µL og a 1 :1000 dilution for each tube). A total of 8 libraries were sequenced (50bp single-end reads) by the Genomeast Sequencing platform, a member of the France Genomique program.

For the epithelium-mesenchyme dataset, lower and upper mouse and hamster first molars

were dissected as above and treated for 15 minutes at 37°C with Dispase (Roche) 10mg/ml

in Hepes/KOH 50mM ph7.7 ; NaCl 150 mM to separate the epithelial from mesenchymal

parts which were stored in RNAlater. For the mouse data, we generated samples for 2

stages in 2 replicates, using embryos from the same litter (stage 15.0 dpc, weight: 350 and

370 mg; stage 16.5 dpc: weight: 788 and 808 mg). Left and right epithelium or

mesenchyme were pooled. For the hamster data, we generated samples for a single stage

without replication. We pooled the left epithelial or mesenchymal parts from 2 embryos

from the same 12.5 dpc litter (413 and 427mg). A total of respectively 16 and 4 libraries

were generated with Truseq V2 kit and sequenced (50bp single-end reads) by the

Genomeast Sequencing platform.

1342

<u>Multivariate analyses</u>

Multivariate analyses were performed using the ade4 package (ade4_1.7-18; (Dray &

Dufour, 2007)). We performed principal component analyses on normalised counts (DESeq

basemeans), and between groups analyses on the resulting components, which allowed us

to quantify the proportion of variance associated with each factor.

We used RAPToR (RAPToR_1.1.4, (Bulteau & Francesconi, 2021)) to estimate the offset

between upper and lower molar development. We staged upper molar samples on

reference made from lower molar samples conjointly for both species. Interpolations were

done with gam models fit on 5 components of an ICA.

1352

<u>Expression levels estimation using RNA-seq and differential expression analysis</u>

For the whole tooth germ data (64 samples) we obtained 100 bp paired-end sequences,

with on average 46.2M (millions) reads per sample. For epithelium/mesenchyme and

bucco/lingual data (respectively 20 and 8 samples), we obtained 50 bp single-end

sequences, with on average 93.7M and 48.6M reads per sample respectively. Raw data are

publically available in ENA with project accession number: PRJEB52633.

1359

These reads were mapped by using Kallisto (version 0.44.0, (Bray et al., 2016)) to custom

reference sequences for hamster and mouse transcriptomes. To generate them, we

retrieved mouse and hamster cDNAs from Ensembl (release 93, July 2018, assemblies

GRCm38.p6 and MesAur1.0 (Howe et al., 2021)), selected 14536 pairs of one-to-one

orthologous transcripts, realigned pairs of sequences with Macse (macse_v2.01, (Ranwez et

al., 2011)) and cropped the alignments to get orthologous segments by using custom

scripts to make expression levels comparable between species.

1367

Differential gene expression analysis (DE analysis) was performed on smoothed expression

profiles over developmental age. Developmental age was estimated with embryo weight

(GAM models above) and stages were homologized based on morphological criteria at

early cap stages and late bell stage (14.5/18.0 days in mouse; 12.2/14.5 days in hamster).

These boundaries, confirmed by PCA analysis of the transcriptome data (Figures 1 and S1)

were used to convert days of development into relative development age (0-10).

1374

Expression profiles were fitted by third degree polynomial splines with 2 interior knots, for

each tooth and species (bs function of spline R package (W. Wang & Yan, 2021),

independently or jointly within tooth and/or species, as explained below. Nested models

were tested by DEseq2 (Love et al., 2014) and the best model was chosen for each gene by

comparing the fit of these nested models (FDR adjusted p-value < 0.05 from DESeq2 LRT

tests). When we compared temporal profiles between species, we accounted for the

average level of expression in each species. This is to focus on changes in regulation over

development, and to discard potential remaining artefacts in species-specific

quantifications. Several tests were performed and are described below with the

corresponding figure number.

1385

To model temporal expression profiles in each species separately (Related to Figure 2A), we

compared a "simple" model with a single curve for both time series to a "complex" model

with one specific curve per tooth. This procedure does not directly provide the sense of the

bias. To estimate it in each species, we computed the values predicted by the upper and

lower models for 100 equally distributed points (i) on the timeline, and measured the

distance point by point as follows : sum((up(i)-low(i))/(up(i)+low(i)) (Mus.dist.sign or

Ham.dist.sign in Table S2). This measure evaluates whether overall, expression levels in

upper molar are above or below lower molar's.

1394

1395 To model the divergence of temporal expression profile in each tooth type separately

1396 (Related to Figure 6D), we compared a "non-divergent" model with a single curve to fit

1397 both time series (with a species-specific offset to only consider the temporal dynamic), to a

1398 "divergent" model with one specific curve per species (with a species-specific offset).

1399

1400 To model temporal expression profiles in the 4 tooth types with tooth-specificity in one

1401 species (related to Figure 3B), the four time series were modelled as in the

1402 "hamster/mouse", plus "mouse tooth-specific" and "hamster tooth-specific" models, which

1403 correspond to distinct curves for upper and lower molars in one species, but not in the

1404 other. Genes were selected by comparison with the "hamster/mouse" model above.

1405

Selection of the temporal expression profile in the 4 tooth types was done as follows

(related to Figure 6A). The "simple" model fits a single curve for the four time series. The

"complex" model fits 4 different curves, one per tooth type. The "hamster/mouse" model

has 2 different curves, one per species. The "upper/lower" model has one curve per tooth,

including the species-specific offset. The best model was selected for each gene by using a

bottom-up approach with the results of four independent tests: t1 compares

"hamster/mouse"*versus* simple model; t2: "upper/lower" *versus* simple; t3: complex *vs*

"upper/lower"; t4: complex *vs* "hamster/mouse". If t1 and t2 are not significant, then the

simple model is chosen. If t1 is significant and not t2, the gene is assigned to:

"hamster/mouse". If t2 but not t1: "upper/lower". Finally, if "lower/upper" or

"hamster/mouse" and t4: complex.

From this selection procedure, percentage of coevolution among genes was computed as

the proportion of selected "hamster/mouse" models among the selected models as follows

(related to Figure 6A): "hamster/mouse"/("hamster/mouse"+"upper/lower"+"complex").

1420

We then computed the intersection of the results with several lists of genes important for

tooth development: 259 genes from the bite-it database (retrieved in July 2019), 187 genes

with a phenotype in tooth development (100 "dispensable" genes, 87 "keystone" genes,

(Hallikas et al., 2021)), and 295 genes belonging to 8 pathways active in tooth development

(17 genes in ACTIVIN pathway, 81 in BMP, 10 in EDA, 69 in FGF, 32 in SHH, 9 in NOTCH,

11 in TGFB, 96 in WNT, courtesy Jukka Jernvall).

1427

1428   <u>Gene Ontology analysis</u>

1429   Gene ontology (GO) analysis was performed and visualised with gprofiler2 (gprofiler2_0.2.0,

1430   (Kolberg et al., 2020)), clusterProfiler (clusterProfiler_3.18.1, (Wu et al., 2021)), and

1431   simplifyGO (simplifyEnrichment_1.2.0 (Gu & Hübschmann, 2021)) using the full list of genes

1432   expressed in the corresponding dataset as a background.

1433

<u>Measure of pathway activation in RNAseq samples</u>

ROMA was used to quantify activation of WNT, BMP and SHH pathways in the bucco-

lingual samples (version rRoma_0.0.4.2000, https://github.com/Albluca/rRoma and

(Martignetti et al., 2016)). ROMA is designed to compare pathway activity in transcriptomic

samples based on expression levels of a list of targets for the pathway. Genes for the SHH

modules were retrieved from GSEA ((Mootha et al., 2003), 41 genes present in our dataset).

Because BMP and WNT pathways are active both in the mesenchyme and the epithelium

and they target different genes in each tissue (O'Connell et al., 2012), we used two

separate lists of targets to estimate both an epithelial and a mesenchymal activity, adapted

from a "regulatory evidence" dataset established for first lower molar development

(O'Connell et al., 2012). Building on literature and their own transcriptomic analysis, the

authors had defined target genes based on their up or downregulation following activation

or inactivation of each pathway. For data consistency, we selected only targets established

in the study from transcriptome analysis, in 13.5 and 14.5 dpc epithelium and 10.5 dpc

mesenchyme. Different modules were built for activities in the mesenchyme and epithelium

compartments. For BMP in the mesenchyme, we considered 15 genes as positive targets

and 4 as negative targets (further noted 15:4). In the epithelium, the numbers of

positive:negative targets were respectively 32:34. For WNT, we built modules with 4:31

positive:negative targets in the mesenchyme, and 33:45 in the epithelium. These in-silico

quantifications were consistent with many known aspects of tooth development. Buccal

compartments all show high levels of pathway activity, consistent with the presence of the

first SEK acting as a source of WNT, BMP and SHH signals. Lingual compartments show

much lower levels of signalling activities than buccal compartments, consistent with their

distance to the first SEK. The lower lingual compartments show BMP and WNT activities

that are higher in epithelium than in mesenchyme, consistent with the fact that epithelial

activation predates mesenchymal activation in tooth development.

1460

<u>Estimating tissue proportions from RNA-seq data by deconvolutions</u>

We used the R package DeconRNASeq (DeconRNASeq_1.32.0 (Gong & Szustakowski,

2013)) to estimate the relative proportions of epithelium and mesenchyme compartments in

bulk tooth germ transcriptomes. We defined lists of marker genes for each compartment by

pairwise differential analysis of tissue-specific transcriptomes (DESeq2, log2 fold change >

3, adjusted p-value < 0.05). We used 1025 mesenchyme and 621 epithelium marker genes

found by comparing 10 epithelium and 10 mesenchyme RNAseq samples, mixing tooth,

stages and species. We estimated the accuracy of the prediction by bootstrapping 1000

times the marker lists. The relative proportions of buccal and lingual compartments was

inferred by a similar procedure. We used 414 buccal and 235 lingual marker genes, from

the differential analysis of 8 samples (DESeq2, log2 fold change > 1, adjusted p-value <

0.05).

1473

<u>Expression levels and transcriptome dynamics in bats</u>

We downloaded all bats and mouse raw RNA-seq samples from a published dataset

(SRP061644, Maier et al. 2017), totalizing 17 samples in mouse and 16 in bat (*Carollia*

Perspicillata) at three consecutive stages: ridge (CS13), bud (CS14) and paddle (CS15)

stage. Bat reads were assembled *de novo* with Trinity v2.14.0 (Grabherr et al., 2011), by

using single end mode and *in silico* normalisation. Bat expression levels were quantified by

Salmon (Patro et al., 2017) with the script provided by Trinity

(align_and_estimate_abundance.pl). Mouse reads were directly mapped with Salmon to the

GENCODE mouse transcriptome reference (gencode.vM29.pc, (Frankish et al., 2021)). Bat

transcripts were assigned to mouse orthologs by blastn (Altschul et al., 1990). Blast and

Trinity were run with prebuilt dockers. Differential analysis was run over smoothed

expression profiles like in the method section "Expression levels estimation using RNA-seq

and differential expression analysis". Code is available here:

https://github.com/msemon/DriftHamsterMouse

1488

1489

1490

1491 # ACKNOWLEDGEMENTS

1492

We acknowledge the contribution of several platforms of SFR Biosciences Gerland-Lyon Sud

(UMS344/US8): the Plateau de Biologie Expérimentale de la Souris (PBES) (many thanks

especially to Jean-Louis Thoumas, Tiphaine Dorel, Céline Angleraux, Marie Teixeira, Myriam

Prudent), the Plateau Technique Imagerie/Microscopie (PLATIM) and the ANIRA/IMMOS

platform with Mathilde Bouchet; as well as the computer resources from PSMN (ENS Lyon).

We acknowledge the technical help of Anne Lambert, Alain Rubod, Mathilde Estevez-Villar,

Claudine Corneloup and the contribution of many students including Coraline Petit, Alice

Lorenc, Margaux Pillon, Ludivine Rotard and Asma Benahmed. We are grateful to several

colleagues and their staff for sending plasmid probes: Irma Thesleff (*Fgf4*) , Hiko Ogura

(*Cv2/Bmper*), D. Duboule (*Fgf10*). We are grateful to Pamela Lockyer, Cam Patterson and

Xinchun Pi for collecting our very first *Bmper* embryos. We are grateful to Jennifer Esser for

kindly providing the *Bmper* strain and advice on genotyping. We kindly thank Mirko

Francesconi, Marie Delattre, Michaelis Averof, Pascal Hagolani for their feedback on the

manuscript.

1507

1508   This work was supported by the Agence Nationale pour la Recherche (ANR 2011 JSV6 00501

1509   "Convergdent"), a grant from the GENOSCOPE - Centre National de Séquençage and a

1510   grant from IDEX Lyon ANR-16-IDEX-0005. Salaries were supported by the Centre National

1511   de la Recherche Scientifique, the Ecole Normale Supérieure de Lyon and the Université de

1512   Lyon, Université Lyon 1. Klara Steklikova benefited from a Barrand Fellowship program.

1513

1514

1515

1516

1517

1518

1519

1520

1521

1522

1523

1524

1525

1526

1527

1528

1529

1530

1531

1532

1533

1534

REFERENCES

Ahnfelt-Rønne, J., Jørgensen, M. C., Hald, J., Madsen, O. D., Serup, P., & Hecksher-

Sørensen, J. (2007). An improved method for three-dimensional reconstruction of

protein expression patterns in intact mouse and chicken embryos and organs. *The*

Journal of Histochemistry and Cytochemistry: Official Journal of the Histochemistry

Society, *55*(9), 925–930.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local

alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Atallah, J., Vurens, G., Mavong, S., Mutti, A., Hoang, D., & Kopp, A. (2014). Sex-specific

repression of dachshund is required for Drosophila sex comb development.

Developmental Biology, *386*(2), 440–447.

Bellusci, S., Grindley, J., Emoto, H., Itoh, N., & Hogan, B. L. (1997). Fibroblast growth factor

10 (FGF10) and branching morphogenesis in the embryonic mouse lung. *Development* ,

124(23), 4867–4878.

Bloomquist, R. F., Parnell, N. F., Phillips, K. A., Fowler, T. E., Yu, T. Y., Sharpe, P. T., &

Streelman, J. T. (2015). Coevolutionary patterning of teeth and taste buds. *Proceedings*

of the National Academy of Sciences of the United States of America, *112*(44), E5954–

E5962.

Booker, B. M., Friedrich, T., Mason, M. K., VanderMeer, J. E., Zhao, J., Eckalbar, W. L.,

Logan, M., Illing, N., Pollard, K. S., & Ahituv, N. (2016). Bat Accelerated Regions

Identify a Bat Forelimb Specific Enhancer in the HoxD Locus. *PLoS Genetics*, *12*(3),

e1005738.

Brawand, D., Soumillon, M., Necsulea, A., Julien, P., Csárdi, G., Harrigan, P., Weier, M.,

Liechti, A., Aximu-Petri, A., Kircher, M., Albert, F. W., Zeller, U., Khaitovich, P.,

Grützner, F., Bergmann, S., Nielsen, R., Pääbo, S., & Kaessmann, H. (2011). The

evolution of gene expression levels in mammalian organs. *Nature*, *478*(7369), 343–348.

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Erratum: Near-optimal

probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(8), 888.

Bulteau, R., & Francesconi, M. (2021). Real age prediction from the transcriptome with

RAPToR. In *bioRxiv* (p. 2021.09.07.459270). https://doi.org/10.1101/2021.09.07.459270

Cardoso-Moreira, M., Halbert, J., Valloton, D., Velten, B., Chen, C., Shao, Y., Liechti, A.,

Ascenção, K., Rummel, C., Ovchinnikova, S., Mazin, P. V., Xenarios, I., Harshman, K.,

Mort, M., Cooper, D. N., Sandi, C., Soares, M. J., Ferreira, P. G., Afonso, S., …

Kaessmann, H. (2019). Gene expression across mammalian organ development.

Nature, *571*(7766), 505–509.

Cardoso-Moreira, M., Sarropoulos, I., Velten, B., Mort, M., Cooper, D. N., Huber, W., &

Kaessmann, H. (2020). Developmental Gene Expression Differences between Humans

and Mammalian Models. *Cell Reports*, *33*(4), 108308.

Carroll, S. B. (2008). Evo-devo and an expanding evolutionary synthesis: a genetic theory of

morphological evolution. *Cell*, *134*(1), 25–36.

Charles, C., Lazzari, V., Tafforeau, P., Schimmang, T., Tekin, M., Klein, O., & Viriot, L.

(2009). Modulation of Fgf3 dosage in mouse and men mirrors evolution of mammalian

dentition. *Proceedings of the National Academy of Sciences of the United States of*

America, *106*(52), 22364–22368.

Cho, S.-W., Lee, H.-A., Cai, J., Lee, M.-J., Kim, J.-Y., Ohshima, H., & Jung, H.-S. (2007).

The primary enamel knot determines the position of the first buccal cusp in developing

mice molars. *Differentiation; Research in Biological Diversity*, *75*(5), 441–451.

Cobourne, M. T., & Sharpe, P. T. (2003). Tooth and jaw: molecular mechanisms of

patterning in the first branchial arch. *Archives of Oral Biology*, *48*(1), 1–14.

Coffinier, C., Ketpura, N., Tran, U., Geissert, D., & De Robertis, E. M. (2002). Mouse

Crossveinless-2 is the vertebrate homolog of a Drosophila extracellular regulator of

BMP signaling. *Mechanisms of Development*, *119 Suppl 1*, S179–S184.

Cooper, L. N., Cretekos, C. J., & Sears, K. E. (2012). The evolution and development of

mammalian flight. In *Wiley Interdisciplinary Reviews: Developmental Biology* (Vol. 1,

Issue 5, pp. 773–779). https://doi.org/10.1002/wdev.50

Correns, A., Zimmermann, L.-M. A., Baldock, C., & Sengle, G. (2021). BMP antagonists in

tissue development and disease. *Matrix Biology plus*, *11*, 100071.

Cretekos, C. J., Wang, Y., Green, E. D., Martin, J. F., Rasweiler, J. J., 4th, & Behringer, R.

R. (2008). Regulatory divergence modifies limb length between mammals. *Genes &*

Development, *22*(2), 141–151.

Cutter, A. D., & Bundus, J. D. (2020). Speciation and the developmental alarm clock. *eLife*,

9. https://doi.org/10.7554/eLife.56276

Davis, B. M. (2011). Evolution of the Tribosphenic Molar Pattern in Early Mammals, with

Comments on the "Dual-Origin" Hypothesis. *Journal of Mammalian Evolution*, *18*(4),

227.

Davis, G. K., Srinivasan, D. G., Wittkopp, P. J., & Stern, D. L. (2007). The function and

regulation of Ultrabithorax in the legs of Drosophila melanogaster. *Developmental*

Biology, *308*(2), 621–631.

Depew, M. J., Simpson, C. A., Morasso, M., & Rubenstein, J. L. R. (2005). Reassessing the

Dlx code: the genetic regulation of branchial arch skeletal pattern and development.

Journal of Anatomy, *207*(5), 501–561.

Dray, S., & Dufour, A.-B. (2007). The ade4 Package: Implementing the Duality Diagram for

Ecologists. *Journal of Statistical Software*, *22*, 1–20.

Duveau, F., & Félix, M.-A. (2012). Role of pleiotropy in the evolution of a cryptic

developmental variation in Caenorhabditis elegans. *PLoS Biology*, *10*(1), e1001230.

Eksi, S. E., Barmina, O., McCallough, C. L., Kopp, A., & Orenic, T. V. (2018). A Distalless-

responsive enhancer of the Hox gene Sex combs reduced is required for segment- and

sex-specific sensory organ development in Drosophila. *PLoS Genetics*, *14*(4),

e1007320.

Félix, M.-A. (2007). Cryptic quantitative evolution of the vulva intercellular signaling network

in Caenorhabditis. *Current Biology: CB*, *17*(2), 103–114.

Félix, M.-A. (2012). Evolution in developmental phenotype space. *Current Opinion in Genetics & Development*, *22*(6), 593–599.

Félix, M.-A., & Wagner, A. (2008). Robustness and evolution: concepts, insights and challenges from a developmental model system. *Heredity*, *100*(2), 132–140.

Fisher, C. R., Wegrzyn, J. L., & Jockusch, E. L. (2020). Co-option of wing-patterning genes underlies the evolution of the treehopper helmet. *Nature Ecology & Evolution*, *4*(2), 250–260.

Frankish, A., Diekhans, M., Jungreis, I., Lagarde, J., Loveland, J. E., Mudge, J. M., Sisu, C., Wright, J. C., Armstrong, J., Barnes, I., Berry, A., Bignell, A., Boix, C., Carbonell Sala, S., Cunningham, F., Di Domenico, T., Donaldson, S., Fiddes, I. T., García Girón, C., … Flicek, P. (2021). GENCODE 2021. *Nucleic Acids Research*, *49*(D1), D916–D923.

Gillis, J. A., Modrell, M. S., & Baker, C. V. H. (2013). Developmental evidence for serial homology of the vertebrate jaw and gill arch skeleton. *Nature Communications*, *4*, 1436.

Gjuvsland, A. B., Vik, J. O., Beard, D. A., Hunter, P. J., & Omholt, S. W. (2013). Bridging the genotype-phenotype gap: what does it take? *The Journal of Physiology*, *591*(8), 2055–2066.

Gong, T., & Szustakowski, J. D. (2013). DeconRNASeq: a statistical framework for deconvolution of heterogeneous tissue samples based on mRNA-Seq data. *Bioinformatics* , *29*(8), 1083–1085.

Gould, S. J. (1977). Ontogeny and phylogeny. 1977. *Cambridge. Belknap*. https://www.academia.edu/download/35249361/Ontogeny_and_Phylogeny_final.pdf

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–652.

Guignard, L., Fiúza, U.-M., Leggio, B., Laussu, J., Faure, E., Michelin, G., Biasuz, K., Hufnagel, L., Malandain, G., Godin, C., & Lemaire, P. (2020). Contact area–dependent cell communication and the morphological invariance of ascidian embryogenesis. *Science*, *369*(6500), eaar5663.

Gu, Z., & Hübschmann, D. (2021). simplifyEnrichment: an R/Bioconductor package for Clustering and Visualizing Functional Enrichment Results. In *bioRxiv* (p. 2020.10.27.312116). https://doi.org/10.1101/2020.10.27.312116

Hallikas, O., Das Roy, R., Christensen, M. M., Renvoisé, E., Sulic, A.-M., & Jernvall, J. (2021). System-level analyses of keystone genes required for mammalian tooth development. *Journal of Experimental Zoology. Part B, Molecular and Developmental Evolution*, *336*(1), 7–17.

Harjunmaa, E., Kallonen, A., Voutilainen, M., Hämäläinen, K., Mikkola, M. L., & Jernvall, J. (2012). On the difficulty of increasing dental complexity. *Nature*, *483*(7389), 324–327.

Harjunmaa, E., Seidel, K., Häkkinen, T., Renvoisé, E., Corfe, I. J., Kallonen, A., Zhang, Z.-Q., Evans, A. R., Mikkola, M. L., Salazar-Ciudad, I., Klein, O. D., & Jernvall, J. (2014). Replaying evolutionary transitions from the dental fossil record. *Nature*, *512*(7512), 44–48.

Hersh, B. M., Nelson, C. E., Stoll, S. J., Norton, J. E., Albert, T. J., & Carroll, S. B. (2007). The UBX-regulated network in the haltere imaginal disc of D. melanogaster. *Developmental Biology*, *302*(2), 717–727.

Hillson, S. (2005). *Teeth*. Cambridge University Press.

Hirschberger, C., Sleight, V. A., Criswell, K. E., Clark, S. J., & Gillis, J. A. (2021). Conserved and unique transcriptional features of pharyngeal arches in the skate (Leucoraja erinacea) and evolution of the jaw. *Molecular Biology and Evolution*, *38*(10), 4187–4204.

Hockman, D., Cretekos, C. J., Mason, M. K., Behringer, R. R., Jacobs, D. S., & Illing, N. (2008). A second wave of Sonic hedgehog expression during the development of the bat limb. *Proceedings of the National Academy of Sciences of the United States of*

America, *105*(44), 16982–16987.

Howe, K. L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M. R., Armean, I.

M., Azov, A. G., Bennett, R., Bhai, J., Billis, K., Boddu, S., Charkhchi, M., Cummins, C.,

Da Rin Fioretto, L., Davidson, C., Dodiya, K., El Houdaigui, B., Fatima, R., … Flicek, P.

(2021). Ensembl 2021. *Nucleic Acids Research*, *49*(D1), D884–D891.

Hu, B., Nadiri, A., Kuchler-Bopp, S., Perrin-Schmitt, F., Peters, H., & Lesot, H. (2006).

Tissue engineering of tooth crown, root, and periodontium. *Tissue Engineering*, *12*(8),

2069–2075.

Ikeya, M., Fukushima, K., Kawada, M., Onishi, S., Furuta, Y., Yonemura, S., Kitamura, T.,

Nosaka, T., & Sasai, Y. (2010). Cv2, functioning as a pro-BMP factor via twisted

gastrulation, is required for early development of nephron precursors. *Developmental*

Biology, *337*(2), 405–414.

Järvinen, E., Salazar-Ciudad, I., Birchmeier, W., Taketo, M. M., Jernvall, J., & Thesleff, I.

(2006). Continuous tooth generation in mouse is induced by activated epithelial

Wnt/beta-catenin signaling. *Proceedings of the National Academy of Sciences of the*

United States of America, *103*(49), 18627–18632.

Jeong, J., Li, X., McEvilly, R. J., Rosenfeld, M. G., Lufkin, T., & Rubenstein, J. L. R. (2008).

Dlx genes pattern mammalian jaw primordium by regulating both lower jaw-specific and

upper jaw-specific genetic programs. *Development* , *135*(17), 2905–2916.

Jernvall, J., & Thesleff, I. (2012). Tooth shape formation and tooth renewal: evolving with the

same signals. *Development* , *139*(19), 3487–3497.

Jia, S., Kwon, H.-J. E., Lan, Y., Zhou, J., Liu, H., & Jiang, R. (2016). Bmp4-Msx1 signaling

and Osr2 control tooth organogenesis through antagonistic regulation of secreted Wnt

antagonists. *Developmental Biology*, *420*(1), 110–119.

Jia, S., Zhou, J., Gao, Y., Baek, J.-A., Martin, J. F., Lan, Y., & Jiang, R. (2013). Roles of

Bmp4 during tooth morphogenesis and sequential tooth formation. *Development* ,

140(2), 423–432.

Johnson, N. A., & Porter, A. H. (2007). Evolution of branched regulatory genetic pathways:

directional selection on pleiotropic loci accelerates developmental system drift.

Genetica, *129*(1), 57–70.

Kalinka, A. T., Varga, K. M., Gerrard, D. T., Preibisch, S., Corcoran, D. L., Jarrells, J., Ohler,

U., Bergman, C. M., & Tomancak, P. (2010). Gene expression divergence recapitulates

the developmental hourglass model. *Nature*, *468*(7325), 811–814.

Kelley, R., Ren, R., Pi, X., Wu, Y., Moreno, I., Willis, M., Moser, M., Ross, M., Podkowa, M.,

Attisano, L., & Patterson, C. (2009). A concentration-dependent endocytic trap and sink

mechanism converts Bmper from an activator to an inhibitor of Bmp signaling. *The*

Journal of Cell Biology, *184*(4), 597–609.

Kim, J., Ahn, Y., Adasooriya, D., Woo, E. J., Kim, H. J., Hu, K. S., Krumlauf, R., & Cho, S.

W. (2019). Shh Plays an Inhibitory Role in Cusp Patterning by Regulation of Sostdc1.

Journal of Dental Research, *98*(1), 98–106.

Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J., & Peterson, H. (2020). gprofiler2 -- an R

package for gene list functional enrichment analysis and namespace conversion toolset

g:Profiler. *F1000Research*, *9*. https://doi.org/10.12688/f1000research.24956.2

Kwon, H.-J. E., Jia, S., Lan, Y., Liu, H., & Jiang, R. (2017). Activin and Bmp4 Signaling

Converge on Wnt Activation during Odontogenesis. *Journal of Dental Research*, *96*(10),

1145–1152.

Lan, Y., Jia, S., & Jiang, R. (2014). Molecular patterning of the mammalian dentition.

Seminars in Cell & Developmental Biology, *25-26*, 61–70.

Lazzari, V., Charles, C., Tafforeau, P., Vianey-Liaud, M., Aguilar, J.-P., Jaeger, J.-J.,

Michaux, J., & Viriot, L. (2008). Mosaic convergence of rodent dentitions. *PloS One*,

3(10), e3607.

Levin, M., Anavy, L., Cole, A. G., Winter, E., Mostov, N., Khair, S., Senderovich, N., Kovalev,

E., Silver, D. H., Feder, M., Fernandez-Valverde, S. L., Nakanishi, N., Simmons, D.,

Simakov, O., Larsson, T., Liu, S.-Y., Jerafi-Vider, A., Yaniv, K., Ryan, J. F., … Yanai, I.

(2016). The mid-developmental transition and the evolution of animal body plans.

Nature, *531*(7596), 637–641.

Liang, C., Musser, J. M., Cloutier, A., Prum, R. O., & Wagner, G. P. (2018). Pervasive

Correlated Evolution in Gene Expression Shapes Cell and Tissue Type Transcriptomes.

Genome Biology and Evolution, *10*(2), 538–552.

Liu, F., Chu, E. Y., Watt, B., Zhang, Y., Gallant, N. M., Andl, T., Yang, S. H., Lu, M.-M.,

Piccolo, S., Schmidt-Ullrich, R., Taketo, M. M., Morrisey, E. E., Atit, R., Dlugosz, A. A., &

Millar, S. E. (2008). Wnt/β-catenin signaling directs multiple stages of tooth

morphogenesis. *Developmental Biology*, *313*(1), 210–224.

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

Mahfooz, N., Turchyn, N., Mihajlovic, M., Hrycaj, S., & Popadić, A. (2007). Ubx regulates

differential enlargement and diversification of insect hind legs. *PloS One*, *2*(9), e866.

Maier, J. A., Rivas-Astroza, M., Deng, J., Dowling, A., Oboikovitz, P., Cao, X., Behringer, R.

R., Cretekos, C. J., Rasweiler, J. J., Zhong, S., & Sears, K. E. (2017). Transcriptomic

insights into the genetic basis of mammalian limb diversity. In *BMC Evolutionary Biology*

(Vol. 17, Issue 1). https://doi.org/10.1186/s12862-017-0902-6

Mann, R. S., & Carroll, S. B. (2002). Molecular mechanisms of selector gene function and

evolution. *Current Opinion in Genetics & Development*, *12*(5), 592–600.

Martignetti, L., Calzone, L., Bonnet, E., Barillot, E., & Zinovyev, A. (2016). ROMA:

Representation and Quantification of Module Activity from Target Expression Data.

Frontiers in Genetics, *7*, 18.

Matsuoka, Y., & Monteiro, A. (2021). Hox genes are essential for the development of

eyespots in Bicyclus anynana butterflies. *Genetics*, *217*(1), 1–9.

Matsuoka, Y., & Monteiro, A. (2022). Ultrabithorax modifies a regulatory network of genes

essential for butterfly eyespot development in a wing sector-specific manner.

Development , *149*(23). https://doi.org/10.1242/dev.200781

McKay, D. J., & Lieb, J. D. (2013). A common set of DNA regulatory elements shapes

Drosophila appendages. *Developmental Cell*, *27*(3), 306–318.

Meguro, F., Porntaveetus, T., Kawasaki, M., Kawasaki, K., Yamada, A., Kakihara, Y., Saeki,

M., Tabeta, K., Kessler, J. A., Maeda, T., & Ohazama, A. (2019). Bmp signaling in molar

cusp formation. *Gene Expression Patterns: GEP*, *32*, 67–71.

Miletich, I., Yu, W.-Y., Zhang, R., Yang, K., Caixeta de Andrade, S., Pereira, S. F. do A.,

Ohazama, A., Mock, O. B., Buchner, G., Sealby, J., Webster, Z., Zhao, M., Bei, M., &

Sharpe, P. T. (2011). Developmental stalling and organ-autonomous regulation of

morphogenesis. *Proceedings of the National Academy of Sciences of the United States*

of America, *108*(48), 19270–19275.

Milocco, L., & Salazar-Ciudad, I. (2020). Is evolution predictable? Quantitative genetics

under complex genotype-phenotype maps. *Evolution; International Journal of Organic*

Evolution, *74*(2), 230–244.

Minelli, A. (2003). The origin and evolution of appendages. *The International Journal of*

Developmental Biology, *47*(7-8), 573–581.

Mitsiadis, T. A., & Drouin, J. (2008). Deletion of the Pitx1 genomic locus affects mandibular

tooth morphogenesis and expression of the Barx1 and Tbx1 genes. *Developmental*

Biology, *313*(2), 887–896.

Mootha, V. K., Lindgren, C. M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J.,

Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M. J.,

Patterson, N., Mesirov, J. P., Golub, T. R., Tamayo, P., Spiegelman, B., Lander, E. S.,

Hirschhorn, J. N., … Groop, L. C. (2003). PGC-1α-responsive genes involved in

oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature*

Genetics, *34*(3), 267–273.

Morgalev, S. Y., Lim, A. G., Morgaleva, T. G., Morgalev, Y. N., Manasypov, R. M., Kuzmina,

D., Shirokova, L. S., Orgogozo, L., Loiko, S. V., & Pokrovsky, O. S. (2023).

Fractionation of organic C, nutrients, metals and bacteria in peat porewater and ice after

freezing and thawing. *Environmental Science and Pollution Research International*,

30(1), 823–836.

Morita, W., Morimoto, N., & Jernvall, J. (2020). Mapping molar shapes on signaling

pathways. *PLoS Computational Biology*, *16*(12), e1008436.

Morita, W., Morimoto, N., Otsu, K., & Miura, T. (2022). Stripe and spot selection in cusp

patterning of mammalian molar formation. *Scientific Reports*, *12*(1), 9149.

Museums Victoria Collections;

https://collections.museumsvictoria.com.au/specimens/116894

Musser, J. M., & Wagner, G. P. (2015). Character trees from transcriptome data: Origin and

individuation of morphological characters and the so-called "species signal." *Journal of*

Experimental Zoology. Part B, Molecular and Developmental Evolution, *324*(7), 588–

604.

Navarro, N., & Murat Maga, A. (2018). Genetic mapping of molar size relations identifies

inhibitory locus for third molars in mice. *Heredity*, *121*(1), 1–11.

O'Connell, D. J., Ho, J. W. K., Mammoto, T., Turbe-Doan, A., O'Connell, J. T., Haseley, P.

S., Koo, S., Kamiya, N., Ingber, D. E., Park, P. J., & Maas, R. L. (2012). A Wnt-bmp

feedback circuit controls intertissue signaling dynamics in tooth organogenesis. *Science*

Signaling, *5*(206), ra4.

Pantalacci, S., Guéguen, L., Petit, C., Lambert, A., Peterkovà, R., & Sémon, M. (2017).

Transcriptomic signatures shaped by cell proportions shed light on comparative

developmental biology. *Genome Biology*, *18*(1), 29.

Pantalacci, S., & Sémon, M. (2015). Transcriptomics of developing embryos and organs: A

raising tool for evo-devo: TRANSCRIPTOMICS OF DEVELOPING EMBRYOS AND

ORGANS. *Journal of Experimental Zoology. Part B, Molecular and Developmental*

Evolution, *324*(4), 363–371.

Pantalacci, S., Sémon, M., Martin, A., Chevret, P., & Laudet, V. (2009). Heterochronic shifts

explain variations in a sequentially developing repeated pattern: palatal ridges of muroid

rodents. *Evolution & Development*, *11*(4), 422–433.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.

Paul, R., Giraud, G., Domsch, K., Duffraisse, M., Marmigère, F., Khan, S., Vanderperre, S., Lohmann, I., Stoks, R., Shashidhara, L. S., & Merabet, S. (2021). Hox dosage contributes to flight appendage morphology in Drosophila. In *Nature Communications* (Vol. 12, Issue 1). https://doi.org/10.1038/s41467-021-23293-8

Pavlicev, M., & Wagner, G. P. (2012). A model of developmental evolution: selection, pleiotropy and compensation. *Trends in Ecology & Evolution*, *27*(6), 316–322.

Pavlopoulos, A., & Akam, M. (2011). Hox gene Ultrabithorax regulates distinct sets of target genes at successive stages of Drosophila haltere morphogenesis. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(7), 2855–2860.

Peter, I. S., & Davidson, E. H. (2011). Evolution of gene regulatory networks controlling body plan development. *Cell*, *144*(6), 970–985.

Peterka, M., Lesot, H., & Peterková, R. (2002). Body weight in mouse embryos specifies staging of tooth development. *Connective Tissue Research*, *43*(2-3), 186–190.

Qiu, M., Bulfone, A., Ghattas, I., Meneses, J. J., Christensen, L., Sharpe, P. T., Presley, R., Pedersen, R. A., & Rubenstein, J. L. (1997). Role of the Dlx homeobox genes in proximodistal patterning of the branchial arches: mutations of Dlx-1, Dlx-2, and Dlx-1 and -2 alter morphogenesis of proximal skeletal and soft tissue structures derived from the first and second arches. *Developmental Biology*, *185*(2), 165–184.

Ranwez, V., Harispe, S., Delsuc, F., & Douzery, E. J. P. (2011). MACSE: Multiple Alignment of Coding SEquences Accounting for Frameshifts and Stop Codons. In *PLoS ONE* (Vol. 6, Issue 9, p. e22594). https://doi.org/10.1371/journal.pone.0022594

Ravisankar, P., Lai, Y.-T., Sambrani, N., & Tomoyasu, Y. (2016). Comparative developmental analysis of Drosophila and Tribolium reveals conserved and diverged

roles of abrupt in insect wing evolution. *Developmental Biology*, *409*(2), 518–529.

Refki, P. N., Armisén, D., Crumière, A. J. J., Viala, S., & Khila, A. (2014). Emergence of

tissue sensitivity to Hox protein levels underlies the evolution of an adaptive

morphological trait. *Developmental Biology*, *392*(2), 441–453.

Sadier, A., Twarogowska, M., Steklikova, K., Hayden, L., Lambert, A., Schneider, P., Laudet,

V., Hovorakova, M., Calvez, V., & Pantalacci, S. (2019). Modeling Edar expression

reveals the hidden dynamics of tooth signaling center patterning. *PLoS Biology*, *17*(2),

e3000064.

Sadier, A., Urban, D. J., Anthwal, N., Howenstine, A. O., Sinha, I., & Sears, K. E. (2021).

Making a bat: The developmental basis of bat evolution. *Genetics and Molecular*

Biology, *43*(1 Suppl 2), e20190146.

Salazar-Ciudad, I. (2012). Tooth patterning and evolution. *Current Opinion in Genetics &*

Development, *22*(6), 585–592.

Salazar-Ciudad, I., & Jernvall, J. (2010). A computational model of teeth and the

developmental origins of morphological variation. *Nature*, *464*(7288), 583–586.

Saxena, A., Sharma, V., Muthuirulan, P., Neufeld, S. J., Tran, M. P., Gutierrez, H. L., Chen,

K. D., Erberich, J. M., Birmingham, A., Capellini, T. D., Cobb, J., Hiller, M., & Cooper, K.

L. (2022). Interspecies transcriptomics identify genes that underlie disproportionate foot

growth in jerboas. *Current Biology: CB*, *32*(2), 289–303.e6.

Serpe, M., Umulis, D., Ralston, A., Chen, J., Olson, D. J., Avanesov, A., Othmer, H.,

O'Connor, M. B., & Blair, S. S. (2008). The BMP-binding protein Crossveinless 2 is a

short-range, concentration-dependent, biphasic modulator of BMP signaling in

Drosophila. *Developmental Cell*, *14*(6), 940–953.

Shimizu, T., Oikawa, H., Han, J., Kurose, E., & Maeda, T. (2004). Genetic analysis of crown

size in the first molars using SMXA recombinant inbred mouse strains. *Journal of Dental*

Research, *83*(1), 45–49.

Siomava, N., Fuentes, J. S. M., & Diogo, R. (2020). Deconstructing the long-standing a priori

assumption that serial homology generally involves ancestral similarity followed by

anatomical divergence. *Journal of Morphology*, *281*(9), 1110–1132.

Stern, D. L. (1998). A role of Ultrabithorax in morphological differences between Drosophila

species. *Nature*, *396*(6710), 463–466.

Tanaka, K., Barmina, O., Sanders, L. E., Arbeitman, M. N., & Kopp, A. (2011). Evolution of

sex-specific traits through changes in HOX-dependent doublesex expression. *PLoS*

Biology, *9*(8), e1001131.

Tiphaine, C., Yaowalak, C., Cyril, C., Helder, G.-R., Jacques, M., Paul, T., Monique, V.-L.,

Laurent, V., & Vincent, L. (2013). Correlated changes in occlusal pattern and diet in

stem Murinae during the onset of the radiation of Old World rats and mice. *Evolution;*

International Journal of Organic Evolution, *67*(11), 3323–3338.

Tomoyasu, Y. (2017). Ultrabithorax and the evolution of insect forewing/hindwing

differentiation. *Current Opinion in Insect Science*, *19*, 8–15.

Tomoyasu, Y., Arakane, Y., Kramer, K. J., & Denell, R. E. (2009). Repeated co-options of

exoskeleton formation during wing-to-elytron evolution in beetles. *Current Biology: CB*,

19(24), 2057–2065.

Torres Cleuren, Y. N., Ewe, C. K., Chipman, K. C., Mears, E. R., Wood, C. G., Al-Alami, C.

E. A., Alcorn, M. R., Turner, T. L., Joshi, P. M., Snell, R. G., & Rothman, J. H. (2019).

Extensive intraspecies cryptic variation in an ancient embryonic gene regulatory

network. *eLife*, *8*. https://doi.org/10.7554/eLife.48220

True, J. R., & Haag, E. S. (2001). Developmental system drift and flexibility in evolutionary

trajectories. *Evolution & Development*, *3*(2), 109–119.

Tschopp, P., Sherratt, E., Sanger, T. J., Groner, A. C., Aspiras, A. C., Hu, J. K., Pourquié,

O., Gros, J., & Tabin, C. J. (2014). A relative shift in cloacal location repositions external

genitalia in amniote evolution. *Nature*, *516*(7531), 391–394.

Tulchinsky, A. Y., Johnson, N. A., Watt, W. B., & Porter, A. H. (2014). Hybrid incompatibility

arises in a sequence-based bioenergetic model of transcription factor binding. *Genetics*,

198(3), 1155–1166.

Urdy, S., Goudemand, N., & Pantalacci, S. (2016). Looking Beyond the Genes: The Interplay Between Signaling Pathways and Mechanics in the Shaping and Diversification of Epithelial Tissues. *Current Topics in Developmental Biology*, *119*, 227–290.

Wang, W., & Yan, J. (2021). Shape-restricted regression splines with R package splines2. *Journal of Data Science: JDS*, 498–517.

Wang, Z., Dai, M., Wang, Y., Cooper, K. L., Zhu, T., Dong, D., Zhang, J., & Zhang, S. (2014). Unique expression patterns of multiple key genes associated with the evolution of mammalian flight. *Proceedings. Biological Sciences / The Royal Society*, *281*(1783), 20133133.

Wang, Z., Young, R. L., Xue, H., & Wagner, G. P. (2011). Transcriptomic analysis of avian digits reveals conserved and derived digit identities in birds. *Nature*, *477*(7366), 583–586.

Weatherbee, S. D., Behringer, R. R., Rasweiler, J. J., 4th, & Niswander, L. A. (2006). Interdigital webbing retention in bat wings illustrates genetic changes underlying amniote limb diversification. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(41), 15103–15107.

Weatherbee, S. D., & Carroll, S. B. (1999). Selector genes and limb identity in arthropods and vertebrates. *Cell*, *97*(3), 283–286.

Wotton, K. R., Jiménez-Guri, E., Crombach, A., Janssens, H., Alcaine-Colet, A., Lemke, S., Schmidt-Ott, U., & Jaeger, J. (2015). Quantitative system drift compensates for altered maternal inputs to the gap gene network of the scuttle fly Megaselia abdita. *eLife*, *4*. https://doi.org/10.7554/eLife.04785

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X., & Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. In *The Innovation* (Vol. 2, Issue 3, p. 100141). https://doi.org/10.1016/j.xinn.2021.100141

1912 Yokomizo, T., & Dzierzak, E. (2010). Three-dimensional cartography of hematopoietic

1913   clusters in the vasculature of whole mouse embryos. *Development* , *137*(21), 3651–

1914   3661.

1915 Zakin, L., Metzinger, C. A., Chang, E. Y., Coffinier, C., & De Robertis, E. M. (2008).

1916   Development of the vertebral morphogenetic field in the mouse: interactions between

1917   Crossveinless-2 and Twisted Gastrulation. *Developmental Biology*, *323*(1), 6–18.

1918 Zhang, Z., Lan, Y., Chai, Y., & Jiang, R. (2009). Antagonistic actions of Msx1 and Osr2

1919   pattern mammalian teeth into a single row. *Science*, *323*(5918), 1232–1234.

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929