**Heritability estimates of antler and body traits in white-tailed deer (*Odocoileus virginianus*)**

**from genomic-relatedness matrices**

Aidan Jamieson[1], Spencer J Anderson[2], Jérémie Fuller[3], Steeve D Côté[3], Joseph M Northrup[2,4],

Aaron BA Shafer[2,5.]

1. Biology Department, Trent University, K9J 7B8. Peterborough, Canada.

2. Environmental and Life Sciences Graduate Program, Trent University, K9J 7B8. Peterborough,

Canada.

3. Département de biologie, Centre d'études nordiques & NSERC Industrial Research Chair in

integrated resource management of Anticosti Island, Université Laval, G1V 0A6. Québec, QC,

Canada

4. Wildlife Research and Monitoring Section, Ontario Ministry of Natural Resources and

Forestry, Trent University, K9J 7B8. Peterborough, Canada.

5. Forensics Program Trent University, K9J 7B8 Peterborough, Canada.

Correspondence: aaronshafer@trentu.ca AND https://orcid.org/0000-0001-7652-225X

**Keywords**

**Abstract**

Estimating heritability ($h^2$) is required to predict the response to selection and is useful in species that are managed or farmed using trait information. Estimating $h^2$ in free-ranging populations is challenging due to the need for pedigrees; genomic-relatedness matrices (GRMs) circumvent this need and can be implemented in nearly any system where phenotypic and SNP data are available. We estimated the heritability of five body and three antler traits in a free-ranging population of white-tailed deer (*Odocoileus virginianus*) on Anticosti Island, Quebec, Canada. We generated GRMs from >10,000 SNPs: dressed body mass and peroneus muscle mass had moderate $h^2$ values of 0.49 and 0.56, respectively. Heritability in male-only antler features ranged from 0.00 to 0.51 and had high standard errors. We explored the influence of minor allele frequency and data completion filters on $h^2$: GRMs derived from fewer SNPs had reduced $h^2$ estimates and the relatedness coefficients significantly deviated from those generated with more SNPs. As a corollary, we discussed limitations to the application of GRMs in the wild, notably how skewed GRMs increase variance around $h^2$ estimates. This is the first study to estimate $h^2$ on a free-ranging population of white-tailed deer and should be informative for breeding designs and management.

How individuals respond to selection is dependent, in part, on the heritability ($h^2$) of the focal traits (Wright, 1920; Falconer and Mackay, 1996; Gillespie, 2004). Quantifying $h^2$ is essential to predicting adaptive responses (e.g. Dziedzic *et al.*, 2019), opening the potential for gene targeted management and conservation (Kardos and Shafer, 2018). Classically, $h^2$ was estimated by comparing the phenotypic variance ($V_P$) of known relatives, which allowed for estimating additive genetic variance ($V_A$) with narrow-sense heritability being $h^2 = V_A / V_P$ (Wray and Visscher, 2008). In free-ranging populations this estimation has required the construction of pedigrees that were then integrated into a mixed-effects model, known as the animal model (Wilson *et al.*, 2010). The animal model disentangles $V_P$ into its additive ($V_A$) and environmental components, enabling $h^2$ to be estimated.

The tracking of individuals and construction of pedigrees in free-ranging populations is time-consuming and costly (Clutton Brock and Sheldon, 2010). As a result, alternative pedigree-free approaches have emerged in which molecular genealogies can be structured into relatedness matrices that replace the pedigree in an animal model (Frentiu *et al.*, 2008). These genealogies have relied on microsatellites, which can be problematic due to biased estimates in small sample sizes (Wang, 2017) and the recommended number of loci exceeding that used in most studies (Kopps *et al.*, 2015). This changed with the advent of high-throughput sequencing that made the generation of thousands of single nucleotide polymorphisms (SNPs) feasible for most organisms (Andrews *et al.*, 2016). Genomic-relatedness matrices (GRM) can be generated and used to estimate the total amount of $V_P$ captured by the panel of SNPs (Yang *et al.*, 2011; Stanton-Geddes *et al.*, 2013). Here, $V_A$ is estimated across SNPs (see Ritland, 1996) and has produced $h^2$ estimates similar to those of sibling studies (Yang *et al.*, 2010). Though few, recent applications to wild populations have paid special attention to the impact of model choice and influence of missing data and minor allele frequencies (MAF) on number of retained SNPs and $h^2$ estimates (e.g. Perrier *et al.*, 2018; Gervais *et al.*, 2019).

Quantifying $h^2$ of specific traits in wild species that are farmed or harvested can be valuable for breeding design (Guo *et al.*, 2018) and management decisions (Coltman, 2008). White-tailed deer (*Odocoileus virginianus*) are a big game species with economic and cultural values across North America. Annual revenue from hunting is in the billions (Cambronne, 2013)

and close to 50 million USD from deer farming (see https://www.nass.usda.gov/AgCensus/).

Antler and body size are positively correlated with male reproductive success (Newbolt *et al.*,

2017), and $h^2$ for these traits have been estimated in captively bred populations. For example,

$h^2$ was estimated using parent-offspring trios at 0.32 for the number of antler points and

for0.45 main beam length, though with high standard errors (Michel *et al.*, 2016). Similarly, $h^2$

was estimated between 0.58 and 0.64 for body mass (Williams *et al.*, 1994). Recent work

suggests a highly polygenic nature of these traits (Anderson *et al.*, 2019), but $h^2$ has yet to be

quantified in a free-ranging population of white-tailed deer. Theoretical and empirical evidence

suggests that bottlenecked populations, such as those we might expect to see in captivity, have

increased $V_A$ (Goodnight, 1988; Whitlock *et al.*, 1993; van Heerwaarden *et al.*, 2008; Jarvis *et al.*,

2011; Taft and Roff, 2012), thus it is unclear how $h^2$ values transfer from captivity to the wild.

Here, we take advantage of an extensive phenotypic database and DNA archive to estimate $h^2$

in antler and body traits of free-ranging white-tailed deer using GRMs.

**Materials and Methods**

**Individual Sampling and phenotypes**

Ear or muscle tissue was obtained and preserved in 95% ethanol between 2003 and 2014 from

harvested white-tailed deer on Anticosti Island, Quebec, Canada (Figure S1). This population

originated from ~220 individuals introduced in the late 1800s and shows no evidence of a

founder effect (Fuller *et al.*, 2020). All individuals had an extensive set of phenotypic metrics

taken (Table S1), were sexed, and aged using cementum layers in incisor teeth (Hamlin *et al.*,

2000) with only animals >1.5 years analyzed. Phenotypes were assessed for correlation and

only one was retained if Spearman's *r* was > 0.70. DNA was isolated from the tissue using a

phenol-chlorophorm-isoamyl procedure and stored at -80 °C.

**Genome relatedness matrix and genotyping**

We used the SNP data generated in Fuller *et al.*, (2020) that applied a rigorous filtering

approach to generate 13,420 SNPs. We took this VCF file and further filtered it to include only

scaffolds that had >1 SNP present, and made a separate data file that only included males. To

quantify the impact of loci missingness (LM) and MAF filters on downstream inferences (e.g.

Shafer *et al.*, 2017) we generated subsets of data with 70, 80, 90 and 100% completion as well

1, 5 and 10% MAF requirements. The SNP genotypes were converted into a binary fileset using

PLINK v.19 (Purcell *et al.*, 2007) that was used to generate a GRM in GCTA v1.92.4 (Yang *et al.*,

2011). The GRMs were input to a mixed-effects linear model that included year of harvest, age,

and sex with a restricted maximum-likelihood (REML) approach to estimate $h^2$ (Meyer, 1989).

An F test of equality of variances was conducted in R v. 3.6.3. to test whether the GRM

distributions differed among the filtered datasets, and we tested whether deviations from a 1:1

ratio, calculated as abs(1-F statistic), correlated to differences in the number of SNPs used to

build the GRMs.

For a subset of male samples we genotyped the SRP54 locus that was previously found

to be associated with antler morphology (Anderson *et al.*, 2019). We custom designed a rhAmp

SNP assay and genotyped individuals via qPCR with a QuantStudio 3 Real-Time PCR System

(Thermo Fisher Scientific, Waltham, MA, USA). The thermocycling condition was 60 °C for 30

seconds followed by 95 °C for 10 minutes, then 40 cycles of 95 °C for 10 seconds, 60 °C for 30

seconds, and 68 °C for 20 seconds, followed by a final 30 seconds at 60 °C. The qPCR had a 5 µL

reaction volume in each 0.1 mL well of a sealed 96-well optical plate. As above the GRMs were

fitted to include year of harvest, age, and the SRP54 locus in an additive framework (0, 1, or 2

alleles).

Scripts have been uploaded to a GIT repository: https://gitlab.com/WiDGeT_TrentU

## Results

We obtained samples from 219 male and 218 female deer. The number of SNPs retained for

body size and antler estimates are provided in Table 1. Increased loci completeness

requirements and higher MAF values reduced the number of SNPs used to build the GRM, with

the final SNP datasets containing between 1,041 and 10,386 SNPs. The distribution of

relatedness values in the GRM showed primarily unrelated individuals, though there was

evidence of full-sibling or parent-offspring pairings (Figure 1). The filtered GRM distributions

differed with increased MAF and LM filters that skewed the relatedness coefficients (Table 2,

Figure 2). Variance between GRMs (i.e. abs(1-F-statistic)) and the difference in number of SNPs

used to build the GRMs were highly correlated ($\beta$ = 1.63e-05, $R^2$ = 0.92, P < 0.001).

Dressed body mass and peroneus muscle mass for males and females together had

heritability estimates of 0.49 and 0.56 respectively; all other body metrics had relatively large

standard errors (SE; Table 3). Likewise, two male-only antler metrics had non-zero $h^2$ estimates

but high SE (Table 3). Smaller $h^2$ estimates were observed from GRMs built with fewer SNPs

(Table S2). Not surprisingly, age and sex had large effects on the body size metrics, with older

individuals and males being larger on average; antler metrics also increased with age (Table 3).

Estimates of $h^2$ were lower when age and sex were not included in the model (Table S2),

whereas removing year of harvest only had a minor effect (Tables S3; S4). We genotyped 74

males at the SRP54 locus (Dryad Accession XXXX); despite a positive effect size of this gene on

antler metrics, all estimates had high SE precluding any meaningful inference (Table S5; S6).

**Discussion**

White-tailed deer are an important big-game species that are hunted and farmed across North

America. If larger antlered or bodied animals are preferentially harvested or bred, there is the

potential for artificial selection provided such traits are heritable. Our data suggest that body

and muscle mass are heritable (Table 2), which is consistent with most vertebrates (Mousseau

and Roff, 1987) including deer (Williams *et al.*, 1994). For antler metrics, despite $h^2$ values

similar to previous estimates (i.e. Lukefahr and Jacobson, 1998; Michel *et al.*, 2016), our

estimates had large standard errors; this is likely a reflection of the smaller male-only data set

and a GRM skewed towards unrelated individuals (Figure 1) as captive studies suggest

moderate levels of heritability. Whether harvest could induce artificial selection in white-tailed

deer is an open question: some data have shown harvested antlers being larger than those

naturally shed (Schoenebeck and Peterson, 2014; but see Ditchkoff *et al.*, 2000), which could

generate an evolutionary response. Still, non-genetic factors are large contributors to body size

and antlers (Table 2) and in our view it is unlikely that harvest is a strong enough selective

pressure to reduce antler size in most managed regions, given the generally large population

size of deer, varied regulations, and often opportunistic nature of the hunt. Small and isolated

populations, including farms, could see a response to selection, especially if $V_A$ is higher (e.g.
Taft and Roff, 2012).

The heritability of traits can vary by sex, age, and across seasons (Réale *et al.*, 1999).
This is an important consideration with GRM approaches for phenotypically stratified
populations that despite circumventing the need for pedigrees, still require basic individual
metadata. In our study this meant aging individuals through cross-sections of incisors (Hamlin
et al. 2000), and not accounting for age and sex in the model resulted in downward biased $h^2$
estimates (Table S2). Likewise, accounting for parturition date and litter size increased $h^2$
estimates in captive white-tailed deer (Michel et al. 2016). Interestingly, including the year of
harvest that presumably reflect environmental variation, particularly during summer, had a
largely negligible effect on $h^2$ (Table S3; S4). The relatively high standard errors surrounding $h^2$,
even after accounting for age and sex, is not uncommon as seen in Michel et al.'s (2016)
analysis of captive deer, and more broadly ungulates (Thomas *et al.*, 2002; Coltman *et al.*,
2005). However, Sim and Coltman (2019) used GRMs and reported moderate $h^2$ estimates with
small SE of horn characteristics of thinhorn sheep (*Ovis dalli*): this difference is likely attributed
to antlers being shed and regrown annually, and horn length being a more easily quantifiable
metric.

Sample size, SNP number, and filtering criteria all must be considered when estimating
$h^2$ with GRMs (Visscher and Goddard, 2015; Gervais *et al.*, 2019). Small sample size increases
sampling variance, typically biasing $V_A$ downwards (Villemereuil *et al.*, 2013). With SNP data,
the GRM variance is inversely proportional to that of $h^2$ (Visscher and Goddard, 2015). Both our
analysis and two other wild ungulate studies (Bérénos *et al.*, 2014; Gervais *et al.*, 2019) had
GRMs skewed towards distant relatives; this undoubtedly played a role in $h^2$ estimates, as
sampling the full spectrum of relatives is difficult in wild populations. The use of SNP data also
involves bioinformatics decisions. High MAF cutoffs should be avoided as this will shift the site-
frequency spectrum towards more common alleles, resulting in reduced $h^2$ estimates (Table S2;
Gervais et al. 2019) and appreciably different GRMs (Table 2; Figure 2). Introducing a stringent
loci missingness criteria in principle should improve estimates (Gervais et al. 2019), though we
saw minimal evidence of this, likely due to the reduced SNP number (Table 1). Collectively,

these filtering parameters all impact the quality and number of SNPs used to build the GRM,

with wild vertebrate studies using a few thousand (Gervais *et al.*, 2019; Sim and Coltman, 2019)

to tens of thousands SNPs (Bérénos *et al.*, 2014; Perrier *et al.*, 2018). While the optimal sample

size and SNP number will depend on demographic history and life history strategies of the focal

population, the generation of GRMs and estimates of $h^2$ in the wild opens up numerous basic

and applied research opportunities, notably predicting the response to selection under harvest

and climate change scenarios. In the case of white-tailed deer, we have shown that body size

and likely antlers are moderately heritable and could respond to selection, which would be

most likely in small and isolated populations.

## Acknowledgements

This work was supported by the Natural Sciences and Engineering Research Council of Canada

Discovery Grants (ABAS and SDC); ComputeCanada Resources for Research Groups (ABAS);

Canadian Foundation for Innovation: John R. Evans Leaders Fund (ABAS); and Industrial Chairs

and Collaborative Research and Development Grants from the Natural Sciences and

Engineering Research Council of Canada (SDC). We thank the outfitters of Anticosti Island and

the Ministère des Forêts, de la Faune et des Parcs du Québec for logistical help associated with

fieldwork, and all of the field assistants and technicians who collected samples and made

phenotypic measurements. Special thanks to Louis Bernatchez for supporting the lab work.

## References

Anderson SJ, Côté SD, Richard JH, Shafer AB (2019). Genomic architecture of artificially and
sexually selected traits in a wild cervid. *bioRxiv*: 841528.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA (2016). Harnessing the power of
RADseq for ecological and evolutionary genomics. *Nat Rev Genet* **17**: 81–92.

Bérénos C, Ellis PA, Pilkington JG, Pemberton JM (2014). Estimating quantitative genetic
parameters in wild populations: a comparison of pedigree and genomic approaches.
*Molecular Ecology* **23**: 3434–3451.

Cambronne A (2013). *Deerland: America's hunt for ecological balance and the essence of*
*wildness*. Rowman & Littlefield.

Clutton Brock T, Sheldon BC (2010). Individuals and populations: the role of long-term,
211           individual-based studies of animals in ecology and evolutionary biology. *Trends Ecol Evol*
212           **25**: 562–573.

213      Coltman DW (2008). Molecular ecological approaches to studying the evolutionary impact of
214           selective harvesting in wildlife. *Molecular Ecology* **17**: 221–235.

215      Coltman DW, O'Donoghue P, Hogg JT, Festa-Bianchet M (2005). Selection and genetic
216           (co)variance in Bighorn sheep. *Evolution* **59**: 1372–1382.

217      Ditchkoff SS, Jr ERW, Lochmiller RL (2000). Using cast antler characteristics to profile quality of
218           white-tailed deer Odocoileus virginianus populations. *wbio* **6**: 53–58.

219      Dziedzic KE, Elder H, Tavalire H, Meyer E (2019). Heritable variation in bleaching responses and
220           its functional genomic basis in reef-building corals (Orbicella faveolata). *Molecular*
221           *Ecology* **28**: 2238–2253.

222      Falconer D, Mackay T (1996). Introduction to quantitative genetics. Essex. *UK: Longman Group.*

223      Frentiu FD, Clegg SM, Chittock J, Burke T, Blows MW, Owens IPF (2008). Pedigree-free animal
224           models: the relatedness matrix reloaded. *Proc Roy Soc B* **275**: 639–647.

225      Fuller J, Ferchaud A-L, Laporte M, Luyer JL, Davis TB, Côté SD, *et al.* (2020). Absence of founder
226           effect and evidence for adaptive divergence in a recently introduced insular population
227           of white-tailed deer (Odocoileus virginianus). *Molecular Ecology* **29**: 86–104.

228      Gervais L, Perrier C, Bernard M, Merlet J, Pemberton JM, Pujol B, *et al.* (2019). RAD-sequencing
229           for estimating genomic relatedness matrix-based heritability in the wild: A case study in
230           roe deer. *Molecular Ecology Resources* **19**: 1205–1217.

231      Gillespie JH (2004). *Population genetics: a concise guide.* JHU Press.

232      Goodnight CJ (1988). Epistasis and the effect of founder events on the additive genetic
233           variance. *Evolution* **42**: 441–454.

234      Guo H, Zeng Q, Li Y, Wang Y, Chen Z, Lin P, *et al.* (2018). Estimating realized heritability for
235           growth in Zhikong scallop (Chlamys farreri) using genome-wide complex trait analysis.
236           *Aquaculture* **497**: 103–108.

237      Hamlin KL, Pac DF, Sime CA, DeSimone RM, Dusek GL (2000). Evaluating the accuracy of ages
238           obtained by two methods for Montana ungulates. *The Journal of Wildlife Management*
239           **64**: 441–449.

240      van Heerwaarden B, Willi Y, Kristensen TN, Hoffmann AA (2008). Population bottlenecks
241           increase additive genetic variance but do not break a selection limit in rain rorest
242           drosophila. *Genetics* **179**: 2135–2146.

Jarvis JP, Cropp SN, Vaughn TT, Pletscher LS, King-Ellison K, Adams-Hunt E, *et al.* (2011). The
effect of a population bottleneck on the evolution of genetic variance/covariance
structure. *Journal of Evolutionary Biology* **24**: 2139–2152.

Kardos M, Shafer AB (2018). The peril of gene-targeted conservation. *Trends in ecology &*
*evolution* **33**: 827–839.

Kopps AM, Kang J, Sherwin WB, Palsbøll PJ (2015). How well do molecular and pedigree
relatedness correspond, in populations with diverse mating systems, and various types
and quantities of molecular and demographic data? *G3: Genes, Genomes, Genetics* **5**:
1815–1826.

Lukefahr SD, Jacobson HA (1998). Variance component analysis and heritability of antler traits
in white-tailed deer. *The Journal of Wildlife Management* **62**: 262–268.

Meyer K (1989). Restricted maximum likelihood to estimate variance components for animal
models with several random effects using a derivative-free algorithm. *Genetics Selection*
*Evolution* **21**: 317.

Michel ES, Demarais S, Strickland BK, Smith T, Dacus CM (2016). Antler characteristics are highly
heritable but influenced by maternal factors. *Jour Wild Mgmt* **80**: 1420–1426.

Mousseau TA, Roff DA (1987). Natural selection and the heritability of fitness components.
*Heredity* **59**: 181–197.

Newbolt CH, Acker PK, Neuman TJ, Hoffman SI, Ditchkoff SS, Steury TD (2017). Factors
influencing reproductive success in male white-tailed deer. *The Journal of Wildlife*
*Management* **81**: 206–217.

Perrier C, Delahaie B, Charmantier A (2018). Heritability estimates from genomewide
relatedness matrices in wild populations: Application to a passerine, using a small
sample size. *Molecular Ecology Resources* **18**: 838–853.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, *et al.* (2007). PLINK: A tool
set for whole-genome association and population-based linkage analyses. *The American*
*Journal of Human Genetics* **81**: 559–575.

Réale D, Festa-Bianchet M, Jorgenson JT (1999). Heritability of body mass varies with age and
season in wild bighorn sheep. *Heredity* **83**: 526–532.

Ritland K (1996). A marker-based method for inferences about quantitative inheritance in
natural populations. *Evolution* **50**: 1062.

Schoenebeck CW, Peterson BC (2014). Evaluation of hunter antler-size selection through an
age-specific comparison of harvested and naturally cast antler metrics. *Journal of Fish*
*and Wildlife Management* **5**: 167–173.

Shafer ABA, Peart CR, Tusso S, Maayan I, Brelsford A, Wheat CW, *et al.* (2017). Bioinformatic
278            processing of RAD-seq data dramatically impacts downstream population genetic
279            inference. *Methods Ecol Evol* **8**: 907–917.

280    Sim Z, Coltman DW (2019). Heritability of horn size in Thinhorn sheep. *Front Genet* **10**.

281    Stanton-Geddes J, Yoder JB, Briskine R, Young ND, Tiffin P (2013). Estimating heritability using
282            genomic data. *Methods in Ecology and Evolution* **4**: 1151–1158.

283    Taft HR, Roff DA (2012). Do bottlenecks increase additive genetic variance? *Conservation
284            Genetics* **13**: 333–342.

285    Thomas SC, Coltman DW, Pemberton JM (2002). The use of marker-based relationship
286            information to estimate the heritability of body weight in a natural population: a
287            cautionary tale. *Journal of Evolutionary Biology* **15**: 92–99.

288    Villemereuil P de, Gimenez O, Doligez B (2013). Comparing parent–offspring regression with
289            frequentist and Bayesian animal models to estimate heritability in wild populations: a
290            simulation study for Gaussian and binary traits. *Methods in Ecology and Evolution* **4**:
291            260–275.

292    Visscher PM, Goddard ME (2015). A General Unified Framework to Assess the Sampling
293            Variance of Heritability Estimates Using Pedigree or Marker-Based Relationships.
294            *Genetics* **199**: 223–232.

295    Wang J (2017). Estimating pairwise relatedness in a small sample of individuals. *Heredity* **119**:
296            302–313.

297    Whitlock MC, Phillips PC, Wade MJ (1993). Gene interaction affects the additive genetic
298            variance in subdivided populations with migration and extinction. *Evolution* **47**: 1758–
299            1769.

300    Williams JD, Krueger W, Harmel DH (1994). Heritabilities for antler characteristics and body
301            weight in yearling white-tailed deer. *Heredity* **73**: 78–83.

302    Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, *et al.* (2010). An
303            ecologist's guide to the animal model. *J Anim Ecol* **79**: 13–26.

304    Wray N, Visscher P (2008). Estimating trait heritability. *Nature education* **1**: 29.

305    Wright S (1920). The relative importance of heredity and environment in determining the
306            piebald pattern of guinea-pigs. *Proceedings of the National Academy of Sciences of the
307            United States of America* **6**: 320.

308    Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, *et al.* (2010). Common SNPs
309            explain a large proportion of the heritability for human height. *Nature genetics* **42**: 565.

Yang J, Lee SH, Goddard ME, Visscher PM (2011). GCTA: a tool for genome-wide complex trait
analysis. *The American Journal of Human Genetics* **88**: 76–82.

**Figure Captions**

**Figure 1.** Genomic-relatedness matrices (GRMs) for white-tailed deer (*Odocoileus virginianus*)

on Anticosti Island, Quebec, Canada. GRMs are divided into the entire dataset (male and

female) and male only.

**Figure 2.** Genomic-relatedness matrices (GRMs) for the sampled white-tailed deer (*Odocoileus*

*virginianus*). GRMs are from the entire dataset (male and female) and were generated with

varying minor allele frequency (MAF) and loci missing (LM) filters.

**Table 1.** Number of SNPs and individuals retained in each genomic-relatedness matrix (GRM)
after standard filtering. Additional filtering was only applied to full male & female datasets
using different minor allele frequency (MAF) required loci completion (LM) criteria.
327

| GRM | No. of SNPs | No. of Individuals |
|---|---|---|
| **Male & female** | | |
| MAF 1%, LM 70% | 10386 | 437 |
| MAF 5%, LM 70% | 9723 | 437 |
| MAF 10%, LM 70% | 2344 | 437 |
| MAF 1%, LM 80% | 10380 | 437 |
| MAF 1%, LM 90% | 10148 | 437 |
| MAF 1%, LM 100% | 1041 | 437 |
| MAF 5%, LM 80% | 9719 | 437 |
| **Male only** | | |
| MAF 1%, LM 70% | 10386 | 219 |

328
329

330  **Table 2.** Pairwise F tests of variances of the filtered genomic-relatedness matrices (GRM) in
331  white-tailed deer (*Odocoileus virginanus*). We used different minor allele frequency (MAF)
332  required loci completion (LM) criteria to generate GRMs. The F statistics and P values are
333  presented, and *n* is 436. ΔSNPs refers to the difference in no. SNPs used to build the GRMs.
334

| GRM 1 | GRM 2 | ΔSNPs | F statistic | P value |
|---|---|---|---|---|
| MAF 1%, LM 70% | MAF 5%, LM 70% | 663 | 0.99 | 0.38 |
| MAF 1%, LM 70% | MAF 10%, LM 70% | 8042 | 0.86 | <0.001 |
| MAF 1%, LM 70% | MAF 1%, LM 80% | 6 | 1.00 | 0.93 |
| MAF 1%, LM 70% | MAF 1%, LM 90% | 238 | 1.01 | 0.002 |
| MAF 1%, LM 70% | MAF 1%, LM 100% | 9345 | 0.88 | <0.001 |
| MAF 1%, LM 70% | MAF 5%, LM 80% | 667 | 1.00 | 0.41 |
| MAF 5%, LM 70% | MAF 10%, LM 70% | 7379 | 0.86 | <0.001 |
| MAF 5%, LM 70% | MAF 1%, LM 80% | 657 | 1.00 | 0.33 |
| MAF 5%, LM 70% | MAF 1%, LM 90% | 425 | 1.02 | <0.001 |
| MAF 5%, LM 70% | MAF 1%, LM 100% | 8682 | 0.88 | <0.001 |
| MAF 5%, LM 70% | MAF 5%, LM 80% | 4 | 1.00 | 0.95 |
| MAF 10%, LM 70% | MAF 1%, LM 80% | 8036 | 1.16 | <0.001 |
| MAF 10%, LM 70% | MAF 1%, LM 90% | 7804 | 1.17 | <0.001 |
| MAF 10%, LM 70% | MAF 1%, LM 100% | 1303 | 1.01 | 0.006 |
| MAF 10%, LM 70% | MAF 5%, LM 80% | 7375 | 1.15 | <0.001 |
| MAF 1%, LM 80% | MAF 1%, LM 90% | 232 | 1.01 | 0.004 |
| MAF 1%, LM 80% | MAF 1%, LM 100% | 9339 | 0.88 | <0.001 |
| MAF 1%, LM 80% | MAF 5%, LM 80% | 661 | 1.00 | 0.36 |
| MAF 1%, LM 90% | MAF 1%, LM 100% | 9107 | 0.86 | <0.001 |
| MAF 1%, LM 90% | MAF 5%, LM 80% | 429 | 0.98 | <0.001 |
| MAF 1%, LM 100% | MAF 5%, LM 80% | 8678 | 1.14 | <0.001 |

335
336

**Table 3.** Estimation of heritability ($h^2$) for body and antler traits in white-tailed deer (*Odocoileus*
*virginianus*) on Anticosti Island, Quebec, Canada. The genomic-relatedness matrix (GRM) was
generated using a minor allele frequency of 1% and required loci completion of 70%. Mean trait
values and effect sizes of age and sex are provided but the effect of year is not shown. Standard
errors for all metrics are included in parentheses. *N* refers to the number individuals retained
after filtering for phenotype completeness.
343

| GRM | *N* | $h^2$ | Mean | Age | Sex |
|---|---|---|---|---|---|
| **Male & female GRM** | | | | | |
| Dressed body mass (kg) | 399 | 0.49 (0.24) | 32 (1.2) | 1.3 (0.1) | 13 (0.8) |
| Total length (cm) | 399 | 0.00 (0.22) | 160 (1.7) | 1.6 (0.2) | 12 (1.1) |
| Hind food length (cm) | 399 | 0.39 (0.26) | 45 (0.3) | 0.26 (0.04) | 3.1 (0.2) |
| Chest circumference (cm) | 399 | 0.24 (0.23) | 84 (1.0) | 1.0 (0.1) | 8.7 (0.7) |
| Peroneus muscle mass (g) | 399 | 0.56 (0.25) | 86 (2.5) | 2.3 (0.3) | 27 (1.7) |
| **Male only GRM** | | | | | |
| Antler spread (cm) | 171 | 0.00 (0.53) | 17 (1.8) | 3.1 (0.3) | - |
| No. of points | 182 | 0.51 (0.51) | 2.1 (0.4) | 0.63 (0.05) | - |
| Main antler beam (cm) | 175 | 0.23 (0.50) | 1.5 (0.1) | 0.17 (0.01) | - |

344
345