1   Contribution Evenness: A functional redundancy metric sensitive to trait stability in microbial

2   communities

4   Taylor M. Royalty (1,2) and Andrew D. Steen (2,1)*

5   University of Tennessee Departments of (1) Earth and Planetary Sciences and (2) Microbiology

7   Running title:

8   Measuring the functional redundancy of microbiomes

10   *Corresponding: asteen1@utk.edu

Abstract

The concept of functional redundancy has received considerable attention in both the

macroecology and microbial ecology literature. As a result, multiple metrics of functional

redundancy have been proposed. These vary in how they weight trait levels, species abundance,

functional richness, and species richness. Here we present a new functional redundancy metric

tailored for community-aggregated traits, which are traits that are quantified at the community

level and can be quantitatively partitioned among species. We call this metric Contribution

Evenness (CE) because it measures how evenly species contribute to a community-aggregated

trait. As CE is an evenness measurement, it ranges from 0 and 1, where 0 corresponds to a single

species contributing to a community-aggregated trait and 1 corresponds to all species

contributing equally. Using *in silico* simulations of species extinctions, we demonstrate that CE

reflects the stability of an ecosystem function to species extinction, a hypothesized ecological

consequence of functional redundancy. As a positive control and to illustrate how CE can be

used with sequence data, we analyzed the functional redundancy of eight nitrogen-transforming

pathways using 2,631 metagenome-assembled genomes from 47 marine sites. CE for marine

nitrogen cycle marker genes was consistent with our qualitative understanding of which nitrogen

pathways are most functionally redundant in the ocean. We found that, on average, the NH_4^+

assimilation pathway was the most functionally redundant (0.44 ± 0.08) while dissimilatory

nitrate reduction was the least redundant (0.005 ± 0.005). As demonstrated here, CE provides a

promising framework for measuring trait stability in microbiomes.

Introduction

Recent advances in sequencing technology and sampling coverage have facilitated the discovery that important ecological functions are frequently replicated across co-occurring species in an ecosystem (1). This phenomenon is referred to as "functional redundancy". The ecological implications of functional redundancy are not fully understood, but one hypothesized consequence is that it increases trait stability, which is the ability of microbial communities to retain a trait or function even when individual members of the community become locally extinct (2–4). To test hypotheses about the ecological and biogeochemical implications of functional redundancy in microbial communities, a quantitative and generally applicable metric is required.

Several metrics for functional redundancy exist in the literature. The diversity of metrics reflects the diverse sources of data used to calculate functional redundancy and the diverse purposes for which this concept has been applied. Functional redundancy has been measured as the ratio of community richness to the theoretical maximum number of functional entities (5), median metabolic overlap of genomic traits (6), expected variability in trait space among community members (7), the degree of covariance between functional diversity and biodiversity (e.g., Galand et al., 2018; Micheli and Halpern, 2005; Miki et al., 2014), and the simple fraction of populations in a community that bear a given function (11–13). These different metrics treat community structure differently. For instance, some metrics are sensitive to species abundances (7, 8), richness (5, 9), or neither (6). Except for the simple fraction of taxa bearing a function, the functional redundancy metrics above represent ecological function in terms of niche, where a niche is thought of as a point in a hyperdimensional trait space (14). Communities with higher taxa density in trait space are interpreted as being more functionally redundant.

This conception of functional redundancy is difficult to apply to microbial communities.

There are many possible, environmentally relevant microbial functions, for most microbes these

functions are known only from a microbe's genome, and the genes for these functions are often

not well-conserved. Thus, it is often impossible to define microbial niches in a meaningful and

tractable way (1). As a result, existing functional redundancy metrics can be challenging to apply

to microbial ecosystem.

Rather than defining functional redundancy in terms of niches, an alternative approach is

to evaluate functional redundancy with respect to individual functions or traits. Here, we propose

a metric designed to quantify the functional redundancy of individual traits in a microbial

community, something that is currently lacking in the literature. We focus on microbial traits

represented as nucleic acid sequences (DNA and RNA), i.e., gene and transcript sequences.

These sequence traits are regularly measured in microbial ecology studies, are used to infer

microbial metabolism, can be meaningfully compared between samples or studies, and are easily

integrated into trait-based models (15–17).

To define this metric, we have treated gene and transcript sequences as "community-

aggregated parameters" (18). These are parameters which are observed at the ecosystem level,

but which can be quantitatively partitioned among individual species. The contribution of each

species to an ecosystem-level process is calculated as the contribution of individuals to the

process, weighted by their species' abundance. Species contributions are then summed to reflect

the community-level processes. Gene and transcript abundances are a natural fit for this concept,

because the abundance of gene sequences can be quantified (e.g. as read depth) and assigned to

specific species (e.g. by mapping reads to reference genomes or metagenome-assembled

genomes) (19).

We use an evenness metric to calculate species' relative contribution to an observed

community-level parameter (20). Briefly, the "true diversity" of contributors is calculated using

species contributions and traditional diversity metrics (21, 22). The observed diversity is

normalized to a theoretical maximum diversity (richness) and made "unrelated", in the sense

used by Chao and Ricotta (20), to community richness. Our approach can be understood as

measuring functional redundancy because it measures how evenly species in a community

contribute to an ecosystem parameter—the more evenly species contribute to an ecosystem

parameter, the more functionally redundant the community is with respect to that process.

Accordingly, we term our measure of functional redundancy "contribution evenness" (CE).

To demonstrate the efficacy of CE, we first performed simulations to optimize the value

of a constant called diversity order which is required in CE calculations (21, 22). Second, we

performed an *in silico* experiment to test how well CE captures trait stability compared to other,

existing functional redundancy metrics. Finally, we demonstrate the applicability of CE with

real-world data by analyzing functional redundancy for eight nitrogen-transforming pathways

found in metagenomic data (23) in *TARA* Oceans draft metagenome-assembled genomes

(MAGs) (24). Overall, we found CE appealing because it is easily calculated from conventional

shotgun genomic sequence data sets, provides insights on the relative contribution by community

members to community-level processes, and is sensitive to the stability of traits to species

extinction.

Methods and Theory

Definition of Functional Redundancy

We define a microbial community, C, with S species, where the ith species has a relative

abundance, p_i. (For simplicity, we refer to ecological units as species, although it would be

possible to perform this analysis at other taxonomic levels – strains, say, or genera – or even to

perform the analysis at the level of guilds in order to evaluate resource competition among

guilds, e.g., sulfate-reducers competing with methanogens for hydrogen electrons). We therefore

represent C as:

$$C = \{p_i > 0 \ \forall \ p_i; \textstyle\sum_{i=1}^{S} p_i = 1\}$$ (1

where p is the set of relative abundances of species, such that any given relative abundance for

the i^{th} species is greater than zero and all species relative abundances sum to one. Equation 1

represents a traditional abundance distribution (25).

Our goal was to calculate how evenly different species in a community, C (equation 1),

contribute to a community-aggregated parameter, or ecosystem function, T, at a fixed time and

space. To do this, we first determined the relative contribution, τ_i, that the i^{th} species makes

towards T such that:

$$\tau_i = \frac{p_i t_i}{\sum_{i=1}^{n} p_i t_i}$$ 2)

where t_i is the species-specific contribution level and p_i is the relative abundance the i^{th} species in

C. In equation 2, τ_i represents the relative contribution that the i^{th} species contributes to a

community-aggregated parameter. We note that the functional analysis and preprocessing of

metagenomes and transcriptomes varies in the literature and public databases. For example, the

TARA Oceans metagenomes have been analyzed as assembled, un-binned metagenomes, where

individual genes were assigned taxonomy, annotation, and abundances (26), and as metagenome-

assembled genomes (24, 27). Although the fundamental meaning of τ does not change across

these different datasets, the *in silico* approach to calculating τ may slightly differ depending on

how sequence data is preprocessed. For un-binned metagenomes, τ_i simply corresponds to the

proportion of reads associated with the i^{th} species which map to function, T. Similarly for

transcriptomes, τ_i simply corresponds to the proportion of transcripts expressed by the i^{th} species

which map to function, T. Assuming the community sequence spaces has been randomly

sampled, these scenarios implicitly account for the number of gene copies (t) and the relative

abundance (p) of the i^{th} species. In the case of metagenome-assembled genomes, relative

contribution level can be calculated with equation 2 such that t_i is gene copies and abundance is

measured as the relative proportion of reads mapping to single-copy marker genes for the i^{th}

species.

As an example, imagine that we have a metagenome that has been binned into MAGs.

Our hypothetical community has three species: a Cyanobacterium, Alphaproteobacterium, and a

Gammaproteobacterium. Mapping metagenome reads to single-copy marker genes reveals that

the Cyanobacteria, the Alphaproteobacterium and Gammaproteobacterium each represent 50%,

25%, and 25% of the community, respectively. We wish to calculate the functional redundancy

of this community with respect to the production of a secondary metabolite. We use the presence

of a gene, *xxxA*, as a proxy for an organisms' potential to produce this secondary metabolite. The

MAGs reveal that the Cyanobacterium and Alphaproteobacterium contain 5 copies of *xxxA* each,

while the Gammaproteobacterium lacks the *xxxA* gene. Thus, the relative contribution from each

species to secondary metabolite production, when using *xxxA* as a proxy, is $\tau=0.67$, $\tau=0.33$, and

$\tau=0$, respectively

Next, we reformulate the abundance distribution for community, C, as an abundance

distribution with respect to τ (relative contribution distribution), such that:

$$C_{\tau,all} = \left\{ \tau \,\middle|\, \tau_i \geq 0 \,\forall\, \tau_i; \sum_{i=1}^{n} \tau_i = 1 \right\}$$

142   (3.1

143   and

$$C_{\tau} = \left\{ \tau \in C_{\tau,all} \,\middle|\, \tau_i \geq 0 \,\forall\, \tau_i; \sum_{i=1}^{\sigma} \tau_i = 1 \right\}$$

144   (3.2

where again, τ_i is the relative contribution by the i^{th} species to T, such that all τ sum to equal 1.

The community represented by equation 3.1 includes species not contributing to the community-

aggregated parameter. The community represented by equation 3.2 is a subset of the one

represented by equation 3.1, where C_{τ} contains only species from $C_{\tau,all}$ contributing towards a

community-aggregated parameter, which we refer to as "contributors." As such, we adopted the

variable, σ, which represents the richness of contributors. The effective richness of contributors

towards T was calculated using equation 3.2 and the Tsallis entropy index (21)(21) such that:

$$D_q = \begin{cases} \left(\sum_{i=1}^{\sigma} \tau_i^q \right)^{\frac{1}{1-q}} & q \neq 1 \\ exp\left(\sum_{i=1}^{\sigma} -\tau_i \log \tau_i \right) & q = 1 \end{cases}$$

152   (4

where τ_i is the relative contribution by i^{th} species in C_{τ} towards T, and D_q is the effective richness

of contributors, and q is the diversity order, which weights the importance of species contributing

more or less towards T when calculating D_q. This is analogous to diversity order weighting the

importance of rare versus abundant species when measuring diversity of species abundances

(22). With diversity order $q=0$, equation 4 simplifies to the number of contributors in equation

3.2 (i.e., σ). Diversity order $q=0$ removes any influence in the variability of τ when calculating

D_q. Increasing the diversity order subsequently decreases the perceived contribution of

contributors with smaller τ. Diversity orders $q=1$ and $q=2$ correspond to Shannon's index. and

Simpson's index.

We then normalize Dq by community richness S, which is equal to Ds if every species

were to contribute to T equally), and adjust evenness to account for the functional dependence

that the lower evenness bound has on community richness (20)(20). As such, functional

redundancy spans the full range from 0 to 1, such that:

$$CE = \frac{D_q - 1}{S - 1}$$ (5

Here, functional redundancy, *CE*, indicates how evenly species contribute to a trait, *T*.

For the hypothetical community described above, the effective richness (D_q, $q=0.5$—see *Results*)

of species possessing *xxxA* was ~1.55, and the functional redundancy, with respect to *xxxA*, was

R=0.47 (*q=0.5*; see *Results*).

Simulations of Functional Redundancy as a Function of Trait Stability

In order to test the fidelity with which CE measures trait stability, or the stability of a

community-aggregated parameter to species extinction, we simulated artificial communities with

varying values of species richness (*S*), richness of contributors (σ), and contribution levels by

contributors (*t)*. First, the richness of contributors was randomly chosen from a uniformly

distributed range of 1 to 100. Next, the richness of species not contributing to the community-

aggregated parameter was randomly chosen from a uniformly distributed range of 0 to 100, so

that some artificial communities were dominated by contributors while others by non-

contributors. The relative abundance, *p*, of individual species was modeled log-normally (28,

29)(28, 29) such that:

$$R(r) = R_0 e^{-a^2 r^2}$$ (6

where R is the r^{th} rank, R_0 is the maximum abundance, and a is the inverse width of the

lognormal distribution. Values of a were randomly selected according to a uniform distribution

from 0 (even) to 0.4 (more uneven), at 0.01 intervals. After establishing species relative

abundances, we randomly chose σ contributing taxa from our lognormally distributed

community. The contribution level, t, was then randomly assigned to these species from a

uniformly distributed range of 0.01 to 1, at intervals of 0.01. We generated and calculated R for

1,000 artificial communities.

The stability of a community-aggregated parameter to extinction was tested by a

simulation in which stability to extinction was quantified as the average fraction of species

richness (S) that must go extinct in order to decrease community-aggregated parameter by 99%.

Species were randomly removed from the community until summed τ was reduced to 1% of its

original level. We used a 99% extinction, rather than 100%, to allow the evaluation of

communities where all species were contributors. In such communities, all species would have to

go extinct to remove 100% of the ecosystem function. We think that it is also justified

ecologically, because dormant microbes in the rare biosphere can serve as a deep reservoir of

traits (30), and it is not clear that fully removing a trait from an ecosystem happens in practice.

The extinction simulation was performed 100 times for each of the individual 1,000 artificial

communities. The stability to extinction generated from the 100 simulations were then averaged

for each of the individual 1,000 artificial communities. We made no *a priori* assumptions about

the optimal diversity order (i.e., q) to use in equations 4 and 5. We performed the simulation

described above varying diversity orders from 0 to 2 at 0.1 intervals. The best diversity order was

determined by fitting generalized additive model (GAM) regressions between R and stability to

extinction, and then choosing the diversity order which resulted in a GAM fit with the lowest

mean absolute error among residuals. All GAM fits were performed using the gam function from

the R package mgcv (31). Last, we performed the same simulations and calculated R using other

methods described in the literature. A summary of these measurements is provided in Table 1.

An R package which calculates functional redundancy (equations 1-5) and performs these

simulations is available at https://github.com/taylorroyalty/funfunfun.

A Case Study of Functional Redundancy in the Global Ocean Nitrogen Cycle

211   Draft metagenome-assembled genomes (MAGs) from Tully, Graham et al. (24) were

downloaded from NCBI (32). Data on metagenome sequencing effort (number of short reads),

MAG relative abundance estimates (pre-corrected for genome size), ocean depths, and ocean

regions, defined in (24), were downloaded from Figshare

(https://doi.org/10.6084/m9.figshare.5188273). MAG relative abundances from each site were

multiplied by the respective site's sequencing effort to convert relative abundances into

abundance count data. Abundance counts for all sites were normalized by the lowest abundance

(greater than 0) among all sites. MAG abundances were then rarefied using the rrarefy function

from the R package vegan (33). We chose a sampling effort of 14,265, as this sampling effort

corresponded to the site with lowest sampling effort (i.e., sequencing effort). We tabulated

nitrogen-cycle marker genes in individual MAGs by blasting MAG proteins against nitrogen-

cycle marker gene databases using the BLASTP algorithm (E-value=10^{-10}) (34). Nitrogen cycle

marker genes were categorized into eight categories: N_2 fixation (*nifH*), nitrification (*amoA*,

hao), NO_3 reduction (*napA*, *narG*), dissimilatory NO_3 reduction (*nrfA*), assimilatory NO_3/NO_2

reduction (*narB*, *nasA*, *nasB*, and *nirB*), NH_4^+ assimilation (*gdhA*, *glnA*), N remineralization

(*gltB*), and denitrification (*nirK*, *nirS*, *norB*, and *nosZ*) (23). Individual BLAST databases were

constructed for each nitrogen cycle category. Gene names associated with individual categories

were searched for in the Swiss-Prot database (35). All proteins matching gene names were

subsequently downloaded and used in BLAST database construction. Functional redundancy was

then calculated for individual stations, at individual depths, for each nitrogen cycle category

using equations 1-5. We used a diversity order, q=0.5, based on the GAM sensitivity analysis.

Table 2 contains protein names, gene names, corresponding EC, and entry name for all genes

used in the above analysis.

Results and Discussion

Representing microbial communities as species abundance, trait abundance, and relative

contribution distributions

CE represents functional redundancy using two distributions: the distribution of species

abundances within a community, and the distribution of contribution levels among species. Here,

we evaluate genomic traits because these are regularly adopted as proxies for microbially-

mediated biogeochemistry (16, 26, 36–38). Because trait levels are measured as non-negative,

real numbers (18), the relative contribution of individual species to a community-aggregated

parameter can be represented as a relative abundance distribution (Figure 1). We determined the

relative contribution of individual species to the community-aggregated parameter from the

product of species relative abundances and their respective trait abundance distributions

(equation 2). We refer to the distribution of the relative contribution of each species to total as

the relative contribution distribution (equation 3). Thus, an abundant species that expresses a

small amount of trait per cell, and a rarer species that expresses a large amount of trait per cell,

can contribute equally with respect to a community-aggregated parameter (Figure 1).

Defining, Quantifying Functional Redundancy, and Choosing a Diversity Order q

We measured functional redundancy as how evenly species in a community contribute to

a community-aggregated parameter and termed this metric as contribution evenness (CE). This

definition is consistent with descriptions of functional redundancy as multiple species sharing

similar ecological roles or functions (2, 3). The relative contribution by species towards a

community aggregated-parameter (equation 2) provides a framework to calculate the effective

number, or richness, of species that contribute to that community-aggregated parameter as an

ecological function. This effective richness is calculated with traditional diversity indices

(equations 3-4) (21, 22). To determine the effective proportion of a community contributing to

the community-aggregated parameter, the richness of contributors is normalized by species

richness (22). Last, to make meaningful comparisons of evenness between communities with

different levels of richness, we made our evenness metric unrelated to community richness

(equation 5)(20).

Numerous diversity indices have been proposed in the literature. Three common

measures of diversity include species richness, Shannon's diversity index, and Simpson's

diversity index, are manifestations of equation 4 when the q term—commonly referred to as the

diversity order—is $q=0$, $q=1$, and $q=2$, respectively. The diversity order changes the relative

weight placed on rare versus common species when determining the effective richness. We

found that GAM fits between functional redundancy and stability to extinction followed positive,

monotonic functions for all diversity orders spanning $q=0$ to $q=2$. However, at $q \approx 0.5$, CE was

most tightly related to trait stability, our benchmark for functional redundancy efficacy (Figure 2;

also compare Fig 3a to Fig 3b). Therefore, we suggest that CE is most effective as an indicator of

trait stability $q=0.5$ makes *CE* the best indicator of how resilient traits are to species extinction.

In that case, equation 4 simplifies to:

$D_q = \left(\sum_{i=1}^{\sigma} \sqrt{\tau_i} \right)^2$ (7

and CE becomes:

$CE = \dfrac{\left(\sum_{i=1}^{\sigma} \sqrt{\tau_i} \right)^2 - 1}{S - 1}$ (8

This is the formulation of CE that we recommend for general use.

Functional Redundancy and Trait Stability to Species Extinction

278   A tangible ecological consequence of functional redundancy is stability of an ecosystem

function to the extinction of species performing that function. When functions are more broadly

distributed across species, a function is less likely to disappear due to the extinction of any subset

of species. This interpretation is often highlighted in macro-ecological studies (2–4, 7, 39–41).

The conceptual relationship between functional redundancy and stability of an ecosystem

function provides an opportunity to demonstrate the fidelity of CE. To do this, CE was calculated

for randomly generated communities. Individual species were randomly assigned relative

contribution levels towards a community-aggregated parameter. We simulated random species

extinction events in the artificial communities until >99% of the original ecosystem function was

removed from the community (see *Methods*).

288   It is important to emphasize that these simulations were not designed to mimic how

extinction occurs in nature. Instead, these simulations empirically demonstrate the degree to

which an ecosystem function is concentrated in species. The more widely an ecosystem function

is distributed across species, the more species must go extinct before the ecosystem function is

lost. We observed a positive, monotonic relationship between the stability to random species

extinction events and CE (Figure 3A). When CE was ~0, stability to species extinction was ~0.5,

meaning ~50% of the community, on average, went extinct before an ecosystem function was

removed from the community. This is equivalent to only one species performing the ecosystem

function. When CE was ~1, stability to species extinct was ~1, meaning that 100% of the

community, on average, had to go extinct to completely remove a function from an ecosystem. In

general, the relationship between CE and trait stability to extinction resembled a saturation

curve—although we opted to use GAMs when calculating mean absolute error (previous section)

as we made no assumption about the true relationship between CE and function stability, other

than it should be a positive, monotonic function.

Here, we recommend the form of CE described in equation 8, as it is optimized for our

intended use case, which is measuring relative stability of traits to species extinctions within

communities. However, we can imagine other applications of the CE concept, for which other

values of q may be appropriate. Even for species diversity (i.e., equation 4), there is still

discussion over the merits of using Simpson's Index ($q=2$), Shannon's Index ($q=1$), and richness

($q=0$) in different situations (42–44). As such, it is important to be explicit about the implications

of behind choosing a trait and the diversity order.

CE Measures Trait Stability More Reliably than Other Functional Redundancy Metrics

Several methods for evaluating community-level functional redundancy exist in the

literature, as detailed in Table 1. While Table 1 certainly does not include all published metrics,

the highlighted metrics reflect the diversity in how traits, community properties (i.e., richness

and evenness), as well as the desired metric utility (by design) vary among different functional

redundancy metrics. Two of the metrics evaluate niche overlap, where niches are defined in

multivariate trait space (6, 7). In the case of Ricotta et al. (7), functional redundancy weights

species abundance, leading to a measure of the expected niche dissimilarity from the mean

community niche. In contrast, Hester et al. (6) evaluates the median metabolic overlap which

considers neither species abundance nor trait levels. One metric evaluates niche occupancy,

where niches are defined in multivariate trait space (5). In this approach, functional redundancy

is taken to be as the ratio of species richness to plausible number niches (unique combinations of

trait states), called functional entities. The last metric is informal but has been utilized by various

authors. In this approach, functional redundancy is simply interpreted as the proportion of

species in a community with a trait (e.g. Wohl et al., 2004; Tully et al., 2017; Trivedi et al.,

2020).

To highlight the difference between our metric and the others described in Table 1, we

repeated the extinction simulations and compared all estimates of functional redundancy with

trait stability to species extinction (Figure 3B-E). In these simulations, only one trait was

considered when calculating functional redundancy with all metrics. For the method presented in

Mouillot et al. (5), we assumed there were two functional entities (trait absence and presence).

Generally, all functional redundancy metrics trended differently to stability to species extinction.

This observation was not surprising, as most of these methods evaluate niche overlap in some

fashion. Among the metrics in Table 1, the fraction of the community possessing a trait behaved

most like CE (Figure 3B). We note that the fraction of the community possessing a trait is a

special case of CE, with diversity order $q=0$. Lastly, one inherent limitation of CE is that it only

evaluates single community-aggregated parameters at a time. To address functional redundancy

with respect to multiple traits perform dimensional reduction (e.g., PCA) on multiple traits and

use the resultant factors as trait levels in equation 2. Factors in this case would be interpreted as

different niches.

Case Study: Redundancy of Nitrogen Cycle Marker Genes in Marine Microbiomes

To demonstrate a real application and provide a positive control for CE results, we

measured CE in the marine nitrogen cycle using genomic data from the *TARA Oceans* project

(47)(45). Recently, (24) published 2,361 draft metagenome-assembled genomes (MAGs) which

represent genomic content from seven oceanographic provinces, three different depths, and four

filter size fractions. Relative abundance data used here were reported in (24) and were

determined by the proportion of sequence reads mapping to assembled MAGs. For simplicity,

we limited our analysis to only MAGs identified in the bacterial size fraction (<3 μm), which

corresponds to free-living microbes. The nitrogen cycle was simplified to eight pathways

(23)(23) and included N_2 fixation, nitrification, NO_3^- reduction, dissimilatory NO_3^- reduction,

assimilatory NO_3^-/NO_2^- reduction, NH_4^+ assimilation, N remineralization, and denitrification. A

general summary of the marker genes (downloaded from Swiss-Prot; see Methods) used for

quantifying nitrogen cycle pathways in MAGs is summarized in Table 2.

Overall, the functional redundancy for individual genomic traits was consistent with our

conceptual understanding of primary nitrogen-transforming pathways in the ocean (Figure 4). A

clear example was the low functional redundancy in the nitrification pathway. We expected this

observation as nitrification metabolisms in oceans are primarily restricted to *Crenarchaeota* (46).

Similarly, the N_2 fixation pathway, which is predominantly facilitated by a small group of

Cyanobacteria, exhibits low functional redundancy. Conversely, the uptake of inorganic N via

NH_4^+ and NO_3^-/NO_2^- assimilation are known to be broadly expressed traits , and accordingly

exhibit high CE. Last, the low functional redundancy in dissimilatory NO_3^- reduction and NO_3^-

reduction were not surprising considering these pathways occur in anaerobic and low-oxygen

environments, respectively (23).

Some variance existed across oceanographic provinces and depth for intra-pathway

functional redundancy. However, inter-pathway variance in functional redundancy was generally

larger than intra-pathway variance (Figure 4). This suggests that the marine nitrogen cycle

marker genes considered here are consistently distributed among communities for MAGs

retrieved in Earth's ocean at middle and lower latitudes. This was consistent with the lack of

relationship between the metagenome companion metadata (e.g., O_2, inorganic N, PO_4^{3-}, latitude,

longitude, etc.) and functional redundancy. The low intra-pathway variance in functional

redundancy was not particularly surprising, as saltwater biomes exhibit the lowest alpha-

diversity and small beta-diversity among free-living microbiomes (47, 49)(45)(47). Minor shifts

in community composition among lower taxonomic ranks (e.g., genus or species) will not

drastically alter redundancy in genomic traits as changes in genomic composition is buffered by

traits conserved at higher taxonomic ranks (e.g., phylum level) (50)(48). This observation was in

agreement with an earlier analysis evaluating cluster of orthologous groups (COGs) composition

across *TARA Oceans* metagenomes (45).

The disproportionate number of lineages that lack cultured representatives highlights the

continued importance of nucleotide sequencing in microbial ecology and the interference of

microbial function (49). Here, we demonstrated that CE provides a framework to evaluate the

functional redundancy for these traits in a novel way. This approach represents an easy-to-apply,

quantitative metric for functional redundancy that faithfully represents the stability of traits to

extinctions of species within a community.

Data Availability

In order to facilitate the use of CE functional redundancy measurements, we created an R

package to calculate CE functional redundancy and run extinction simulations. The package is

available at https://github.com/taylorroyalty/funfunfun.

Acknowledgments

We gratefully acknowledge the comments of three anonymous reviewers, whose

constructive comments substantially improved the manuscript. Funding for this project was

provided by a C-DEBI graduate fellowship to TMR and by the U.S. Department of Energy,

Office of Science, Office of Biological and Environmental Research, Genomic Science Program

under Award Number DE-SC0020369 for TMR and ADS. This is C-DEBI contribution number

[to be assigned upon manuscript acceptance].

Author Contribution

TMR designed the research, performed research, analyzed data, developed the R

package, and wrote the paper. ADS designed the research, developed the R package, and wrote

the paper.

References

1. Louca S, Polz MF, Mazel F, Albright MBN, Huber JA, O'Connor MI, Ackermann M,
Hahn AS, Srivastava DS, Crowe SA, Doebeli M, Parfrey LW. 2018. Function and
functional redundancy in microbial systems. Nat Ecol Evol 2:936–943.
2. Naeem S. 1998. Species Redundancy and Ecosystem Reliability 12:39–45.
3. Rosenfeld JS, Mall M. 2002. Functional redundancy in ecology and conservation.
OIKOS 1.
4. Fonseca CR, Ganade G. 2001. Species Functional Redundancy, Random Extinctions and
the Stability of Ecosystems. J Ecol 89:118–125.
5. Cottingham KL, Brown BL, Lennon JT. 2001. Biodiversity may regulate the temporal
variability of ecological systems. Ecol Lett 4:72–85.
6. Mouillot D, Villéger S, Parravicini V, Kulbicki M, Arias-González JE, Bender M,
Chabanet P, Floeter SR, Friedlander A, Vigliola L, Bellwood DR. 2014. Functional over-
redundancy and high functional vulnerability in global fish faunas on tropical reefs. Proc
Natl Acad Sci U S A 111:13757–13762.
7. Hester ER, Jetten MSM, Welte CU, Lücker S. 2019. Metabolic overlap in
environmentally diverse microbial communities. Front Genet 10:1–10.
8. Ricotta C, de Bello F, Moretti M, Caccianiga M, Cerabolini BEL, Pavoine S. 2016.
Measuring the functional redundancy of biological communities: a quantitative guide.
Methods Ecol Evol 7:1386–1395.
9. Galand PE, Pereira O, Hochart C, Auguet JC, Debroas D. 2018. A strong link between
marine microbial community composition and function challenges the idea of functional
redundancy. ISME J 12:2470–2478.
10. Micheli F, Halpern BS. 2005. Low functional redundancy in coastal marine assemblages.
Ecol Lett 8:391–400.
11. Miki T, Yokokawa T, Matsui K. 2013. Biodiversity and multifunctionality in a microbial
community: A novel theoretical approach to quantify functional redundancy. Proc R Soc
B Biol Sci 281.
12. Tully BJ, Wheat CG, Glazer BT, Huber JA. 2018. A dynamic microbial community with
high functional redundancy inhabits the cold, oxic subseafloor aquifer. ISME J 12:1–16.

13.   Wohl DL, Arora S, Gladstone JR. 2004. Functional Redundancy Supports Biodiversity
and Ecosystem Function in a Closed and Constant Environment. Ecology 85:1534–1540.

14.   Trivedi CB, Stamps BW, Lau GE, Grasby SE, Templeton AS, Spear JR. 2020. Microbial
Metabolic Redundancy Is a Key Mechanism in a Sulfur-Rich Glacial Ecosystem.
mSystems 5:1–16.

15.   Chase JM, Leibold MA. 2003. Ecological Niches, 1st ed. The University of Chicago
Press, Chicago.

16.   Green JL, Bohannan BJM, Whitaker RJ. 2008. Microbial biogeography: From taxonomy
to traits. Science (80-) 320:1039–1043.

17.   Violle C, Navas M, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E, Oikos S, May
N, Violle C, Navas M, Vile D, Kazakou E, Fortunel C. 2007. Let the concept of trait be
functional! Oikos 116:882–892.

18.   Krause S, Le Roux X, Niklaus PA, van Bodegom PM, Lennon T. JT, Bertilsson S,
Grossart HP, Philippot L, Bodelier PLE. 2014. Trait-based approaches for understanding
microbial biodiversity and ecosystem functioning. Front Microbiol 5:1–10.

19.   Fierer N, Barberán A, Laughlin DC. 2014. Seeing the forest for the genes: Using
metagenomics to infer the aggregated traits of microbial communities. Front Microbiol
5:1–6.

20.   Chao A, Ricotta C. 2019. Quantifying evenness and linking it to diversity , beta diversity
, and similarity 100:1–15.

21.   Jost L. 2006. Entropy and diversity. Oikos 2.

22.   Hill MO. 1973. Diversity and Evenness : A Unifying Notation and Its Consequences.
Ecology 54:427–432.

23.   Pajares S, Ramos R. 2019. Processes and Microorganisms Involved in the Marine
Nitrogen Cycle: Knowledge and Gaps. Front Mar Sci 6.

24.   Tully BJ, Graham ED, Heidelberg JF. 2018. The reconstruction of 2,631 draft
metagenome-assembled genomes from the global oceans. Sci Data 5:1–8.

25.   McGill BJ, Etienne RS, Gray JS, Alonso D, Anderson MJ, Benecha HK, Dornelas M,
Enquist BJ, Green JL, He F, Hurlbert AH, Magurran AE, Marquet PA, Maurer BA,
Ostling A, Soykan CU, Ugland KI, White EP. 2007. Species abundance distributions:

Moving beyond single prediction theories to integration within an ecological framework.

Ecol Lett 10:995–1015.

26.   Salazar G, Paoli L, Alberti A, Huerta-Cepas J, Ruscheweyh HJ, Cuenca M, Field CM,

Coelho LP, Cruaud C, Engelen S, Gregory AC, Labadie K, Marec C, Pelletier E, Royo-

Llonch M, Roux S, Sánchez P, Uehara H, Zayed AA, Zeller G, Carmichael M, Dimier C,

Ferland J, Kandels S, Picheral M, Pisarev S, Poulain J, Acinas SG, Babin M, Bork P,

Boss E, Bowler C, Cochrane G, de Vargas C, Follows M, Gorsky G, Grimsley N, Guidi

L, Hingamp P, Iudicone D, Jaillon O, Kandels-Lewis S, Karp-Boss L, Karsenti E, Not F,

Ogata H, Pesant S, Poulton N, Raes J, Sardet C, Speich S, Stemmann L, Sullivan MB,

Sunagawa S, Wincker P. 2019. Gene Expression Changes and Community Turnover

Differentially Shape the Global Ocean Metatranscriptome. Cell 179:1068-1083.e21.

27.   Delmont TO, Quince C, Shaiber A, Esen ÖC, Lee ST, Rappé MS, MacLellan SL, Lücker

S, Eren AM. 2018. Nitrogen-fixing populations of Planctomycetes and Proteobacteria are

abundant in surface ocean metagenomes. Nat Microbiol 3:804–813.

28.   Magurran AE. 1988. Ecological Diversity and Its Measurement, 1st ed. Croom Helm Ltd.

29.   Galand PE, Casamayor EO, Kirchman DL, Lovejoy C. 2009. Ecology of the rare

microbial biosphere of the Arctic Ocean. Proc Natl Acad Sci 106:22427–22432.

30.   Lennon JT, Jones SE. 2011. Microbial seed banks: The ecological and evolutionary

implications of dormancy. Nat Rev Microbiol 9:119–130.

31.   Wood S. 2017. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML

Smoothness Estimation.

32.   Federhen S. 2012. The NCBI Taxonomy database. Nucleic Acids Res 40:136–143.

33.   Philip D. 2003. Computer program review VEGAN , a package of R functions for

community ecology. J Veg Sci 14:927–930.

34.   Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2019. GenBank. Nucleic

Acids Res 47:D94–D99.

35.   Bairoch A, Apweiler R. 2000. The SWISS-PROT protein sequence database and its

supplement TrEMBL in 2000. Nucleic Acids Res 28:45–48.

36.   Falkowski PG, Fenchel T, Delong EF. 2008. The Microbial Engines That Drive Earth's

Biogeochemical Cycles. Science (80-) 320:1034–1039.

37. Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD. 2019.

Defining trait-based microbial strategies with consequences for soil carbon cycling under

climate change. ISME J 445866.

38. Baker BJ, Lazar CS, Teske AP, Dick GJ. 2015. Genomic resolution of linkages in carbon,

nitrogen, and sulfur cycling among widespread estuary sediment bacteria. Microbiome

3:14.

39. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, Thomas BC,

Singh A, Wilkins MJ, Karaoz U, Brodie EL, Williams KH, Hubbard SS, Banfield JF.

2016. Thousands of microbial genomes shed light on interconnected biogeochemical

processes in an aquifer system. Nat Commun 7:13219.

40. Teeling H, Fuchs BM, Becher D, Klockow C, Gardebrecht A, Bennke CM, Kassabgy M,

Huang S, Mann AJ, Waldmann J, Weber M, Klindworth A, Otto A, Lange J, Bernhardt J,

Reinsch C, Hecker M, Peplies J, Bockelmann FD, Callies U, Gerdts G, Wichels A,

Wiltshire KH, Glöckner FO, Schweder T, Amann R. 2012. Substrate-Controlled

Succession of Marine Bacterioplankton Populations Induced by a Phytoplankton Bloom.

Science (80-) 336:608–611.

41. McPherson JM, Yeager LA, Baum JK. 2018. A simulation tool to scrutinise the

behaviour of functional diversity metrics. Methods Ecol Evol 9:200–206.

42. Tilman D. 2001. Functional Diversity. Encycl Biodivers Second Ed 3:587–596.

43. Tilman D, Reich PB, Knops JMH. 2006. Biodiversity and ecosystem stability in a

decade-long grassland experiment. Nature 441:629–632.

44. Patil GP, Taillie C. 1982. Diversity as a concept and its measurement. J Am Stat Assoc

77:548–561.

45. Jost L. 2007. Partitioning Diversity into Independent Alpha and Beta Components.

Ecology 88:2427–2439.

46. Morris EK, Caruso T, Buscot F, Fischer M, Hancock C, Maier TS, Meiners T, Müller C,

Obermaier E, Prati D, Socher SA, Sonnemann I, Wäschke N, Wubet T, Wurst S, Rillig

MC. 2014. Choosing and using diversity indices: insights for ecological applications

from the German Biodiversity Exploratories. Ecol Evol 4:3514–24.

47.   Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, Djahanschiri B,
Zeller G, Mende DR, Alberti A, Cornejo-Castillo FM, Costea PI, Cruaud C, D'Ovidio F,
Engelen S, Ferrera I, Gasol JM, Guidi L, Hildebrand F, Kokoszka F, Lepoivre C, Lima-
Mendez G, Poulain J, Poulos BT, Royo-Llonch M, Sarmento H, Vieira-Silva S, Dimier
C, Picheral M, Searson S, Kandels-Lewis S, Boss E, Follows M, Karp-Boss L, Krzic U,
Reynaud EG, Sardet C, Sieracki M, Velayoudon D, Bowler C, De Vargas C, Gorsky G,
Grimsley N, Hingamp P, Iudicone D, Jaillon O, Not F, Ogata H, Pesant S, Speich S,
Stemmann L, Sullivan MB, Weissenbach J, Wincker P, Karsenti E, Raes J, Acinas SG,
Bork P. 2015. Structure and function of the global ocean microbiome. Science (80-)
348:1–10.
48.   Wuchter C, Abbas B, Coolen MJL, Herfort L, Van Bleijswijk J, Timmers P, Strous M,
Teira E, Herndl GJ, Middelburg JJ, Schouten S, Damsté JSS. 2006. Archaeal nitrification
in the ocean. Proc Natl Acad Sci U S A 103:12317–12322.
49.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi
A, Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-
Baeza Y, González A, Morton JT, Mirarab S, Xu ZZ, Jiang L, Haroon MF, Kanbar J, Zhu
Q, Song SJ, Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM,
Hampton-Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA, Clauset A,
Stevens RL, Shade A, Pollard KS, Goodwin KD, Jansson JK, Gilbert JA, Knight R,
Agosto Rivera JL, Al-Moosawi L, Alverdy J, Amato KR, Andras J, Angenent LT,
Antonopoulos DA, Apprill A, Armitage D, Ballantine K, Bárta J, Baum JK, Berry A,
Bhatnagar A, Bhatnagar M, Biddle JF, Bittner L, Boldgiv B, Bottos E, Boyer DM, Braun
J, Brazelton W, Brearley FQ, Campbell AH, Caporaso JG, Cardona C, Carroll JL, Cary
SC, Casper BB, Charles TC, Chu H, Claar DC, Clark RG, Clayton JB, Clemente JC,
Cochran A, Coleman ML, Collins G, Colwell RR, Contreras M, Crary BB, Creer S,
Cristol DA, Crump BC, Cui D, Daly SE, Davalos L, Dawson RD, Defazio J, Delsuc F,
Dionisi HM, Dominguez-Bello MG, Dowell R, Dubinsky EA, Dunn PO, Ercolini D,
Espinoza RE, Ezenwa V, Fenner N, Findlay HS, Fleming ID, Fogliano V, Forsman A,
Freeman C, Friedman ES, Galindo G, Garcia L, Garcia-Amado MA, Garshelis D, Gasser
RB, Gerdts G, Gibson MK, Gifford I, Gill RT, Giray T, Gittel A, Golyshin P, Gong D,

Grossart HP, Guyton K, Haig SJ, Hale V, Hall RS, Hallam SJ, Handley KM, Hasan NA,

Haydon SR, Hickman JE, Hidalgo G, Hofmockel KS, Hooker J, Hulth S, Hultman J,

Hyde E, Ibáñez-Álamo JD, Jastrow JD, Jex AR, Johnson LS, Johnston ER, Joseph S,

Jurburg SD, Jurelevicius D, Karlsson A, Karlsson R, Kauppinen S, Kellogg CTE,

Kennedy SJ, Kerkhof LJ, King GM, Kling GW, Koehler A V., Krezalek M, Kueneman J,

Lamendella R, Landon EM, Lanede Graaf K, LaRoche J, Larsen P, Laverock B, Lax S,

Lentino M, Levin II, Liancourt P, Liang W, Linz AM, Lipson DA, Liu Y, Lladser ME,

Lozada M, Spirito CM, MacCormack WP, MacRae-Crerar A, Magris M, Martín-Platero

AM, Martín-Vivaldi M, Martínez LM, Martínez-Bueno M, Marzinelli EM, Mason OU,

Mayer GD, McDevitt-Irwin JM, McDonald JE, McGuire KL, McMahon KD, McMinds

R, Medina M, Mendelson JR, Metcalf JL, Meyer F, Michelangeli F, Miller K, Mills DA,

Minich J, Mocali S, Moitinho-Silva L, Moore A, Morgan-Kiss RM, Munroe P, Myrold

D, Neufeld JD, Ni Y, Nicol GW, Nielsen S, Nissimov JI, Niu K, Nolan MJ, Noyce K,

O'Brien SL, Okamoto N, Orlando L, Castellano YO, Osuolale O, Oswald W, Parnell J,

Peralta-Sánchez JM, Petraitis P, Pfister C, Pilon-Smits E, Piombino P, Pointing SB,

Pollock FJ, Potter C, Prithiviraj B, Quince C, Rani A, Ranjan R, Rao S, Rees AP,

Richardson M, Riebesell U, Robinson C, Rockne KJ, Rodriguezl SM, Rohwer F,

Roundstone W, Safran RJ, Sangwan N, Sanz V, Schrenk M, Schrenzel MD, Scott NM,

Seger RL, Seguinorlando A, Seldin L, Seyler LM, Shakhsheer B, Sheets GM, Shen C,

Shi Y, Shin H, Shogan BD, Shutler D, Siegel J, Simmons S, Sjöling S, Smith DP, Soler

JJ, Sperling M, Steinberg PD, Stephens B, Stevens MA, Taghavi S, Tai V, Tait K, Tan

CL, Taş N, Taylor DL, Thomas T, Timling I, Turner BL, Urich T, Ursell LK, Van Der

Lelie D, Van Treuren W, Van Zwieten L, Vargas-Robles D, Thurber RV, Vitaglione P,

Walker DA, Walters WA, Wang S, Wang T, Weaver T, Webster NS, Wehrle B,

Weisenhorn P, Weiss S, Werner JJ, West K, Whitehead A, Whitehead SR, Whittingham

LA, Willerslev E, Williams AE, Wood SA, Woodhams DC, Yang Y, Zaneveld J,

Zarraonaindia I, Zhang Q, Zhao H. 2017. A communal catalogue reveals Earth's

multiscale microbial diversity. Nature 551:457–463.

50.   Royalty TM, Steen AD. 2019. Quantitatively Partitioning Microbial Genomic Traits

among Taxonomic Ranks across the Microbial Tree of Life. mSphere 4:e00446-19.

51.   Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel

Uncultured Microbial Cells Dominate Earth Microbiomes. mSystems 3:1–12.

Tables

Table 1. A summary of metrics of community-level functional redundancy.

Authors	Interpretation	Equation	Notes
This work	Effective proportion of species with a trait	$$R = \frac{D_q - 1}{S - 1}$$	• Accounts for species abundances • Continuous trait states • Individual traits • R spans 0 to 1
(8)	Expected trait dissimilarity among species in a community	$$R = 1 - \frac{\sum_i p_i \sum_j p_j \delta_{ji}}{D_2}$$ δ_{ji} = trait distance D_2 = taxonomic diversity order (q=2)	• Accounts for species abundances • Continuous trait states • Individual or multivariate traits • Normalized by a maximum theoretical dissimilarity • R spans 0 to 1
(6)	Ratio of species to number of functional entities (states)	$$R = \frac{S}{FE};$$ FE = # of possible combinations of trait states for all traits under consideration	• Binary trait states • Individual or multivariate traits • R spans ~0 ($\frac{S}{FE} = 0$) to S
Various (see text for examples)	Fraction of species with a trait	$$R = \frac{v}{S}$$	• Binary trait states • Individual traits • R spans 0 to 1
(7)	Median overlap of traits between species	$$R = med(O);$$ $$O = \{o_{i,j} = |g_i \cap g_j|; i \neq j\};$$ $$g = taxa;$$ O = set containing # of overlaps among species	• Binary trait states • $|g_i \cap g_j|$ represents the number of overlapping traits g between species i and species j • Individual and multivariate traits • R spans 0 to ∞

585   Table 2. A summary of marker gene sequences used in the BLASTP analysis of *TARA Oceans*

586   MAGs. All marker genes were downloaded from Swiss-Prot.

Pathway	Gene Names	EC Number	# Marker Genes
N Remineralization	gltB	1.4.1.13	14
N Fixation	nifH	1.18.6.1	127
Nitrification	amoA	1.14.99.39	2
	hoa	1.7.2.6	1
NO_3^- Reduction	napA	1.9.6.1	136
	narG	1.7.5.1	13
Dissimilatory NO_3^- Reduction	nrfA	1.7.2.2	77
Assimilatory NO_3^-/NO_2^- Reduction	narB	1.7.5.1	3
	nasA	1.7.7.2	1
	nasB	1.7.1.4	3
	nirB	1.7.1.15	5
NH_4^+ Assimilation	gdhA	1.4.1.3	14
	glnA	6.3.1.2	173
Denitrification	nirK	1.7.2.1	7
	nirS	1.7.2.1; 1.7.99.1	4
	norB	1.7.2.5	2
	nosZ	1.7.2.4	11

Figures

Figure 1: Community taxon abundance and relative contribution level are scalable and provide a

measure for the contribution that different species have towards a community-aggregated

parameter. The evenness of species-level contribution to the community-aggregated parameter is

taken as the functional redundancy.

Figure 2: The sensitivity of mean absolute error between functional redundancy and trait stability

to species extinction in response changing diversity order q in equation 4.

Figure 3: A comparison of the relationship between selected different functional redundancy

metrics and trait stability (see methods). Functional redundancy calculated with equation 5

(q=0.5) (**A**), equation 5 (q=0) (**B**), methodology from Ricotta et al. (2016) (**C**), methodology

from Mouillot et al. (2014) (**D**), and methodology from Hester et al. (2019). Black lines in panels

A-D correspond to best fit GAMs. The stars in panel **E** correspond to the average trait stability at

R=0 and R=1. The color scale corresponds to the fraction of a community with a trait versus.

Nitrogen-Transforming Pathway

Figure 4: Boxplots showing functional redundancy calculated for eight nitrogen-transforming

pathways in *TARA Oceans* draft MAGs. Different colors correspond to sample depth. Whiskers

correspond to 150% of the inter-quantile region.