Contribution Evenness: A functional redundancy metric for microbially-mediated

biogeochemical rates and processes

Taylor M. Royalty (1,2) and Andrew D. Steen (2,1)*

University of Tennessee Departments of (1) Earth and Planetary Sciences and (2) Microbiology

Running title:

Measuring the functional redundancy of microbiomes

*Corresponding: asteen1@utk.edu

Abstract

The concept of functional redundancy has received considerable attention in both the macroecology and microbial ecology literature. As a result, multiple metrics of functional redundancy have been proposed. These vary in how they weight trait levels, species abundance, functional richness, and species richness. Trait-based models are also increasingly popular ways to represent the function of microbial ecosystems. Here we present a new functional redundancy measurement tailored to be useful for community-aggregated parameters, which are summed ecosystem processes contributed by species, such as rates of chemical transformations. We focus in particular on community-aggregated parameters mediated by microbial communities. We call this approach contribution evenness (CE). CE measures how evenly species contribute to a community-aggregated parameter based on species contribution levels and abundances. As CE is an evenness measurement, quantitative representation of functional redundancy spans between 0 and 1, where 0 corresponds to a single species contributing and 1 corresponds to all species contributing equally to a community-aggregated parameter. CE is sensitive to the resilience of an ecosystem function to species extinction—an ecological consequence of functional redundancy. To illustrate how CE can be used, we analyzed the functional redundancy of eight nitrogen-transforming pathways using 2,631 metagenome-assembled genomes from 47 marine sites. We found that, on average, NH_4^+ assimilation pathway was the most functionally redundant (0.44 ± 0.08) while dissimilatory nitrate reduction had the lowest functional redundancy (0.005 ± 0.005). As demonstrated here, CE is a metric that is sensitive to trait resilience and allows for comparative analysis of functional redundancy for different traits.

Introduction

Functional redundancy was originally proposed as a community property that reflected the resilience of an ecosystem function (Naeem, 1998). Species performing a similar function could compensate for one another in the event of local species extinction, and thus, those functions are more resilient to taxa extinction. The purpose of measuring functional redundancy was to focus conservation efforts on vulnerable ecological groups that provide essential and unique ecological functions—i.e., keystone species (Walker, 1992). Since its proposal, interest in measuring functional redundancy has extended beyond conservation biology. In both the macroecology and microbial ecology literature, functional redundancy has measured in studies evaluating community succession, intra-community resource competition, biogeochemical fluxes, and metabolism biodiversity (Louca et al., 2018). Since measurements of functional redundancy have been used for diverse purposes in the ecological literature, it is helpful to identify metrics for functional redundancy that are tailored to purpose.

Several metrics for functional redundancy exist in the literature. Functional redundancy has been taken as the ratio of community richness and the theoretical maximum number of functional entities (Mouillot et al., 2014), median metabolic overlap of genomic traits (Hester, Jetten, Welte, & Lücker, 2019), expected variability in trait space among community members (Ricotta et al., 2016), the degree of covariance in functional diversity and biodiversity (e.g., Galand et al., 2018; Micheli & Halpern, 2005; Miki, Yokokawa, & Matsui, 2013), and the simple fraction of a community with a given trait (Trivedi et al., 2020; Tully, Wheat, Glazer, & Huber, 2018; Wohl, Arora, & Gladstone, 2004). Notably, these different metrics treat community structure differently. For instance, some metrics are sensitive to species abundances (Galand et al., 2018; Ricotta et al., 2016), richness (Micheli & Halpern, 2005; Mouillot et al., 2014), or

neither (Hester et al., 2019). The diversity of metrics reflects the diverse sources of data used to

calculate functional redundancy and the diverse research purposes for how this concept is

applied in research.

With exception of measuring a fraction of a community with a trait, all the functional

redundancy metrics listed above represent ecological function in terms of niche, where a niche is

thought of as a point in a hyperdimensional trait space. Communities with higher species density

in trait space are interpreted as being more functionally redundant. For the metrics mentioned

above, high species richness relative to unique combinations of traits, high overlap of metabolic

pathways, low variance of community members' position in trait space, and resilience of

community function to disturbance of community structure all suggest functionally redundant

communities.

An ecological function we are interested in understanding is the relative contribution by

species towards community-aggregated parameters. Community-aggregated parameters are

ecosystem functions (e.g. primary productivity, carbon turnover, nitrogen fixation, etc.) observed

on the community level, where the contribution by individual species is weighted by species

abundance and summed together by species-level contributions (Violle et al., 2007). Thus, we

argue one way to understand a species' ecological function is its contribution towards a

community-aggregated parameter, or the community-level ecosystem function. Describing

ecological function in terms of community-aggregated parameters would be particularly valuable

to metagenomic studies of microbial communities (Fierer, Barberán, & Laughlin, 2014; Krause

et al., 2014).

However, it is challenging to assign functions to microbes. Most microbial cells on Earth

belong to genera which lack cultured representatives, so the phenotypes for these organisms are

often unknown (Lloyd, Steen, Ladau, Yin, & Crosby, 2018; Staley & Konopka, 1985). Function

can sometimes be inferred experimentally (e.g., Orsi et al., 2018), but in most cases, information

about microbial traits comes exclusively from sequencing studies, where genomic traits are

proxies for microbial metabolisms. For microbial communities, we are often most concerned

with traits that relate to environmental processes, such as elemental transformations or secondary

metabolite production (Louca et al., 2018). Thus, it may be preferential to evaluate a small subset

of specific traits versus the full hyperdimensional trait space (thousands of genes, and a difficult-

to-describe number of metabolic processes, per genome) of a community.

As functional redundancy is a measure of species density with respect to an ecological

function, enumerating the number of "contributors" towards a community-aggregated parameter

may suffice as a measure of functional redundancy. Although not explicitly stated as such, some

methods we mentioned before use this approach (Trivedi et al., 2020; Tully, Wheat, et al., 2018;

Wohl et al., 2004). We note that simple enumeration does not weight the variable contributions

by different species, which may be desirable. Here, we propose a functional redundancy

measurement that accounts for variable contributions by different species towards community-

aggregated parameters. With this approach, the "true diversity" of contributors to a community-

aggregated parameter is calculated with traditional diversity metrics, where entropy is converted

into an effective richness of contributors (Hill, 1973; Jost, 2006). The observed diversity is

normalized to a theoretical maximum diversity and made "unrelated" to community richness

(Chao & Ricotta, 2019). Importantly, the normalization steps allow for meaningful comparisons

between communities with different species richness. Our measure of functional redundancy,

which we term contribution evenness (CE) for conciseness, is a measure of how evenly species

contribute to a community-aggregated parameter.

To demonstrate the efficacy of CE, we performed two *in silico* experiments with artificial

communities in parallel. Both experiments evaluated the extent that CE recapitulated the

resilience of a community-aggregated parameter to extinction of species. The first experiment

sought to minimize variance between CE and trait resilience to species extinction by changing

the diversity order (Hill, 1973; Jost, 2006) used to calculate CE. The next experiment compared

CE with other functional redundancy metrics as a demonstration that CE is fundamentally

different from previously proposed metrics. Last, we analyzed functional redundancy for eight

nitrogen-transforming pathways found in metagenomic data (Pajares & Ramos, 2019) in *TARA*

Oceans draft metagenome-assembled genomes (MAGs) (Tully, Graham, & Heidelberg, 2018).

We found CE appealing because it related to trait resilience to species extinction and was easily

generalizable—the functional redundancy among different genomic traits were comparable to

one another while not making assumptions about multi-trait guilds. We hope that the formulation

of functional redundancy fills a void in the genomic ecology literature by providing a generalized

tool for generating higher quality and comparable measurements of functional redundancy.

Methods and Theory

Definition of Functional Redundancy

Here, we define a microbial community, C, with S species, where the ith species has a

relative abundance, p_i. (For simplicity, we refer to ecological units as species, although it would

be possible to perform this analysis at other taxonomic levels – strains, say, or genera – or even

to perform the analysis at the level of guilds in order to evaluate resource competition among

guilds, e.g., sulfate-reducers competing with methanogens for hydrogen electrons). We therefore

represent C as:

$C = \{p | p_i > 0 \ \forall \ p_i; \sum_{i=1}^{S} p_i = 1\}$ (1

where p is the set of relative abundances of species, such that any given relative abundance for

the ith species is greater than zero and all species relative abundances sum to one. Equation 1

represents a traditional abundance distribution (McGill et al., 2007).

Our goal was to calculate how evenly different species in a community, C (equation 1),

contribute to a community-aggregated parameter, or net ecosystem function, T, at a fixed time

and space. To do this, we first determined the relative contribution that the ith species makes

towards T, defined here as τ_i, such that:

$\tau_i = \frac{p_i t_i}{\sum_{i=1}^{n} p_i t_i}$ 2)

where t_i is the species-specific contribution level and p_i is the relative abundance the ith species in

C. In equation 2, τ_i represents the relative contribution that the ith species, or population,

contributes to a community-aggregated parameter. Units of t are flexible such that t can be

expressed in terms of rates (e.g., g C m^{-2} year^{-1} for net primary productivity), quantities (e.g., net

gene copies in a metagenome), or dimensionless numbers (e.g., a relative proportion of rates or

quantities); however, T must be quantitative.

As an example, imagine a hypothetical community with three species, a Cyanobacterium

(p=0.5), an Alphaproteobacterium (p=0.25), and a Gammaproteobacterium (p=0.25). We wish to

calculate the functional redundancy of this community with respect to production of some

secondary metabolite. We use the presence of a gene, $xxxA$, as a proxy for an organisms'

potential to produce this secondary metabolite. The average number of $xxxA$ gene copies in the

genomes of a Cyanobacterium, Alphaproteobacterium, and Gammaproteobacterium, are 5, 5,

and 0, respectively. Thus, the relative contribution from each species to secondary metabolite

production, when using *xxxA* as a proxy, is $\tau=0.67$, $\tau=0.33$, and $\tau=0$, respectively

146   Next, we reformulate the abundance distribution for community, *C*, as an abundance

distribution with respect to τ (relative contribution distribution), such that:

$$C_{\tau,all} = \{\tau | \tau_i \geq 0 \ \forall \ \tau_i; \sum_{i=1}^{n} \tau_j = 1\}$$ (3.1

and

$$C_\tau = \{\tau \in C_{\tau,all} | \tau_i > 0 \ \forall \ \tau_i; \sum_{i=1}^{\sigma} \tau_i = 1\}$$ (3.2

where again, τ_i is the relative contribution by the i^{th} species to *T*, such that all τ sum to equal 1.

The community represented by equation 3.1 includes species not contributing to the community-

aggregated parameter. The community represented by equation 3.2 is a subset of equation 3.1,

where C_τ contains only species from $C_{\tau,all}$ contributing towards a community-aggregated

parameter, which we refer to as "contributors." As such, we adopted the variable, σ, which

represents the richness of contributors. The effective richness of contributors towards *T* was

calculated using equation 3.2 and the HCDT entropy index (Jost, 2006) such that:

$$D_q = \begin{cases} (\sum_{i=1}^{\sigma} \tau_i{}^q)^{\frac{1}{(1-q)}} & q \neq 1 \\ exp(\sum_{i=1}^{\sigma} -\tau_i \log \tau_i) & q=1 \end{cases}$$ (4

where τ_i is the relative contribution by i^{th} species in C_τ towards *T*, and D_q is the effective richness

of contributors, and *q* is the diversity order, which weights the importance of species contributing

more or less towards *T* when calculating D_q. This is analogous to diversity order weighting the

importance of rare versus abundant species when measuring diversity of species abundances

(Hill, 1973). With diversity order $q=0$, equation 4 simplifies to the number of contributors in

equation 3.2 (i.e., σ). Diversity order q=0 removes any influence in the variability of τ when

calculating D_q. Increasing the diversity order subsequently decreases the perceived contribution

of contributors with smaller τ. Diversity orders $q=1$ and $q=2$ correspond to Shannon's index. and

Simpson's index.

We then normalize D_q by community richness (equal to assuming every species

contributes to T equally), S, and adjust evenness to range from 0 to 1 (Chao & Ricotta, 2019),

such that:

$R = \frac{D_q - 1}{S - 1}$ (5

Here, functional redundancy, R, is taken as how evenly species contribute to a community-

aggregated parameter, T. For the hypothetical community described above, the effective richness

(D_q, $q=0.5$—see *Results*) of species possessing *xxxA* was ~1.55, and the functional redundancy,

with respect to *xxxA*, was $R=0.47$ ($q=0.5$; see *Results*).

Simulations of Functional Redundancy as a Function of Resilience to Species Extinction

In order to test the fidelity with which CE measures resilience of a community-

aggregated parameter, we simulated artificial communities with varying values of species

richness (S), richness of contributors (σ), and contribution levels by contributors (t). First, the

richness of contributors was randomly chosen from a uniformly distributed range of 1 to 100.

Next, the richness of species not contributing to the community-aggregated parameter was

randomly chosen from a uniformly distributed range of 0 to 100, so that some artificial

communities were dominated by contributors while others by non-contributors. The relative

abundance, p, of individual species was modeled log-normally (Galand, Casamayor, Kirchman,

& Lovejoy, 2009; Magurran, 1988) such that:

$R(r) = R_0 e^{-a^2 r^2}$　　　　　　　　　　　　　　　　　　　　　　　　　　(6

where R was the r^{th} rank, R_0 was the maximum abundance, and a was the inverse width of the

lognormal distribution. Values of a were randomly selected from 0 (even) to 0.4 (more uneven),

at 0.01 intervals. After establishing species relative abundances, we randomly chose σ

contributing taxa from our lognormally distributed community. The contribution level, t, was

then randomly assigned to these species from a uniformly distributed range of 0.01 to 1, at

intervals of 0.01. We generated and calculated R for 1,000 artificial communities.

The resilience of a community-aggregated parameter to extinction was tested by a

simulation in which resilience to extinction was quantified as the average fraction of species

richness (S) that must go extinct in order to decrease community-aggregated parameter by 99%.

Species were randomly removed from the community until summed τ was reduced to 1% of its

original level. The extinction simulation was performed 100 times for each of the individual

1,000 artificial communities. The resilience to extinction generated from the 100 simulations

were then averaged for each of the individual 1,000 artificial communities. We made no *a priori*

assumptions about the optimal diversity order (i.e., q) to use in equations 4 and 5. We performed

the simulation described above varying diversity orders from 0 to 2 at 0.1 intervals. The best

diversity order was determined by fitting generalized additive model (GAM) regressions

between R and resilience to extinction, and then choosing the diversity order which resulted in a

GAM fit with the lowest mean absolute error among residuals. All GAM fits were performed

using the gam function from the R package mgcv (Wood, 2017). Last, we performed the same

simulations and calculated R using other methods described in the literature. A summary of these

measurements is provided in Table 1. An R package which calculates functional redundancy

(equations 1-5) and performs these simulations is available at

https://github.com/taylorroyalty/functional_redundancy_script.

A Case Study of Functional Redundancy in the Global Ocean Nitrogen Cycle

Draft metagenome-assembled genomes (MAGs) from (Tully, Graham, et al., 2018) were

downloaded from NCBI (Federhen, 2012). Data on metagenome sequencing effort (number of

short reads), MAG relative abundance estimates (pre-corrected for genome size), ocean depths,

and ocean regions, defined in (Tully, Graham, et al., 2018), were downloaded from Figshare

(https://doi.org/10.6084/m9.figshare.5188273). MAG relative abundances from each site were

multiplied by the respective site's sequencing effort to convert relative abundances into

abundance count data. Abundance counts for all sites were normalized by the lowest abundance

(greater than 0) among all sites. MAG abundances were then rarefied using the rrarefy function

from the R package vegan (Philip, 2003). We chose a sampling effort of 14,265, as this sampling

effort corresponded to the site with lowest sampling effort (i.e., sequencing effort). We tabulated

nitrogen-cycle marker genes in individual MAGs by blasting MAG proteins against nitrogen-

cycle marker gene databases using the BLASTP algorithm (E-value=10^{-10}) (Clark, Karsch-

Mizrachi, Lipman, Ostell, & Sayers, 2019). Nitrogen cycle marker genes were categorized into

eight categories: N_2 fixation (*nifH*), nitrification (*amoA*, *hao*), NO_3 reduction (*napA*, *narG*),

dissimilatory NO_3 reduction (*nrfA*), assimilatory NO_3/NO_2 reduction (*narB*, *nasA*, *nasB*, and

nirB), NH_4^+ assimilation (*gdhA*, *glnA*), N remineralization (*gltB*), and denitrification (*nirK*, *nirS*,

norB, and *nosZ*) (Pajares & Ramos, 2019). Individual BLAST databases were constructed for

each nitrogen cycle category. Gene names associated with individual categories were searched

for in the Swiss-Prot database (Bairoch & Apweiler, 2000). All proteins matching gene names

were subsequently downloaded and used in BLAST database construction. Functional

redundancy was then calculated for individual stations, at individual depths, for each nitrogen

cycle category using equations 1-5. We used a diversity order, q=0.5, based on the GAM

sensitivity analysis. Table 2 contains protein names, gene names, corresponding EC, and entry

name for all genes used in the above analysis.

Results and Discussion

Representing microbial communities as species abundance, trait abundance, and relative

contribution distributions

We represent microbial communities using two distributions: the distribution of species

abundances within a community, and the distribution of contribution levels among species. Here,

we evaluate genomic traits as these are regularly adopted as proxies for microbially-mediated

biogeochemistry (Anantharaman et al., 2016; Baker, Lazar, Teske, & Dick, 2015; Falkowski,

Fenchel, & Delong, 2008; Krause et al., 2014; Malik et al., 2019; Salazar et al., 2019; Teeling et

al., 2012). Because trait levels are measured as non-negative, real numbers (Violle et al., 2007),

the relative contribution of individual species to a community-aggregated parameter can be

represented as a relative abundance distribution (Figure 1). We determined the relative

contribution of individual species to the community-aggregated parameter from the product of

species relative abundances and their respective trait abundance distributions (equation 2). We

refer to the distribution of the relative contribution of each species to total as the relative

contribution distribution (equation 3). Thus, an abundant species that expresses a small amount

of trait per cell, and a rarer species that expresses a large amount of trait per cell, can contribute

equally with respect to a community-aggregated parameter (Figure 1).

Defining, Quantifying Functional Redundancy, and Choosing a Diversity Order q

We measured functional redundancy as how evenly species in a community contribute to

a community-aggregated parameter and termed this metric as contribution evenness (CE). This

definition is consistent with descriptions of functional redundancy as multiple species sharing

similar ecological roles or functions (Naeem, 1998; Rosenfeld & Mall, 2002). The relative

contribution by species towards a community aggregated-parameter (equation 2) provides a

framework to calculate the effective number, or richness, of species that contribute to that

community-aggregated parameter as an ecological function. This effective richness is calculated

with traditional diversity indices (equations 3-4) (Hill, 1973; Jost, 2006). To determine the

effective proportion of a community contributing to the community-aggregated parameter, the

richness of contributors is normalized by species richness (Hill, 1973). Last, to make meaningful

comparisons of evenness between communities with different levels of richness, we made our

evenness metric unrelated to community richness (equation 5) (Chao & Ricotta, 2019).

Numerous diversity indices have been proposed in the literature. Three common

measures of diversity include species richness, Shannon's diversity index, and Simpson's

diversity index, are manifestations of equation 4 when the q term—commonly referred to as the

diversity order—is $q=0$, $q=1$, and $q=2$, respectively. The diversity order changes the relative

weight placed on rare versus common species when determining the effective richness. Here, we

adopted a diversity order $q=0.5$, based on empirical results from our numerical extinction

simulations (see *Functional Redundancy and Trait Resilience to Species Extinction* for details).

We found that GAM fits between functional redundancy and resilience to extinction followed

positive, monotonic functions for all diversity orders spanning $q=0$ to $q=2$. However, at $q\approx0.5$,

CE was most constrained (mean absolute error) with respect to resilience to extinction, our

benchmark for functional redundancy efficacy (Figure 2). As such, we thought it appropriate to

select the diversity order maximizing efficacy with respect to our benchmark.

For $0.1<q\leq2$, diversity order only weakly affected mean absolute error. Diversity orders

$q=0$ (richness), $q=1$ (Shannon's Index), and $q=2$ (Simpson's Index) correspond to the reciprocal

harmonic, geometric, and arithmetic means of the taxa proportions (p_i or τ_i), respectively. These

diversity orders are perceived as both intuitive and preferential diversity orders relative to

empirically derived diversity orders (Hill, 1973). Given diversity orders $0.1<q\leq2$ do not alter our

conclusions, we allow investigators to select a custom diversity order in the R package.

Functional Redundancy and Trait Resilience to Species Extinction

A tangible ecological consequence of functional redundancy is resilience of an ecosystem

function to the extinction of species performing that function. When functions are more broadly

distributed across species, a function is less likely to disappear due to the extinction of any subset

of species. This interpretation is often highlighted in macro-ecological studies (McPherson,

Yeager, & Baum, 2018; Naeem, 1998; Ricotta et al., 2016; Rosenfeld & Mall, 2002; Tilman,

2001). The conceptual relationship between functional redundancy and resilience of an

ecosystem function provides an opportunity to demonstrate the fidelity of CE. To do this, CE

was calculated for randomly generated communities. Individual species were randomly assigned

a contribution levels to a community aggregated parameter. We simulated random species

extinction events in the artificial communities until >99% of the original ecosystem function was

removed from the community. This choice of threshold allowed for the evaluation of

communities where all species were contributors. In such communities, a 100% extinction rate

was necessary to remove the ecosystem function.

It is important to emphasize that these simulations were not designed to mimic how

extinction occurs in nature. Instead, these simulations empirically demonstrate the degree to

which an ecosystem function is concentrated in species. The more widely an ecosystem function

is distributed across species, the more species must go extinct before the ecosystem function is

lost. We observed a positive, monotonic relationship between the resilience to random species

extinction events and CE (Figure 3A). When CE was ~0, resilience to species extinction was

~0.5, meaning ~50% of the community, on average, went extinct before an ecosystem function

was removed from the community. This is equivalent to only one species performing the

ecosystem function. When CE was ~1, resilience to species extinct was ~1. Again, this means

100% of the community, on average, had to go extinct to completely remove a function from an

ecosystem. In general, the relationship between CE and trait resilience to extinction resembled a

saturation curve—although we opted to use GAMs when calculating mean absolute error

(previous section) as we made no assumption about the true relationship between CE and

function resilience, other than it should be a positive, monotonic function.

CE Measures Resilience Differently than Other Functional Redundancy Metrics

Several methods for evaluating community-level functional redundancy exist in the

literature, as detailed in Table 1. While Table 1 certainly does not include all published metrics,

the highlighted metrics reflect the diversity in how traits, community properties (i.e., richness

and evenness), as well as the desired metric utility (by design) vary among different functional

redundancy metrics. Two of the metrics evaluate niche overlap, where niches are defined in

multivariate trait space (Hester et al., 2019; Ricotta et al., 2016). In the case of Ricotta et al.

(2016), functional redundancy weights species abundance, leading to a measure of the expected

niche dissimilarity from the mean community niche. In contrast, Hester et al. (2019) evaluates

the median metabolic overlap which considers neither species abundance nor trait levels. One

metric evaluates niche occupancy, where niches are defined in multivariate trait space (Mouillot

et al., 2014). In this approach, functional redundancy is taken to be as the ratio of species

richness to plausible number niches (unique combinations of trait states), called functional

entities. The last metric is informal but has been utilized by various authors. In this approach,

functional redundancy is simply interpreted as the proportion of species in a community with a

trait (e.g., Trivedi, 2020; Tully, Wheat, et al., 2018; Wohl, Arora, & Gladstone, 2004).

327   To highlight the difference between our metric and the others described in Table 1, we

repeated the extinction simulations and compared all estimates of functional redundancy with

trait resilience to species extinction (Figure 3B-E). In these simulations, only one trait was

considered when calculating functional redundancy with all metrics. For the method presented in

Mouillot et al. (2014), we assumed there were two functional entities (trait absence and

presence). Generally, all functional redundancy metrics trended differently to resilience to

species extinction. This observation was not surprising, as most of these methods evaluate niche

overlap in some fashion. Among the metrics in Table 1, the fraction of the community

possessing a trait behaved most similar to CE (Figure 3B). We note that the fraction of the

community possessing a trait is a special case of CE, with diversity order q=0. Lastly, one

inherent limitation of CE is that it only evaluates single community-aggregated parameters at a

time. To address functional redundancy with respect to multiple traits perform dimensional

reduction (e.g., PCA) on multiple traits and use the resultant factors as trait levels in equation 2.

Factors in this case would be interpreted as different niches.

Case Study: Redundancy of Nitrogen Cycle Marker Genes in Marine Microbiomes

To demonstrate a use of CE, we measured functional redundancy in the marine nitrogen

cycle using genomic data from the *TARA Oceans* project (Sunagawa et al., 2015). Recently,

(Tully, Graham, et al., 2018) published 2,361 draft metagenome-assembled genomes (MAGs)

which represent genomic content from seven oceanographic provinces, three different depths,

and four filter size fractions. Relative abundance data used here were reported in (Tully, Graham,

et al., 2018) and were determined by the proportion of sequence reads mapping to assembled

MAGs. For simplicity, we limited our analysis to only MAGs identified in the bacterial size

fraction (<3 μm), which corresponds to free-living microbes. The nitrogen cycle was simplified

to eight pathways (Pajares & Ramos, 2019) and included N_2 fixation, nitrification, NO_3^-

reduction, dissimilatory NO_3^- reduction, assimilatory NO_3^-/NO_2^- reduction, NH_4^+ assimilation, N

remineralization, and denitrification. A general summary of the marker genes (downloaded from

Swiss-Prot; see Methods) used for quantifying nitrogen cycle pathways in MAGs is summarized

in Table 2.

Overall, the functional redundancy for individual genomic traits was consistent with our

conceptual understanding of primary nitrogen-transforming pathways in the ocean (Figure 4). A

clear example was the low functional redundancy in the nitrification pathway. We expected this

observation as nitrification metabolisms in oceans are primarily restricted to *Crenarchaeota*

(Wuchter et al., 2006). Similarly, the N_2 fixation pathway, which is predominantly facilitated by

a small group of *Cyanobacteria*, exhibits low functional redundancy. Conversely, the uptake of

inorganic N via NH_4^+ and NO_3^-/NO_2^- assimilation are known to be broadly expressed traits

(Pajares & Ramos, 2019), and accordingly exhibit high CE. Last, the low functional redundancy

in dissimilatory NO_3^- reduction and NO_3^- reduction were not surprising considering these

pathways occur in anaerobic and low-oxygen environments, respectively (Pajares & Ramos,

2019).

Some variance existed across oceanographic provinces and depth for intra-pathway

functional redundancy. However, inter-pathway variance in functional redundancy was generally

larger than intra-pathway variance (Figure 4). This suggests that the marine nitrogen cycle

marker genes considered here are consistently distributed among communities for MAGs

retrieved in Earth's ocean at middle and lower latitudes. This was consistent with the lack of

relationship between the metagenome companion metadata (e.g., O_2, inorganic N, PO_4^{3-}, latitude,

longitude, etc.) and functional redundancy. The low intra-pathway variance in functional

redundancy was not particularly surprising, as saltwater biomes exhibit the lowest alpha-

diversity and small beta-diversity among free-living microbiomes (Sunagawa et al., 2015;

Thompson et al., 2017). Minor shifts in community composition among lower taxonomic ranks

(e.g., genus or species) will not drastically alter redundancy in genomic traits as changes in

genomic composition is buffered by traits conserved at higher taxonomic ranks (e.g., phylum

level) (Royalty & Steen, 2019). This observation was in agreement with an earlier analysis

evaluating cluster of orthologous groups (COGs) composition across *TARA Oceans*

metagenomes (Sunagawa et al., 2015).

A more recent *TARA Oceans* analysis expanded sampling coverage to include more

metagenomes at polar latitudes. Community composition was found to be significantly different

at polar latitudes relative to lower and middle latitudes (Salazar et al., 2019). In this analysis,

eggNOG functional richness was compared to OTU richness across all metagenomes, leading to

the conclusion that metagenomes had low functional redundancy—the opposite conclusion from

Sunagawa et al. (2015). The discrepancy between these two analyses probably reflects

differences in how genes were annotated (COGs versus eggNOG). eggNOG orthologous groups

are defined by sequence homology and phylogeny, so gene clusters from different clades can

have the same function but different eggNOG annotation (Huerta-Cepas et al., 2019). Thus,

changes in biodiversity should give rise to a corresponding change in functional diversity if using

eggNOG annotations—again, annotations should correlate strongly with taxonomy. More

broadly speaking, this emphasizes the importance of properly contextualizing genomic trait

functional redundancy to account for annotation scheme.

In our analysis of *TARA Oceans* metagenomes, we used genomic traits as proxies for

nitrogen metabolisms. As a point of emphasis, measurable properties (e.g., nitrogen uptake)

provide direct measures of ecosystem function that are more easily integrated into predictive

models. In an ideal scenario, the latter would be discernible in all ecological studies, for all

ecosystem functions. However in the case of microbes, a disproportionate number of lineages

that lack cultured representatives highlights the continued importance of nucleotide sequencing

in microbial ecology (Lloyd et al., 2018). Consequently, the microbiological community relies

on 'omics technology (genes, transcripts, or proteins) to characterize community function.

Conceivably, other 'omics techniques such as metatranscriptomics and metaproteomics can be

evaluated with CE as well. In these approaches, the observed metatranscript and metapeptide

sequence abundances implicitly incorporate relative abundances of species, and do not require

scaling lineage abundances as is done in equation 2. Furthermore, technological advances with

single-cell nanoSIMS may provide a method for quantifying lineage-specific metabolic rates for

in situ inoculation studies (Musat, Foster, Vagner, Adam, & Kuypers, 2012). Coupling these

metabolic rates with abundance data, we believe CE could measure functional redundancy of

phenotypes.

Data Availability

In order to facilitate the use of CE functional redundancy measurements, we created an R package to calculate CE functional redundancy and run extinction simulations. The package is available at https://github.com/taylorroyalty/functional_redundancy_script.

Acknowledgments

We gratefully acknowledge the comments of three anonymous reviewers, whose constructive comments substantially improved the manuscript. Funding for this project was provided by a C-DEBI graduate fellowship to TMR and by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science Program under Award Number DE-SC0020369 for TMR and ADS. This is C-DEBI contribution number [to be assigned upon manuscript acceptance].

Author Contribution

TMR designed the research, performed research, analyzed data, developed the R package, and wrote the paper. ADS designed the research, developed the R package, and wrote the paper.

References

Anantharaman, K., Brown, C. T., Hug, L. A., Sharon, I., Castelle, C. J., Probst, A. J., …

Banfield, J. F. (2016). Thousands of microbial genomes shed light on interconnected

biogeochemical processes in an aquifer system. *Nature Communications*, *7*, 13219. doi:

10.1038/ncomms13219

Bairoch, A., & Apweiler, R. (2000). The SWISS-PROT protein sequence database and its

supplement TrEMBL in 2000. *Nucleic Acids Research*, *28*(1), 45–48. doi:

10.1093/nar/28.1.45

Baker, B. J., Lazar, C. S., Teske, A. P., & Dick, G. J. (2015). Genomic resolution of linkages in

carbon, nitrogen, and sulfur cycling among widespread estuary sediment bacteria.

Microbiome, *3*(1), 14. doi: 10.1186/s40168-015-0077-6

Chao, A., & Ricotta, C. (2019). *Quantifying evenness and linking it to diversity , beta diversity ,*

and similarity. *100*(12), 1–15. doi: 10.1002/ecy.2852

Clark, K., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2019). GenBank.

Nucleic Acids Research, *47*(D1), D94–D99. doi: 10.1093/nar/gky989

Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The Microbial Engines That Drive Earth's

Biogeochemical Cycles. *Science*, *320*(5879), 1034–1039. doi: 10.1126/science.1153213

Federhen, S. (2012). The NCBI Taxonomy database. *Nucleic Acids Research*, *40*(D1), 136–143.

doi: 10.1093/nar/gkr1178

Fierer, N., Barberán, A., & Laughlin, D. C. (2014). Seeing the forest for the genes: Using

metagenomics to infer the aggregated traits of microbial communities. *Frontiers in*

Microbiology, *5*(NOV), 1–6. doi: 10.3389/fmicb.2014.00614

Galand, P. E., Casamayor, E. O., Kirchman, D. L., & Lovejoy, C. (2009). Ecology of the rare

microbial biosphere of the Arctic Ocean. *Proceedings of the National Academy of Sciences*,

106(52), 22427–22432. doi: 10.1073/pnas.0908284106

Galand, P. E., Pereira, O., Hochart, C., Auguet, J. C., & Debroas, D. (2018). A strong link

between marine microbial community composition and function challenges the idea of

functional redundancy. *ISME Journal*, *12*(10), 2470–2478. doi: 10.1038/s41396-018-0158-1

Hester, E. R., Jetten, M. S. M., Welte, C. U., & Lücker, S. (2019). Metabolic overlap in

environmentally diverse microbial communities. *Frontiers in Genetics*, *10*(OCT), 1–10. doi:

10.3389/fgene.2019.00989

Hill, M. O. (1973). Diversity and Evenness : A Unifying Notation and Its Consequences.

Ecology, *54*(2), 427–432.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., …

Bork, P. (2019). EggNOG 5.0: A hierarchical, functionally and phylogenetically annotated

orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Research*,

47(D1), D309–D314. doi: 10.1093/nar/gky1085

Jost, L. (2006). *Entropy and diversity*. *2*.

Krause, S., Le Roux, X., Niklaus, P. A., van Bodegom, P. M., Lennon T., J. T., Bertilsson, S., …

Bodelier, P. L. E. (2014). Trait-based approaches for understanding microbial biodiversity

and ecosystem functioning. *Frontiers in Microbiology*, *5*(MAY), 1–10. doi:

10.3389/fmicb.2014.00251

Lloyd, K. G., Steen, A. D., Ladau, J., Yin, J., & Crosby, L. (2018). Phylogenetically Novel

Uncultured Microbial Cells Dominate Earth Microbiomes. *MSystems*, *3*(5), 1–12. doi:

10.1128/mSystems.00055-18

Louca, S., Martiny, A. C., Polz, M. F., Doolittle, W. F., Nelson, M. B., Hahn, A. S., … Hallam,

S. J. (2018). Function and functional redundancy in microbial communities. *Nature Ecology*

and Evolution, *2*, 936–943.

Magurran, A. E. (1988). *Ecological Diversity and Its Measurement* (1st ed.). doi: 10.1007/978-

94-015-7358-0

Malik, A. A., Martiny, J. B. H., Brodie, E. L., Martiny, A. C., Treseder, K. K., & Allison, S. D.

(2019). Defining trait-based microbial strategies with consequences for soil carbon cycling

under climate change. *ISME Journal*, 445866. doi: 10.1101/445866

McGill, B. J., Etienne, R. S., Gray, J. S., Alonso, D., Anderson, M. J., Benecha, H. K., … White,

E. P. (2007). Species abundance distributions: Moving beyond single prediction theories to

integration within an ecological framework. *Ecology Letters*, *10*(10), 995–1015. doi:

10.1111/j.1461-0248.2007.01094.x

McPherson, J. M., Yeager, L. A., & Baum, J. K. (2018). A simulation tool to scrutinise the

behaviour of functional diversity metrics. *Methods in Ecology and Evolution*, *9*(1), 200–

206. doi: 10.1111/2041-210X.12855

Micheli, F., & Halpern, B. S. (2005). Low functional redundancy in coastal marine assemblages.

Ecology Letters, *8*(4), 391–400. doi: 10.1111/j.1461-0248.2005.00731.x

Miki, T., Yokokawa, T., & Matsui, K. (2013). Biodiversity and multifunctionality in a microbial

community: A novel theoretical approach to quantify functional redundancy. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1776). doi: 10.1098/rspb.2013.2498

Mouillot, D., Villéger, S., Parravicini, V., Kulbicki, M., Arias-González, J. E., Bender, M., … Bellwood, D. R. (2014). Functional over-redundancy and high functional vulnerability in global fish faunas on tropical reefs. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(38), 13757–13762. doi: 10.1073/pnas.1317625111

Musat, N., Foster, R., Vagner, T., Adam, B., & Kuypers, M. M. M. (2012). Detecting metabolic activities in single cells, with emphasis on nanoSIMS. *FEMS Microbiology Reviews*, *36*(2), 486–511. doi: 10.1111/j.1574-6976.2011.00303.x

Naeem, S. (1998). *Species Redundancy and Ecosystem Reliability*. *12*(1), 39–45.

Orsi, W. D., Wilken, S., del Campo, J., Heger, T., James, E., Richards, T. A., … Santoro, A. E. (2018). Identifying protist consumers of photosynthetic picoeukaryotes in the surface ocean using stable isotope probing. *Environmental Microbiology*, *20*(2), 815–827. doi: 10.1111/1462-2920.14018

Pajares, S., & Ramos, R. (2019). Processes and Microorganisms Involved in the Marine Nitrogen Cycle: Knowledge and Gaps. *Frontiers in Marine Science*, *6*(November). doi: 10.3389/fmars.2019.00739

Philip, D. (2003). Computer program review VEGAN , a package of R functions for community ecology. *Journal of Vegetation Science*, *14*, 927–930. doi: 10.1111/j.1654-1103.2003.tb02228.x

Ricotta, C., de Bello, F., Moretti, M., Caccianiga, M., Cerabolini, B. E. L., & Pavoine, S. (2016).

Measuring the functional redundancy of biological communities: a quantitative guide. *Methods in Ecology and Evolution*, *7*(11), 1386–1395. doi: 10.1111/2041-210X.12604

Rosenfeld, J. S., & Mall, M. (2002). Functional redundancy in ecology and conservation. *OIKOS*, *1*.

Royalty, T. M., & Steen, A. D. (2019). Quantitatively Partitioning Microbial Genomic Traits among Taxonomic Ranks across the Microbial Tree of Life. *MSphere*, *4*(4), e00446-19. doi: 10.1128/mSphere.00446-19

Salazar, G., Paoli, L., Alberti, A., Huerta-Cepas, J., Ruscheweyh, H. J., Cuenca, M., … Wincker, P. (2019). Gene Expression Changes and Community Turnover Differentially Shape the Global Ocean Metatranscriptome. *Cell*, *179*(5), 1068-1083.e21. doi: 10.1016/j.cell.2019.10.014

Staley, J., & Konopka, A. (1985). Measurement of In Situ Activities of Nonphotosynthetic Microorganisms in Aquatic and Terrestrial Habitats. *Annual Review of Microbiology*, *39*(1), 321–346. doi: 10.1146/annurev.micro.39.1.321

Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G., … Bork, P. (2015). Structure and function of the global ocean microbiome. *Science*, *348*(6237), 1–10. doi: 10.1126/science.1261359

Teeling, H., Fuchs, B. M., Becher, D., Klockow, C., Gardebrecht, A., Bennke, C. M., … Amann, R. (2012). Substrate-Controlled Succession of Marine Bacterioplankton Populations Induced by a Phytoplankton Bloom. *Science*, *336*(May), 608–611. doi: 10.1126/science.1218344

Thompson, L. R., Sanders, J. G., McDonald, D., Amir, A., Ladau, J., Locey, K. J., … Zhao, H.

(2017). A communal catalogue reveals Earth's multiscale microbial diversity. *Nature*,

551(7681), 457–463. doi: 10.1038/nature24621

Tilman, D. (2001). Functional Diversity. *Encyclopedia of Biodiversity: Second Edition*, *3*, 587–

596. doi: 10.1016/B978-0-12-384719-5.00061-7

Trivedi, C. B., Stamps, B. W., Lau, G. E., Grasby, S. E., Templeton, A. S., & Spear, J. R. (2020).

Microbial Metabolic Redundancy Is a Key Mechanism in a Sulfur-Rich Glacial Ecosystem.

MSystems, *5*(4), 1–16. doi: https://doi.org/10.1128/mSystems .00504-20.

Tully, B. J., Graham, E. D., & Heidelberg, J. F. (2018). The reconstruction of 2,631 draft

metagenome-assembled genomes from the global oceans. *Scientific Data*, *5*, 1–8. doi:

10.1038/sdata.2017.203

Tully, B. J., Wheat, C. G., Glazer, B. T., & Huber, J. A. (2018). A dynamic microbial

community with high functional redundancy inhabits the cold, oxic subseafloor aquifer.

ISME Journal, *12*(1), 1–16. doi: 10.1038/ismej.2017.187

Violle, C., Navas, M., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., … Fortunel, C. (2007).

Let the concept of trait be functional! *Oikos*, *116*(5), 882–892. doi:

https://doi.org/10.1111/j.0030-1299.2007.15559.x

Walker, B. H. (1992). Biodiversity and Ecological Redundancy. *Conservation Biology*, *6*(1), 18–

23. doi: 10.1046/j.1523-1739.1992.610018.x

Wohl, D. L., Arora, S., & Gladstone, J. R. (2004). Functional Redundancy Supports Biodiversity

and Ecosystem Function in a Closed and Constant Environment. *Ecology*, *85*(6), 1534–

1540.

Wood, S. (2017). *mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML Smoothness*

Estimation.

Wuchter, C., Abbas, B., Coolen, M. J. L., Herfort, L., Van Bleijswijk, J., Timmers, P., …

Damsté, J. S. S. (2006). Archaeal nitrification in the ocean. *Proceedings of the National*

Academy of Sciences of the United States of America, *103*(33), 12317–12322. doi:

10.1073/pnas.0600756103

Data Accessibility

We provided a function for calculating functional redundancy with CE and an example script at

https://github.com/taylorroyalty/functional_redundancy_script.

Author Contribution

TMR designed the research, performed research, analyzed data, and wrote the paper. ADS

designed the research, performed research, and wrote the paper.

Tables

Table 1. A summary of metrics of community-level functional redundancy.

Authors	Interpretation	Equation	Notes
Here	Effective proportion of species with a trait	$R = \dfrac{D_q - 1}{S - 1}$	• Accounts for species abundances • Continuous trait states • Individual traits • R spans 0 to 1
(Ricotta et al., 2016)	Expected trait dissimilarity among species in a community	$R = 1 - \dfrac{\sum_i p_i \sum_j p_j \delta_{ji}}{D_2}$; $\delta_{ji} = trait\ distance$; $D_2 = $ Taxonomic diversity order $(q = 2)$	• Accounts for species abundances • Continuous trait states • Individual or multivariate traits • Normalized by a maximum theoretical dissimilarity • R spans 0 to 1
(Mouillot et al., 2014)	Ratio of species to number of functional entities (states)	$R = \dfrac{S}{FE}$; FE = # of possible combinations of trait states for all traits under consideration	• Binary trait states • Individual or multivariate traits • R spans ~0 ($\lim\limits_{FE \to \infty} \dfrac{S}{FE} = 0$) to S
Various (see text for examples)	Fraction of species with a trait	$R = \dfrac{v}{S}$	• Binary trait states • Individual traits • R spans 0 to 1
(Hester et al., 2019)	Median overlap of traits between species	$R = med(O)$; $O = \{o \mid o_{i,j} = \lvert g_i \cap g_j \rvert; i \neq j\}$; $g = taxa$; O = set containing # of overlaps among species	• Binary trait states • $\lvert g_i \cap g_j \rvert$ represents the number of overlapping traits g between species i and species j • Individual and multivariate traits • R spans 0 to ∞

Table 2. A summary of marker gene sequences used in the BLASTP analysis of *TARA*
Oceans MAGs. All marker genes were downloaded from Swiss-Prot.

Pathway	Gene Names	EC Number	# Marker Genes
N Remineralization	*gltB*	1.4.1.13	14
N Fixation	*nifH*	1.18.6.1	127
Nitrification	*amoA*	1.14.99.39	2
	hoa	1.7.2.6	1
NO$_3^-$ Reduction	*napA*	1.9.6.1	136
	narG	1.7.5.1	13
Dissimilatory NO$_3^-$ Reduction	*nrfA*	1.7.2.2	77
Assimilatory NO$_3^-$/NO$_2^-$ Reduction	*narB*	1.7.5.1	3
	nasA	1.7.7.2	1
	nasB	1.7.1.4	3
	nirB	1.7.1.15	5
NH$_4^+$ Assimilation	*gdhA*	1.4.1.3	14
	glnA	6.3.1.2	173
Denitrification	*nirK*	1.7.2.1	7
	nirS	1.7.2.1; 1.7.99.1	4
	norB	1.7.2.5	2
	nosZ	1.7.2.4	11

Figures

Figure 1.

Community taxon abundance and relative contribution level are scalable and provide a measure

for the contribution that different species have towards a community-aggregated parameter. The

evenness of species-level contribution to the community-aggregated parameter is taken as the

functional redundancy.

Figure 2.

The sensitivity of mean absolute error between functional redundancy and trait resilience to

species extinction in response changing diversity order q in equation 4.

Figure 3.

A comparison of the relationship between selected different functional redundancy metrics and trait resilience (see methods). Functional redundancy calculated with equation 5 (q=0.5) (**A**), equation 5 (q=0) (**B**), methodology from Ricotta et al. (2016) (**C**), methodology from Mouillot et al. (2014) (**D**), and methodology from Hester et al. (2019). Black lines in panels **A-D** correspond to best fit GAMs. The stars in panel **E** correspond to the average trait resilience at R=0 and R=1. The color scale corresponds to the fraction of a community with a trait versus.

Figure 4.

Boxplots showing functional redundancy calculated for eight nitrogen-transforming pathways in

TARA Oceans draft MAGs. Different colors correspond to sample depth. Whiskers correspond to

150% of the inter-quantile region.