Structural basis for PRC2 decoding of active histone methylation marks H3K36me2/3

Ksenia Finogenova(1), Jacques Bonnet (1), Simon Poepsel(2,3), Ingmar B. Schäfer(4), Katja

Finkl(1), Katharina Schmid(1), Claudia Litz(1), Mike Strauss(5,6), Christian Benda(4) and Jürg

Müller(1)

1)   Max Planck Institute of Biochemistry, Laboratory of Chromatin Biology, Martinsried,
Germany
2)   California Institute for Quantitative Biology (QB3), University of California, Berkeley,
Molecular Biophysics and Integrative Bio-Imaging Division, Lawrence Berkeley
National Laboratory, Berkeley, CA
3)   University of Cologne, Center for Molecular Medicine Cologne (CMMC), Faculty of
Medicine and University Hospital Cologne, Cologne, Germany, Cologne Excellence
Cluster for Cellular Stress Responses in Ageing-Associated Diseases (CECAD),
University of Cologne, Cologne, Germany
4)   Max Planck Institute of Biochemistry, Department of Structural Cell Biology,
Martinsried, Germany
5)   Max Planck Institute of Biochemistry, cryoEM Facility, Martinsried, Germany
6)   Current address: McGill University, Department of Anatomy and Cell Biology,
Montreal, Canada
21

22 Correspondence should be addressed to J.M. (muellerj@biochem.mpg.de)

ABSTRACT

Repression of genes by Polycomb requires that PRC2 modifies their chromatin by trimethylating lysine 27 on histone H3 (H3K27me3). At transcriptionally active genes, di- and trimethylated H3K36 inhibit PRC2. Here, the cryo-EM structure of PRC2 on dinucleosomes reveals how binding of its catalytic subunit EZH2 to nucleosomal DNA orients the H3 N-terminus via an extended network of interactions to place H3K27 into the active site. Unmodified H3K36 occupies a critical position in the EZH2-DNA interface. Mutation of H3K36 to arginine or alanine inhibits H3K27 methylation by PRC2 on nucleosomes *in vitro*. Accordingly, *Drosophila* H3K36A and H3K36R mutants show reduced levels of H3K27me3 and defective Polycomb repression of HOX genes. The relay of interactions between EZH2, the nucleosomal DNA and the H3 N-terminus therefore creates the geometry that permits allosteric inhibition of PRC2 by methylated H3K36 in transcriptionally active chromatin.

INTRODUCTION

Many post-translational modifications on histone proteins are essential for processes in the underlying chromatin. Typically, histone modifications themselves do not alter chromatin structure directly but function by binding effector proteins which alter chromatin or by interfering with such interactions. The histone methyltransferase Polycomb Repressive Complex 2 (PRC2) and its regulation by accessory proteins and histone modifications represent a prime example for understanding these interaction mechanisms (Laugesen et al., 2019; Yu et al., 2019). PRC2 trimethylates lysine 27 in histone H3 (H3K27me3), a modification that is essential for the transcriptional repression of developmental regulator genes that control cell fate decisions in metazoans (McKay et al., 2015; Pengelly et al., 2013). H3K27me3 marks chromatin for interaction with PRC1, an effector which compacts chromatin (Francis et al., 2004; Grau et al., 2011). H3K27me3 is also recognized by PRC2 itself, and this interaction allosterically activates the PRC2 enzyme complex to facilitate deposition of H3K27me3 across

extended domains of chromatin (Hansen et al., 2008; Jiao and Liu, 2015; Margueron et al.,

2009). Genetic studies and subsequent biochemical work established that PRC2 is in addition

subject to negative regulation. In particular, the H3K4me3, H3K36me2 and H3K36me3 marks

present on nucleosomes in transcriptionally active chromatin directly inhibit H3K27

methylation by PRC2 (Gaydos et al., 2012; Klymenko and Müller, 2004; Schmitges et al., 2011;

Streubel et al., 2018; Yuan et al., 2011). Importantly, while stimulation of PRC2 activity by

H3K27me3 acts *in trans*, inhibition of PRC2 by H3K4me3, H3K36me2 and H3K36me3

requires that these modifications are present *in cis*, that is, on the same H3 molecule containing

the K27 substrate lysine (Schmitges et al., 2011; Yuan et al., 2011). While recent structural

studies have uncovered the allosteric activation mechanism for PRC2 (Jiao and Liu, 2015;

Justin et al., 2016), the molecular basis of PRC2 inhibition by active chromatin marks has

remained enigmatic. In particular, in nucleosome binding assays, PRC2-DNA interactions

make the largest contribution to the nucleosome-binding affinity of PRC2 (Choi et al., 2017;

Wang et al., 2017) and H3K4me3, H3K36me2 and H3K36me3 do not seem to have a major

effect on this binding affinity (Guidotti et al., 2019; Jani et al., 2019; Schmitges et al., 2011).

Instead, these three modification were found to reduce the k_{cat} of PRC2 for H3K27 methylation

(Jani et al., 2019; Schmitges et al., 2011). Recent cross-linking studies led to the suggestion of

a possible sensing pocket for H3K36 on the surface of EZH2 (Jani et al., 2019) but there is no

structural data how that proposed interaction might occur. Similarly, a recent structure of PRC2

bound to a dinucleosome revealed how the catalytic lobe of PRC2 contacts nucleosomes

through DNA interactions but provided no structural insight into how the H3 N-termini might

be recognized (Poepsel et al., 2018). Here, a refined structure of PRC2 bound to a

dinucleosome allowed us to visualize how the histone H3 N-terminus on substrate nucleosomes

is threaded into the EZH2 active site. Our analyses reveal that H3K36 assumes a critical

position in the PRC2-nucleosome interaction interface that permits the complex to gauge the

H3K36 methylation state.

 RESULTS

 EZH2 interaction with nucleosomal DNA orients the H3 N-terminus for H3K27 binding

 to the active site

We assembled recombinant full-length human PRC2 in complex with its accessory factor PHF1

(i.e. PHF1-PRC2) (Choi et al., 2017) on a heterodimeric dinucleosome (di-Nuc), which

consisted of a 'substrate' nucleosome with unmodified histone H3 and an 'allosteric'

nucleosome containing H3 with a trimethyllysine analog (Simon et al., 2007) at K27, separated

by a 35 base pair (bp) DNA linker (Poepsel et al., 2018) (**Figure 1A, B**). Single particle cryo-

electron microscopy analysis yielded a reconstruction of the PHF1-PRC2:di-Nuc assembly

with an overall resolution of 5.2 Å (**Supplementary Figures 1-3**). The map showed clear

density for the catalytic lobe of PRC2 with similar chromatin interactions and binding geometry

as previously described for the catalytic lobe of AEBP2-PRC2 (Poepsel et al., 2018) where

PRC2 contacts the two nucleosomes via interactions with the DNA gyres (**Figure 1C**).

Specifically, the substrate nucleosome is bound by the EZH2$_{CXC}$ domain residues K563, Q565,

K569 and Q570 (**Figure 1D, Supplementary Figure 4A**, cf. (Poepsel et al., 2018)), while the

allosteric nucleosome is contacted by EED and by the SBD and SANT1 domains of EZH2

(**Figure 1E**, cf. (Poepsel et al., 2018)). We could not detect density for the 'bottom lobe' of

PRC2 (Chen et al., 2018; Kasinath et al., 2018) or for the N-terminal winged-helix and tudor

domains of PHF1 that bind DNA and H3K36me3, respectively (Ballaré et al., 2012; Cai et al.,

2013; Choi et al., 2017; Li et al., 2017; Musselman et al., 2013).

Using particle signal subtraction and focused refinement on the interface of EZH2 and

the substrate nucleosome (**Supplementary Figures 2-3**), we then obtained an improved map

at an apparent overall resolution of 4.4 Å which revealed well-defined density for the H3 N-

terminus (**Figure 1F, Supplementary Figure 3B-D**). The visible sidechain density combined

with the crystallographic models of the PRC2 catalytic lobe and of the mononucleosome

enabled us to build a pseudo-atomic model of the histone H3 N-terminus spanning residues

R26 to K37 (**Figure 1F**). This model revealed that EZH2 recognizes the H3 N-terminus via an

extended network of contacts besides the previously described ionic interactions near the active

site where H3 R26 interacts with EZH2 Q648/D652, and H3 K27 with the aromatic cage above

the EZH2 catalytic center (Justin et al., 2016) (**Figure 1F**). Specifically, our structure suggests

two hydrophobic hotspots, the first one involving H3 A29/P30 and EZH2 residues F667, A697,

V699, I708 and F724 and the second one involving H3 V35 and F542, F557 and P558 of EZH2

(**Figure 1F**). H3 G33/G34 is likely not recognized by PRC2 but might act as a flexible hinge

between the two hydrophobic interaction sites (**Figure 1F**). H3K36 is directly juxtaposed to the

EZH2$_{CXC}$-DNA interaction surface and appears to be involved in the EHZ2-DNA interface. The

side chain density of H3K36 suggests that the epsilon-amino group of H3K36 engages in a

polar interaction with the carbonyl group of Q570 and in long-range electrostatic interactions

with the phosphate backbone of the nucleosomal DNA (**Figure 1F, Supplementary Figure 4**

C-E). Taken together, our analyses reveal that an extensive network of interactions between

EZH2, the nucleosomal DNA and the H3 N-terminus. This complex geometric arrangement

orients the H3 N-terminus into an extended conformation, threading H3K27 into the EZH2

active site. In this context, it should be noted that a previously postulated H3K36-binding

pocket centered on E579 of EZH2 (Jani et al., 2019) is located approximately 19 Å away from

H3K36 in our structure (**Supplementary Figure 4F**). An interaction of H3K36 with E579 of

EZH2 as proposed by Muir and co-workers (Jani et al., 2019) would require a very different

binding geometry of PRC2 on the nucleosome and major structural rearrangements of PRC2 or

the nucleosome in order to avoid steric clashes.

The EZH2 CXC contact with DNA is essential for H3K27 methylation

We next analysed how the PRC2 surfaces contacting the substrate and the allosteric nucleosome

contribute to the formation of productive PRC2-chromatin interactions. For these experiments,

we used PHF1$_C$-PRC2, which contains the minimal 5-kDa PRC2-interaction domain of PHF1

(**Figure 1A**, (Chen et al., 2020; Choi et al., 2017)) but lacks the H3K36me3-binding tudor and

the DNA-binding winged-helix domains of PHF1 (Choi et al., 2017; Li et al., 2017; Musselman

et al., 2013). PHF1$_C$-PRC2 therefore only retains the DNA-binding surfaces of the 4-subunit

PRC2 core complex and was used because it generally behaved better in purifications than the

4-subunit PRC2 core complex. For simplicity we shall, in the following, refer to the PHF1$_C$-

PRC2 complex as PRC2. We generated three mutant versions of PRC2. In PRC2$^{CXC>A}$ (K563A

Q565A K569A Q570A), the EZH2$_{CXC}$ interface is mutated (**Figure 1D**), in PRC2$^{EED>A}$ (K77A

K83A K385A K398A K400A K408A), the EED interface contacting the allosteric nucleosome

(**Figure 1E**), is mutated, and PRC2$^{CXC>A/EED>A}$ carries the combination of these mutations. We

first used electromobility shift assays (EMSA) to measure the binding affinity of wild-type and

mutant PRC2 complexes on mononucleosomes. These mononucleosomes were assembled on

a 215 bp long DNA fragment containing the 147-bp 601 nucleosome-positioning sequence

(Lowary and Widom, 1998) in the center and linker DNA on both sides. Wild-type PRC2

bound this mononucleosome with an apparent K$_d$ in the mid-nanomolar range (**Figure 2A, B**,

cf. (Choi et al., 2017)). The binding affinities of PRC2$^{CXC>A}$ or PRC2$^{EED>A}$ were two- to three-

fold lower than that of wild-type PRC2 and that of PRC2$^{CXC>A/EED>A}$ was about five-fold lower

compared to the wild-type complex (**Figure 2A, B**, compare lanes 11-30 with 1-10, and

Supplementary Figure 5A). The residual nucleosome binding shown by PRC2$^{CXC>A/EED>A}$

(**Figure 2A**, lanes 21-30) could in part be due to incomplete disruption of the mutated interfaces

but in part is likely also due to the previously identified nucleosome-binding activity of the

PRC2 bottom lobe (Chen et al., 2018; Nekrasov et al., 2005). The overall binding affinity of

the PRC2 core complex for chromatin therefore appears to result from interactions of at least

three distinct surfaces of this complex with nucleosomes. Among those, binding of the

EZH2$_{CXC}$ domain to the nucleosomal DNA contributes only modestly to the total binding

affinity.

We then analysed the histone methyltransferase (HMTase) activity of PRC2$^{CXC>A}$. On

the same mononucleosomes used above, PRC2$^{CXC>A}$ showed almost no detectable HMTase

activity compared to wild-type PRC2 (**Figure 2C**, compare lanes 5-7 with 2-4, see also

Supplementary Figure 5B). On dinucleosomes, EED binding to one nucleosome might be

expected to facilitate interaction of the mutated EZH2$^{CXC>A}$ domain with the H3 N-termini on

the juxtaposed second nucleosome. Indeed, on dinucleosomes, the PRC2$^{CXC>A}$ complex does

generate H3K27me1 and -me3 but less efficiently than wild-type PRC2 (**Figure 2C**, compare

lanes 12-14 with 9-11). When comparing the activities of the different complexes, it should be kept in mind that the interpretation of H3K27me3 formation as read-out of complex activity on dinucleosomes is more complicated than on mononucleosomes, because H3K27me3, once placed on one of the nucleosomes, will allosterically activate PRC2 to methylate H3K27 on the linked second nucleosome (Margueron et al., 2009) (Jiao and Liu, 2015). The main conclusion to be drawn from our analyses here is that the DNA-binding interaction of the EZH2$_{CXC}$ domain with substrate nucleosomes is critical for engaging the H3 N-terminus in a manner that allows H3K27 methylation.

Unmodified H3K36 in the EZH2$_{CXC}$-DNA interaction interface is critical for H3K27 methylation in nucleosomes

The architecture of the EZH2$_{CXC}$-DNA interface around H3K36 (**Figure 1F**) suggested that a bulkier side chain, such as that of a tri- or di-methylated lysine or an arginine may not be accommodated in this interface. In EMSAs, the affinity of PRC2 for binding to mononucleosomes containing a trimethyllysine analog at H3K36 (H3Kc36me3) was indistinguishable from that for binding to unmodified mononucleosomes (**Figure 3A, B**). However, as previously reported (Schmitges et al., 2011; Yuan et al., 2011), H3K27 mono- and trimethylation by PRC2 was strongly inhibited on H3Kc36me3-containing mononucleosomes (**Figure 3C**, compare lanes 5-7 with 2-4, see also **Supplementary Figure 6A**). Methylation of H3K27 was also inhibited on mononucleosomes where H3K36 had been mutated to arginine or alanine (H3^{K36R} and H3^{K36A} mononucleosomes, respectively) (**Figure 3C**, compare lanes 8-13 with 2-4). PRC2 inhibition on H3^{K36R} and H3^{K36A} mononucleosomes was, however, less severe than on H3Kc36me3 mononucleosomes (**Figure 3C**, compare lanes 8-13 with 5-7). We note that the quantitative analyses here show inhibition of PRC2 HMTase activity on H3^{K36A} mononucleosomes, consistent with earlier studies (Jani et al., 2019), whereas other studies previously had failed to detect inhibition on H3^{K36A} mononucleosomes (Schmitges et al., 2011). Taken together, our structural and biochemical analyses suggest that the unmodified side chain of H3K36 is critical for the productive positioning of H3K27 in the catalytic center of PRC2.

Neither the bulkier side chains of trimethyllysine or arginine nor the short apolar side chain of

alanine appear to provide the correct fit at the position of H3K36. Finally, we compared PRC2

HMTase activity on histone $H3_{18-42}$ peptides that were either unmodified or contained

H3K36me3 using a Masspectrometry-based methylation assay. Importantly, on this isolated

peptide, H3K36me3 did not inhibit H3K27 monomethylation by PRC2   (**Figure 3D,**

Supplementary Figure 6B). The allosteric inhibition of PRC2 by H3K36me3 therefore only

occurs in the context of the geometric constraints of the nucleosome.

H3K36me3 inhibits H3K27 methylation by PHF1-PRC2

DNA-binding by the winged-helix domain of PHF1 increases the binding affinity and residence

time of PHF1-PRC2 on nucleosomes about two- to three-fold, resulting in more efficient

H3K27 methylation by this complex compared to PRC2 (Choi et al., 2017). Furthermore, the

PHF1 tudor domain binds to H3K36me3 in the context of a nucleosome (Musselman et al.,

2013). To investigate whether these interactions might modulate PRC2 inhibition by

H3K36me3, we compared the HMTase activity of full-length PHF1-PRC2 (**Figure 1B**) on

unmodified and H3Kc36me3 mononucleosomes. H3K27 mono- and tri-methylation by PHF1-

PRC2 was strongly inhibited on H3Kc36me3 mononucleosomes (**Supplementary Figure 6C**).

H3K36me3 therefore inhibits H3K27 methylation by PHF1-PRC2 even though this complex

has higher binding-affinity and a prolonged residence time on nucleosomes (Choi et al., 2017).

Further analyses will be needed to assess whether and how interaction of the PHF1 tudor

domain with H3K36me3 might change H3K27 methylation by PHF1-PRC2 on more complex

oligonucleosome substrates containing H3K36me3- and unmodified nucleosomes.

***Drosophila* with H3^{K36R} or H3^{K36A} mutant chromatin arrest development at different**

stages

The observation that PRC2 is not only inhibited on H3K36me2/3-modified nucleosomes but

also on H3^{K36R} and on H3^{K36A} mutant nucleosomes prompted us to investigate how H3K27

trimethylation is affected in *Drosophila* with H3^{K36R} or H3^{K36A} mutant chromatin. H3K27me3

is primarily found on canonical histone H3 (McKay et al., 2015; Pengelly et al., 2013). We used the following strategy to replace the canonical histone H3 gene copies encoded in the *HisC* gene cluster with H3[K36R] or H3[K36A] mutant versions. Animals that are homozygous for a deletion of the *HisC* gene cluster (i.e. *Df(2L)HisC* homozygotes) arrest development at the blastoderm stage after exhaustion of the pool of maternally-deposited histones but transgene cassettes providing 12 copies of the wild-type histone gene unit (*12xHisGU^WT*) rescue *Df(2L)HisC* homozygotes into viable adults (Günesdogan et al., 2010; McKay et al., 2015). We therefore generated *Df(2L)HisC* homozygotes carrying *12xHisGU^H3K36R* or *12xHisGU^H3K36A* transgene cassettes and shall refer to these animals as *H3^K36R* and *H3^K36A* mutants, respectively. The *Drosophila* strains to generate *H3^K36R* mutant animals had been described previously (McKay et al., 2015). The strain for generating *H3^K36A* mutants was constructed in this study.

H3^K36R mutant animals complete embryogenesis and their cuticle morphology is indistinguishable from wildtype (**Supplementary Figure 7**). In agreement with the results from Matera and colleagues (McKay et al., 2015), we found that these animals arrest development during the larval or pupal stages. Specifically, 81% of *H3^K36R* mutant animals arrested development at variable time points during larval growth, 18% develop to form pupae that die prior to metamorphosis, and 1% develop into late pupae that complete metamorphosis but then arrest as pharate adults (**Supplementary Figure 7**). Like Matera and colleagues (McKay et al., 2015), we have not observed any *H3^K36R* mutants that eclose from the pupal case, and both our studies therefore disagree with a report from the Schwartz lab who claimed that *H3^K36R* mutants would be able to develop into adults (Dorafshan et al., 2019). When we dissected the rare *H3^K36R* mutant pharate adults from their pupal cases and examined their epidermal structures, we found that they consistently showed homeotic transformations reminiscent of Polycomb group (PcG) mutants. These PcG mutant phenotypes included antenna-to-leg transformations and extra sex comb teeth on meso- and metathoracic legs in males (**Supplementary Figure 7**). A molecular analysis of these PcG phenotypes will be presented below.

$H3^{K36A}$ mutants also complete embryogenesis and the morphology of their embryonic

cuticle also appeared indistinguishable from wildtype (**Supplementary Figure 7**). 96% of the

$H3^{K36A}$ mutant animals arrest development before hatching from the eggshell and the 4% that

hatch die during the first larval instar (**Supplementary Figure 7**). $H3^{K36A}$ mutants therefore

arrest development earlier and in a narrower time window than $H3^{K36R}$ mutants. The molecular

basis for the earlier lethality of $H3^{K36R}$ mutant animals remains to be determined. However, it

is not unusual that mutations changing the chemical properties of a particular histone lysine

residue result in phenotypes with different severity (e.g. (Copur et al., 2018))

***Drosophila* with H3^{K36R} or H3^{K36A} mutant chromatin show diminished H3K27me3 levels**

at canonical PcG target genes

We next performed Western blot analyses to examine H3K36me2, H3K36me3 and H3K27me3

bulk levels in $H3^{K36R}$ and $H3^{K36A}$ mutant animals. In the case of $H3^{K36R}$ mutants, we used

extracts from diploid imaginal disc and central nervous system (CNS) tissues dissected from

third instar larvae, and in the case of $H3^{K36A}$ mutants we used total nuclear extracts from late-

stage embryos. For the interpretation of the following experiments, it is important to keep in

mind that $H3^{K36R}$ and $H3^{K36A}$ zygotic mutant animals initially also contain a pool of maternally-

deposited wild-type canonical H3 molecules that, together with H3^{K36R} and H3^{K36A}, become

incorporated into chromatin during the pre-blastoderm cleavage cycles. Even though these

wild-type H3 molecules then become diluted during every cell cycle and are eventually fully

replaced by mutant H3, they are probably still present in the chromatin of late-stage embryos

because of the few cell divisions that take place prior to the end of embryogenesis. In diploid

tissues from larvae, replacement by mutant H3 is expected to be much more complete because

of the extensive cell proliferation that occurs in these tissues during larval growth.

In $H3^{K36R}$ mutant larvae, H3K36me2 and H3K36me3 bulk levels were reduced more

than 4-fold compared to wildtype (**Figure 4A**). The residual H3K36me2 and H3K36me3

signals (**Figure 4A**, lane 4) probably represent the methylated versions of the histone variant

H3.3 that are encoded by the genes *H3.3A* and *H3.3B* that are not located in the *HisC* locus and

had not been mutated in these animals. Intriguingly, $H3^{K36R}$ mutant animals also showed an about two-fold reduction in H3K27me3 bulk levels compared to wildtype (**Figure 4A**, compare lanes 4-6 with 1-3). The reduction of not only H3K36me2 and H3K36me3 but also of H3K27me3 bulk levels in $H3^{K36R}$ mutant larvae was previously also noted by Matera and colleages (Meers et al., 2017). H3K27 tri-methylation by PRC2 therefore appears to be compromised in *Drosophila* chromatin consisting of H3^K36R nucleosomes.

In $H3^{K36A}$ mutant embryos, H3K36me2 and H3K36me3 bulk levels were reduced about 3- to 4-fold comprared to wildtype (**Figure 4B**, compare lanes 5-8 with 1-4). Part of the residual H3K36me2 and H3K36me3 methylation signals in $H3^{K36A}$ mutant embryos probably represents the methylated versions of the histone variant H3.3. However, considering that the reduction is less pronounced than in $H3^{K36R}$ mutant larvae, at least some of the H3K36me2 and H3K36me3 signal might also represent maternally-deposited wild-type canonical H3. H3K27me3 bulk levels appeared largely unchanged compared to wildtype (**Figure 4B**, compare lanes 5-8 with 1-4).

We next performed ChIP-seq experiments to examine how the genome-wide profiles of H3K36me2 and H3K27me3 are changed in $H3^{K36R}$ and $H3^{K36A}$ mutants. In the case of $H3^{K36R}$ mutants, we compared these profiles in cells from imaginal disc and CNS tissues dissected from late-stage third instar $H3^{K36R}$ and wildtype larvae, and in the case of $H3^{K36A}$ mutants we compared the profiles in late-stage $H3^{K36A}$ and wildtype embryos. As expected from the Western blot analyses (**Figure 4A, B**), H3K36me2 levels across the genome were strongly diminished, both in $H3^{K36R}$ mutant larvae and in $H3^{K36A}$ mutant embryos (**Figure 4C-F, Supplementary Figure 8, Table S2**). The H3K27me3 profiles confirmed that the levels of this modification were reduced in $H3^{K36R}$ mutant larvae (**Figure 4C**). While the average reduction was only about two-fold, H3K27me3 levels were particularly strongly diminished at canonical PRC2 target genes such as the HOX genes that in wildtype animals are decorated with high-levels of H3K27me3 (**Figure 4C, E, Supplementary Figure 8, Table S2**). Specifically, at the HOX genes *Ultrabithorax* (*Ubx*), *abdominal-A* (*abd-A*), *Abdominal-B* (*Abd-B*) or *Antennapedia* (*Antp*), H3K27me3 levels in $H3^{K36R}$ mutants were between 3- and 4-fold

lower than in wildtype (**Figure 4C, E**). As expected from the Western blot analyses (**Figure**

4B), $H3^{K36A}$ mutant embryos did not show a general reduction in the H3K27me3 profile (**Figure**

4D). However, H3K27me3 levels were 1,5-fold reduced across the HOX genes (**Figure 4D,**

F). In *Drosophila* with $H3^{K36R}$ or $H3^{K36A}$ chromatin, PRC2 therefore appears to be unable to

generate high levels of H3K27me3 at Polycomb target genes.

Polycomb repression of HOX genes is impaired in *Drosophila* with H3^{K36R} or H3^{K36A}

mutant chromatin

The PcG-like phenotypes in the rare $H3^{K36R}$ mutant animals that survive into pharate adults and

the reduction of H3K27me3 levels at HOX genes in these mutants prompted us to analyse

whether and how expression of these genes is altered in $H3^{K36R}$ and $H3^{K36A}$ mutants. In a first

set of experiments, we analysed HOX gene expression in embryos. Both mutants showed

stochastic misexpression of *Abd-B* in single cells in late stage embryos (**Figure 5A**). *Abd-B*

misexpression in $H3^{K36R}$ and $H3^{K36A}$ mutant embryos was however clearly less widespread than

in $H3^{K27R}$ mutant embryos or in embryos lacking the PRC2 subunit Esc that are shown for

comparison (**Figure 5A**). Moreover, we were unable to detect misexpression of *Antp* or *Ubx*

in $H3^{K36R}$ or $H3^{K36A}$ mutant embryos.

We next analysed HOX gene expression in imaginal discs and CNS tissues from third

instar $H3^{K36R}$ mutant larvae. In the CNS of every single mutant individual, *Ubx* was widely

misexpressed in many single cells in an apparently stochastic pattern (**Figure 5B**). 50% of the

$H3^{K36R}$ mutant larvae also showed stochastic misexpression of *Ubx* in individual cells in the

wing pouch of the wing imaginal disc (**Figure 5C**). *Ubx* misexpression in $H3^{K36R}$ mutant wing

discs was less widespread than in $H3^{K27R}$ mutants that are shown for comparison (**Figure 5C**).

Finally, we found that 100% of the $H3^{K36R}$ mutant larvae showed misexpression of *Antp* in the

antenna primordium of the eye-anntennal disc (**Figure 5D**). We also observed this

misexpression in clones of $H3^{K36A}$ homozygous cells that we had induced in $H3^{K36A}$

heterozygous animals (**Figure 5D**) and in $H3^{K27R}$ mutant clones that were induced as control

(**Figure 5D**). *Drosophila* with chromatin consisting of $H3^{K36R}$ or $H3^{K36A}$ nucleosomes therefore

show stochastic misexpression of multiple HOX genes. The most straightforward interpretation

of this misexpression is that it is caused by defective Polycomb repression as a result of the

reduced H3K27me3 levels in HOX gene chromatin.

DISCUSSION

Understanding how PRC2 binds chromatin and how it is regulated is essential for understanding

how the complex marks genes for Polycomb repression to maintain cell fate decisions. The

work in this study leads to the following main conclusions. First, the structure of nucleosome-

bound PHF1-PRC2 allowed to visualize how interaction of the catalytic lobe of the complex

with the substrate nucleosome threads the histone H3 N-terminus into the active site of EZH2

through a relay of contacts. Second, structure-guided mutational analyses showed that DNA-

binding by the EZH2$_{CXC}$ domain is critical for productive PRC2-nuclesome interactions. Third,

unmodified H3K36 is accommodated in a key position in the EZH2$_{CXC}$-DNA interface and

while H3K36 provides the correct fit, the methylated forms H3K36me2/3, or mutated H3K36R

or H3K36A do not seem to fit because they strongly diminish H3K27 methylation. Fourth,

H3K36 is also critical for normal H3K27 methylation *in vivo* because *Drosophila* with H3^{K36R}

or H3^{K36A} mutant chromatin show reduced levels of H3K27me3 and fail to fully maintain

Polycomb repression at HOX target genes. In the following, we shall discuss key aspects of

these new findings in the context of our previous knowledge of PRC2 regulation and function.

Different forms of PRC2 use the same molecular interactions for binding the H3 N-

terminus on substrate nucleosomes

Unlike many other histone-modifying enzymes (e.g. (McGinty et al., 2014; Worden et al.,

2019)), PRC2 does not recognize the nucleosome by docking on its acidic patch (Luger et al.,

1997) to engage with the histone substrate. Instead, the complex interacts with chromatin by

binding to the DNA gyres on the nucleosome ((Poepsel et al., 2018), this study). Prevous

studies that had measured the binding affinity and residence time of PRC2 on nucleosomes and

free DNA had found that DNA-binding makes the largest make contribution to the chromatin-

binding affinity of PRC2 (Choi et al., 2017; Wang et al., 2017). The mutational analyses here

establish that interaction of highly conserved residues in the EZH2$_{CXC}$ domain with the DNA

on the substrate nucleosome is critical for H3K27 methylation (**Figure 2C**). Moreover, this

interaction sets the register for a network of interactions of the H3 N-terminus with the EZH2

surface that permits H3K27 to reach into the active site (**Figure 1 D, F**). Consistent with our

findings here, an independent recent study of a cryo-EM structure of PRC2 with co-factors

JARID2 and AEBP2 bound to a mononucleosome with monoubiquitylated H2A (Kasinath et

al., 2020) identified very similar interactions of EZH2 with the nucleosomal DNA and the H3

N-terminus. Different forms of PRC2 that contain different accessory proteins and dock in

different ways on chromatin therefore contact the substrate H3 N-terminus in the nucleosome

through similar interactions.

The position of H3K36 in the EZH2$_{CXC}$-nucleosome interface enables allosteric regulation

by H3K36 methylation

Important novel insight from our structure came from the observation that unmodified H3K36

is located in a critical position in the EZH2$_{CXC}$-DNA interface. Unmodified H3K36 has the

right fit for interaction of the H3 N-terminus with the EZH2 surface and placement of H3K27

in the active site. The inhibition of H3K27 mono-, di- and tri-methylation on nucleosomes

carrying H3K36me2 or -me3 (Schmitges et al., 2011; Yuan et al., 2011) or on H3^{K36R} or H3^{K36A}

nucleosomes (**Figure 3C**) suggests that these alterations of the H3K36 side chain impair the

interaction of H3K27 with the active site of EZH2. On isolated H3 N-terminal peptides,

H3K36me3 did not inhibit the formation of H3K27me1 (**Figure 3D**), consistent with earlier

findings that on peptide substrates H3K36me3 only has a minor effect on the kcat of H3K27

methylation (Schmitges et al., 2011) (Jani et al., 2019). Also, H3K36me3 does not diminish

the affinity of PRC2 for binding to mononucleosomes (**Figure 3A, B**) and does not reduce the

residence time of PRC2 on nucleosome arrays (Guidotti et al., 2019). Taken together, a possible

scenario would therefore be that within the time frame of the PRC2 nucleosome binding and

reaction cycle, docking of the H3K36 side chain in the $EZH2_{CXC}$-DNA interface is critical for

rapid alignment of the H3 N-terminus on the EZH2 surface into a catalytically competent state.

According to this view, H3K36me2/3 does not locally disrupt nucleosome binding but

allosterically inhibits H3K27 from interacting with the EZH2 active site.

H3K27 methylation and Polycomb repression are defective in Drosophila with H3^{K36R} or

H3^{K36A} chromatin

The finding that PRC2 is inhibited on H3^{K36R} and H3^{K36A} nucleosomes *in vitro* had prompted

us to use a genetic histone replacement strategy in *Drosophila* (Günesdogan et al., 2010)

(McKay et al., 2015) to assess PRC2 inhibition on H3^{K36R} or H3^{K36A} chromatin *in vivo*. Previous

studies had found that *Drosophila H3^{K36R}* mutants are able to develop into the pupal stages and,

consistent with this late develpomental arrest, whole third instar larvae were found to show only

relatively minor changes in their transriptome compared to wildtype animals (McKay et al.,

2015; Meers et al., 2017). Here, we found that a few rare *H3^{K36R}* mutant animals even survive

into pharate adults and that these show remarkably little morphological defects apart from

homeotic transformations characteristic of Polycomb mutants (**Figure S7**). We show that these

phenotypes are caused by defective Polycomb repression of multiple HOX genes (**Figure 5**)

and that they are linked to reduced levels of H3K27me3 at these genes (**Figure 4**). A simple

straightforward explanation for these phenotypes in *H3^{K36R}* or *H3^{K36A}* mutant animals is that

PRC2 is unable to effectively deposit high levels of H3K27me3 on the H3^{K36R} or H3^{K36A}

nucleosomes, respectively, in their chromatin. Accordingly, H3K27me3 levels at HOX genes

are below the threshold needed to reliably maintain Polycomb repression and consequently,

HOX genes become stochastically misexpressed in a fraction of cells. Finally, we note that in

H3^{K36R} mutant larvae, the experimental setting where we have been able to generate the most

complete replacement of H3 by H3^{K36R}, H3K27me3 levels at HOX genes were only about 3- to

4-fold reduced compared to wildtype (**Figure 4C**). However, as shown in **Figure 3C**, on

nucleosomes *in vitro*, H3K36me3 inhibited PRC2 more effectively than H3^{K36R} or H3^{K36A}. It

therefore seems likely that in contrast to the *H3^{K36R}* and *H3^{K36A}* mutants that we have used as

proxy, H3K36me2 and H3K36me3 *in vivo* also inhibit PRC2 more effectively from depositing

H3K27me3 on H3K36me2- or H3K36me3-modified nucleosomes in transcriptionally active

chromatin.

Concluding remark

The structural, biochemical and genetic work reported in this study shows that it is the exquisite

geometry formed by a relay of interactions between the PRC2 enzyme, nucleosomal DNA and

the H3 N-terminus that enable the histone methylation marks H3K36me2 and H3K36me3 in

transcriptionally active chromatin to allosterically prevent PRC2 from depositing the repressive

histone methylation mark H3K27me3 at transcribed genes.

MATERIALS AND METHODS
428
Protein expression and purification
Human PHF1-PRC2 wild-type (wt) complex was expressed and purified as previously
described (Choi et al., 2017). In brief, an optimized ratio of the baculoviruses for the different
PHF1-PRC2 subunits was used to infect HiFive cells (Invitrogen). Cell were lysed using a glass
Dounce homogenizer and the complex was purified using affinity chromatography (Ni-NTA
and Strep-tag), followed by simultaneous TEV mediated protease tag cleavage and Lambda
Phosphatase treatment (obtained from the MPI of Biochemistry Protein Core facility) and a
final size exclusion chromatography (SEC) step in a buffer containing 25 mM Hepes, pH 7.8,
150 mM NaCl, 10% glycerol, 2 mM DTT.
$PRC2^{CXC>A}$, $PRC2^{EED>A}$ and $PRC2^{CXC>A/EED>A}$ mutants were generated by PCR with primers
containing the desired mutations, subsequent ligation and transformation. Expression and
purification were performed as above.
Xenopus laevis (*X.l.*) and *Drosophila melanogaster* (*D.m.*) histones were expressed and purified
from inclusion bodies as described in (Luger et al., 1999). To mimic the inhibitory mark
H3K36me3 or the allosteric activating mark H3K27me3, the cysteine side chain of a mutated
D.m. histone $H3^{C110A\ K36C}$ or *X.l.* histone $H3^{C110A\ K27C}$ was alkylated with (2-bromoethyl)
trimethylammonium bromide (Sigma Aldrich) as described previously (Simon et al., 2007).
Nucleosomes containing these modifications are abbreviated with e.g. H3Kc36me3.
447
448 For histone octamers, equimolar amounts of histones H2A, H2B, H4 and H3 (wt, $H3^{K36A}$,
449 $H3^{K36R}$, H3Kc27me3 or H3Kc36me3) were mixed and assembled into octamers in high salt
450 buffer containing 10 mM Tris-HCL pH 7.5, 2 M NaCl, 1 mM EDTA, 5 mM β-mercaptoethanol.
451 Subsequent SEC was performed to separate octamers from H3/H4 tetramers or H2A/H2B
452 dimers (Luger et al., 1999).
453
454 Reconstitution of nucleosomes:
455 For *X.l.* and *D.m* mononuclesomes used in biochemical assays, 6-carboxyfluorescein (6-FAM)-
456 labeled 215 bp 601 DNA (Lowary and Widom, 1998) was PCR amplified from the p601

plasmid, purified on a MonoQ column (GE Healthcare), precipitated with ethanol and dissolved in the same high salt buffer used for octamers. Optimized ratios of octamer to DNA (usually ranging between 0.8-1.3 : 1) were mixed and nucleosomes were reconstituted by gradient and stepwise dialysis against low salt buffers to a final buffer containing 25 mM Hepes, pH 7.8, 60 mM NaCl, 2 mM DTT.

X.l. asymmetrical dinucleosomes for cryo-EM studies containing one unmodified substrate nucleosome and one H3K27me3-modified (allosteric) nucleosome connected with a 35 bp linker DNA were reconstituted using the protocol described in (Poepsel et al., 2018). In brief, substrate nucleosomes and allosteric nucleosomes were separately assembled on the respective *DraIII* digested nucleosomal DNA. The latter was generated by PCR with primers introducing the desired linker and *DraIII* recognition sites and purified as described above. The assembled nucleosomes were purified on a preparative native gel system (Biorad 491 prep cell). After ligation using T4 ligase (Thermo Fisher Scientific) the resulting dinucleosomes were purified from aberrant or non-ligated mononucleosomes by a second preparative native gel system (Biorad 491 prep cell). In contrast to (Poepsel et al., 2018), the dinucleosome DNA used in this study contained an additional 30 bp overhang on the substrate nucleosome, thus resulting in the following DNA sequence:

5′–601 binding (allosteric nucleosome) – agcgatctCACCCCGTGatgctcgatactgtcata – 601 binding (substrate nucleosome) – atgcatgcatatcattcgatctgagctcca –3′ (after DraIII digestion, assembly of substrate/allosteric nucleosome and ligation to dinucleosomes).

X.l. symmetrical unmodified dinucleosomes used for the HMTase assays with the PRC2^{CXC} mutants were obtained by reconstituting octamers with a 377 bp DNA containing two 601 sequences connected by a 35 bp linker DNA. A vector containing the 377 bp sequence was ordered from Invitrogen GeneArt and was used for PCR resulting in:

5′–atatctcgggcttatgtgatggac – 601 binding (substrate nucleosome 1) – agcgatctcaacgagtgatgctcgatactgtcata – 601 binding (substrate nucleosome 2) – gtattgaacagcgactcgggatat–3′.

The PCR products were purified as described above. Optimized ratios of octamer : DNA (usually ranging between 1.8-2.3 : 1) were mixed and nucleosomes were reconstituted by gradient and stepwise dialysis against low salt buffers to a final buffer containing 25 mM Hepes, pH 7.8, 60 mM NaCl, 2 mM DTT.

Cryo-EM Data acquisition
Complexes of PHF1-PRC2 and asymmetrically modified 35 bp dinucleosomes were assembled and grids were prepared as described previously, with the difference of using 0.005% NP40 instead of 0.01% (Poepsel et al., 2018). Cryo-EM data were collected on an FEI Titan Krios microscope operated at 300 kV and equipped with a post-column GIF and a K2 Summit direct detector (Gatan) operated in counting mode. A total of 3467 movies were collected at a nominal magnification of 81,000x (1.746 Å/pixel) at the specimen level using a total exposure of 53 e$^-$ / Å^2 distributed over 60 frames and a target defocus range from 1.5–3 μm. Data acquisition was carried out with SerialEM.

Cryo-EM Data processing
Movies were aligned and corrected for beam-induced motion as well as dose compensated using MotionCor2 (Zheng et al., 2017). CTF estimation of the summed micrographs was performed with Gctf (Zhang, 2016) and particles were picked in Gautomatch (http://www.mrc-lmb.cam.ac.uk/kzhang/ K. Zhang, MRC LMB, Cambridge, UK) using templates created from the AEBP2-PRC2-dinucleosome cryo-EM structure (EMD-7306, (Poepsel et al., 2018). All subsequent image processing steps were performed in Relion 3.0 (Zivanov et al., 2018) as

shown in Fig. S2. A total of 1,028,229 candidate particles were subjected to two rounds of
initial 3D classification against a reference map (AEBP2-PRC2-dinucleosome low-pass filtered
to 60 Å) and the Bayesian fudge factor (T value) set to 8. 330,482 remaining particles were
subjected to two more rounds of 3D classification, this time using the best 3D model from the
previous run as reference. Finally, the two best 3D models were 3D refined and further
classified into 10 classes without translational and rotational sampling, using a T value of 4.
From this run, the best 3D classes with the highest nominal overall resolution and rotational
and translational accuracies were subjected to iterative rounds of 3D refinement, this time
applying a soft mask for solvent flattening, per particle CTF refinement and Bayesian polishing.
The highest nominal resolution was only achieved by combining several models from the
previous 3D run, likely due to missing particle views in one or the other individual model. The
final map after postprocessing had an overall nominal resolution of 5.2 Å, as determined from
the gold-standard FSC criterion of 0.143 (Rosenthal and Henderson, 2003) (Fig. S1D). The
density (Overall PHF1-PRC2:di-Nuc) with fitted models is shown in Fig.1A and in Fig. S1E
using UCSF ChimeraX (Goddard et al., 2018). Local resolution estimation was performed in
Relion 3.0 and is shown in Fig. S1B. The spherical angular distribution of all particles in the
final model is shown in Fig. S1C.
To further improve the resolution and map details of the region around the H3 N-terminus,
particle subtraction and focused 3D refinement was applied (Bai et al., 2015; Ilca et al., 2015;
Zhou et al., 2015). Using a mask generated with UCSF Chimera (Pettersen et al., 2004) and
Relion 3.0 the signal of the allosteric nucleosome as well as parts of PRC2 (EED and $EZH2_{allo}$)
was subtracted from all particle images. These signal subtracted particles were then subjected
to focused 3D refinement using a soft mask around the substrate nucleosome and $EZH2_{sub}$. This
yielded a 4.4 Å map ($EZH2_{sub}$-Nuc_{sub}) (Fig. S3B). Local resolution estimation is shown in
Fig.S3A. For model building and depiction, the final density was further sharpened (applied b
– factor: - 66) using the Multisharpen function in Coot (Emsley et al., 2010) (e.g. in Figs. 1E,
S3D, E and F).
To confirm the side chain information visible in the Coot sharpened map, Phenix Resolve
density modification was run on the two half maps generated from the 3D refinement of the
$EZH2_{sub}$-Nuc_{sub} map (Terwilliger et al., 2019). The resolution of the map according to Phenix
cryo EM density modification output improved to 4 Å and the resulting map was used as an
additional guideline for model building as well as for depiction (in Figs. S4 A-D.).
543
Cryo-EM data fitting, modeling and refinement
Available crystal structures were fitted into the final maps using rigid-body fitting in UCSF
Chimera and all manual remodeling, morphing and building was performed in Coot. For PRC2,
the crystal structure of the catalytic lobe of human PRC2 (PDB: 5HYN (Justin et al., 2016))
was used. Since the SBD helix and the SANT1 helix bundle of the crystal structure was not
accommodated well by the corresponding EM density, this region was fitted separately. A
model of a dinucleosome with linker DNA (Supplementary dataset 1 in (Poepsel et al., 2018),
including crystal structures of nucleosomes, PDB 3LZ1, also PDB 1AOI, also PDB 6T9L) was
fitted.
553
The above described overall model was then used as a starting model for fitting and building
$EZH2_{sub}$-Nuc_{sub} into the focused map. Where possible, missing parts in the model were built de-
novo, i.e. the H3 N-terminal tail (residues 30-37) between the catalytic site of PRC2 and the
substrate histone. Available information from crystal structures was used as a guide (PRC2 with
H3 peptide bound: PDB: 5HYN (Justin et al., 2016), and high resolution crystal structures of
nucleosomes (PDB 1AOI and PDB 6T9L) (Luger et al., 1997). Parts of $EZH2_{sub}$-Nuc_{sub} model
were then fitted using the morph fit routine in Coot or manually (Casañal et al., 2020).
Secondary structure restraints for real-space refinement were generated automatically with
phenix.secondary_structure_restraints (Sobolev et al., 2015) and manually curated. Hydrogens
were added and the model was real-space refined with a resolution- cutoff of 4.4 Å with Phenix
(Afonine et al., 2018) (phenix-1.18rc1-3777), using reference structures (PDB 6T9L and PDB

1AOI for nucleosome and one copy of the human PRC2 crystal structure generated from PDB
5HYN), applying strict secondary structure and Ramachandran restraints.
Our final model includes the modelled side chains of the fitted crystal/cryo-EM structures. This
is in our opinion supported by the data as the substrate nucleosome protein core is resolved to
app. 4 Å (Fig. S3A) and the map in these regions shows clear bulky side chain information (Fig
S3D). The EZH2 density is of worse quality however even at lower resolution side chains likely
contribute to the signal in the particle images and thereby an overall good model to map fit (in
our case given by the high CC values as well as $FSC_{modelvsmap}$) is arguably only ensured in the
presence of side chains. However we caution readers against in interpreting our model at side
chain resolution in poorly resolved regions.
Structures were visualized with UCSF ChimeraX (Goddard et al., 2018) and PyMOL2
(https://pymol.org/2/).
577
<u>Electrophoretic mobility shift assay (EMSA)</u>
EMSAs on a 1.2% agarose gel in 0.4x TBE Buffer with 45 nM 6-FAM - labeled
mononucleosomes (unmodified wt *X.l.* for bandshifts with the PRC2^{CXC} mutants, unmodified
wt *D.m.* and *D.m* H3Kc36me3 trimethyllysine analog containing nucleosomes) and increasing
PRC2 concentrations (concentrations indicated in the figures above the gels) were performed
in triplicates as described in (Choi et al., 2017). A Typhoon FLA 9500 scanner and the Fiji
software was used for densitometric analysis of the 6-FAM signal (Schindelin et al., 2012).
Background correction and calculation of the fractions of bound nucleosomes was performed
with R using tidyverse (https://www.r-project.org/). In detail: two parts were boxed out in each
lane: 1. unbound nucleosomes ('unbound' box) and 2. shifted nucleosomes ('bound',
everything above 'unbound'). The boxed-out signals were integrated and background corrected
by subtracting the respective control ('bound' background of lane 1 for 'bound' boxes and
'unbound' background of lane 10 for 'unbound' boxes). To calculate the fraction of bound vs.
unbound nucleosomes, the value for 'bound' nucleosome in each lane was divided by the total
signal (sum of bound and unbound) of the same lane. Hill function fitting and illustration of
the plot were subsequently performed with Prism 8 (GraphPad).
594
<u>Histonemethyltransferase (HMTase) assay</u>
For all HMTase assays, 446 nM of mononucleosomes or 223 nM of dinucleosomes were
incubated with indicated amounts of the different PRC2 complexes, in a reaction buffer
containing 20 mM HEPES pH 7.8, 50 mM NaCl, 2.5 mM $MgCl_2$, 5% glycerol, 0.25 mM EDTA,
0.5 mM DTT and 80 μM S-adenosylmethionine (SAM). Reactions were allowed to proceed for
90 min at RT before quenching by the addition of 1x (final concentration) SDS loading buffer
and heat inactivation at 95 °C for 5 min. Proteins were separated by electrophoresis on a 16%
(w/v) SDS gel, transferred to a nitrocellulose membrane and probed with antibodies against
H3K27me3 (Millipore, 07-449), H3K27me1 (Millipore, 07-448) and H4 (Abcam, ab10158).
For quantification, HMTase reactions and the corresponding western blots on *D.m.* unmodified,
H3Kc36me3, H3$^{K36A/R}$ mononucleosomes were performed in triplicates and subjected to
densitometric analysis (Chemiluminescence signal, ImageQuant LAS 4000). The integrated
densitometric signal (band) in each lane was background corrected against the control lane (lane
1, no PRC2 in the reaction) and normalized with respect to the lane containing the highest
amount (i.e. 100%) of PRC2 on unmodified nucleosomes (lane 4). The relative amounts of
trimethylation/monomethylation for all other lanes were calculated with respect to lane 4.
Graphical representations were made with Prism 8 (GraphPad).
612
<u>Mass Spectrometry (MS)</u>
500 nM of PRC2 were incubated with 2 μM of either unmodified or H3$_{18-42}$ peptide containing
the K36me3 modification in HMTase reaction buffer (described above) and methyltransferase
activity was allowed to proceed over night at RT. Reactions were then quenched with 1%
trifluoroacetic acid (TFA). Home-made stage tips with poly(styrenedivinylbenzene) copolymer
(SDB-XC) were used to remove PRC2 from the reactions (Rappsilber et al., 2007). First, stage
tips were washed with methanol, followed by a second wash with buffer B (0.1% (v/v) formic

acid, 80% (v/v) acetonitrile). The SDB-XC material was then equilibrated with buffer A (0.1%
(v/v) formic acid) and 40 µl of sample was applied and washed several times. Finally, samples
were eluted using buffer B and introduced into the Bruker maXis II ETD mass spectrometer by
flow injection of 20 µl sample using an Agilent HPLC at a flow rate of 250 µl/min and 0.05%
TFA in 70% acetonitril:H2O as solvent for ESI-MS time-of-flight analysis.
Peptides were ionized at a capillary voltage of 4500 V and an end plate offset of 500 V.
Full scan MS spectra (200-1600 m/z) were acquired at a spectra rate of 1 Hz and a collision cell
energy of 15 eV.
Raw data files were processed using Bruker Compass DataAnalysis. The m/z spectra were
deconvoluted (maximum entropy method) with an instrument resolving power of 10,000 and
the resulting neutral spectra peaks were integrated. For quantification, the experiment was
performed in triplicates. The sum of the monomethylation peak areas was divided by the sum
of the first 4 peaks of the input peptide together with the sum of the monomethylation peak
areas. Illustration of the quantification was subsequently performed with Prism 8 (GraphPad).
A Welch's t-test was calculated to show the nonsignificant difference between the activity of
PRC2 on unmodified or H3K36me3 peptide.
636
<u>Construction of histone transgenes to generate $H3^{K36A}$ and $H3^{K36R}$ strains</u>
Site directed mutagenesis on *pENTR221-HisGU.WT, pENTRL4R1-HisGU.WT* and
pENTRR2L3-HisGU.WT (Günesdogan et al., 2010) was used to mutate histone H3K36 to
alanine or arginine. The final constructs *pfC31-attB-3xHisGU.H3K36A* and *pfC31-attB-*
3xHisGU.H3K36R were generated by Gateway LR recombination of above vectors and
integrated at attP sites VK33 (BDSC 9750) and 86Fb (BDSC 130437). The full genotypes of
animals used in the study is described below.
644
645 <u>*Drosophila* strains and genotypes</u>
646 The following strains were used in this study:
647
648 *Oregon-R*
649
650 *w; Df(2L)His^C FRT40A/ Df(2L)His^C FRT40A; 12xHisGU^wt/ 12xHisGU^wt* (McKay et al., 2015)
651
652 *w; Df(2L)His^C FRT40A/ CyO ubi-GFP; 12xHisGU^H3K36R/TM6B* (McKay et al., 2015)
653
654 *w; Df(2L)His^C FRT40A/ CyO twi:Gal4 UAS:GFP; 3xHisGU^H3K36A(VK33) 3xHisGU^H3K36A(86Fb)/ 3xHisGU^H3K36A(VK33) 3xHisGU^H3K36A(86Fb)*
655 (generated in this study)
656
657 *w; Df(2L)His^C FRT40A/ CyO ubi:GFP; 3xHisGU^H3K27R(68E) 3xHisGU^H3K27R (86Fb)/ 3xHisGU^H3K27R (68E) 3xHisGU^H3K27R (86Fb)* (Pengelly et al.,
658 2013)
659
660 *w hs-flp; w; hs-nGFP FRT40A/ hs-nGFP FRT40; 3xHisGU^H3K27R(68E) 3xHisGU^H3K27R(86Fb)/ 3xHisGU^H3K27R(68E) 3xHisGU^H3K27R (86Fb)* (Pengelly
661 et al., 2013)
662
663 *w hs-flp; M(2)25A ubi-GFP FRT40A/CyO*
664
665 *yw; esc^6 b pr / CyO, P[esc^+]*
666
667 *In(2LR) Gla / CyO, esc^2*
668
669
670 The following genotypes were used for the experiments shown in:
671
672 **Figure 4A, C, E**
673 *wt:*   *Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^wt(VK33)/ 12xHisGU^wt(VK33)*
674 *H3^K36R:*   *Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^H3K36R(VK33)/TM6B*
675
676 **Figure 4B, D, F**
677 *wt:*   *Oregon-R*
678 *H3^K36A:*   *w; Df(2L)His^C FRT40A/ Df(2L)His^C FRT40A; 3xHisGU^H3K36A(VK33) 3xHisGU^H3K36A(86Fb)/ 3xHisGU^H3K36A(VK33) 3xHisGU^H3K36A(86Fb)*
679
Figure 5A
wt: *Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^wt(VK33)/ 12xHisGU^wt(VK33)*

$H3^{K36R}$: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{H3K36R}(VK33)/TM6B
$H3^{K36A}$: w; Df(2L)His^C FRT40A/ Df(2L)His^C FRT40A; 3xHisGU^{H3K36A}(VK33) 3xHisGU^{H3K36A}(86Fb)/ 3xHisGU^{H3K36A}(VK33) 3xHisGU^{H3K36A}(86Fb)
$H3^{K27R}$: w; Df(2L)His^C FRT40A/ Df(2L)His^C FRT40A; 3xHisGU^{H3K27R}(68E) 3xHisGU^{H3K27R} (86Fb)/ 3xHisGU^{H3K27R} (68E) 3xHisGU^{H3K27R} (86Fb)
esc^-: esc^6 b pr / CyO, esc^2 ($esc^{mat-zyg-}$ obtained as progeny from esc^6 b pr/CyO, esc^2 parents)
686

687 **Figure 5B**
688 wt: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{wt}(VK33)/ 12xHisGU^{wt}(VK33)
689 $H3^{K36R}$: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{H3K36R}(VK33)/TM6B
690

691 **Figure 5C**
692 wt: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{wt}(VK33)/ 12xHisGU^{wt}(VK33)
693 $H3^{K36R}$: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{H3K36R}(VK33)/TM6B
694 $H3^{K27R}$: w hs-flp; Df(2L)His^C FRT40A/hs-nGFP FRT40A; 3xHisGU^{H3K27R}(68E)3xHisGU^{H3K27R}(86Fb)/3xHisGU^{H3K27R}(68E)3xHisGU^{H3K27R}(86Fb)
695

696 **Figure 5D**
697 wt: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{wt}(VK33)/ 12xHisGU^{wt}(VK33)
698 $H3^{K36R}$: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{H3K36R}(VK33)/ TM6B
699 $H3^{K36A}$: w hs-flp; Df(2L)HisC FRT40A/ M(2)25AubiGFP FRT40; 3xHisGU^{H3K36A}(VK33) 3xHisGU^{H3K36R}(86Fb)/ +
700 $H3^{K27R}$: w hs-flp; Df(2L)HisC FRT40A/ M(2)25A ubi-GFP FRT40; 3xHisGU^{H3K27R}(68E) 3xHisGU^{H3K27R} (86Fb)/ +
701

702 **Figure S7**
703 wt: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{wt}(VK33)/ 12xHisGU^{wt}(VK33)
704 $H3^{K36R}$: Df(2L) HisC FRT40/ Df(2L) HisC FRT40; 12xHisGU^{H3K36R}(VK33)/TM6B
705 $H3^{K36A}$: Df(2L)His^C FRT40A/ Df(2L)His^C FRT40A; 3xHisGU^{H3K36A}(VK33) 3xHisGU^{H3K36A}(86Fb)/ 3xHisGU^{H3K36A}(VK33) 3xHisGU^{H3K36A}(86Fb)
706

<u>Immunohistochemistry and immunofluorescence stainings</u>
Embryos of the appropriate genotypes listed above were identified by the lack of GFP marked
balancer chromosomes, fixed and stained with Abd-B antibody, following standard protocols.
Imaginal discs from third instar larvae were stained with Antp and Cy3-labeled secondary
antibodies following standard protocols. For clonal analysis (**Fig. 3D**), clones were induced 96
hrs before analyses by heat-shocked induced expression of Flp recombinase in the genotypes
listed above.
715

<u>ChIP-seq analysis in *Drosophila* embryos and in larval tissues</u>
Embryo collection, chromatin preparation and ChIP: 21-24 hr old *wt, H3^{K36A}* embryos (see
above for details of genotypes) were dechorionated, quick-frozen in liquid N2 and stored at -
80˚C. 5 µL of thawed embryos were homogenized in 5 mL of fixing solution (60 mM KCl, 15
mM NaCl, 4 mM MgCl$_2$, 15 mM Hepes pH 7.6, 0.5% Triton X-100, 0.5 mM DTT, protease
inhibitors, 0.9% Formaldehyde) at r.t.. The homogenate was filtered through a strainer (Greiner
Bio-One, EASYstrainer^{TM} 100 µm, #542 000) and incubated for 10 min with frequent gentle
shaking. Cross-linking was stopped by the addition of 450 µL of 2.5 M Glycine. Fixed nuclei
were washed with 1 mL of buffer A1 (60 mM KCl, 15 mM NaCl, 4 mM MgCl$_2$, 15 mM Hepes
pH 7.6, 0.5% Triton X-100, 0.5 mM DTT, protease inhibitors), washed with 1 mL of pre-lysis
buffer (140 mM NaCl, 15 mM Hepes pH 7.6, 1 mM EDTA, 0.5 mM EGTA, 1% Triton X-100,
0.5 mM DTT, 0.1% Na Deoxycholate, protease inhibitors), resuspended in 1 mL of lysis buffer
(140 mM NaCl, 15 mM Hepes pH 7.6, 1 mM EDTA, 0.5 mM EGTA, 1% Triton X-100, 0.5
mM DTT, 0.1% Na Deoxycholate, protease inhibitors, 0.1% SDS, 0.5% N-laurylsarcosine),
incubated at least 10 min at 4˚C with shaking, and transferred into milliTUBES 1 mL AFA
Fiber (100) (Covaris, #520130) for sonication. Sonication was performed in a Covaris S220
AFA instrument using the following setup: 140W (peak incident power) / 5% (duty cycle) / 200
(cycle per burst) / 15 min. Insoluble material was removed by centrifugation in an Eppendorf
centrifuge at 14000 rpm (10 min at 4°C). Input chromatin was quantified by measuring DNA
concentration after decrosslinking using Qubit (Thermo Scientific) and 250 ng of chromatin
were used for each ChIP experiment. 250 ng of an independently prepared batch of *D.*
pseudoobscura chromatin were spiked-in in each ChIP experiment for subsequent
normalization of the ChIP-seq datasets. The rest of the ChIP protocol was performed as
described (Bonnet et al., 2019). For each condition, the ChIP experiment was performed in
duplicates from two biologically independent chromatins. ChIP on hand-dissected CNS and
imaginal disc tissues from 3rd instar *wt* or *H3^{K36R}* homozygous larvae (see above for details on

genotypes) was performed as described (Laprell et al., 2017) with the difference *D.*
pseudoobscura chromatin was spiked in at a 1:1 ratio of dm / dp chromatin.
Library preparation and sequencing: Library preparation for sequencing was performed with
TruSeq kits from Illumina. Illumina systems (NextSeq 500) were used for paired-end DNA
sequencing. All reads were aligned using STAR (Dobin et al., 2013) to the *D. melanogaster*
dm6 genome assembly (Santos et al., 2015) and to the *D. pseudoobscura* dp3 genome assembly
(Nov. 2004, FlyBase Release 1.03). Only sequences that mapped uniquely to the genome with
a maximum of two mismatches were considered for further analyses.
Identification of H3K36me2 and H3K27me3 enriched regions: The Bioconductor STAN-
package (Zacher et al., 2017) was used to define the location of H3K36me2-enriched regions.
The seven chromosome arms (X, 2L, 2R, 3L, 3R, 4 and Y) defined in the dm6 genome assembly
were segmented in 200 bp bins. STAN annotated each of these bins into 1 of 3 'genomic states'
based on the number of H3K36me2 ChIP-seq reads and the number of input reads overlapping
with each bin, in 21-24 hr wild-type embryos. These 3 'genomic states' corresponded to:
'H3K36me2 enriched' regions; 'low or no H3K36me2' regions and 'no input' regions. The
Poisson Lognormal distribution was selected and fitting of hidden Markov models was
performed with a maximum number of 100 iterations. Stretches of consecutive bins annotated
as 'H3K36me2 enriched' regions were sometimes separated by a few bins showing another
type of annotation (i.e. 'no input'). To define a relevant set of H3K36me2 enriched regions, we
considered that if stretches of consecutive bins annotated as 'H3K36me2 enriched' regions are
not separated by more than 7 Kb, they can be fused. High-level H3K27me3 domains previously
defined using the same Bioconductor STAN-package in Bonnet et al (Bonnet et al., 2019) were
used in this study.
Normalization and visualisation of H3K27me3 and H3K36me2 ChIP-Seq datasets: The
proportion of *D. pseudoobscura* reads as compared to *D. melanogaster* reads in input and in
samples was used to normalize the H3K36me2 and H3K27me3 ChIP-seq datasets from $H3^{K36A}$
and $H3^{K36R}$ mutants to the corresponding wild-type H3K36me2 and H3K27me3 ChIP-seq
datasets respectively (see Table S2). Chip-seq tracks shown in Fig. 4 show the average of the
two biological replicates that were performed for each condition. Y-axes of ChIP-seq tracks
correspond to normalized numbers of mapped reads per million reads per 200 bp bin.
Calculation of read coverage: In wild-type and $H3^{K36A \ and \ R}$ mutant conditions, H3K36me2 and
H3K27me3 ChIP-seq read coverages across gene bodies were computed on genomic intervals
starting 750 bp upstream transcription start sites and ending 750 bp downstream transcription
termination sites. Read coverage is defined as the normalized number of mapped reads per
million reads from a ChIP-seq dataset divided by the number of mapped reads per million reads
from the corresponding input dataset across a genomic region. Among the *D. melanogaster*
Refseq genes, approximately 10800 and 9200 are overlapping with H3K36me2 enriched
regions, approximately 1030 and 1030 genes are overlapping with high-level H3K27me3
domains and 5400 and 6300 are localized in other genomic regions in embryos and larvae,
respectively.
782
<u>*Drosophila* nuclear and cell extracts for western blot analysis</u>
For embryonic total nuclear extracts, nuclei from 21-24 hr old *wt*, $H3^{K36A}$ or $H3^{K36A}$ mutant
embryos were purified and quantified as described (Bonnet et al., 2019). Pellets of nuclei were
resuspended in appropriate volumes of SDS sample buffer proportional to the number of nuclei
in each pellet. Extracts were then sonicated in a Bioruptor instrument (Diagenode) (8 cycles
(30 sec ON / 30 sec OFF), high power mode), incubated at 75˚C for 5 min and insoluble material
was removed by centrifugation at 14000 rpm for 1 mn at r.t..
Total cell extracts from imaginal disc tissues were prepared by resuspending hand-dissected
disc tissues in SDS sample buffer. Extracts were then sonicated, incubated at 75˚C for 5 min
and insoluble material was removed by centrifugation.
793
794   <u>Antibodies</u>
795   For ChIP analysis:
796   Rabbit monoclonal anti-H3K27me3   Cell Signaling Technology   #9733

797	Rabbit polyclonal anti-H3K36me2	Abcam	#9049
798			
799	For Western blot analysis on embryonic and larval extracts:		
800	Rabbit monoclonal anti-H3K27me3	Cell Signaling Technology	#9733
801	Rabbit polyclonal anti- H3K27me3	Millipore	#07-449
802	Rabbit polyclonal anti-H3K27me1	Millipore	#07-448
803	Rabbit monoclonal anti-H3K36me3	Cell Signaling Technology	#4909
804	Rabbit monoclonal anti-H3K36me2	Cell Signaling Technology	#2901
805	Rabbit polyclonal anti-H2B	(against full-length recombinant D.m. H2B)	
806	Rabbit polyclonal anti-H4	Abcam	#10158
807	Rabbit polyclonal anti-Caf1	(Gambetta et al., 2009)	
808			
809	For immunohistochemistry and immunofluorescence analysis:		
810	Mouse monoclonal anti-Abd-B	DSHB (1A2E9)	
811	Mouse monoclonal anti-Antp	DSHB (8C11)	
812			

ACKNOWLEDGMENTS

We thank Eva Nogales for generous advice, sharing of expertise and for hosting K.F. for grid

preparation. We thank Tom Cech for stimulating discussions. We thank J.R. Prabu for

excellent computing support, S.Uebel, E.Weyher, R.Kim and A.Yeroslaviz of the MPIB core

facilities for excellent technical support and S.Schkoelziger and S.Schmähling for help with

some of the experiments. This work was supported by the Deutsche Forschungsgemeinschaft

(SFB1064) and the MPG. ChIP-seq data have been deposited in GEO (accession

number: GSE148254).

REFERENCES

Afonine, P.V., Poon, B.K., Read, R.J., Sobolev, O.V., Terwilliger, T.C., Urzhumtsev, A.,
Adams, P.D., 2018. Real-space refinement in PHENIX for cryo-EM and crystallography.
Acta Crystallogr D Struct Biol 74, 531–544. doi:10.1107/S2059798318006551

Bai, X.-C., Rajendra, E., Yang, G., Shi, Y., Scheres, S.H.W., 2015. Sampling the
conformational space of the catalytic subunit of human γ-secretase. Elife 4, 1485.
doi:10.7554/eLife.11182

Ballaré, C., Lange, M., Lapinaite, A., Martin, G.M., Morey, L., Pascual, G., Liefke, R.,
Simon, B., Shi, Y., Gozani, O., Carlomagno, T., Benitah, S.A., Di Croce, L., 2012. Phf19
links methylated Lys36 of histone H3 to regulation of Polycomb activity. Nat. Struct.
Mol. Biol. 19, 1257–1265. doi:10.1038/nsmb.2434

Bonnet, J., Lindeboom, R.G.H., Pokrovsky, D., Stricker, G., Çelik, M.H., Rupp, R.A.W.,
Gagneur, J., Vermeulen, M., Imhof, A., Müller, J., 2019. Quantification of Proteins and
Histone Marks in Drosophila Embryos Reveals Stoichiometric Relationships Impacting
Chromatin Regulation. Dev. Cell 51, 632–644.e6. doi:10.1016/j.devcel.2019.09.011

Cai, L., Rothbart, S.B., Lu, R., Xu, B., Chen, W.-Y., Tripathy, A., Rockowitz, S., Zheng, D.,
Patel, D.J., Allis, C.D., Strahl, B.D., Song, J., Wang, G.G., 2013. An H3K36
methylation-engaging Tudor motif of polycomb-like proteins mediates PRC2 complex
targeting. Mol. Cell 49, 571–582. doi:10.1016/j.molcel.2012.11.026

Casañal, A., Lohkamp, B., Emsley, P., 2020. Current developments in Coot for
macromolecular model building of Electron Cryo-microscopy and Crystallographic Data.
Protein Sci. 29, 1069–1078. doi:10.1002/pro.3791

Chen, S., Jiao, L., Liu, X., Yang, X., Liu, X., 2020. A Dimeric Structural Scaffold for PRC2-
PCL Targeting to CpG Island Chromatin. Mol. Cell 77, 1265–1278.e7.
doi:10.1016/j.molcel.2019.12.019

Chen, S., Jiao, L., Shubbar, M., Yang, X., Liu, X., 2018. Unique Structural Platforms of
Suz12 Dictate Distinct Classes of PRC2 for Chromatin Binding. Mol. Cell 69, 840–
852.e5. doi:10.1016/j.molcel.2018.01.039

Choi, J., Bachmann, A.L., Tauscher, K., Benda, C., Fierz, B., Müller, J., 2017. DNA binding
by PHF1 prolongs PRC2 residence time on chromatin and thereby promotes H3K27
methylation. Nat. Struct. Mol. Biol. 298, 1039. doi:10.1038/nsmb.3488

Copur, Ö., Gorchakov, A., Finkl, K., Kuroda, M.I., Müller, J., 2018. Sex-specific phenotypes
of histone H4 point mutants establish dosage compensation as the critical function of
H4K16 acetylation in Drosophila. Proc Natl Acad Sci USA 115, 13336–13341.
doi:10.1073/pnas.1817274115

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29,
15–21. doi:10.1093/bioinformatics/bts635

Dorafshan, E., Kahn, T.G., Glotov, A., Savitsky, M., Walther, M., Reuter, G., Schwartz, Y.B.,
2019. Ash1 counteracts Polycomb repression independent of histone H3 lysine 36
methylation. EMBO Rep. e46762. doi:10.15252/embr.201846762

Emsley, P., Lohkamp, B., Scott, W.G., Cowtan, K., 2010. Features and development of Coot.
Acta Crystallogr. D Biol. Crystallogr. 66, 486–501. doi:10.1107/S0907444910007493

Francis, N.J., Kingston, R.E., Woodcock, C.L., 2004. Chromatin compaction by a polycomb
group protein complex. Science 306, 1574–1577. doi:10.1126/science.1100576

Gambetta, M.C., Oktaba, K., Müller, J., 2009. Essential role of the glycosyltransferase
sxc/Ogt in polycomb repression. Science 325, 93–96. doi:10.1126/science.1169727

Gaydos, L.J., Rechtsteiner, A., Egelhofer, T.A., Carroll, C.R., Strome, S., 2012. Antagonism
between MES-4 and Polycomb repressive complex 2 promotes appropriate gene
expression in C. elegans germ cells. Cell Rep 2, 1169–1177.
doi:10.1016/j.celrep.2012.09.019

Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., Ferrin,
T.E., 2018. UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci. 27, 14–25. doi:10.1002/pro.3235
Grau, D.J., Chapman, B.A., Garlick, J.D., Borowsky, M., Francis, N.J., Kingston, R.E., 2011.
Compaction of chromatin by diverse Polycomb group proteins requires localized regions
of high charge. Genes Dev. 25, 2210–2221. doi:10.1101/gad.17288211
Guidotti, N., Lechner, C.C., Bachmann, A.L., Fierz, B., 2019. A Modular Ligation Strategy
for Asymmetric Bivalent Nucleosomes Trimethylated at K36 and K27. Chembiochem
20, 1124–1128. doi:10.1002/cbic.201800744
Günesdogan, U., Jäckle, H., Herzig, A., 2010. A genetic system to assess in vivo the functions
of histones and histone modifications in higher eukaryotes. EMBO Rep. 11, 772–776.
doi:10.1038/embor.2010.124
Hansen, K.H., Bracken, A.P., Pasini, D., Dietrich, N., Gehani, S.S., Monrad, A., Rappsilber,
J., Lerdrup, M., Helin, K., 2008. A model for transmission of the H3K27me3 epigenetic
mark. Nat. Cell Biol. 10, 1291–1300. doi:10.1038/ncb1787
Henderson, R., Sali, A., Baker, M.L., Carragher, B., Devkota, B., Downing, K.H., Egelman,
E.H., Feng, Z., Frank, J., Grigorieff, N., Jiang, W., Ludtke, S.J., Medalia, O., Penczek,
P.A., Rosenthal, P.B., Rossmann, M.G., Schmid, M.F., Schröder, G.F., Steven, A.C.,
Stokes, D.L., Westbrook, J.D., Wriggers, W., Yang, H., Young, J., Berman, H.M., Chiu,
W., Kleywegt, G.J., Lawson, C.L., 2012. Outcome of the first electron microscopy
validation task force meeting., in:. Presented at the Structure (London, England : 1993),
pp. 205–214. doi:10.1016/j.str.2011.12.014
Ilca, S.L., Kotecha, A., Sun, X., Poranen, M.M., Stuart, D.I., Huiskonen, J.T., 2015.
Localized reconstruction of subunits from electron cryomicroscopy images of
macromolecular complexes. Nat Commun 6, 8843–8. doi:10.1038/ncomms9843
Jani, K.S., Jain, S.U., Ge, E.J., Diehl, K.L., Lundgren, S.M., Müller, M.M., Lewis, P.W.,
Muir, T.W., 2019. Histone H3 tail binds a unique sensing pocket in EZH2 to activate the
PRC2 methyltransferase. Proc. Natl. Acad. Sci. U.S.A. 116, 8295–8300.
doi:10.1073/pnas.1819029116
Jiao, L., Liu, X., 2015. Structural basis of histone H3K27 trimethylation by an active
polycomb repressive complex 2. Science 350, aac4383–aac4383.
doi:10.1126/science.aac4383
Justin, N., Zhang, Y., Tarricone, C., Martin, S.R., Chen, S., Underwood, E., De Marco, V.,
Haire, L.F., Walker, P.A., Reinberg, D., Wilson, J.R., Gamblin, S.J., 2016. Structural
basis of oncogenic histone H3K27M inhibition of human polycomb repressive complex
2. Nat Commun 7, 11316. doi:10.1038/ncomms11316
Kasinath, V., Beck, C., Sauer, P., Poepsel, S., Kosmatka, J., Faini, M., Toso, D., Aebersold,
R., Nogales, E., 2020. JARID2 and AEBP2 regulate PRC2 activity in the presence of
H2A ubiquitination or other histone modifications. bioRxiv 2020.04.20.049213.
Kasinath, V., Faini, M., Poepsel, S., Reif, D., Feng, X.A., Stjepanovic, G., Aebersold, R.,
Nogales, E., 2018. Structures of human PRC2 with its cofactors AEBP2 and JARID2.
Science 359, 940–944. doi:10.1126/science.aar5700
Klymenko, T., Müller, J., 2004. The histone methyltransferases Trithorax and Ash1 prevent
transcriptional silencing by Polycomb group proteins. EMBO Rep. 5, 373–377.
doi:10.1038/sj.embor.7400111
Laprell, F., Finkl, K., Müller, J., 2017. Propagation of Polycomb-repressed chromatin requires
sequence-specific recruitment to DNA. Science 356, 85–88. doi:10.1126/science.aai8266
Laugesen, A., Højfeldt, J.W., Helin, K., 2019. Molecular Mechanisms Directing PRC2
Recruitment and H3K27 Methylation. Mol. Cell 74, 8–18.
doi:10.1016/j.molcel.2019.03.011
Li, H., Liefke, R., Jiang, J., Kurland, J.V., Tian, W., Deng, P., Zhang, W., He, Q., Patel, D.J.,
Bulyk, M.L., Shi, Y., Wang, Z., 2017. Polycomb-like proteins link the PRC2 complex to
CpG islands. Nature. doi:10.1038/nature23881

Lowary, P.T., Widom, J., 1998. New DNA sequence rules for high affinity binding to histone
octamer and sequence-directed nucleosome positioning. J. Mol. Biol. 276, 19–42.
doi:10.1006/jmbi.1997.1494
Luger, K., Mäder, A.W., Richmond, R.K., Sargent, D.F., Richmond, T.J., 1997. Crystal
structure of the nucleosome core particle at 2.8 A resolution. Nature 389, 251–260.
doi:10.1038/38444
Luger, K., Rechsteiner, T.J., Richmond, T.J., 1999. Preparation of nucleosome core particle
from recombinant histones. Meth. Enzymol. 304, 3–19.
Margueron, R., Justin, N., Ohno, K., Sharpe, M.L., Son, J., Drury, W.J., Voigt, P., Martin,
S.R., Taylor, W.R., De Marco, V., Pirrotta, V., Reinberg, D., Gamblin, S.J., 2009. Role
of the polycomb protein EED in the propagation of repressive histone marks. Nature 461,
762–767. doi:10.1038/nature08398
McGinty, R.K., Henrici, R.C., Tan, S., 2014. Crystal structure of the PRC1 ubiquitylation
module bound to the nucleosome. Nature 514, 591–596. doi:10.1038/nature13890
McKay, D.J., Klusza, S., Penke, T.J.R., Meers, M.P., Curry, K.P., McDaniel, S.L., Malek,
P.Y., Cooper, S.W., Tatomer, D.C., Lieb, J.D., Strahl, B.D., Duronio, R.J., Matera, A.G.,
2015. Interrogating the function of metazoan histones using engineered gene clusters.
Dev. Cell 32, 373–386. doi:10.1016/j.devcel.2014.12.025
Meers, M.P., Henriques, T., Lavender, C.A., McKay, D.J., Strahl, B.D., Duronio, R.J.,
Adelman, K., Matera, A.G., 2017. Histone gene replacement reveals a post-
transcriptional role for H3K36 in maintaining metazoan transcriptome fidelity. Elife 6,
1191. doi:10.7554/eLife.23249
Musselman, C.A., Gibson, M.D., Hartwick, E.W., North, J.A., Gatchalian, J., Poirier, M.G.,
Kutateladze, T.G., 2013. Binding of PHF1 Tudor to H3K36me3 enhances nucleosome
accessibility. Nat Commun 4, 2969–9. doi:10.1038/ncomms3969
Nekrasov, M., Wild, B., Müller, J., 2005. Nucleosome binding and histone methyltransferase
activity of Drosophila PRC2. EMBO Rep. 6, 348–353. doi:10.1038/sj.embor.7400376
Pengelly, A.R., Copur, Ö., Jäckle, H., Herzig, A., Müller, J., 2013. A histone mutant
reproduces the phenotype caused by loss of histone-modifying factor Polycomb. Science
339, 698–699. doi:10.1126/science.1231382
Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C.,
Ferrin, T.E., 2004. UCSF Chimera--a visualization system for exploratory research and
analysis. J Comput Chem 25, 1605–1612. doi:10.1002/jcc.20084
Poepsel, S., Kasinath, V., Nogales, E., 2018. Cryo-EM structures of PRC2 simultaneously
engaged with two functionally distinct nucleosomes. Nat. Struct. Mol. Biol. 25, 154–162.
doi:10.1038/s41594-018-0023-y
Rappsilber, J., Mann, M., Ishihama, Y., 2007. Protocol for micro-purification, enrichment,
pre-fractionation and storage of peptides for proteomics using StageTips. Nat Protoc 2,
1896–1906. doi:10.1038/nprot.2007.261
Rosenthal, P.B., Henderson, R., 2003. Optimal determination of particle orientation, absolute
hand, and contrast loss in single-particle electron cryomicroscopy. J. Mol. Biol. 333,
721–745. doi:10.1016/j.jmb.2003.07.013
Rosenthal, P.B., Rubinstein, J.L., 2015. Validating maps from single particle electron
cryomicroscopy. Curr. Opin. Struct. Biol. 34, 135–144. doi:10.1016/j.sbi.2015.07.002
Santos, dos, G., Schroeder, A.J., Goodman, J.L., Strelets, V.B., Crosby, M.A., Thurmond, J.,
Emmert, D.B., Gelbart, W.M., FlyBase Consortium, 2015. FlyBase: introduction of the
Drosophila melanogaster Release 6 reference genome assembly and large-scale migration
of genome annotations. Nucleic Acids Res. 43, D690–7. doi:10.1093/nar/gku1099
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source
platform for biological-image analysis. Nat. Methods 9, 676–682.
doi:10.1038/nmeth.2019
Schmitges, F.W., Prusty, A.B., Faty, M., Stützer, A., Lingaraju, G.M., Aiwazian, J., Sack, R.,
Hess, D., Li, L., Zhou, S., Bunker, R.D., Wirth, U., Bouwmeester, T., Bauer, A., Ly-

Hartig, N., Zhao, K., Chan, H., Gu, J., Gut, H., Fischle, W., Müller, J., Thomä, N.H.,
2011. Histone methylation by PRC2 is inhibited by active chromatin marks. Mol. Cell
42, 330–341. doi:10.1016/j.molcel.2011.03.025
Simon, M.D., Chu, F., Racki, L.R., la Cruz, de, C.C., Burlingame, A.L., Panning, B.,
Narlikar, G.J., Shokat, K.M., 2007. The site-specific installation of methyl-lysine analogs
into recombinant histones. Cell 128, 1003–1012. doi:10.1016/j.cell.2006.12.041
Sobolev, O.V., Afonine, P.V., Adams, P.D., Urzhumtsev, A., 2015. Programming new
geometry restraints: parallelity of atomic groups. J Appl Crystallogr 48, 1130–1141.
doi:10.1107/S1600576715010432
Streubel, G., Watson, A., Jammula, S.G., Scelfo, A., Fitzpatrick, D.J., Oliviero, G., McCole,
R., Conway, E., Glancy, E., Negri, G.L., Dillon, E., Wynne, K., Pasini, D., Krogan, N.J.,
Bracken, A.P., Cagney, G., 2018. The H3K36me2 Methyltransferase Nsd1 Demarcates
PRC2-Mediated H3K27me2 and H3K27me3 Domains in Embryonic Stem Cells. Mol.
Cell 70, 371–379.e5. doi:10.1016/j.molcel.2018.02.027
Tan, Y.Z., Baldwin, P.R., Davis, J.H., Williamson, J.R., Potter, C.S., Carragher, B., Lyumkis,
D., 2017. Addressing preferred specimen orientation in single-particle cryo-EM through
tilting. Nat. Methods 14, 793–796. doi:10.1038/nmeth.4347
Terwilliger, T.C., Ludtke, S.J., Read, R.J., Adams, P.D., Afonine, P.V., 2019. Improvement
of cryo-EM maps by density modification 58, 214–31. doi:10.1101/845032
Wang, X., Paucek, R.D., Gooding, A.R., Brown, Z.Z., Ge, E.J., Muir, T.W., Cech, T.R.,
2017. Molecular analysis of PRC2 recruitment to DNA in chromatin and its inhibition by
RNA. Nat. Struct. Mol. Biol. 24, 1028–1038. doi:10.1038/nsmb.3487
Worden, E.J., Hoffmann, N.A., Hicks, C.W., Wolberger, C., 2019. Mechanism of Cross-talk
between H2B Ubiquitination and H3 Methylation by Dot1L. Cell 176, 1490–1501.e12.
doi:10.1016/j.cell.2019.02.002
Yu, J.-R., Lee, C.-H., Oksuz, O., Stafford, J.M., Reinberg, D., 2019. PRC2 is high
maintenance. Genes Dev. 33, 903–935. doi:10.1101/gad.325050.119
Yuan, W., Xu, M., Huang, C., Liu, N., Chen, S., Zhu, B., 2011. H3K36 methylation
antagonizes PRC2-mediated H3K27 methylation. J. Biol. Chem. 286, 7983–7989.
doi:10.1074/jbc.M110.194027
Zacher, B., Michel, M., Schwalb, B., Cramer, P., Tresch, A., Gagneur, J., 2017. Accurate
Promoter and Enhancer Identification in 127 ENCODE and Roadmap Epigenomics Cell
Types and Tissues by GenoSTAN. PLoS ONE 12, e0169249.
doi:10.1371/journal.pone.0169249
Zhang, K., 2016. Gctf: Real-time CTF determination and correction. J. Struct. Biol. 193, 1–
12. doi:10.1016/j.jsb.2015.11.003
Zheng, S.Q., Palovcak, E., Armache, J.-P., Verba, K.A., Cheng, Y., Agard, D.A., 2017.
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. Nat. Methods 14, 331–332. doi:10.1038/nmeth.4193
Zhou, Q., Huang, X., Sun, S., Li, X., Wang, H.-W., Sui, S.-F., 2015. Cryo-EM structure of
SNAP-SNARE assembly in 20S particle. Cell Res. 25, 551–560. doi:10.1038/cr.2015.47
Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., Scheres,
S.H., 2018. New tools for automated high-resolution cryo-EM structure determination in
RELION-3. Elife 7, 163. doi:10.7554/eLife.42166
1027
1028

1029

Figure 1

 FIGURE 1

 Interaction of the PRC2 catalytic lobe with nucleosomal DNA orients the H3 N-terminus

 for H3K27 binding to the active site.

(**A**) Domain organization in the five subunits of PHF1-PRC2. Dashed boxes indicate protein

portions visible in the PHF1-PRC2:di-Nuc cryo-EM reconstruction and fitted in the structural

model. In PHF1, C corresponds to the short C-terminal fragment used in PHF1$_C$-PRC2.

(**B**) Coomassie-stained SDS PAGE analysis of representative PHF1-PRC2 (upper panel)

and *X.l.* octamer preparations (lower panel) after size-exclusion chromatography (SEC)

purification. Pooled fractions of PHF1-PRC2, incubated with heterodimeric dinucleosomes

generated by DNA ligation of a reconstituted unmodified and a H3Kc27me3-modified

mononucleosome were used as input material for cryo-EM analysis.

(**C**) Cryo-EM reconstruction of PHF1-PRC2:di-Nuc in two orientations with fitted crystal

structures of human PRC2 catalytic lobe (PDB: 5HYN, (Justin et al., 2016)) and nucleosomes

(1AOI, (Luger et al., 1997)) in a di-Nuc model with 35 bp linker DNA (see also

Supplementary Figures 1-4, Table S1, Movie S1). Density is colored as in (**A**) to show PRC2

subunits, DNA (blue) and octamers of substrate (pink) and allosteric (yellow) nucleosomes.

Boxes indicate regions shown in (**D**), (**E**) and (**F**), respectively.

(**D**) Interaction of EZH2$_{CXC}$ residues with the DNA gyres of the substrate nucleosome; residues

mutated in PRC2$^{CXC>A}$ are indicated. For the H3 N-terminus (pink), only the peptide backbone

is shown in this view (see **F**).

(**E**) Interface formed by EED and the EZH2 SBD domain with DNA gyres on the allosteric

nucleosome; residues mutated in PRC2$^{EED>A}$ are indicated. Asterisk indicates the approximate

location of a residue, which is not built in the model.

(**F**) The H3 N-terminus (pink), shown as a pseudoatomic model fitted into the 4.4 Å density

map, is recognized by EZH2 through an extensive interaction network (see text). Note the well-

defined side-chain density of H3K36 (see also **Supplementary Figure 3D and 4C-E**).

1056

1057

A

FIGURE 2

The EZH2$_{CXC}$-DNA interaction interface is critical for H3K27 methylation on

nucleosomes.

(**A**) Binding reactions with indicated concentrations of PRC2 (lanes 1-10), PRC2$^{CXC>A}$ (lanes

11-20) or PRC2$^{CXC>A/EED>A}$ (lanes 21-30) and 45 nM 6-carboxyfluorescein-labeled

mononucleosomes, analyzed by EMSA on 1.2% agarose gels; for analysis of PRC2$^{EED>A}$

binding, see **Supplementary Figure 5A**.

(**B**) Quantitative analysis of EMSA data in A by densitometry of 6-carboxyfluorescein signals

from independent experiments (n=3); error bars, SEM.

(**C**) Western Blot (WB) analysis of H3K27me1 and H3K27me3 formation in HMTase reactions

with indicated concentrations of PRC2 and PRC2$^{CXC>A}$ on 446 nM mononucleosomes (lanes 1-

7) or 223 nM dinucleosomes (lanes 8-14). Note that these concentrations result in equal

numbers of nucleosomes and therefore equal numbers of H3 substrate molecules in the

reactions on mono- and dinucleosomes, as can be seen from the Coomassie-stained gel of the

reactions in **Supplementary Figure 5B**. H4 WB signal served as control for Western blot

processing.

1074

1075

1076

Figure 3

FIGURE 3

The unmodified H3K36 side chain in the EZH2$_{CXC}$-DNA interaction interface is critical

for H3K27 methylation on nucleosomes.

(**A**, **B**) EMSA analysis and quantification as in **Figure 2A and B**, using PRC2 and

mononucleosomes that were unmodified (lanes 1-10) or contained a trimethyllysine analog at

H3K36 (H3Kc36me3, lanes 11-20).

(**C**) Western Blot (WB) analysis of HMTase reactions with PRC2 as in **Figure 2C** on

unmodified (lanes 1-4), H3Kc36me3 (lanes 5-7), H3^{K36R} (lanes 8-10) or H3^{K36A} (lanes 11-13)

mononucleosomes (446 nM). Coomassie stained gel of reactions is shown in **Supplementary**

Figure 6A. Bottom: quantification of H3K27me3 and H3K27me1 chemiluminescence signals,

respectively, by densiometry analysis from three independent experiments. In each experiment,

the methylation signal in lane 4 was defined as 100% and used to quantify the corresponding

H3K27 methylation signals in the other lanes on the same membrane. Circles show individual

data points and error bars SEM.

(**D**) HMTase reactions monitoring H3K27me1 formation by PRC2 on H3$_{18-42}$ peptides that were

unmodified (top) or contained K36me3 (bottom). Left: Deconvoluted ESI-MS spectra from

data shown in **Supplementary Figure 6B**. On both substrates, areas of the four colored peaks

of H3K27me1-modified and unmodified substrate peptides were used for quantification of

H3K27me1 formation. Right: Symbols represent percentages of peptides carrying H3K27me1

in three independent experiments, error bars show SEM; Welch's t-test showed no significant

(ns) difference between H3K27 monomethylation on the two peptide substrates.

1098

1099

1100

Figure 4

 FIGURE 4

 H3^{K36A} and *H3^{K36R}* mutants show reduced levels of H3K27me3.

 (**A**) Western blot analysis on serial dilutions (4:2:1) of total cell extracts from wing, haltere

 and 3rd leg imaginal disc tissues dissected from *wildtype* (*wt*, lanes 1-3) and *H3^{K36R}* mutant

 (lanes 4-6) third instar larvae. Blots were probed with antibodies against H3K36me3,

 H3K36me2 or H3K27me3; in each case, probing of the same membranes with antibodies

 against Caf1-55 and H2B served as controls for loading and western blot processing. Note the

 reduced levels of H3K36me3 and H3K36me2 but also of H3K27me3 in *H3^{K36R}* mutants

 compared to *wildtype* (*wt*) (see text). See Materials and Methods for details of all genotypes.

 (**B**) Western blot analysis on serial dilutions (8:4:2:1) of total nuclear extracts from 21-24 hr

 old *wt* (lanes 1-4) and *H3^{K36A}* mutant (lanes 5-8) embryos, probed with antibodies against

 H3K36me3, H3K36me2 or H3K27me3; and with antibodies against Caf1-55 and H2B as

 controls. Note that H3K36me3 and H3K36me2 levels are reduced in *H3^{K36R}* mutants compared

 to *wt* but that H3K27me3 levels appear undiminished in the mutant (see text).

 (**C**) Left, violin plot showing the fold-change of H3K36me2 coverage in *H3^{K36R}* mutant larvae

 relative to *wt* at genes that in wildtype larval CNS and imaginal disc tissues are decorated with

 H3K36me2 (see Materials and Methods). The dashed line marks the median reduction (4.5-

 fold), the dotted lines indicate indicated the interval comprising 80% of regions. Right, Bee

 plot showing the fold-change of H3K27me3 coverage in *H3^{K36R}* mutant larvae relative to *wt* at

 genes that in wildype larval CNS and imaginal disc tissues are associated with high-level

 H3K27me3 regions (see Materials and Methods). The dashed line marks the median reduction

 (2.1-fold), the dotted lines indicate the interval comprising 80% of regions. Note that

 H3K27me3 coverage at the HOX genes *abd-A*, *Abd-B*, *Ubx* and *Antp* is between 3- and 4-fold

 reduced.

 (**D**) Analysis and representation as in (**C**) but showing fold-changes in H3K36me2 and

 H3K27me3 coverage in *H3^{K36A}* mutant late-stage embryos relative to *wt* at genes that in

 wildtype embryos are decorated with H3K36me2 and H3K27me3, respectively. Note that

 H3K27me3 coverage at the HOX genes *abd-A*, *Abd-B*, *Ubx* and *Antp* is about 1.5-fold reduced.

See also **Supplementary Figure 8**.

(**E**) H3K27me3 and H3K36me2 ChIP-seq profiles in larval CNS and imaginal disc tissues

from *wt* (blue) and *H3^{K36R}* mutant (purple) third instar larvae; in the tracks showing the profiles

in the *H3^{K36R}* mutant, the *wt* profile is superimposed as reference (see **Table S2** and Materials

and Methods for information about normalization). Top: genomic interval containing the

Bithorax-Complex harbouring the HOX genes *Ubx*, *abd-A* and *Abd-B*; bottom: genomic interval

containing the *Antennapedia-Complex* with the HOX genes *lab*, *pb*, *Dfd*, *Scr* and *Antp*. Note

the 3- to 4-fold reduction of H3K27me3 levels across the *Bithorax* and *Antennapedia* loci in

H3^{K36R} mutants. Also note that the analyzed tissues (CNS, thoracic imaginal discs and eye-

antenna discs) represent a mixed population of cells with respect to transcriptionally active and

repressed states of each HOX gene. For each HOX gene there is a substantially larger

proportion of cells in which the gene is decorated with H3K27me3 and repressed by Polycomb

and only a small proportion of cells in which the gene is active and decorated with H3K36me2.

(**F**) H3K27me3 and H3K36me2 ChIP-seq profiles at the *Bithorax* and *Antennapedia* loci as in

(**E**) but from *wt* (blue) and *H3^{K36A}* mutant (yellow) late-stage embryos with the *wt* profile

superimposed in the tracks showing the profiles in the *H3^{K36A}* mutant. H3K27me3 levels across

the *Bithorax* and *Antennapedia* loci in *H3^{K36A}* mutants are only about 1.5-fold reduced

compared to *wt*.

1147

1148

FIGURE 5

***Drosophila* with H3^{K36R} or H3^{K36A} chromatin show defective Polycomb repression at HOX**

genes

(**A**) Ventral views of stage 16 wildtype (*wt*), *H3^{K36A}*, *H3^{K36R}*, *H3^{K27R}* or *esc* (*esc$^-$*) mutant

embryos, stained with antibody against Abd-B protein; the *esc* mutant embryo lacked both

maternal and zygotic expression of *esc* (see Materials and Methods for details of all genotypes).

The vertical bar marks the anterior boundary of *Abd-B* expression in parasegment (ps) 10 in *wt*

embryos. Note the stochastic misexpression of Abd-B protein in single cells anterior to ps10

in *H3^{K36R}* and *H3^{K36A}* mutant embryos (arrowheads). *H3^{K27R}* and *esc* mutant embryos show

widespread misexpression of Abd-B protein in the head-to-tail pattern characteristic of PcG

mutants. For reasons that are not well understood, *H3^{K36A}* and *H3^{K36R}* mutants also show partial

loss of Abd-B expression in cells in ps10 (empty arrowheads).

(**B**) Larval CNS and brain lobe tissues from wildtype (*wt*) or *H3^{K36R}* mutant third instar larvae,

stained with antibody against Ubx protein (red) and Hoechst (DNA) to label all nuclei; location

of CNS and brain lobes (BL) are indicated in the right panel. The white bars mark the anterior

boundary of *Ubx* expression in ps5 in *wt* embryos, the asterisk marks the Ubx-expressing cells

in the central midline of ps4 that are part of the wild-type Ubx pattern. Note the stochastic

misexpression of Ubx protein in many single cells anterior to ps5 in the CNS and in the brain

lobes (arrowheads).

(**C**) Imaginal wing (W), haltere (H) and 3rd leg (3L) discs from wildtype (*wt*) or *H3^{K36R}* mutant

third instar larvae and, as reference, discs from a larvae with clones of *H3^{K27R}* mutant cells that

are marked by the absence of GFP. In all cases discs were stained with antibody against Ubx

protein (red) and Hoechst (DNA) to label all nuclei. In *wt* animals, Ubx is expressed in the

halter and 3rd leg disc but not in the wing disc where it is repressed by the PcG machinery. Note

that in *H3^{K36R}* mutants, Ubx is misexpressed in small clusters of cells in the pouch area of the

wing disc (arrowheads) but remains repressed in the rest of the wing disc. Such misexpression

was detected in 50% of wing discs (*n* = 28). As reference, a wing discs with *H3^{K27R}* mutant

clones is shown, where all cells in the clones in the wing pouch (arrowheads) show

misexpression of Ubx and only mutant cells in the notum and hinge show no misexpression

(empty arrowheads) (cf. (Pengelly et al., 2013)). Also note that in $H3^{K36R}$ mutants ($n > 30$

mutant animals analysed), Ubx expression in haltere and leg discs appears unperturbed

(asterisks).

(**D**) Eye-antennal imaginal discs from wildtype (*wt*) or $H3^{K36R}$ mutant larvae and below discs

from larvae with clones of $H3^{K36A}$ or $H3^{K27R}$ mutant cells that are marked by the absence of

GFP. All animals were stained with antibody against Antp protein (red) and Hoechst (DNA)

to label all nuclei. Antp is not expressed in the eye-antennal disc of *wt* animals. Note that in

$H3^{K36R}$ mutant discs, Antp is misexpressed in large clusters of cells (arrowheads) in the antenna

primordium (Ant). Note that Antp is also misexpressed in $H3^{K36A}$ or $H3^{K27R}$ mutant cell clones

in the antenna primordium (arrowheads) and that in these cases misexpression also only occurs

in a subset of the mutant cells and not in all clones.

1189

1190

1191

1192

1193

Supplementary Figure 1

SUPPLEMENTARY FIGURE 1

Initial Cryo-EM analysis of the PHF1-PRC2:di-Nuc complex (related to Fig. 1).

(**A**) Representative micrograph of the cryo-EM dataset (left) and reference-free 2D classes from particles picked without templates (right) (performed to ensure that no bias was introduced through templates picking and references in 3D classification).. Circles indicate particles, which were picked with templates and directly subjected to 3D analysis (see **fig. S2**).

(**B**) Local resolution estimation of the 5.2 Å overall PHF1-PRC2:di-Nuc map. The substrate nucleosome and the adjacent part of EZH2 are well resolved (colors red to yellow).

(**C**) Spherical angular distribution of particles included in the final reconstruction of PHF1-PRC2:di-Nuc.

(**D**) Output from the 3DFSC Processing Server (https://3dfsc.salk.edu/ (Tan et al., 2017)) showing the Fourier Shell Correlation (FSC) as a function of spatial frequency, generated from masked independent half maps of PRC2:diNuc: global FSC (red), directional FSC (blue histogram) and deviation from mean (spread, green dotted line). The nominal overall resolution of 5.24 Å was estimated according to the gold standard FSC cutoff of 0.143 (grey dotted line) (Rosenthal and Henderson, 2003). Sphericity is an indication for anisotropy and amounts to 0.806 in this data. The minor directional anisotropy of the data can be explained by the slightly preferred orientation and missing views as seen in (**C**).

(**E**) Top: Refined and postprocessed cryo-EM density map of overall PHF1-PRC2:di-Nuc colored according to the subunit organization. Bottom: pseudoatomic model of fitted crystal structure of the human PRC2 catalytic lobe (PDB: 5HYN) and a di-Nuc model with 35 bp linker DNA (Poepsel et al., 2018), including PDB 1AOI.

Motion correction and dose weighting with MotionCor 2
Nr. aligned micrographs: 3466
Pixel size (all steps of processing): 1.746 Å/pix

PRC2-Dinucleosome structure (EMD-7306),
low-pass filtered to 60 Å

Template-based Autopicking in Gautomatch
Nr. of initial particle candidates: 1,028,229

2 rounds of 3D classification to separate particles from contaminations in subsets of ca. 300,000 particles
each with T= 8, 3-4 classes, 25 iterations. 330,482 particles were further subjected to 2 rounds of 3D classification
with T=8, 3 classes. 25 iterations to separate well-aligning PRC2-Dinucleosome 3D models
from "empty" or not well-aligning PRC2-Dinucleosomes.

26,487 particles   28,088 particles   29,414 particles
16.0 Å   15.6 Å   19.5 Å

3D refinement

10.0 Å, 54,575 particles

3D classification wtihout translational and spatial sampling (10 classes, T=4, 25 iterations)

1715 particles   1645 particles   1656 particles   2249 particles   6649 particles   1461 particles   27,014 particles   3971 particles   5562 particles   2653 particles
23.7 Å   23.7 Å   24.8 Å   20.2 Å   16.0 Å   24.8 Å   11.1 Å   18.16 Å   16.5 Å   19.5 Å

45,849 particles

Best classes with
highest resolution

Iterative rounds of focused 3D refinement, CTFrefine and polishing

Overall PHF1-PRC2:di-Nuc
5.4 Å / 5.2 Å after postprocessing, b-factor=-90.7;
Mask used for postprocessing:

Signal subtraction of the allosteric nucleosome
as well as EED/EZH2$_{allo}$

Focused 3D refinement

EZH2$_{sub}$-Nuc$_{sub}$
4.8 Å / 4.4 Å after postprocessing, b-factor=-76.5;
Mask used for postprocessing:

1218   **SUPPLEMENTARY FIGURE 2**
1219
Overview of the cryo-EM Data-Processing and Particle Sorting Scheme (related to Figure
1).
Processing and particle sorting scheme, also described in Methods. Squares indicate 3D classes
(and corresponding particles) chosen for further processing steps based on their nominal global
resolution values, translational and rotational accuracy and the presence of detailed structural
information. Two final reconstructions were obtained in this study: Overall PHF1-PRC2:di-
Nuc, and $EZH2_{sub}$-Nuc_{sub} after performing signal subtraction (mask indicated in pink) and
focused refinement. Masks used for postprocessing are shown in yellow.
1228

Supplementary Figure 3

 SUPPLEMENTARY FIGURE 3

1230

Cryo-EM analysis of the focused EZH2$_{sub}$-Nuc$_{sub}$ map (related to Fig. 1).

(**A**) Local resolution estimation of the focused 4.4 Å EZH2$_{sub}$:Nuc$_{sub}$ reconstruction. Regions

in the nucleosome core as well as the adjacent regions including parts of the H3 N-terminus

close to the exit side of the nucleosome are well resolved (4.0 – 5.5 Å). Regions close to the

mask, especially the nucleosomal DNA and parts of EZH2, are less well resolved (colors green

to blue).

(**B**) Global FSC generated from masked independent half maps of EZH2$_{sub}$-Nuc$_{sub}$ (Focused,

blue line) and the overall PHF1-PRC2:di-Nuc (Overall, violet line) were plotted against spatial

frequency. The resolution of 4.4 Å for for EZH2$_{sub}$-Nuc$_{sub}$ map and 5.2 Å for the overall PHF1-

PRC2:di-Nuc map were estimated according to the gold standard FSC cutoff of 0.143 (brown

dotted line) (Rosenthal and Henderson, 2003)

(**C**) FSC between the atomic model and the masked (applied in Phenix) map of EZH2$_{sub}$-Nuc$_{sub}$

after real-space refinement (Afonine et al., 2018). Green line represents the cut-off at 0.5 (4.6

Å) and blue line represents the cut-off at 0.143 (4.3 Å) (see also Table 1) (Henderson et al.,

2012; Rosenthal and Henderson, 2003; Rosenthal and Rubinstein, 2015).

(**D**) Selected regions within EZH2$_{sub}$-Nuc$_{sub}$ showing side chain density, e.g. K36 (red arrow).

A red asterisk indicates the last residue of the H3 tail visible in known crystal structures (usually

P38 or H39). The quality of the map around K36 is shown as a separate zoom-out below and

in three different views to demonstrate the lack of anisotropy present in the density.

1250

1251

1252

Supplementary Figure 4

A. Front view/CXC

B. Top view/Bridge Helix

C. Bottom view/K36 position

D. Back view/Bridge helix and K36 position

E.

F.

H3 tail/E579 pocket

SUPPLEMENTARY FIGURE 4

The improved map of the interaction between EZH2 and the substrate nucleosome after focused refinement reveals location of H3K36 and it's environment (related to Fig. 1).

(**A**) The front view of EZH2$_{sub}$-Nuc$_{sub}$ cryo-EM density and model shows details of the EZH2$_{CXC}$ interaction with nucleosomal DNA.

(**B**) The top view of EZH2$_{sub}$-Nuc$_{sub}$ cryo-EM shows a tubular density into which based on recent findings of (Kasinath et al., 2020) a helix was built. The bridge helix, which based on this study is likely constituted of the EZH2 residues 497-511, is located above V35 of the H3 tail. As can be seen when observing the density-modified map (Terwilliger et al., 2019) of EZH2$_{sub}$-Nuc$_{sub}$ at lower threshold, it presumably engages in interactions with the nucleosomal DNA, the H3 tail and EZH2, as described in greater detail in (Kasinath et al., 2020).

(**C**) The bottom view of EZH2$_{sub}$-Nuc$_{sub}$ cryo-EM density and model shows details of the vicinity of K36 with the corresponding density for the H3 tail, EZH2 and nucleosomal DNA. The orange square indicates the region shown as a zoom-in in (**E**).

(**D**) The back view of EZH2$_{sub}$-Nuc$_{sub}$ cryo-EM density and model shows details of the location of K36 and the bridge helix.

(**E**) Zoom-in views of H3K36 and it's chemical environment. Approximate distances of the epsilon-amino group of H3K36 to the nearest residues are indicated with a dotted grey line.

(**F**) Location of the Glu-579 pocket (Jani et al., 2019) in the EZH2$_{sub}$:Nuc$_{sub}$ reconstruction and it's distance to H3K36 (app. 19 Å). The described mechanism by Jani et al (Jani et al., 2019) involving recognition of H3K36 by Glu-579 is incompatible with the presented structural data as the location differs significantly and major rearrangements as the relocation of the helix-loop region between residues 564-576 would be necessary to avoid the given steric and geometric hindrance and allow for potential interaction.

A

B

 SUPPLEMENTARY FIGURE 5

 The EZH2$_{CXC}$-DNA interaction interface is critical for H3K27 methylation on
 nucleosomes (related to Fig. 2).
 (**A**) Binding reactions with indicated concentrations of PRC2 (lanes 1-10) or PRC2$^{EED>A}$ (lanes
 11-20) and 45 nM 6-carboxyfluorescein-labeled mononucleosomes, analyzed by EMSA on
 1.2% agarose gels.
 (**B**) Coomassie-stained 4-12% SDS-PAGE of the HMTase reactions shown in **Fig. 2C**. *Xenopus*
 laevis (*X.l.*) nucleosomes were used for these experiments. The short 5-kDa PHF$_C$ fragment is
 not visible on this gel.

A

B

C

 SUPPLEMENTARY FIGURE 6

 Accommodation of unmodified H3K36 in the EZH2$_{CXC}$-DNA interaction interface is
 essential for H3K27 methylation on nucleosomes and PHF1-PRC2 (related to Fig. 3).
 (**A**)   Coomassie-stained 4-12% SDS-PAGE of the HMTase reactions shown in **Fig. 3C**.
 Drosophila melanogaster (*D.m.*) nucleosomes were used for these experiments. The short 5-
 kDa PHF$_C$ fragment is not visible on this gel.
 (**B**) Full ESI MS spectra (upper part) and full deconvoluted MS spectra (lower part) shown for
 input peptides without PRC2 as a control (left) and with PRC2 (right) to ensure no overlapping
 between possible adduct peaks and monomethylation peaks.
 (**C**) Western Blot (WB) analysis of HMTase reactions with full-length PHF1-PRC2 on
 unmodified (lanes 1-4) or H3Kc36me3 (lanes 5-7) mononucleosomes (446 nM).

wt

H3^{K36R}

H3^{K36A}

A

From embryos (as above):
83% develop into pupae
75% develop into viable and fertile
adults (as below)

From embryos (as above):
81% † at variable time points
between 1st and 3rd instar
18% † as pupae
1% † as pharate adults (as below)

From embryos (as above):
96% † as unhatched embryos
4% † as 1st instar larvae

B

adult ♀

pharate adult ♀

C

adult ♂   adult ♂   A

pharate adult ♂   A   pharate adult ♂   A

D

L2   L3   L1   adult ♂

L2   L3   L1   pharate adult ♂

1308 **SUPPLEMENTARY FIGURE 7**
1309
***Drosophila* with H3^{K36R} or H3^{K36A} mutant chromatin arrest development at different**
stages
(**A**) Ventral views of cuticles from wildtype (*wt*), *H3^{K36R}*, or *H3^{K36A}* mutant embryos. Note that
the cuticle pattern of the mutant animals is indistinguishable from that of the *wt* embryo.
Below: for each genotype, the fraction of embryos that developed into larve, pupae, pharate
adults or viable adults is listed. The fraction was determined by monitoring the development
of collected hatched 1st instar larvae (*wt*: *n*=300, *H3^{K36R}*: *n*=2000) or unhatched embryos
(*H3^{K36A}*: *n*=200). The GFP marker on the Balancer chromosomes was used for identifying
H3^{K36R} and or *H3^{K36A}* mutants.
(**B**) Dorsal views of the posterior portion of the thorax and of the abdomen. From 2000 hatched
H3^{K36R} mutant 1st instar larvae, a total of 18 pharate adults was recovered. Most *H3^{K36R}* mutant
pharate adults showed a relatively normal overall body patterning apart from the homeotic
transformations illustrated below.
(**C**) Frontal view of adult heads illustrating the antenna-to-leg transformation in *H3^{K36R}* mutant
pharate adults. The antenna-to-leg transformation in *H3^{K36R}* mutant animals ranged from mild
(arrows) to more extensive transformations with formation of leg-like structures such as in this
extreme case (arrowheads).
(**D**) The sex comb in males is normally only present on the protoracic (L1) legs (arrowheads).
Among the *H3^{K36R}* mutant pharate adult males recovered (*n*=13), five showed one or several
extra sex comb teeth (arrow) on the meso- (L2) or metathoracic (L3) legs. Extra sex comb teeth
in adults are a hallmark phenotype of Polycomb mutants.
1331
1332

Supplementary Figure 8

1333   **SUPPLEMENTARY FIGURE 8**
1334
H3^{K36R} and *H3^{K36A}* **mutants show altered H3K36me2 and H3K27me3 profiles (related to**
Fig. 4)
(**A**) Top, scatter plots showing H3K36me2 coverage in relation to H3K27me3 coverage in *wt*
larvae. Green dots represent 9200 gene bodies overlapping with genomic intervals showing
H3K36me2 enrichment, blue dots represent 1030 gene bodies overlapping with genomic
intervals defined as high-level H3K27me3 domains (Bonnet et al., 2019), and grey dots
represent 6300 gene bodies showing no enrichment for either methylation mark in larvae (see
Materials and Methods). Bottom, scatter plot showing the H3K36me2 read coverage in relation
to H3K27me3 read coverage in *H3^{K36R}* mutant larvae.
(**B**) As in (**A**) but showing H3K36me2 coverage in relation to H3K27me3 coverage in *wt*
embryos (top) and in *H3^{K36A}* mutant embryos (bottom). Green dots represent 10800 gene bodies
overlapping with genomic intervals showing H3K36me2 enrichment, blue dots represent 1030
gene bodies overlapping with genomic intervals defined as high-level H3K27me3 domains
(Bonnet et al., 2019), and grey dots represent 5400 gene bodies showing no enrichment for
either methylation mark in embryos (see Materials and Methods).
1350
1351
1352
1353

Table S1

Cryo electron microscopy data collection summary, processing statistics and model

Cryo electron microscopy data collection (applicable to Overall PHF1-PRC2:diNuc and EZH2$_{sub}$-Nuc$_{sub}$ maps)	
Microscope	FEI Titan Krios GII
Voltage (kV)	300
Camera	Gatan K2-Summit
Energy Filter	Gatan Quantum-LS (GIF)
Pixel size (Å/pix) (calibrated)	1.75
Nominal magnification (x)	81000
Preset target global defocus range (μm)	0.5 - 3.5
Total electron exposure (fluence, e$^-$/Å$^{2)}$	52,96
Exposure rate (flux) (e$^-$/ Å^2/s)	3,47
Nr. of frames collected per micrograph	60
Energy filter slit width (eV)	20
Automation Software	SerialEM

3D reconstruction (applicable to Overall PHF1-PRC2:diNuc and EZH2$_{sub}$-Nuc$_{sub}$ maps)		
Number of movies	3466	
Initially selected particle candidates	1,028,229	
Final number of particles	45,849	
	Overall PHF1-PRC2:diNuc	**EZH2$_{sub}$-Nuc$_{sub}$**
Resolution $_{\text{FSC independent halfmaps (0.143) masked}}$ (Å)^a	5.24	4.36
Local resolution range (Å)	4.01 - 24.97	4.01 – 15.00
Sharpening B-factor (Å^2)	-90.7	-76.5
Refinement EZH2$_{sub}$-Nuc$_{sub}$		
No. atoms		28866 (Hydrogens: 13086)
Residues		Protein: 1171 Nucleotide: 312
Ligands		ZN: 8
CC$_{mask,}$ CC$_{box}$, CC$_{peaks}$, CC$_{volume}$^{b}		0.75, 0.83, 0.66, 0.74
Mean CC for ligands		0.72
Resolution$_{\text{FSC masked map vs. model (0/0.143/0.5)}}$ (Å)^b		4.3/4.3/4.6
R.m.s. deviations		
Bond lengths (Å)		0.004
Bond angles (°)		0.642
Ramachandran favored (%)		97.51
Ramachandran gen. allowed (%)		2.32
Ramachandran disallowed (%)		0.17

MolProbity score		1.53
Clash score		7.97
ADP (B factors)		
Iso/Aniso (#)		15925/0
Min/max/mean		
Protein		65.89/264.21/121.82
Nucleotide		97.31/221.62/132.65
Ligand		151.54/251.77/201.51
Rotamer outliers (%)		0.00
Cβ outliers (%)		0.00
CaBLAM outliers (%)		1.59

1357 [a]according to the Fourier Shell Correlation (FSC) cut-off criterion of 0.143 defined in (Rosenthal and Henderson, 2003)
1358 [b]according to the map-vs.-model Correlation Coefficient definitions in (Afonine et al., 2018a)
1359

1360   **Table S2.**

1361

1362   **Number of aligned reads to the *D. melanogaster* and *D. pseudoobscura* genomes from**
1363   **ChIP and input datasets and normalization process (related to Figure 4 and**
1364   **Supplementary Figure 8).**

1365

1366

1367   **Movie S1**

1368

1369   **Cryo-EM structure of the PHF1-PRC2:di-Nuc complex (related to Figure 1).**

1370