Microbiological safety of industrially reared insects for food: Identification of bacterial endospores and targeted detection of foodborne viruses

Vandeweyer[a], D., Lievens[b], B., Van Campenhout[a*], L.

[a] *KU Leuven, Department of Microbial and Molecular Systems (M²S), Lab4Food, Geel Campus, 2440 Geel, Belgium*

[b] *KU Leuven, Department of Microbial and Molecular Systems (M²S), Laboratory for Process Microbial Ecology and Bioinspirational Management (PME&BIM), Group T Campus, 3000 Leuven, Belgium*

[*] *Corresponding author:* Leen Van Campenhout, KU Leuven, Geel Campus, Kleinhoefstraat 4, 2440 Geel, Belgium. E-mail: leen.vancampenhout@kuleuven.be

Abstract

Edible insects are characterised by high microbial numbers of which the bacterial endospores fraction can survive thermal processing. It is unknown, however, which bacterial species occur as endospore in edible insects and what impact they have on food safety. Additionally, edible insects have never been explored for the presence of foodborne viruses so far. In this study, we found that the bacterial endospore fraction in a collection of mealworm and cricket samples obtained from commercial insect producers can comprise a large amount of *Bacillus cereus* group members that can pose insect or human health risks. Monitoring and effective mitigation of these risks are urged. By contrast, none of the foodborne viruses hepatitis A virus, hepatitis E virus and norovirus genogroup II were detected in the sample collection. Therefore, food safety risks originating from these viral pathogens can be concluded to be low.

Introduction

As from 2014, edible insects and insect-derived food products started to enter the European markets. According to the International Platform of Insects for Food and Feed (IPIFF), 6,000 tonnes of insect proteins were produced by its European members in 2019. The organisation estimates production to be around 3 mio tonnes in 2030. Due to the expansion of the insect sector risk assessments and food safety research regarding insects as a new food matrix were initiated, e.g. by EFSA[1]. In the past five years, many basic microbiological food safety questions were formulated and investigated, as reviewed by Garofalo *et al.* (2019)[2]. In summary, freshly reared insects may be highly contaminated with diverse microorganisms. Even though insects are not likely to be consumed in a raw status, it is important to know which organisms should be targeted by heat and other processing treatments. It was demonstrated both for fungi and bacteria for yellow mealworms (*Tenebrio molitor*)[3], lesser mealworms (*Alphitobius diaperinus*)[4] and tropical house crickets (*Gryllodes sigillatus*)[5] that high numbers of these microorganisms can easily be reduced, e.g. by applying a mild heat treatment.. However, it was repeatedly observed that a fraction of the bacteria, i.e. the endospores, can survive those mild heat treatments[3,6]. *Bacillus* and *Clostridium* are two relevant bacterial genera that are able to produce endospores. Since several members of these genera are food pathogens (e.g. *B. cereus*, *C. perfringens*, *C. botulinum*), the survival of endospores in the insect matrix may pose food safety risks. DNA-based studies have already identified, among others, the spore-forming genera *Bacillus*, *Brevibacillus, Clostridium, Lysinibacillus* and *Paenibacillus* in edible insects and insect products produced in Europe[7–12]. Yet, it is generally not clear which specific species were found and whether or not they were present in spore form. Consequently, the first goal of this research (experiment 1) was to identify which bacterial endospores can be present in raw samples of the two edible insect species reared mostly for human consumption

nowadays: the yellow mealworm (*Tenebrio molitor*) and the house cricket (*Acheta domesticus*).

Specific attention was given to *B. cereus*-related isolates, which were further characterised by

exploring specific genes with qPCR.

Another aspect that has not been explored so far is the presence of pathogenic foodborne

viruses in insects for food. The possibility that foodborne viruses such as hepatitis A virus

(HAV), hepatitis E virus (HEV) or norovirus (NoV) can be present in insects or derived

foodstuffs has been proposed by EFSA already in 2015[1]. The three said viruses are the most

commonly transmitted viruses through foodstuffs[13] and are relevant to be investigated for

insects and insect-based food matrices. This research was the first to elucidate whether the

selected human foodborne viruses HAV, HEV and NoV genogroup II (GII, responsible for the

majority of norovirus cases[14]) are present in raw mealworm and cricket species, and if so, in

what quantity (experiment 2).

When studying the microbiota of insects, it is very plausible that the organisms present

in their gut and on their exoskeleton differ between insects cultivated in the laboratory and the

same species reared at industrial scale, where the hygiene levels may be lower and hence the

house flora different. Since the aim of this work was to study bacterial endospores and viruses

with respect to safety for the consumer, insect samples obtained from several commercial insect

producers were studied and multiple batches per producer were included in the experimental

set-up.

Results and discussion

Bacterial endospore counts

In experiment 1, the raw insect samples investigated were first subjected to bacterial

endospore counts and, as a comparison, total viable counts (Figure 1). The obtained values for

total viable aerobic (between 7.5 and 8.7 log cfu/g) and anaerobic (7.4 to 8.7 log cfu/g) counts

are similar to the typically high counts for raw edible insects reported earlier[2,15]. Additionally,

the values for the aerobic and anaerobic counts are highly comparable. This may suggest that a

large fraction of both counts exists of facultative anaerobic species. When comparing both

insect species investigated, it is clear that the total viable counts for house crickets are higher

than those for yellow mealworms. This confirms what was proposed earlier[16], being that

microbiological findings of particular insect species should not be generalised for all edible

insects, as nevertheless is done in literature occasionally. For bacterial endospore counts, which

represent the fraction of the total viable counts present as spore form significant variation was

observed between samples from the same insect species, compared to the total viable counts.

This variation may result from differences in (hygiene in) the rearing environment and/or feed,

and is commonplace for insects[2,9,11]. Again, anaerobic and aerobic endospore counts displayed

comparable trends (Figure 1).

Currently in the insect sector, edible insects are typically subjected to a heat treatment

prior to processing and consumption, such as boiling and/or oven drying. It has been

demonstrated[3,5,6] that bacterial endospores can survive these treatments and remain present and

viable in insects for human consumption. The highest endospore count observed in this study

was 4.2 log cfu/g (for a house cricket sample). While no European microbiological criteria exist

specifically for edible insects, this count does surpass the lower action limit for *Bacillus cereus*

in edible insects as published by the Belgian Federal Agency for the Safety of the Food Chain[17].

In the case that a large fraction of these spores found would consist of *B. cereus*, this would

pose a health risk for human consumption. For clostridia, no criteria or action limits exist, but

it is advised to pay attention to these potential pathogens as well.

Identification of bacterial endospores

Microbiological health risks of food products are determined not only by population densities of microbial (sub)groups, but also by the particular species present in the foodstuff. In this study, we focussed on bacterial endospores, which are resistant against several treatments that are harmful for vegetative cells. Pathogenic bacteria present in their spore form are of major concern for edible insects.

In total, 142 endospore-forming isolates were collected from the bacterial endospore count plates obtained from yellow mealworm and house cricket samples. These isolates were present as endospore in the raw insect matrix and survived the pasteurisation treatment during analysis (as part of the ISO-based endospore count), germinated and grew on Plate Count Agar. By homogenising the sample prior to the isolation of endospores, it was possible to obtain a complete view on the presence of endospores both in and on the surface of the insect[16]. A complete view is necessary, since the insects considered are processed and consumed completely. On the other hand, the approach used in this study impedes the possibility to locate the microorganisms on or in the insect body. Likewise, it is not possible to track the origin of the microorganisms, i.e. whether they were introduced via the substrate, via the rearing environment or during packaging or transport.

The isolates collected were assigned to four different genera of endospore-forming bacteria, including *Bacillus*, *Lysinibacillus*, *Brevibacillus* and *Clostridium*. Based on further identification, the *Bacillus* genus was further divided into *Bacillus cereus* group (also *B. cereus* sensu lato (s.l.)) and *Bacillus* (non-cereus) (Figure 2A; Supplementary Table 1). Of the 50 isolates obtained from yellow mealworms, 20 (i.e. 40%) were identified as a member of the *Bacillus cereus* group. For house crickets, even 79 out of 92 isolates (86%) were identified as

B. cereus s.l. This may indicate that house crickets are highly susceptible to colonisation with *B. cereus* group members.

The *B. cereus* group consists, among a few recently described species, of the closely related bacteria *B. cereus* sensu stricto, *B. thuringiensis*, *B. weihenstephanensis*, *B. wiedmannii*, *B. anthracis*, *B. mycoides*, *B. pseudomycoides*, *B. cytotoxicus*, and *B. toyonensis*[18–21]. Most isolates assigned to the *B. cereus* group were identified as *B. cereus* s.s. or *B. paramycoides* (proposed novel member[22]). Due to the close genetic relationships within the *B. cereus* group, however, it is hard to distinguish between the different species compiled in this group based on the 16 S rRNA gene[21]. Several species of the *B. cereus* group are relevant with respect to human health, agriculture or food safety. In the case of edible insects, presence of the well-known pathogen *B. cereus* s.s. may pose a severe food safety risk for consumers, but also some *B. weihenstephanensis* and *B. cytotoxicus* strains were reported as potential human (foodborne) pathogens[23,24]. For insect rearing, *B. thuringiensis* may be harmful because of its insecticidal properties[25]. Whatever *B. cereus* group species were present in the insects, the *B. cereus* group poses an important threat for the edible insect sector and should be monitored. Also, given the fact that the *B. cereus* group contains psychrotrophic strains[20,26], chilled preservation of heat-treated insects can still allow the growth (and toxin production) of food pathogens such as *B. cereus* s.s. Mitigation strategies to lower the risks regarding pathogenic bacterial endospores include (i) the use of substrates that do not contain these species, which requires a thorough quality control, (ii) decontamination of the insects via the hurdle or combination strategy (e.g. steam, combining heat with pressure, pulsed electric fields, irradiation, …) or (iii) prevent spores from germinating (e.g. by reducing the water activity, pH and/or temperature of the matrix, …).

A few previous studies on the microbiota harboured by edible insects also report

members of the *B. cereus* group. For example, *B. cereus* s.s. and *B. weihenstephanensis* were

detected by PCR-DGGE (Denaturing Gradient Gel Electrophoresis) in edible processed insects

bought in Belgium and the Netherlands[10]. In another study[11], processed edible insects were

described to contain up to 6.6 log cfu/g presumptive *B. cereus*, of which *B. cytotoxicus*, *B.*

cereus s.s. and *B. thuringiensis* isolates could be identified using sequencing and biomolecular

identification.

The second most observed endospore-forming bacteria were members of the genus

Lysinibacillus (21 out of 142 isolates, or 15%; Figure 2A) and the majority was identified as *L.*

fusiformis (Supplementary Table 1). This spore former, renamed from *Bacillus fusiformis*[27],

rarely acts as a human pathogen, but especially its relative *L. sphaericus* has been reported as a

potent insect pathogen. *L. sphaericus* preferably targets mosquitos, but activity against other

insects species has been observed as well[28]. Furthermore, *Lysinibacillus* spp. were described as

potential proteolytic spoilage organisms that can thus alter product quality[29,30]. Also

previously[11], *Lysinibacillus* sp. was detected in edible insects.

Besides *B. cereus* members, also other, non-cereus *Bacillus* spp. were detected (16/142

isolates, 11%; Figure 2A). The genus *Bacillus* has regularly been found in edible

insects[6,10,11,31,32], but it has rarely been described at species level. According to the identification

results (Supplementary Table 1), non-cereus *Bacillus* isolates may correspond to several species

including *B. pumilus*, *B. altitudinis*, *B. siamensis*, *Bacillus licheniformis*, *B. vallismortis* and *B.*

subtilis. The latter three species are closely related and, comparably to the *B. cereus* group, they

are members of the *B. subtilis* group[33]. As was also the case for the *B. cereus* group, the 16 S

rRNA gene does not allow for species-level identification for *B. subtilis* group members[33]. *B.*

subtilis and its relatives are generally considered as spoilage organisms rather than human food

pathogens[34], but were linked to toxin production and a few cases of foodborne illness in the

past[35,36]. On the other hand, they are also frequently described as beneficial for plants and/or

animals e.g. as biocontrol organism or probiotic[37,38], which may pose opportunities for

industrial valorisation, also in the insect sector.

Six isolates (3%; Figure 2A) were ascribed to the genus *Brevibacillus* and identified as

either *B. laterosporus* or *B. halotolerans* (Supplementary Table 1). They were only obtained

from yellow mealworms and not from house crickets. In previous research[7], a *Brevibacillus* sp.

was detected by Illumina sequencing in yellow mealworms, in abundances up to 28%. *B.*

laterosporus has been reported as insect pathogen[39]. Together with *B. thuringiensis*, a member

of the *B. cereus* group, and *Lysinibacillus sphaericus*, a realistic risk for edible insect rearing

exists when these entomopathogenic bacteria are present. Further, *Brevibacillus* spp. can act as

spoilage organism and impact the quality of insect food products[29]. On the other hand, *B.*

laterosporus has also been described as a broad-spectrum antimicrobial species against

phytopathogenic bacteria and fungi[39], which may involve agricultural benefits.

While both aerobic and anaerobic bacterial endospore counts were performed and

isolates were picked randomly from all plates from all samples, it is striking that almost solely

aerobic and/or facultative anaerobic organisms were identified. (Strict) anaerobic spore formers

such as specific *Clostridium* spp. were encountered in edible insects before, including yellow

mealworms and house crickets[7–9,40]. In this study, however, only one isolated was assigned to

the genus *Clostridium*, yet, with a low sequence identity (69.4 %, Supplementary Table 1). The

genus *Clostridium* is of concern regarding food safety, since it contains the food pathogens *C.*

perfringens and *C. botulinum*. They may both produce potent toxins that cause gastroenteris

and botulism, respectively. Food poisoning caused by *C. perfringens*, however, generally

requires the ingestion of 10^8 cells that can then sporulate (and release their toxin) in the

intestine[41]. The total viable anaerobic counts observed in this study may exceed 8 log cfu/g, but

probably do not consist solely of *Clostridium* cells. A selective *Clostridium* count may be useful

to confirm this. All previous observations of (possible) anaerobic spore-forming bacteria were,

however, based on culture-independent methods. The random selection of isolates and/or the

strict anaerobic conditions necessary for cultivating certain *Clostridium* species may have

impeded to obtain anaerobic species that might have been present in the samples investigated.

Accordingly, the specific isolation and identification of anaerobic bacterial spore formers would

form an interesting addition to this research.

qPCR detection of B. cereus-related genes

It has been demonstrated that the 16 S rRNA gene is extremely similar (97.34 to 100%

inter-species similarity) between all *B. cereus* group members [20,42]. The 16 S rRNA gene-based

identification method applied in this study was therefore complemented by studying additional

genetic markers to further characterise and discriminate the *B. cereus* s.l. isolates. Firstly, all *B.*

cereus s.l. were subjected to a qPCR assay targeting the *panC* gene encoding a pantothenate-β-

alanine ligase[43]. The presence of this gene was used to confirm the assignment of the isolates

to the *B. cereus* group, since *panC* genotyping is a typical method to group *B. cereus* s.l. species

based on phenotypic characteristics[44]. Next, by targeting the *cesA* gene[45], each isolate was

assessed for plasmid presence encoding for the production of the heat-resistant cereulide toxin,

the most severe food safety risk[46,47]. This allows to evaluate the possibility for each isolate to

act as a human pathogen of the emetic type, regardless of its taxonomic classification.

Additionally, to date, this virulence plasmid was only encountered in *B. cereus* s.s. and *B.*

weihenstephanensis, while it was reported not to be present in a large collection of *B.*

thuringiensis, *B. mycoides* and *B. pseudomycoides* strains[18,47]. In the (edible) insect sector, the

distinction between foodborne pathogen (*B. cereus* s.s. or *B. weihenstephanensis*) and insect-

borne pathogen (*B. thuringiensis*) is of major interest, and presence or absence of the *cesA* gene

provides additional information in this regard.

Of the 98 *B. cereus* group isolates assessed for the *panC* gene, only three were found to

be negative. Consequently, the identification of 95 *B. cereus* group isolates was considered

reliable. Next, in 64 of the 98 isolates (65 %), the *cesA* gene was detected (Figure 2B). This

indicates that those isolates may be able to produce the cereulide toxin under suitable

conditions. Influencing factors are for instance temperature, the food matrix (e.g. water activity

(a_w)) and the number of *B. cereus* cells present[48,49]. While the precise impact of water activity

is not fully understood yet, a minimal a_w-value of 0.953 has been proposed for the production

of cereulide[50]. The water activity of raw edible mealworms and crickets, as were used in this

study, ranges between 0.95 and 0.98[16] and is consequently suitable for cereulide production.

The amount of *B. cereus* required to be present for toxin production to take place is typically

reported as 3 to 5 log cfu/g[18]. *B. cereus* counts were not assessed in this study, but in literature,

presumptive *B. cereus* were already detected up to 6.6 log cfu/g in processed edible mole

crickets[11].

The presence of the cereulide plasmid in *B. cereus* in its endospore form poses an

increased food safety risk. The number of bacterial endospores in the samples investigated was

maximally 4.2 log cfu/g and did not relate solely to *B. cereus* s.l. Despite this observation, a

risk is still present because after a heat treatment (which is survived by the spores and can also

even activate them), the endospores can germinate in the insect matrix, and in absence of

competition, multiply and produce cereulide. A second heating step will not be sufficient to

destroy the toxin and thus the risk. Consequently, the exact conditions for and the extent of

cereulide production that can occur in edible insects and derived food products is a future

research need.

Presence of foodborne viruses

Three different RT-qPCR assays were optimised to detect and quantify the RNA viruses

HAV, HEV and NoV GII in insect RNA extracts and assessed for their reliability. The addition

of an Internal Amplification Control (IAC) to the RT-qPCR assay has proven to be a robust

control in virus analysis in food matrices[51] and has become common practice[52,53]. Similarly, in

this study, IACs were included in all samples investigated and were systematically recovered

in the three assays (C_t values $\pm$ 35), while negative IAC controls showed no amplification after

40 cycles, proving the reactions to be successful. In order to quantify the viral titer, standard

curves were constructed and showed to be of good quality, according to their calculated

parameters R^2, slope and efficiency. R^2 was determined at 0.998 for HAV and 0.999 for HEV

and NoV GII. Next, the slope was calculated as -3.737, -3.287 and -3.625 for HAV, HEV and

NoV GII, respectively. Finally the efficiency showed values of 85.192%, 101.487% and

88.739% for HAV, HEV and NoV GII respectively. Based on the recovery of the reference

virus cDNA fragments, the detection limit for each assay was set at 100 copies/µl undiluted

RNA extract. Also the consistent amplification of positive controls (C_t values $\pm$ 31) and absence

of qPCR signal for negative controls contributed to the conclusion that the RT-qPCR assays

were reliable.

In none of the samples any of the viruses HAV, HEV or NoV GII could be detected,

meaning that the viruses were either not present or the samples contained less than 100 viral

RNA copies/µl extract. Consequently, the transmission risk for HAV, HEV and NoV GII via

yellow and lesser mealworm and (tropical) house cricket to humans is low. However, other

insect species and/or other foodborne viruses should still be investigated in future research, and

it should involve industrial samples.

Conclusions

In an extensive collection of industrially produced edible insects, this study uncovers a

large contrast between the high risk for foodborne illnesses originating from bacterial spores

and the low risk from human viruses in edible mealworms and crickets. For the bacterial

endospores, the largest fraction of isolates was identified as members of the *Bacillus cereus*

group, which contains foodborne and insect-borne pathogens. Sixty-five % of the *B. cereus*

group isolates contained the cereulide plasmid that can allow the production of a heat-resistant

toxin. The presence of the insect pathogens *Brevibacillus laterosporus* and *Lysinibacillus sp.,*

on the other hand, may involve economic risks for the insect rearer due to production losses.

Altogether, the possibility that batches are rejected due to microbiological problems may

increase, with concomitantly an increasing economic impact, since the sector is growing as

outlined above. Appropriate mitigation strategies to control the presence and outgrowth of

bacterial endospores should hence be adopted. In literature, limited information is available,

but insect producing and processing companies pay efforts in this field.

The presence of foodborne viruses in edible insects was investigated here for the first

time. Using methods shown to be reliable, hepatitis A virus, hepatitis E virus and norovirus

genogroup II were not detected (detection limit of 100 copies/µl extract). Hence, food safety

risks related to these viruses can be stated to be very low. However, as industrial insect

production methods are evolving due to automation and upscaling, virus detection is advised to

be continued and also performed for other species than those considered in this study.

Methods

Sample collection and preparation

For experiment 1 (identification of bacterial endospores in edible insects), four samples of living yellow mealworms (*Tenebrio molitor*) and four samples of living house crickets (*Acheta domesticus*) were collected at the end of their rearing cycle, without further processing. For each insect species, two different industrial rearing companies in Belgium (yellow mealworms) or Belgium and the Netherlands (house crickets) were sampled two times each (2 samples x 2 rearers x 2 sampling moments or batches). All living insect samples were packed by the producer in a closed tray or box and transported at ambient temperature.

For experiment 2 (detection of foodborne viruses in edible insects), the same 17 untreated (raw) mealworm and cricket samples that were investigated previously[7,16] for their bacterial composition (yellow mealworm, house cricket and tropical house cricket (*Gryllodes sigillatus)* samples from different rearing companies and batches) were employed, as well as an extra sample of lesser mealworms (*Alphitobius diaperinus*) obtained in another study[4] where it was used to study microbial dynamics during rearing (larvae day 35, post-harvest).

Samples were treated aseptically as from arrival in the laboratory. Prior to the analyses, dead specimens were removed with sterile tweezers and remaining insects were sedated by cooling them for approximately 1 h at 4 °C. Subsequently, the samples were homogenised by pulverisation with a sterilised hand-held mixer (Bosch CNHR 25, Belgium) as described earlier[9]. This way, the whole insect microbiota can be considered.

Bacterial endospore counts and isolation

All samples employed in experiment 1 were subjected to aerobic and anaerobic bacterial endospore counts in threefold. Following the procedure described earlier[16], for each analysis, 5

g of pulverised insects were diluted in 45 g peptone physiological salt solution (0.85% NaCl,

0.1% peptone, Biokar Diagnostics, Beauvais, France) and homogenised in a Bagmixer®

(Interscience, Saint Nom, France) for 1 min. This primary dilution was given a pasteurisation

treatment at 80 °C for 10 min and was then further diluted and plated on Plate Count Agar

(PCA, Biokar diagnostics) using the spread plate technique to easily pick colonies later. Both

aerobic and anaerobic endospore counts were determined after incubation for 48 h at 37 °C[54].

As a comparison, a similar but unpasteurised dilution series was used to determine the total

viable aerobic and anaerobic counts of the samples (pour plate technique, PCA, 72 h at 30 °C).

Anaerobic conditions were generated in Anaerocult containers (2.5 L, VWR International,

Leuven, Belgium) using 'AnaeroGen 2.5 L atmosphere generation systems' (Thermo Fisher

Scientific, Asse, Belgium) and evaluated using resazurin indicators (BR0055N, Thermo Fisher

Scientific).

For each sample, after incubation, several colonies with various morphologies were

picked from the countable plates and streaked on nutrient agar (NA, Biokar Diagnostics, 24 h

at 37 °C, (an)aerobic incubation depending on their origin) to form axenic cultures. An

individual colony was subsequently incubated overnight in nutrient broth (NB, Biokar

Diagnostics, 18 h at 37 °C) and stored at - 80 °C after addition of glycerol to a final

concentration of 50 % (v/v). This way, in total 67 and 95 spore-forming isolates were collected

from the yellow mealworm and house cricket samples, respectively.

Identification of bacterial endospores

All 162 endospore isolates were, after recultivation in NB, subjected to

phenol/chlorophorm genomic DNA extraction as described earlier[55] and subsequently to PCR

(T100™ Thermal Cycler, Bio-Rad, Temse, Belgium), amplifying the 16 S ribosomal RNA

(rRNA) gene. The PCR reaction (20 µl) contained 1.25 U of TaKaTa ExTaq Polymerase and 1

x ExTaq Buffer (Clontech Laboratories, Palo Alto, CA, USA), 312.5 µM of each dNTP, 1.0

µM of each primer (27F and 1492R, Table 1), and 5 ng genomic DNA (measured by a Nanodrop

spectrophotometer, Thermo Fisher Scientific). PCR conditions included initial denaturation for

2 min at 95 °C, followed by 34 cycles of 45 s denaturation at 95 °C, 45 s annealing at 59 °C

and 45 s elongation at 72 °C, and final elongation for 10 min at 72 °C. Obtained amplicons

were sequenced using the same reverse primer as used in the PCR (1492R, Table 1) by

Macrogen Europe (Amsterdam, the Netherlands). Resulting sequences were aligned and

trimmed to an average read length of 861 bp. Isolates were subsequently classified using the

EzBioCloud 16S rRNA gene database[56]. This way, a total of 142 isolates (50 from yellow

mealworms, 92 from house crickets) could be identified.

qPCR detection of B. cereus-related genes panC and cesA

For 99 isolates classified as *B. cereus* s.l., two additional genes were traced using qPCR.

For each extract, firstly, a qPCR assay targeting the *panC* gene was applied in order to confirm

the association with *B. cereus* s.l. Next, presence of the cereulide toxin pathway was

investigated by assessing the *cesA* gene. All qPCR reactions (10 µl) were executed in a

QuantStudio 3 qPCR system (Applied Biosystems, Thermo Fisher Scientific) and contained 5

µl PowerUp SYBR Green Master Mix with UNG (uracil-N glycosylase to prevent carryover

contamination; Thermo Fisher Scientific), 1 µM (*panC*) or 0.4 µM (*cesA*) of each primer (Table

1) and 60 ng of DNA (measured by a MySpec spectrophotometer, VWR International). qPCR

reaction conditions for the *panC* detection followed the master mix supplier's protocol: 2 min

at 50 °C (UNG activation) and 2 min of initial denaturation at 95 °C, followed by 40 cycles of

15 s denaturation at 95 °C and combined annealing and elongation for 1 min at 60 °C. For the

cesA detection, the same conditions were applied, except for the initial denaturation step which

was shortened to 20 seconds[57]. Each qPCR was followed by a melt curve analysis by raising

the temperature gradually from 60 to 95 °C (0.1 °C/s) while continuously monitoring

fluorescence. Positive (*B. cereus* emetic type, LMG 17603, Belgian Coordinated Collections of

Microorganisms (BCCM)) and negative (no template, sterile ultra-pure water instead) controls

were included. Data analysis was executed using the online application "Design and Analysis"

in the qPCR DataConnect cloud software (Thermo Fisher Scientific). Successful detection was

considered when the melt curve analysis revealed a single peak at the expected melting

temperature, and when obtained C_t values did not exceed these of the negative control (33 for

cesA and 35 for *panC*).

RT-qPCR quantification of selected viruses

In order to detect and quantify HAV, HEV and NoV GII, previously described

methods[52] were optimised and used to screen the insect samples in experiment 2. Standard

quantification curves and Internal Amplification Controls (IACs)[51] were included to assess

quantification and reliability. Corresponding virus reference cDNA was used as standard, either

obtained from an RNA transcript (NoV GII after RT-PCR, see below) or plasmids containing

virus genome fragments (HAV and HEV after PCR targeting the reference fragment). IAC

RNAs were generated by transforming IAC-containing plasmids into chemically competent

Escherichia coli cells (One Shot™ TOP10, Thermo Fisher Scientific) and transcription to RNA

using the T7 High Yield Transcription kit (Thermo Fisher Scientific).

Homogenised insect samples (experiment 2) were subjected in twofold to a total RNA

extraction, using the Mo Bio RNA PowerSoil kit (manufacturer's protocol, Carlsbad, CA,

USA). Next, each RNA extract was subjected to a two-step reverse transcription real-time PCR

(RT-qPCR) assay for each target virus separately. First, a tenfold RNA extract dilution was

used to translate target virus RNA into cDNA (Table 1). In this step, the IAC corresponding to

the target virus was added to exclude false negative results. Each reaction (20 µl) was performed

using the High Capacity cDNA Reverse Transcription kit (Applied Biosystems, Thermo Fisher

Scientific) and contained 2.0 µl RT Buffer, 1.0 µl MultiScribe™ Reverse Transcriptase, 2 µM

reverse primer (Table 1), 0.8 µl dNTP Mix (100 mM), 0.5 µl RNase Inhibitor, 1000 IAC copies,

and 40 ng RNA (as measured by MySpec spectrophotometer). Reaction conditions (iCycler

Thermal Cycler, Bio-Rad) were as described by the supplier: 10 min at 25 °C, 2 h at 37 °C, and

5 min at 85 °C. Positive (10^8 IAC copies) and negative (no template, sterile ultra-pure water

instead) controls were included for all virus assays. Next, for each sample, the obtained cDNA

was used as a template in the subsequent qPCR which aimed to detect and quantify HAV, HEV

or NoV GII as well as to detect the corresponding IAC at the same time. The three virus assays

were conducted based on protocols described earlier[52]. All analyses were performed in

duplicate and each reaction (20 µl) contained 1x TaqMan Universal Master Mix with UNG

(Applied Biosystems, Thermo Fisher Scientific), 0.25 µM of each primer (Table 1), 0.25 µM

virus probe (FAM-labelled, Table 1) and 0.10 µM IAC probe (VIC-labelled, Table 1). All

reactions were executed using a QuantStudio 3 qPCR system. qPCR conditions were as follows:

2 min at 50 °C, 10 min at 95 °C and 40 cycles of 15 s denaturation at 95 °C and 60 s combined

annealing and elongation at 60 °C. Virus quantities and standard curve parameters were

calculated in the qPCR DataConnect cloud software. Again, positive (1000 virus cDNA copies)

and negative (no template, sterile ultra-pure water instead) controls were included. Only when

the IAC was detected below the negative control C_t, the RT-qPCR assay was considered

successful.

Statistical analyses

To determine differences in microbial counts between samples in experiment 1, results were subjected to one-way ANOVA with all-pairs Tukey-Kramer HSD post hoc test. Normality and homoscedasticity assumptions were tested for the analysed data sets using the Shapiro-Wilk W test and Levene's test, respectively. In the cases that homoscedasticity was not confirmed, Welch's ANOVA was used instead. Results were displayed as a Mean Diamonds plots with proportional x-axis. JMP pro 14.0.0 (SAS Institute Inc., Cary, NC, USA) was used with significance level 0.05 for the statistical tests and creation of the plots.

Data availability

Sequencing data obtained in this study have been deposited in GenBank (National Center for Biotechnology Information, NCBI) with the accession codes MN508485 to MN508527 (Supplementary Table 1). All other relevant data are available from the corresponding author on request.

Acknowledgements

Virus references and IACs were kindly provided by I. Di Bartolo from the Italian Istituto Superiore di Sanità (ISS, Rome, Italy) and N. Cook from Fera Science Ltd. (York, United Kingdom). Additionally, the authors would like to thank A. Paeleman (Scientia Terrae Research Institute, Sint-Katelijne-Waver, Belgium) for her expertise and assistance in designing and optimising the qPCR protocols, S. Crauwels (KU Leuven, Leuven, Belgium) for processing the sequencing results and R. Smets (KU Leuven, Geel) for the statistical help. Also J. Franciotti, L. De Vrindt, N. Huybrechts, M. Gerits, S. Machtajiw, J. Plas and E. Van Vossole are acknowledged for their assistance in the lab. This research was financially supported by

Author Contributions

445   D.V. designed, prepared and executed all experiments, including sample collection and

446 preparation, microbiological analyses, DNA and RNA extractions and PCR and RT-qPCR

447 reactions. D.V. analysed the data, constructed tables and figures and wrote the main text. B.L.

448 and L.V. supervised the study, provided additional insight in data analysis and revised the

449 manuscript.

Competing Interests statement

452   The authors declare no competing interests.

References

1. EFSA Scientific Committee. Risk profile related to production and consumption of

insects as food and feed. *EFSA J. 2015* **13**, 60 pp. (2015).

2. Garofalo, C. *et al.* Current knowledge on the microbiota of edible insects intended for

human consumption: A state-of-the-art review. *Food Res. Int.* **125**, 108527 (2019).

3. Vandeweyer, D., Lenaerts, S., Callens, A. & Van Campenhout, L. Effect of blanching

followed by refrigerated storage or industrial microwave drying on the microbial load

of yellow mealworm larvae (*Tenebrio molitor*). *Food Control* **71**, 311–314 (2017).

4. Wynants, E. *et al.* Microbial dynamics during production of lesser mealworms

(*Alphitobius diaperinus*) for human consumption at industrial scale. *Food Microbiol.*

70, 181–191 (2018).

5. Vandeweyer, D. *et al.* Microbial Dynamics during Industrial Rearing, Processing, and Storage of Tropical House Crickets (*Gryllodes sigillatus*) for Human Consumption. *Appl. Environ. Microbiol.* **84**, AEM.00255-18 (2018).

6. Klunder, H. C., Wolkers-Rooijackers, J., Korpela, J. M. & Nout, M. J. R. Microbiological aspects of processing and storage of edible insects. *Food Control* **26**, 628–631 (2012).

7. Vandeweyer, D., Crauwels, S., Lievens, B. & Van Campenhout, L. Metagenetic analysis of the bacterial communities of edible insects from diverse production cycles at industrial rearing companies. *Int. J. Food Microbiol.* **261**, 11–18 (2017).

8. Garofalo, C. *et al.* The microbiota of marketed processed edible insects as revealed by high-throughput sequencing. *Food Microbiol.* **62**, 15–22 (2017).

9. Stoops, J. *et al.* Microbial community assessment of mealworm larvae (*Tenebrio molitor*) and grasshoppers (*Locusta migratoria migratorioides*) sold for human consumption. *Food Microbiol.* **53**, **Part B**, 122–127 (2016).

10. Osimani, A. *et al.* Insight into the proximate composition and microbial diversity of edible insects marketed in the European Union. *Eur. Food Res. Technol.* **243**, 1157–1171 (2017).

11. Fasolato, L. *et al.* Edible processed insects from e-commerce: Food safety with a focus on the Bacillus cereus group. *Food Microbiol.* **76**, 296–303 (2018).

12. Stoops, J. *et al.* Minced meat-like products from mealworm larvae (*Tenebrio molitor* and *Alphitobius diaperinus*): Microbial dynamics during production and storage. *Innov. Food Sci. Emerg. Technol.* **41**, 1–9 (2017).

13. Petrović, T. & D'Agostino, M. Viral Contamination of Food. in *Antimicrobial Food*

Packaging (ed. Barros-Velázquez, J.) 65–79 (Academic Press, 2016).

doi:10.1016/B978-0-12-800723-5.00005-X

14. Da Silva, A. K. *et al.* Evaluation of removal of noroviruses during wastewater

treatment, using real-time reverse transcription-PCR: Different behaviors of

genogroups I and II. *Appl. Environ. Microbiol.* **73**, 7891–7897 (2007).

15. De Smet, J. *et al.* Stability assessment and laboratory scale fermentation of pastes

produced on a pilot scale from mealworms (*Tenebrio molitor*). *LWT* **102**, 113–121

(2019).

16. Vandeweyer, D., Crauwels, S., Lievens, B. & Van Campenhout, L. Microbial counts of

mealworm larvae (*Tenebrio molitor*) and crickets (*Acheta domesticus* and *Gryllodes*

sigillatus) from different rearing companies and different production batches. *Int. J.*

Food Microbiol. **242**, 13–18 (2017).

17. FASFC. Appendix: Table with regulatory criteria and action limits. *Action limits for*

microbiological contaminants in foods (2019). Available at:

http://www.favv.be/thematischepublicaties/_documents/Narval_microbio_VCT_JWS_

ADK__CKS_27_feb_2019_nl.xlsx.

18. EFSA Panel on Biological Hazards. Risks for public health related to the presence of

Bacillus cereus and other *Bacillus* spp. including *Bacillus thuringiensis* in foodstuffs.

EFSA J. **14**, 93 (2016).

19. Rasigade, J. P., Hollandt, F. & Wirth, T. Genes under positive selection in the core

genome of pathogenic *Bacillus cereus* group members. *Infect. Genet. Evol.* **65**, 55–64

(2018).

20. Liu, Y. *et al.* Genomic insights into the taxonomic status of the *Bacillus cereus* group.

Sci. Rep. **5**, 14082 (2015).

21.   Ehling-Schulz, M., Lereclus, D. & Koehler, T. M. The *Bacillus cereus* Group: *Bacillus*

Species with Pathogenic Potential. *Microbiol. Spectr.* **7**, (2019).

22.   Liu, Y. *et al.* Proposal of nine novel species of the *Bacillus cereus* group. *Int. J. Syst.*

Evol. Microbiol. **67**, 2499–2508 (2017).

23.   Guinebretière, M.-H. *et al. Bacillus cytotoxicus* sp. nov. is a novel thermotolerant

species of the *Bacillus cereus* group occasionally associated with food poisoning. *Int. J.*

Syst. Evol. Microbiol. **63**, 31–40 (2013).

24.   Stenfors, L. P., Mayr, R., Scherer, S. & Granum, E. Pathogenic potential of fifty

Bacillus weihenstephanensis strains. *FEMS Microbiol. Lett.* **215**, 47–51 (2002).

25.   Bravo, A., Gill, S. S. & Soberón, M. Mode of action of *Bacillus thuringiensis* Cry and

Cyt toxins and their potential for insect control. *Toxicon* **49**, 423–435 (2007).

26.   Martínez, S., Borrajo, R., Franco, I. & Carballo, J. Effect of environmental parameters

on growth kinetics of *Bacillus cereus* (ATCC 7004) after mild heat treatment. *Int. J.*

Food Microbiol. **117**, 223–7 (2007).

27.   Ahmed, I., Yokota, A., Yamazoe, A. & Fujiwara, T. Proposal of *Lysinibacillus*

boronitolerans gen. nov. sp. nov., and transfer of *Bacillus fusiformis* to *Lysinibacillus*

fusiformis comb. nov. and *Bacillus sphaericus* to *Lysinibacillus sphaericus* comb. nov.

Int. J. Syst. Evol. Microbiol. **57**, 1117–1125 (2007).

28.   Berry, C. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. *J. Invertebr.*

Pathol. **109**, 1–10 (2012).

29.   Lücking, G., Stoeckel, M., Atamer, Z., Hinrichs, J. & Ehling-Schulz, M.

Characterization of aerobic spore-forming bacteria associated with industrial dairy

processing environments and product spoilage. *Int. J. Food Microbiol.* **166**, 270–279

(2013).

30.   De Jonghe, V. *et al.* Toxinogenic and spoilage potential of aerobic spore-formers

isolated from raw milk. *Int. J. Food Microbiol.* **136**, 318–325 (2010).

31.   Wang, Y. & Zhang, Y. Investigation of gut-associated bacteria in *Tenebrio molitor*

(Coleoptera: Tenebrionidae) larvae using culture-dependent and DGGE methods. *Ann.*

Entomol. Soc. Am. **108**, 941–949 (2015).

32.   Osimani, A. *et al.* The bacterial biota of laboratory-reared edible mealworms (*Tenebrio*

molitor L.): From feed to frass. *Int. J. Food Microbiol.* **272**, 49–60 (2018).

33.   Wang, L. T., Lee, F. L., Tai, C. J. & Kasai, H. Comparison of *gyrB* gene sequences,

16S rRNA gene sequences and DNA-DNA hybridization in the *Bacillus subtilis* group.

Int. J. Syst. Evol. Microbiol. **57**, 1846–1850 (2007).

34.   De Vos, P. *et al. Bergey's Manual of Systematic Bacteriology Volume 3*. (Springer

Science+Business Media, 2009). doi:10.1007/978-0-387-68489-5

35.   Kramer, J. M. & Gilbert, R. J. *Bacillus cereus* and other *Bacillus* species. in *Foodborne*

Bacterial Pathogens (ed. Doyle, M.) 21–70 (Marcel Dekker, 1989).

36.   From, C., Pukall, R., Schumann, P., Hormazábal, V. & Granum, P. E. Toxin-producing

ability among *Bacillus* spp. outside the *Bacillus cereus* group. *Appl. Environ.*

Microbiol. **71**, 1178–1183 (2005).

37.   Serrano, L., Manker, D., Brandi, F. & Cali, T. The Use of *Bacillus subtilis* QST 713

and *Bacillus pumilus* QST 2808 as Protectant Fungicides in Conventional Application

Programs for Black Leaf Streak Control. *Acta Hortic.* **986**, 149–156 (2013).

38.   Tactacan, G. B., Schmidt, J. K., Miille, M. J. & Jimenez, D. R. A *Bacillus subtilis*

(QST 713) spore-based probiotic for necrotic enteritis control in broiler chickens. *J.*

Appl. Poult. Res. **22**, 825–831 (2013).

39.   Ruiu, L. *Brevibacillus laterosporus*, a Pathogen of Invertebrates and a Broad-Spectrum

Antimicrobial Species. *Insects* **4**, 476–492 (2013).

40.   Osimani, A. *et al.* Occurrence of transferable antibiotic resistances in commercialized

ready-to-eat mealworms (*Tenebrio molitor* L.). *Int. J. Food Microbiol.* **263**, 38–46

(2017).

41.   Madigan, M. T., Martinko, J. M., Dunlap, P. V & Clark, D. P. *Brock Biology of*

microorganisms. (Pearson/Benjamin Cummings, 2009).

42.   Ceuppens, S., Boon, N. & Uyttendaele, M. Diversity of *Bacillus cereus* group strains is

reflected in their broad range of pathogenicity and diverse ecological lifestyles. *FEMS*

Microbiology Ecology **84**, 433–450 (2013).

43.   Zhuang, K. *et al.* Typing and evaluating heat resistance of *Bacillus cereus* sensu stricto

isolated from the processing environment of powdered infant formula. *J. Dairy Sci.*

102, 7781–7793 (2019).

44.   Guinebretière, M. H. *et al.* Ecological diversification in the *Bacillus cereus* Group.

Environ. Microbiol. **10**, 851–865 (2008).

45.   Fricker, M., Messelhäußer, U., Busch, U., Scherer, S. & Ehling-Schulz, M. Diagnostic

real-time PCR assays for the detection of emetic *Bacillus cereus* strains in foods and

recent food-borne outbreaks. *Appl. Environ. Microbiol.* **73**, 1892–1898 (2007).

46.   Ehling-Schulz, M., Frenzel, E. & Gohar, M. Food-bacteria interplay: Pathometabolism

of emetic *Bacillus cereus*. *Front. Microbiol.* **6**, 704 (2015).

47.   Hoton, F. M. *et al.* Family portrait of *Bacillus cereus* and *Bacillus weihenstephanensis*

cereulide-producing strains. *Environ. Microbiol. Rep.* **1**, 177–183 (2009).

48.   Agata, N., Ohta, M. & Yokoyama, K. Production of *Bacillus cereus* emetic toxin

(cereulide) in various foods. *Int. J. Food Microbiol.* **73**, 23–27 (2002).

49.   Finlay, W. J. J., Logan, N. A. & Sutherland, A. D. *Bacillus cereus* produces most

emetic toxin at lower temperatures. *Lett. Appl. Microbiol.* **31**, 385–389 (2000).

50.   Rouzeau-Szynalski, K., Stollewerk, K., Messelhäusser, U. & Ehling-Schulz, M. Why

be serious about emetic *Bacillus cereus*: Cereulide production and industrial

challenges. *Food Microbiol.* **85**, 103279 (2020).

51.   Diez-Valcarce, M., Kovač, K., Cook, N., Rodríguez-Lázaro, D. & Hernández, M.

Construction and Analytical Application of Internal Amplification Controls (IAC) for

Detection of Food Supply Chain-Relevant Viruses by Real-Time PCR-Based Assays.

Food Anal. Methods **4**, 437–445 (2011).

52.   Diez-Valcarce, M. *et al.* Occurrence of Human Enteric Viruses in Commercial Mussels

at Retail Level in Three European Countries. *Food Environ. Virol.* **4**, 73–80 (2012).

53.   Di Bartolo, I., Angeloni, G., Ponterio, E., Ostanello, F. & Ruggeri, F. M. Detection of

hepatitis E virus in pork liver sausages. *Int. J. Food Microbiol.* **193**, 29–33 (2015).

54.   Dijk, R. *et al. Microbiologie van voedingsmiddelen: Methoden, principes en criteria*.

(MYbusinessmedia, 2015).

55.   Lievens, B. *et al.* Design and development of a DNA array for rapid detection and

identification of multiple tomato vascular wilt pathogens. *FEMS Microbiol. Lett.* **223**,

113–122 (2003).

56.   Yoon, S. H. *et al.* Introducing EzBioCloud: A taxonomically united database of 16S

rRNA gene sequences and whole-genome assemblies. *Int. J. Syst. Evol. Microbiol.* **67**,

1613–1617 (2017).

57.   Ueda, S., Yamaguchi, M., Iwase, M. & Kuwabara, Y. Detection of Emetic *Bacillus*

cereus by Real-Time PCR in Foods. *Biocontrol Sci.* **18**, 227–232 (2013).

Figures

Figure 1 Mean Diamonds plots representing the microbial counts in log cfu/g of the samples investigated in experiment 1. Figure 1A and 1B display the total viable aerobic and anaerobic counts, respectively, Figure 1C and 1D display aerobic and anaerobic bacterial endospore counts, respectively. ● and ▲ represent the three replicates of yellow mealworm (YM) and house cricket (HC) samples, respectively. Sample names consist of the insect species followed by the rearing company number and the batch number. Anaerobic bacterial endospore counts for YM 1.1 were not determined. The horizontal black line represents the mean of all samples. The top and bottom of each diamond represent the 100% confidence interval for each sample and the line across the middle of each diamond represents the sample mean. 95% confidence marks appear as lines above and below the sample mean.

[a,b,c]Mean counts from different samples with the same superscript are not statistically different (p > 0.05). [$]Only two replicates were displayed.

Figure 2 (2A) Number of bacterial endospore isolates from yellow mealworms and house crickets corresponding to the bacterial genera identified by 16 S rRNA gene sequencing and (2B) presence of the *cesA* gene, *in B. cereus* group isolates, as determined by qPCR.

 Tables

Table 1 Primers and probes employed in this study.

Target	Oligonucleotide (name)	Sequence (5' – 3')
16 S rRNA gene	Forward primer (27F)	AGA GTT TGA TCC TGG CTC AG
	Reverse primer (1492R)	TAC GGY TAC CTT GTT ACG ACT T
B. cereus genes		
panC[43]	Forward primer (panCF)	TYG GTT TTG TYC AAC RA TGG
	Reverse primer (panCR)	CAT AAT CTA CAG TGC CTT TCG
cesA[45]	Forward primer (cesAF)	CAC GCC GAA AGT GAT TAT ACC AA
	Reverse primer (cesAR)	CAC GAT AAA ACC ACT GAG ATA GTG
Viruses		
HAV[52]	Forward primer (HAV68)	TCA CCG CCG TTT GCC TAG
	Reverse primer (HAV240)	GGA GAG CCC TGG AAG AAA G
	Probe (HAV150)	6FAM-CCT GAA CCT GCA GGA ATT AA-MGBNFQ*
HEV[52]	Forward primer (JVHEVF)	GGT GGT TTC TGG GGT GAC
	Reverse primer (JVHEVR)	AGG GGT TGG TTG GAT GAA
	Probe (JVHEVP)	6FAM-TGA TTC TCA GCC CTT CGC-BHQ*
NoV GII[52]	Forward primer (QNIF2d)	ATG TTC AGR TGG ATG AGR TTC TCW GA
	Reverse primer (COG2R)	TCG ACG CCA TCT TCA TTC ACA
	Probe (QNIFS)	6FAM-AGC ACG TGG GAG GGC GAT CG-BHQ*
IAC[51]	IAC probe (PrfAP)	VIC-CCA TAC ACA TAG GTC AGG-MGBNFQ*

 *6FAM: 6-carboxyfluorescein, MGB-NFQ: minor groove binder non-fluorescent quencher, BHQ:
black hole quencher, VIC: 2'-chloro-7'-phenyl-1,4-dichloro-6-carboxyfluorescein.