Different immunomodulatory role of Ly6G[hi] and Ly6G[int] neutrophils during antiviral antibody therapy

Authors:

Jennifer Lambour[1], Mar Naranjo-Gomez[1,2], Myriam Boyer-Clavel[3] and Mireia Pelegrin[1,2] *

Address:

[1] IGMM, Univ Montpellier, CNRS, Montpellier, France
[2] Current address: IRMB, Univ Montpellier, INSERM, CNRS, Montpellier, France
[3] Montpellier Ressources Imagerie, Biocampus, Univ Montpellier, CNRS, Montpellier, France

*Corresponding author and Lead contact:

Mireia Pelegrin

Institute of Molecular Genetics of Montpellier

1919, route de Mende

34293 Montpellier Cedex 5

France

mireia.pelegrin@inserm.fr

Keywords : antiviral immune responses, monoclonal antibodies, immunotherapy, immunomodulation, vaccinal effects, immune complexes, neutrophils heterogeneity, monocytes, FcγR.

SUMMARY

Antiviral monoclonal antibodies (mAbs) can generate protective immunity through immune complexes (IC)-FcγRs interactions. We have shown the essential role of neutrophils in mAb-induced immunity of retrovirus-infected mice. Using this model, here we addressed how viral infection, with or without mAb therapy, affects neutrophils' functional activation. We found that neutrophils activated by viral ICs secreted high levels of chemokines able to recruit monocytes and neutrophils themselves. Moreover, inflammatory cytokines potentiated chemokines and cytokines release by IC-activated cells and induced FcγRs upregulation. Similarly, infection and mAb-treatment upregulated FcγRs expression on neutrophils and enhanced their cytokines and chemokines secretion. FcγRs upregulation allowed to identify *in vivo* two splenic neutrophils subpopulations with distinct phenotypic and functional properties that differentially and sequentially collaborate with inflammatory monocytes to induce Th1-type responses in mAb-treated mice. Our work provides novel findings on the heterogeneity and the immunomodulatory role of neutrophils in the enhancement of immune responses upon antiviral mAb therapy.

INTRODUCTION

The development of powerful antiviral monoclonal antibodies (mAbs) has provided new therapeutic opportunities to treat severe viral infections (Dibo et al., 2019; Hooft van Huijsduijnen et al., 2020; Salazar et al., 2017; Wang and Zhang, 2020). Fc-dependent mechanisms are crucial for efficient antiviral activity of neutralizing mAbs through the engagement of IgG receptors (FcγRs) expressed on immune cells. These Fc-FcγR interactions lead to the elimination of viral particles and virus-infected cells through phagocytic and cytotoxic mechanisms (i.e. antibody-dependent cellular phagocytosis (ADCP), antibody-dependent cell-mediated cytotoxicity (ADCC),...) (Bournazos and Ravetch, 2017a; Lu et al., 2018). Moreover, studies in different animal models of viral infection, including ours, have provided evidence that mAbs can also enhance antiviral immune responses (so called "vaccinal effects") in a Fc-dependent manner (Pelegrin et al., 2015). These vaccinal effects have been recently reported in HIV-infected patients treated with broadly neutralizing mAb (bnAbs) (Naranjo-Gomez and Pelegrin, 2019; Niessl et al., 2020; Schoofs et al., 2016) although the mechanisms involved have not been identified thus far. The elucidation of the molecular and cellular mechanisms driving Fc-dependent, mAb-mediated immunomodulation is therefore an important issue that will be key to achieving protective immunity against severe viral infections by mAbs.

While several Fc-mediated effector functions (i.e. ADCC, ADCP, ...) have been shown to be required for antibody-mediated antiviral protection (Bournazos et al., 2014, 2019; Earnest et al., 2019; Fox et al., 2019; Gunn et al., 2018; Liu et al., 2017) whether and how FcγR engagement by antiviral mAbs affects the immunomodulatory properties of different FcγR-expressing cells (i.e. cytokines/chemokines secretion, activation markers expression, …) has been little studied. In addition, the specific contribution of different FcγRs-expressing cells in the induction of vaccinal effects by mAbs still remains ill-understood. These issues are all the more important to address given that both virus-driven inflammation and Fc-mediated clearance of opsonized virus/infected cells by immune effector cells generate multiple danger signals able to further modulate the host immune response. However, multiple restrictions (i.e. technical and ethical issues, costs, …) largely limit those studies in humans and non-human primates (NHP). As an alternative, *in vivo* studies in immunocompetent mice infected with the Murine Leukemia Virus FrCasE allowed the identification of several immunological mechanisms that drive protective immunity upon mAb therapy (Lambour et al., 2016; Pelegrin et al., 2015). We have previously showed that treatment of FrCasE-infected mice with the neutralizing mAb 667 (recognizing the retroviral envelope glycoprotein, expressed in virions and infected cells) elicits protective adaptive antiviral immunity through the engagement of FcγRs. Notably, mAbs form immune complexes (ICs) with viral determinants that enhance antiviral T-cell responses through FcγR-mediated binding to dendritic cells (DCs) (Celis and Chang, 1984; Michaud et al., 2010; Nasser et al., 2010; Yamamoto et al., 2009). We have recently shown a key

immunomodulatory role of neutrophils in the induction of protective humoral responses *via* the
acquisition of B-cell helper functions upon engagement of FcγRs by the therapeutic mAb (Naranjo-
Gomez et al., 2018). Thus, the immunomodulatory actions of mAbs rely on their Fc fragment binding
to multiple FcγR-expressing cells such as DCs and neutrophils. While the role of IC-activated DCs in
the enhancement of antiviral immune responses has been addressed in several studies (Lambour et al.,
2016; Wang et al., 2019; Wen et al., 2016) the role of IC-activated neutrophils has mostly been
overlooked. Evidence shows that neutrophils, in addition to being key effector cells to fight against
invading pathogens, are also endowed with immunomodulatory properties through the secretion of a
plethora of chemokines and cytokines (Mantovani et al., 2011; Tamassia et al., 2018; Tecchio and
Cassatella, 2016). Yet, the functional activation of neutrophils by viral ICs and the resulting effect on
their immunomodulatory properties have poorly been studied in the context of antiviral mAbs
therapies. Furthermore, recent studies have highlighted a vast heterogeneity of neutrophils phenotypes
and functions in both homeostatic and pathological conditions (Deniset and Kubes, 2018; Ng et al.,
2019; Scapini et al., 2016; Silvestre-Roig et al., 2016, 2019). As these phenotypic and functional
variants of neutrophils depend on their context-dependent stimulation (inflammatory environment,
timing, disease progression, …), it is important to dissect how the viral infection and mAb therapy
shape the phenotype and functional properties of neutrophils.

Here, we used the FrCasE retroviral model to address how viral infection, with or without mAb
therapy, affects the phenotypical and functional activation of neutrophils. Neutrophils activated by
viral determinants secreted high levels of monocytes- and neutrophils-recruiting chemokines. We have
shown that the inflammatory environment resulting from the ongoing infection and mAb-treatment
modulates the functional properties of neutrophils and monocytes. We have also identified two
neutrophils subpopulations with distinct functional activation states that differentially participate in the
induction of Th1-type responses in mAb-treated mice in collaboration with inflammatory monocytes.
Our work provides hitherto unreported findings on both the heterogeneity and immunomodulatory
properties of neutrophils in a context of antiviral mAb-therapy. These findings might help to improve
mAb-based antiviral therapies by tailoring therapeutic interventions aiming at harnessing the
immunomodulatory properties of neutrophils.

RESULTS

Neutrophils activated by viral determinants, free or in the form of ICs, secrete high levels of chemokines able to recruit monocytes and neutrophils. We have previously shown a key role of neutrophils in the induction of long-term protective antiviral immunity upon mAb therapy of infected mice (Naranjo-Gomez et al., 2018). To better characterize the phenotypical and functional activation of neutrophils by viral determinants, free or in the form of ICs, we isolated bone marrow (BM) neutrophils from naive mice and stimulated them for 24h *in vitro* with free FrCasE virions or opsonized with the 667 mAb (ICs) (Figure 1A). Free 667 mAb was used as control. Both, FrCasE virions and ICs induced a strong activation of neutrophils as shown by a higher expression of the CD11b molecule as well as an increased frequency of CD11b^{hi} CD62L^{lo} neutrophils (Figure 1B). However, IC-mediated phenotypic activation was significantly higher.

We next assessed the functional activation of virus- and IC-stimulated neutrophils by measuring their chemokines and cytokines secretion capacity. Both stimuli poorly induced the secretion of the 12 cytokines analyzed, with no significant differences between unstimulated neutrophils *versus* virus or IC-stimulated neutrophils (Supplemental Figure 1A). This contrasted with the high secretion of IL-6 and TNFα pro-inflammatory cytokines observed upon lypopolisaccharide (LPS) neutrophils stimulation despite similar CD11b upregulation induced by LPS and viral determinants (Supplemental Figures 1B-1C). These data show that bacterial-related pathogen-associated molecular patterns (PAMPs) induce a functional activation of neutrophils different from that of viral stimuli. In contrast to cytokine secretion, both virus and ICs lead to high secretion levels of several chemokines such as CCL2, CXCL1, CXCL5 and to a lesser extent CCL3 and CCL4 (Figure 1C). In addition, LPS-stimulated neutrophils only secreted a low amount of CCL3 and CCL4 chemokines, with no secretion of CCL2, CXCL1, CXCL5 chemokines thus providing further evidence of a pathogen-dependent secretion profile of neutrophils (Supplemental Figure 1D). Interestingly, the chemokines which were more strongly secreted by neutrophils upon viral stimuli (but not by LPS) have been shown to be involved in the recruitment of monocytes (CCL2) (Fan et al., 2018; Kim et al., 2011; Leonard and Yoshimura, 1990; Robben et al., 2005) and neutrophils themselves (CXCL1, CXCL5) (Haider et al., 2019; Pruenster et al., 2009; Ritzman et al., 2010; Rovai et al., 1998).

Inflammatory monocytes activated by viral determinants, free or in the form of ICs, secrete high levels of chemokines able to recruit neutrophils and monocytes. Taking into account the high amounts of the monocytes-recruiting CCL2 chemokine secreted by viral determinants-activated neutrophils, we next assessed the phenotypical and functional activation of monocytes by these stimuli. Ly6C^{hi} monocytes were isolated from naive mice BM and stimulated for 24h with virus or ICs (Figure 2A). Phenotypical activation was evaluated by measuring the CD86 co-stimulatory molecule expression on the cell

surface. Free virions and ICs significantly increased CD86 expression but no significant differences were observed between both stimuli (Figure 2B).

Virus- and ICs induced a weak secretion of most of the 12 cytokines analyzed, except for IL6 and TNFα. However, no significant differences between unstimulated monocytes *versus* virus- or IC-stimulated monocytes were detected (Supplemental Figure 2A). This contrasted with high level secretion of IL-6, TNFα and IFNγ observed upon LPS stimulation (Supplemental Figures 2B-2D). The quantification of the chemokine release by activated monocytes showed some interesting similarities with neutrophils. Following virus or ICs stimulation, monocytes produced CXCL5, CXCL1 and CCL2, with no significant differences between virus *versus* ICs (Figure 2C). As compared to neutrophils, higher amounts of the neutrophil-recruiting chemokine CXCL1 were detected as well as lower amounts of CXCL5 and CCL2. Interestingly, LPS stimulation induced a wider and different panel of chemokine release (Supplemental Figure 2D), notably with the secretion of high amounts of CCL3, CCL4, CCL5, and to a lesser extent CXCL1 and CXCL10. As observed in neutrophils, these data show that viral stimuli induce a functional activation of monocytes different from that of bacterial-related PAMPs.

Inflammatory conditions potentiate the activation of neutrophils and monocytes by viral ICs. As the inflammatory microenvironment resulting from the viral infection and mAb therapy might affect the antiviral immune response, we next assessed the phenotypic and functional activation of neutrophils and monocytes by viruses and ICs in the presence of proinflammatory/immunomodulatory cytokines, such as TNFα, IFN-I and IFNγ.

As depicted by an increased expression of CD11b, TNFα and IFNγ significantly enhanced the phenotypic activation of neutrophils by ICs (but not by free virus) (Figure 3A). Priming conditions also modulated the secretion profile of virus- and IC-activated cells in a cytokine-specific manner. TNFα enhanced the secretion of CXCL1 and TNFα itself by IC-activated neutrophils but not by virus-activated neutrophils (Figure 3B, Supplemental figure 3A). IFN-I and IFNγ (Figure 3B, Supplemental figure 3A) significantly potentiated CCL4 secretion by IC-activated neutrophils. Both types of IFN induced the production of CXCL10, consistent with the IFN-dependent induction of this chemokine. However, CXCL10 secretion was significantly increased in neutrophils activated by virus or ICs upon IFN priming. IFNγ also significantly enhanced CCL5 and CXCL1 secretion by IC-stimulated neutrophils. In contrast, TNFα-, IFN-I- or IFNγ priming did not modulate the secretion of CCL2, CXCL5 and CCL3 by neither virus- nor IC-activated neutrophils (not shown).

As for monocytes, IFN-I and IFNγ significantly potentiated the phenotypic activation of IC-stimulated

monocytes (as depicted by an increased expression of CD86 molecule) (Figure 3C) but not that of

virus-activated cells. TNFα, IFN-I or IFNγ significantly potentiated the secretion of CCL2 and

CXCL1 (Figures 3D and Supplemental figures 3B) by IC-activated monocytes. Similar to neutrophils,

IFNγ and IFN-I enhanced the secretion of CXCL10 by IC-stimulated monocytes and IFN-I

significantly increased the production of TNFα. Priming conditions also modulated the

cytokine/chemokine secretion profile of virus-activated monocytes, but in a more restricted way than

in IC-activated cells. Thus, CXCL1 secretion was enhanced by TNFα and IFNγ and CCL2 by IFN-I.

Both types of IFN also increased the release of CXCL10 by virus-activated monocytes (Figure 3D).

Similar secretion levels of CXCL5 virus- and IC-activated monocytes were observed in the presence

or in the absence of TNFα-, IFN-I- and IFNγ- priming (not shown).

Altogether, these data highlight that inflammatory conditions strongly potentiate the functional

activation of IC-stimulated neutrophils and monocytes, although with different immune outcomes

depending on the priming condition and the responding cell type (Figure 3). Our data describe a

distinct enhancement of neutrophils and monocytes secretion profile by TNFα-, IFN-I- and IFNγ-

priming showing thus a cytokine-specific effect on the functional activation of these IC-stimulated

cells. Such cytokine/chemokine secretion enhancement is also cell type specific (i.e. increased

secretion of CCL4, CCL5, CXCL1 and TNFα by IC-activated neutrophils and CCL2, CXCL1 and

TNFα by IC-activated monocytes). In contrast, inflammatory conditions hardly modify the secretion

profile of free virus-activated cells, except for the secretion of the CXCL10 IFN-inducible chemokine.

Inflammatory conditions upregulate the expression of FcγRIV on in vitro activated neutrophils and

monocytes. We next assessed whether the activation of neutrophils and monocytes by virus and IC

affected their FcγRs expression, both in the absence and in the presence of inflammatory cytokines.

Indeed, FcγRs expression might differ between steady-state *versus* inflammatory/pathological

conditions (Wang and Jönsson, 2019), however little is known about modulation of FcγRs expression

in the context of viral infections and mAb-therapy. Neither virus nor IC upregulated the expression of

FcγRIV on neutrophils and monocytes (Figure 4A-4D). On the contrary, IFNγ and IFN-I priming (but

not TNFα) upregulated the expression of FcγRIV on neutrophils (IFNγ) (Figure 4B) and monocytes

(IFNγ and IFN-I) (Figure 4D). Both IFNs also enhanced FcγRI expression on monocytes whereas a

decreased FcγRII-III expression was observed following IC stimulation and cytokine-priming

(Supplemental Figure 4). No FcγRI expression was detected on neutrophils neither upon viral

determinants stimulation nor cytokine priming.

Viral infection and mAb therapy upregulate FcγRIV expression on two neutrophils subpopulations and on inflammatory monocytes. We next assessed *in vivo* whether the inflammatory environment resulting from FrCasE viral infection and 667 mAb therapy modulated the expression of FcγRIV. The modulation of this FcγR is all the more relevant to be studied in this experimental model as (i) it is a high affinity receptor for IgG2a, (which is the isotype of the 667 mAb) and (ii) it is strongly expressed in neutrophils, the latter having a key immunomodulatory role in mAb-mediated protection of retrovirus-infected mice (Naranjo-Gomez et al., 2018).

To this end, mice were infected and treated, or not, with the therapeutic mAb (infected/treated and infected/non-treated, respectively). Then, the expression of FcγRIV on neutrophils and inflammatory monocytes from the spleen (one of the main sites of viral replication) was evaluated at different time points post-infection (p.i.). Age-matched naive mice were used as controls. The cell populations of interest were defined by flow cytometry (Figure 5A) based on the expression of CD11b, Ly6G and Ly6C to gate neutrophils (CD11b$^+$, Ly6G^{hi}) and inflammatory monocytes (CD11b$^+$Ly6G$^-$Ly6C^{hi}). Interestingly, we identified another population of neutrophils displaying a lower expression of the cell surface marker Ly6G (CD11b$^+$Ly6G^{int}) (Figure 5A). No expression of the eosinophils cell surface marker Siglec F was detected in this population (not shown). The frequency of both populations of neutrophils CD11b$^+$Ly6G^{hi} and CD11b$^+$Ly6G^{int} was significantly increased in the spleen of infected/non-treated mice at 8 and 14 days p.i. This increased neutrophil frequency was associated with a higher percentage of spleen infected cells as compared to infected/treated mice (Naranjo-Gomez et al., 2018). On the contrary, the frequency of inflammatory monocytes (CD11b$^+$Ly6G$^-$Ly6C^{hi}) remained unchanged in infected mice at day 8 p.i. but was significantly increased at day 14 p.i. (Figure 5B), suggesting an earlier recruitment of neutrophil than inflammatory monocytes upon infection.

In steady-state conditions, FcγRIV was highly expressed on CD11b$^+$Ly6G^{hi} and CD11b$^+$Ly6G^{int} neutrophils. Lower FcγRIV expression was detected in inflammatory monocytes at the different time points assessed (Figure 5C). Notably, upon viral infection and mAb-treatment, FcγRIV expression was upregulated in the three myeloid cells populations although, to a different magnitude depending on the cell type, time p.i. and mAb-treatment (Figure 5C). At day 8 p.i. FcγRIV expression was significantly upregulated on neutrophils and monocytes in infected/treated and infected/non-treated mice, the latter showing a stronger FcγRIV upregulation. In contrast, at day 14 p.i. infected/treated mice showed a stronger upregulation of FcγRIV on the three myeloid cells populations as compared to infected/non-treated and control mice. CD11b$^+$Ly6G^{int} neutrophils displayed the strongest expression of FcγRIV as well as the highest magnitude of FcγRIV upregulation after infection and mAb-treatment. No detectable FcγRI expression was observed at day 8 p.i. in any of the three cell types assessed.

However, at day 14 p.i. infected/treated mice showed a significant upregulation of this inducible FcγR

receptor in CD11b$^+$Ly6G^{int} neutrophils and monocytes (Supplemental Figure 5).

These results highlight that viral infection and mAb therapy results in an evolving inflammatory

environment that dramatically modulates the expression of FcγRs in neutrophils and monocytes,

leading to a stronger upregulation of FcγRIV at 8 days p.i. in infected/non treated mice and at 14 days

p.i. in infected/treated mice.

High FcγRIV-expressing CD11b$^+$Ly6G^{int} neutrophils subpopulation displays phenotypic properties

different from those of CD11b$^+$Ly6G^{hi} neutrophils. High level expression and strong upregulation of

FcγRIV on CD11b$^+$Ly6G^{int} (Figure 5C and 5D) suggested different and specific properties of this

neutrophil subpopulation. We thus characterized it further. We observed neutrophils Ly6G^{int} to be

smaller than neutrophils Ly6G^{hi} and displaying a lower granulosity (Figure 5D). To determine the

specificities of this population, we studied the expression pattern of neutrophils and monocytes cell-

surface markers in these cells at steady-state conditions (Figure 5E). Similar to CD11b$^+$Ly6G^{hi}

neutrophils, CD11b$^+$Ly6G^{int} cells expressed low levels of the monocyte cell surface markers Ly6C and

CCR2 (Figure 5E). They also expressed the CXCR4 chemokine receptor although at intermediate

levels as compared to CD11b$^+$Ly6G^{hi} neutrophils and inflammatory monocytes. However, they weakly

express the chemokine receptor CXCR2 which is highly expressed on CD11b$^+$Ly6G^{hi} neutrophils.

These results revealed that Ly6G^{int} cells display a neutrophil-like phenotype but with specific

characteristics as compared to classical Ly6G^{hi} neutrophils, notably (i) smaller size and granulosity,

(ii) higher viral-induced FcγRIV expression, and (iii) a pattern expression of cell surface markers

characteristic of neutrophils (Ly6C^{lo}, CCR2$^-$, CXCR4^{int}) except for the lack of expression of the

CXCR2 receptor. This suggests that CD11b$^+$Ly6G^{int} cells are a phenotypic variant of neutrophils.

However, whether CD11b$^+$Ly6G^{int} cells represent a *bona fide* novel neutrophils subset or neutrophils

displaying a different activation/maturation state could not be discriminated by these phenotypic

properties as this will require integrative high-dimensional analyses including not only functional and

phenotypic characterization but also deep genomic profiling of the neutrophil

subpopulations identified (Ng et al., 2019).

FcγRIV upregulation in infected/treated mice is associated with enhanced expression of MHC-II and

co-stimulatory molecules on neutrophils and monocytes. We next addressed *in vivo* the functional

activation of splenic neutrophils and inflammatory monocytes in infected mice with or without

immunotherapy at days 8 and 14 p.i. by monitoring cell surface activation markers. We were able to

show that the activation state of the cells evolved over time. At day 8 p.i., (Figure 6A) CD11b was upregulated on neutrophils and monocytes upon viral infection either in the absence or in the presence of immunotherapy. However, infected/treated mice showed a significantly higher CD86 upregulation in CD11b+Ly6G^{hi} and CD11b+Ly6G^{int} neutrophils than infected/non-treated mice. MHC-II molecule was also upregulated on CD11b+Ly6G^{int} neutrophils and monocytes upon viral infection. At day 14 p.i. (Figure 6B), CD11b+Ly6G^{hi} splenic neutrophils were similarly activated in infected/treated- and infected/non-treated mice, as deduced from an increased expression of CD11b and CD86. However, CD86 expression on CD11b+Ly6G^{int} neutrophils was significantly increased in infected/treated mice, suggesting a mAb-mediated upregulation of these costimulatory molecules on this neutrophil subpopulation. Likewise, inflammatory monocytes from infected/treated mice also showed a significantly increased expression of CD86 co-stimulatory molecules as compared to infected/non-treated mice. Notably, MHC-II expression was significantly increased in infected/treated mice in both neutrophil populations as well as in monocytes as compared to infected/non-treated mice.

Consistent with a FcγRIV upregulation in neutrophils and monocytes in infected/treated mice (Figure 5C), these data demonstrate a mAb-mediated upregulation of MHC-II and costimulatory molecules in CD11b+Ly6G^{int} neutrophils and monocytes as well as a significant upregulation of MHC-II in CD11b+Ly6G^{hi} neutrophils at 14 days p.i. (Figure 6B).

CD11b+Ly6G^{hi}, CD11b+Ly6G^{int} neutrophils and inflammatory monocytes are differentially and sequentially activated upon antiviral mAb treatment. To further characterize the functional activation state of the two neutrophils populations (CD11b+Ly6G^{hi} and CD11b+Ly6G^{int}) and inflammatory monocytes upon viral infection and mAb-therapy, we next addressed the cytokine and chemokine secretion profile of splenic neutrophils and monocytes sorted from infected mice treated, or not, with the 667 mAb, at days 8 and 14 p.i. Cells sorted from age-matched naive mice were used as controls. We found that the secretion profile evolved over time and was cell type- and stimulus specific, with globally enhanced cytokine/chemokines secretion in cells isolated from infected/treated mice (Figure 7A and 7B), in particular at day 14 p.i.

At day 8 p.i., the CD11b+Ly6G^{hi} neutrophils subpopulation from both infected/non-treated and infected/treated mice showed a broad chemokine and cytokine secretion profile as deduced from an increased secretion of most of the 13 chemokines and 12 cytokines assessed, although at weak levels (Figure 7B and 7C). CD11b+Ly6G^{int} neutrophils showed enhanced expression of several chemokines mostly in cells isolated from infected/non-treated mice notably with a strong enhanced secretion of CCL3, CCL5 and CXCL5. In addition, CD11b+Ly6G^{int} neutrophils also showed a slight secretion of multiples cytokines in infected/treated mice, but to a lesser extent than CD11b+Ly6G^{hi} neutrophils

(lower fold-increase) (Figure 7B and 7C). In contrast, infection and mAb-treatment hardly affected the

chemokine/cytokine secretion of Ly6C^{hi} monocytes. They only showed a weak increase in CXCL1

secretion in infected/treated mice as well as a slight increase in cytokines secretion (both in terms of

diversity and fold increase) mainly in infected/non-treated mice (Figure 7B).

Interestingly, at day 14 p.i., the functional activation of the three different cell types assessed

completely differed from that observed at day 8 p.i. (Figure 7B and 7C). We found a more restricted

but stronger induction of chemokines/cytokines secretion, mostly in cells isolated from

infected/treated mice. In contrast to the broad cytokines and chemokines secretion profile observed at

day 8 p.i., CD11b$^+$Ly6G^{hi} neutrophils only secreted CCL4 and CCL5 chemokines and showed no

cytokine secretion at day 14 p.i. In addition, CD11b$^+$Ly6G^{int} neutrophils showed a strong enhanced

expression of 6 chemokines (CCL3, CCL4, CCL5, CCL22, CXCL9 and CXCL10) and 3 Th1-

polarizing cytokines (IL-6, TNFα, IFNγ) in infected/treated mice. Ly6C^{hi} monocytes showed a

proinflammatory chemokine/cytokine secretion profile similar to CD11b$^+$Ly6G^{int} neutrophils, except

for the secretion of CXCL9, that was specific to CD11b$^+$Ly6G^{int} neutrophils and CXCL1 that was only

secreted by monocytes.

In summary, FrCasE infection and 667 mAb-treatment induce the secretion of multiple chemokines

and cytokines by neutrophils and monocytes. Interestingly, the secretion profile evolves over time and

differs among the three different cell types assessed, providing further evidence of a different

functional activation state of the two neutrophils subpopulations (Ly6G^{hi} and Ly6G^{int}). In addition,

these data show an enhanced secretion of Th1-polarizing cytokines and chemokines by cells isolated

from infected/treated mice at 14 p.i, showing a role for the therapeutic mAb in the functional

activation of these FcγRIV highly-expressing cells.

DISCUSSION

We have previously shown that neutrophils have a key immunomodulatory role in the induction of protective immunity by antiviral mAbs through the acquisition of B-cell helper functions (Naranjo-Gomez et al., 2018). Here we provide a new insight into both the heterogeneity and immunomodulatory role of neutrophils in a context of antiviral mAb-therapy. Our work shows that two functionally distinct neutrophil subpopulations can secrete multiple cytokines and chemokines able to recruit and activate multiple cells of the innate and the adaptive immune system. The cytokine/chemokine secretion profile: differs between both neutrophil subpopulations, is stimulus-dependent, evolves over time upon viral infection and mAb-treatment and suggests a potential cooperation with inflammatory monocytes in the induction of Th1 immune responses upon mAb-treatment of infected mice.

Our data show that neutrophils cytokine/chemokine secretion profile differs between viral *versus* bacterial stimulus. This is important to highlight as most of the studies investigating the immunomodulatory role of neutrophils have been conducted in bacterial infection models. Thus, *in vitro* stimulation of neutrophils by viral determinants led to a poor production of cytokines but to a wide and strong release of chemokine, with notably high secretion levels of the monocytes-and neutrophils-recruiting chemokines (CCL2, CXCL1, CXCL5) that was not observed upon LPS stimulation. On the contrary, LPS-stimulated neutrophils produced high amounts of proinflammatory cytokines but a narrower and weaker chemokine release. As for neutrophils, stimulation of monocytes by viral determinants (but not LPS) led to the secretion of high amounts of the neutrophils-recruiting chemokine CXCL1 and to a lesser extent the monocytes-recruiting chemokine CCL2. These results suggest a self-sustaining mechanism of neutrophils and monocytes recruitment upon viral infection, and raise the hypothesis of an early cooperation between neutrophils and monocytes in the modulation/induction of the antiviral immune response. This secretion profile of neutrophils induced by viral determinants is in agreement with an increased CCL2 release observed upon *in vitro* HIV stimulation of neutrophils (Yoshida et al., 2009). Increased levels of CCL2 and CXCL1 have also been reported in neutrophil- and monocytes-infiltrated tissues in different viral infections, (Hokeness et al., 2005; Miller et al., 2004; Rubio and Sanz-Rodriguez, 2007; Seo et al., 2011). However, chemokine increase was mostly assessed in total tissue extracts but not directly in neutrophils or monocytes isolated cells, which prevented the identification of the cell origin of chemokines. It is worth noting that a very early but transient expression has also been reported (i.e. 24-48h p.i.) of CCL2 and CXCL1 upon infection by RSV, CMV and influenza virus (Hokeness et al., 2005; Miller et al., 2004; Wareing et al., 2004). Such early and transient expression would be in agreement with the very low secretion of these chemokines detected at day 8 p.i. in infected mice treated, or not, with the therapeutic mAb.

Our data also highlight the effect of inflammatory conditions on the modulation of the phenotypic and functional activity of neutrophils and monocytes. Thus, *in vitro* priming with inflammatory/immunomodulatory cytokines clearly potentiated the release of several chemokines and cytokines by IC-activated neutrophils and monocytes (Figure 3) while having a less pronounced or no effect on virus-activated cells. This suggests that cytokine-priming associated to FcγR-triggering leads to enhanced immune responses. Although not formally shown, these results support a role for the inflammatory environment in the enhancement of chemokine/cytokine secretion by neutrophils and monocytes observed in infected-treated mice. Furthermore, our *in vitro* observations also allowed us to dissect the specific effect of IFN-γ, IFN-I and TNF-α on the enhancement of the functional properties of IC-activated neutrophils and monocytes. Notably, IFN-γ and IFN-I priming (but not TNF-α), led to the upregulation of FcγRs in neutrophils and/or monocytes, in agreement with previous reports in other experimental settings (Dahal et al., 2017; Lehmann et al., 2017; Morquin et al., 2017; Pricop et al., 2001). Consistent with these results, our *in vivo* work shows that the inflammatory environment resulting from ongoing infection and mAb-treatment also modulates FcγRs expression on neutrophils and monocytes (in particular FcγRIV). Thus, FcγRIV expression is more strongly enhanced in infected/non-treated mice at day 8 p.i. and in infected/treated mice at day 14 p.i., suggesting a different and evolving inflammatory environment in both groups of mice. These results are in agreement with previous work showing FcγRs modulation by the inflammatory conditions resulting from bacterial and IC-mediated autoimmune pathologies (Bruhns, 2012; Wang and Jönsson, 2019); and provide new evidence on the regulation of FcγRs expression in the specific inflammatory context of antiviral mAb-based immunotherapies, not hitherto reported. Notably, our work shows that the magnitude of FcγRIV upregulation upon viral infection and mAb-treatment is different in neutrophils and monocytes. This differential FcγRIV expression together with a different expression level of the Ly6G neutrophil cell surface marker (Ly6G^{hi} and Ly6G^{int}) contributed to the identification and characterization of two neutrophils subpopulation displaying distinct functional properties as deduced from their different (i) phenotype (size, granulosity, cell surface molecules) (Figures 5 and 6), (ii) magnitude of expression and upregulation of FcγRIV, costimulatory molecules and MHC-II (Figures 5 and 6) and (iii) cytokine/chemokine secretion profile (Figure 7) upon viral infection and mAb-treatment. Recent studies have shown neutrophils heterogeneity in healthy tissues as well as in different pathological conditions (Ng et al., 2019; Silvestre-Roig et al., 2019) through the identification of different neutrophils subpopulations. Such neutrophil heterogeneity is context-dependent as it is influenced by the nature and magnitude of the stimuli. In the particular context of infectious diseases, neutrophils heterogeneity is still poorly characterized and it has mostly been assessed in bacterial infections (Deniset et al., 2017; Tsuda et al., 2004). Similar to our observations Deniset and collaborators (Deniset et al., 2017) reported two distinct splenic neutrophils subpopulations displaying different

Ly6G expression levels (Ly6G^{hi} and Ly6G^{int}) in *Streptococcus pneumoniae* infected mice. This study

evidenced some phenotypic differences (cell surface markers of maturation, nuclear morphology, ...)

between both neutrophil subpopulations but their functional activation was not assessed. Importantly,

we describe here two phenotypic and functional variants of neutrophils displaying distinct

immunomodulatory properties that might be key and complementary for inducing antiviral protective

immunity. Our work provides thus novel findings on the heterogeneity and function of neutrophils

upon viral infection and mAb-treatment.

Our work provides new mechanistic insights into the hitherto underestimated role of neutrophils as

key cells in the modulation of adaptive antiviral immunity upon mAb treatment. Interestingly, at

steady-state conditions, neutrophils express high levels of FcγRIV. Its expression is higher than that

observed in inflammatory monocytes. Thus, it is tempting to speculate that upon viral infection and

mAb-treatment, the early neutrophils recruitment, together with their high expression of FcγRIV,

result in potent Fc-triggering by ICs in these cells. This might be key to initiate the modulation of

immune responses through the production of multiple cytokines and chemokines able to recruit and

activate multiple innate immune cells such as monocytes, NK cells, DC and neutrophils themselves

(Tecchio and Cassatella, 2016). This suggest a potential role for neutrophils as early drivers of the

induction of vaccinal effects by mAbs. Supporting this hypothesis, at day 8 p.i. both neutrophils

populations showed a higher and a wider induction of chemokines and cytokines release than

monocytes. Subsequently, the inflammatory environment resulting from viral infection and mAb-

mediated antiviral effects induces the upregulation of FcγRs in the different innate immune cells that

might further increase Fc-triggering by ICs leading to improved immune responses, both in terms of

magnitude and quality. In agreement with this, the FcγRIV upregulation in neutrophils and monocytes

in infected/treated mice observed at 14 days p.i. is associated with (i) higher expression of

costimulatory molecules and MHC-II and (ii) higher secretion of cytokines and chemokines. In

keeping with this, enhanced secretion of Th1-polarizing cytokines (IL-6, TNFα, IFNγ) (Czarniecki,

1993; Hildenbrand et al., 2008; Jin et al., 2016; Uciechowski and Dempke, 2020; Wang et al., 2005)

and chemokines (CXCL9 and CXCL10) (Gaylo-Moynihan et al., 2019; Kelly-Scumpia et al., 2010;

Tannenbaum et al., 1998; Tokunaga et al., 2018) by CD11b$^+$Ly6G^{int} neutrophils and monocytes from

infected/treated mice at 14 p.i, argues in favor of a mAb-mediated Th1 polarization of the antiviral

immune response by these cells. Consistent with such Th1 polarization, we previously reported strong

CD8 T-cells responses at day 14 p.i. in infected/treated mice (Gros et al., 2008; Michaud et al., 2010)

as well as long-lasting protective humoral immune responses predominantly of the IgG2a isotype

(Gros et al., 2005; Michaud et al., 2010; Naranjo-Gomez et al., 2018). Finally, it is also worth

mentioning the upregulation of MHC-II molecules observed in CD11b$^+$Ly6G^{hi} and CD11b$^+$Ly6G^{int}

neutrophils sorted from infected/treated mice. This observation broadens the immunomodulatory role

of neutrophils upon mAb-treatment as it suggests that neutrophils might acquire antigen-presenting

cell features and induce antigen-specific T-cells responses (Lok et al., 2019; Meinderts et al., 2019;

Vono et al., 2017). In agreement with this, it has been shown that upon *ex vivo* stimulation with IgG

immune complex (IC) or viral cognate antigens, neutrophils upregulated expression of MHCII and

costimulatory molecules and increased T cell activation.

Our work show that mAb-mediated activation of neutrophils and monocytes might be essential to

induce strong Th1-polarized early immune responses that might be key to induce protective immunity

by mAb. In addition, the distinct secretion profile of Ly6G^{hi} and Ly6G^{int} neutrophils and Ly6C^{hi}

monocytes, in terms of type, amount and kinetics of chemokines/cytokines secretion, shows a

differentially and sequentially activation of theses FcγRIV-expressing cells upon antiviral mAb

treatment and suggests their key and complementary action in the induction of protective immune

responses by mAb. Thus, the identification of neutrophils as key players in the induction of protective

immunity by antiviral mAbs and their potential collaboration with inflammatory monocytes might

have therapeutic implications. These findings might help to improve mAb-based antiviral therapies by

tailoring therapeutic interventions aiming at harnessing the immunomodulatory properties of these

cells. To this end, different approaches could be envisaged: (i) the use of appropriate

immunostimulatory host-directed therapies and (ii) the design of antiviral mAbs engineered to

enhance their affinity for FcγRs expressed on human neutrophils and monocytes, such as FcγRIIa and

FcγRIIIa (Bournazos et al., 2019; Chen et al., 2017; DiLillo and Ravetch, 2015) .Thus, in addition to

allow superior antibody-mediated phagocytosis, these Fc-engineered mAbs could also modulate

cytokine/chemokine production to ultimately lead to more effective adaptive immune responses.

METHODS

Mice: Inbred 129/Sv/Ev mice (H-2D^b haplotype) were used and maintained under conventional, pathogen-free facilities at the Institut de Génétique Moléculaire de Montpellier (RAM-ZEFI). They have been used without distinction as to sex and at different ages according to experiments.

Viral stocks: FrCas^E (Portis et al., 1990) viral stocks were produced and stored as previously described in (Gros et al., 2005).

Viral infection and immunotherapy: Eight-day-old 129/Sv/Ev mice were infected by intraperitoneal (i.p.) administration with 50 μl of a viral suspension containing 50,000 focus-forming units (FFU) and treated, or not, with 30 μg 667 mAb targeting gp70 protein of viral envelope, (McAtee and Portis, 1985), 1-hour p.i. and on days 2 and 5 p.i. by i.p. administration. Mice were euthanized and spleens collected at days 8 and 14 p.i.

Phenotypical and functional activation of FcγRIV-expressing cell from spleen ex vivo: Single-cell suspensions of splenocytes were obtained from naive, infected/non-treated and infected/treated mice at 8 and 14 days p.i. and immunotherapy. Spleen cell suspensions were obtained by mechanical dissociation in PBS, then filtered in 0,70 μm strainer. 20 % of each spleen was used for immunophenotyping by FACS, 80% left was dedicated to main FcγRIV-expressing cell sorting. Red blood cells were lysed (ACK, Lonza) and an enrichment with biotinylated anti-B220 (BD Biosciences), anti-CD3 (BD Biosciences), following of anti-biotin Ab magnet-bead coupled (Miltenyi) and magnetic LS-columns (Miltenyi) was performed to remove spleen lymphocytes and increase the sorting efficacy. Cells were stained with specific marker of populations of interest (Ly6G-BD BioSciences, Ly6C-BioLegend, CD11b-BD BioSciences) and neutrophils (Ly6G^{hi} and L6G^{int}) and inflammatory monocytes (Ly6C^{hi}) were sorted (>97-98% pure) using the BD Biosciences FACSAria device. Sorted cells were cultured in 96-well plates at a density of $4x10^6$ cells/ml ($1x10^6$ cells/well) in 10% FBS-containing RPMI medium for 24h, and cell-free supernatants were collected and stored at -20°C, to allow cytokines and chemokines protein release quantification.

Flow cytometry: Organs of interest were collected to realize immunophenotyping of immune cells. Spleen cell suspensions were obtained by mechanical dissociation in PBS. BM cell suspensions were obtained by dissection and PBS-2%-FBS flushing of tibias and femurs. Cells were stained at 4°C using fluorochrome-conjugated antibodies against CCR2 (SA203G11, BioLegend), CD11b (M1/70, eBioscience), CD11c (HL3, BD Biosciences), CD16/32 (93, BD Biosciences), CD45.2 (104, BD Biosciences), CD62L (MEL-14, BD Biosciences), CD64 (x54-5/7.1, BD Biosciences), CD86 (GL1, BD Biosciences), CXCR2 (SA044G4, BioLegend), CXCR4 (2B11, Life Technologies), Ly6G (1A8,

BD Biosciences), Ly6C (AL-21, BioLegend) and Siglec F (E50-2440, BD Pharmingen). FcγRIV expression was determined using 9E9 antibody (kindly provided by Dr.Pierre Bruhns, Institut Pasteur), produced by BioXcell, and then labeled with Alexa Fluor 647. FrCasE-infected cells were assayed using an anti-Gag mAb (H34) (Chesebro et al., 1981; kindly provided by Pr. Ulf Dittmer, University of Duisburg-Essen) labeled with Alexa Fluor 647 or incubated with an anti-IgG2b labeled PE. Forward scatter area and forward scatter time-of-flight, as well as side scatter, were used to remove doublets from flow cytometry analyses. Cells were analyzed on FACS LSR Fortessa (BD Bioscience), and the data were analyzed using the FlowJo software.

Neutrophils and monocytes isolated from BM used in in vitro experiments*:* Neutrophils and monocytes were purified from 8 to 11-week-old naive mice BM. After dissection of lower limbs, BM cell suspensions were collected by PBS-2% FBS EDTA (2 mM) flushing (25G needle) of tibias and femurs. BM cell suspensions were filtered with a 0,40 μm nylon strainer. Two magnetic-based cell sorting (MACS) isolation kits were used to purified either neutrophils (Miltenyi Biotec) and monocytes, (Miltenyi Biotec), both with high purity (>97-98%), determined by FACS (LSR Fortessa, BD Bioscience). Cells were placed in culture in U bottom 96-well plates at a concentration of 1 million/ml in 10% FBS-containing RPMI medium.

Stimulation of neutrophils and monocytes in vitro: Purified neutrophils and monocytes were seeded at 150 000 cells/ well in 150 μl of RPMI, then incubated for 24h with LPS (1μg/ml, Sigma), or FrCas^E virus (MOI 5: 5 viral particles /cell), or ICs (virus FrCas^E and 1μg 667 mAb), or 1μg 667 mAb alone. In parallel, the same experiments were performed adding inflammatory cytokines, TNFα 100 UI/ml (Peprotech), IFN γ 100 UI/ml (eBioScience), IFNα11 a 1000 UI/ml, produced and generously provided by Dr. Gilles Uzé (DIMP, CNRS). After 24h of stimulation, supernatants were collected and stored at -20°C to quantify chemokines and cytokines protein release secretion (describe below). Phenotypical activation of both neutrophils and monocytes was measured using surface markers by flow cytometry.

Chemokines and cytokines protein release quantification: Soluble chemokines and cytokines secretion were quantified from cell-free collected supernatants of *in vitro* cultured neutrophils and monocytes and of sorted splenic neutrophils Ly6G^{hi} and Ly6G^{int}, and inflammatory monocytes (of naive, infected/non-treated and infected/treated mice 8 and 14 days p.i. and immunotherapy), using bead-based immunoassays (LegendPLex, BioLegend) and analyzed on the BD Bioscience-LSR Fortessa device. The protein release quantification was established by the appropriate software (LEGENDplex^{TM} data analysis).

Statistics: Statistical analyses were performed using GraphPad Prism 5 (GraphPad Software). Data
were expressed as mean ± SEM, and statistical significance was established using a parametric 1-way
ANOVA test or non-parametric Kruskal-Wallis test for multiple comparisons or paired Student's *t*
tests when two groups were compared. *P* values lower than 0.05 were considered as statistically
significant.

Study approval. All experimental procedures were performed in accordance with the French national
animal care guidelines (CEEA-LR-12146 approval, Ethics committee of the Languedoc-Roussillon
Region, Montpellier).

AUTHOR CONTRIBUTIONS

586
587 Jennifer Lambour (JL), Mar Naranjo-Gomez (MN-G) and Mireia Pelegrin (MP) defined the research
588 program. JL, MN-G, and Myriam Boyer-Clavel (MB-C) performed the experiments. JL, MN-G and
589 MP carried out the data analyses. JL and MP wrote the manuscript. Grants to MP funded the study.
590

591 **ACKNOWLEDGMENTS**
592
This work was supported by grants from the ANRS (France REcherche Nord&sud Sida-hiv
Hépatites), the Ligue Régionale Contre le Cancer, Sidaction and the Fondation pour la Recherche
Médicale. M. Naranjo-Gomez, J. Lambour, and M. Pelegrin are members of the "MabImprove
Labex", a public grant overseen by the French National Research Agency (ANR) as part of the
"Investments for the future" program (reference: ANR-10-LABX -53-01) that also supported this
work. We thank the imaging facility MRI, which is part of the UMS BioCampus Montpellier and a
member of the national infrastructure France-BioImaging, supported by the French National Research
Agency (ANR-10-INBS-04, "Investments for the future"). We are grateful to the animal facility of the
Institut de Génétique Moléculaire de Montpellier (RAM-ZEFI) which is part of the "Réseau des
Animaleries Montpelliéraines" RAMIBiSA Facility for animal experiments. We are grateful to S.
Gailhac from MRI for support in cytometry experiments, to Thierry Gostan (SERENAD Complex
Biological Data Analysis Service) for support in statistical analyses, to Helen Phillips Bevis for
English editing services and to Drs. Valerie Dardalhon, Gilles Uzé and Pascale Plence for critical
reading of the manuscript.

CONFLICT OF INTEREST DISCLOSURES
609
610 The authors declare no competing financial interests.

REFERENCES

Bournazos, S., and Ravetch, J.V. (2017). Fcγ Receptor Function and the Design of Vaccination Strategies. Immunity *47*, 224–233.

Bournazos, S., Klein, F., Pietzsch, J., Seaman, M.S., Nussenzweig, M.C., and Ravetch, J.V. (2014). Broadly Neutralizing Anti-HIV-1 Antibodies Require Fc Effector Functions for In Vivo Activity. Cell *158*, 1243–1253.

Bournazos, S., DiLillo, D.J., Goff, A.J., Glass, P.J., and Ravetch, J.V. (2019). Differential requirements for FcγR engagement by protective antibodies against Ebola virus. Proc Natl Acad Sci USA *116*, 20054–20062.

Bruhns, P. (2012). Properties of mouse and human IgG receptors and their contribution to disease models. Blood *119*, 5640–5649.

Celis, E., and Chang, T.W. (1984). HBsAg-serum protein complexes stimulate immune T lymphocytes more efficiently than do pure HBsAg. Hepatology *4*, 1116–1123.

Chen, T.F., Sazinsky, S.L., Houde, D., DiLillo, D.J., Bird, J., Li, K.K., Cheng, G.T., Qiu, H., Engen, J.R., Ravetch, J.V., et al. (2017). Engineering Aglycosylated IgG Variants with Wild-Type or Improved Binding Affinity to Human Fc Gamma RIIA and Fc Gamma RIIIAs. Journal of Molecular Biology *429*, 2528–2541.

Chesebro, B., Collins, J.K., Wehrly, K., Nishio, J., and Cloyd, M. (1981). Expression of cell surface Friend virus gp70 does not block reinfection by ecotropic murine leukemia viruses. Virology *115*, 125–129.

Czarniecki, C.W. (1993). The role of tumor necrosis factor in viral disease. Antiviral Research *22*, 223–258.

Dahal, L.N., Dou, L., Hussain, K., Liu, R., Earley, A., Cox, K.L., Murinello, S., Tracy, I., Forconi, F., Steele, A.J., et al. (2017). STING Activation Reverses Lymphoma-Mediated Resistance to Antibody Immunotherapy. Cancer Res *77*, 3619–3631.

Deniset, J.F., and Kubes, P. (2018). Neutrophil heterogeneity: Bona fide subsets or polarization states? J. Leukoc. Biol. *103*, 829–838.

Deniset, J.F., Surewaard, B.G., Lee, W.-Y., and Kubes, P. (2017). Splenic Ly6Ghighmature and Ly6Gintimmature neutrophils contribute to eradication ofS. pneumoniae. J. Exp. Med. *214*, 1333–1350.

Dibo, M., Battocchio, E.C., dos Santos Souza, L.M., da Silva, M.D.V., Banin-Hirata, B.K., Sapla, M.M.M., Marinello, P., Rocha, S.P.D., and Faccin-Galhardi, L.C. (2019). Antibody Therapy for the Control of Viral Diseases: An Update. CPB *20*, 1108–1121.

DiLillo, D.J., and Ravetch, J.V. (2015). Differential Fc-Receptor Engagement Drives an Anti-tumor Vaccinal Effect. Cell *161*, 1035–1045.

Earnest, J.T., Basore, K., Roy, V., Bailey, A.L., Wang, D., Alter, G., Fremont, D.H., and Diamond, M.S. (2019). Neutralizing antibodies against Mayaro virus require Fc effector functions for protective activity. Journal of Experimental Medicine *216*, 2282–2301.

Fan, Y., Wang, L., and Dou, X. (2018). Serum Monocyte Chemoattractant Protein-1 Predicts Liver Inflammation of Patients with Chronic Hepatitis B. Clin. Lab. *64*.

Fox, J.M., Roy, V., Gunn, B.M., Huang, L., Edeling, M.A., Mack, M., Fremont, D.H., Doranz, B.J., Johnson, S., Alter, G., et al. (2019). Optimal therapeutic activity of monoclonal antibodies against chikungunya virus requires Fc-FcγR interaction on monocytes. Sci. Immunol. *4*, eaav5062.

Gaylo-Moynihan, A., Prizant, H., Popović, M., Fernandes, N.R.J., Anderson, C.S., Chiou, K.K., Bell, H., Schrock, D.C., Schumacher, J., Capece, T., et al. (2019). Programming of Distinct Chemokine-Dependent and -Independent Search Strategies for Th1 and Th2 Cells Optimizes Function at Inflamed Sites. Immunity *51*, 298-309.e6.

Gros, L., Dreja, H., Fiser, A.L., Plays, M., Pelegrin, M., and Piechaczyk, M. (2005). Induction of long-term protective antiviral endogenous immune response by short neutralizing monoclonal antibody treatment. J. Virol. *79*, 6272–6280.

Gros, L., Pelegrin, M., Michaud, H.-A., Bianco, S., Hernandez, J., Jacquet, C., and Piechaczyk, M. (2008). Endogenous cytotoxic T-cell response contributes to the long-term antiretroviral protection induced by a short period of antibody-based immunotherapy of neonatally infected mice. J. Virol. *82*, 1339–1349.

Gunn, B.M., Yu, W.-H., Karim, M.M., Brannan, J.M., Herbert, A.S., Wec, A.Z., Halfmann, P.J., Fusco, M.L., Schendel, S.L., Gangavarapu, K., et al. (2018). A Role for Fc Function in Therapeutic Monoclonal Antibody-Mediated Protection against Ebola Virus. Cell Host & Microbe *24*, 221-233.e5.

Haider, C., Hnat, J., Wagner, R., Huber, H., Timelthaler, G., Grubinger, M., Coulouarn, C., Schreiner, W., Schlangen, K., Sieghart, W., et al. (2019). Transforming Growth Factor-β and Axl Induce CXCL5 and Neutrophil Recruitment in Hepatocellular Carcinoma: HEPATOBILIARY MALIGNANCIES. Hepatology *69*, 222–236.

Hildenbrand, B., Lorenzen, D., Sauer, B., Hertkorn, C., Freudenberg, M.A., Peters, J.H., Nesselhut, T., Unger, C., and Azemar, M. (2008). IFN-y enhances T(H)1 polarisation of monocyte-derived dendritic cells matured with clinical-grade cytokines using serum-free conditions. Anticancer Res. *28*, 1467–1476.

Hokeness, K.L., Kuziel, W.A., Biron, C.A., and Salazar-Mather, T.P. (2005). Monocyte Chemoattractant Protein-1 and CCR2 Interactions Are Required for IFN- / -Induced Inflammatory Responses and Antiviral Defense in Liver. The Journal of Immunology *174*, 1549–1556.

Hooft van Huijsduijnen, R., Kojima, S., Carter, D., Okabe, H., Sato, A., Akahata, W., Wells, T.N.C., and Katsuno, K. (2020). Reassessing therapeutic antibodies for neglected and tropical diseases. PLoS Negl Trop Dis *14*, e0007860.

Jin, P., Zhao, Y., Liu, H., Chen, J., Ren, J., Jin, J., Bedognetti, D., Liu, S., Wang, E., Marincola, F., et al. (2016). Interferon-γ and Tumor Necrosis Factor-α Polarize Bone Marrow Stromal Cells Uniformly to a Th1 Phenotype. Sci Rep *6*, 26345.

Kelly-Scumpia, K.M., Scumpia, P.O., Delano, M.J., Weinstein, J.S., Cuenca, A.G., Wynn, J.L., and Moldawer, L.L. (2010). Type I interferon signaling in hematopoietic cells is required for survival in mouse polymicrobial sepsis by regulating CXCL10. The Journal of Experimental Medicine *207*, 319–326.

Kim, Y.-G., Kamada, N., Shaw, M.H., Warner, N., Chen, G.Y., Franchi, L., and Núñez, G. (2011). The Nod2 Sensor Promotes Intestinal Pathogen Eradication via the Chemokine CCL2-Dependent Recruitment of Inflammatory Monocytes. Immunity *34*, 769–780.

Lambour, J., Naranjo-Gomez, M., Piechaczyk, M., and Pelegrin, M. (2016). Converting monoclonal
antibody-based immunotherapies from passive to active: bringing immune complexes into play.
Emerg Microbes Infect *5*, e92.

Lehmann, B., Biburger, M., Brückner, C., Ipsen-Escobedo, A., Gordan, S., Lehmann, C., Voehringer,
D., Winkler, T., Schaft, N., Dudziak, D., et al. (2017). Tumor location determines tissue-specific
recruitment of tumor-associated macrophages and antibody-dependent immunotherapy response. Sci.
Immunol. *2*, eaah6413.

Leonard, E.J., and Yoshimura, T. (1990). Human monocyte chemoattractant protein-1 (MCP-1).
Immunol. Today *11*, 97–101.

Liu, Q., Fan, C., Li, Q., Zhou, S., Huang, W., Wang, L., Sun, C., Wang, M., Wu, X., Ma, J., et al.
(2017). Antibody-dependent-cellular-cytotoxicity-inducing antibodies significantly affect the post-
exposure treatment of Ebola virus infection. Sci Rep *7*, 45552.

Lok, L.S.C., Dennison, T.W., Mahbubani, K.M., Saeb-Parsy, K., Chilvers, E.R., and Clatworthy, M.R.
(2019). Phenotypically distinct neutrophils patrol uninfected human and mouse lymph nodes. Proc
Natl Acad Sci USA *116*, 19083–19089.

Lu, L.L., Suscovich, T.J., Fortune, S.M., and Alter, G. (2018). Beyond binding: antibody effector
functions in infectious diseases. Nat. Rev. Immunol. *18*, 46–61.

Mantovani, A., Cassatella, M.A., Costantini, C., and Jaillon, S. (2011). Neutrophils in the activation
and regulation of innate and adaptive immunity. Nat. Rev. Immunol. *11*, 519–531.

McAtee, F.J., and Portis, J.L. (1985). Monoclonal antibodies specific for wild mouse neurotropic
retrovirus: detection of comparable levels of virus replication in mouse strains susceptible and
resistant to paralytic disease. J. Virol. *56*, 1018–1022.

Meinderts, S.M., Baker, G., van Wijk, S., Beuger, B.M., Geissler, J., Jansen, M.H., Saris, A., ten
Brinke, A., Kuijpers, T.W., van den Berg, T.K., et al. (2019). Neutrophils acquire antigen-presenting
cell features after phagocytosis of IgG-opsonized erythrocytes. Blood Advances *3*, 1761–1773.

Michaud, H.-A., Gomard, T., Gros, L., Thiolon, K., Nasser, R., Jacquet, C., Hernandez, J.,
Piechaczyk, M., and Pelegrin, M. (2010). A crucial role for infected-cell/antibody immune complexes
in the enhancement of endogenous antiviral immunity by short passive immunotherapy. PLoS Pathog.
6, e1000948.

Miller, A.L., Bowlin, T.L., and Lukacs, N.W. (2004). Respiratory Syncytial Virus–Induced
Chemokine Production: Linking Viral Replication to Chemokine Production In Vitro and In Vivo. J
INFECT DIS *189*, 1419–1430.

Morquin, D., Tuaillon, E., Makinson, A., Bendriss, S., Le Moing, V., and Reynes, J. (2017). Impact of
T cell activation, HIV replication and hepatitis C virus infection on neutrophil CD64 expression.
Cytometry B Clin Cytom *92*, 492–497.

Naranjo-Gomez, M., and Pelegrin, M. (2019). Vaccinal effect of HIV-1 antibody therapy: Current
Opinion in HIV and AIDS *14*, 325–333.

Naranjo-Gomez, M., Lambour, J., Piechaczyk, M., and Pelegrin, M. (2018). Neutrophils are essential
for induction of vaccine-like effects by antiviral monoclonal antibody immunotherapies. JCI Insight *3*.

Nasser, R., Pelegrin, M., Michaud, H.-A., Plays, M., Piechaczyk, M., and Gros, L. (2010). Long-
lasting protective antiviral immunity induced by passive immunotherapies requires both neutralizing
and effector functions of the administered monoclonal antibody. J. Virol. *84*, 10169–10181.

Ng, L.G., Ostuni, R., and Hidalgo, A. (2019). Heterogeneity of neutrophils. Nat Rev Immunol *19*,
255–265.

Niessl, J., Baxter, A.E., Mendoza, P., Jankovic, M., Cohen, Y.Z., Butler, A.L., Lu, C.-L., Dubé, M.,
Shimeliovich, I., Gruell, H., et al. (2020). Combination anti-HIV-1 antibody therapy is associated with
increased virus-specific T cell immunity. Nat Med *26*, 222–227.

Pelegrin, M., Naranjo-Gomez, M., and Piechaczyk, M. (2015). Antiviral Monoclonal Antibodies: Can
They Be More Than Simple Neutralizing Agents? Trends Microbiol. *23*, 653–665.

Portis, J.L., Czub, S., Garon, C.F., and McAtee, F.J. (1990). Neurodegenerative disease induced by the
wild mouse ecotropic retrovirus is markedly accelerated by long terminal repeat and gag-pol
sequences from nondefective Friend murine leukemia virus. J. Virol. *64*, 1648–1656.

Pricop, L., Redecha, P., Teillaud, J.L., Frey, J., Fridman, W.H., Sautès-Fridman, C., and Salmon, J.E.
(2001). Differential modulation of stimulatory and inhibitory Fc gamma receptors on human
monocytes by Th1 and Th2 cytokines. J. Immunol. *166*, 531–537.

Pruenster, M., Mudde, L., Bombosi, P., Dimitrova, S., Zsak, M., Middleton, J., Richmond, A.,
Graham, G.J., Segerer, S., Nibbs, R.J.B., et al. (2009). The Duffy antigen receptor for chemokines
transports chemokines and supports their promigratory activity. Nat Immunol *10*, 101–108.

Ritzman, A.M., Hughes-Hanks, J.M., Blaho, V.A., Wax, L.E., Mitchell, W.J., and Brown, C.R.
(2010). The Chemokine Receptor CXCR2 Ligand KC (CXCL1) Mediates Neutrophil Recruitment and
Is Critical for Development of Experimental Lyme Arthritis and Carditis. Infection and Immunity *78*,
4593–4600.

Robben, P.M., LaRegina, M., Kuziel, W.A., and Sibley, L.D. (2005). Recruitment of Gr-1+
monocytes is essential for control of acute toxoplasmosis. Journal of Experimental Medicine *201*,
1761–1769.

Rovai, L.E., Herschman, H.R., and Smith, J.B. (1998). The murine neutrophil-chemoattractant
chemokines LIX, KC, and MIP-2 have distinct induction kinetics, tissue distributions, and tissue-
specific sensitivities to glucocorticoid regulation in endotoxemia. J Leukoc Biol *64*, 494–502.

Rubio, N., and Sanz-Rodriguez, F. (2007). Induction of the CXCL1 (KC) chemokine in mouse
astrocytes by infection with the murine encephalomyelitis virus of Theiler. Virology *358*, 98–108.

Salazar, G., Zhang, N., Fu, T.-M., and An, Z. (2017). Antibody therapies for the prevention and
treatment of viral infections. NPJ Vaccines *2*, 19.

Scapini, P., Marini, O., Tecchio, C., and Cassatella, M.A. (2016). Human neutrophils in the saga of
cellular heterogeneity: insights and open questions. Immunol. Rev. *273*, 48–60.

Schoofs, T., Klein, F., Braunschweig, M., Kreider, E.F., Feldmann, A., Nogueira, L., Oliveira, T.,
Lorenzi, J.C.C., Parrish, E.H., Learn, G.H., et al. (2016). HIV-1 therapy with monoclonal antibody
3BNC117 elicits host immune responses against HIV-1. Science *352*, 997–1001.

Seo, S.-U., Kwon, H.-J., Ko, H.-J., Byun, Y.-H., Seong, B.L., Uematsu, S., Akira, S., and Kweon, M.-
N. (2011). Type I interferon signaling regulates Ly6C(hi) monocytes and neutrophils during acute
viral pneumonia in mice. PLoS Pathog. *7*, e1001304.

Silvestre-Roig, C., Hidalgo, A., and Soehnlein, O. (2016). Neutrophil heterogeneity: implications for
homeostasis and pathogenesis. Blood *127*, 2173–2181.

Silvestre-Roig, C., Fridlender, Z.G., Glogauer, M., and Scapini, P. (2019). Neutrophil Diversity in
Health and Disease. Trends in Immunology *40*, 565–583.

Tamassia, N., Bianchetto-Aguilera, F., Arruda-Silva, F., Gardiman, E., Gasperini, S., Calzetti, F., and
Cassatella, M.A. (2018). Cytokine production by human neutrophils: revisiting the "dark side of the
moon." Eur. J. Clin. Invest. e12952.

Tannenbaum, C.S., Tubbs, R., Armstrong, D., Finke, J.H., Bukowski, R.M., and Hamilton, T.A.
(1998). The CXC chemokines IP-10 and Mig are necessary for IL-12-mediated regression of the
mouse RENCA tumor. J. Immunol. *161*, 927–932.

Tecchio, C., and Cassatella, M.A. (2016a). Neutrophil-derived chemokines on the road to immunity.
Semin. Immunol. *28*, 119–128.

Tecchio, C., and Cassatella, M.A. (2016b). Neutrophil-derived chemokines on the road to immunity.
Seminars in Immunology *28*, 119–128.

Tokunaga, R., Zhang, W., Naseem, M., Puccini, A., Berger, M.D., Soni, S., McSkane, M., Baba, H.,
and Lenz, H.-J. (2018). CXCL9, CXCL10, CXCL11/CXCR3 axis for immune activation – A target
for novel cancer therapy. Cancer Treatment Reviews *63*, 40–47.

Tsuda, Y., Takahashi, H., Kobayashi, M., Hanafusa, T., Herndon, D.N., and Suzuki, F. (2004). Three
Different Neutrophil Subsets Exhibited in Mice with Different Susceptibilities to Infection by
Methicillin-Resistant Staphylococcus aureus. Immunity *21*, 215–226.

Uciechowski, P., and Dempke, W.C.M. (2020). Interleukin-6: A Masterplayer in the Cytokine
Network. Oncology *98*, 131–137.

Vono, M., Lin, A., Norrby-Teglund, A., Koup, R.A., Liang, F., and Loré, K. (2017). Neutrophils
acquire the capacity for antigen presentation to memory CD4+ T cells in vitro and ex vivo. Blood *129*,
1991–2001.

Wang, Q., and Zhang, L. (2020). Broadly neutralizing antibodies and vaccine design against HIV-1
infection. Front. Med. *14*, 30–42.

Wang, Y., and Jönsson, F. (2019). Expression, Role, and Regulation of Neutrophil Fcγ Receptors.
Front. Immunol. *10*, 1958.

Wang, J.P., Kurt-Jones, E.A., Shin, O.S., Manchak, M.D., Levin, M.J., and Finberg, R.W. (2005).
Varicella-Zoster Virus Activates Inflammatory Cytokines in Human Monocytes and Macrophages via
Toll-Like Receptor 2. Journal of Virology *79*, 12658–12666.

Wang, X.-Y., Wang, B., and Wen, Y.-M. (2019). From therapeutic antibodies to immune complex
vaccines. Npj Vaccines *4*, 2.

Wareing, M.D., Lyon, A.B., Lu, B., Gerard, C., and Sarawar, S.R. (2004). Chemokine expression
during the development and resolution of a pulmonary leukocyte response to influenza A virus
infection in mice. Journal of Leukocyte Biology *76*, 886–895.

Wen, Y.-M., Mu, L., and Shi, Y. (2016). Immunoregulatory functions of immune complexes in
vaccine and therapy. EMBO Mol Med *8*, 1120–1133.

Yamamoto, T., Iwamoto, N., Yamamoto, H., Tsukamoto, T., Kuwano, T., Takeda, A., Kawada, M.,
Tsunetsugu-Yokota, Y., and Matano, T. (2009). Polyfunctional CD4+ T-cell induction in neutralizing
antibody-triggered control of simian immunodeficiency virus infection. J. Virol. *83*, 5514–5524.

Yoshida, T., Kobayashi, M., Li, X.-D., Pollard, R.B., and Suzuki, F. (2009). Inhibitory effect of
glycyrrhizin on the neutrophil-dependent increase of R5 HIV replication in cultures of macrophages.
Immunol Cell Biol *87*, 554–558.

    FIGURE LEGENDS

Figure 1: **Phenotypic and functional activation of neutrophils stimulated with FrCas^E virions**

and ICs. A. *Experimental scheme*. **B**. *Phenotypic activation of neutrophils stimulated by FrCas^E*

virions or viral ICs made with the 667 mAb. V, free virions; IC, viral ICs; M, culture medium. Free

667 mAb was used as control. Activation was assessed by monitoring CD11b expression and

frequency of CD11b^{hi} CD62L^{lo} neutrophils. The data represent 12 independent experiments. **C**.

Functional activation of neutrophils stimulated by FrCas^E virions (V) or viral ICs made with the 667

mAb (IC). Chemokines release was assessed in supernatants of neutrophils isolated from BM of naive

mice (>97-98% purity) and stimulated for 24 h by FrCas^E virions (red) or viral ICs (blue) or left

unstimulated (grey). The data represent 5 independent experiments. Data are expressed as means +/-

SEM. Statistical significance was established using a parametric 1-way ANOVA test (*p < 0.05; **p <

0.01; ***p < 0.001).

Figure 2: **Phenotypic and functional activation of monocytes stimulated with FrCas^E virions and**

ICs. **A.** *Experimental scheme*. **B**. *Phenotypic activation of monocytes stimulated by FrCas^E virions or*

viral ICs made with the 667 mAb. Free 667 mAb was used as control. V, free virions; IC, viral ICs; M,

culture medium. Activation was assessed by monitoring CD86 expression. The data represent 7

independent experiments **C**. *Functional activation of monocytes stimulated by FrCas^E virions (V) or*

viral ICs made with the 667 mAb (IC). Chemokines release was assessed in supernatants of monocytes

isolated from BM of naive mice (>97-98% purity) and stimulated for 24 h by FrCas^E virions (red) or

viral ICs (blue) or left unstimulated (grey). The data represent 5 independent experiments. Data are

expressed as means +/- SEM. Statistical significance was established using a parametric 1-way

ANOVA test (*p < 0.05; **p < 0.01; ***p < 0.001).

Figure 3: **Cytokine priming potentialize the functional activation of neutrophils and monocytes**

by ICs. BM-derived neutrophils and monocytes were isolated from naive mice and activated as in

Figures 1 and 2 in the presence, or in the absence, of *TNFα, IFN-I or IFNγ.* **(A, C).** *Phenotypic*

activation of activated neutrophils (A) and monocytes (C) by TNFα, IFN-I or IFNγ. Activation was

assessed by monitoring CD11b **(A)** and CD86 expression **(C)**. *Modulation of the functional activation*

of neutrophils and monocytes by TNFα, IFN-I or IFNγ. **(B, D).** Chemokines and cytokines release

were assessed in supernatants of activated neutrophils and monocytes. Histograms depict the increase

in cytokine/chemokine release upon priming of virus- and IC-activated neutrophils by *TNFα, IFN-I or*

IFNγ (calculated as the difference in the amount of chemokine/cytokine secretion by V- or IC-

stimulated neutrophils in the absence or in the presence of priming; **Supplemental figure 3**). Only

those chemokines/cytokines significantly enhanced by cytokine priming are depicted. The data

represent 6 independent experiments for **A**, 4 independent experiments for **B** and 3 independent experiments for **C** and **D**. Data are expressed as means +/- SEM. Diamonds indicate significant differences to all the other stimulation conditions (black diamond) or to corresponding medium without virus or IC stimuli (grey diamond) as determined by Kruskal-Wallis tests ($p < 0.05$).

Figure 4: **FcγRIV is upregulated by IFN priming on both neutrophils and monocytes cell surface.** Neutrophils and monocytes were isolated from naive mice and stimulated as in **Figure 3**. FcγRIV expression was evaluated on neutrophils **(A, B)** and monocytes **(C, D)** in absence **(A, C)** or in the presence of TNFα, IFN-I or IFNγ **(B, D)**. The data represent 5 independent experiments for neutrophils (**A, B**) and 3 independent experiments for monocytes (**C, D**). Data are expressed as means +/- SEM. Diamonds indicate significant differences to the corresponding stimuli in the absence of cytokine priming (open diamond) or to corresponding medium without virus or IC stimuli (grey diamond) as determined by Kruskal-Wallis tests ($p < 0.05$).

Figure 5: **FcγRIV is upregulated *in vivo* on two different splenic neutrophils subpopulations and on inflammatory monocytes.** Splenocytes from naive (grey), infected/non-treated (I/NT; red) and infected/treated (I/T;blue) mice were analyzed on days 8 (D8) and 14 (D14) p.i. for (i) neutrophils and monocytes recruitment and (ii) for FcγRIV expression. **(A)** Gating strategy used to define neutrophils and monocytes populations. **(B)** Frequencies of *CD11b+Ly6G^hi* and *CD11b+Ly6G^int* neutrophils subpopulations as well as *Ly6C^hi* monocytes in the CD45.2+ leukocytic population. The data represent 5 independent experiments at D8 p.i. and 6 independent experiments at D14 p.i with at least 6-9 mice per group (I/NT and I/T) and 3-5 mice per group (naive mice). **(C-D)** FcγRIV expression on *CD11b+Ly6G^hi* and *CD11b+Ly6G^int* neutrophils subpopulations and *Ly6C^hi* monocytes. The data represent 5 independent experiments at D8 p.i. and 6 independent experiments at D14 p.i with at least 6-9 mice per group (I/NT and I/T) and 3-5 mice per group (naive mice). **(E)** Cell surface markers expression of *CD11b+Ly6G^hi* and *CD11b+Ly6G^int* neutrophils and *Ly6C^hi* monocytes isolated from 15-days old naive mice, with 3-5 mice per group. Data are expressed as means +/- SEM. Statistical significance was established using a parametric 1-way ANOVA test (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$).

Figure 6: **Cell surface activation markers expression on two different splenic neutrophils subpopulations and on inflammatory monocytes.** Splenocytes from naive (grey), infected/non-treated (I/NT; red) and infected/treated (I/T;blue) mice were analyzed on days 8 **(A)** and 14 p.i. **(B)** for neutrophils and monocytes expression of CD11b, CD86 and MHC-II. Data represent 5 independent experiments at D8 p.i. and 6 independent experiments at D14 p.i with at least 6-9 mice per group (I/NT and I/T) and 3-5 mice per group (naive mice). Data are expressed as means +/- SEM. Statistical

significance was established using a parametric 1-way ANOVA test (*p < 0.05; **p < 0.01; ***p <
0.001).

Figure 7: **Cytokine and chemokine secretion profile of two different splenic neutrophils**
subpopulations and on inflammatory monocytes. Splenic neutrophils subpopulations and
inflammatory monocytes from naive, infected/non-treated (I) and infected/treated mice (IT) were
isolated at days 8 p.i. and 14 p.i. for assaying their chemokine and cytokine secretion profile in
supernatants of sorted cells cultured at a density of $4x10^6$ cells/ml ($1x10^6$ cells/well) for 24 h. Data
represent 3 independent experiments on D8 p.i. and 4 independent experiments on D14 p.i. with at
least 6-8 mice per group. (**A**) Total quantity of chemokines and cytokines secreted by neutrophils
subpopulations and inflammatory monocytes from infected/non-treated and infected/treated mice. (**B**)
Chemokine and cytokine secretion of neutrophils subpopulations and inflammatory monocytes
isolated from naive, infected/non-treated (I) and infected/treated mice (IT). The color code shows fold-
increase compared to secretion detected in cells isolated from naive mice (the raw chemokine and
cytokine release values are given in brackets, expressed in pg/ml). (**C**) Chemokine and cytokine
secretion of neutrophils subpopulations and inflammatory monocytes isolated from infected/non-
treated (I) and infected/treated mice (IT) expressed as percentage of chemokine and cytokine showing
a fold-increase ≥1.4 as compared to cells isolated from naive mice.

SUPPLEMENTAL FIGURE LEGENDS
Supplemental Figure 1. **Functional activation of neutrophils stimulated with viral determinants**
or LPS. BM-derived neutrophils were isolated from naive mice and stimulated for 24h *in vitro* with
viral determinants (virus or ICs) or LPS (1μg/ml). **A**. *Cytokine secretion profile of neutrophils*
stimulated by virus (red) o viral ICs (blue) or unstimulated (grey). Cytokine release was assessed in
supernatants of neutrophils isolated from BM of naive mice (>97-98% purity). Data represent 5
independent experiments and are expressed as means +/- SEM. Statistical significance was established
using a parametric 1-way ANOVA test (*p < 0.05; **p < 0.01; ***p < 0.001). **B-D**. *Functional*
activation of neutrophils stimulated by LPS. Activation was assessed by monitoring CD11b expression
and frequency of CD11b^{hi} CD62L^{lo} neutrophils (**B**) as well as the cytokines (**C**) and chemokines (**D**)
released in supernatants of neutrophils stimulated for 24 h by LPS (green) or left unstimulated (grey).
Data represent 12 independent experiments and are expressed as means +/- SEM. Statistical
significance was established using a paired Student's t test (*p < 0.05; **p < 0.01; ***p < 0.001).

Supplemental Figure 2: **Functional activation of monocytes stimulated with viral determinants**
or LPS. BM-derived monocytes were isolated from naive mice and stimulated for 24h *in vitro* with
viral determinants (virus or IC) or LPS (1μg/ml). **A**. *Cytokine secretion profile of monocytes*

stimulated by virus (red) o viral ICs (blue) or unstimulated (grey). Cytokine release was assessed in

supernatants of monocytes isolated from BM of naive mice (>97-98% purity) and stimulated for 24 h.

The data represent 6 independent experiments and are expressed as means +/- SEM. Statistical

significance was established using a parametric 1-way ANOVA test (*p < 0.05; **p < 0.01; ***p <

0.001). **B-D**. *Functional activation of monocytes stimulated by LPS*. Activation was assessed by

monitoring the CD86 expression level (**B**) as well as the cytokines (**C**) and chemokines (**D**) release

monitored in supernatants of monocytes stimulated for 24 h by LPS (green) or left unstimulated

(grey). Data represent 5 independent experiments and are expressed as means +/- SEM. Statistical

significance was established using a paired Student's t test (*p < 0.05; **p < 0.01; ***p < 0.001).

Supplemental Figure 3. **Cytokine priming enhance chemokine/cytokine release by IC-activated**

neutrophils and monocytes. BM-derived neutrophils and monocytes were isolated from naive mice

and activated as in **Figures 1 and 2** in the presence, or in the absence, of TNFα, IFN-I or IFNγ.

Chemokines and cytokines release were assessed in supernatants of activated neutrophils (**A**) and

monocytes (**B**). Only those chemokines/cytokines significantly enhanced by cytokine priming are

depicted. The data represent 4 independent experiments for **A** and 3 independent experiments for **B**.

Diamonds indicate significant differences to all the other stimulation conditions (black diamond), or to

corresponding medium without virus or ICs stimuli (grey diamond), or to the corresponding stimuli in

the absence of cytokine priming (open diamond), as determined by Kruskal Wallis test (p < 0.05).

Supplemental Figure 4: **FcγRI and FcγRII-III** *in vitro* **expression on neutrophils and monocytes**

stimulated with viral determinants, in the presence and the absence of TNFα, IFN-I or IFNγ.

FcγRII-III and FcγRI expression was monitored on neutrophils (**A**) and monocytes (**B**). No FcγRI

expression was detected on neutrophils. The data represent 6 independent experiments for **A** et 4

independent experiments for **B**. Data are expressed as means +/- SEM. Diamonds indicate significant

differences from the corresponding stimuli in the absence of cytokine priming (open diamond), or

from all the other stimulation conditions (black diamond) or from corresponding medium without

virus or ICs stimuli (grey diamond) as determined by Kruskal Wallis test (p < 0.05).

Supplemental Figure 5: **FcγRI and FcγRII-III** *in vivo* **expression on two different splenic**

neutrophils subpopulations and on inflammatory monocytes. Splenocytes from naïve (grey,

infected/non-treated (I/NT;red) and infected/treated (I/T; blue) mice were analyzed on days 8 (**A**) and

14 p.i. (**B**) for FcγRII-III and FcγRI expression on $CD11b^+Ly6G^{hi}$ and $CD11b^+Ly6G^{int}$ neutrophils

subpopulations and $Ly6C^{hi}$ monocytes. Data represent 5 independent experiments at D8 p.i. and 6

independent experiments at D14 p.i with at least 6-9 mice per group (I/NT and I/T) and 3-5 mice per

group (naive mice). Data are expressed as means +/- SEM. Statistical significance was established

using a parametric 1-way ANOVA test (*p < 0.05; **p < 0.01; ***p < 0.001).

Figure 1

A

B

C

Figure 2

Figure 3

Figure 5

Figure 6

Figure 7