Item recognition and lure discrimination in younger and older adults are supported by

alpha/beta desynchronization

Anna E. Karlsson[1,2], Claudia C. Wehrspaun[1], and Myriam C. Sander[1]

[1] Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin,

Germany

[2] Department of Psychology, Humboldt Universität zu Berlin, Germany

Running head: Item recognition and lure discrimination in younger and older adults

Correspondence concerning this article should be addressed to: Myriam C. Sander,

sander@mpib-berlin.mpg.de, Max Planck Institute for Human Development, Lentzeallee 94,

14195 Berlin

Abstract

Our episodic memories vary in their specificity, ranging from a mere sense of familiarity to

detailed recollection of the initial experience. Recent work suggests that alpha/beta

desynchronization promotes information flow through the cortex, tracking the richness in

detail of recovered memory representations. At the same time, as we age, memories

become less vivid and detailed, which may be reflected in age-related reductions in

alpha/beta desynchronization during retrieval. To understand age differences in the

specificity of episodic memories, we investigated differences in alpha/beta desynchronization

between younger (18–26 years, $n = 31$) and older (65–76 years, $n = 27$) adults during item

recognition and lure discrimination.

Alpha/beta desynchronization increased linearly with the demand for memory specificity, i.e.,

the requirement to retrieve details for an accurate response, across retrieval situations

(correct rejections < item recognition < lure discrimination). Stronger alpha/beta

desynchronization was related to memory success, as indicated by reliable activation

differences between correct and incorrect memory responses. In line with the assumption of

a loss of mnemonic detail in older age, older adults had more difficulties than younger adults

to discriminate lures from targets. Importantly, they also showed a reduced modulation of

alpha/beta desynchronization across retrieval demands. Together, these results extend

previous findings by demonstrating that alpha/beta desynchronization dissociates between

item recognition and the retrieval of highly detailed memories as required in lure

discrimination, and that age-related impairments in episodic retrieval are accompanied by

attenuated modulations in the alpha/beta band. Thus, we provide novel findings suggesting

that alpha/beta desynchronization tracks mnemonic specificity and that changes in these

oscillatory mechanisms may underlie age-related declines in episodic memory.

Keywords: aging, alpha/beta oscillations, EEG, episodic memory, lure discrimination

1. Introduction

Episodic memory, the ability to remember events from the past, allows us to mentally travel

back in time to re-visit places and people we have encountered (Tulving, 1985). It is crucial

for human cognition and behavior, since it allows us to learn from experience and to detect

novel elements in our surroundings. Memory retrieval can range from a mere sense of

familiarity, that is, when we fail to recover specifying details but experience the event as

familiar, to the recollection of highly detailed memories accompanied by the retrieval of

specifying information (Jacoby, 1998; Mandler, 1980; for review, see Yonelinas, 2002). With

advancing age, the ability to retrieve episodic memories declines (Naveh-Benjamin et al.,

2003; Nyberg et al., 2012) and older adults' memories seem to be less vivid and detailed

(Folville et al., 2020; Greene & Naveh-Benjamin, 2020) than younger adults'. Higher levels of

memory errors in older adults indicate the importance of retrieving memories with a high level

of detail (Bowman et al., 2019; Fandakova et al., 2018; Koutstaal & Schacter, 1997).

Nevertheless, the neural mechanisms underlying the loss of memory specificity with

advancing age are still poorly understood.

A growing body of research has established the involvement of neural oscillations in

memory processes (Düzel et al., 2010; Hanslmayr et al., 2016; Herweg et al., 2020; Nyhus &

Curran, 2010). Brain oscillations provide a temporal window for the synchronous inhibitory or

excitatory activity of neural cell assemblies. Such synchronous activity shapes synaptic

weights within these assemblies (Markram et al., 1997) and thereby serves information

processing related to memory formation and retrieval. Recent work has demonstrated that

successful memory encoding and retrieval are associated with decreases in power in the

alpha/beta frequency range (8–20 Hz; Griffiths et al., 2019; Waldhauser et al., 2016; for

review, see Hanslmayr et al., 2012), reflecting the desynchronization of neural activity

(Buzsáki et al., 2012). According to recent theoretical and empirical work, alpha/beta

desynchronization is critical for efficient information processing (Hanslmayr et al., 2012;

Parish et al., 2018) serving as a gating mechanism allowing for the flow of information

through relevant neural assemblies. In support of this notion, alpha/beta desynchronization

has been shown to reflect the level of processing at encoding (Hanslmayr et al., 2009; for

review, see Hanslmayr & Staudigl, 2014) and seems to track the reactivation of mnemonic

content at retrieval. Specifically, alpha/beta desynchronization was linearly related to the

amount of information retrieved (Khader & Rösler, 2011; Waldhauser et al., 2016) and to the

fidelity of information represented in cortex (Griffiths et al., 2019). Importantly, alpha/beta

desynchronization does not seem to represent mnemonic information per se, but initiates

neural conditions beneficial for cortical reinstatement, presumably by promoting information

flow to the neural substrates conveying information about mnemonic content. Accordingly,

two recent studies have reported that alpha/beta desynchronization increased systematically

across conditions characterized by different levels of retrieval demands. In these studies,

alpha/beta desynchronization showed a linear increase across item recognition and

associative memory (Martín-Buro et al., 2020; Staresina et al., 2016), suggesting that the

level of desynchronization might vary depending on the level of mnemonic detail needed for

a correct response. In turn, the level of alpha/beta desynchronization may be considered as

an index of the specificity of a recovered memory trace. However, the extent to which

alpha/beta desynchronization is modulated by the amount of detail ingrained in a memory

trace is still unclear, advocating the investigation of situations in which the retrieval of highly

detailed, specific memories is needed, e.g., when the correct response depends on the

detection of subtle changes in a test item.

The neural mechanisms supporting the ability to discriminate a similar lure from a

previously encountered old item have typically been investigated in studies that extended the

traditional old/new recognition memory paradigm by including similar lures in the test list,

targeting the ability to remember details (Stark et al., 2019). These studies have provided

evidence for the necessity to lay down orthogonalized memories in order to reduce

interference, a process called pattern separation that relies primarily on the hippocampus (for

reviews, see Stark et al., 2019; Yassa & Stark, 2011). However, other brain regions have

also been shown to be important for the detection of similar lures. Specifically, regions

throughout the ventral visual stream as well as earlier visual regions have been shown to

dissociate between targets and perceptually and/or semantically similar lures (Bowman et al.,

2019; Chouinard et al., 2008; Kim et al., 2009; Pidgeon & Morcom, 2016). Given the known

role of these regions in the reactivation of mnemonic content (e.g., St-Laurent et al., 2015;

Wing et al., 2015), successful target–lure discrimination presumably also relies on the

recovery of fine-grained memory representations. As elaborated above, the level of

alpha/beta desynchronization during memory retrieval may relate to the detailedness of

episodic memories, in particular, and may reflect underlying differences in memory

specificity. Thus, we predicted that alpha/beta desynchronization contributes not only to

successful item recognition, as indexed by power differences between new and old

information, but also supports the discrimination between highly similar mnemonic

information underlying successful lure discrimination.

While the discrimination between highly similar mnemonic information is usually a

difficult, yet feasible task for younger adults, older adults often show an impairment in tasks

that require the retrieval of detailed information. Age differences are rarely observed when

simple item recognition is sufficient, but older adults show difficulties when the retrieval of

episodic detail is paramount to the task at hand, such as context or source information (e.g.,

Bastin et al., 2014; Luo & Craik, 2009; for review, see Old & Naveh-Benjamin, 2008).

Similarly, the ability to discriminate similar lures from old items, which relies on the ability to

recover memories with high specificity, is negatively affected by age (Stark et al., 2015; Stark

& Stark, 2017; for review, see Stark et al., 2019) and is accompanied by an increased

reliance on gist information and familiarity-based processes when making mnemonic

judgments (Kensinger & Schacter, 1999; Koen & Yonelinas, 2014; Stark et al., 2015). The

loss of mnemonic detail and reduced specificity of memory content in older adults is also

reflected in age differences in the specificity of neural activation, so-called neural

dedifferentiation (Bowman et al., 2019; Koen et al., 2019; Sommer et al., 2020; for review,

see Koen & Rugg, 2019). Early functional magnetic resonance imaging (fMRI) studies

demonstrated that while in younger adults, processing of category-specific information, such

as faces or houses, is supported by category-selective regions (fusiform face area and

parahippocampal place area respectively), older adults show a decrease in this neural

selectivity (Park et al., 2004). Recent studies using multivariate pattern analysis support this

initial finding and underline the relevance of neural specificity for memory performance

(Kobelt et al. 2020; Koen et al., 2019; Zheng et al., 2018). Age differences in oscillatory

mechanisms of memory formation (Werkle-Bergner et al., 2006) and retrieval may further

contribute to the reduced memory specificity in older adults, but remarkably few studies have

addressed this question as yet. In particular, given the evidence that the level of alpha/beta

desynchronization during retrieval may track differences in memory specificity, a reduction in

alpha/beta desynchronization in older adults during retrieval may be related to the observed

memory decline. Among the very few studies investigating age differences in oscillatory

underpinnings of episodic memory, a recent study showed that alpha/beta desynchronization

during learning discriminates between subsequently remembered and subsequently

forgotten item–picture associations in both younger and older adults (Sander et al., 2020;

see also Strunk & Duarte, 2019). Importantly, the size of the subsequent memory effect

depended on the structural integrity of the inferior frontal gyrus, a region that is known to be

relevant for deep elaboration of information (Hanslmayr et al., 2011; Kim, 2011), and whose

integrity was reduced in the group of older adults. Thus, the study by Sander and colleagues

(2020) supports the notion that episodic memories differ between younger and older adults

as a result of age differences during encoding (Craik, 2020). Regarding retrieval, alpha/beta

desynchronization has been shown to distinguish between correct rejections of novel foils

and correct pair memory judgments in both younger and older adults (Strunk et al., 2017).

Together, these studies suggest that alpha/beta desynchronization, as a signature of the

specificity of memories laid down at encoding and recovered at retrieval, represents a

promising candidate for understanding age differences in episodic memory.

To summarize, we put forward the idea that the specificity of episodic memories

determines the level of alpha/beta desynchronization during retrieval. Based on the

observations that desynchronization in the alpha/beta frequency range is related to the level

of processing at encoding (Hanslmayr & Staudigl, 2014) and to the quality of the recovered

memory trace at retrieval (Griffiths et al., 2019), we hypothesized that also alpha/beta

desynchronization at retrieval varies with retrieval demand, i.e. as a function of the level of

mnemonic detail required to make an accurate memory judgment. The distinction between

previously learned information and newly encountered information should be reflected in

stronger alpha/beta desynchronization for old responses compared to new responses (see

Martín-Buro et al., 2020). An even more fine-grained mnemonic representation is necessary

when new information is highly similar to the learned information. Therefore, correct "similar"

responses in a recognition memory task should be accompanied by even stronger alpha/beta

desynchronization than correct "old" responses. Thus, we predicted a linear relationship

between alpha/beta desynchronization at retrieval and the level of mnemonic detail required

to make an accurate memory judgment (New < Old < Lure). In accordance with evidence for

an age-related reduction in episodic detail of retrieved memories, we predicted age

differences between younger and older adults when the discrimination between highly similar

information is required, i.e., during lure discrimination. Furthermore, we hypothesized that

older adults' reduced memory performance would be accompanied by smaller modulations in

alpha/beta desynchronization across experimental situations with varying retrieval demands.

To this end, younger and older adults were measured with electroencephalography (EEG)

while performing an episodic memory task. After studying pictures of everyday items,

participants performed a surprise recognition memory test, where they were required to

discriminate between old items, new items, and perceptually similar lures (see Fig.1). This

design allowed us to contrast neural activity between conditions with varying retrieval

demands requiring different levels of mnemonic detail (correctly rejecting a new item,

correctly remembering an old item, and correctly endorsing a lure as similar) in an age-

comparative setting.

2. Methods

2.1. Participants

Younger (n = 41) and older (n = 42) right-handed adults were recruited via a

database of the Max-Planck Institute for Human Development to participate in the study for a

compensation of €12/hour. One older and three younger adults were excluded due to

behavioral performance below chance level (see section 2.5) and one older adult was

excluded based on the MMST (Mini-Mental-Status Test; Folstein et al., 1975; see below). In

addition, after preprocessing of the EEG data, another 7 younger and 13 older adults were

excluded due to too few trials across the trial bins (see section 2.6). The remaining sample

consisted of 31 younger adults (M_{age} = 21.84, SD_{age} = 2.6, range 18–26 years, 16 females)

and 27 older adults (M_{age} = 70.74, SD_{age} = 2.73, range 65–76 years, 17 females).

To further characterize the sample, the participants completed a demographic

questionnaire and a series of test batteries. Older adults were assessed using the MMST

measuring cognitive impairment. A MMST score of 27 or higher indicates normal cognition,

whereas a MMST score of 19–23 indicates mild cognitive impairment (O'Bryant et al., 2008).

One older adult did not pass the cut off (MMST score = 23) and was excluded from further

analysis. Although the MMST is not sufficient to with certainty rule out mild cognitive

impairment, it is noteworthy that the remaining sample scored relatively high above the cut

off (range: 28-30), and thus most likely represents relatively healthy older adults. All

participants completed the Digit Symbol substitution test (DST; Wechsler, 1955). Younger

adults showed a significantly higher DST score (M = 74.67; SD = 10.76) than older adults (M

= 51.41; SD = 10.66; W = 787, α = 0.05, p < .001). Working memory (WM) was tested using

the WM digit-sorting task (Kray, 2000), and younger adults (M = 9.77; SD = 2.64) did not

score significantly higher than older adults (M = 8.44; SD = 3.04; W = 515, α = 0.05, p = .13).

Performance in the marker tasks was similar to other cognitive neuroscience studies

previously run at our research center (e.g., Fandakova et al., 2014; Sander et al., 2011) and

showed a typical pattern of age differences. The ethics committee of the Max Planck Institute

for Human Development, Berlin, approved the study.

2.2. Stimuli

The experiment and recording of behavioral responses were programmed in

MATLAB (version 2016b; The MathWorks Inc., Natick, MA, USA), using the Psychophysics

Toolbox (Brainard, 1997). The stimulus pool consisted of 250 real world asymmetric objects

(500 x 500 pixels) from a publicly available data base (Brady et al., 2008). They were

presented on the center of the screen.

2.3. Experimental paradigm

The experiment was performed in a dimly lit room that was electromagnetically and

acoustically shielded. Prior to the task, participants received instructions and completed 15

practice trials that could be repeated if necessary. In addition, each experimental session

started with written instructions on the screen and the participant initiated each session with

a button press. Prior to starting the memory task, participants engaged in five minutes of

wakeful rest. During encoding (Fig. 1), randomly drawn objects were presented against (*i*) a

white background (50 stimuli), (*ii*) a background that flickered between white and black at a

frequency of 36 Hz (50 stimuli), or (*iii*) a background that flickered between white and black

at a frequency of 28.8 Hz (50 stimuli). Before the experiment, participants were informed of

the flickering background and instructed to pay no attention to it. Effects of the flickering

background at encoding are beyond the scope of this manuscript and will be ignored for the

analysis of the recognition phase. However, to test for any effects of the flickering

background on behavioral performance, we computed an ANOVA with flicker (36/28.8 Hz)

and behavioral measure (lure discrimination/corrected recognition; see section 2.5 for the

calculation of these measures) as within-subject factors, and age (older/younger) as

between-subject factor. No significant main effect of (F(1,168) = 0.18, *p* = .675) or

interactions with (*p* > .250) flicker were found. Therefore, the retrieval data were collapsed

across flicker frequencies for the analysis.

Each encoding trial began with a black screen with a centered white fixation cross

(henceforth referred to as ISI) with a jittered presentation time (1–1.5 s). Subsequently, the

object was presented (2 s) followed by first another ISI (0.7 s) and second the presentation of

a retro-cue (0.5 s), indicating either to remember (green dot), or not to remember (red dot)

the previously shown item. The presentation times were chosen based on previous work that

successfully used retro-cues (Mok et al., 2016). An ANOVA with cue (remember/don't

remember) and behavioral measure (lure discrimination/corrected recognition) as within-

subject factors, and age (younger/older) as between-subject factor was used to assess

whether cue condition had any effects on behavioral performance. No significant main effect

of (F(1,168) = 2.99, p = .085) or interactions with (p > .320) cue were found and thus the data

were collapsed across cue condition for the purpose of the current analysis.

After the encoding phase, participants completed a short sham recognition test aimed

to prevent active rehearsal during the following 10-minute break. For ten randomly selected

items (i.e., four new items and six old items that had been followed by a green retro cue),

participants had to indicate whether the item was "old" (i.e., shown during the encoding

phase), "mirrored" (i.e., the mirrored image was shown during the encoding phase), or "new"

(i.e., the item had not been seen before). Following the sham recognition, subjects were

randomly assigned to one of two different 10-minute break conditions, either engaging in

wakeful rest (n = 30) while fixating a black screen with a centered white fixation cross or

performing a distraction task (n = 28) during which they were instructed to imagine walking in

a well-known public area in Berlin while imagining the scenery as vividly as possible. In line

with previous evidence (Dewar et al., 2012), we expected higher performance after wakeful

rest than after a distraction task. An ANOVA with behavioral measure (lure

discrimination/corrected recognition) as within-subject factor, and rest (wakeful

rest/distraction task) and age (younger/older) as between-subject factors, yielded no

significant main effect of rest (F(1,54) = 0.82, p = .368) but a significant behavioral measure x

rest x age interaction (F(1,54) = 5.74, p = .020, η^2 = 0.10). However, no pairwise

comparisons were significant (p > .083; see also Varma et al., 2017). We therefore collapsed

across both groups in the current analysis.

Following the break, participants performed a recognition task including all remaining

objects (n = 140) that had been shown during the encoding phase. These were randomly

presented intermixed with pictures of new objects (n = 100). For 20% of the old objects (i.e.,

n = 28), the respective mirror image was shown and participants were asked to indicate by

key press whether the object was (*i*) old, (*ii*) mirrored, or (*iii*) new (Fig. 1; Motley & Kirwan,

2012). Each trial started with a jittered ISI (1–1.5 s) followed by the presentation of an object

(2 s). To prevent potential confounds of motor responses in the EEG signal, participants

withheld their response until a subsequently presented screen provided the response

alternatives and their mapping onto the cursor keys on the keyboard. Responses were made

with the right hand and the next trial began when a response was given. Given that

participants had to withhold their responses, the use of response times as indicators of

cognitive dynamics is clearly limited. Accordingly, an ANOVA on median response times with

retrieval condition as within-subject factor (old/new/lure) and age (younger/older) as

between-subject factor yielded no significant effects (all ps > .067).

Figure 1. Experimental paradigm. During the encoding phase participants viewed different
objects. Participants were instructed to either remember the object if it was followed by a
green retro cue, or to not remember the object, if followed by a red retro cue. Objects were
presented on flickering or white backgrounds. Note that neither the flicker nor the retro cue
manipulations are relevant to the current analysis. After a 10-minute break, participants were
asked to take a surprise memory recognition test. In this test, old objects either identical to
the presentation during encoding or as their own mirror image (20% of the old objects) were
presented together with completely new objects. Participants had to indicate whether the
presented object was old, mirrored, or new.

2.4. EEG recording and preprocessing

EEG data were continuously recorded with BrainVision Recorder (BrainVision

Products GmbH, Gilching, Germany) using elastic caps with 61 Ag/Ag–Cl passive electrodes

placed according to the 10-10 system. AFz was used as ground electrode and the right

mastoid served as reference during recording. Additional electrodes were placed on the left

mastoid for later re-referencing, and on the outer canthi (horizontal EOG) and below the left

eye (vertical EOG) to monitor eye movements. Electrode preparation ensured impedances

below 5 kΩ before recording. A band-pass filter of 0.1 to 250 Hz was applied and data were

recorded with a sampling rate of 1000 Hz and a sampling interval of 1000 μS. During

recording, EEG was monitored online, and channels with extremely noisy signal or excessive

impedance were marked to be excluded from further analysis. In addition, heart rate was

recorded via electrocardiogram (ECG) to remove possible cardiovascular artifacts from the

EEG signal (see below).

The EEG data were preprocessed and analyzed with the FieldTrip toolbox (Oostenveld et

al., 2011) and custom-written MATLAB code. Offline, EEG data were filtered (4th filter order)

with a pass-band of 1–125 Hz, and re-referenced to the linked mastoid channels. The ECG

data were filtered with a band-stop filter (48–52 Hz) and appended with the preprocessed

EEG data. Next, the data were segmented into 6-second epochs (-2000–4000 ms) and all

trials were visually inspected. Trials with extreme artifacts were removed for the ICA only.

Saccade-related transient spike potentials were identified using the COSTRAP algorithm and

removed from the signal as independent components (Hassler et al., 2011). Blink, eye-

movement, muscle, and heartbeat artifacts were detected using ICA (Bell & Sejnowski, 1995)

and removed from the signal. Artifact-contaminated channels and trials (determined across

all epochs) were automatically identified (1) using the FASTER algorithm (Nolan et al., 2010)

and (2) by detecting outliers exceeding four standard deviations of the kurtosis of the

distribution of power values in each epoch within low- (0.2–2 Hz) or high-frequency (30–100

Hz) bands respectively. Channels labeled as artifact-contaminated were interpolated using

spherical splines (Perrin et al., 1989). Finally, a second visual inspection of each trial per

participant was performed and any remaining artifact-contaminated trials were excluded from

further analysis, resulting in an average of 15.3% of trials rejected. The preprocessed EEG

data were subjected to time-frequency decomposition of each epoch (-2000–4000 ms, 50 ms

sliding window) using a Morlet wavelet approach (wavelet width = 7) as implemented in

FieldTrip, estimating spectral power from 1 to 20 Hz in steps of 1 Hz. Finally, data was

segmented into the time-window of interest (0–2000 ms) and the power spectrum was log

transformed on the single trial level (Smulders et al., 2018). Note that no baseline correction

was applied to the data.

2.5. Behavioral analysis

To determine performance above chance level, a balanced accuracy score was

calculated by taking the mean over the proportion of correct responses to new items, old

items, and lures respectively, yielding a chance level of 33%.

Responses were categorized as *old hits* (responding "old" to an old item), *old misses*

("new"/"mirrored" to an old item), *lure hits* ("mirrored" to a lure), *lure misses* ("old" to a lure),

correct rejections (CR; correctly rejecting a new item), and *false alarms* (FA; endorsing a new

item as "old"). The different proportions of responses given to the three test item categories

are presented in Fig. 2. To assess younger and older adults' ability to recognize an old item

while correcting for response bias, we computed corrected recognition scores by subtracting

FA rates from old hit rates for each participant (Snodgrass & Corwin, 1988). Similarly, to

evaluate participants' abilities to discriminate mirrored lures from old items, we computed a

lure discrimination index (LDI) by subtracting the lure miss rate from the lure hit rate

("Mirrored"|Lure – "Old"|Lure; Ngo et al., 2018; Toner et al., 2009).

To investigate age differences in item recognition and lure discrimination, we

performed a mixed effects ANOVA with age (younger/older) as between-subject, and

measure (corrected recognition/LDI) as within-subject factors. Any significant interactions

were followed up with pairwise comparisons. Bonferroni-corrected p-values are reported.

2.6. EEG group level analysis

To quantify the differences in power between conditions of different retrieval

demands, trials associated with correctly rejecting a new item (CR), correctly recognizing an

old item (old hits), and correctly discriminating a lure (lure hits), were binned and averaged

within participants. Participants with less than a minimum of 6 trials per bin were excluded

from the analysis (see section 2.1 for sample information), resulting in an average of 84.9

(range: 68–94) and 82 (range: 59–93) CR trials, 72.5 (range: 28–108) and 82 (range: 35–

109) old-hit trials, and 12.6 (range: 6–26) and 11.1 (range: 6–20) lure-hit trials for younger

and older adults respectively.

The averaged time-frequency power values were subjected to a non-parametric,

cluster-based, random permutation test (FieldTrip toolbox; Maris & Oostenveld, 2007),

contrasting the time-frequency power spectrum between conditions with different retrieval

demands (CR < old hits < lure hits) at each channel, frequency band (2–30 Hz), and time

point (0–2 s). Initially, clusters were formed based on univariate two-sided, dependent

samples regression coefficient *t*-statistics for each electrode–frequency–time point. The

threshold for data points to be included in a cluster was set to $p = .05$ and the spatial

constraint was set to a minimum of two neighboring channels. Next, the significance of the

cluster-level statistic (i.e., the summed t-values) was assessed by comparison to a

permutation null-distribution, which was obtained by randomly switching the condition labels

and re-computing the *t*-test 1000 times. The final cluster *p*-value (i.e., the Monte Carlo

significance probability) is the proportion of random partitions in which the cluster-level

statistics were exceeded. The cluster-level significance threshold was set to *p*-values below

.025 (two-sided significance threshold). This analysis yielded one significant negative cluster

in the alpha/beta frequency range, comprising most channels from ~600–2000 ms (see

section 3.2). This electrode–frequency–time cluster was used for the following subsequent

analyses:

1. To explore potential age differences in the strength of the observed alpha/beta

desynchronization effect (i.e., the slope), we computed individual linear

regressions and averaged the individual regression coefficients within the

electrode–frequency–time cluster that was identified across younger and older

adults (see above). We tested for differences between the age groups using an

independent samples *t*-test.

2. To examine the relevance of the level of alpha/beta desynchronization for memory

performance, we contrasted neural activity associated with successful and

unsuccessful responses for item recognition and lure discrimination. To this end, we

extracted and averaged the individual log-transformed power values within the

identified time–frequency–electrode cluster for each condition of interest (i.e., old hits,

old misses, lure hits, and lure misses) and participant. Power values were entered in

two ANOVAs with age (younger, older) as between-subject, and accuracy (Hit, Miss)

as within-subject factors for old and lure items respectively. Significant interactions

were followed-up with pairwise comparisons. Reported p-values are Bonferroni-

corrected.

Note that while the electrode–frequency–time cluster was based on trials with

correct responses only, this analysis contrasts correct and incorrect responses to

assess the behavioral relevance of the effect. We would like to point out that whereas

lure misses includes only "old" responses and thus allows us to dissociate lure

discrimination from item recognition, old misses includes all incorrect responses (i.e.,

"new" and "mirrored"). Although "mirrored" and "new" responses to old items most

likely involve different cognitive processes (i.e. familiarity and the absence of

familiarity respectively; see Yonelinas, 2002), both responses are characterized by an

absence of recollection. Thus, by categorizing "mirrored" and "new" responses as

misses, the contrast aim to target pure recognition as compared to familiarity-based

memory.

Some participants contributed fewer than 6 trials per condition and were

excluded from this analysis, resulting in a sub-sample of 45 participants (23 younger

adults, 22 older adults) with an average of 75.3 (range: 47–108) and 82.3 (range: 35–

105) old-hit trials, 33 (range: 6–62) and 27.4 (range: 6–76) old-miss trials, 11.7

(range: 6–20) and 10.6 (range: 6–17) lure-hit trials, and 8.9 (range: 6–17) and 12.7

(range: 7–20) lure-miss trials for younger and older adults respectively.

Data Accessibility. Data and custom Matlab code of the main analyses are available on

the Open Science Framework repository (https://osf.io/dt4ax/).

3. Results

3.1. Older adults show reduced lure discrimination

To investigate age differences in item recognition and lure discrimination, a mixed

ANOVA with age (younger, older) as between, and performance measure (corrected

recognition, LDI) as within-subject factors was implemented. The results (see Fig. 2)

demonstrated a marginally significant main effect of age ($F_{(1;56)}$ = 3.82; p = 0.056, $np2$ =

0.1) and a main effect of measure ($F_{(1;56)}$ = 249.19; p < .001, $np2$ = 0.82), reflecting lower

performance in older (*M* = 0.33, *SD* = 0.41) than younger (*M* = 0.42, *SD* = 0.31) adults as

well as better recognition memory (*M* = 0.66, *SD* = 0.18) than lure discrimination (*M* = 0.1,

SD = 0.28). In addition, a significant age x measure interaction was found ($F_{(1;56)}$ = 13.81; *p*

< .001, *np2* = 0.2), reflecting comparable recognition memory across age groups (*t*(51.8) = -

0.9, *p* > .250) but reduced lure discrimination in older compared to younger adults (*t*(55.71) =

3.3, *p* = 0.003, d = 0.86).

Figure 2. A: Response rates (y-axis) for the different responses, "Old" (green), "New"
(purple), and "Mirrored" (red), across the different test item categories (x-axis) for younger
(top) and older (bottom) adults. Bars represent the means with error bars showing standard
deviations. Dots indicate individual data points. B: Illustration of the age difference evident for
lure discrimination. Memory performance score (y-axis) for the two memory measures
(corrected recognition, lure discrimination index; x-axis) across age groups (younger in blue,
older in orange). Boxplots represent the interquartile range (first and third quantile) with dots
representing individual participants. The rhombus indicates the mean, the horizontal bar
indicates the median, and violin plots illustrate the sample density. The asterisk reflects
significance below .05.

3.2. Alpha/beta desynchronization varies as a function of retrieval demand

Time-frequency representations (Fig. 3A) of the retrieval-related neural activity averaged

over posterior channels (TP7, CP5, CP3, CP1, CPz, CP2, CP4, CP6, TP8, P7, P5, P3, P1,

Pz, P2, P4, P6, P8, PO7, PO3, POz, PO4, PO8, O1, Oz, O2) display a clear increase in the

theta frequency range (~3–7 Hz) following stimulus onset. This early increase in power is

followed by a strong power decrease in the alpha/beta frequency range (~8–20 Hz). A similar

oscillatory pattern is evident across the three conditions with different retrieval demands. In

order to test for differences in alpha/beta desynchronization between the conditions, we

conducted linear regressions across levels of retrieval demand (correct rejections < item

recognition < lure discrimination), collapsed across age groups. A negative cluster was

identified, reflecting a reliable linear relationship between alpha/beta desynchronization and

retrieval demand ($p < .001$; see Fig. 3B & D). This negative cluster showed a wide-spread

topographical distribution with a maximum over posterior channels (see Fig. 3C), temporally

encompassed most of the epoch (~0.6–2 s), and comprised a large frequency range within

the alpha and beta frequency bands (~8–25 Hz). To assess age differences in the strength of

the observed alpha/beta desynchronization effect (i.e., the slope), we compared younger and

older adults' regression coefficients within the electrode–frequency–time cluster. Importantly,

older and younger adults differed with regard to the strength of the effect ($t(54.51) = -2.35$, *p*

$= 0.022$, d = 0.62), reflected in a steeper slope in younger (*M* = -0.82, *SD* = 0.41) as

compared to older (*M* = -0.56, *SD* = 0.42) adults (see Fig. 3E).

Figure 3. A: Time (x-axis) – frequency (y-axis) representations of log-transformed power for conditions of different retrieval demands (correct rejections, item recognition, lure discrimination) averaged over participants. Note that the data is plotted with baseline correction (absolute baseline, -0.5–0.2 s) for illustrative purposes only. For analysis, no baseline correction was applied. B: Results of the cluster-permutation analysis using dependent regression to test for differences in alpha/beta power across retrieval conditions. The highlighted negative time-frequency cluster indicates a reliable linear increase in alpha/beta desynchronization across retrieval demands. C: Topographical distribution of the effect from 400-2000 ms in steps of 200 ms. The effect is wide-spread across all channels but with a posterior maximum throughout the epoch. The asterisks highlight significant channels. D: For visualization purposes only, we display log-transformed power (y-axis) averaged within the negative cluster by age group (x-axis). This illustrates that the linear

increase in alpha/beta desynchronization across retrieval demands is evident in both age
groups. However, as displayed in (E): the slope of alpha/beta desynchronization (expressed
as t-values of the individual linear regression averaged within the cluster) is significantly
steeper in younger (blue) as compared to older (orange) adults. Boxplots represent the
interquartile range (first and third quantile) with dots representing individual participants. The
rhombus indicates the mean, the horizontal bar indicates the median, and violin plots
illustrate the sample density. The asterisk reflects significance below .05.

3.3. Larger alpha/beta desynchronization predicts within-person memory performance

If the observed modulation of alpha/beta desynchronization by retrieval demand and

age group contributes to memory performance, we would expect it to vary with regard to

memory success. Therefore, we contrasted log-transformed power values within the

observed cluster for successful versus unsuccessful item recognition as well as successful

and unsuccessful lure discrimination. To this end, we conducted two separate ANOVAs,

including age group (younger, older) as between-subject, and accuracy (Hits, Misses) as

within-subject factors, for old and lure items respectively (see Fig. 4). For old items, a main

effect of accuracy ($F(1;43) = 6.723$; $p = 0.013$, $np2 = 0.14$) was found, indicating stronger

alpha/beta desynchronization for hits ($M = 2.48$, $SD = 0.24$) as compared to misses ($M = 2.5$,

$SD = 0.25$) regardless of age. However, a significant interaction of age and accuracy ($F(1;43)$

$= 10.32$; $p = 0.002$, $np2 = 0.19$) followed up with pairwise comparisons revealed that

alpha/beta desynchronization was stronger for old item hits as compared to old item misses

in younger adults ($t(22) = -5.02$, $p < .001$, d = 0.19), whereas no significant difference was

evident for older adults ($t(21) = 0.42$, $p > .250$). For lure items, we found a main effect of

accuracy ($F(1;43) = 37.266$; $p < .001$, $np2 = 0.46$), reflecting stronger alpha/beta

desynchronization for hits ($M = 2.42$, $SD = 0.22$) as compared to misses ($M = 2.48$, $SD = $

0.25) regardless of age. The interaction between age group and accuracy was not significant

($F(1;43) = 2.29$; $p > 0.137$).

Figure 4. A: Log-transformed power (y-axis) averaged within the previously determined
cluster (see Fig. 3B) by age group (left: younger adults, right: older adults) and accuracy
(Hits: green, Misses: red). B: For illustrative purposes only, difference scores (Hits – Misses)
as a function of retrieval demand (Old, Lure; x-axis) and age (younger: blue; older: orange),
demonstrates the age differences in the modulation of alpha/beta desynchronization by
retrieval success across retrieval conditions. Boxplots represent the interquartile range (first
and third quantile) with dots representing individual participants. The rhombus indicates the
mean and the horizontal bar indicates the median. Asterisks reflect significance below .05.

4. Discussion

Memory performance depends on the encoding and retrieval of detailed information.

However, our memories are not all equally precise, but vary in their level of specificity,

determining memory success and failure. Becoming older is accompanied by an increased

loss of mnemonic specificity which may contribute to the commonly observed episodic

memory decline (Nyberg et al., 2012; Wang et al., 2017). In light of recent work proposing

that alpha/beta desynchronization relates to the quality or specificity of retrieved memories

(Griffiths et al., 2019; Khader & Rösler, 2011; Martín-Buro et al., 2020), we investigated

whether alpha/beta desynchronization was modulated by different retrieval demands in

younger and older adults. Specifically, we asked whether the level of alpha/beta

desynchronization is not only modulated by the need to detect old (compared to new)

information as in successful item recognition, but also varies with the level of detail required

for the discrimination between targets and lures, which provide highly similar information.

4.1. Alpha/beta desynchronization supports item recognition and lure discrimination

As hypothesized, alpha/beta desynchronization showed a linear relationship with

retrieval demand, that is, the amount of detail required to make an accurate mnemonic

judgment (New < Old < Lure; see Fig. 3.B). This finding supports the notion that the level of

alpha/beta desynchronization reflects the level of mnemonic evidence available during

retrieval and can serve as an index of memory specificity. Our findings parallel those of a

recent study with a sample of young adults only by Martín-Buro and colleagues (2020). That

study showed that alpha power decreases across retrieval conditions that differ in the

amount of information needed for a correct response (correct rejections < item recognition <

associative recall). Using source localization, the authors were able to link the observed

alpha desynchronization to the posterior parietal cortex (PPC). Accordingly, they proposed

that parietal alpha desynchronization may reflect the accumulation of mnemonic evidence

governed by the PPC, serving as an 'episodic buffer' representing recollected information

(Rugg & Vilberg, 2013). Although the effect reported here shows a widespread topography, it

is seemingly maximal over posterior channels (see figure 3.C). Thus, our results extend

these findings by demonstrating that alpha/beta desynchronization discriminates between

item recognition and lure discrimination in a similar manner. Since target–lure discrimination

depends on the detection of subtle differences, the retrieval of a highly detailed, specific

memory representation is necessary for a correct memory judgement. We suggest that this

higher memory specificity is supported by stronger alpha/beta desynchronization in lure

conditions, indexing the larger amount of detail engrained in the recovered memory trace.

Importantly, by contrasting trials associated with correct and incorrect memory responses we

demonstrated reliably stronger desynchronization for correct as compared to incorrect

retrieval. Thus, we expand previous research further by providing evidence that this gradual

increase in alpha/beta desynchronization contributes to memory performance and predicts

memory success both during item recognition and during discrimination of highly similar

lures. In particular the difference in desynchronization between correct and incorrect lure

responses provides further support for the view that alpha/beta desynchronization tracks the

level of detail of retrieved memories (Hanslmayr et al., 2016), rather than the disinhibition of

task relevant networks as posed by competing models of the functional role of alpha/beta

oscillations (Jensen & Mazaheri, 2010; Klimesch et al., 2007). Given that incorrect trials were

defined as "old" responses to lures, this contrast builds on the fine difference between the

retrieval of the target memory without sufficient detail to detect the perceptual difference of

the mirrored image, and the retrieval of a fine-grained memory representation. Thus, together

with previous observations, our findings provide compelling evidence that alpha/beta

desynchronization supports the gradual accumulation of mnemonic evidence and supports

the successful recovery of highly detailed memories.

The observed effect of alpha/beta desynchronization during retrieval also parallels

findings from earlier event-related potential (ERP) studies that aimed to dissociate memory

processes associated with successful and unsuccessful lure discrimination. In particular, a

left-parietal old/new effect (late positive potential: LPP) has been established as a neural

correlate of successful recollection (see Rugg & Curran, 2007 for review). Some studies

observed the LPP only for true memories (e.g., Curran, 2000; Poch et al.,2019), suggesting

that lure false alarms are driven by familiarity. However, others have demonstrated that the

LPP is observable during lure discrimination independent of memory success (e.g., Curran &

Cleary, 2003; Morcom, 2015), suggesting that lure discrimination requires the recovery of

mnemonic detail and evokes recollection. While we cannot clearly dissociate the relative

contribution of familiarity and recollection in the present paradigm, the onset of the present

effect falls within a similar time range (500–800 ms) as the well-established ERP marker of

recollection and demonstrates a maximum over posterior channels, supporting our

interpretation of alpha/beta desynchronization during retrieval as an index of the specificity of

the retrieved content.

However, it is important to note that with the current design we cannot dissociate

processes related to the recovery of mnemonic content, i.e. retrieval itself, and processes

operating on retrieved content, such as cue specification and monitoring of retrieved content

(for reviews on the involvement of the PFC in episodic memory see e.g. Simon & Spiers,

2003; Eichenbaum, 2017). Thus, it is possible that the effect observed here involves a

mixture of these processes unfolding over time and jointly contributing to the accumulation of

mnemonic evidence.

Previous work has highlighted the crucial role of the hippocampus for successful lure

discrimination. The hippocampus is proposed to lay down non-overlapping memories during

learning (cf. Stark et al., 2019, for review) and to prompt the recovery of the cortical memory

traces at retrieval (Norman & O'Reilly, 2003; Rissman & Wagner, 2012). Accordingly, a

recent intracranial EEG study (Griffiths et al., 2019) demonstrated that cortical alpha/beta

desynchronization (in the anterior temporal lobe) was preceded by hippocampal gamma

increases by a few hundred milliseconds during retrieval, suggesting a directionality in the

flow of information from the hippocampus to the cortex. Furthermore, it is important to note

that although the effect reported here were prominent in the alpha/beta band, activity in other

frequency bands, especially in the theta frequency range (~3–7 Hz), have been shown to

play a crucial role in memory processes. For instance, theta power increases have been

associated with recollection (Gruber et al., 2008) and with hippocampal connectivity with

cortical regions during retrieval (Herweg et al., 2016; for reviews, see Herweg et al., 2020;

Nyhus & Curran, 2010; Staresina & Wimber, 2019). Thus, the recovery of detailed mnemonic

content may rely on the interactive contribution of discrete oscillatory mechanisms,

orchestrating the communication of information through functionally distinct parts of the

episodic memory network. It seems that alpha/beta desynchronization supports the efficiency

by which mnemonic information, as initiated by the hippocampus, is conveyed throughout the

network and represented in 'associative hubs' such as the PPC (Rugg & King, 2017; Rugg &

Vilberg, 2013), thereby ultimately supporting successful item recognition and lure

discrimination.

4.2. Reduced alpha/beta desynchronization in older adults may indicate a loss in memory

specificity

The strength of the observed alpha/beta desynchronization effect across retrieval

demands (i.e., the slope) differed between age groups (see Fig. 3.E), with a smaller effect in

older adults. This age difference in the modulation of alpha/beta power was accompanied by

differences in lure discrimination abilities between younger and older adults (see Fig. 2),

pointing to the potential contribution of age-related changes in alpha/beta desynchronization

to episodic memory decline. Our findings are in line with converging evidence of detrimental

effects of aging on the ability to retrieve episodic details (Bowman et al., 2019; Stark et al.,

2019). We add to this evidence by showing that older adults' difficulties to discriminate

between highly overlapping mnemonic information may relate to less efficient

desynchronization in the alpha/beta frequency bands, presumably reflecting less stable

recovery of detailed memories. We further show that, in contrast to younger adults,

alpha/beta desynchronization in older adults distinguished between correct and incorrect

memories only when reliance on highly detailed memories, i.e., recollection, was required

(lure items), but not when a correct response could rely on familiarity alone, i.e., in the case

of old items (see Fig. 4). When older adults successfully identified a similar lure, alpha/beta

desynchronization was stronger as compared to when they incorrectly endorsed a similar

lure as old (and this effect was comparable across age groups). However, no differences in

alpha/beta desynchronization were observed between correct and incorrect responses to old

target items. Thus, we speculate that less efficient alpha/beta desynchronization in the aging

brain indicates a reduction in information flow (Hanslmayr et al., 2012) which promotes

impairments in the retrieval of detailed memories. As a consequence, older adults are more

likely to rely on gist information and familiarity-based processing, making them more prone to

incorrectly endorse a similar lure as an old item.

In the present study, we did not disentangle the degree to which age differences

occurring at retrieval are a consequence of age differences during encoding. Recent findings

suggest that age differences in episodic memory could also be attributed to encoding-related

neural activity modulated by differences in the integrity of functionally relevant brain regions

(Sander et al., 2020), and that memory quality (defined based on encoding success)

influences retrieval-related processes differently in younger and older adults (Fandakova et

al., 2018). Thus, it is possible that the age differences observed here are downstream effects

of impoverished encoding rendering constructed memories less detailed. Investigations of

the relation between alpha/beta desynchronization and the quality of represented mnemonic

content across encoding and retrieval (see Griffiths et al., 2019 for an approach in this

direction in younger adults) would be a valuable next step for future studies. While in the

current study, we can only infer the level of mnemonic detail available to a person from the

probability of correct responses to targets and lures, we propose that future studies could

map within-person variations of single-trial alpha/beta power in younger and older adults

directly to the representational properties of neural patterns of stimulus-related activity, for

example, their distinctiveness (Koen & Rugg, 2019).

Finally, it is important to note that not only differences in retrieval, but also post-

retrieval processes may have contributed to the observed age differences. The need to

recruit monitoring processes increase with retrieval difficulty (e.g. Dobbins et al., 2002), and

older adults show reductions in the recruitment of these processes, especially when

evaluating information highly similar to previously learned material (e.g. Gutchess et al.,

2007; Pidgeon & Morcom, 2014; Trelle et al., 2019). As in the present design, the lures are

perceptually very similar to the target items, they potentially cause interference and require

additional control processes to make a judgment. Thus, we cannot rule out the possibility that

the age differences reported here result from the decline of a combination of processes that

are necessary for successful lure discrimination, e.g. recovery of a highly detailed memory

representation as well as monitoring of the retrieved content (e.g. Trelle et al., 2017; see

Devitt & Schacter, 2016 for review).

To summarize, we showed that alpha/beta desynchronization increases across

retrieval situations in which correct memory responses require different levels of detail. Age-

related impairments in the retrieval of highly detailed memories were accompanied by

reduced modulations of alpha/beta desynchronization. We further demonstrated that the

level of alpha/beta desynchronization was predictive of memory success: In young adults,

trials with correct and incorrect memory judgements differed reliably for both old targets and

lures. However, in older adults, these effects of memory accuracy were only observed for

lures, thus, when memory success depends on the retrieval of precise details. Taken

together, our results suggest that alpha/beta desynchronization tracks the specificity of

retrieved memories. Importantly, our findings also provide support for the notion that age

differences in oscillatory mechanisms of retrieval contribute to age-related declines in

episodic memory.

Acknowledgements

This research was conducted within the "Lifespan Age Differences in Memory

Representations" (LIME) project at the Center for Lifespan Psychology, Max Planck Institute

for Human Development, Berlin, Germany. MCS was supported by the Minerva program of the

Max Planck Society.

AEK is a doctoral fellow of the International Max Planck Research School on the Life Course

(LIFE). We thank Gabriele Faust (LIME) and all student assistants and interns who helped with

participant recruitment and data collection, project members for helpful feedback on the

analysis, and all study participants for their time. We thank Julia Delius for editorial assistance.

References

Bastin, C., Diana, R. A, Simon, J., Collette, F., Yonelinas, A. P., & Salmon, E. (2014).

Associative memory in aging: The effect of unitization on source memory. *Psychology*

and Aging, *28*(1), 275–283. https://doi.org/10.1037/a0031566

Bell, A. J., Sejnowski, T. J., 1995. An information-maximization approach to blindseparation

and blind deconvolution. *Neural Computation*, *7*(6), 1129–1159.

https://doi.org/10.1162/neco.1995.7.6.1129

Bowman, C. R., Chamberlain, J. D., & Dennis, N. A. (2019). Sensory representations

supporting memory specificity: Age effects on behavioral and neural discriminability.

Journal of Neuroscience, *39*(12), 2265–2275.

https://doi.org/10.1523/JNEUROSCI.2022-18.2019

Brady, T. F., Konkle, T., Alvarez, G. A., & Oliva, A. (2008). Visual long-term memory has a

massive storage capacity for object details. *Proceedings of the National Academy of*

Sciences of the United States of America, *105*(38), 14325–14329.

https://doi.org/10.1073/pnas.0803390105

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897x00357

Buzsáki, G., Anastassiou, C. A., & Koch, C. (2012). The origin of extracellular fields and

currents — EEG, ECoG, LFP and spikes. *Nature Reviews Neuroscience*, *13*, 407–420.

https://doi.org/10.1038/nrn3241

Chouinard, P. A., Morrissey, B. F., Köhler, S., & Goodale, M. A. (2008). Repetition

suppression in occipital–temporal visual areas is modulated by physical rather than

semantic features of objects. *NeuroImage*, *41*(1), 130–144.

https://doi.org/10.1016/j.neuroimage.2008.02.011

Craik, F. I. M. (2020). Remembering: An activity of mind and brain. *Annual Review of*

Psychology, *71*, 1–24.

Curran, T. (2000). Brain potentials of recollection and familiarity. *Memory & Cognition*, *28*(6),

923–938. https://doi.org/10.3758/bf03209340

Curran, Tim, & Cleary, A. M. (2003). Using ERPs to dissociate recollection from familiarity in picture recognition. *Cognitive Brain Research*, *15*(2), 191–205. https://doi.org/10.1016/S0926-6410(02)00192-1

Devitt, A. L., & Schacter, D. L. (2016). False memories with age: Neural and congitive underpinnings. *Neuropsychologia, 91,* 346–359. https://doi.org/10.1016/j.neuropsychologia.2016.08.030

Dewar, M., Alber, J., Butler, C., Cowan, N., & Della Sala, S. (2012). Brief wakeful resting boosts new memories over the long term. *Psychological Science*, *23*(9), 955–960. https://doi.org/10.1177/0956797612441220

Dobbins, I. G., Foley, H., Schacter, D. L., & Wagner, A. D. (2002). Executive control during episodic retrieval: Multiple prefrontal processes subserve source memory. *Neuron, 35,* 989–996. 10.1016/s0896-6273(02)00858-9

Düzel, E., Penny, W. D., & Burgess, N. (2010). Brain oscillations and memory. *Current Opinion in Neurobiology, 20*(2)*,* 143–149. https://doi.org/10.1016/j.conb.2010.01.004

Eichenbaum, H. (2017). Memory: Organization and Control. *Annual Review of Psychology, 68,* 19–45. 10.1146/annurev-psych-010416-044131

Fandakova, Y., Sander, M. C., Grandy, T. H., Cabeza, R., Werkle-Bergner, M., & Shing, Y. L. (2018). Age differences in false memory: The importance of retrieval monitoring processes and their modulation by memory quality. *Psychology And Aging*, *33*(1), 119–133. https:// doi.org/10.1037/pag0000212

Fandakova, Y., Sander, M. C., Werkle-Bergner, M., & Shing, Y. L. (2014). Age differences in short-term memory binding are related to working memory performance across the lifespan. *Psychology and Aging*, *29*(1), 140–149. https://doi.org/10.1037/a0035347

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, *12(3)*, 189–198. https://doi.org/10.1016/0022-3956(75)90026-6

Folville, A., Bahri, M. A., Delhaye, E., Salmon, E., D'Argembeau, A., & Bastin, C. (2020).

Age-related differences in the neural correlates of vivid remembering. *NeuroImage*, *206*,

Article 116336. https://doi.org/10.1016/j.neuroimage.2019.116336

Greene, N. R., & Naveh-Benjamin, M. (2020). A specificity principle of memory: Evidence

from aging and associative memory. *Psychological Science*, *31*(3), 316–331.

https://doi.org/10.1177/0956797620901760

Griffiths, B. J., Mayhew, S. D., Mullinger, K. J., Charest, I., Wimber, M., & Hanslmayr, S.

(2019). Alpha/beta power decreases track the fidelity of stimulus-specific information.

eLife, 8, Article e49562. https://doi.org/10.7554/eLife.49562

Griffiths, B. J., Parish, G., Roux, F., Michelmann, S., & Van der Plas, M., Kolibius, L. D.,

Chelvarajah, R., Rollings, D. T., Sawlani, V., Hamer, H., Gollwitzer, S., Kreiselmeyer,

G., Staresina, B., Wimber, M., & Hanslmayr, S. (2019). Directional coupling of slow and

fast hippocampal gamma with neocortical alpha/beta oscillations in human episodic

memory. *Proceedings of the National Academy of Sciences of the Untied States of*

America, 116(43), 21834–21842. https://doi.org/10.1073/pnas.1914180116

Gruber, T., Tsivilis, D., Giabbiconi, C-M., & Müller, M. M. (2008). Induced

electroencephalogram oscillations during source memors: familiarity is reflected in the

gamma band and recollection in the theta band. *Journal of Cognitive Nueroscience,*

20(6), 1043–1053. https://doi.org/10.1162/jocn.2008.20068

Gutchess, A. H., Hebrank, A., Sutton, B. P., Leshikar, E., Chee, M. W. L., Tan, J. C., . . .

Park, D. C. (2007). Contextual interference in recognition memory with age.

NeuroImage, 35, 1338–1347. http://dx.doi.org/10.1016/j.neuroimage.2007.01.043

Hanslmayr, S., Volberg, G., Wimber, M., Raabe, M., Greenlee, M. W., & Bäuml, K. H. T.

(2011). The relationship between brain oscillations and BOLD signal during memory

formation: A combined EEG-fMRI study. *Journal of Neuroscience*, *31*(44), 15674–

15680. https://doi.org/10.1523/JNEUROSCI.3140-11.2011

Hanslmayr, S., Spitzer, B., & Bäuml, K. H. T. (2009). Brain oscillations dissociate between

semantic and nonsemantic encoding of episodic memories. *Cerebral Cortex*, *19*(7),

1631–1640. https://doi.org/10.1093/cercor/bhn197

Hanslmayr, S., Staresina, B. P., & Bowman, H. (2016). Oscillations and episodic memory:

Addressing the synchronization/desynchronization conundrum. *Trends in*

Neurosciences, *39*(1), 16–25. https://doi.org/10.1016/j.tins.2015.11.004

Hanslmayr, S., & Staudigl, T. (2014). How brain oscillations form memories: A processing

based perspective on oscillatory subsequent memory effects. *NeuroImage*, *85*, 648–

655. http://dx.doi.org/10.1016/j.neuroimage.2013.05.121

Hanslmayr, S., Staudigl, T., & Fellner, M.C. (2012). Oscillatory power decreases and long-

term memory: the information via desynchronization hypothesis. *Frontiers in Human*

Neuroscience, *6*, Article 74. https://doi.org/10.3389/fnhum.2012.00074

Hassler, U., Trujillo Berreto, N., & Gruber, T. (2011). Induced gamma band responses in

human EEG after the control of miniature saccadic artifacts. *NeuroImage*, *57*(4), 1411–

1421. https://doi.org/10.1016/j.neuroimage.2011.05.062

Herweg, N. A., Solomon, E. A., & Kahana, M. J. (2020). Theta oscillations in human memory.

Trends in Cognitive Sciences, *24*(3), 208–227. https://doi.org/10.1016/j.tics.2019.12.006

Herweg, N. A., Apitz, T., Leicht, G., Mulert, C., Fuentemilla, L., & Bunzeck, N. (2016). Theta

– Alpha oscillations bind the hippocampus, prefrontal cortex, and striatum during

recollection: evidence from simultaneous EEG-fMRI. *The Journal of Neuroscience,*

36(12), 3579–3587. 10.1523/JNEUROSCI.3629-15.2016

Jacoby, L. L. (1998). Invariance in automatic influences of memory: Toward a user's guide

for the process-dissociation procedure. *Journal of Experimental Psychology: Learning,*

Memory, and Cognition, *24*(1), 3–26. https://doi.org/10.1037/0278-7393.24.1.3

Jensen, O., Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity:

gating by inhibition. *Frontiers in Human Neuroscience, 4*, 186.

doi:10.3389/fnhum.2010.00186

Kensinger, E. A., & Schacter, D. L. (1999). When true memories suppress false memories:

Effects of ageing. *Cognitive Neuropsychology*, *16*(3–5), 399–415.

https://doi.org/10.1080/026432999380852

Khader, P. H., & Rösler, F. (2011). EEG power changes reflect distinct mechanisms during

long-term memory retrieval. *Psychophysiology*, *48*(3), 362–369. https://doi.org/10.1111/j.1469-8986.2010.01063.x

Kim, H. (2011). Neural activity that predicts subsequent memory and forgetting: A meta-analysis of 74 fMRI studies. *NeuroImage*, *54*(3), 2446–2461. https://doi.org/10.1016/j.neuroimage.2010.09.045

Kim, J. G., Biederman, I., Lescroart, M. D., & Hayworth, K. J. (2009). Adaptation to objects in the lateral occipital complex (LOC): Shape or semantics□? *Vision Research*, *49*(18), 2297–2305. https://doi.org/10.1016/j.visres.2009.06.020

Klimesch, W., Sauseng, P., Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews, 53*, 63-88. https://doi.org/10.1016/j.brainresrev.2006.06.003

Kobelt, M., Sommer, V. R., Keresztes, A., Werkle-Bergner, M., & Sander, M. C. (2020). Tracking age differences in neural distinctiveness across representational levels. *bioRxiv* preprint. https://doi.org/10.1101/2020.07.06.187187

Koen, J. D., Hauck, N., & Rugg, M. D. (2019). The relationship between age, neural differentiation, and memory performance. *Journal of Neuroscience*, *39*(1), 149–162. https://doi.org/10.1523/JNEUROSCI.1498-18.2018

Koen, J. D., & Rugg, M. D. (2019). Neural dedifferentiation in the aging brain. *Trends in Cognitive Sciences*, *23*(7), 547–559. https://doi.org/10.1016/j.tics.2019.04.012

Koen, J. D., & Yonelinas, A. P. (2014). The effects of healthy aging, amnestic mild cognitive impairment, and Alzheimer's disease on recollection and familiarity: A meta-analytic review. *Neuropsychology Review, 24*(3), 332–354. https://doi.org/10.1007/s11065-014-9266-5

Koutstaal, W., & Schacter, D. L. (1997). Gist-based false recognition of pictures in older and younger adults. *Journal of Memory and Language*, *37*(4)*,* 555–583. https://doi.org/10.1006/jmla.1997.2529

Kray, J. (2000). *Adult age differences in task switching: Components, generalizability, and modifiability*. Pabst Science Publishers.

Luo, L., & Craik, F. I. M. (2009). Age differences in recollection: Specificity effects at retrieval. *Journal of Memory and Language*, *60*(4), 421–436. https://doi.org/10.1016/j.jml.2009.01.005

Mandler, G. (1980). Recognizing: The judgment of previous occurrence. *Psychological Review*, *87*(3), 252–271. https://doi.org/10.1037//0033-295X.87.3.252

Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Markram, H., Lübke, J., Frotscher, M., & Sakmann, B. (1997). Regulation of synaptic efficacy by coincidence of postsynaptic APs and EPSPs. *Science*, *275*(5297), 213–216. https://doi.org/10.1126/science.275.5297.213

Martín-Buro, M. C., Wimber, M., Henson, R. N., & Staresina, B. P. (2020). Alpha rhythms reveal when and where item and associative memories are retrieved. *Journal of Neuroscience, 40*(12), 2510–2518. https://doi.org/10.1523/jneurosci.1982-19.2020

Mok, R. M., Myers, N. E., Wallis, G., & Nobre, A. C. (2016). Behavioral and neural markers of flexible attention over working memory in aging. *Cerebral Cortex*, *26*(4), 1831–1842. https://doi.org/10.1093/cercor/bhw011

Morcom, A. M. (2015). Resisting false recognition : An ERP study of lure discrimination. *Brain Research*, *1624*, 336–348. https://doi.org/10.1016/j.brainres.2015.07.049

Motley, S. E., & Kirwan, C. B. (2012). A parametric investigation of pattern separation processes in the medial temporal lobe. *Journal of Neuroscience*, *32*(38), 13076–13084. https://doi.org/10.1523/JNEUROSCI.5920-11.2012

Naveh-Benjamin, M., Hussain, Z., Guez, J., & Bar-On, M. (2003). Adult age differences in episodic memory: further support for an associative-deficit hypothesis. *Journal of Experimental Psychology. Learning, Memory, and Cognition*, *29*(5), 826–837. https://doi.org/10.1037/0278-7393.29.5.826

Ngo, C. T., Newcombe, N. S., & Olson, I. R. (2018). The ontogeny of relational memory and pattern separation. *Developmental Science*, *21*(2), Article e12556.

https://doi.org/10.1111/desc.12556

Nolan, H., Whelan, R., Reilly, R.B., 2010. FASTER: fully automated statisticalthresholding for EEG artifact rejection. *Journal of Neuroscience Methods, 192*(1), 152–162. https://doi.org/10.1016/j.jneumeth.2010.07.015

Norman, K. A., & O'Reilly, R. C. (2003). Modeling hippocampal and neocortical contributions to recognition memory: a complementary-learning-systems approach. *Psychological Review, 110*(4), 611–646. https://doi.org/10.1037/0033-295X.110.4.611

Nyberg, L., Lövdén, M., Riklund, K., Lindenberger, U., & Bäckman, L. (2012). Memory aging and brain maintenance. *Trends in Cognitive Sciences, 16*(5), 292–305. https://doi.org/10.1016/j.tics.2012.04.005

Nyhus, E., & Curran, T. (2010). Functional role of gamma and theta oscillations in episodic memory. *Neuroscience and Biobehavioral Reviews, 34*(7), 1023–1035. https://doi.org/10.1016/j.neubiorev.2009.12.014

O'Bryant, S. E., Humphreys, J. D., Smith, G. E., Ivnik, R. J., Graff-Radford, N. R., Petersen, R. C., & Lucas, J. A. (2008). Detecting dementia with the mini-mental state examination in highly educated individuals. *Archives of Neurology, 65*(7), 963-967. doi:10.1001/archneur.65.7.963

Old, S. R., & Naveh-Benjamin, M. (2008). Differential effects of age on item and associative measures of memory: a meta-analysis. *Psychology and Aging, 23*(1), 104–118. https://doi.org/10.1037/0882-7974.23.1.104

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience, 2011,* Article 156869. https://doi.org/10.1155/2011/156869

Parish, G., Hanslmayr, S., & Bowman, H. (2018). The sync/desync model: How a synchronized hippocampus and a desynchronized neocortex code memories. *Journal of Neuroscience, 38*(14), 3428–3440. https://doi.org/10.1523/JNEUROSCI.2561-17.2018

Park, D. C., Polk, T. A., Park, R., Minear, M., Savage, A., & Smith, M. R. (2004). Aging

reduces neural specialization in ventral visual cortex. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(35), 13091–13095. https://doi.org/10.1073/pnas.0405148101

Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F., 1989. Spherical splines for scalp potential and current density mapping. *Electroencephalography and Clinical Neurophysiology, 72*(2), 184–187. https://doi.org/10.1016/0013-4694(89)90180-6

Pidgeon, L. M., & Morcom, A. M. (2016). Cortical pattern separation and item-specific memory encoding. *Neuropsychologia*, *85*, 256–271. https://doi.org/10.1016/j.neuropsychologia.2016.03.026

Pidgeon, L.M., & Morcom, A.M. (2014). Age-related increases in false recognition: The role of perceptual and conceptual similarity. *Frontiers in Aging Neuroscience, 6*, 1–17.

Poch, C., Prieto, A., Hinojosa, J. A., & Campo, P. (2019). The impact of increasing similar interfering experiences on mnemonic discrimination: Electrophysiological evidence. *Cognitive Neuroscience*, *10*(3), 129–138. https://doi.org/10.1080/17588928.2019.1571484

Rissman, J., & Wagner, A. D. (2012). Distributed representations in memory: Insights from functional brain imaging. *Annual Review of Psychology*, *63*, 101–128. https://doi.org/10.1146/annurev-psych-120710-100344

Rugg, M. D., & Curran, T. (2007). Event-related potentials and recognition memory. *Trends in Cognitive Sciences*, *11*(6), 251–257. https://doi.org/10.1016/j.tics.2007.04.004

Rugg, M. D., & King, D. R. (2018). Ventral lateral parietal cortex and episodc memory retrieval. *Cortex, 107,* 238–250. https://doi.org/10.1016/j.cortex.2017.07.012

Rugg, M. D., & Vilberg, K. L. (2013). Brain networks underlying episodic memory retrieval. *Current Opinion in Neurobiology*, *23*(2), 255–260. https://doi.org/10.1016/j.conb.2012.11.005

Sander, M. C., Fandakova, Y., Grandy, T. H., Shing, Y. L., & Werkle-Bergner, M. (2020). Oscillatory mechanisms of successful memory formation in younger and older adults are related to structural integrity. *Cerebral Cortex.* Advance online publication.

https://doi.org/10.1093/cercor/bhz339

Sander, M. C., Werkle-bergner, M., & Lindenberger, U. (2011). Binding and strategic selection in working memory: A lifespan dissociation. *Psychology and Aging*, *26*(3), 612–624. https://doi.org/10.1037/a0023055

Simons, S. J., & Spiers, H. J. (). Prefrontal and medial temporal lobe interactions in long-term memory. *Nature Reviews Neuroscience, 4*(8), 637–48. 10.1038/nrn1178

Smulders, F. T. Y., Oever, S. ten, Donkers, F. C. L., Quaedflieg, C. W. E. M., & Ven, V. Van De. (2018). Single-trial log transformation is optimal in frequency analysis of resting EEG alpha. *European Journal of Neuroscience*, *48*(7), 2585–2598. https://doi.org/10.1111/ejn.13854

Snodgrass, J. G., & Corwin, J. (1988). Pragmatics of measuring recognition memory: Applications to dementia and amnesia. *Journal of Experimental Psychology. General*, *117*(1), 34–50. https://doi.org/10.1037/0096-3445.117.1.34

Sommer, V. R., Fandakova, Y., Grandy, T. H., Shing, Y. L., Werkle-bergner, M., & Sander, M. C. (2020). Neural pattern similarity differentially relates to memory performance in younger and older adults. *Journal of Neuroscience, 39*(41), 8089–8099. https://doi.org/10.1523/JNEUROSCI.0197-19.2019.

St-Laurent, M., Abdi, H., & Buchsbaum, B. R. (2015). Distributed patterns of reactivation predict vividness of recollection. *Journal of Cognitive Neuroscience*, *27*(10), 2000–2018. https://doi.org/10.1162/jocn_a_00839

Staresina, B. P., Michelmann, S., Bonnefond, M., Jensen, O., Axmacher, N., & Fell, J. (2016). Hippocampal pattern completion is linked to gamma power increases and alpha power decreases during recollection. e*Life*, *5,* Article e17397. https://doi.org/10.7554/eLife.17397.001

Staresina, B. P., Wimber, M. (2019). A Neural Chronometry of Memory Recall. *Trends in Cognitive Sciences, 23*(12), 1071–1085. https://doi.org/10.1016/j.tics.2019.09.011

Stark, S. M., Kirwan, C. B., & Stark, C. E. L. (2019). Mnemonic similarity task: A tool for assessing hippocampal integrity. *Trends in Cognitive Sciences, 23*(11), 938–951.

https://doi.org/10.1016/j.tics.2019.08.003

Stark, S. M., & Stark, C. E. L. (2017). Age-related deficits in the mnemonic similarity task for objects and scenes. *Behavioural Brain Research*, *333*, 109–117. https://doi.org/10.1016/j.bbr.2017.06.049

Stark, S. M., Stevenson, R., Wu, C., Rutledge, S., & Stark, C. E. L. (2015). Stability of age-related deficits in the mnemonic similarity task across task variations. *Behavioral Neuroscience*, *129*(3), 257–268. https://doi.org/10.1037/bne0000055

Strunk, J., & Duarte, A. (2019). Prestimulus and poststimulus oscillatory activity predicts successful episodic encoding for both young and older adults. *Neurobiology of Aging*, *77*, 1–12. https://doi.org/10.1016/j.neurobiolaging.2019.01.005

Strunk, J., James, T., Arndt, J., & Duarte, A. (2017). Age-related changes in neural oscillations supporting context memory retrieval. *Cortex*, *91*, 40–55. https://doi.org/10.1016/j.cortex.2017.01.020

Toner, C. K., Pirogovsky, E., Kirwan, C. B., & Gilbert, P. E. (2009). Visual object pattern separation deficits in nondemented older adults. *Learning & Memory*, *16*, 338–342. https://doi.org/10.1101/lm.1315109

Trelle, A. N., Henson, R. N. A., Green, D. A. E., Simons, J. S., 2017. Declines in representational quality and strategic retrieval processes contribute to age-related increases in false recognition. *Journal of Experimental Psychology: Learning, Memory, and Cognition, 43*(12), 1883–1897. 10.1037/xlm0000412

Trelle, A. N., Henson, R. N., Simons J. S. (2019). Neural evidence for age-related differences in representational quality and strategic retrieval processes. *Neurobiology of Aging, 84*, 50–60. https://doi.org/10.1016/j.neurobiolaging.2019.07.012

Tulving, E. (1985). Memory and consciousness. *Canadian Psychology/Psychologie canadienne*, *26*(1), 2–12. https://doi.org/10.1037/h0080017

Varma, S., Takashima, A., Krewinkel, S., Kooten, M. Van, Fu, L., Medendorp, P. W., Kessels, R. P. C., & Daselaar, S. M. (2017). Non-interfering effects of active post-encoding tasks on episodic memory consolidation in humans. *Frontiers in Behavioral*

Neuroscience, *11,* (Article 54). https://doi.org/10.3389/fnbeh.2017.00054

Waldhauser, G. T., Braun, V., & Hanslmayr, S. (2016). Episodic memory retrieval functionally

relies on very rapid reactivation of sensory information. *Journal of Neuroscience*, *36*(1),

251–260. https://doi.org/10.1523/JNEUROSCI.2101-15.2016

Wang, W., Daselaar, S. M., & Cabeza, R. (2017). Episodic memory decline and healthy

aging. In J. H. Byrne (Ed.), *Learning and memory: A comprehensive reference* (2nd ed.)

(pp. 475–497). Elsevier. https://doi.org/10.1016/B978-0-12-809324-5.21093-6

Wechsler, D. (1955). *Wechsler Adult Intelligence Scale manual.* New York: Psychological

Corporation.

Werkle-Bergner, M., Müller, V., Li, S.-C., & Lindenberger, U. (2006). Cortical EEG correlates

of successful memory encoding: Implications for lifespan comparisons. *Neuroscience*

and   Biobehavioral   Reviews,   *30*(6),   839–854.

https://doi.org/10.1016/j.neubiorev.2006.06.009

Wing, E. A., Ritchey, M., & Cabeza, R. (2015). Reinstatement of individual past events

revealed by the similarity of distributed activation patterns during encoding and retrieval.

Journal   of   Cognitive   Neuroscience,   *27*(4),   679–691.

https://doi.org/10.1162/jocn_a_00740

Yassa, M. A., & Stark, C. E. L. (2011). Pattern separation in the hippocampus. *Trends in*

Neurosciences, *34*(10), 515–525. https://doi.org/10.1016/j.tins.2011.06.006

Yonelinas, A. P. (2002). The nature of recollection and familiarity: A review of 30 years of

research.   *Journal   of   Memory   and   Language*,   *46*(3),   441–517.

https://doi.org/10.1006/jmla.2002.2864

Zheng, L., Gao, Z., Xiao, X., Ye, Z., Chen, C., & Xue, G. (2018). Reduced fidelity of neural

representation underlies episodic memory decline in normal aging. *Cerebral Cortex*,

28(7), 2283–2296. https://doi.org/10.1093/cercor/bhx130