Regulation of blood cell transdifferentiation by oxygen sensing neurons in *Drosophila*

Sean Corcoran[2], Anjeli Mase[2], Yousuf Hashmi[2], Debra Ouyang[2], Jordan Augsburger[2], Thea Jacobs[2], Katelyn Kukar[2], Katja Brückner[1, 2, 3, 4]

[1]Eli and Edythe Broad Center of Regeneration Medicine and Stem Cell Research

[2]Department of Cell and Tissue Biology

[3]Cardiovascular Research Institute, University of California San Francisco, San Francisco, CA

[4] Corresponding Author:

35 Medical Center Way

San Francisco, CA 94143-066

e-mail: katja.brueckner@ucsf.edu

Highlights

- Functional lineage tracing reveals in vivo transdifferentiation in a *Drosophila* model of hematopoiesis

- Active sensory neurons of the caudal sensory cones promote blood cell transdifferentiation in the *Drosophila* larva

- Environmental oxygen sensing through atypical guanylyl cyclases in sensory cone neurons drives blood cell transdifferentiation

Keywords

Drosophila melanogaster, transdifferentiation, hematopoiesis, microenvironment, hemocyte, macrophage, plasmatocyte, crystal cell, sensory neurons, sensory cones, atypical guanylyl cyclase, oxygen sensing, hypoxia

Summary

Transdifferentiation generates functionally specialized cell types independent of stem or progenitor

cells. Despite the unique nature of the process, it remains poorly understood how transdifferentiation

is regulated in vivo. Here we reveal a mechanism of environmental control of blood cell

transdifferentiation in a *Drosophila melanogaster* model of hematopoiesis. Using functional lineage

tracing, we find in vivo evidence for transdifferentiation from macrophage-like plasmatocytes to

crystal cells that execute melanization. Interestingly, this transdifferentiation is promoted by neuronal

activity of a specific subset of sensory neurons, in the sensory cones at the caudal end of the larva, as

is evidenced through genetic ablation, and manipulation of neuronal activity by *Kir2.1* and *TrpA1*.

Crystal cells develop from plasmatocyte clusters surrounding the sensory cones. Strikingly,

environmental conditions trigger this process: oxygen sensing, through atypical guanylyl cyclases

(Gyc88E, Gyc89Da, Gyc89Db) that are specifically expressed in sensory cone neurons, drives

plasmatocyte-to-crystal cell transdifferentiation, as hypoxia or *Gyc* silencing cause crystal cell

reduction and loss of transdifferentiation. Our findings reveal an unexpected functional and molecular

link of environment-monitoring sensory neurons that govern blood cell transdifferentiation in vivo,

suggesting similar principles in vertebrate systems where environmental sensors and blood cell

populations coincide.

Introduction

A Drosophila model of blood cell transdifferentiation

The phenomenon of transdifferentiation has been noted in a variety of species (Cieslar-Pobuda et al., 2017; Graf, 2011; Reid and Tursun, 2018), yet its in vivo regulation remains poorly understood. In the vertebrate blood cell system, reports of transdifferentiation have for the most part been limited to cell culture systems and experimental manipulations. For example, C/EBP (CCAAT/enhancer-binding protein) transcription factors drive transdifferentiation of vertebrate B cells to macrophages (Xie et al., 2004) (Di Tullio et al., 2011), B lymphoma and leukemia cell lines to macrophages (Rapino et al., 2013), and B cells to Granulocyte-Macrophage Precursors (Cirovic et al., 2017). Similarly, manipulation of key transcription factors such as FLI1 and ERG results in transdifferentiation of erythroblasts to megakaryocytes (Siripin et al., 2015), and deletion of the BAF Chromatin Remodeling Complex Subunit Bcl11b triggers T cell transition to NK cells (Li et al., 2010). Transdifferentiation of lymphoid and myeloid cells has been modeled mathematically (Collombet et al., 2017). However, in vivo, the underlying cellular and molecular mechanisms of blood cell transdifferentiation during development and homeostasis and the role of the environment remain elusive.

To investigate principles of in vivo transdifferentiation in the hematopoietic system, we turned to a model in *Drosophila melanogaster*. *Drosophila* offers proven parallels to the two major lineages that produce myeloid blood cells in vertebrates (Davies et al., 2013; Perdiguero and Geissmann, 2016; Sieweke and Allen, 2013), with its two myeloid lineages of blood cells, or hemocytes (Gold and Brückner, 2014, 2015; Holz et al., 2003), (1) the embryonic lineage of hemocytes that proliferate as differentiated cells in hematopoietic pockets of the larval body wall, and resemble vertebrate tissue macrophages, and (2) the progenitor-based lymph gland lineage (Banerjee et al., 2019). Both *Drosophila* blood cell lineages produce at least three differentiated blood cell types: macrophage-like plasmatocytes, crystal cells that mediate melanization, and lamellocytes, large immune cells specialized for encapsulation (Banerjee et al., 2019; Gold and Brückner, 2014, 2015). During larval stages, embryonic-lineage plasmatocytes show signs of fate changes to other blood cell types: Transdifferentiation to lamellocytes occurs in response to immune challenges (Markus et al., 2009), and transdifferentiation to crystal cells was suggested even under steady state conditions (Leitao and Sucena, 2015). However, it remains unclear what are the anatomical requirements and environmental inputs that regulate plasmatocyte-to-crystal cell transdifferentiation in vivo. Hematopoietic sites of

the *Drosophila* larva contain sensory neuron clusters of the peripheral nervous system (PNS) that
serve as microenvironments for plasmatocyte survival, proliferation and localization (Gold and
Brückner, 2014, 2015; Makhijani et al., 2017; Makhijani et al., 2011; Makhijani and Brückner, 2012).
Considering this, we investigated the role of these specialized hematopoietic pockets in hemocyte
transdifferentiation.

Results

Phagocytic plasmatocytes transdifferentiate to crystal cells

First we examined the formation of crystal cells and their anatomical locations during larval
development. We quantified crystal cells using a traditional way of labeling crystal cells based on
their expression of prophenoloxidases (Corcoran and Brückner, 2020; Rizki and Rizki, 1959),
enzymes responsible for melanization, the main immune function of crystal cells (Bidla et al., 2009;
Dudzic et al., 2015; Lu et al., 2014). Induction of melanization blackens crystal cells (Corcoran and
Brückner, 2020; Rizki and Rizki, 1959) and marks similar cell populations as the crystal cell reporter
lozenge-GAL4 (*lz-GAL4; UAS-GFP*) (SupplFig. 1 A, B). During larval development, crystal cell
numbers increase slowly over the first and second larval instar stages, but expand more rapidly
during the third instar stage (Fig. 1 A), largely mirroring the exponential increase of plasmatocytes
during larval development (Fig 1 B) (Makhijani et al., 2011; Petraki et al., 2015). To visualize the
locations of crystal cells and plasmatocytes, we coexpressed two fluorescent protein reporters (for
crystal cells *BcF2-GFP* (Tokusumi et al., 2009), and for plasmatocytes *HmlΔ-DsRed* (Makhijani et
al., 2011)). Interestingly, crystal cells are strongly enriched in a cluster in the terminal segment of the
Drosophila larva, a region where plasmatocytes are also known to accumulate (Fig. 1 C-C'', D-D')
(Makhijani et al., 2011). Crystal cells are occasionally also found in other hematopoietic pockets, and
in dorsal vessel-associated clusters where floating hemocytes accumulate (Cevik et al., 2019; Petraki
et al., 2015).

Since previous studies suggested plasmatocyte-to-crystal cell transdifferentiation solely based on live
imaging (Leitao and Sucena, 2015), we sought an independent functional lineage tracing approach.
Asking whether crystal cells derive from undifferentiated progenitors or differentiated,
phagocytically active plasmatocytes, we traced cells based on the unique ability of differentiated
plasmatocytes to phagocytically uptake fluorescently labeled beads (Fig. 1 E-E'''). We injected
Drosophila larvae expressing fluorescent reporters for plasmatocytes and crystal cells with blue

fluorescent latex beads. Injected larvae were incubated in a time course, followed by the release of

hemocytes and quantification of the relative fractions of phagocytosis-labeled plasmatocytes and -

crystal cells (Fig. 1 F, G). The fraction of blue bead positive plasmatocytes quickly reached saturation

(~50% at 1h, ~90% at 4h and 22h). In contrast, crystal cells were labeled by blue beads with a

significant time delay (<10% at 1h, ~ 50% at 4h, ~70% at 22h) (Fig. 1 G). Together with previous

reports that suggested crystal cells are not capable of phagocytosis or proliferation (Lanot et al., 2001;

Leitao and Sucena, 2015; Tattikota et al., 2019) this supports a model of plasmatocyte-to-crystal cell

transdifferentiation, in which crystal cells derive from phagocytically active plasmatocyes, rather than

from undifferentiated, phagocytosis-incompetent progenitors.

Sensory neuron activity promotes crystal cell transdifferentiation

Next, we investigated the anatomical locations of crystal cells and plasmatocytes relative to sensory

neurons, using fluorescent reporters and live imaging, including the sensory neuron specific driver

21-7-GAL4, UAS-CD8-GFP (Song et al., 2007) (Fig. 2 A). We found that both plasmatocytes and

crystal cells colocalize with sensory neurons, with one important difference: plasmatocytes are found

in all hematopoietic pockets (Fig. 2 A), while crystal cells are mainly localized in the terminal

hematopoietic pocket of the larva (Fig. 2 B).

Given this colocalization with sensory neurons, we asked whether sensory neuron activity has an

effect on crystal cell generation. To mimic activation of sensory neurons, we exposed larvae to the

acetylcholine receptor agonist carbamoycholine (carbachol). Carbachol exposure over 4 hours

resulted in a moderate increase of crystal cell numbers (Fig. 2 C). More specifically and

complementary to this, we silenced sensory neurons by transient expression of *Kir2.1*, an inward

rectifying K+ channel that causes neuron hyperpolarization (Baines et al., 2001). Interestingly, *Kir2.1*

expression over 22 hours caused a dramatic drop in crystal cell numbers (Fig. 2 D). There was no

significant effect of neuronal silencing on total hemocyte numbers under comparable conditions

(SupplFig. 2 and (Makhijani et al., 2017)). To address whether sensory neuron silencing by Kir2.1

affects plasmatocyte to crystal cell transdifferentiation, we performed phagocytosis lineage tracing.

Indeed, transient neuronal silencing affects the fraction of crystal cells derived from blue bead labeled

plasmatocytes, while the ability of plasmatocytes to phagocytose remains the same (Fig. 2 E, F).

Taken together, our findings suggest that sensory neuron activity promotes transdifferentiation of

plasmatocytes to crystal cells.

Crystal cells colocalize with, and require, sensory neurons of the sensory cones

To gain more insight, we focused on the caudal cluster of crystal cells. Closer inspection of combined

fluorescent reporters for crystal cells and sensory neurons showed that crystal cells colocalize

particularly well with sensory organs of the sensory cones (Fig. 3 A-A'' and C-D''), protruding

structures that are grouped around the posterior spiracles, the terminal tubes of the tracheal system

(Hayashi and Kondo, 2018; Kuhn et al., 1992). Plasmatocytes accumulate in a large cluster around

the sensory cones already in the 2^{nd} instar larva (Fig. 3 B-B''), which seems to foreshadow the pattern

and abundance of crystal cells in the 3^{rd} instar larva (Fig. 3 D-D''). We therefore hypothesized that

many plasmatocytes of the 2^{nd} instar larva may transdifferentiate to crystal cells, as is apparent in the

3^{rd} instar.

Given the intriguing colocalization of crystal cells and sensory neurons and the dependence of blood

cell transdifferentiation on neuronal activity, we investigated the specific requirement of sensory cone

neurons for crystal cell production. A group of genes specifically expressed in sensory cone neurons

are atypical guanylyl cyclases, *Gyc88E*, *Gyc89Da*, and *Gyc89Db* (Vermehren-Schmaedick et al.,

2010). Using the driver *Gyc89Db-GAL4* (Vermehren-Schmaedick et al., 2010) (Fig. 3 E, F), we

ablated sensory cone neurons by expressing the proapoptotic gene *head involution defect (Hid)* in its

non-repressible version (*Hid ^{ala5}*) (Bergmann et al., 2002). Ablation of sensory cone neurons did not

affect larval viability, but it strongly reduced crystal cell numbers (Fig. 3 G). Conversely, ectopic

activation of sensory cone neurons by specific expression and transient induction of the heat-induced

cation channel TrpA1 (Hamada et al., 2008) caused a significant increase in crystal cells (Fig. 3 H).

Taken together, we conclude that crystal cells and their precursor plasmatocytes are strongly enriched

at the sensory cones of the larva. Sensory cone neurons, and their activity, are required for crystal

cells.

Oxygen sensing through atypical guanylyl cyclases drives plasmatocyte-to-crystal cell

transdifferentiation

The atypical guanylyl cyclases Gyc88E, Gyc89Da, and Gyc89Db are cytoplasmic oxygen sensors

that form heterodimers of Gyc88E and either Gyc89Da or Gyc89Db, and Gyc88E homodimers

(Huang et al., 2007; Morton, 2004; Morton et al., 2005; Vermehren et al., 2006). Activated Gyc

complexes generate the second messenger cyclic GMP (cGMP), which in turn activates neurons

(Morton, 2004; Morton et al., 2008). Since we found that sensory cone neurons expressing these

genes are required for crystal cell formation, we decided to test Gyc function itself in relation to

crystal cell transdifferentiation. When the obligatory subunit, Gyc88E, was silenced in the sensory

cone neurons, crystal cell numbers were reduced (Fig. 4 A). Phagocytosis lineage tracing confirmed

that Gyc function is required for transdifferentiation of plasmatocytes to crystal cells, as the fraction

of crystal cells carrying blue beads was significantly reduced in *Gyc88E* RNAi knockdowns, while

phagocytosis by plasmatocytes remained the same (Fig. 4 B). Given the role of Gycs as oxygen

sensors, we next investigated the effect of varying atmospheric oxygen concentrations on crystal cell

formation. Assessing crystal cell numbers per larva as readout, we exposed larvae to 8% or 5%

oxygen for 6 hours. Compared to normoxia (~21% oxygen), 8% hypoxia caused mild reduction in

crystal cells and 5% oxygen caused a significant drop in crystal cells, while total hemocyte numbers

stayed constant (SupplFig. 3 A, B, Fig. 4 C). Following up on the robust results obtained under 5%

oxygen, we determined whether hypoxia affects blood cell transdifferentiation, performing

phagocytosis lineage tracing for 6 hours under hypoxic conditions (5% O_2) and normoxia.

Interestingly, hypoxia phenocopies silencing of Gyc function in sensory cone neurons, resulting in a

significant reduction of blue beads in crystal cells, while phagocytosis levels of plasmatocytes remain

unaffected (Fig. 4 D). We therefore conclude that oxygen sensing through atypical guanylyl cyclases

(Gycs) that are linked to activation of the sensory cone neurons drives plasmatocyte-to-crystal cell

transdifferentiation (Fig. 4 E). This model supports the kinetics of crystal cells following

plasmatocyte expansion, based on the transition of plasmatocytes to crystal cells in particular in the

proximity of sensory cone neurons (SupplFig. 4 A).

Discussion

Our work has identified an unexpected mechanistic link between oxygen sensing and blood cell

transdifferentiation, which is facilitated through a particular set of sensory neurons and intracellular

Gyc oxygen sensors. This new paradigm inspires the search for similar principles of neuronally

controlled blood cell transdifferentiation that responds to environmental conditions in other species

including humans.

Transdifferentiation results in the conversion of one differentiated cell type to another. In some

systems, new differentiated cell types arise after de-differentiation to a transient pluripotent

intermediate (Graf, 2011; Pesaresi et al., 2019; Reid and Tursun, 2018). Our work supports a model

of direct transdifferentiation based on fluorescent reporters and phagocytosis lineage tracing.

Independent approaches also support a model of continuous blood cell transdifferentiation from

plasmatocytes to crystal cells through progressive states, based on single cell RNAseq pseudotime lineage analysis (Tattikota et al., 2019). Transdifferentiation may be the fastest and most efficient way for animals to shape the composition of their blood cell pool according to environmental conditions such as oxygen levels and potentially other inputs.

Gyc intracellular oxygen sensors mediate oxygen detection in sensory neurons (Vermehren et al., 2006), similar to other oxygen sensing mechanisms known in vertebrate neurons and other sensory cells (Caravagna and Seaborn, 2016; Pokorski et al., 2016). In contrast, HIF (hypoxia inducible factor) transcription factors regulate target genes in response to low oxygen conditions in a variety of cell types (Gorr et al., 2006; Majmundar et al., 2010). Hypoxia, through HIF, regulates mammalian hematopoiesis, lymphopoiesis and erythropoiesis (Chabi et al., 2019; Haase, 2013; Imanirad and Dzierzak, 2013). The *Drosophila* HIF1a *sima* (*similar*) plays a role in crystal cell formation in the *Drosophila* lymph gland (Mukherjee et al., 2011), however this effect is independent of HIF1β and hypoxia target genes. Instead, formation of *Drosophila* crystal cells has been linked to Notch signaling at various developmental stages (Duvic et al., 2002; Krzemien et al., 2007; Lebestky et al., 2000; Lebestky et al., 2003; Leitao and Sucena, 2015; Mukherjee et al., 2011). It remains to be determined whether oxygen sensing neurons and/or the signal receiving plasmatocytes are connected to Notch signaling or an independent pathway that governs the fate switch of plasmatocytes to crystal cells.

Linking oxygen sensing to crystal cell formation via Gycs may have several advantages. (1) Oxygen sensing by neurons is more sensitive and immediate, and respond to a variety of environmental conditions. Sensory cone neurons are at all times in contact with the surrounding atmosphere, based on social burying behaviors during feeding, when larvae expose their caudal ends with the sensory cones, also allowing air intake to the tracheal system through the neighboring posterior spiracles (Hayashi and Kondo, 2018; Wu et al., 2003) (SupplFig. 4 B); eventually larvae exit the food source to pupariate (Wu et al., 2003) (SupplFig. 4 B). With their exposed nature, the sensory neurons may also integrate other inputs such as chemical cues or even light from the environment (Stewart et al., 2015; Vermehren-Schmaedick et al., 2011; Xiang et al., 2010) when communicating to their targets. (2) Activation of the sensory cone neurons coordinates blood cell transdifferentiation with other responses to hypoxia, such as a behavioral escape response to hypoxic conditions (Morton, 2011; Vermehren-Schmaedick et al., 2010). In this context, Gycs generate cGMP, which activates cyclic nucleotide gated channels (CNG) (Morton et al., 2008; Vermehren-Schmaedick et al., 2010). CNG

channels mediate influx of calcium ions, resulting in consecutive activation of calmodulin/CaMK

signaling and sensory transduction (Kaupp and Seifert, 2002; Pifferi et al., 2006).

Neuronal regulation of the hematopoietic system and other organs is an important paradigm in

biology, which is starting to come to light in a variety of species (Kumar and Brockes, 2012). In

Drosophila, neuronal regulation of hematopoiesis is well established. Embryonic-lineage

plasmatocytes depend on sensory neurons for their survival, proliferation and localization (Gold and

Brückner, 2014, 2015; Makhijani et al., 2017; Makhijani et al., 2011; Makhijani and Brückner, 2012).

Activin-β, a TGF-β family ligand, is a key signal produced by active sensory neurons that promotes

plasmatocyte proliferation and adhesion (Makhijani et al., 2017). Identification of the signal/s from

active sensory cone neurons that trigger transdifferentiation will be the focus of intense future study.

We postulate secreted factor/s, which could potentially, albeit at reduced efficiency, also act at a

distance, promoting transdifferentiation of smaller numbers of crystal cells in the dorsal vessel

associated hemocyte clusters (Leitao and Sucena, 2015) and segmental hematopoietic pockets. In this

context it is interesting to note that in long term memory formation, Activin expression is induced

downstream of calmodulin/CaMK/CREB signaling in both *Drosophila* and vertebrates (Inokuchi et

al., 1996; Miyashita et al., 2012), suggesting potential parallels of this signaling cassette in neuron-

induced blood cell transdifferentiation.

In vertebrates, functional links of hematopoiesis with sensory neurons or other sensing systems

remain largely unknown, despite some aspects of bone marrow hematopoiesis and inflammatory

responses being regulated by the autonomic nervous system (Hanoun et al., 2015; Pavlov and Tracey,

2012). Oxygen sensing could be an important regulatory factor in recently identified hematopoietic

sites such as the vertebrate lung, which provides a microenvironment for limited blood cell

progenitors and megakaryocytes that are active in platelet production (Lefrancais et al., 2017; Martin

et al., 1983). The lung, like many other organs, also harbors tissue macrophages that proliferate in

local microenvironments and bear evolutionary parallels with *Drosophila* embryonic-lineage

plasmatocytes (Gold and Brückner, 2014, 2015; Perdiguero and Geissmann, 2016). Interestingly, the

lung and airways are rich in vagal afferent nerves that sense chemical and mechanical cues (Chang et

al., 2015; Mazzone and Undem, 2016), and neuroendocrine cells of the lung that sense oxygen early

in life and later provide a microenvironment for airway epithelial cells (Caravagna and Seaborn,

2016; Cutz et al., 2007). Investigation of sensory neurons and other sensors may therefore open a new

chapter in the regulation of vertebrate hematopoiesis, transdifferentiation, and immune cell fate and

function.

Materials and Methods

Drosophila Strains

Drosophila drivers, reporters and related lines used were *HmlΔ-GAL4, UAS-GFP* (Sinenko and Mathey-Prevot, 2004)*;* combination driver *HmlΔGAL4, UAS-GFP; He-GAL4* ((Yang et al., 2015) gift from Dan Hultmark), *lz-GAL4; UAS-GFP* (J. Pollock, Bloomington), *21-7-GAL4* (Makhijani et al., 2011; Song et al., 2007) *Gyc89Db-GAL4 (Morton et al., 2008); BcF6-GFP* (Tokusumi et al., 2009), *BcF6-mCherry* (Tokusumi et al., 2009), *BcF2-GFP* (Tokusumi et al., 2009); *HmlΔ-DsRed* (Makhijani et al., 2011); and *tubGAL80ts* (McGuire et al., 2003). UAS lines used were *UAS-CD8-GFP* (Song et al., 2007); *UAS-Kir2.1* (Baines et al., 2001) (Bloomington); *UAS-TrpA1* (Bloomington); *UAS-Hid ala5* (Bergmann et al., 2002); *UAS-Gyc88E RNAi* (Bloomington). Control lines used were *w1118* (Bloomington) or *yw* (Bloomington). Unless otherwise stated, fly crosses were set and maintained at 25° Celsius. Crosses with the driver *Gyc89Db-GAL4* were performed at 29°C to enhance expression.

Hemocyte Quantification

Total hemocyte quantification was performed essentially as described in (Corcoran and Brückner, 2020; Petraki et al., 2015). All fluorescently-marked hemocytes of single larvae were released into wells marked by a hydrophobic PAP pen (Beckman Coulter) on glass slides filled with 20-30 μL PBS. Cells were allowed to settle for 15-20 min, and were imaged by fluorescence tile scan microscopy on a Leica DMI4000B microscope with Leica DFC350FX camera and 20x objective. Cell numbers in images were analyzed by particle quantification using Fiji/ImageJ (Corcoran and Brückner, 2020; Petraki et al., 2015; Schindelin et al., 2012).

Crystal cell quantification was performed using fluorescent protein reporters or melanization. To quantify crystal cells in live animals, larvae were placed on a slide in a small drop of 10-15 μl PBS with a coverslip on top. Fluorescent crystal cells in each segment were manually counted, rolling the larva by gently moving the coverslip. For phagocytosis lineage tracing, fluorescent crystal cells were quantified ex vivo (see below). To quantify crystal cells based on their ability to melanize (due to their hallmark expression of functional Prophenoloxidase 1 and 2) larvae were placed in 250 μl of PBS in Eppendorf tubes and heated at 65° Celsius for 22 minutes in a heat block, and melanized (black) cells were quantified under a stereoscope by manual counting (Corcoran and Brückner, 2020; Rizki and Rizki, 1959).

Larvae were analyzed at various developmental times after egg laying (AEL) corresponding to the

following size ranges: 1st instar: ~0.5-1.4mm (22-46h AEL; 2nd instar: ~1.5-2.6mm (47-77h AEL); 3rd

instar: ~2.7- >3.5mm (from 78h AEL). For the crosses used, no developmental delays were observed;

therefore selection of specified larval size ranges from 24h embryo collections was used in lieu of

more tightly timed embryo collections.

Manipulation of Neuronal Activity

To mimic sensory neuron stimulation, larvae were exposed to 10mg/ml carbamylcholine (carbachol,

Sigma Aldrich) in fly food for 4 hours allowing direct cuticle contact with carbachol (Makhijani et

al., 2017).

To activate specific neuron populations, the heat-inducible cation channel TrpA1 was ectopically

expressed and transiently heat induced to mimic neuron activation (Hamada et al., 2008).

Sopecifically, *TrpA1* crosses were set at RT and shifted to 29°C for 4h. Larvae were analyzed within

30 minutes following this period.

Sensory neuron silencing was achieved by transiently expressing a transgenic of the inward rectifying

potassium channel Kir2.1 (Baines et al., 2001), under control of a sensory neuron specific GAL4

driver and a temperature-sensitive GAL4 inhibitor GAL80ts (McGuire et al., 2003); genotypes for

neuronal silencing experiments were *21-7-GAL4, UAS-GFP, HmlΔDsRed/UAS-Kir2.1;*

tubGAL80ts/+ with controls *21-7-GAL4, UAS-GFP, HmlΔDsRed/+; tubGAL80ts/+*. F1 from crosses

were raised at 18° Celsius. To temporarily silence sensory neurons, larvae were shifted from 18° to

29° Celsius for 22 hours to destabilize GAL80ts, allowing for expression of Kir2.1, which

hyperpolarizes sensory neuron preventing firing. Larvae were analyzed within 1 hour post-silencing.

Phagocytosis Lineage Tracing

For lineage tracing of crystal cells that derive from actively phagocytic plasmatocytes, larvae with

genotype *HmlΔGAL4, UAS-GFP; BcF6-mCherry* were injected with 69 nl of blue fluorescent

FluoSphere Carboxylate-modified 0.2 um beads (Invitrogen) diluted 1:100 in PBS using a Nanoject

injector (Drummond Scientific). After injection, larvae were placed on food with yeast for 4 hours.

Hemocytes of individual larvae were then released into a well marked on a glass slide by PAP pen

and filled with 20-30 µl PBS; hemocytes were allowed to settle for 15-20 minutes in a wet chamber

(Corcoran and Brückner, 2020; Petraki et al., 2015). Cells were imaged with a Leica DMI4000

microscope with tilescan function. Cell quantification was conducted manually from a representative

central field of each image (containing ~150-300 plasmatocytes). For genotypes that give rise to substantially reduced numbers of crystal cells, crystal cells of the whole tilescan area were analyzed. Transdifferentiation was quantified by comparing ratios of crystal cells positive for blue particles/ total crystal cells (mCherry positive) for various experimental and control conditions. The same quantification for plasmatocytes (GFP positive) with blue beads relative to all plasmatocytes was conducted as an internal control for injection efficiency and phagocytic fitness.

Hypoxia experiments

Hypoxia experiments were conducted using a hypoxic chamber (Biospherix, Inc., Laconia, NY). Oxygen concentrations were set as indicated, supplementing reduced O_2 with N_2. Incubations were performed at room temperature (RT) for 6h. Crystal cell melanization assays were conducted with larvae of the genotype w^{1118}, and phagocytic lineage tracing and total hemocyte experiments used *HmlΔ-GAL4, UAS-GFP; BcF6-mCherry* larvae. For all experiments, 15-25 larvae of the desired age/size and genotype were placed on inactivated yeast paste in cell strainer snap cap tubes, allowing for rapid gas exchange (Corning #352235). Six cell strainer tubes were held by a fitted rack that was inserted into a large cylindrical polystyrene *Drosophila* population cage with steel mesh on one side (custom construction). During transport from the hypoxic chamber to the bench, population cages were sealed in a plastic bag to maintain hypoxic conditions up to the point when larvae were analyzed. Control samples were placed in an identical setup but left at normoxic conditions at comparable temperature (RT) and time (6h). Following hypoxia/normoxia conditions, assays were conducted as described in the respective paragraphs.

Microscopy

Released hemocytes were imaged on a Leica DMI4000B microscope as described above. Live imaging of larvae was done as described previously, immobilizing larvae on an ice-cooled metal block (Makhijani et al. *Development* 2011). Imaging was performed using a Leica M205FA fluorescent stereoscope with DFC300FX color digital camera and Leica LAS Montage module, combining z-stacks into single in-focus images.

Statistical Analysis

For all experiments numbers of larvae per genotype and condition are indicated in the Figure Legends. For each genotype and condition the mean and standard deviation were determined and significance was tested by 2-way ANOVA (Prism). For hemocyte counts over the course of larval

329 development, regression analysis was performed (Excel). For phagocytosis lineage tracing, the mean
330 and standard deviation of percentages of blue bead positive cells were determined and assessed by 2-
331 way ANOVA (Prism). P-value cutoffs for significance were as follows: $* = p < 0.05$, $** = p < 0.01$,
332 and $*** = p < 0.001$. Pools of both male and female larvae were analyzed.

Authors' contributions

335 KB conceived and supervised the study. SC, AM and KB planned the experiments. SC, AM, JA, YH,
336 DO, TJ, KK carried out the experiments. SC, AM, JA, YH, DO, TJ, KK and KB analyzed the data.
337 KB wrote the manuscript with input from all authors.

Competing interests

340 The authors have no competing interests.

Funding

343 This work was supported by grants from the American Cancer Society RSG DDC-122595, American
344 Heart Association 13BGIA13730001, National Science Foundation 1326268, and National Institutes
345 of Health 1R01GM112083, 1R56HL118726 and 1R01GM131094 (to KB).

Acknowledgements

348 We thank D. Morton, R. Schulz, U. Banerjee, A. Bergmann, C. Evans, D. Hultmark, Y.N. Jan, L.
349 Kockel, the Bloomington Stock Center, and TRiP for fly stocks. We are grateful to Khalida Sabeur
350 and Manideep Chavali for help to access and use hypoxia chamber equipment. Thanks to Ava
351 Brückner-Kockel for advice on regression analysis. We thank members of the Brückner lab for
352 discussion and feedback on the manuscript.

References

Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T., and Bate, M. (2001). Altered electrical properties in *Drosophila* neurons developing without synaptic transmission. J Neurosci *21*, 1523-1531.

Banerjee, U., Girard, J.R., Goins, L.M., and Spratford, C.M. (2019). *Drosophila* as a Genetic Model for Hematopoiesis. Genetics *211*, 367-417.

Bergmann, A., Tugentman, M., Shilo, B.Z., and Steller, H. (2002). Regulation of cell number by MAPK-dependent control of apoptosis: a mechanism for trophic survival signaling. Dev Cell *2*, 159-170.

Bidla, G., Hauling, T., Dushay, M.S., and Theopold, U. (2009). Activation of insect phenoloxidase after injury: endogenous versus foreign elicitors. J Innate Immun *1*, 301-308.

Caravagna, C., and Seaborn, T. (2016). Oxygen Sensing in Early Life. Lung *194*, 715-722.

Cevik, D., Acker, M., Michalski, C., and Jacobs, J.R. (2019). Pericardin, a *Drosophila* collagen, facilitates accumulation of hemocytes at the heart. Dev Biol *454*, 52-65.

Chabi, S., Uzan, B., Naguibneva, I., Rucci, J., Fahy, L., Calvo, J., Arcangeli, M.L., Mazurier, F., Pflumio, F., and Haddad, R. (2019). Hypoxia Regulates Lymphoid Development of Human Hematopoietic Progenitors. Cell Rep *29*, 2307-2320 e2306.

Chang, R.B., Strochlic, D.E., Williams, E.K., Umans, B.D., and Liberles, S.D. (2015). Vagal Sensory Neuron Subtypes that Differentially Control Breathing. Cell *161*, 622-633.

Cieslar-Pobuda, A., Knoflach, V., Ringh, M.V., Stark, J., Likus, W., Siemianowicz, K., Ghavami, S., Hudecki, A., Green, J.L., and Los, M.J. (2017). Transdifferentiation and reprogramming: Overview of the processes, their similarities and differences. Biochim Biophys Acta Mol Cell Res *1864*, 1359-1369.

Cirovic, B., Schonheit, J., Kowenz-Leutz, E., Ivanovska, J., Klement, C., Pronina, N., Begay, V., and Leutz, A. (2017). C/EBP-Induced Transdifferentiation Reveals Granulocyte-Macrophage Precursor-like Plasticity of B Cells. Stem Cell Reports *8*, 346-359.

Collombet, S., van Oevelen, C., Sardina Ortega, J.L., Abou-Jaoude, W., Di Stefano, B., Thomas-Chollier, M., Graf, T., and Thieffry, D. (2017). Logical modeling of lymphoid and myeloid cell specification and transdifferentiation. Proc Natl Acad Sci U S A *114*, 5792-5799.

Corcoran, S., and Brückner, K. (2020). Quantification of blood cells in *Drosophila* and other insects. Springer Protocols Handbooks: Immunity in Insects *Sandrelli F, Tettamanti G, editors*, 65-77.

Cutz, E., Yeger, H., and Pan, J. (2007). Pulmonary neuroendocrine cell system in pediatric lung disease-recent advances. Pediatr Dev Pathol *10*, 419-435.

Davies, L.C., Jenkins, S.J., Allen, J.E., and Taylor, P.R. (2013). Tissue-resident macrophages. Nat Immunol *14*, 986-995.

Di Tullio, A., Vu Manh, T.P., Schubert, A., Castellano, G., Mansson, R., and Graf, T. (2011). CCAAT/enhancer binding protein alpha (C/EBP(alpha))-induced transdifferentiation of pre-B cells into macrophages involves no overt retrodifferentiation. Proc Natl Acad Sci U S A *108*, 17016-17021.

Dudzic, J.P., Kondo, S., Ueda, R., Bergman, C.M., and Lemaitre, B. (2015). *Drosophila* innate immunity: regional and functional specialization of prophenoloxidases. BMC Biol *13*, 81.

Duvic, B., Hoffmann, J.A., Meister, M., and Royet, J. (2002). Notch signaling controls lineage specification during *Drosophila* larval hematopoiesis. Curr Biol *12*, 1923-1927.

Gold, K.S., and Brückner, K. (2014). *Drosophila* as a model for the two myeloid blood cell systems in vertebrates. Exp Hematol *42*, 717-727.

Gold, K.S., and Brückner, K. (2015). Macrophages and cellular immunity in *Drosophila melanogaster*. Semin Immunol *27*, 357-368.

Gorr, T.A., Gassmann, M., and Wappner, P. (2006). Sensing and responding to hypoxia via HIF in model invertebrates. J Insect Physiol *52*, 349-364.

Graf, T. (2011). Historical origins of transdifferentiation and reprogramming. Cell Stem Cell *9*, 504-516.

Haase, V.H. (2013). Regulation of erythropoiesis by hypoxia-inducible factors. Blood Rev *27*, 41-53.

Hamada, F.N., Rosenzweig, M., Kang, K., Pulver, S.R., Ghezzi, A., Jegla, T.J., and Garrity, P.A. (2008). An internal thermal sensor controlling temperature preference in *Drosophila*. Nature *454*, 217-220.

Hanoun, M., Maryanovich, M., Arnal-Estape, A., and Frenette, P.S. (2015). Neural regulation of hematopoiesis, inflammation, and cancer. Neuron *86*, 360-373.

Hayashi, S., and Kondo, T. (2018). Development and Function of the *Drosophila* Tracheal System. Genetics *209*, 367-380.

Holz, A., Bossinger, B., Strasser, T., Janning, W., and Klapper, R. (2003). The two origins of hemocytes in *Drosophila*. Development *130*, 4955-4962.

Huang, S.H., Rio, D.C., and Marletta, M.A. (2007). Ligand binding and inhibition of an oxygen-sensitive soluble guanylate cyclase, Gyc-88E, from *Drosophila*. Biochemistry *46*, 15115-15122.

Imanirad, P., and Dzierzak, E. (2013). Hypoxia and HIFs in regulating the development of the hematopoietic system. Blood Cells Mol Dis *51*, 256-263.

Inokuchi, K., Kato, A., Hiraia, K., Hishinuma, F., Inoue, M., and Ozawa, F. (1996). Increase in activin beta A mRNA in rat hippocampus during long-term potentiation. FEBS Lett *382*, 48-52.

Kaupp, U.B., and Seifert, R. (2002). Cyclic nucleotide-gated ion channels. Physiol Rev *82*, 769-824.

Krzemien, J., Dubois, L., Makki, R., Meister, M., Vincent, A., and Crozatier, M. (2007). Control of blood cell homeostasis in *Drosophila* larvae by the posterior signalling centre. Nature *446*, 325-328.

Kuhn, D.T., Sawyer, M., Packert, G., Turenchalk, G., Mack, J.A., Sprey, T.E., Gustavson, E., and Kornberg, T.B. (1992). Development of the D. *melanogaster* caudal segments involves suppression of the ventral regions of A8, A9 and A10. Development *116*, 11-20.

Kumar, A., and Brockes, J.P. (2012). Nerve dependence in tissue, organ, and appendage regeneration. Trends in neurosciences *35*, 691-699.

Lanot, R., Zachary, D., Holder, F., and Meister, M. (2001). Postembryonic hematopoiesis in *Drosophila*. Dev Biol *230*, 243-257.

Lebestky, T., Chang, T., Hartenstein, V., and Banerjee, U. (2000). Specification of *Drosophila* hematopoietic lineage by conserved transcription factors. Science *288*, 146-149.

Lebestky, T., Jung, S.H., and Banerjee, U. (2003). A Serrate-expressing signaling center controls
Drosophila hematopoiesis. Genes Dev *17*, 348-353.

Lefrancais, E., Ortiz-Munoz, G., Caudrillier, A., Mallavia, B., Liu, F., Sayah, D.M., Thornton, E.E.,
Headley, M.B., David, T., Coughlin, S.R*., et al.* (2017). The lung is a site of platelet biogenesis and a
reservoir for haematopoietic progenitors. Nature *544*, 105-109.

Leitao, A.B., and Sucena, E. (2015). *Drosophila* sessile hemocyte clusters are true hematopoietic
tissues that regulate larval blood cell differentiation. Elife *4*.

Li, P., Burke, S., Wang, J., Chen, X., Ortiz, M., Lee, S.C., Lu, D., Campos, L., Goulding, D., Ng,
B.L*., et al.* (2010). Reprogramming of T cells to natural killer-like cells upon Bcl11b deletion.
Science *329*, 85-89.

Lu, A., Zhang, Q., Zhang, J., Yang, B., Wu, K., Xie, W., Luan, Y.X., and Ling, E. (2014). Insect
prophenoloxidase: the view beyond immunity. Front Physiol *5*, 252.

Majmundar, A.J., Wong, W.J., and Simon, M.C. (2010). Hypoxia-inducible factors and the response
to hypoxic stress. Mol Cell *40*, 294-309.

Makhijani, K., Alexander, B., Rao, D., Petraki, S., Herboso, L., Kukar, K., Batool, I., Wachner, S.,
Gold, K.S., Wong, C*., et al.* (2017). Regulation of *Drosophila* hematopoietic sites by Activin-beta
from active sensory neurons. Nat Commun *8*, 15990.

Makhijani, K., Alexander, B., Tanaka, T., Rulifson, E., and Brückner, K. (2011). The peripheral
nervous system supports blood cell homing and survival in the *Drosophila* larva. Development *138*,
5379-5391.

Makhijani, K., and Brückner, K. (2012). Of blood cells and the nervous system: Hematopoiesis in the
Drosophila larva. Fly (Austin) *6*, 254-260.

Markus, R., Laurinyecz, B., Kurucz, E., Honti, V., Bajusz, I., Sipos, B., Somogyi, K., Kronhamn, J.,
Hultmark, D., and Ando, I. (2009). Sessile hemocytes as a hematopoietic compartment in *Drosophila*
melanogaster. Proc Natl Acad Sci U S A *106*, 4805-4809.

Martin, J.F., Slater, D.N., and Trowbridge, E.A. (1983). Abnormal intrapulmonary platelet
production: a possible cause of vascular and lung disease. Lancet *1*, 793-796.

Mazzone, S.B., and Undem, B.J. (2016). Vagal Afferent Innervation of the Airways in Health and
Disease. Physiol Rev *96*, 975-1024.

McGuire, S.E., Le, P.T., Osborn, A.J., Matsumoto, K., and Davis, R.L. (2003). Spatiotemporal rescue
of memory dysfunction in *Drosophila*. Science *302*, 1765-1768.

Miyashita, T., Oda, Y., Horiuchi, J., Yin, J.C., Morimoto, T., and Saitoe, M. (2012). Mg(2+) block of
Drosophila NMDA receptors is required for long-term memory formation and CREB-dependent gene
expression. Neuron *74*, 887-898.

Morton, D.B. (2004). Atypical soluble guanylyl cyclases in *Drosophila* can function as molecular
oxygen sensors. J Biol Chem *279*, 50651-50653.

Morton, D.B. (2011). Behavioral responses to hypoxia and hyperoxia in *Drosophila* larvae:
molecular and neuronal sensors. Fly (Austin) *5*, 119-125.

Morton, D.B., Langlais, K.K., Stewart, J.A., and Vermehren, A. (2005). Comparison of the properties
of the five soluble guanylyl cyclase subunits in *Drosophila melanogaster*. J Insect Sci *5*, 12.

Morton, D.B., Stewart, J.A., Langlais, K.K., Clemens-Grisham, R.A., and Vermehren, A. (2008).
Synaptic transmission in neurons that express the *Drosophila* atypical soluble guanylyl cyclases,
Gyc-89Da and Gyc-89Db, is necessary for the successful completion of larval and adult ecdysis. J
Exp Biol *211*, 1645-1656.

Mukherjee, T., Kim, W.S., Mandal, L., and Banerjee, U. (2011). Interaction between Notch and Hif-
alpha in development and survival of *Drosophila* blood cells. Science *332*, 1210-1213.

Pavlov, V.A., and Tracey, K.J. (2012). The vagus nerve and the inflammatory reflex--linking
immunity and metabolism. Nature reviews. Endocrinology *8*, 743-754.

Perdiguero, E.G., and Geissmann, F. (2016). The development and maintenance of resident
macrophages. Nat Immunol *17*, 2-8.

Pesaresi, M., Sebastian-Perez, R., and Cosma, M.P. (2019). Dedifferentiation, transdifferentiation and
cell fusion: in vivo reprogramming strategies for regenerative medicine. Febs J *286*, 1074-1093.

Petraki, S., Alexander, B., and Brückner, K. (2015). Assaying Blood Cell Populations of the
Drosophila melanogaster Larva. J Vis Exp.

Pifferi, S., Boccaccio, A., and Menini, A. (2006). Cyclic nucleotide-gated ion channels in sensory
transduction. FEBS Lett *580*, 2853-2859.

Pokorski, M., Takeda, K., and Okada, Y. (2016). Oxygen Sensing Mechanisms: A Physiological
Penumbra. Adv Exp Med Biol *952*, 1-8.

Rapino, F., Robles, E.F., Richter-Larrea, J.A., Kallin, E.M., Martinez-Climent, J.A., and Graf, T.
(2013). C/EBPalpha induces highly efficient macrophage transdifferentiation of B lymphoma and
leukemia cell lines and impairs their tumorigenicity. Cell Rep *3*, 1153-1163.

Reid, A., and Tursun, B. (2018). Transdifferentiation: do transition states lie on the path of
development? Curr Opin Syst Biol *11*, 18-23.

Rizki, M.T., and Rizki, R.M. (1959). Functional significance of the crystal cells in the larva of
Drosophila melanogaster. J Biophys Biochem Cytol *5*, 235-240.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source platform for biological-image
analysis. Nat Methods *9*, 676-682.

Sieweke, M.H., and Allen, J.E. (2013). Beyond stem cells: self-renewal of differentiated
macrophages. Science *342*, 1242974.

Sinenko, S.A., and Mathey-Prevot, B. (2004). Increased expression of *Drosophila* tetraspanin,
Tsp68C, suppresses the abnormal proliferation of ytr-deficient and Ras/Raf-activated hemocytes.
Oncogene *23*, 9120-9128.

Siripin, D., Kheolamai, P., Y, U.P., Supokawej, A., Wattanapanitch, M., Klincumhom, N.,
Laowtammathron, C., and Issaragrisil, S. (2015). Transdifferentiation of erythroblasts to
megakaryocytes using FLI1 and ERG transcription factors. Thromb Haemost *114*, 593-602.

Song, W., Onishi, M., Jan, L.Y., and Jan, Y.N. (2007). Peripheral multidendritic sensory neurons are
necessary for rhythmic locomotion behavior in *Drosophila* larvae. Proc Natl Acad Sci U S A *104*,
5199-5204.

Stewart, S., Koh, T.W., Ghosh, A.C., and Carlson, J.R. (2015). Candidate ionotropic taste receptors
in the *Drosophila* larva. Proc Natl Acad Sci U S A *112*, 4195-4201.

Tattikota, S.G., Hu, Y., Liu, Y., Cho, B., Barrera, V., Steinbaugh, M., Yoon, S.-H., Comjean, A., Li, F., Dervis, F.*, et al.* (2019). A single-cell survey of *Drosophila* blood. bioRxiv, 2019.2012.2020.884999.

Tokusumi, T., Shoue, D.A., Tokusumi, Y., Stoller, J.R., and Schulz, R.A. (2009). New hemocyte-specific enhancer-reporter transgenes for the analysis of hematopoiesis in *Drosophila*. Genesis *47*, 771-774.

Vermehren, A., Langlais, K.K., and Morton, D.B. (2006). Oxygen-sensitive guanylyl cyclases in insects and their potential roles in oxygen detection and in feeding behaviors. J Insect Physiol *52*, 340-348.

Vermehren-Schmaedick, A., Ainsley, J.A., Johnson, W.A., Davies, S.A., and Morton, D.B. (2010). Behavioral responses to hypoxia in *Drosophila* larvae are mediated by atypical soluble guanylyl cyclases. Genetics *186*, 183-196.

Vermehren-Schmaedick, A., Scudder, C., Timmermans, W., and Morton, D.B. (2011). *Drosophila* gustatory preference behaviors require the atypical soluble guanylyl cyclases. J Comp Physiol A Neuroethol Sens Neural Behav Physiol *197*, 717-727.

Wu, Q., Wen, T., Lee, G., Park, J.H., Cai, H.N., and Shen, P. (2003). Developmental control of foraging and social behavior by the *Drosophila* neuropeptide Y-like system. Neuron *39*, 147-161.

Xiang, Y., Yuan, Q., Vogt, N., Looger, L.L., Jan, L.Y., and Jan, Y.N. (2010). Light-avoidance-mediating photoreceptors tile the *Drosophila* larval body wall. Nature *468*, 921-926.

Xie, H., Ye, M., Feng, R., and Graf, T. (2004). Stepwise reprogramming of B cells into macrophages. Cell *117*, 663-676.

Yang, H., Kronhamn, J., Ekstrom, J.O., Korkut, G.G., and Hultmark, D. (2015). JAK/STAT signaling in *Drosophila* muscles controls the cellular immune response against parasitoid infection. EMBO Rep *16*, 1664-1672.

Figure Legends

Figure 1. Crystal cells in the *Drosophila* larva are generated by transdifferentiation

(A) Development of crystal cell numbers over time. Crystal cells per larva, assessed by melanization, relative to larval size and developmental stage. Genotype is w^{1118}, n=155. Mean and standard deviation, regression analysis.

(B) Development of total hemocyte numbers over time. Total hemocytes per larva, genotype is *HmlΔGAL4, UAS-GFP; He-GAL4*; n=107. Mean and standard deviation, regression analysis.

(C-C'') Localization of plasmatocytes and crystal cells in the 3rd instar larva,; genotype is *BcF2-GFP/ HmlΔ-DsRed*; plasmatocytes labeled by *HmlΔ-DsRed* (red); crystal cells labeled by *BcF2-GFP* (green); lateral view, posterior right.

(D-D') Model of distribution of plasmatocyte (red) and crystal cells (orange) in *Drosophila* larva, relevant sensory neuron clusters of the hematopoietic pockets (green); lateral view, posterior right.

(E-E''') Scheme of phagocytosis lineage tracing assay. Blue fluorescent latex beads are injected into early 3rd instar larvae, cells are released after indicated incubation times; fraction of cells containing phagocytosed beads are determined. Possible outcomes are depicted with crystal cells in orange, plasmatocytes in red, progenitors in grey.

(F) Sample image of analyzed cells; genotype *BcF2-GFP/ HmlΔ-DsRed*; crystal cells in green, plasmatocytes in red; blue arrowheads indicate crystal cells with incorporated blue beads.

(G) Quantification of samples as in (G); fraction of plasmatocytes and crystal cells containing blue beads at time 1h, 4, 22h after injection; n=21; mean and standard deviation.

Figure 2. Sensory Neuron activity regulates transdifferentiation to crystal cells

(A) Plasmatocytes colocalize with sensory neurons in all hematopoietic pockets; genotype *21-7-GAL4, UAS-CD8-GFP, HmlΔDsRed/CyO* ; lateral view, posterior right.

(B) Crystal cells also colocalize with sensory neurons but are mainly found in a cluster at the caudal end of the larva; genotype *21-7-GAL4, UAS-CD8-GFP/+; BCF6-mCherry/+*; lateral view, posterior right.

(C) Treatment of larvae with the AchR agonist carbachol to mimic sensory neuron activation, treatment for 4h; genotype is *yw*; quantification of crystal cells by melanization; control n=65 (+45); carbachol n=34 (+50). Individual value plot with mean and standard deviation, two-way ANOVA.

(D) Transient silencing of sensory neurons, quantification of crystal cells by melanization; genotypes are experiment *21-7-GAL4, UAS-CD8GFP, HmlΔ-DsRed/ UAS-Kir2.1; tubGAL80ts/ +* and control *21-7-GAL4, UAS-CD8GFP, HmlΔ-DsRed/ +; tubGAL80ts/ +*. Larvae induced at 29˚C for 22h. Mean and standard deviation, two-way ANOVA.

(E, F) Phagocytosis lineage tracing, genotypes are experiment *21-7-GAL4, UAS-CD8GFP, HmlΔ-DsRed/ UAS-Kir2.1; BcF6-GFP/ tubGAL80ts*, and control *21-7-GAL4, UAS-CD8GFP, HmlΔ-DsRed/ +; BcF6-GFP/ tubGAL80ts*.

(E) Sample image of analyzed cells, plasmatocytes (red), crystal cells (green), injected beads (blue).

(F) Quantification of samples as in (E); fraction of plasmatocytes and crystal cells containing blue beads, experiment n=12 and control n=12. Mean and standard deviation, two-way ANOVA.

Figure 3. Crystal cells are clustered around the sensory cones and are promoted by sensory cone neurons

(A-D) Localization of sensory neurons, plasmatocytes and crystal cells, caudal view of larvae; scale bars 0.25mm.

(A) Sensory neurons (green), genotype *21-7-GAL4, UAS-CD8-GFP, HmlΔ-DsRed/CyO*; 3rd instar larva.

(B) Plasmatocytes (green) and crystal cells (red), genotype *HmlΔ-GAL4, UAS-GFP; BcF6-mCherry*; 2nd instar larva.

(C) Crystal cells (red) and sensory neurons (green), genotype *21-7-GAL4, UAS-CD8-GFP/+; BcF6-mCherry/+*; 2nd instar larva.

(D) Crystal cells (red) and sensory neurons (green), *21-7-GAL4, UAS-CD8-GFP/+; BcF6-mCherry/+*; 3rd instar larva.

(A'-D') Models corresponding to (A-D), respectively, plasmatocytes red, crystal cells orange, sensory neurons green; caudal view.

(A''-D'') Models, lateral view, corresponding to (A-C and A'-C'), respectively; lateral view.

(E) *Gyc89Db-GAL4* driver expressing GFP in sensory cone neurons (green), lateral view, genotype is *UAS-GFP/+; Gyc89Db-GAL4/+*; lateral view, posterior right; scale bar 0.5mm.

(F) Larva as in (E), caudal view.

(G) Ablation of sensory cone neurons affects crystal cells; quantification of crystal cells per larva by melanization. Genotypes are experiment *UAS-Hid ala5/+; Gyc89Db-GAL4/tubGAL80ts*, n=34 and control *Gyc89Db-GAL4/tubGAL80ts*, n=34. Crosses were raised at 18˚C and temperature shifted to 29 ˚C for 16 h. Individual value plot with mean and standard deviation, two-way ANOVA.

(H) Transient activation of TrpA1 in sensory cone neurons; quantification of crystal cells per larva by melanization. Genotypes are experiment *UAS-TrpA1/ +; Gyc89Db-GAL4, / +*, n=46, and control *Gyc89Db-GAL4/+*, n=48; in addition, one experiment F1 cohort *UAS-TrpA1/ +; Gyc89Db-GAL4, / +*, n=47, was maintained as uninduced control at RT. Crosses were raised at RT and temperature shifted to 29˚C for 4 hours;. Individual value plot with mean and standard deviation, two-way ANOVA.

Figure 4. Oxygen sensing through Gycs in sensory cone neurons drives plasmatocyte-to-crystal cell transdifferentiation

(A) RNAi silencing of *Gyc88E* in sensory cone neurons results in reduced crystal cell numbers determined by melanization; genotyes are experiment *Gyc89Db-GAL4/UAS-Gyc88ERNAi*, n=45 and control *Gyc89Db-GAL4/+*. n=45. Individual value plot with mean and standard deviation, two-way ANOVA.

(B) Phagocytosis lineage tracing, effect of *Gyc88E* RNAi in sensory cone neurons on transdifferentiation; genotypes are experiment *21-7-GAL4, UA5-CD8-GFP, HmlΔ-DsRed/+; BcF6-GFP/ UAS-Gyc88ERNAi*, n=11 and control *21-7-GAL4, UA5-CD8-GFP, HmlΔ-DsRed/+; BcF6-GFP/+*, n=16. Bar chart with mean and standard deviation, two-way ANOVA.

(C) Effect of hypoxia (5% O2) on crystal cell number per larva determined by melanization; genotype is w^{1118}; hypoxia n=46 and normoxia n=48. Individual value plot with mean and standard deviation, two-way ANOVA.

(D) Phagocytosis lineage tracing, effect of hypoxia (5% O2) on transdifferentiation; genotype is *HmlΔ-GAL4, UAS-GFP; BcF6-mCherry*; hypoxia n=14 and normoxia n=15. Bar chart with mean and standard deviation, two-way ANOVA.

 (E) Model. Sensory cone neurons detect oxygen by cytoplasmic Gyc heterodimeric oxygen sensors. Gycs convert GTP to cGMP, which activates CNG channels, resulting in influx of calcium (Ca2+) leading to downstream signaling and neuronal activation. Active neurons induce plasmatocyte-to-crystal cell transdifferentiation.

Supplemental Materials

Supplemental Figure 1. Methods of crystal cell labeling

(A) Heat induced melanization of crystal cells. Genotype is *lz-GAL4;UAS-GFP*

(B) Fluorescent reporter labeling to visualize, and quantify, crystal cells. Genotype is *lz-GAL4;UAS-GFP*. Note that the labeled crystal cell pattern by both methods is very similar.

Supplemental Figure 2. Limited transient sensory neuron silencing does not affect total hemocyte numbers

(A) Transient silencing of sensory neurons, quantification of total hemocytes; genotypes are *21-7-GAL4, UAS-CD8-GFP, HmlΔ-DsRed/ UAS-Kir2.1; tubGAL80ts/ +* , n=12 and control *21-7-GAL4, UAS-CD8-GFP, HmlΔ-DsRed/ +; tubGAL80ts/ +*, n=13. Larvae induced at 29˚C for 22h. Mean and standard deviation, two-way ANOVA

Supplemental Figure 3. Hypoxia affects crystal cell counts but not total hemocyte numbers

 (A) Effect of 8% hypoxia on crystal cell number determined by melanization; genotype is *w1118*; hypoxia n=19 and normoxia control n=18. Individual value plot with mean and standard deviation, two-way ANOVA.

(B) Total hemocyte number under conditions of hypoxia (5% O2) and normoxia; genotype is *HmlΔ-GAL4, UAS-GFP; BcF6-mCherry*; hypoxia n=14 and normoxia control n=14. Mean and standard deviation, two-way ANOVA.

Supplemental Figure 4. Models

(A) Stimulated by oxygen, activated sensory cone neurons produce signal/s that drive transdifferentiation of a fraction of plasmatocytes to crystal cells. According to this model, transdifferentiation is triggered by contacting the sensory neuron signal, therefore plasmatocytes in anatomical proximity to the sensory cone neurons are most likely to convert into crystal cells.

666   (B) Model illustrating exposure of the caudal end of *Drosophila* larvae including the sensory cones
667   and posterior spiracles to the air, while burying in food. Mature larvae leave the food in preparation
668   of pupariation, now fully exposed to the air.

A

Crystal Cells

$y = 27.291e^{0.3521x}$
$R^2 = 0.97876$

Crystal Cells/larva

1.4-1.6 mm
1.7-1.9 mm
2.0-2.2 mm
2.3-2.5 mm
2.6-2.8 mm
2.9-3.1 mm
3.2-3.4 mm

1st · 2nd · 3rd

B

Total Hemocytes

$y = 414.17e^{0.3673x}$
$R^2 = 0.97852$

Plasmatocytes/larva

1.4-1.6 mm
1.7-1.9 mm
2.0-2.2 mm
2.3-2.5 mm
2.6-2.8 mm
2.9-3.1 mm
3.2-3.4 mm

1st · 2nd · 3rd

C *BcF2-GFP* *HmlΔ-DsRed*

C' *HmlΔ-DsRed* Plasmatocytes

C'' *BcF2-GFP* Crystal Cells

D Plasmatocytes

D' Sensory Neurons Crystal Cells

E Inject larvae with blue beads → Analyse Hemocytes

E' Plasmatocyte → Transdifferentiation

E'' Crystal Cell → Homeostasis

E''' Progenitor → Differentiation

F Latex Beads *BcF2-GFP* *HmlΔ-DsRed*

G Transdifferentiation timecourse

percent bead+ cells

1h 4h 22h

plasmatocytes
crystal cells

Corcoran et al., Figure 1

Corcoran et al. Figure 2

Corcoran et al. Figure 3

Corcoran et al. Figure 4

Corcoran et al. Supplemental Figure 1

Corcoran et al. Supplemental Figure 2

A   Crystal Cells
     Hypoxia (8% O$_2$)

B   Total Hemnocytes
     Hypoxia (5% O$_2$)

Corcoran et al. Supplemental Figure 3

A

O₂

Sensory Neurons

Plasmatocytes   Crystal Cells

Influence of
Sensory Neurons

B

Atmosphere

wandering
3rd instar

exposure to air

Posterior Spiracles

Sensory Cones

1st   2nd   3rd instar

Food Source

Corcoran et al. Supplemental Figure 4