1   **Multiple environmental parameters impact core lipid cyclization in Sulfolobus acidocaldarius**
2

3   Alec Cobban[a,b,*], Yujiao Zhang[a,c], Alice Zhou[a,#], Yuki Weber[^,d], Ann Pearson[d], William D. Leavitt[a,b,e,*]
4
[a.] Department of Earth Sciences, Dartmouth College, Hanover, NH 03755 USA
[b.] Department of Biological Sciences, Dartmouth College, Hanover, NH 03755 USA
[c.] State Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy
of Sciences, Guangzhou 510640 China
[d.] Department of Earth & Planetary Sciences, Harvard University, Cambridge, MA 02138 USA
[e.] Department of Chemistry, Dartmouth College, Hanover, NH 03755 USA
11
12   *Correspondence authors e-mail addresses: alec.bradley.cobban@dartmouth.edu,
13   william.d.leavitt@dartmouth.edu
14   Key Words: GDGT; ring index; thermoacidophile; temperature; pH; growth rate
15   Current Address: [#]Department of Earth Science, University of Michigan; [^]Greenlight Biosciences Inc.,
16   Medford, MA
17

ABSTRACT

Environmental reconstructions based on microbial lipids require understanding the coupling

between environmental conditions and membrane physiology. The paleotemperature proxy TEX$_{86}$ is

built on the observation that archaea alter the number of five- and six-membered rings in the

hydrophobic core of their glycerol dibiphytanyl glycerol tetraether (GDGT) membrane lipids when

growing at different temperatures. However, recent work with these archaea also highlights a role

for other factors, such as pH or energy availability in determining the degree of core lipid cyclization.

To better understand the role of these variables we cultivated a model Crenarchaeon, *Sulfolobus*

***acidocaldarius*, over a range in temperature, pH, oxygen flux, or agitation speed, and quantified the**

changes in growth rate, biomass yield, and core lipid compositions. The average degree of cyclization

in core lipids correlated with growth rate under most conditions. When considered alongside other

experimental findings from both the thermoacidophilic and mesoneutrophilic archaea, the results

suggest the cyclization of archaeal lipids records a universal response to energy availability at the

cellular level. Although we isolated the effects of individual parameters, there remains a need for

multi-factor experiments (e.g., pH + temperature + redox) to establish a robust framework to

interpret biomarker records of environmental change.

1.0 Introduction

Key taxa from the domain Archaea produce membranes partly or mostly composed of the geostable

lipid biomarkers, glycerol dibiphytanyl glycerol tetraether (GDGTs (1, 2). Archaea that produce GDGTs

maintain optimal membrane fluidity and permeability by altering the number of cyclopentyl and/or

cyclohexyl rings within the tetraether core (3). Early work with thermoacidophilic archaea showed that

temperature influences GDGT ring abundance (4, 5), which formed the basis for using GDGT distributions

as a paleotemperature proxy, known as TEX_{86} (6). Early studies also implicated pH and oxidant availability

in altering ring abundance (5, 7). Recent work with marine mesoneutrophilic Thaumarchaeota showed that

electron donor or acceptor availability is an important determinant of GDGT cyclization (8, 9), and work

with thermoacidophilic Crenarchaeota demonstrated that shifts in pH, temperature, growth phase, or

electron donor availability also resulted in different degrees of GDGT cyclization (10–14). These changes

in GDGT composition highlight the need for cells to maintain membrane fluidity and permeability within

an operative range (15, 16). Modelling studies show that cyclization of GDGTs increases the degree of

membrane packing by bringing both the isoprenoid chains and polar head groups closer together (17).

Tighter membrane packing may be one of the many strategies archaea evolved to survive chronic energy

stress (18). It follows that not only temperature, but also the other major environmental variables that play

a significant role in determining membrane composition, must be considered in order to understand GDGT-

based reconstructions of past conditions.

To better interpret GDGT ring abundance in response to environmental change we cultivated the

model thermoacidophile, *Sulfolobus acidocaldarius* DSM639 (hereafter DSM639), under different regimes

of pH, temperature, physical perturbation, or oxygen availability. This model organism grows quickly and

to high densities, has the ability to grow under different culture conditions, has a suite of genetic tools

available for downstream manipulation (19), and has a known biochemical mechanism of ring cyclization

(20). In this study we quantified core GDGTs and the average abundance of cyclopentyl rings, as well as

growth parameters such as doubling time and biomass yield. We observed clear trends between growth rate

and ring abundance across all conditions. Under cultivation conditions farthest from the organismal

optimum, DSM639 produced several additional GDGT isomers in addition to the typical suite of GDGTs.

We discuss the implications for batch and bioreactor-based cultivation efforts and highlight the need for

multiparameter studies going forward.

2.0   Methods

Axenic cultures of DSM639 were cultivated in either batch cultures exchanging with the

atmosphere or in closed, gas-fed batches. Medium was prepared following Wagner et al. (2012), with

sucrose (0.2% w/v) and oxygen as the electron donor/acceptor pair. In closed batch experiments a single

parameter was varied per experiment – temperature (65, 70, 75 and 80 °C), pH (2, 3, 4), or shaking speed

(0, 50, 61, 75, 97, 125, 200 or 300 RPM) – where default "optimal" parameters were 70 °C, pH 3, and 200

RPM. Five biological replicates were cultivated per condition. Atmospheric batch experiments were

performed in temperature-controlled shaking incubators (Innova42 New Brunswick, Eppendorf,

Hauppauge NY, USA), using the incubator's digital control for maintenance of both temperature and

shaking speed. Evaporation was minimized by maintaining large trays of water in the incubator to saturate

the atmosphere. Gas fed-batch experiments were performed in three 1-L glass bioreactors using myControl

PID controllers (Applikon, Delft, Netherlands). In all gas-fed batch experiments the following parameters

were held constant: temperature of 70 °C, impeller speed of 200 RPM, gas flux of 200 mL per min, and pH

3. Strain DSM639 was pre-cultured in batch prior to inoculation into gas fed-batch reactors. The sparge gas

contained oxygen partial pressures of either 0.2, 0.5, 1.0, 2.0 or 20 % (balance N_2). When DSM639 was

grown on 0.2% O_2 it underwent fewer than 2 generations before entering stationary phase. Due to the

concern that the harvested biomass was significantly influenced by the inoculum, we performed a second

(subculture) run on 0.2% O_2, where two reactors of fresh 0.2% O_2-sparged media were inoculated with late-

logarithmic phase cells from an initial 0.2% O_2 experiment. Growth was determined by optical density

measurements at 600 nm.

Biomass for lipids was harvested in early stationary phase for atmospheric batch conditions, and

both late logarithmic and early stationary phase for gas-fed batch (*Figure 1*). Biomass samples were pelleted

by centrifugation, frozen at -80°C, and freeze-dried prior to lipid extraction. Core lipid fractions were

obtained and analyzed as previously detailed (13, 21). Specific growth rates were calculated by a modified

implementation of the algorithm in Hall et al. (2014). Rate calculation script is available on GitLab (see

Supplement). GDGT abundances were measured by ultra-high-performance liquid chromatography

(UHPLC) coupled to an Agilent 6410 triple-quadrupole mass spectrometer (MS) as previously described

(13). Ring indices (RI) were calculated using Eq 1. based on peak areas of GDGTs as modified from ref.

(22).

$$RI=\frac{1*[GDGT\text{-}1] + 2*[GDGT\text{-}2] + 3*[GDGT\text{-}3] + 4*[GDGT\text{-}4] + 5*[GDGT\text{-}5] + 6*[GDGT\text{-}6] + 7*[GDGT\text{-}7] + 8*[GDGT\text{-}8]}{[GDGT\text{-}0]+[GDGT\text{-}1]+[GDGT\text{-}2]+[GDGT\text{-}3]+[GDGT\text{-}4]+[GDGT\text{-}5]+[GDGT\text{-}6]+[GDGT\text{-}7]+[GDGT\text{-}8]} \quad (1)$$

Multiple isomers of GDGTs- 3, 4 and 5 were detected in some experiments, similar to samples from ref.

(13). The peak areas of the minor isomers were summed with their respective major components (e.g.,

GDGT-3 + GDGT-3') before calculating RI.

A three-way Type II analysis of variance (ANOVA) was run on GDGT relative abundances for

the atmospheric batch experiments. A one-way Type II ANOVA was run on relative abundances of each

GDGT for oxygen concentration experiments. Non-metric multidimensional scaling (NMDS) analyses

were conducted for both atmospheric batch and gas-fed batch experiments using the Bray-Curtis

dissimilarity of the absolute abundances of each GDGT, and the first two dimensions were plotted. Code

for all statistical analyses is available on GitLab (https://git.dartmouth.edu/leavitt_lab/cobban-saci-lipids-

batch-and-fed-batch-2020).

3.0   Results

DSM639 grew faster and to generally higher terminal densities at higher shaking speeds (Figure

1A), higher temperatures (Figure 1B), and at higher oxygen partial pressures (Figure 1D). DSM639 was

less noticeably sensitive to pH (Figure 1C); its growth optimum is thought to be pH 3 (23), although we

observed similar growth rates at all three pH values and higher biomass yields at pH 2 and 4. The GDGT

distributions from all replicates are shown in Figure 2 and Table S1. Some conditions yielded multiple

isomers of GDGTs 3, 4 and 5, previously seen only in chemostats under energy limitation (13). These extra

isomers were observed at higher temperatures (75 and 80°C; Figure 2A), low pH (Figure 2B), shaking

speeds above 125 RPM (Figure 2C), and under oxygen partial pressures below 0.5% O_2.

We performed statistical analyses to determine how each GDGT was individually impacted by

changing culture conditions. Main effects ANOVA was performed on GDGT relative abundances for each

experimental condition (Table 1). Temperature correlated with significant variation in relative abundances

of all GDGTs except for GDGT-0, 1 and the isomer of GDGT-3. Shaking speed was associated with

significant variation in all relative abundances except GDGT-3, the GDGT-3 isomer, GDGT-7, and GDGT-

8. In response to pH, all GDGTs varied significantly except for GDGT-0, -1, -2, -7 and -8. Oxygen flux

caused variation in all observed GDGTs, except GDGT-0, -3, -3 isomer, and -4. We also reanalyzed data

from a recent chemostat experiment we performed with DSM639 (13), finding that growth rate

significantly correlated with the relative abundance of all GDGTs, except GDGT-3 isomer. Considering in

aggregate our atmospheric batch, gas-fed batch, and chemostat experiments with DSM639, our results

demonstrate that each physical variable can independently influence different subsets of GDGT core lipids.

Doubling time and RI were determined from each replicate under each condition (Figure 3). As

temperature increased, doubling time decreased and RI increased (Figure 3A and 3E). Doubling was fastest

at pH 3, slightly slower as pH 2, and slowest at pH 4, while RI was highest at pH 3, lower at pH 2 and

lowest at pH 4 (Figure 3B and 3F). Growth was faster and RI increased along with shaking speed from 50

to 200 RPM, with exceptions at 0 and 300RPM (Figure 3C and 3G). In gas-fed experiments, growth was

the slowest at 0.2% O_2, faster at 0.5, 1.0 and 2 % O_2, and fastest at 20% O_2. In these experiments RI was

highest at 20% O_2, then dropped as % O_2 decreased to 0.5 %, but then increased again at the lowest O_2

concentration of 0.2 % (Figure 3D and 3H).

The RI results from each experimental condition were compared using doubling time as the

reference variable (Figure 4). Interestingly, linear regressions for all of these different experimental

conditions coalesce around a pattern of negative correlation between RI and doubling time in atmospheric

batch experiments (Figure 4A-C), but positive correlation between RI and doubling time in gas fed-batch

and chemostat experiments (Figure 4D and Figure 4E).

Differences in core GDGT composition between samples was compared via NMDS using Bray-

Curtis dissimilarity, separately for atmospheric batch, fed-batch, and chemostat cultivation experiment

(Figures 5 and 6). In batch conditions there is visually distinct clustering of samples based on temperature,

pH, or shaking speed (Figure 5). Closer points on the NMDS are more similar in overall GDGT composition

than they are to points farther away. This approach can identify unique GDGT distributions in samples

which may otherwise appear similar in terms of consolidated metrics, such as RI. In gas fed-batch

experiments, separate clusters can be seen based on both growth phase and oxygen concentration (Figure

6A). In chemostat experiments from Zhou et al. (2019) clusters seem to arise based on doubling times

(Figure 6B). Stress values for all conditions are plotted on each NMDS plot, and were all below 0.05,

indicating a good fit of the original data to the two-dimensional ordination.

4.0   Discussion

Our studies of the model thermoacidophile DSM639 are broadly consistent with prior data, including

work on other thermoacidophilic Crenarchaeota as well as the more distantly related mesoneutrophilic

Thaumarchaeota. The results support the previously demonstrated roles of pH, temperature, and oxygen

(electron acceptor) availability in affecting core GDGT composition. We further examine, for the first time,

the impact of agitation on GDGT cyclization. Higher temperatures yielded higher average cyclization,

which is consistent with most experimental studies of both Thaum- and Crenarchaeota (9–12, 24, 25), yet

opposes recent environmental findings in alkaline hotsprings (26). When pH deviated from the optimum

pH of ~3 for DSM639, cyclization decreased. Whereas prior thermoacidophile studies showed that

cyclization primarily decreased with increases in pH (10, 11, 25), we observed decreased cyclization when

the pH was both above and below optimal. Despite the similarity in RI values at pH 2 and pH 4, the

composition of GDGTs at each pH also was distinct (Figure 2, Figure 5). GDGT profiles contained more

of both the higher and lower ring-numbered GDGTs at pH 2, but showed more of the moderately cyclized

(GDGT-3, -4) compounds at pH 4. These observations highlight that differences in GDGT composition are

not always captured by RI values. The results of ANOVA show that all conditions tested had a significant

effect on a unique subset of GDGTs, as opposed to unidirectional or single-pattern control over the total

lipid composition (Table 1). This makes it difficult to assess the robustness of uniform metrics that

consolidate GDGT composition data into a single value, e.g., RI or TEX_{86}. This complication affects

application of such index values to environments in which there may be multiple variables experiencing

simultaneous shifts caused by different environmental drivers (e.g., ref. (27)) .

169   To our knowledge, this study is the first to specifically address the effects of agitation speed on

GDGT composition. Shaking of microbial cultures is classically believed to provide aeration and increase

oxygen availability. However, we saw contradictory trends in RI as a function of shaking speed, with

outlying data at 0 and 300 RPM (Figure 3G). Between these ranges, however, RI increased as growth rates

and shaking speed increased. The inconsistent response to shaking, along with discrepancies within

the % O_2 experiments, as discussed below, suggests that the effects of shaking are not solely tied to O_2

availability in the growth medium. A previous study in another model taxon suggested that shaking,

independent of aeration, affects growth by inducing stress caused by physical interaction with the

environment (28). Physical stress may have caused the anomalous trend at 300 RPM, while the trend at 0

RPM may reflect O_2-limitation (similar to the 0.2% O_2 experiment, Figure 3H).

179   The oxygen partial pressure experiments expand on the existing understanding of GDGT response

to electron acceptor availability. Our data are consistent with recent batch experiments performed on

mesoneutrophilic Thaumarchaeota, where RI generally decreased with initial headspace % O_2 (9). All such

experiments (ours and ref. (9)) were isothermal, of constant pH, and tested similar ranges of % O_2. Taken

together, these data show that two distinct groups of archaea both exhibit consistent responses of lipid

composition to trends in O_2 partial pressures. Importantly, both our data and the results from Qin and

colleagues (9) differ from the agitation speed experiments, which show the opposite relationship between

RI and growth rate. The oxygen limitation experiments performed here were mixed with an impeller, while

those from Qin and colleagues (9) were performed without shaking – in both cases, electron-acceptor

availability was modified without altering the other environmental factors that may otherwise change in the

agitation speed experiments. From the variation in RI in these experiment types, it seems possible that

higher ring indices are associated with higher metabolic energy stress and mechanical/environmental

stresses.

Growth rate and ring index broadly covary, independent of which variable is causing this forcing

(Figure 4). The temperature, pH and shaking speed experiments show inverse trends between growth rate

and RI, consistent with batch experiments in the existing Thaumarchaeota (Figure S1) and Crenarchaeota

(Figure S2) literature (11, 12, 29). However, these trends directly oppose results from other experiments

characterizing GDGT response to pH in thermoacidophiles (11). It is possible that the fastest growth

conditions in batch culture are those with high environmental or energetic stress (high temperature/shaking

speed), and this increase in stress necessitates generally higher cyclization for the cells to maintain

homeostasis. For pH, it is possible that GDGT distribution may be changing in a way that is not best

recorded by RI, leading to a lack of consistent trends when only these simple index values are reported. The

gas-fed batch experiments showed consistent positive trends between RI and doubling time (more rings at

slower growth rate) as in other bioreactor-based experiments with both thermoacidophiles and ammonia

oxixidizing archaea (8, 13). Because the bioreactor-based experiments vary only in electron donor or

acceptor availability, – i.e., have constant physical conditions, pH, and growth stage – the environmental

stresses across each set of experiments should be equivalent. Thus, the growth rates and RI values of

bioreactor experiments specifically are responding to variations in energy supply and demand. Cell

responses, including extent of GDGT cyclization, should be tied to energy availability. This disconnect

between energy availability versus a mixture of environmental stressors may be at the root of the opposing

trends of RI vs. growth rates for batch (Figure 4 A, B, C) and bioreactor (Figure 4 D, E; refs. 8, 13)

experiments.

The occurrence of multiple isomers of a GDGT (e.g., GDGT-3 and GDGT-3’) consistently

occurred when DSM639 was cultivated under conditions we interpret as the most physiologically stressful.

Namely, the isomers were most abundant at the lowest pH, highest temperature, highest shaking speed, and

lowest oxygen concentrations (Figure 2). It is possible that these isomers were produced as part of a generic

cellular stress response, or that they have different physical properties that are advantageous in less

hospitable living conditions.

Finally, the overall distribution of lipids changed dramatically in response to shifts in temperature,

pH, shaking speed or oxygen concentration in directions that were not reflected by RI alone. The clearest

example of this is the response to pH (Figure 2B), where the pH 2 and 4 experiments were more similar in

RI value than they were to the RI value for pH 3, but differed markedly in their GDGT distributions. This

may indicate that despite having similar RI values, their associated lipid profiles incorporated different

mixtures of GDGTs that could have opposing effects on membrane fluidity and permeability. This

observation highlights how taking the weighted average of GDGTs in a measurement like RI can mask

variance. It is critical to note this when considering shifts in RI in natural systems where independent

measures of pH, temperature, or oxygen availability, for example, may not be available.

226   GDGT cyclization by both thermoacidophile and mesoneutrophile archaea reflects strain-specific

membrane optimization to a wide array of environmental perturbations. Insights from experiments such as

our present work are supported by observations from redox-stratified marine and terrestrial systems where

either electron donor or acceptor availability, or other major parameters such as temperature or pH, are

statistically correlated with shifts in GDGT composition (30–35). Resolving how environmental parameters

trigger archaea to vary their degree of GDGT cyclization in complex systems remains a core challenge.

5.0   Conclusions

234   Our experiments indicate that Crenarchaeota alter their membrane lipid composition in response to

temperature,   pH,   shaking   speed,   and   oxygen   concentration.   Higher   physical   stress   (high

temperatures/shaking speeds) and lower energy availability (low % O_2) were mostly associated with higher

RI values, while sub- and super-optimal pH caused a variety of changes to lipid composition that were not

visible in consolidated RI values. This physiological plasticity likely allows cells to maintain optimal

membrane fluidity and permeability, as well as minimize energy loss, when confronted with chemical and

physical stress. This survival strategy of varying GDGT cyclization in response to a variety of

environmental challenges is common to the Cren- and Thaumarchaeota; such observations ultimately

support Valentine's hypothesis (17) that the unifying physiological and evolutionary feature of the archaeal

domain is the ability to persist under conditions of chronic energy stress. Expanding the experimental matrix

tested here is a priority going forward to better constrain the covariate interactions of these environmental

conditions in affecting GDGT profiles. This is more easily done in the thermoacidophiles due to their ease

of cultivation, but ultimately must be extended to the Thaumarchaeota and to mixed assemblages, in order

to better understand paleoenvironmental proxies such as TEX_{86}.

6.0 Supplementary information, Data and Code.

All supplemental code and data available online:

https://git.dartmouth.edu/leavitt_lab/cobban-saci-lipids-batch-and-fed-batch-2020

Figures and dataframes are also available at:

https://doi.org/10.6084/m9.figshare.c.4863426.v1

7.0 Acknowledgments

Funding was provided by the American Chemical Society PRF #57209-DNI2 (WDL), the Walter and

Constance Burke Fund at Dartmouth College (WDL), and Dartmouth College Undergraduate Advising and

Research (UGAR) at Dartmouth (AC); the Visiting PhD Research program Fund at Guangzhou Institute of

Geochemistry (YZ); the Swiss National Science Foundation P2BSP2_168716 (YW); NSF OCE-1843285

and OCE-1702262 (AP). We thank Sonja Albers at the University of Freiburg for providing *S.*

acidocaldarius DSM639, Beverly Chiu for editorial and laboratory assistance, and Dr. Felix Elling for

assistance with GDGT analyses.

263
264
265

8.0 References

1.   Pearson A, Ingalls AE. 2013. Assessing the Use of Archaeal Lipids as Marine

Environmental Proxies. Annu Rev Earth Planet Sci 41:359–384.

2.   Schouten S, Hopmans EC, Sinninghe Damsté JS. 2013. The organic geochemistry of

glycerol dialkyl glycerol tetraether lipids: A review. Org Geochem 54:19–61.

3.   Gliozzi A, Relini A, Chong PL-G. 2002. Structure and permeability properties of

biomimetic membranes of bolaform archaeal tetraether lipids. J Membr Sci 206:131–147.

4.   De Rosa M, Esposito E, Gambacorta A, Nicolaus B, Bu'Lock JD. 1980. Effects of

temperature on ether lipid composition of Caldariella acidophila. Phytochemistry 19:827–

831.

5.   De Rosa M, Gambacorta A. 1988. The lipids of archaebacteria. Prog Lipid Res 27:153–175.

6.   Schouten S, Hopmans EC, Schefuß E, Sinninghe Damsté JS. 2002. Distributional variations

in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water

temperatures? Earth Planet Sci Lett 204:265–274.

7.   Zillig W, Holz I, Janekovic D, Schäfer W, Reiter WD. 1983. The Archaebacterium

Thermococcus celer Represents, a Novel Genus within the Thermophilic Branch of the

Archaebacteria. Syst Appl Microbiol 4:88–94.

8.   Hurley SJ, Elling FJ, Könneke M, Buchwald C, Wankel SD, Santoro AE, Lipp JS, Hinrichs

K-U, Pearson A. 2016. Influence of ammonia oxidation rate on thaumarchaeal lipid

composition and the TEX86 temperature proxy. Proc Natl Acad Sci 113:7762–7767.

9.   Qin W, Carlson LT, Armbrust EV, Devol AH, Moffett JW, Stahl DA, Ingalls AE. 2015.

Confounding effects of oxygen and temperature on the TEX86 signature of marine

Thaumarchaeota. Proc Natl Acad Sci 112:10979–10984.

10.   Boyd ES, Pearson A, Pi Y, Li W-J, Zhang YG, He L, Zhang CL, Geesey GG. 2011.

Temperature and pH controls on glycerol dibiphytanyl glycerol tetraether lipid composition

in the hyperthermophilic crenarchaeon Acidilobus sulfurireducens. Extremophiles 15:59–

65.

11.   Feyhl-Buska J, Chen Y, Jia C, Wang J-X, Zhang CL, Boyd ES. 2016. Influence of growth

phase, pH, and temperature on the abundance and composition of tetraether lipids in the

Thermoacidophile Picrophilus torridus. Front Microbiol 7:1323.

12.   Jensen SM, Neesgaard VL, Skjoldbjerg SLN, Brandl M, Ejsing CS, Treusch AH. 2015. The

Effects of Temperature and Growth Phase on the Lipidomes of Sulfolobus islandicus and

Sulfolobus tokodaii. Life 5:1539–1566.

13.   Zhou A, Weber Y, Chiu BK, Elling FJ, Cobban AB, Pearson A, Leavitt WD. 2019. Energy

flux controls tetraether lipid cyclization in Sulfolobus acidocaldarius. Environ Microbiol

22:343–353.

14.   Quehenberger J, Pittenauer E, Allmaier G, Spadiut O. 2020. The influence of the specific

growth rate on the lipid composition of Sulfolobus acidocaldarius. Extremophiles.

15.   Oger PM, Cario A. 2013. Adaptation of the membrane in Archaea. Biophys Chem 183:42–

56.

16.   van de Vossenberg JL, Driessen AJ, Konings WN. 1998. The essence of being

extremophilic: the role of the unique archaeal membrane lipids. Extremophiles 2:163–170.

17.   Shimada H, Nemoto N, Shida Y, Oshima T, Yamagishi A. 2008. Effects of pH and

Temperature on the Composition of Polar Lipids in Thermoplasma acidophilum HO-62. J

Bacteriol 190:5404–5411.

18.   Valentine DL. 2007. Opinion: Adaptations to energy stress dictate the ecology and

evolution of the Archaea. Nat Rev Microbiol 5:316–323.

19.   Wagner M, van Wolferen M, Wagner A, Lassak K, Meyer BH, Reimann J, Albers S-V.

2012. Versatile Genetic Tool Box for the Crenarchaeote Sulfolobus acidocaldarius. Front

Microbiol 3:214.

20.   Zeng Z, Liu X-L, Farley KR, Wei JH, Metcalf WW, Summons RE, Welander PV. 2019.

GDGT cyclization proteins identify the dominant archaeal sources of tetraether lipids in the

ocean. Proc Natl Acad Sci 116:22505.

21.   Elling FJ, Könneke M, Nicol GW, Stieglmeier M, Bayer B, Spieck E, de la Torre JR,

Becker KW, Thomm M, Prosser JI. 2017. Chemotaxonomic characterisation of the

thaumarchaeal lipidome. Environ Microbiol 19:2681–2700.

22.   Hall BG, Acar H, Nandipati A, Barlow M. 2014. Growth Rates Made Easy. Mol Biol Evol

31:232–238.

23.   Grogan DW. 1989. Phenotypic characterization of the archaebacterial genus Sulfolobus:

comparison of five wild-type strains. J Bacteriol 171:6710–6719.

24.   Elling FJ, Könneke M, Mußmann M, Greve A, Hinrichs K-U. 2015. Influence of

temperature, pH, and salinity on membrane lipid composition and TEX86 of marine

planktonic thaumarchaeal isolates. Geochim Cosmochim Acta 171:238–255.

25.   Kaur G, Mountain BW, Stott MB, Hopmans EC, Pancost RD. 2015. Temperature and pH

control on lipid composition of silica sinters from diverse hot springs in the Taupo Volcanic

Zone, New Zealand. Extremophiles 19:327–344.

26.   Boyer GM, Schubotz F, Summons RE, Woods J, Shock EL. 2020. Carbon Oxidation State

in Microbial Polar Lipids Suggests Adaptation to Hot Spring Temperature and Redox

Gradients. Front Microbiol 11.

27.   Polik CA, Elling FJ, Pearson A. 2018. Impacts of Paleoecology on the TEX86 Sea Surface

Temperature Proxy in the Pliocene-Pleistocene Mediterranean Sea. Paleoceanogr

Paleoclimatology 33:1472–1489.

28.   Juergensmeyer MA, Nelson ES, Juergensmeyer EA. 2007. Shaking alone, without

concurrent aeration, affects the growth characteristics of Escherichia coli. Lett Appl

Microbiol 45:179–183.

29.   Elling FJ, Könneke M, Lipp JS, Becker KW, Gagen EJ, Hinrichs K-U. 2014. Effects of

growth phase on the membrane lipid composition of the thaumarchaeon Nitrosopumilus

maritimus and their implications for archaeal lipid distributions in the marine environment.

Geochim Cosmochim Acta 141:579–597.

30.   Hurley SJ, Lipp JS, Close HG, Hinrichs K-U, Pearson A. 2018. Distribution and export of

isoprenoid tetraether lipids in suspended particulate matter from the water column of the

Western Atlantic Ocean. Org Geochem 116:90–102.

31.   Pearson A, Pi Y, Zhao W, Li W, Li Y, Inskeep W, Perevalova A, Romanek C, Li S, Zhang

CL. 2008. Factors Controlling the Distribution of Archaeal Tetraethers in Terrestrial Hot

Springs. Appl Environ Microbiol 74:3523–3532.

32.   Sollich M, Yoshinaga MY, Häusler S, Price RE, Hinrichs K-U, Bühring SI. 2017. Heat

stress dictates microbial lipid composition along a thermal gradient in marine sediments.

Front Microbiol 8:1550.

33.   Weber Y, Damsté JSS, Zopfi J, De Jonge C, Gilli A, Schubert CJ, Lepori F, Lehmann MF,

Niemann H. 2018. Redox-dependent niche differentiation provides evidence for multiple

bacterial sources of glycerol tetraether lipids in lakes. Proc Natl Acad Sci 115:10926–

10931.

34. Xie W, Zhang CL, Wang J, Chen Y, Zhu Y, de la Torre JR, Dong H, Hartnett HE, Hedlund

BP, Klotz MG. 2015. Distribution of ether lipids and composition of the archaeal

community in terrestrial geothermal springs: impact of environmental variables. Environ

Microbiol 17:1600–1614.

35. Zhang Z, Smittenberg RH, Bradley RS. 2016. GDGT distribution in a stratified lake and

implications for the application of TEX_{86} in paleoenvironmental reconstructions. Sci Rep

6:34465.

366
367

Table 1: Results of ANOVA for GDGT relative abundances based on variation in experimental conditions

Factor	GDGT.0	GDGT.1	GDGT.2	GDGT.3	GDGT.3.iso	GDGT.4	GDGT.4.iso	GDGT.5	GDGT.5.iso	GDGT.6	GDGT.7	GDGT.8
Temperature	NS	NS	**	***	NS	***	***	***	***	***	***	***
RPM	***	***	***	NS	NS	***	**	***	**	**	NS	NS
pH	NS	NS	NS	***	***	***	***	**	***	***	NS	NS
O_2 Sparge	NS	*	**	NS	NS	~	**	*	**	*	NA	NA
Doubling Time	***	***	***	***	NS	***	***	***	***	***	***	***

374 Results of type-II three-way ANOVAs (Temperature, RPM, and pH) and one-way ANOVAs (O_2 sparge and doubling time individually). Data for targeted
375 doubling time are from Zhou et al. (2019). Notations: ***, $p<0.001$; **, $p<0.01$; *, $p<0.05$; ~, $0.05<p<0.06$; NS, $p>0.06$; NA, below detection limit.

376
Figure 1:

Figure 1. Averaged growth curves from each experiment, plotted as the $Log_2(OD_{600})$ versus time
in hours. (A, B, C) Atmospheric-batch only and (D) gas-fed batch experiments (see methods for
details). Error bars show the mean ±1 SE of $Log_2(OD_{600})$ values for the five replicates (A, B, C),
or two to three replicates (D).

Figure 2:

385
Figure 2. The core GDGT distribution from each replicate in each experiment. The batch
experiments for temperature (A), pH (B) or shaking speed (C) each had five replicates. The fed-
batch experiments had two or three replicate reactors per sampling, and were sampled in both
Late Logarithmic (LL) and Early Stationary (ES) growth phases. (D). Also in (D), the 0.2*
denotes the serially transferred 0.2% O_2 experiment (procedure detailed in the methods).

Figure 3: Error! Reference source not found.

Figure 3. The doubling time (A,B,C,D) or ring index (E,F,G,H) for each set of conditions. Each
point represents a single replicate. The RI in fed-batch experiments (H) was determined both in
late-log phase and early stationary phase and are separated by color coding, with blue and pink
corresponding to Late Log and Early Stationary respectively. At 0.2% O_2, the serial transfer
experiment is denoted as solid circles for both RI and doubling time
399

Figure 4:

Figure 4. Ring index vs. doubling time. All biological replicates are shown, with experiments
differentiated by shape in each condition. Batch experiments (A-C) only varied a single
parameter relative to standard conditions at 70°C, pH 3, 200 RPM. In gas-fed batch experiments
(D), lipid samples taken for GDGT analysis during each of the two sampled growth phases are
identified by color, and 0.2* represents the serially transferred 0.2% O_2 experiment. Data from
constant-rate experiments performed with the same strain (E) are provided for comparison (data
from Zhou et al. 2019). Linear regressions for each experiment have grey shaded regions
showing 95% confidence interval (A: slope = -0.10 ± 0.01, p < 0.0001, R^2 =0.64; B: slope
= -0.147 ± 0.03, p < 0.001, R^2 =0.62; C: slope = -0.011 ± 0.0017, p < 0.0001, R^2 =0.53; D: slope
= 0.120 ± 0.0015, p < 0.0001, R^2 =0.68; E: slope = 0.031 ± 0.0031, p < 0.0001, R^2 =0.82).

413
Figure   5:

Figure 5. Clustering of GDGT composition in batch experiments color coded by change in
growth condition. Experiments were performed at 70°C, pH=3, 200 RPM, with a change in only
the individual condition described in the legend. NMDS is based on abundance of hydrolyzed
core GDGT lipids using Bray Curtis dissimilarity (stress = 0.043).

Figure 6:

422
Figure 6. Clustering of GDGT composition in fed-batch and chemostat experiments. Clustering
based on NMDS of abundance of hydrolyzed core GDGT lipids using Bray Curtis dissimilarity.
Stress for each NMDS is reported in the top right of a given panel. Gas-fed batch experiments
are arranged by symbol for experimental type and color for growth phase of collected sample
(A). Data in (B) are from Zhou et al. (2019) and show effects of different doubling times as
controlled by chemostat (B).

Figure S1

Figure S1. Ring Index vs. Doubling Time (GDGT 0-4, Cren and Cren-regioisomer treated as
GDGT-5) in *Nitrosopumulis maritimus* based on growth phase experiments in Elling et al., 2014.
Calculated growth rates are slightly different depending on the phase, but do not exhibit a large
spread. Early growth may show the highest rate due to the few measurements that comprise
that data (slope = -0.027 ± 0.017, *p* = 0.25, R^2 =0.34).

Figure S2

438
439
440   **Figure S2**. Ring Index vs. Growth Rate(GDGT 0-8) in *Picrophilus torridus*, based on work of
441   Feyhl-Buska 2016. Temperature experiments were performed at a single pH and pH
442   experiments were performed isothermally (A: slope = -0.05 ± 0.03, p = 0.19, R^2 = 0.23; B: slope
443   = 0.07 ± 0.05, p = 0.19, R^2 = 0.11).

444   **Table S1: GDGT compositions across all experiment conditions and sampling times**

Temp	RPM	pH	O2	Phase	GDGT-0	GDGT-1	GDGT-2	GDGT-3	GDGT-3 iso	GDGT-4	GDGT-4 iso	GDGT-5	GDGT-5 iso	GDGT-6	GDGT-7	GDGT-8	RI
70	200	2	Atmospheric	ES	1.17	5.14	22.65	19.44	7.31	7.88	14.01	11.37	4.27	6.72	0.05	0.00	3.32
70	200	2	Atmospheric	ES	1.20	5.20	23.44	19.77	7.41	7.40	14.55	10.71	4.15	6.14	0.04	0.00	3.32
70	200	2	Atmospheric	ES	1.42	5.94	23.39	19.09	6.85	8.28	13.69	11.18	3.88	6.23	0.04	0.00	3.31
70	200	2	Atmospheric	ES	1.09	4.65	20.59	19.27	6.24	8.85	14.45	13.30	4.02	7.49	0.05	0.00	3.47
70	200	2	Atmospheric	ES	1.08	4.83	23.31	20.05	6.79	8.49	13.49	11.74	3.98	6.20	0.04	0.00	3.35
65	200	3	Atmospheric	ES	0.35	1.42	9.43	21.62	0.00	62.25	0.00	4.78	0.00	0.15	0.01	0.00	3.62
65	200	3	Atmospheric	ES	0.25	1.30	8.19	18.53	0.00	62.38	0.00	9.03	0.00	0.31	0.00	0.00	3.59
65	200	3	Atmospheric	ES	0.32	1.62	11.19	23.53	0.00	56.42	0.00	6.68	0.00	0.23	0.00	0.00	3.70
65	200	3	Atmospheric	ES	0.29	1.43	10.08	22.13	0.00	58.26	0.00	7.55	0.00	0.25	0.00	0.00	3.55
65	200	3	Atmospheric	ES	0.27	1.46	10.50	22.38	0.00	57.16	0.00	7.96	0.00	0.27	0.00	0.00	3.60
70	200	3	Atmospheric	ES	0.29	0.88	5.88	17.15	0.00	55.21	2.76	16.67	0.00	1.14	0.01	0.00	3.86
70	200	3	Atmospheric	ES	0.28	1.24	8.34	17.43	0.00	54.17	3.13	15.30	0.00	0.00	0.11	0.00	3.76
70	200	3	Atmospheric	ES	0.39	1.67	10.63	22.22	0.00	44.72	3.89	15.23	0.00	1.23	0.01	0.00	3.66
70	200	3	Atmospheric	ES	0.30	1.24	9.01	21.68	0.00	51.19	2.85	13.61	0.00	0.11	0.00	0.00	3.68
70	200	3	Atmospheric	ES	0.32	1.45	9.68	20.80	0.00	50.07	2.16	15.47	0.00	0.00	0.05	0.00	3.69
75	200	3	Atmospheric	ES	0.34	1.28	7.92	11.20	1.10	33.09	6.47	30.52	0.84	6.84	0.35	0.05	4.14
75	200	3	Atmospheric	ES	0.28	1.04	6.73	11.54	1.16	32.97	4.95	30.90	1.62	8.35	0.40	0.06	4.21
75	200	3	Atmospheric	ES	0.32	1.26	7.19	12.19	0.00	32.29	5.56	32.35	0.80	7.61	0.38	0.06	4.18
75	200	3	Atmospheric	ES	0.51	2.30	16.76	23.93	3.19	20.99	10.49	17.26	0.97	3.60	0.00	0.00	3.51
75	200	3	Atmospheric	ES	0.30	1.11	6.71	12.65	0.00	33.73	6.86	30.84	0.81	6.52	0.43	0.06	4.16
80	200	3	Atmospheric	ES	0.38	1.18	6.78	9.40	1.73	15.84	7.19	36.55	3.52	17.16	0.25	0.04	4.50
80	200	3	Atmospheric	ES	0.38	0.98	5.88	7.78	1.48	15.86	7.37	37.08	3.66	19.48	0.01	0.03	4.60
80	200	3	Atmospheric	ES	0.39	1.05	5.73	8.79	1.43	14.40	6.74	36.05	4.41	20.71	0.25	0.05	4.61
80	200	3	Atmospheric	ES	0.48	1.30	6.81	10.79	1.64	14.72	9.26	29.98	5.27	19.74	0.00	0.00	4.47
80	200	3	Atmospheric	ES	0.51	2.91	0.43	11.14	1.72	21.94	8.10	34.31	3.43	15.40	0.09	0.01	4.67
70	200	4	Atmospheric	ES	0.98	4.23	24.57	32.16	0.00	35.74	0.00	2.28	0.00	0.05	0.00	0.00	3.08
70	200	4	Atmospheric	ES	1.07	4.65	26.94	32.58	0.00	33.01	0.00	1.71	0.00	0.04	0.00	0.00	2.98
70	200	4	Atmospheric	ES	0.87	3.72	23.92	31.90	0.00	37.60	0.00	1.94	0.00	0.05	0.00	0.00	3.08
70	200	4	Atmospheric	ES	1.05	4.68	26.64	32.33	0.00	33.28	0.00	2.00	0.00	0.04	0.00	0.00	3.04
70	200	4	Atmospheric	ES	0.85	3.87	23.53	32.38	0.00	37.06	0.00	2.27	0.00	0.04	0.00	0.00	2.97

70	0	3	Atmospheric	ES	5.59	16.99	31.53	25.16	1.16	15.73	1.75	1.95	0.04	0.10	0.00	0.00	2.39
70	0	3	Atmospheric	ES	4.31	14.17	31.30	25.07	1.66	17.34	2.34	3.55	0.07	0.20	0.00	0.00	2.55
70	0	3	Atmospheric	ES	5.44	16.63	33.73	24.14	1.36	14.98	1.66	1.95	0.04	0.09	0.00	0.00	2.38
70	0	3	Atmospheric	ES	4.45	15.00	33.16	25.01	1.35	16.20	1.87	2.74	0.05	0.17	0.00	0.00	2.48
70	0	3	Atmospheric	ES	3.61	12.95	32.80	26.78	1.42	16.98	2.14	3.07	0.07	0.19	0.00	0.00	2.56
70	50	3	Atmospheric	ES	8.27	21.98	35.25	22.62	1.19	9.15	0.90	0.59	0.02	0.03	0.00	0.00	2.07
70	50	3	Atmospheric	ES	9.24	23.82	35.76	20.93	1.09	7.74	0.86	0.52	0.02	0.03	0.00	0.00	1.99
70	50	3	Atmospheric	ES	10.42	24.40	36.04	19.87	1.11	6.84	0.80	0.48	0.02	0.03	0.00	0.00	1.93
70	50	3	Atmospheric	ES	8.27	22.62	36.60	21.53	1.15	8.32	0.87	0.58	0.02	0.03	0.00	0.00	2.04
70	50	3	Atmospheric	ES	9.49	23.22	35.71	20.89	1.09	8.13	0.87	0.56	0.02	0.03	0.00	0.00	2.00
70	61	3	Atmospheric	ES	6.87	21.29	36.88	22.47	1.12	9.92	0.81	0.60	0.01	0.02	0.00	0.00	2.12
70	61	3	Atmospheric	ES	7.24	21.93	37.33	22.30	1.06	8.82	0.77	0.52	0.01	0.02	0.00	0.00	2.08
70	61	3	Atmospheric	ES	7.47	22.71	37.24	21.84	1.15	8.13	0.85	0.55	0.02	0.03	0.00	0.00	2.05
70	61	3	Atmospheric	ES	8.66	23.23	37.85	20.93	0.55	7.55	0.72	0.45	0.01	0.02	0.00	0.00	1.99
70	61	3	Atmospheric	ES	7.07	20.74	37.37	22.77	0.69	9.65	0.97	0.69	0.02	0.03	0.00	0.00	2.12
70	75	3	Atmospheric	ES	4.65	15.80	35.48	25.77	0.97	14.15	1.37	1.71	0.03	0.07	0.00	0.00	2.38
70	75	3	Atmospheric	ES	4.48	15.62	35.95	25.73	0.88	14.56	1.26	1.43	0.02	0.06	0.00	0.00	2.38
70	75	3	Atmospheric	ES	4.48	15.82	37.55	25.17	0.80	13.78	1.11	1.23	0.02	0.05	0.00	0.00	2.35
70	75	3	Atmospheric	ES	4.02	13.87	35.45	27.28	0.94	15.31	1.43	1.60	0.03	0.07	0.00	0.00	2.45
70	75	3	Atmospheric	ES	4.99	17.71	38.96	24.47	0.79	11.17	0.97	0.90	0.01	0.04	0.00	0.00	2.25
70	97	3	Atmospheric	ES	3.41	12.03	39.65	27.11	2.16	10.42	2.81	2.20	0.08	0.14	0.00	0.00	2.44
70	97	3	Atmospheric	ES	3.14	12.31	40.38	26.67	2.61	8.71	3.64	2.28	0.09	0.17	0.00	0.00	2.43
70	97	3	Atmospheric	ES	3.44	12.26	39.59	26.68	2.38	9.48	3.39	2.52	0.09	0.18	0.00	0.00	2.44
70	97	3	Atmospheric	ES	3.54	12.17	40.11	26.74	2.16	9.62	3.09	2.33	0.08	0.16	0.00	0.00	2.43
70	97	3	Atmospheric	ES	3.20	13.09	38.37	27.22	2.11	10.51	2.95	2.33	0.08	0.15	0.00	0.00	2.45
70	125	3	Atmospheric	ES	1.49	6.79	30.21	29.63	2.85	14.82	6.52	6.83	0.18	0.69	0.00	0.00	2.89
70	125	3	Atmospheric	ES	1.46	5.97	28.73	30.29	3.01	14.33	7.18	7.94	0.24	0.83	0.00	0.00	2.95
70	125	3	Atmospheric	ES	1.23	5.57	28.13	30.60	2.94	14.45	7.39	8.52	0.27	0.90	0.00	0.00	2.99
70	125	3	Atmospheric	ES	1.42	5.78	29.67	30.70	2.81	14.25	6.70	7.62	0.23	0.83	0.00	0.00	2.94
70	125	3	Atmospheric	ES	1.42	6.15	29.80	30.59	2.78	14.41	6.57	7.31	0.22	0.74	0.00	0.00	2.92
70	200	3	Atmospheric	ES	0.72	3.94	26.50	33.67	1.86	22.81	3.96	6.14	0.09	0.31	0.00	0.00	3.04
70	200	3	Atmospheric	ES	1.12	5.43	28.37	32.97	1.84	19.90	4.16	5.81	0.08	0.32	0.00	0.00	2.94
70	200	3	Atmospheric	ES	0.74	3.34	23.37	32.52	1.66	24.86	4.70	8.22	0.10	0.48	0.00	0.00	3.15

70	200	3	Atmospheric	ES	0.76	3.65	24.80	33.96	1.47	24.28	4.09	6.57	0.08	0.34	0.00	0.00	3.08
70	200	3	Atmospheric	ES	0.84	4.04	26.51	33.72	1.96	21.85	4.29	6.31	0.09	0.37	0.00	0.00	3.03
70	300	3	Atmospheric	ES	1.00	6.04	33.31	33.81	0.99	20.83	1.74	2.20	0.02	0.06	0.00	0.00	2.79
70	300	3	Atmospheric	ES	0.93	5.78	32.38	32.46	1.03	23.16	1.80	2.39	0.02	0.07	0.00	0.00	2.83
70	300	3	Atmospheric	ES	1.12	6.64	33.12	32.91	0.94	21.33	1.75	2.11	0.02	0.06	0.00	0.00	2.78
70	300	3	Atmospheric	ES	1.11	6.84	32.93	33.45	0.91	21.22	1.59	1.89	0.02	0.05	0.00	0.00	2.77
70	300	3	Atmospheric	ES	1.22	6.45	33.47	33.15	0.99	20.71	1.74	2.19	0.02	0.06	0.00	0.00	2.77
70	NA	3	0.2	ES	1.46	4.14	12.21	26.77	1.68	42.60	2.74	8.02	0.06	0.32	0.00	0.00	3.38
70	NA	3	0.2	ES	1.59	4.58	12.19	23.99	3.87	45.87	1.37	6.31	0.04	0.20	0.00	0.00	3.34
70	NA	3	0.2	ES	1.10	3.10	10.27	22.84	4.12	47.89	2.38	7.91	0.08	0.30	0.00	0.00	3.47
70	NA	3	0.2	LL	1.83	5.32	15.77	31.08	1.96	37.60	1.57	4.61	0.08	0.19	0.00	0.00	3.17
70	NA	3	0.2	LL	2.39	6.81	17.46	27.46	2.64	37.78	1.32	3.96	0.06	0.15	0.00	0.00	3.09
70	NA	3	0.2	LL	1.72	5.25	16.18	29.23	2.83	38.02	2.20	4.29	0.09	0.19	0.00	0.00	3.18
70	NA	3	0.2*	ES	2.54	6.87	16.42	23.70	31.34	3.27	3.89	11.18	0.16	0.64	0.00	0.00	2.94
70	NA	3	0.2*	ES	3.29	8.78	20.57	26.96	22.93	4.82	4.16	7.98	0.13	0.37	0.00	0.00	2.78
70	NA	3	0.2*	LL	3.76	9.24	22.09	29.82	25.00	1.84	2.68	5.18	0.08	0.30	0.00	0.00	2.64
70	NA	3	0.2*	LL	4.46	12.55	27.32	28.45	19.44	1.56	2.41	3.58	0.05	0.19	0.00	0.00	2.46
70	NA	3	0.5	ES	4.87	15.97	35.74	25.44	1.57	13.49	1.47	1.35	0.04	0.07	0.00	0.00	2.36
70	NA	3	0.5	ES	3.42	12.54	32.88	27.73	1.39	18.83	1.45	1.64	0.04	0.07	0.00	0.00	2.56
70	NA	3	0.5	ES	4.50	15.32	35.92	26.29	0.91	14.49	1.31	1.16	0.03	0.06	0.00	0.00	2.38
70	NA	3	0.5	LL	4.01	11.78	27.72	31.72	0.65	21.99	0.73	1.32	0.02	0.07	0.00	0.00	2.62
70	NA	3	0.5	LL	3.12	10.54	27.63	31.06	0.92	24.58	0.77	1.29	0.02	0.07	0.00	0.00	2.70
70	NA	3	0.5	LL	4.29	14.17	31.27	29.69	1.06	17.49	0.92	1.03	0.04	0.06	0.00	0.00	2.48
70	NA	3	1	ES	4.39	13.56	31.55	28.59	1.35	14.74	2.71	2.87	0.09	0.16	0.00	0.00	2.52
70	NA	3	1	ES	5.58	16.64	35.64	25.02	1.54	12.09	1.60	1.75	0.06	0.09	0.00	0.00	2.32
70	NA	3	1	ES	5.90	17.87	39.24	22.47	1.98	9.21	1.93	1.28	0.05	0.06	0.00	0.00	2.21
70	NA	3	1	LL	5.99	17.26	32.81	25.75	1.83	12.08	2.32	1.79	0.07	0.10	0.00	0.00	2.33
70	NA	3	1	LL	5.96	17.41	33.72	25.50	1.28	12.54	1.93	1.53	0.05	0.07	0.00	0.00	2.31
70	NA	3	1	LL	6.04	18.18	37.27	24.03	1.67	9.59	1.83	1.27	0.05	0.06	0.00	0.00	2.22
70	NA	3	2	ES	10.55	25.40	39.17	18.08	1.05	4.69	0.72	0.31	0.01	0.01	0.00	0.00	1.84
70	NA	3	2	ES	10.02	23.81	36.71	19.73	1.49	6.46	1.17	0.57	0.02	0.02	0.00	0.00	1.94
70	NA	3	2	ES	8.36	23.14	39.46	19.47	1.40	6.57	1.02	0.54	0.02	0.02	0.00	0.00	1.98
70	NA	3	2	LL	14.86	30.32	37.55	13.07	0.25	3.37	0.31	0.24	0.01	0.02	0.00	0.00	1.61

70	NA	3	2	LL	17.76	31.65	34.97	12.23	0.32	2.70	0.24	0.12	0.00	0.01	0.00	0.00	1.52
70	NA	3	2	LL	16.10	30.65	36.95	12.99	0.35	2.59	0.26	0.11	0.00	0.01	0.00	0.00	1.57
70	NA	3	20	ES	4.77	18.70	40.66	25.81	0.61	8.50	0.61	0.32	0.01	0.01	0.00	0.00	2.17
70	NA	3	20	ES	5.01	19.22	37.79	26.21	0.74	9.83	0.74	0.42	0.01	0.01	0.00	0.00	2.20
70	NA	3	20	ES	3.89	16.56	37.41	27.59	0.90	11.81	1.05	0.77	0.01	0.02	0.00	0.00	2.32
70	NA	3	20	LL	8.88	24.84	39.95	20.53	0.34	4.99	0.30	0.15	0.00	0.01	0.00	0.00	1.89
70	NA	3	20	LL	10.64	24.69	37.17	21.26	0.38	5.38	0.32	0.14	0.00	0.01	0.00	0.00	1.88
70	NA	3	20	LL	8.16	22.50	38.00	22.94	0.48	7.12	0.53	0.25	0.00	0.01	0.00	0.00	2.01

445
446
447
448   GDGT relative abundances. In the O_2 column, "Atmospheric" is used to describe experiments without gas sparge, and 0.2* is used to describe the
449   serially transferred 0.2% O_2 gas-fed batch experiment. ES and LL in the Phase column describe Early Stationary and Late Log phase sampling
450   respectively.