Shifting Baselines: Physiological legacies contribute to the response of reef coral to

frequent heat waves

Christopher B. Wall[1,2*], Contessa A. Ricci[3], Alexandra D. Wen[1,4], Bren E. Ledbetter[3], Delania E.

Klinger[3], Laura D. Mydlarz[3], Ruth D. Gates[1], Hollie M. Putnam[5]

[1]University of Hawaiʻi at Mānoa, Hawaiʻi Institute of Marine Biology, PO Box 1346, Kāneʻohe,

HI 96744, USA

[2]Pacific Biosciences Research Center, University of Hawaiʻi at Mānoa, 1933 East West Road,

Honolulu, HI 96816, USA

[3]University of Texas at Arlington, Department of Biology, Arlington, TX 76019, USA

[4]University of Miami, Rosenstiel School of Marine and Atmospheric Science, Miami, FL 33149,

USA

[5]University of Rhode Island, Department of Biological Sciences, Kingston, RI USA

*corresponding author: chris.wall@hawaii.edu

Abstract

1. Global climate change is altering coral reef ecosystems. Notably, marine heat waves are producing widespread coral bleaching events that are increasing in frequency, with projections for annual bleaching events on reefs worldwide by mid-century.

2. The response of corals to elevated seawater temperatures can be modulated by abiotic factors at site of origin and dominant endosymbiont type, which can result in a shift in multiple coral traits and drive physiological legacy effects that influence the trajectory of reef corals under subsequent thermal stress events. It is critical, therefore, to evaluate the potential for shifting physiological and cellular baselines driven by these factors in *in situ* bleaching (and recovery) events. Here, we use the back-to-back regional bleaching events of 2014 and 2015 in the Hawaiian Islands and subsequent recovery periods to test the hypothesis that coral multivariate trait space (here termed physiotype, sensu (Van Straalen, 2003) shift in multiple bleaching events, modulated by both environmental histories and symbiotic partnerships (Symbiodiniaceae).

3. Despite fewer degree heating weeks in the first-bleaching event relative to the second (7 *vs*. 10), bleaching severity in a dominant reef building coral on Hawaiian reefs, *Montipora capitata,* was greater (~70% *vs*. 50% bleached cover) and differences due to environmental history (reef site) were more pronounced. Melanin, an immune cytotoxic response, provided an initial defense during the first event, potentially priming antioxidant activity, which peaked in the second-bleaching event (i.e., a legacy effect). While magnitude of bleaching differed, immune response patterns were shared among corals harboring heat-sensitive and heat-tolerant Symbiodiniaceae. This supports a pattern of increased constitutive immunity in corals resulting from repeat bleaching

events, with greater specialized enzymes (catalase, peroxidase, superoxide dismutase)

and attenuated melanin synthesis.

4. This study demonstrates bleaching events have implications for reef corals beyond

shaping their ecological assemblages. These events can change the magnitude and/or

identity of response variables contributing to physiotype, thus generating physiological

legacies carried over into the future. Quantifying baseline coral physiotypes and tracking

their shifts will be critical to understanding and forecasting the effects of increased

bleaching frequency on coral biology and ecology in the Anthropocene.

1. Introduction

Environmental change at local and global scales is degrading coral reef ecosystems. Notably,

long-term trends in ocean warming and punctuated marine heat waves are destabilizing the

symbiosis between corals and their endosymbiont algae (Symbiodiniaceae) and increasing coral

bleaching events (Hughes et al., 2018a). The cumulative impacts of these episodic heat waves

not only shift ecological and functional baselines (Hughes et al., 2019; McWilliam, Pratchett,

Hoogenboom, & Hughes, 2020), but may also have legacy effects on coral biology, such that

responses to bleaching events can be modulated by prior exposure (e.g., Brown, Dunne,

Goodson, & Douglas, 2000). It is growing increasingly clear that susceptibility to bleaching in

reef corals is based on the interaction of environmental histories (Brown et al., 2000; Safaie et

al., 2018) and holobiont traits, including endosymbiont communities and genetics and

physiology of the coral host (Barshis et al., 2013; Palmer, Bythell, & Willis, 2010; Palumbi,

Barshis, Traylor-Knowles, & Bay, 2014; Sampayo, Ridgway, Bongaerts, & Hoegh-Guldberg,

2008).

Thermotolerant Symbiodiniaceae can confer bleaching resistance in some corals (Sampayo et al.,

2008), and the Hawaiian coral *Montipora capitata* exhibits reduced bleaching when dominated

by *Durusdinium* as opposed to *Cladocopium* Symbiodiniaceae (Cunning, Ritson-Williams, &

Gates, 2016). However, corals are metaorganisms, and host properties such as constitutive

immunity (Palmer, 2018a) and antioxidant capacity (Barshis et al., 2013) are also central to

maintaining cellular homeostasis and preventing bleaching mortality (Palmer et al., 2010). As

such, immunological processes are implicated as targets for natural selection and are integral to

the future of reef-building corals in the face of climate change (Mydlarz, McGinty, & Harvell,

2010; Palmer, 2018a; Pinzón, Beach-Letendre, Weil, & Mydlarz, 2014). The nexus of these

influential properties of coral stress responses – symbiont community and host immunity – and

their role in repetitive natural bleaching and recovery events remains to be fully understood,

especially with respect to the role of environmental history (Palumbi et al., 2014; Safaie et al.,

2018). As global climate change increases the frequency and severity of coral bleaching events

(Hughes et al., 2018), understanding the symbiotic, immune, and antioxidant responses of corals

will be central to understanding the cumulative and latent effects of bleaching on the function of

reef corals through time and enhancing our capacity to project community responses to thermal

stress events.

The widely documented recent major bleaching of the Great Barrier Reef (2016-2017) reduced

coral cover by >50% (Stuart-Smith, Brown, Ceccarelli, & Edgar, 2018), altered coral (Hughes et

al., 2018b) and fish (Robinson, Wilson, Jennings, & Graham, 2019) assemblages, and disturbed

the stock-recruitment capacity of reef corals (Hughes et al., 2019). As climate change continues

to intensify, it is now critical to track shifts in the biology and function of corals through

examination of multivariate trait-space, defined as 'physiotypes' by Van Straalen (2003; e.g.,

Figure 1a), the term which we use throughout this manuscript. Trait-space approaches have

recently been applied to track changes in reef coral communities and their representative trait

diversity following bleaching (Hughes et al., 2018b; McWilliam et al., 2020), and the generation

of physiotype time series data will provide a critical match to further understand and interpret the

growing collection of coral reef ecological time series datasets (De'ath, Fabricius, Sweatman, &

Puotinen, 2012; Edmunds et al., 2014; Edmunds et al., 2014; McClanahan, Ateweberhan, &

Omukoto, 2007). Together, these timeseries will allow us to integrate across cellular, ecological

and evolutionary scales and improve our mechanistic understanding of these scale linkages.

Importantly, tracking of multiple, interactive coral physiological traits provides the opportunity

to reveal the potential for beneficial acclimatization or negative legacy effects that may underlie

shifts in physiotype (e.g., Figure 1a) and testable mechanistic hypotheses underpinning resistance

or susceptibility to repeated stress (Figure 1b).

This integrative multivariate approach to tracking coral performance was utilized to test specific

hypotheses regarding targeted response variables and the linkage between environmentally-

driven physiological, symbiotic, and cellular (i.e., antioxidants, immunity) legacies as

mechanisms underpinning bleaching outcomes (Palmer et al., 2010; Palmer & Traylor-Knowles,

2012; Venn, Loram, & Douglas, 2008). Both the host and symbiont have enzymatic defenses to

mitigate oxidative stress (e.g., peroxidase, catalase, superoxide dismutase) (Venn et al., 2008)

caused by host (e.g., heat-damaged mitochondrial membranes, Dunn et al., 2012) and symbiont

(e.g., damage to the D1 protein of photosystem II (PSII)) (Jones, Hoegh-Guldberg, Larkum, &

Schreiber, 1998; Lesser, 1997) sources, ultimately leading to apoptosis and dysbiosis when

severe enough (Weis, 2008) (Figure 1b). Thermal stress also triggers host immune responses

(Mydlarz, Couch, Weil, Smith, & Harvell, 2009; Mydlarz et al., 2010; Pinzón et al., 2015), and

higher constitutive immunity (i.e., immune activity necessary for cellular homeostasis) reduces

coral thermal sensitivity and disease susceptibility (Palmer et al., 2010). In addition, the

melanin-synthesis pathway is an important component of host immunity that is active in wound

healing and pathogen invasion while also serving as a photoprotectant that may mitigate

bleaching stress (Mydlarz & Palmer, 2011; Palmer et al., 2010) (Figure 1b). However, corals

may mount different cellular mechanisms to combat bleaching due to energetic requirements

(Fuess, Mann, Jinks, Brinkhuis, & Mydlarz, 2018; Palmer, 2018b; Pinzón et al., 2015), stress

frequency (i.e, acute, chronic, repeated stress) (Ainsworth et al., 2016; Schoepf et al., 2015), or

as a function of histories of biotic and abiotic challenges that modulate constitutive immunity

(Mydlarz et al., 2009; Palmer, 2018b; Wall et al., 2018). For instance, there is evidence corals

may rely on different antioxidant and immunity mechanisms based on their environmental

histories, as lower melanin synthesis and higher antioxidant activity was observed in heat-

stressed corals from a reef with high pCO_2-variability compared to low pCO_2-variability (Wall et

al., 2018). Thus, pairing the measurement of bleaching metrics like cell density and chlorophyll

concentrations with immune and antioxidants provides a powerful and tractable approach within

a multivariate integrative framework.

The Hawaiian Islands, once thought to be a bleaching refuge for corals (Jokiel & Brown, 2004),

experienced severe bleaching in 2014 and 2015 across the populated Main and remote

Northwestern Hawaiian Islands (Bahr, Rodgers, & Jokiel, 2017; Couch et al., 2017). In

Kāneʻohe Bay, Oʻahu, Hawaiʻi, 7.1 degree heating weeks (DHW) were observed in 2014 and

10.2 DHW in 2015, beginning in August and July, respectively (*see* Methods), resulting in 45 ±

2% (mean ± SE) bleaching in the first event and 30 ± 4% bleaching in the second (Bahr et al.,

2017). An ecologically dominant reef coral, *Montipora capitata* (Dana 1846), showed

significant bleaching in both events (Figure 2a), with differences in bleaching responses

observed at the site and event level.

Using the opportunity presented by these natural bleaching events (Figure 2), we tracked coral

percent bleaching and physiotypes using a multivariate trait-space approach. We quantified the

response of *M. capitata* in Kāneʻohe Bay to the back-to-back bleaching events of 2014 and 2015

(i.e., bleaching in October 2014, recovery in February 2015, bleaching in October 2015, and

recovery in February 2016) at two Kāneʻohe Bay reefs (Lilipuna and Reef 14) with contrasting

environmental histories relating to seawater residence (10 – 20 d *vs*. >30 d), pCO_2 variability (ca.

300 *vs*. 600 μatm pCO_2 diel flux), and proximity to shore (Drupp, De Carlo, Mackenzie,

Bienfang, & Sabine, 2011; Drupp et al., 2013; Lowe, Falter, Monismith, & Atkinson, 2009).

We predicted that: (H1) immunity and antioxidant contributions to bleaching responses will

influenced by site environmental histories and symbiont communities such that greater bleaching

prevalence and attenuated immune responses will be observed in corals from the high variable-

pCO_2 site (Reef 14) and those association with thermally sensitive Symbiodiniaceae. We also

expected (H2) immunity and antioxidant activity would not occur with the same magnitude or

trajectory after repeated bleaching events due to legacy effects shaping acclimatory or stress

response pathways over time. Considering the wide-ranging functions of melanin in host

immunity (Palmer et al., 2008, 2010), we expected melanin synthesis to act as an acute and

broad-spectrum defense in physiologically stressed corals, with antioxidants having a more

specialized response after chronic or repeat bleaching stress. Finally, while the thermal tolerance

of symbiont communities is vital to coral bleaching sensitivity, evidence for differential

immunity or antioxidant capacities in coral holobionts associated with thermally tolerant

symbiont species is lacking. Given that the coral *M. capitata* is known to be dominated by two

genetically and physiologically contrasting Symbiodiniaceae, *Cladocopium* sp. and *Durusdinium*

glynnii (Cunning et al., 2016; Wall, Kaluhiokalani, Popp, Donahue, & Gates, 2020), we

predicted (H3) that holobionts harboring heat-tolerant *D. glynnii* would show higher immune and

antioxidant responses linked to greater bleaching resistance. To test these hypotheses, we

measured bleaching severity and recovery at each location using benthic surveys and measured

coral traits in coral fragments, including: physiological measurements of cellular bleaching

magnitude (symbiont density, areal- and cell-specific chlorophyll *a*, holobiont total protein and

total biomass), and completed enzymatic assays for mechanisms contributing to bleaching

resistance through coral host antioxidant capacity (peroxidase, catalase, superoxide dismutase),

and the host immune response of the melanin cascade (prophenoloxidase, melanin). This

approach allows for a more holistic description of coral physiotype that we tracked through two

bleaching and recovery periods.

2. Materials and Methods

2.1 Site description

Coral bleaching and recovery at two reef systems: a fringing reef (hereafter, 'Lilipuna'

[21°25′36.8″N, 157°47′24.0″W]) in southern Kāneʻohe Bay adjacent to the Hawaiʻi Institute of

Marine Biology (HIMB) on Moku o Loʻe, and an inshore patch reef (hereafter, 'Reef 14'

[21°27′08.6″N, 157°48′04.7″W]) in central Kāneʻohe Bay. These reefs sites were chosen due to

their unique environmental history of seawater pCO_2 and hydrodynamics (Drupp et al., 2013;

Wall et al., 2018). Seawater pCO_2 adjacent to the both locations (fringing reef at Lilipuna and

the patch reef Reef 14) is comparable (ca. 450 μatm), however, diel pCO_2 flux is significantly

higher (196 – 976 μatm pCO_2) in central Kāneʻohe Bay near Reef 14 relative to the Bay's

southern basin proximate to Lilipuna (225 – 671 μatm) (P. Drupp et al., 2011; Drupp et al.,

2013).

2.2 Environmental monitoring

Temperature (Hobo pendants, ± 0.53 °C accuracy, 0.14 °C resolution, Onset Computer Corp.,

Bourne, MA) and photosynthetic active radiation (PAR) loggers (Odyssey, Dataflow Systems

Limited, Christchurch, New Zealand) were placed at Lilipuna and Reef 14 at a depth of 1 m.

Loggers were periodically removed from the reef, cleaned, and replaced monthly. Temperature

and PAR were recorded continuously at 15 min intervals from October 2014 – February 2016,

however, gaps in environmental data do exist as a consequence of instrument failure and loss.

Gaps in logger temperature data were supplemented with NOAA temperature data from the

Moku o Lo'e station at HIMB (NOAA, 2019). PAR loggers were calibrated against a LI-1400

quantum meter (Li-Cor, Lincoln, Nebraska, USA) attached to a cosine LI-192 underwater

quantum sensor. Temperature loggers were calibrated with a certified digital thermometer (5-

077-8, ± 0.05 °C accuracy, Control Company, USA) and cross-calibrated against each other for

standardization. Degree heating weeks (DHW) for the southern portion of Kāneʻohe Bay where

our corals were collected were calculated in R using *in situ* Moku o Loʻe temperature data

(NOAA, 2019), with the difference between mean half-week temperatures (i.e., mean hourly

temp over 3.5 d) and the maximum monthly mean temperature of 27.7 °C, (Jokiel & Brown,

2004)] to determine 'hotspots'. DHW were determined as the number of hotspots >1 across a

rolling 12 weeks window (i.e., 24 half-weeks) (NOAA, 2020). DHW for windward Oʻahu and

the Hawaiian Islands during the 2014 and 2015 bleaching events have been previously reported

(Bahr et al., 2017; Sale, Marko, Oliver, & Hunter, 2019). The purpose of our calculations are to

quantify the incurred heat stress using empirical calculation of DHW from temperature data

collected proximate to our two reef locations.

2.3 Benthic surveys and coral collections

Four sampling periods were identified as corresponding to a 'bleaching period' following the

point of maximum thermal stress (10 October 2014 and 12 October 2015) and a post-bleaching

'recovery period' approximately 4 months after peak seawater warming (11 February 2015 and

26 February 2016). In each time period, benthic surveys were conducted at each reef site using a

20 m transect and a line-point-intersect at 1 m intervals. Transects were positioned parallel to

natural contours of the reef, being the north-south axis of the fringing reef (Lilipuna) and the

east-west axis of patch reef (Reef 14). At each reef, transects ($n = 2$) were placed along the reef

crest (1 – 2 m), and benthic community cover was recorded at the species level for reef corals

(*Montipora capitata, Pocillopora* spp. [*P. acuta*, *P. damicornis*]*, Porites compressa*), and either

crustose coralline algae (CCA), macroalgae, or sand/bare/turf. For corals, bleaching state was

quantified categorically, being either non-bleached (i.e., appearing fully pigmented), or bleached

(i.e., exhibiting degrees of tissue paling/pigment variegation or being wholly white). *M. capitata*

cover was calculated as the proportion of total benthic cover, and *M. capitata* bleaching extent

was calculated as proportion of total *M. capitata* colonies bleached. The relative position of

transects at each reef was recorded to allow for repeat surveys within the same general location

at each reef across survey periods.

In each sampling period, coral branch tips (<4 cm length) were collected from forty *M. capitata*

coral colonies ($n = 1$ fragment colony^{-1}) along the reef crest at each reef site at a depth of ca. 1 m

(State of Hawai'i Department of Land and Natural Resources, Special Activity Permit 2015-17

and 2016-69). Immediately post collection, corals were snap-frozen in liquid nitrogen and

returned to HIMB and stored at -80 °C. While remaining frozen, each colony was split in half

along its longitudinal axis. One-half of each coral fragment was stored at HIMB (-80 °C) until

processing for physiological assays and qPCR. The corresponding fragment-halves were used

for immunological assays and were shipped to the University of Texas at Arlington using a dry-

shipper charged with liquid nitrogen.

2.4 DNA extraction and symbiont community analysis

Symbiodiniaceae DNA was extracted by adding an isolate of coral tissue (500 μl) to 500 ul DNA

buffer (0.4 M NaCl, 0.05 M EDTA) with 2% (w/v) sodium dodecyl sulfate, following a modified

CTAB-chloroform protocol (Cunning et al., 2016) (dx.doi.org/10.17504/protocols.io.dyq7vv).

To determine the dominant genera of Symbiodiniaceae hosted by each *M. capitata* fragment, a

quantitative PCR (qPCR) (Cunning et al., 2016) was used to amplify Symbiodiniaceae actin gene

loci, with the corresponding number of actin gene copies in the Symbiodiniaceae genera

Cladocopium and *Durusdinium* (namely, ITS2 types C31, with C17 and C21 and D1-4-6

[*Durusdinium glynnii* (Wham, Ning, & LaJeunesse, 2017)]) known to be numerically dominant

in Kāneʻohe Bay *M. capitata* (Cunning et al., 2016). Specificity of genus-level primers have

been previously validated using a combination of Symbiodiniaceae internal transcribed spacer

(ITS2) region of rDNA and actin gene sequencing (Cunning & Baker, 2013). Duplicate qPCR

reactions (10 μl) were run for each coral sample using a StepOnePlus platform (Applied

Biosystems) set to 40 cycles, a relative fluorescence (ΔR_n) threshold of 0.01, and internal cycle

baseline of 3 – 15. Symbiont genera detected in only one technical replicate were considered

absent. The relative abundance *of Cladocopium and Durusdinium* symbionts (i.e., C:D ratio) in

each sample was determined from the ratio of amplification threshold cycles (C_T) for

Cladocopium and *Durusdinium* (i.e., C_T^C, C_T^D) using the formula C:D = $2^{\wedge}(C_T^C - C_T^D)$, where

genus-specific C_T values are normalized according to gene locus copy number and fluorescence

intensity (Cunning et al., 2016). Coral colonies were determined to be *Cladocopium*- or

Durusdinium-dominated based on numerical abundance (>0.5 proportion) of each genus from

qPCR analysis (Innis, Cunning, Ritson-Williams, Wall, & Gates, 2018).

2.5 Physiological metrics

The extraction and processing of coral and symbiont tissues was performed following established

methods (summarized in Wall et al., 2018). Briefly, coral tissue was removed from the skeleton

using an airbrush filled with 0.2 μm filtered seawater, yielding ~10 – 30 ml of tissue slurry.

Extracted tissues were briefly homogenized and subsampled for the following physiological

metrics: symbiont cell densities, chlorophyll *a* concentrations, protein biomass, and the total

organic biomass determined from the ash-free dry weight (AFDW) of coral + algae tissues.

Coral tissue slurries were stored at -20 °C.

All physiological metrics were normalized to the surface area (cm^2) of coral skeleton using the

paraffin wax-dipping technique (Stimson & Kinzie, 1991). Symbiont cell counts were measured

by replicate cell counts ($n = 6 – 10$) on a haemocytometer, and expressed as symbiont cells cm^{-2}.

Chlorophyll *a* was quantified by concentrating algal cells through centrifugation ($3,000 \times g$ for 3

min) and extracting pigments in the algal pellet in 100% acetone for 36 h in darkness at -20 °C.

Spectrometric absorbance were measured ($\lambda = 630$ and 663 nm) using a 96-well quartz plate with

two technical replicates; chlorophyll concentrations quantified using the equations for

dinoflagellates (Jeffrey & Humphrey, 1975), normalized for path length and expressed as ug

chlorophyll *a* cm^{-2} and pg chlorophyll *a* symbiont $cell^{-1}$. Total protein concentration (soluble +

insoluble in the holobiont) was quantified using the Pierce BCA (bicinchoninic acid) Protein

Assay Kit (Pierce Biotechnology, Waltham, MA). Protein solubilization was achieved by adding

1 M NaOH and heating (90 °C) for 60 min, followed by the neutralizing to ca. pH 7.5 with 1 N

HCl. Protein was measured spectrophotometrically ($\lambda = 562$ nm) in a 96-well plate with three

technical replicates against a bovine serum albumin standard and expressed as mg protein cm^{-2}.

Total fraction of organic biomass was measured by drying a subsample of coral tissue at 60 °C

(48 h) in pre-burned aluminum pans followed by burning in a muffle furnace (450 °C) for 4 h.

The difference between the dried and burned masses is the AFDW and expressed as mg cm^{-2}.

2.6 Immunity and oxidative stress metrics

Immunology and oxidative stress metrics were determined using previously published protocols

for coral host tissues (Mydlarz et al., 2009; Mydlarz & Palmer, 2011; Palmer et al., 2011; Wall et

al., 2018). A 3 – 4 ml aliquot of coral tissue slurry was obtained by airbrushing with our coral

extraction buffer at pH 7.0 (100 mM TRIS buffer + 0.05 mM dithiothreitol). Tissue was

homogenized for 1 min on ice (hand held homogenizer, Powergen 125, Fisher Scientific,

Waltham, MA), and 1 ml of the resulting slurry was freeze-dried for 24 h (VirTis BTK freeze-

dryer, SP Scientific, Warminster, PA) and used for melanin concentration estimates. The

remaining slurry was centrifuged at 4 °C at 2,500 × *g* (Eppindorf 5810 R centrifuge, Hamburg,

Germany) for 5 min to remove cellular debris and most Symbiodiniaceae cells to achieve a host-

enriched cell-free extract. All colorimetric measurements were calculated using a Synergy 2

multi-Detection microplate reader with Gen5 software (BioTek, Winooski, VT, USA). All assays

were run in duplicate or triplicate on separate 96-well microtiter plates. Total protein

concentration of each coral cell-free extract was determined using the RED660 protein assay (G

Biosciences, Saint Louis, MO) with a bovine serum albumin standard curve.

To determine the concentration of melanin within each sample, the freeze-dried aliquots of

weighed and dried tissue were gently vortexed with 400 μl of 10 M sodium hydroxide (NaOH)

and left to extract for 48 h. Samples were bead-beaten with 1 mm glass beads and then vortexed

for 10 s and then centrifuged at 7,000 × g for 5min. For each sample, 65 μl of supernatant were

aliquoted in duplicate into a 96-well g-area microtiter plate (Greiner Bioone, Monroe, NC,

USA). The plates were read at an absorbance of 490 nm and the concentration of melanin was

determined using a standard curve (0 – 2 mg) of commercial melanin (Sigma-Aldrich, St. Louis,

MO) dissolved in 10 M NaOH for 48 h and treated the same as the samples on each microtiter

plate. Data presented are converted to mg melanin normalized to mg of tissue (Fuess et al.,

2018).

PPO activity was determined for each sample using duplicate 20 μl aliquots of coral cell-free

extract in clear 96-well microtiter plates, with 50 μl of 10 mM phosphate buffered saline at pH

7.0. To each well, 20 μl of trypsin (0.2 mg ml^{-1} concentration in deionized filtered water) was

added to activate PPO and the reaction was initiated by the addition of 20 μl of 25 mM L-1,3-

dihydroxyphenylalanine (L-DOPA; Sigma-Aldrich). The absorbance at 490 nm was measured

over 25 min and the change in absorbance during the linear portion of the reaction (typically 10 –

15 min) was normalized to mg protein and time for each sample (ΔAbs490 nm mg protein^{-1} min^{-}

^{1}) (Mydlarz & Palmer, 2011; Fuess et al., 2018).

Coral host oxidative stress was determined by measuring the scavenging activity of the coral

cell-free extracts to different substrates comparable to the antioxidants: peroxidase (POX),

catalase (CAT), and superoxide dismutase (SOD). Peroxidase activity (EC 1.11.1.7) was

determined for each sample using 10 μl aliquots of coral cell-free extract, in duplicate within 96-

well microtiter plates, diluted with 50 μl phosphate buffer (pH6.0). To each well, of 25 μl of 25

mM guaiacol in 10 mM PBS (pH 6.0) was added and the reaction was initiated with the addition

of 12.5 μl of 25 mM H_2O_2 high purity 30% hydrogen peroxide (H_2O_2; Sigma Aldrich). Guaiacol

is a phenolic compound with scavenging properties that is commonly used as a substrate for

peroxidase activity (Mydlarz & Harvell, 2007). The absorbance was read at 470 nm every min

for 15 min and the change over time was calculated for the linear part of the reaction (0 – 10

min) and normalized to mg protein in each sample the units for peroxidase activity are (ΔAbs470

nm mg protein^{-1} min^{-1}) (Mydlarz & Harvell, 2008).

Catalase activity (EC 1.11.1.6) was determined using the H_2O_2 depletion assay using 10 μl coral

cell free extract and 50 μl of 10 mM of pH 6.0 phosphate buffered saline (PBS) in duplicate

wells of a UV-transparent 96-well microtiter plate. The assay was activated by the addition of

10 μl 25 mM H_2O_2. The absorbance was measured at 240 nm over 15 min and the change in

absorbance over time (e.g., initial - final) was calculated for the linear part of the reaction

(typically 0 – 8 min) and converted to μmol H_2O_2 using a standard curve of known

concentrations, normalized per mg protein. CAT activity is presented as μmol H_2O_2 scavenged

mg protein^{-1} min^{-1} (Palmer et al., 2011).

Superoxide dismutase (EC 1.15.1.1) activity was calculated using the SOD Assay Determination

Kit-WST (Fluka, Switzerland) which includes Dojindo's highly water-soluble tetrazolium salt,

WST-1(2-(4- iodophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)- 2H-tetrazolium, monosodium

salt), that produces a water-soluble formazan dye with an absorbance peak at 450 nm upon

reduction with superoxide anion. 10 μL of extract was incubated with WST-1 and xanthine

oxidase, which produces superoxide anion. Inhibition of absorbance at 450 nm (scavenging of

the superoxide anion) was monitored in wells containing coral extracts and SOD standards and

compared to untreated samples. One SOD unit of activity (U) equates to 50% inhibition of

superoxide anion, and data are presented as SOD U mg protein^{-1} (Krueger et al., 2015).

2.7 Statistical analysis

Permutation analysis of variance (PERMANOVA) and non-metric multidimensional scaling

(NMDS) were performed using a balanced matrix of all physiology and immunity/antioxidant

responses (*n* = 10 responses) in the package *vegan* (Oksanen et al., 2019). PERMANOVA used

a scaled and centered matrix with Euclidean calculations of pairwise distances using *adonis2*.

The NMDS data matrix was double standardized using a Wisconsin and square root

transformation with a euclidean distance using *metaMDS*. Response variables showing

significant correlations (*p*<0.05) with NMDS axes were plotted as vectors using envfit command

in *vegan*. Coral 'physiotypes' were defined by convex hulls, with borders defined by the range

of points in the multivariate trait space (i.e., NMDS1 and NMDS2). Physiotypes were grouped

categorically by periods, or the interactions of Site, Symbiont, and Period. Physiotype centroids

used in trajectory plots were calculated as the mean of NMDS1 and NMD2 for each category.

Ecological benthic data (*M. capitata* total cover and bleached cover) were tested in a linear

model with Periods (two bleaching and two recovery events) and Sites (Lilipuna, Reef 14) as

fixed effects. Physiology, antioxidant, and immunity response variables were analyzed using a

linear model with Periods, Sites, and Symbiont community composition (*Cladocopium*- or

Durusdinium-dominated) as fixed effects. Normal distribution and equal variance assumptions

of ANOVA were examined by graphical representation of residuals and quantile:quantile plots.

Where assumptions were not met, Box-Cox tests were performed (Box & Cox, 1964) and data

transformations applied in the package *MASS* (Venable & Ripley, 2002). Analysis of variance

tables for linear models were generated using type-II sum or squares with multiple comparisons

using the package *emmeans* (Lenth, 2019). All analyses were performed in *R* version 3.6.1 (R

Core Team, 2019). All data and code to generate figures and perform analyses are archived and

openly available at Github (https://github.com/cbwall/Gates-Mydlarz-bleaching-

recovery/releases/tag/v4).

3. Results

Coral cover at the two sites ranged from (mean ± SD) 63 ± 11% to 93 ± 11% from 2014-2016.

Greater total bleached coral cover was observed in the first bleaching event (62 – 75 ± 9%)

relative to the second (43 – 55 ± 16%) (i.e., during thermal stress) (Figure 2). Mean *M. capitata*

cover was stable across time at Lilipuna (~40%) but was much more variable at Reef 14, falling

from 91 ± 4% in October 2014 to 51 ± 7% in February 2016. In the two bleaching events

(October 2014 and 2015) greater bleaching of *M. capitata* corals was observed at Reef 14,

compared to Lilipuna (*p*=0.020).

PERMANOVA results testing effects of Period (bleaching event), Site (Lilipuna *vs*. Reef 14) and

dominant Symbiodiniaceae species (*Cladocopium* sp. or *D. glynnii*) revealed significant

differences of all main effects (*p*<0.001) and the interaction of Period-by-Site (*p*<0.001) and

Period-by-Symbiont community ($p<0.001$) (Figure 3 and 4, Table 1). In each of the four

periods, coral physiotypes occupied unique positions in multidimensional trait space (Figure 2),

with the location and trajectories of coral physiotypes in each period principally being a function

of symbiont community, and to a lesser extent, site of collection (i.e., environmental history and

physiological legacy) (Figure 3, 4).

Physiotypes of bleached corals relative to recovered corals showed stronger separation in 2014-

2015, which was apparent at both reef sites (Figure 3, top row) and paralleled patterns of greater

bleaching prevalence in the first-bleaching event (Figure 2c). Similarly, site effects on coral

physiotypes were most pronounced in the first bleaching event (Table 1, Figure 4). Despite

greater bleaching at Reef 14 in October 2014 (Figure 2b), shifts in bleaching and recovery

performance envelopes were most distinct at Lilipuna (Figure 4). At the physiological level,

separation of physiotypes during and after thermal stress was driven by bleaching sensitivity and

symbiont cells/chlorophyll concentrations ($p<0.001$, Figure 4 & S2, Table S1). "Bleached"

physiotypes observed in periods of temperature stress (October 2014 and 2015) generally aligned

with colonies hosting the more thermally sensitive *Cladocopium* sp., although in the first event

coral physiotypes at Reef 14 symbiont communities did not separate according to symbiont

communities (Figure 4). Nevertheless, across all sampling periods *M. capitata* colonies

dominated by *Durusdinium* symbionts had higher symbiont cell densities, less variable areal

chlorophyll concentrations, and lower chlorophyll per symbiont cell (i.e., pg chlorophyll cell^{-1})

compared to colonies dominated by *Cladocopium* symbionts ($p<0.001$, Figure S2). While

Cladocopium chlorophyll cell^{-1} oscillated across bleaching-and-recovery periods, *Durusdinium*

pg chlorophyll cell^{-1} remained stable ($p=0.010$) and was also slightly higher at Reef 14 relative to

Lilipuna across all timepoints ($p=0.001$, Table S1). Coral protein and total biomass was variable

across the study, but each showed positive correlations with *Durusdinium*-dominated

physiotypes from Lilipuna in bleaching and recovery periods (Figure 4). Overall, protein was

9% higher in *Durusdinum*-dominated colonies (*p=*0.031) and 32% higher in Lilipuna corals

during first bleaching (October 2014) but equivalent at all sampling points thereafter (*p=*0.002).

Total biomass was lower during recovery periods (*p<*0.001) in addition to being 20 – 40% higher

at Lilipuna compared to Reef 14 in all periods (except in February 2015 recovery) (*p<*0.001) and

~30% higher in *Durusdinium*-dominated colonies during the second bleaching, but equivalent

across all colonies in other periods (*p=*0.012) (Figure S2).

Immunity and antioxidant metrics (i.e., MEL, PPO, POX, CAT, SOD) differed through time in

response to repeat bleaching and recovery (Figure 3, Figure 4) (*p<*0.001), while the

environmental influence from the Sit and Symbiont communities were less pronounced (Figure

S2, S3, and Table S1, S2). In the first event, corals responded to thermal stress by increasing

melanin synthesis (*p<*0.001) (with corresponding declines in PPO precursors) and increasing

catalase (*p<*0.001). During recovery, prophenoloxidase (*post-hoc*: *p<*0.001) and superoxide

dismutase increased (*post-hoc*: *p<*0.001) as melanin and catalase declined (Figure 4, 5, S2, & S3,

Table S1, S2). Notably, the melanin pathway increased in all corals in the 2014-bleaching event

regardless of bleaching sensitivity (i.e., loss or retention of symbionts/chlorophyll), or symbiont

community, and was a significant cellular response that shaped coral physiotypes in the first

bleaching event (Figure 5, first row). Corals in the second bleaching experienced a 6-fold

increase in melanin (October 2015 *vs.* February 2015) along with corresponding reductions in

prophenoloxidase, however, melanin activity was significantly less compared to the high

melanin activity observed in the first bleaching period (0.015 *vs.* 0.003 mg melanin mg tissue^{-1} in

first- versus second-bleaching period). Conversely to lower melanin in the second bleaching was

a peak in catalase activity, which reached its highest level in 2015 bleaching (*post-hoc*, *p<0.001*)

and was most pronounced in corals from Lilipuna compared to Reef 14 (*post-hoc*, *p<0.001*)

(Table S2). The peak in catalase during 2015 bleaching corresponded with the lowest observed

peroxidase activity, particularly for *Cladocopium*-dominated colonies (*post-hoc*, *p*=0.007).

Subsequently in the 2016 recovery period, catalase declined by ~70% and peroxidase activity

doubled, reaching peak activity similar to those observed in time periods. In contrast to other

immunity and antioxidant responses, superoxide dismutase showed a unique trajectory,

increasing progressively through time with each sampling period, being at its highest constitutive

activity after repeat bleaching in the 2016 recovery, which was double that observed in the first

bleaching period (October 2014) (*p<0.001*). Similar to catalase, superoxide dismutase was also

slightly higher in corals from Lilipuna compared to those at Reef 14 (*p*=0.028).

4. Discussion

Ocean thermal anomalies are reshaping the structure and function of coral reefs the world over.

Three mass bleaching events occurred since 2014 (Hughes et al., 2017), and climate models

predict that by mid-century bleaching will be an annual phenomena (van Hooidonk, Maynard, &

Planes, 2013). The stress responses of corals to ocean warming is based on a network of dynamic

interactions at biological and environmental levels, that can influence how they respond to the

physiological challenges posed by a warming planet (Suggett & Smith, 2019). Accordingly,

there is a need for holistic measures of coral response that incorporate the influence of and

environmental history (Palumbi et al., 2014) on physiological legacy effects. For example

assessment of symbiont community (Suggett, Warner, & Leggat, 2017) and cellular memory

(Brown, Dunne, Edwards, Sweet, & Phongsuwan, 2015) in the coral holobiont during and after

thermal stress, to test for shifting physiological status that may ameliorate or exacerbate coral

bleaching. Thus, physiotype tracking – or multivariate physiological and molecular time series

data – provide the capacity for identification of the cellular mechanisms that contribute to

resilience, acclimatization, and/or resistance that can allow for better predictions of coral

ecological responses to intensifying climate change and other regional disturbances (Vercelloni

et al., 2020).

Symbiosis – via differential performance of genera and types (Sampayo et al., 2008; Stat et al.,

2008) and the potential for switching and shuffling (Buddemeier & Fautin 1993; Baker et al.,

2003) – provide one clear example of how environmental history can shift performance

baselines. In this study, the role of dominant Symbiodiniaceae on holobiont physiotype was

present, but the symbiont community effects were often Period- or Site-dependent. The high

symbiont and chlorophyll *a* densities in *M. capitata* colonies dominated by *D. glynnii* across

Periods aligns with the paradigm of high thermotolerance within the genus *Durusdinium*

(Cunning et al., 2016; Lesser, Stat, & Gates, 2013; Silverstein, Cunning, & Baker, 2017; Wham

et al., 2017). However, antioxidant and immunity metrics were equivalent in colonies differing

in dominant symbiont partner, demonstrating negligible effects on antioxidant and immune traits

counter to the expectation that these would be modulated by symbiont-derived bleaching

resistance. In fact, this similarity was despite a greater loss of symbiont cells and

photopigmentation in colonies dominated by *Cladocopium* relative to *Durusdinium*. Symbiont

PSII photodamge is thought to lead the cascade of cellular events that culminate in bleaching

(Weis, 2008). However, some Symbiodiniaceae (*Symbiodinium* sp., *Durusdinium trenchii*) have

been shown to resist expulsion from the host despite incurring photodamage (Kemp et al., 2011;

Silverstein et al., 2017). Therefore, host mechanisms regulating redox status may be equally

important in coral thermal stress responses and decoupled from symbiont photophysiological

function (Krueger et al., 2015). In our study, the similarities observed between the antioxidant

and immune responses of *Cladocopium*- and *Durusdinium*-dominated *M. capitata* colonies

during both bleaching and recovery periods implicate an integral role of host mechanisms in

coral responses to bleaching and recovery despite the presence of functionally distinct symbiont

communities.

Shifts in physiotypes across bleaching and recovery periods were influenced by symbiont

communities and the physical environmental at each site, however, the significance of these

predictors were not uniform across time and varied during first- and second-bleaching events.

For instance, the effects of reef site were clear in the first event, possibly relating to greater

bleaching in 2014, which reduced symbiont cells in both corals dominated by *Cladocopium* sp.

and thermotolerant *D. glynii* at Reef 14. In the context of previous works, our data agrees with

findings that DHW in Kāneʻohe Bay were lower in the first bleaching event relative to the

second (5 – 7 DHW [2014] *vs*. 10 – 12 DHW [2015]) (Bahr et al., 2017), while the percentage of

bleached coral cover was higher in 2014 at our study sites (70% [2014] *vs*. 50% [2015]) and in

Kāneʻohe Bay as a whole (45 – 77% [2014] *vs*. 30 – 55% [2015]) (Bahr et al., 2017; Ritson-

Williams & Gates, 2020) possibly due to greater cumulative temperature stress in 2014 (Bahr et

al., 2017; Ritson-Williams & Gates, 2020). While this study and others agree that lower

bleaching prevalence was observed in Kāneʻohe Bay during the second bleaching event, we

show this corresponded with attenuated Site × Symbiont effects on coral physiotypes (Figure 4

bottom row), suggesting a positive feedback between this interaction and bleaching severity.

Therefore, the symbiont community harbored by corals is integral to bleaching responses and

coral physiotypes (Suggett et al., 2017), but the relative importance of symbiont effects are also

tempered by present environmental conditions and site-specific environmental histories.

Environmental conditions between Lilipuna and Reef 14 did not differ in terms of light

availability, seawater temperature, or DHW (Figures 2 & S1). Nevertheless, the hydrodynamics

between these reefs are substantial, producing considerable differences in seawater residence and

pCO_2 variability (Drupp et al. 2011, 2013). In addition, the fringing reef habitat of Lilipuna with

a close proximity to shore and silt-dominated backreef benthos is in stark contrast to Reef 14,

which is patch reef pinnacle in the middle of the Kāneʻohe Bay lagoon. Together, the

persistence of long-term hydrodynamic and biogeochemistry conditions between these reefs have

manifested as physiological legacies in resident *M. capitata*, which we demonstrate influence the

response of corals to regional bleaching events. In the laboratory, we showed *M. capitata* from

Reef 14 exhibited greater antioxidant activity and F_v/F_m but lower melanin compared to Lilipuna

colones (Wall et al., 2018). We hypothesize these patterns observed in laboratory and field

studies may be underpinned by energetic limitations due to pCO_2 histories. Immunity is

energetically expensive (Palmer et al. 2018b) and studies show corals exposed to high-pCO_2

upregulate genes for energy reserve metabolism (Vidal-Dupiol et al., 2013) and can have lower

lipid biomass compared to corals at ambient-pCO_2 (Wall, Mason, Ellis, Cunning, & Gates,

2017). In this case flow and pCO_2-history may be an important factor in the physiological

legacies of Kāneʻohe Bay *M. capitata*, which had a greater impact on during the first bleaching

event relative to the second.

Coral physiotypes are influenced by the physiological state and integrity of the coral-

Symbiodiniaceae symbiosis, which includes (but is not limited to) pigmentation and symbiont

densities. As expected, we observed symbiont loss/repopulation to drive changes in coral

physiotypes between bleaching and recovery periods, and in support of our hypothesis, prior

events had cumulative impacts on physiotype trajectories. Notably, coral physiotypes occupied

distinct spaces in each sampling period and showed changing trajectory through time (2014-

2016) with the terminus of the trajectory in 2016-recovery resulting in a coral trait space

intermediate between prior bleaching and recovery periods (Figure 1). These trajectories may be

driven by those responses active in either the first or second bleaching-recovery periods (such as

melanin), or by changes in constitutive immune activity that are cyclical (e.g., PPO and CAT), or

continuously increasing through time (e.g., SOD). For instance, changes in total biomass

generally declined in the aftermath of bleaching, being lower in recovery periods, but did not

follow patterns in symbiont or photopigment concentrations. Coral tissue biomass also increased

through time and was highest in the second year (on average pooled across all samples). The

progressive increases in constitutive antioxidant activity exemplified by superoxide dismutase,

may also have contributed to maintenance of biomass and therefore enhanced the potential for

overall coral colony survival (Thornhill et al., 2011). Therefore, the moderate increase in tissue

biomass through time, despite repeat stress events, may be a result of greater resilience in corals

surviving repeat bleaching events, in addition to seasonal and stress-dependent fluctuations in

coral tissue (Fitt, McFarland, Warner, & Chilcoat, 2000; Wall, Ritson-Williams, Popp, & Gates,

2019).

The observed differences in antioxidant and immune activity in corals during repeat bleaching

and recovery reveal that shifts in cellular priorities and mechanisms for dealing with bleaching

stress exist, which ultimately shapes coral physiotypes and their trajectories through time and

between stress states (Figure 5). In this case, increased catalase activity during bleaching may be

linked to reactive oxygen and nitrogen species and increased host apoptosis-like pathways

(Hawkins, Krueger, Becker, Fisher, & Davy, 2014; Krueger et al., 2015). Conversely, the

accumulation of superoxide dismutase is decoupled from the onset and subsidence of thermal

stress seen in catalase. This response may reflect the "cellular memory" of corals to thermal

stress, acting as a buffer to future oxidative stress (Barshis et al., 2013; Brown et al., 2015) to

reduce bleaching effects while also contributing to greater tissue retention and post-bleaching

survival. Both corals and Symbiodiniaceae have a broad array of superoxide dismutase isoforms

(Krueger et al., 2015; Richier, Sabourault, & Courtiade, 2006), which can be used to combat

reactive oxygen species originating from damage to photomachinery and host mitochondria that

together can trigger apoptosis (Dunn et al., 2012). The persistent increase of superoxide

dismutase through time thus suggests antioxidant "frontloading" (Barshis et al., 2013) is an

important strategy used by corals during repetitive thermal stress to reduce damage.

Engagement of the melanin pathway (collectively here as the prophenoloxidase reservoir and the

melanin product) is known to be an important generalized stress response for mitigating disease

and thermal stress effects in corals (Mydlarz, Holthouse, Peters, & Harvell, 2008; Palmer et al.,

2010; Wall et al., 2018). Likewise, exposure to thermal stress increases antioxidant activity in

corals and Symbiodiniaceae (Gardner et al., 2017). In the first-bleaching event, the melanin

pathway was observed as the primary cellular response to warming in all corals, regardless of

bleaching sensitivity or symbiont community. This initial peak in melanin synthesis further

supports the central role of this pathway as a generalized cellular response to periodic stress,

including wound repair, disease, photodamage, and bleaching, and/or a photoprotective role to

reduce excess excitation energy (Mydlarz et al., 2008; Palmer, Traylor-Knowles, Willis, &

Bythell, 2011; Palmer et al., 2010; Wall et al., 2018). Its subsequent decline suggests the

melanin cascade may i) prime the antioxidant response and no longer is required to act as a

primary stress response role, or ii) had its capacity overwhelmed in the second-bleaching,

resulting in corals primarily using antioxidant stress response mechanisms. Our findings are

more in line with the former. Under this thermal-priming hypothesis, sublethal thermal stress

and high variable temperature regimes can contribute to protective mechanisms in the coral

holobiont and bleaching resistance (Ainsworth et al., 2016; Barshis et al., 2013; Bellantuono,

Granados-Cifuentes, Miller, Hoegh-Guldberg, & Rodriguez-Lanetty, 2012; Safaie et al., 2018).

However, an increased frequency or magnitude of thermal stress may overcome acquired cellular

responses (or symbiont partnerships) that support coral resistance to bleaching and mortality

(Ainsworth et al., 2016).

In our study, the hierarchy of cellular stress responses shifts away from melanin-synthesis in

favor of antioxidant activity (Figure 5), which may reflect the specialized nature of antioxidants

in mitigating cellular damage and maintaining coral holobiont homeostasis (Murphy, Collier, &

Richmond, 2019). Antioxidant proteins are central to cellular stress response network

(Scandalios, 2002) and in corals there is clear evidence that antioxidant enzymes have a stress

inducible role, driven by both environmental stress and pathogen elicitors (Mydlarz & Harvell,

2007; Mydlarz et al., 2008; Palmer et al., 2011). Indeed, *M. capitata* exposed to thermal stress

showed greater antioxidant activity but lower melanin when primed by a history of high pCO_2-

variability (Wall et al., 2018). The increasing role of catalase, peroxidase, and superoxide

dismutase activity through time and in the second-bleaching event, compared to attenuated

melanin-synthesis pathway (Figure 5, S1, S2), further demonstrates that specialized antioxidative

enzymes are important for coping with chronic or persistent stress, as often reported in thermally

stressed corals (Gardner et al., 2017; Lesser, 1997). Such constitutive upregulation of cellular

defenses have also been observed in corals from areas of high thermal variance (Barshis et al.,

2013) and those acclimatized to warmer conditions before the onset of bleaching stress

(Bellantuono et al., 2012). In particular, the continued increases in superoxide dismutase and the

persistence of high levels of peroxidase in the final 2016-recovery period indicate that increasing

constitutive levels of select antioxidants is key to maintaining cellular homeostasis after repeated

or chronic thermal perturbations (Figure 5, S2). Such shifts in homeostatic mechanisms through

strategies like gene plasticity (Kenkel & Matz, 2016) or constitutive frontloading (Barshis et al.,

2013) reveal the important role of immunity, antioxidants, and their interplay in shaping

ecological and evolutionary trajectories of reef corals and the future of coral reefs.

In conclusion, we show coral physiotypes were distinct during each period of bleaching and

recovery from 2014-2016, and this indicates physiological legacy effects driven by chronic stress

or beneficial acclimation influence both bleaching and recovery trajectories. Bleaching was

strongly related to symbiont communities and their sensitivity to thermal stress, which was

influenced by site environmental history. However, physiotypes were shaped by the cascade of

immune and antioxidant activity in the bleaching and recovery periods irrespective of symbiont

communities, providing a mechanistic hypothesis for the shifting coral physiotype baseline.

Specifically, we observed changes in constitutive immunity from one year to the next in response

to extreme stress events, indicating a fundamental change in the homeostatic mechanisms

employed by corals. In an era of increasing frequency and magnitude of thermal stress, the

ongoing study of mechanisms such as frontloading of genes like heat shock proteins and

antioxidants to attenuate bleaching effects and support thermal resilience (Barshis et al., 2013) is

of great importance. Additionally, the examination of acclimatization of populations to

environmental stress that may be gained through gene expression plasticity (Kenkel & Matz,

2016), or epigenetic mechanisms (Durante, Baums, Williams, Vohsen, & Kemp, 2019; Eirin-

Lopez & Putnam, 2019; Liew et al., 2018) will be critical. Our study provides details of cellular

and physiological legacy effects in corals and evidence that environmental memory shapes the

homeostatic strategies of corals, which ultimately dictates a coral's ability to respond to future

stress events.

Acknowledgements

The authors acknowledge funding support from an Environmental Protection Agency STAR

Fellowship Assistance Agreement (FP-91779401-1) to C.B.W and NSF 1756623 (Biological

Oceanography, Integrative and Ecological Physiology, and EPSCoR) to H.M.P. The views

expressed in this publication have not been reviewed or endorsed by the EPA and are solely

those of the authors. We also thank P.J. Edmunds for insightful comments on an earlier

manuscript draft. This is SOEST contribution number xxx and HIMB contribution number xxx.

Author contributions

646   C.B.W., C.A.R., L.D.M., R.D.G., and H.M.P. designed the project. C.B.W., C.A.R., L.D.M., and

647   H.M.P. wrote the manuscript, and C.B.W. statistically analyzed the data. Coral collections were

648   performed by C.B.W. and A.D.W. Laboratory analyses were performed by C.B.W., C.A.R.,

649   A.D.W., B.E.L., and D.E.K.

Competing interests

652   The authors declare they have no competing interests.

Data availability

655   All data and code to generate figures and perform analyses are archived and openly available at

656   Github (https://github.com/cbwall/Gates-Mydlarz-bleaching-recovery/releases/tag/v4).

657   **Tables**

Table 1. Results of PERMANOVA testing the effects of repeated bleaching and recovery periods

on *Montipora capitata* corals hosting two distinct symbiont communities at two reef locations.

Factor	df	SS	R^2	F	p
Period	3	855.740	0.297	48.491	**<0.001**
Site	1	60.996	0.021	10.369	**<0.001**
Symbiont	1	202.349	0.070	34.399	**<0.001**
Period × Site	3	69.003	0.024	3.910	**<0.001**
Period × Symbiont	3	60.101	0.021	3.406	**<0.001**
Site × Symbiont	1	6.861	0.002	1.166	0.293
Period × Site × Symbiont	3	19.048	0.007	1.079	0.360
Residual	273	1605.902	0.558		
Total	288	2880.000	1.000		

Period = sequential bleaching and recovery events from October 2014 – February 2016, *Site* = Lilipuna or Reef 14, *Symbiont* = *Cladocopium* sp. or *Durusdinium glynnii* dominated symbiont community. *SS* = sum of squares; *df* = degrees of freedom; bold *p* values represent significant effects ($p < 0.05$).

663 **Figures**
664
665
666

667
668
Figure 1. (a) Multivariate analyses identify changes in organism physiotype through space and
time by accounting for the variation in multiple traits (Van Straalen, 2003). Transitions of
organisms from initial (α^0) to stressed states (β^1), such as during bleaching in response to a heat
wave, and their recovery to initial (α^0) (i.e., physiological resilience) or altered states (α^1).
Alternatively the absence of physiotype shifts may be observed in stress resistant individuals
(i.e., maintained at α^0). (b) Conceptual model of melanin and antioxidant activity in coral and
Symbiodiniaceae under heat stress (indicated by lightning bolts). Reactive oxygen species
(ROS), oxygen singlets ($^1O_2$) and superoxide (O_2^-), generated by symbiont photochemical
dysfunction and host mitochondria membrane damage, are neutralized by a combination of
antioxidant scavenger enzymes (e.g., superoxide dismutase [SOD], catalase [CAT], and
peroxidase [POX]). Host immunity via the melanin-synthesis pathway can also scavenge ROS
during intermediate steps that lead to the synthesis of melanin (MEL). Oxidative bursts during
the melanin-synthesis pathway can also create ROS that act as antimicrobials, which may also
lead to priming, or antioxidant enzymes up-regulation (shown in bold letters and dark lines).

Figure 2. *Montipora capitata* cover and bleaching patterns at two reefs in relation to thermal regimes. (a) A bleached *M. capitata* colony and (b) benthic surveys of absolute *M. capitata* at two sites (Lilipuna and Reef 14) during and after thermal stress in 2014-2015 and 2015-2016. (c) The proportion of bleached *M. capitata* colonies scored during (October) and after (February) thermal stress (values are mean ± SD, *n* = duplicate transects). Symbols indicate differences among sites within a period (*) and among sites (†). (d) Overlay of mean daily temperatures at Lilipuna and Reef 14 and the NOAA-HIMB Moku o Loʻe buoy from January 2014 - January 2016 and (e) a comparison of temperature ramping and cooling between the first and second bleaching events relative to our collection dates. Vertical dashed lines indicate coral collections during bleaching and recovery periods in the first (*black,* 2014-2015) and second (*gray*, 2015-2016) events.

Figure 3. Non-metric multidimensional scaling (NMDS) analyses of coral performance
envelopes during bleaching stress and post-bleaching recovery. *Montipora capitata* corals
dominated by *Cladocopium* sp. (red) or *Durusdinium glynnii* (blue) symbionts from two sites
(*solid lines* Lilipuna, *dashed lines* Reef 14). Convex hulls represent the coral physiotype (i.e.,
NMDS point clusters) of all corals in each time period, with points indicating mean centroids of
physiotype polygon for each group. Lines show trajectories of mean centroids for each group
across the four periods (Bleaching 2014, Recovery 2015, Bleaching 2015, Recovery 2016) and
triangle arrowheads indicate trajectory termini.
776
777

Figure 4. Non-metric multidimensional scaling (NMDS) analyses identify coral physiotypes
separated by dominant symbiont type (colors), bleaching-recovery periods (lighter and darker
shades), sites (columns), and events (rows). *Montipora capitata* corals dominated by
Cladocopium sp. or *Durusdinium glynnii* symbionts from Lilipuna (left panel) and Reef 14 (right
panel) during bleaching (Bleach) and recovery (Recov) periods. Biplot vectors (black arrows)
represent significant physiology and immunity responses ($p < 0.05$) according to squared
correlation coefficients (r^2). AFDW = ash-free dry weight biomass (mg gdw^{-1}), CAT = catalase,
MEL = melanin, POX = peroxidase, PPO = prophenoloxidase, SOD = superoxide dismutase,
chla = chlorophyll *a µg* cm^{-2}, chla cell = chlorophyll *a* per symbiont cell, prot = protein mg cm^{-2},
symb cells = symbionts cm^{-2}.

792
793
Figure 5. Schematic of mechanistic coral responses to repeat bleaching and recovery leading to
shifted physiotypes. In the first event (*top row*) thermal stress leads to a substantial increase in
MEL along with modest spikes in CAT and SOD. Corals in the first recovery showed increases
in PPO (as the precursor to the melanin-pathway) as well as SOD and POX, while CAT activity
declined. In the second event (*bottom row*), CAT and SOD spike with modest contributions of
MEL and a general decline in POX. Corals in the second recovery had the highest levels of SOD
across all time points, an increase in POX and a sharp decline in CAT (*see* Figure 4 for
abbreviations).
802

References

Ainsworth, T. D., Heron, S. F., Ortiz, J. C., Mumby, P. J., Grech, A., Ogawa, D., … Leggat, W.

(2016). Climate change disables coral bleaching protection on the Great Barrier Reef.

Science, *352*(6283), 338–342. doi: 10.1126/science.aac7125

Bahr, K. D., Rodgers, K. S., & Jokiel, P. L. (2017). Impact of Three Bleaching Events on the

Reef Resiliency of Kāneʻohe Bay, Hawaiʻi. *Frontiers in Marine Science*, *4*, 435. doi:

10.3389/fmars.2017.00398

Baker, A. C. (2003). Flexibility and Specificity in Coral-Algal Symbiosis: Diversity, Ecology,

and Biogeography of *Symbiodinium*. *Annual Review of Ecology, Evolution, and*

Systematics, *34*, 661–689.

Barshis, D. J., Ladner, J. T., Oliver, T. A., Seneca, F. O., Traylor-Knowles, N., & Palumbi, S. R.

(2013). Genomic basis for coral resilience to climate change. *Proceedings of the National*

Academy of Sciences of the United States of America, *110*(4), 1387–1392. doi:

10.1073/pnas.1210224110

Bellantuono, A. J., Granados-Cifuentes, C., Miller, D. J., Hoegh-Guldberg, O., & Rodriguez-

Lanetty, M. (2012). Coral Thermal Tolerance: Tuning Gene Expression to Resist Thermal

Stress. *PloS One*, *7*(11), e50685–14. doi: 10.1371/journal.pone.0050685

Box, G. E. P., & Cox, D. R. (1964). An Analysis of Transformations. *Journal of the Royal*

Statistical Society. Series B, Statistical Methodology, *26*(2), 211–243. doi: 10.1111/j.2517-

6161.1964.tb00553.x

Brown, B. E., Dunne, R. P., Edwards, A. J., Sweet, M. J., & Phongsuwan, N. (2015). Decadal

environmental "memory" in a reef coral? *Marine Biology*, *162*(2), 479–483.

http://link.springer.com/10.1007/s00227-014-2596-2

Brown, B. E., Dunne, R. P., Goodson, M. S., & Douglas, A. E. (2000). Bleaching patterns in reef

corals. *Nature*, *404*(6774), 142–143. doi: 10.1038/35004657

Buddemeier, R. W., & Fautin, D. G. (1993). Coral bleaching as an adaptive mechanism.

Bioscience, *43*(5), 320–326.

Couch, C. S., Burns, J. H. R., Liu, G., Steward, K., Gutlay, T. N., Kenyon, J., … Kosaki, R. K.

(2017). Mass coral bleaching due to unprecedented marine heatwave in

Papahānaumokuākea Marine National Monument (Northwestern Hawaiian Islands). *PloS*

One, *12*(9), e0185121. doi: 10.1371/journal.pone.0185121

Cunning, R., & Baker, A. C. (2013). Excess algal symbionts increase the susceptibility of reef

corals to bleaching. *Nature Climate Change*, *3*, 259–262. doi: 10.1038/nclimate1711

Cunning, R., Ritson-Williams, R., & Gates, R. D. (2016). Patterns of bleaching and recovery of

Montipora capitata in Kāne 'ohe Bay, Hawai 'i, USA. *Marine Ecology Progress Series*,

551, 131–139. http://www.int-res.com/abstracts/meps/v551/p131-139/

De'ath, G., Fabricius, K. E., Sweatman, H., & Puotinen, M. (2012). The 27-year decline of coral

cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of*

Sciences of the United States of America, *109*(44), 17995–17999. doi:

10.1073/pnas.1208909109

Drupp, P. S., De Carlo, E. H., Mackenzie, F. T., Sabine, C. L., Feely, R. A., & Shamberger, K. E.

(2013). Comparison of CO_2 Dynamics and Air–Sea Gas Exchange in Differing Tropical

Reef Environments. *Aquatic Geochemistry*, *19*(5-6), 371–397. doi: 10.1007/s10498-013-

9214-7

Drupp, P., De Carlo, E. H., Mackenzie, F. T., Bienfang, P., & Sabine, C. L. (2011). Nutrient

Inputs, Phytoplankton Response, and CO_2 Variations in a Semi-Enclosed Subtropical

Embayment, Kaneohe Bay, Hawaii. *Aquatic Geochemistry*, *17*(4), 473–498. doi:

10.1007/s10498-010-9115-y

Dunn, S. R., Pernice, M., Green, K., Hoegh-Guldberg, O., & Dove, S. G. (2012). Thermal stress

promotes host mitochondrial degradation in symbiotic cnidarians: are the batteries of the

reef going to run out? *PloS One*, *7*(7), e39024. doi: 10.1371/journal.pone.0039024

Durante, M. K., Baums, I. B., Williams, D. E., Vohsen, S., & Kemp, D. W. (2019). What drives

phenotypic divergence among coral clonemates of *Acropora palmata*? *Molecular Ecology*,

28(13), 3208–3224. doi: 10.1111/mec.15140

Edmunds, P. J., Adjeroud, M., Baskett, M. L., Baums, I. B., Budd, A. F., Carpenter, R. C., …

Gates, R. D. (2014). Persistence and Change in Community Composition of Reef Corals

through Present, Past, and Future Climates. *PloS One*, *9*(10), e107525.

http://dx.plos.org/10.1371/journal.pone.0107525.s001

Edmunds, P. J., Pochon, X., Levitan, D. R., Yost, D. M., Belcaid, M., Putnam, H. M., & Gates,

R. D. (2014). Long-term changes in *Symbiodinium* communities in *Orbicella annularis* in

St. John, US Virgin Islands. *Marine Ecology Progress Series*, *506*, 129–144. doi:

10.3354/meps10808

Eirin-Lopez, J. M., & Putnam, H. M. (2019). Marine Environmental Epigenetics. *Annual Review*

of Marine Science, *11*, 335–368. doi: 10.1146/annurev-marine-010318-095114

Fitt, W. K., McFarland, F. K., Warner, M. E., & Chilcoat, G. C. (2000). Seasonal patterns of

tissue biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral

bleaching. *Limnology and Oceanography*, *45*(3), 677–685. doi: 10.4319/lo.2000.45.3.0677

Fuess, L. E., Mann, W. T., Jinks, L. R., Brinkhuis, V., & Mydlarz, L. D. (2018). Transcriptional

analyses provide new insight into the late-stage immune response of a diseased Caribbean

coral. *Royal Society Open Science*, *5*(5), 172062. doi: 10.1098/rsos.172062

Gardner, S. G., Raina, J.-B., Nitschke, M. R., Nielsen, D. A., Stat, M., Motti, C. A., … Petrou,

K. (2017). A multi-trait systems approach reveals a response cascade to bleaching in corals.

BMC Biology, *15*(1), 117. doi: 10.1186/s12915-017-0459-2

Hawkins, T. D., Krueger, T., Becker, S., Fisher, P. L., & Davy, S. K. (2014). Differential nitric

oxide synthesis and host apoptotic events correlate with bleaching susceptibility in reef

corals. *Coral Reefs* , *33*(1), 141–153. doi: 10.1007/s00338-013-1103-4

Hughes, T. P., Anderson, K. D., Connolly, S. R., Heron, S. F., Kerry, J. T., Lough, J. M., …

Wilson, S. K. (2018a). Spatial and temporal patterns of mass bleaching of corals in the

Anthropocene. *Science*, *359*(6371), 80–83. doi: 10.1126/science.aan8048

Hughes, T. P., Kerry, J. T., Álvarez-Noriega, M., Álvarez-Romero, J. G., Anderson, K. D.,

Baird, A. H., … Wilson, S. K. (2017). Global warming and recurrent mass bleaching of

corals. *Nature*, *543*(7645), 373–377. doi: 10.1038/nature21707

Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Chase, T. J., Dietzel, A., … Woods, R.

M. (2019). Global warming impairs stock–recruitment dynamics of corals. *Nature*,

568(7752), 387–390. doi: 10.1038/s41586-019-1081-y

Hughes, T. P., Kerry, J. T., Baird, A. H., Connolly, S. R., Dietzel, A., Eakin, C. M., … Torda, G.

(2018b). Global warming transforms coral reef assemblages. *Nature*, *556*(7702), 492–496.

doi: 10.1038/s41586-018-0041-2

Innis, T., Cunning, R., Ritson-Williams, R., Wall, C. B., & Gates, R. D. (2018). Coral color and

depth drive symbiosis ecology of *Montipora capitata* in Kāneʻohe Bay, Oʻahu, Hawaiʻi.

Coral Reefs , *37*(2), 423–430. doi: 10.1007/s00338-018-1667-0

Jeffrey, S. W., & Humphrey, G. F. (1975). New spectrophotometric equations for determining

chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. *Biochemie*

Und Physiologie Der Pflanzen: BPP, *167*(2), 191–194. doi: 10.1016/S0015-

3796(17)30778-3

Jokiel, P. L., & Brown, E. K. (2004). Global warming, regional trends and inshore environmental

conditions influence coral bleaching in Hawaii. *Global Change Biology*, *10*(10), 1627–

1641. doi: 10.1111/j.1365-2486.2004.00836.x

Jones, R. J., Hoegh-Guldberg, O., Larkum, A. W. D., & Schreiber, U. (1998). Temperature-

induced bleaching of corals begins with impairment of the CO_2 fixation mechanism in

zooxanthellae. *Plant, Cell & Environment*, *21*(12), 1219–1230.

Kenkel, C. D., & Matz, M. V. (2016). Gene expression plasticity as a mechanism of coral

adaptation to a variable environment. *Nature Ecology & Evolution*, *1*(1), 14. doi:

10.1038/s41559-016-0014

Krueger, T., Fisher, P. L., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., … Davy, S.

K. (2015). Transcriptomic characterization of the enzymatic antioxidants FeSOD, MnSOD,

APX and KatG in the dinoflagellate genus *Symbiodinium*. *BMC Evolutionary Biology*, *15*,

48. doi: 10.1186/s12862-015-0326-0

Krueger, T., Hawkins, T. D., Becker, S., Pontasch, S., Dove, S., Hoegh-Guldberg, O., … Davy,

S. K. (2015). Differential coral bleaching—Contrasting the activity and response of

enzymatic antioxidants in symbiotic partners under thermal stress. *Comparative*

Biochemistry and Physiology, Part A, *190*(C), 15–25. doi: 10.1016/j.cbpa.2015.08.012

Lenth, R. V. (2019). Russell Lenth (2019). emmeans: Estimated Marginal Means, aka Least-

Squares Means. R package version 1.3.5. https://CRAN.R-project.org/package=emmeans.

Journal of Statistical Software, *69*(1), 1–33. https://CRAN.R-

project.org/package=emmeans

Lesser, M. P. (1997). Oxidative stress causes coral bleaching during exposure to elevated

temperatures. *Coral Reefs* , *16*(3), 187–192. doi: 10.1007/s003380050073

Lesser, M. P., Stat, M., & Gates, R. D. (2013). The endosymbiotic dinoflagellates (*Symbiodinium*

sp.) of corals are parasites and mutualists. *Coral Reefs* , *32*(3), 603–611. doi:

10.1007/s00338-013-1051-z

Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., Michell, C. T., … Aranda, M.

(2018). Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-

building coral. *Science Advances*, *4*(6), eaar8028. doi: 10.1126/sciadv.aar8028

Lowe, R. J., Falter, J. L., Monismith, S. G., & Atkinson, M. J. (2009). A numerical study of

circulation in a coastal reef-lagoon system. *Journal of Geophysical Research*, *114*(C6), 997.

doi: 10.1029/2008JC005081

McClanahan, T. R., Ateweberhan, M., & Omukoto, J. (2007). Long-term changes in coral colony

size distributions on Kenyan reefs under different management regimes and across the 1998

bleaching event. *Marine Biology*, *153*(5), 755–768. doi: 10.1007/s00227-007-0844-4

McWilliam, M., Pratchett, M. S., Hoogenboom, M. O., & Hughes, T. P. (2020). Deficits in

functional trait diversity following recovery on coral reefs. *Proceedings. Biological*

Sciences / The Royal Society, *287*(1918), 20192628. doi: 10.1098/rspb.2019.2628

Murphy, J. W. A., Collier, A. C., & Richmond, R. H. (2019). Antioxidant enzyme cycling over

reproductive lunar cycles in *Pocillopora damicornis*. *PeerJ*, *7*, e7020. doi:

10.7717/peerj.7020

Mydlarz, L. D., & Harvell, C. D. (2007). Peroxidase activity and inducibility in the sea fan coral

exposed to a fungal pathogen. *Comparative Biochemistry and Physiology. Part A,*

Molecular & Integrative Physiology, *146*(1), 54–62. doi: 10.1016/j.cbpa.2006.09.005

Mydlarz, L. D., & Palmer, C. V. (2011). The presence of multiple phenoloxidases in Caribbean

reef-building corals. *Comparative Biochemistry and Physiology. Part A, Molecular &*

Integrative Physiology, *159*(4), 372–378. doi: 10.1016/j.cbpa.2011.03.029

Mydlarz, L. D., Couch, C. S., Weil, E., Smith, G., & Harvell, C. D. (2009). Immune defenses of

healthy, bleached and diseased *Montastraea faveolata* during a natural bleaching event.

Diseases of Aquatic Organisms, *87*(1-2), 67–78. doi: 10.3354/dao02088

Mydlarz, L. D., Holthouse, S. F., Peters, E. C., & Harvell, C. D. (2008). Cellular Responses in

Sea Fan Corals: Granular Amoebocytes React to Pathogen and Climate Stressors. *PloS One*,

3(3), e1811–e1819. doi: 10.1371/journal.pone.0001811

Mydlarz, L. D., McGinty, E. S., & Harvell, C. D. (2010). What are the physiological and

immunological responses of coral to climate warming and disease? *The Journal of*

Experimental Biology, *213*(6), 934–945. doi: 10.1242/jeb.037580

NOAA. (2019, August 14). Retrieved August 14, 2019, from National Data Buoy Center, Station

MOKH1 - 1612480 - Mokuoloe, HI website:

https://www.ndbc.noaa.gov/station_page.php?station=mokh1

NOAA. (2020, April 20). Coral Reef Watch, NOAA Satellite and Information Services,

https://coralreefwatch.noaa.gov/satellite/methodology/methodology.php

Oksanen, J., Guillaume, F. B., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner,

H. (2019). *vegan: Community Ecology Package*. 631–637. https://cran.r-

project.org/web/packages/vegan/index.html

Palmer, C. V. (2018a). Immunity and the coral crisis. *Communications Biology*, *1*, 91. doi:

10.1038/s42003-018-0097-4

Palmer, C. V. (2018b). Warmer Water Affects Immunity of a Tolerant Reef Coral. *Frontiers in*

Marine Science, *5*, 253. doi: 10.3389/fmars.2018.00253

Palmer, C. V., & Traylor-Knowles, N. (2012). Towards an integrated network of coral immune

mechanisms. *Proceedings. Biological Sciences / The Royal Society*, *279*(1745), 4106–4114.

doi: 10.1098/rspb.2012.1477

Palmer, C. V., Bythell, J. C., & Willis, B. L. (2010). Levels of immunity parameters underpin

bleaching and disease susceptibility of reef corals. *FASEB Journal: Official Publication of*

the Federation of American Societies for Experimental Biology, *24*(6), 1935–1946. doi:

10.1096/fj.09-152447

Palmer, C. V., McGinty, E. S., Cummings, D. J., Smith, S. M., Bartels, E., & Mydlarz, L. D.

(2011). Patterns of coral ecological immunology: variation in the responses of Caribbean

corals to elevated temperature and a pathogen elicitor. *The Journal of Experimental*

Biology, *214*(Pt 24), 4240–4249. doi: 10.1242/jeb.061267

Palmer, C. V., Traylor-Knowles, N. G., Willis, B. L., & Bythell, J. C. (2011). Corals Use Similar

Immune Cells and Wound-Healing Processes as Those of Higher Organisms. *PloS One*,

6(8), e23992–11. doi: 10.1371/journal.pone.0023992

Palumbi, S. R., Barshis, D. J., Traylor-Knowles, N., & Bay, R. A. (2014). Mechanisms of reef

coral resistance to future climate change. *Science*, *344*(6186), 895–898. doi:

10.1126/science.1251336

Pinzón C, J. H., Beach-Letendre, J., Weil, E., & Mydlarz, L. D. (2014). Relationship between

Phylogeny and Immunity Suggests Older Caribbean Coral Lineages Are More Resistant to

Disease. *PloS One*, *9*(8), e104787. doi: 10.1371/journal.pone.0104787

Pinzón, J. H., Kamel, B., Burge, C. A., Harvell, C. D., Medina, M., Weil, E., & Mydlarz, L. D.

(2015). Whole transcriptome analysis reveals changes in expression of immune-related

genes during and after bleaching in a reef-building coral. *Royal Society Open Science*, *2*(4),

140214. doi: 10.1098/rsos.140214

R Core Team. (2020). *R: A language and environment for statistical computing*. https://www.R-

project.org

Richier, S., Sabourault, C., Courtiade, J., Zucchini, N., Allemand, D., & Furla, P. (2006).

Oxidative stress and apoptotic events during thermal stress in the symbiotic sea anemone,

Anemonia viridis. *The FEBS Journal*, *273*(18), 4186–4198. doi: 10.1111/j.1742-

4658.2006.05414.x

Ritson-Williams, R., & Gates, R. D. (2020). Coral community resilience to successive years of

bleaching in Kāneʻohe Bay, Hawaiʻi. *Coral Reefs* . doi: 10.1007/s00338-020-01944-4

Robinson, J. P. W., Wilson, S. K., Jennings, S., & Graham, N. A. J. (2019). Thermal stress

induces persistently altered coral reef fish assemblages. *Global Change Biology*, *25*(8),

2739–2750. doi: 10.1111/gcb.14704

Safaie, A., Silbiger, N. J., McClanahan, T. R., Pawlak, G., Barshis, D. J., Hench, J. L., … Davis,

K. A. (2018). High frequency temperature variability reduces the risk of coral bleaching.

Nature Communications, *9*(1), 1671. doi: 10.1038/s41467-018-04074-2

Sale, T. L., Marko, P. B., Oliver, T. A., & Hunter, C. L. (2019). Assessment of acclimatization

and subsequent survival of corals during repeated natural thermal stress events in Hawaiʻi.

Marine Ecology Progress Series, *624*, 65–76. doi: 10.3354/meps13031

Sampayo, E. M., Ridgway, T., Bongaerts, P., & Hoegh-Guldberg, O. (2008). Bleaching

susceptibility and mortality of corals are determined by fine-scale differences in symbiont

type. *Proceedings of the National Academy of Sciences of the United States of America*,

105(30), 10444–10449. doi: 10.1073/pnas.0708049105

Scandalios, J. G. (2002). Oxidative stress responses--what have genome-scale studies taught us?

Genome Biology, *3*(7), REVIEWS1019. doi: 10.1186/gb-2002-3-7-reviews1019

Schoepf, V., Grottoli, A. G., Levas, S. J., Aschaffenburg, M. D., Baumann, J. H., Matsui, Y., &

Warner, M. E. (2015). Annual coral bleaching and the long-term recovery capacity of coral.

Proceedings. Biological Sciences / The Royal Society, *282*(1819), 20151887. doi:

10.1098/rspb.2015.1887

Silverstein, R. N., Cunning, R., & Baker, A. C. (2017). Tenacious D: *Symbiodinium* in clade D

remain in reef corals at both high and low temperature extremes despite impairment. *The*

Journal of Experimental Biology, *220*(Pt 7), 1192–1196. doi: 10.1242/jeb.148239

Stat, M., Morris, E., & Gates, R. D. (2008). Functional diversity in coral–dinoflagellate

symbiosis. *Proceedings of the National Academy of Sciences of the United States of*

America, *105*(27), 9256–9261.

Stimson, J., & Kinzie, R. A. (1991). The temporal pattern and rate of release of zooxanthellae

from the reef coral *Pocillopora damicornis* (Linnaeus) under nitrogen-enrichment and

control conditions. *Journal of Experimental Marine Biology and Ecology*, *153*(1), 63–74.

doi: 10.1016/S0022-0981(05)80006-1

Stuart-Smith, R. D., Brown, C. J., Ceccarelli, D. M., & Edgar, G. J. (2018). Ecosystem

restructuring along the Great Barrier Reef following mass coral bleaching. *Nature*,

560(7716), 92–96. doi: 10.1038/s41586-018-0359-9

Suggett, D. J., & Smith, D. J. (2019). Coral bleaching patterns are the outcome of complex

biological and environmental networking. *Global Change Biology*. doi: 10.1111/gcb.14871

Suggett, D. J., Warner, M. E., & Leggat, W. (2017). Symbiotic Dinoflagellate Functional

Diversity Mediates Coral Survival under Ecological Crisis. *Trends in Ecology & Evolution*,

1–11. doi: 10.1016/j.tree.2017.07.013

Thornhill, D. J., Rotjan, R. D., Todd, B. D., Chilcoat, G. C., Iglesias-Prieto, R., Kemp, D. W., …

Fitt, W. K. (2011). A Connection between Colony Biomass and Death in Caribbean Reef-

Building Corals. *PloS One*, *6*(12), e29535. doi: 10.1371/journal.pone.0029535

van Hooidonk, R., Maynard, J. A., & Planes, S. (2013). Temporary refugia for coral reefs in a

warming world. *Nature Climate Change*, *3*(5), 508–511. doi: 10.1038/nclimate1829

Van Straalen, N. M. (2003). Ecotoxicology becomes stress ecology. *Environmental Science &

Technology*, *37*(17), 324A – 330A. doi: 10.1021/es0325720

Venable, W. N., & Ripley, B. D. (2002). Modern Applied Statistics with S. Springer. *New York*.

Venn, A. A., Loram, J. E., & Douglas, A. E. (2008). Photosynthetic symbioses in animals.

Journal of Experimental Botany, *59*(5), 1069–1080.

Vercelloni, J., Liquet, B., Kennedy, E. V., González-Rivero, M., Caley, M. J., Peterson, E. E., …

Mengersen, K. (2020). Forecasting intensifying disturbance effects on coral reefs. *Global

Change Biology*. doi: 10.1111/gcb.15059

Vidal-Dupiol, J., Zoccola, D., Tambutté, E., Grunau, C., Cosseau, C., Smith, K. M., …

Tambutté, S. (2013). Genes related to ion-transport and energy production are upregulated

in response to CO_2-driven pH decrease in corals: new insights from transcriptome analysis.

PloS One, *8*(3), e58652. doi: 10.1371/journal.pone.0058652

Wall, C. B., Kaluhiokalani, M., Popp, B. N., Donahue, M. J., & Gates, R. D. (2020). Divergent

symbiont communities determine the physiology and nutrition of a reef coral across a light-

availability gradient. *The ISME Journal*. doi: 10.1038/s41396-019-0570-1

Wall, C. B., Mason, R. A. B., Ellis, W. R., Cunning, R., & Gates, R. D. (2017). Elevated pCO_2

affects tissue biomass composition, but not calcification, in a reef coral under two light

regimes. *Royal Society Open Science*, *4*(11), 170683. doi: 10.1098/rsos.170683

Wall, C. B., Ricci, C. A., Foulds, G. E., Mydlarz, L. D., Gates, R. D., & Putnam, H. M. (2018).

The effects of environmental history and thermal stress on coral physiology and immunity.

Marine Biology, Vol. 165. doi: 10.1007/s00227-018-3317-z

Wall, C. B., Ritson-Williams, R., Popp, B. N., & Gates, R. D. (2019). Spatial variation in the

biochemical and isotopic composition of corals during bleaching and recovery. *Limnology*

and Oceanography, *64*(5), 2011–2028. doi: 10.1002/lno.11166

Weis, V. M. (2008). Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of

symbiosis. *The Journal of Experimental Biology*, *211*(Pt 19), 3059–3066. doi:

10.1242/jeb.009597

Wham, D. C., Ning, G., & LaJeunesse, T. C. (2017). *Symbiodinium glynnii* sp. nov., a species of

stress-tolerant symbiotic dinoflagellates from pocilloporid and montiporid corals in the

Pacific Ocean. *Phycologia*, *56*, 396–409. doi: 10.2216/16-86.1

1071