Italian Odonates in the Pandora's Box: A Comprehensive DNA Barcoding Inventory Shows Taxonomic Warnings at the Holarctic Scale

Running Title: DNA barcoding and taxonomic warnings in Odonata

Andrea Galimberti[1][^]*, Giacomo Assandri[2][^], Davide Maggioni[3,4], Fausto Ramazzotti[1], Daniele Baroni[5], Gaia Bazzi[6], Ivan Chiandetti[7], Andrea Corso[8], Vincenzo Ferri[9], Mirko Galuppi[10], Luca Ilahiane[11], Gianandrea La Porta[12], Lorenzo Laddaga[13], Federico Landi[14], Fabio Mastropasqua[15], Samuele Ramellini[16], Roberto Santinelli[17], Giovanni Soldato[18], Salvatore Surdo[19], Maurizio Casiraghi[1]

ZooPlantLab, Department of Biotechnology and Biosciences, University of Milano - Bicocca, P.za Della Scienza 2, 20126-I Milan, Italy.
Area per l'Avifauna Migratrice, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), Via Ca' Fornacetta 9, 40064-I Ozzano Emilia BO, Italy.
Department of Environmental and Earth Sciences (DISAT), University of Milano - Bicocca, P.za Della Scienza 1, 20126-I Milan, Italy.
Marine Research and High Education (MaRHE) Center, University of Milano - Bicocca, 12030 Faafu Magoodhoo, Maldives.
Section of Ecology, Department of Biology, University of Turku, FI-20014 Turku, Finland.
CROS Varenna, via Venini 17, 23829 Varenna (LC), Italy.
Via Braide Podé 8, 33010 Colloredo di Monte Albano (UD), Italy.
Via Camastra 10, 96100 Siracusa, Italy.
Department of Biology, University of Rome 2 - Tor Vergata, Via della Ricerca Scientifica 1, 00133-I Rome, Italy.
Via Federico Borromeo 24, 20822 Seveso (MB), Italy.
Department of Sciences and Technological Innovation (DISIT), University of Eastern Piedmont, Viale Teresa Michel 11, 15121 Alessandria, Italy.
Department of Chemistry, Biology and Biotechnology (DCBB), University of Perugia, Via Elce di Sotto 8, 06123 Perugia, Italy.
Società di Scienze Naturali del Verbano Cusio Ossola, Natural Science Museum Collegio Mellerio Rosmini, Domodossola (VB), Italy.
Via G. Mameli 14- I, 62100 Macerata, Italy.
Association Centro Studi de Romita, Via G. Postiglione 9, 70126-I Bari, Italy.
Department of Ecology and Environmental Policies, University of Milan, Via Celoria 26, 20133 Milan, Italy.
via Alfieri 24, 20855 Lesmo (MB), Italy.
Via Ormea 130, 10126 Torino, Italy.
Department of Agriculture, Food and Forest Sciences, University of Palermo, Viale delle Scienze, 90128 Palermo, Italy.

^ A. Galimberti and G. Assandri should be considered joint first authors.

* Corresponding author andrea.galimberti@unimib.it

Abstract

The Odonata are considered among the most endangered freshwater faunal taxa. Their DNA-based monitoring relies on validated reference datasets that are often lacking or do not cover important biogeographical centres of diversification. This study presents the results of a DNA barcoding campaign on Odonata, based on the standard 658 bp 5' end region of the mitochondrial COI gene, involving the collection of 812 specimens (409 of which barcoded) from peninsular Italy and its main islands (328 localities), belonging to all the 88 species (31 Zygoptera and 57 Anisoptera) known from the country. Additional BOLD and GenBank data from Holarctic samples expanded the dataset to 1294 DNA barcodes. A multi-approach species delimitation analysis involving two distance (OT and ABGD) and four tree-based (PTP, MPTP, GMYC, bGMYC) methods were used to explore these data. Of the 88 investigated morphospecies, 75 (85%) unequivocally corresponded to distinct Molecular Operational Units, whereas the remaining ones were classified as 'warnings' (i.e., showing a mismatch between morphospecies assignment and DNA-based species delimitation). These results are in contrast with other DNA barcoding studies on Odonata showing up to 95% of identification success. The species causing warnings were grouped in three categories depending on if they showed low, high, or mixed genetic divergence patterns. The analysis of haplotype networks revealed unexpected intraspecific complexity at the Italian, Palearctic, and Holarctic scale, possibly indicating the occurrence of cryptic species. Overall, this study provides new insights into the taxonomy of odonates and a valuable basis for future DNA and eDNA-based monitoring studies.

Keywords: Anisoptera, BOLD, Cryptic species, Odonata, Species delimitation, Zygoptera

Introduction

Modern conservation efforts rely on the availability of regional or local comprehensive species inventories (Morinière et al., 2019; Weigand et al., 2019; Altermatt et al., 2020). However, in most cases, morphology alone makes it hard to complete such inventories, so that alternative taxonomic methods, based on technological and analytical advances, have become fundamental to support and integrate the study of biodiversity (Padial et al., 2010; DeSalle & Goldstein, 2019; Schmid-Egger et al., 2019). Since the early 2000s, DNA barcoding has become a reliable basis for assembling the reference sequence libraries necessary to identify specimens of known species, also enhancing species discovery in neglected or poorly investigated taxonomic groups (e.g., Hebert et al., 2003; Galimberti et al., 2012; Dapporto et al., 2019). Curated and comprehensive DNA barcode reference libraries such as the Barcode of Life Data System BOLD (Ratnasingham & Hebert, 2007; 2013) often allow fast and reliable species identification with a considerable saving in terms of time and resources when personnel and taxonomic expertise are limited.

DNA barcoding has certainly revolutionized modern taxonomy, but some caveats dealing for example with pseudogenes (Berthier et al., 2011), introgression, and incomplete lineage sorting phenomena may lead to misidentification or wrong taxonomic assumptions (Eberle et al., 2020). However, even considering these possible pitfalls, the use of a variety of sequence analysis methods and parameters can be adopted to detect warnings (i.e., mismatches between morphospecies assignment and DNA-based species delimitation) of possible taxonomic relevance from DNA barcoding data (De Salle & Goldstein, 2019; Matos-Maraví et al., 2019). Such warnings could be then used to plan subsequent research aimed at assessing presumptive new species detected with these methods (Carstens et al., 2013). For example, by integrating morphology, multi-locus genetics, ecology or other information, the molecularly defined

species could be either confirmed or rejected (Kajitoch, Montagna, & Wanat, 2018; Dufresnes

et al., 2019). On the other hand, morphological species that are inconsistently delimited at the

molecular level could be designated for further analyses, such as geometric morphometrics

(Solano et al., 2018) or population genomics (Dufresnes et al., 2020).

Another possible weakness of DNA barcoding regards the geographical distribution of the

samples. Some studies observed a diminished identification accuracy due to the increasing

intraspecific genetic divergence when a wider geographical sampling scale was considered

(Meyer & Paulay 2005; Bergsten et al. 2012) or when genetic diversity hotspots are ignored or

undersampled (Gaytán et al., 2020). The availability of reliable reference databases covering

taxonomic diversity and including the whole intraspecific geographic variability of the

considered taxa constitutes a keystone element for large biodiversity investigations associated

with DNA and environmental DNA (eDNA) metabarcoding (Porter & Hajibabaei, 2018; Piper

et al., 2019; Creedy et al., 2020).

In this context, freshwater ecosystems are gaining more and more attention concerning

eDNA-based studies (Pawloski et al., 2018; Bush et al., 2019; Weigand et al., 2019) since they

are globally threatened and there is an increasing need to improve the monitoring of their

biodiversity and changes over time (Collen et al., 2014; Darwall et al., 2018). Among

freshwater bioindicators, the Odonata is a group of growing social and scientific interest. It is

a small order, by insect standards, counting about 6300 species worldwide (Schorr & Paulson,

2019); 145 species have been reported in Europe, with 51 and 94 species belonging to the

suborders Zygoptera (damselflies) and Anisoptera (dragonflies), respectively (Boudot &

Kalkman, 2015; Viganò, Janni, & Corso, 2017; Lopez-Estrada et al., 2020). Traditionally,

odonate identification has relied on morphological data that could be biased by difficulties in

larval/exuviae recognition, the occurrence of cryptic species, and even high introgression rates

(Marinov 2001; Bried, D'Amico, & Samways, 2012; Solano et al., 2018; Bried & Hinchliffe,

2019). The integration of traditional field protocols with alternative strategies, such as DNA

and eDNA-based approaches, may support fast and accurate species identification and

monitoring of this insects order.

Although the Odonata, together with Trichoptera, Hemiptera, and crustaceans are the

best-covered taxa in the BOLD System database (Weigand et al., 2019), the currently available

data do not take into account possible biases due to the absence of representatives from areas

of biogeographical/speciation interest (Gaytán et al., 2020), and to the absence of genetic

information (including DNA barcoding data) for some species of conservation concern (Sahlén

et al., 2010; Boudot & Kalman, 2015; Kalkman et al., 2018). Europe can be considered the

cradle of odonate systematics, but the evolutionary history and radiation of many genera,

especially in Mediterranean glacial refugia, is still unresolved (e.g., *Calopteryx*, *Ischnura* and

Onychogomphus, see Dijkstra & Kalkman, 2012). In the Western Mediterranean, at least three

areas have been recognized as distinct glacial refugia during Pleistocene glaciations: Iberia,

peninsular Italy (and its islands) and the Maghreb (Schmitt & Varga, 2012; Husemann et al

2013; Heiser, Dapporto, & Schmitt, 2014). Similarly, the Alps acted as a barrier and a refuge

through climatically induced range changes, and for this reason are today recognized as a major

geographical feature in shaping the phylogeography of European species (Hewitt, 2004). For

this reason, the Italian peninsula, which is a natural bridge between the Mediterranean basin

and the Alps, is a major biodiversity hotspot for odonates with 89 breeding species (Riservato

et al., 2014a; http://www.odonata.it/libe-italiane/), of which 19 are listed as threatened or near-

threatened in the Italian Red List (Riservato et al. 2014b) and 10 listed in the European

"Habitat" Directive. Nevertheless, almost no genetic information is available in public

databases for Italian odonates with less than 100 COI barcode sequences deposited there,

mostly belonging to a few *Cordulegaster* and *Coenagrion* species. Thus, the claim that Italy

has not played an important biogeographical and evolutionary role for the odonate

diversification (Heiser & Schmitt, 2010) appears questionable, given the limited amount of

molecular investigation occurred to date.

In order to increase the availability of genetic information on European Odonata, in this

study we aimed at *i*) creating the first comprehensive and well-curated reference DNA

barcoding library of Italian species, based on intense national sampling encompassing the main

regions of biogeographical interest; *ii*) investigating the diversity and distribution of

mitochondrial lineages of Italian Odonata using multiple species delimitation approaches,

based on both character-explicit and distance-based methods (*sensu* DeSalle & Goldstein,

2019). Moreover, since almost all the species occurring in Italy are more widely distributed in

Europe, Asia and North America, the same approach has also been performed at the Holarctic

scale by creating a second comprehensive dataset including a mixture of Italian and non-Italian

publicly available sequences.

Materials and Methods

Specimen collection, identification and archiving

During 2018-2019, a network of 20 professional and voluntary taxonomists (including

the authors), involved in the "DNA barcoding Italian odonates" project, collected odonates in

six macro-regions of Italy (see Figure 1). Such regions correspond to the main cradles of

intraspecific genetic divergence previously identified for other taxa (e.g., Dapporto, 2010;

Bernini et al., 2016; Wauters et al., 2017; Dufresnes et al., 2019; Scalercio et al., 2020).

Sampling sites included 328 localities scattered in most of the Italian administrative

regions and were mostly wet ecosystems belonging to protected or not protected areas from the

Alpine (up to 2300 m a.s.l.), peninsular and insular (i.e., Sardinia and Sicily) regions. For the

sampling of protected species, the Italian Ministry of the Environment issued a special permit

(Prot. DPN 0031783.20-11-2019).

Only adults were sampled by hand netting and identified according to Dijkstra & Lewington

(2006). In some cases, especially for damselflies belonging to *Coenagrion* and *Chalcolestes*

genera, fine characters diagnostic for species identification (e.g., pronotum, cerci and

ovipositor) were checked using a stereomicroscope (Olympus SZX7). The nomenclature used

in this study follows the last update of the World Odonata List (Schorr and Paulson, 2019). For

each individual, one or two middle legs were stored in 96% EtOH before DNA extraction. All

samples and most specimens were vouchered and stored in the MIB:ZPL collection (University

of Milano - Bicocca, Milan, Italy) whereas the remaining specimens are conserved in GA

private collection.

A total of 812 specimens belonging to 88 morphospecies (31 Zygoptera and 57 Anisoptera)

were collected. Sampling localities and details are available in BOLD under the project code

ZPLOD

(https://www.boldsystems.org/index.php/MAS_Management_DataConsole?codes=ZPLOD)

and are also reported in Appendix S1. Sampling location coordinates were given in decimal

degrees (EPSG: 4326) and approximated at the first decimal unit (i.e., $\approx$ 11 km). Since many

species (or biotopes) are of conservation concern, we chose not to give details on exact locations

(Lunghi et al., 2019).

178   **[FIGURE 1]**

DNA extraction, amplification, and sequencing

For each morphospecies, samples for genetic analysis were selected from distant sites

to maximize the chance of observing intraspecific geographic variation. Overall, 409 specimens

were selected for DNA barcoding characterization. DNA was extracted from a middle leg

(Zygoptera) or a coxa (Anisoptera) using the Qiagen DNeasy Blood and Tissue Kit (Qiagen,

Milan, Italy) following the manufacturer's instructions.

For each sample, the standard DNA barcode 5'-end region of the mitochondrial COI gene (658

bp) was amplified with the M13-tailed primers ODOF1_t1 and ODOR1_t1 (Semotok,

unpublished, Source: BOLD Systems primer database), previously used by Koroiva and co-

workers (2017). In case of unsuccessful amplification, the alternative COI primers

ODO_LCO1490d/ ODO_HCO2198d (Dijkstra et al., 2014) were adopted to amplify the

selected region. In both cases, PCR amplification conditions were as follows: 94°C for 5 min,

35 cycles at 94°C for 60 s, 50°C for 90 s and 72°C for 60 s, and a final extension at 72°C for 7

min. PCRs were conducted as in Mazzamuto et al. (2016) and sequencing was performed bi-

directionally at Eurofins Genomics (Milan, Italy). Standard M13F and M13R oligos were used

for sequencing the amplicons obtained with the ODOF1_tl/ODOR1_tl primer pair. Consensus

sequences were obtained by editing the electropherograms with Bioedit 7.2 (Hall, 1999). After

primer trimming, the presence of open reading frame was verified for the obtained consensus

sequences   by   using   the   on-line   tool   EMBOSS   Transeq

(http://www.ebi.ac.uk/Tools/st/emboss_transeq/), then sequences were aligned with MAFFT

7.110 (Katoh & Standley, 2013) using the *E-INS-i* option. Consensus sequences were deposited

in the BOLD Systems (project code ZPLOD) and GenBank (accession numbers reported in

Appendix S1 and Appendix S2).

Sequence mining and datasets assembling

Orthologous COI barcode sequences, belonging to the species included in the Italian

odonate list (Riservato et al., 2014a) were downloaded from public repositories (i.e., BOLD

and GenBank). Sequences showing insertions/deletions, > 1% of missing sites or overlapping

less than 500 bp with the COI region amplified and sequenced for the Italian samples, were

discarded. Overall, a total of 885 COI barcode sequences were retrieved from public

repositories. These nucleotide sequences and those obtained in the present study were grouped

in two datasets: (i) dataset DS1, composed of the DNA barcoding sequences produced in this

study on Italian samples (i.e., 409) plus 69 sequences from Italian samples of 10 odonate species

analysed in previous studies (see Appendix S1); (ii) dataset DS2, composed of the sequences

mined from online databases plus dataset DS1 and other COI barcode sequences obtained from

additional samples retrieved by the authors in other western Palearctic countries (See Appendix

S2). The two datasets were kept separated in order to evaluate first the efficiency of the multi-

approach species delimitation and to detect taxonomic warnings (e.g., reduced interspecific

genetic variability or high intraspecific structure at the COI barcode locus) within Italian

reference specimens only (DS1). Then, to better investigate such warnings on a wider

geographic scale and find new cases of possible taxonomic relevance, the same species

delimitation strategy was adopted on sequence data from across the Holarctic range of the taxa

occurring in Italy (DS2).

Species delimitation analysis

Phylogenetic trees were reconstructed separately for Zygoptera and Anisoptera. First,

identical sequences were collapsed into unique haplotypes (See Appendix S1 and Appendix

S2) and appropriate substitution models were determined using JModelTest 2 (Darriba et al.,

2012) using the Akaike Information Criterion (AIC), the corrected Akaike Information

Criterion (AICc), and the Bayesian Information Criterion (BIC). For Anisoptera, the best-fitting

substitution models were GTR+I+G (AIC, BIC) and HKY+I+G (AICc), whereas for Zygoptera

GTR+I+G (AIC) and HKY+I+G (BIC, AICc), and the use of different models did not influence

downstream analyses. Phylogenetic inference analyses were performed using two optimality

criteria: Bayesian inference (BI) and maximum likelihood (ML). BI was conducted by using

BEAST 1.8.2 (Drummond et al., 2012), setting a coalescent tree prior and an uncorrelated

lognormal relaxed clock. Three replicate analyses were run for 10^8 million generations with a

sampling frequency of 10,000 and were combined using LogCombiner 1.8.2 (Drummond et al.,

2012) with a burn-in set to 25%, after checking the stationarity for effective sampling size (ESS)

and unimodal posterior distribution using Tracer v.1.6 (Rambaut et al., 2014), making sure ESS

values were higher than 200 for all parameters. Maximum clade credibility trees were computed

using TreeAnnotator 1.8.2 (Drummond et al., 2012). ML analyses were performed using

RAxML 8.2.9 (Stamatakis, 2014) with 1000 bootstrap replicates, applying the GTR+I+G

substitution model. Each sub-order dataset was rooted using two sequences belonging to the

other dataset (*Nehalennia speciosa* (Charpentier, 1840) and *Pyrrhosoma nymphula* (Sulzer,

1776) for Anisoptera, and *Libellula quadrimaculata* Linnaeus, 1758 and *Lindenia tetraphylla*

(Vander Linden, 1825) for Zygoptera). The obtained trees were visualised using FigTree v.1.4.3

(Rambaut, 2012) and customized using CorelDRAW X7. All analyses were run on CIPRES

server (Miller, Pfeiffer, & Schwartz, 2010).

Distance-based and tree-based species delimitation approaches (*sensu* DeSalle &

Goldstein, 2019) were employed to explore species boundaries in DS1, after removing the

outgroups.

Distance-based methods rely on genetic distances to explore the presence of barcoding gaps

and allow for species non-monophyly (Fontaneto, Flot, & Tang, 2015). In this work, we used

the Automatic Barcoding Gap Discovery (ABGD; Puillandre et al., 2012) and the Optimal

Threshold (OT) (Brown et al., 2012). ABGD delimitations were run on the website

https://bioinfo.mnhn.fr/abi/public/abgd/. Parameters were set keeping Pmin = 0.001, Steps =

100, X = 1, Nb bins = 20, distance = Kimura 2-parameter, and varying Pmax between 0.01 to

0.1, in order to test the consistency of delimitations by changing the *a priori* maximum level of

intraspecific distance. Only results obtained with Pmax = 0.1 have been reported. OT was

calculated from the K2P genetic distance matrices with the 'LocalMinima' and 'treshOpt'

functions of the SPIDER package (Brown et al., 2012) in R (R Core Team, 2019). In particular,

OT is the genetic distance value that minimizes the cumulative identification error, that is the

sum of false positive (no conspecific matches within the threshold of the query) and false

negative (sequences from multiple species within the threshold) cases (Galimberti et al., 2012).

Tree-based methods require species monophyly and are based on the analysis of branching rates

(Fontaneto, Flot, & Tang, 2015). Here, the Poisson Tree Process (PTP; Zhang et al., 2013) and

Generalized Mixed Yule-Coalescent (GMYC; Pons et al., 2006; Fujisawa & Barraclough,

2013) models were used. Single-threshold Bayesian PTP analyses (Zhang et al., 2013) were

performed on the website http://species.h-its.org/ptp, running the analyses for 400,000 MCMC

generations, with thinning value = 100 and burn-in = 0.25, and checking the trace file for

convergence of the MCMC, whereas multiple-threshold PTP (MPTP) analyses (Kapli et al.,

2017) were run on the website https://mptp.h-its.org. Single-threshold GMYC (Pons et al.,

2006) and bGMYC (Reid & Carstens, 2012) analyses were run using BI ultrametric trees and

were performed in R (R Core Team, 2019) using the packages SPLITS (Ezard, Tomochika, &

Barraclough, 2009), APE (Paradis, Claude & Strimmer, 2004), and bGMYC (Reid & Carstens,

2012). bGMYC analyses were performed on a subset of 100 trees retrieved from the 10,000

trees obtained with each BI analysis, after assessing the convergence of the MCMC. Support

values for the species delimitation hypotheses obtained with the Bayesian implementation of

PTP and GMYC are shown in Appendix S3.

Apart from OT, all the other species delimitation approaches were independent, requiring no *a*

priori information on the existing morphospecies. Overall, we considered a delimitation to be

reliable when the different methods were mostly in agreement. The same pipeline was applied

to the Zygoptera and Anisoptera COI sequences alignments of DS2. Conversely, OT values

remained the same inferred from DS1 because reliable *a priori* information on morphospecies

was available only for the reference samples collected in this study.

When mismatches occurred between delimitation based on DNA barcoding data and

morphospecies assignment based on samples collected in this study (DS1) and/or inferred from

public data (DS2), we identified a 'warning'. Such mismatches have been grouped into three

distinct categories (i.e., no interspecific-, intraspecific- and mixed- delimitation), depending on

if they occur in spite of low, high or mixed genetic divergence patterns.

Considering the possible pitfalls in using a single mitochondrial marker to investigate species

boundaries (Lohse, 2009; Petit & Excoffier, 2009), additional information from molecular

repositories, literature and unpublished data from the entomologists participating in this study

were considered to support the reliability of delimited species and legitimate the necessity of

better investigating the detected warnings.

Warnings investigation

To better support the main categories of warnings resulting from species delimitation

analyses and to explore patterns of intraspecific geographic genetic variation, multi-locus

genetic distance comparisons and/or haplotype networks were generated for selected species

groups. In particular, public nucleotide sequences of the mitochondrial 16s rDNA and the

nuclear AgT, PRMT and MLC genes were retrieved from GenBank for *Coenagrion mercuriale*

(Charpentier, 1840), *C. puella* (Linnaeus, 1758), *C. pulchellum* (Vander Linden, 1825) and *C.*

ornatum (Sélys, 1850). These nuclear markers were selected among the others as they were

found to show a higher phylogeographic resolution in damselflies (Ferreira et al., 2014a).

Genetic distances among species or lineages were computed using MEGA X (Kumar et al.,

2018), and intra- and inter-clade uncorrected *p*-distances were calculated with 1000 bootstrap

replicates.

In case of intraspecific delimitation, and when at least 20 COI barcode sequences were available

from specimens collected in Italy and across the species Holarctic range, unrooted minimum

spanning COI networks were obtained using the median-joining algorithm implemented in

PopART 1.7 (Leigh & Bryant 2015).

Results

A reference DNA barcoding database of Italian odonates

We obtained high-quality full-length COI DNA barcode sequences (658 bp) for all the

409 selected specimens (see Appendix S1 for BOLD Process IDs and GenBank accession

numbers), representing 88 morphospecies belonging to 36 genera and 10 families. No

sequences contained insertions/deletions (indels), stop codons, or were biased by NUMT

interference, with the only exception of one *Platycnemis pennipes* (Pallas, 1771) specimen

(ZPLOD687-20) and one *Orthetrum cancellatum* (Linnaeus, 1758) specimen (ZPLOD613-20)

with the primer pair ODOF1_t1/ODOR1_t1. Re-amplification and sequencing with the other

primer set (i.e., ODO_LCO1490d/ODO_HCO2198d) allowed obtaining the correct DNA

barcode sequence for these samples.

The number of newly barcoded specimens per species ranged from 1 to 34 (mean $\pm$ SD = 4.6 $\pm$

4.6); 8 morphospecies were represented by a single accession (Appendix S1). The obtained

DNA barcode sequences allowed to add two Italian (and Palearctic) species (i.e., *Orthetrum*

nitidinerve (Sélys, 1841) and *Somatochlora meridionalis* Nielsen, 1935) not represented at all

in the BOLD and GenBank databases, and novel haplotypes for almost all the investigated taxa.

The DNA barcodes of DS1 were assigned to 83 distinct Barcode Index Numbers (BINs,

Ratnasingham & Hebert, 2013) by BOLD and at least three of these (i.e., BOLD:AEC5518,

BOLD:AEC4388, and BOLD:AEC4264 corresponding to *E. lindenii*, *L. dryas* and *O.*

nitidinerve, respectively) were unique to the system at the moment of dataset submission

(February 2020).

Considering the 69 Italian DNA barcode records mined from GenBank, a DNA

barcoding reference library of 478 COI sequences from 88 species was established (i.e, DS1,

see Appendix S1). The analysis of this dataset showed that genetic K2P-distance variation

within morphospecies ranged from 0% to 9.17% (mean $\pm$ SD= 0.48 $\pm$ 0.62 %) in Zygoptera and

from 0% to 2.64% (mean $\pm$ SD = 0.33 $\pm$ 0.29 %) in Anisoptera. Interspecific K2P-distance

values ranged from 0% to 27.29% (mean $\pm$ SD= 19.41 $\pm$ 3.71%) and from 0% to 25.28% (mean

$\pm$ SD= 18.25 $\pm$ 3.18 %) in Zygoptera and Anisoptera, respectively.

The distributions of intraspecific and interspecific K2P distances overlap, thus resulting in the

absence of a complete barcode gap in DS1 (Appendix S4), similarly to what observed by

Koroiva and Kvist (2018). Both ABGD and OT approaches identified a number of groups quite

lower than the collected morphospecies (Table 1). In particular, the optimal threshold that

minimises the number of false positive and false negative identifications resulted in 3.16% and

1.96% of K2P distance for Zygoptera and Anisoptera respectively, with associated cumulative

errors 24.1% (57 sequences out of 237) and 9.96% (24 sequences out of 241) and mainly due

to false negative cases. Overall, the Maximum intraspecific vs. Nearest Neighbour and Mean

intraspecific vs. Nearest Neighbour analyses conducted in BOLD, confirmed the occurrence of

a barcode gap for most investigated species groups (Appendix S5).

We obtained further 816 DNA barcode records of the same species included in DS1

from BOLD and GenBank referred to Holarctic specimens. These were assembled in a

comprehensive DNA barcoding inventory of 1294 COI sequences (i.e, DS2, see Appendix S2).

In DS2, Genetic K2P-distance variation within morphospecies ranged from 0% to 8.81% (mean

$\pm$ SD = 0.69% $\pm$ 0.78%) in Zygoptera and from 0% to 6.72% (mean $\pm$ SD = 0.59% $\pm$ 0.63%)

in Anisoptera. Interspecific K2P-distance values ranged from 0% to 27.29% (mean $\pm$ SD =

19.51 $\pm$ 3.67%) and from 0% to 26.69% (mean $\pm$ SD = 18.09% $\pm$ 3.17%) in Zygoptera and

Anisoptera, respectively. In contrast to DS1, the two genetic distance-based approaches (i.e.,

ABGD and OT) identified in DS2 a number of groups equal or quite higher than the collected

morphospecies (Table 1), nevertheless not showing a complete barcode gap (Appendix S4). To

better investigate this mismatch, multiple species delimitation approaches were conducted on

both datasets.

Species delimitation

Species delimitation analyses were conducted first on the reference DNA barcoding

DS1 dataset and were based on separate phylogenetic trees encompassing, respectively, the

damselfly and dragonfly morphospecies occurring in Italy (see Figure 2 and Figure 3). As

highlighted in the trees and reported in Table 1, the overall number of groups delimited upon

the criteria explained in Material and Methods, was quite lower than the number of recognized

morphospecies (i.e., 28/31 in Zygoptera and 56/57 in Anisoptera). In general, the congruence

between the delimited groups and the morphology-based identifications ('no warnings' in Table

1) was higher in Anisoptera, with 52 out of 57 (91.2%) cases, than in Zygoptera, with 21 out of

31 (67.7%) cases. Most of the warnings were due to reduced or no interspecific genetic diversity

with seven and four morphospecies involved in Zygoptera and Anisoptera, respectively.

Conversely, one case in both suborders was related to high genetic divergence at the

intraspecific level (i.e., *Erythromma lindenii* (Sélys, 1840) and *Onychogomphus forcipatus*

(Linnaeus, 1758)). Finally, in two damselfly species (i.e., *Chalcolestes viridis* (Vander Linden,

1825) and *C. parvidens* Artobolevskii, 1929), the pattern of genetic divergence was a mix of

the other two warning categories with one sample (ZPLOD168-20), morphologically identified

as *C. viridis*, included in the group of *C. parvidens* (see Figure 2, Figure 3 and Table 1).

383   **[FIGURE 2]**

385   **[FIGURE 3]**

387   **[TABLE 1]**

In contrast to DS1, the overall number of delimited groups in DS2 was higher than the

number of investigated species (i.e., 32/31 in Zygoptera and 61/57 in Anisoptera). Overall, the

warnings found in DS1 were almost all confirmed in DS2 (Table 1 and Appendix S6) and new

17 cases, belonging to 7 species (3 damselflies and 4 dragonflies) were found due to the new

BOLD and GenBank DNA barcode sequences included in this dataset. Table 2 provides a list

of the warning cases outlined in DS2.

In Zygoptera, several changes occurred in species delimitation between DS1 and DS2 involving

all the three categories of warnings (Table 1, Table 2 and Appendix S6). Warnings related to a

reduced interspecific genetic divergence slightly decreased because *Coenagrion puella,* which in DS1 was delimited in a single group with *C. pulchellum* and *C. ornatum*, in DS2 showed a mixed delimitation pattern due to the inclusion of some North African highly divergent DNA barcode sequences, already reported by a previous study (Ferreira et al., 2016). The remaining two cases belonging to this category involved the species pairs *Ischnura elegans* (Vander Linden, 1820) */ I. genei* (Rambur, 1842) and *Calopteryx splendens* (Harris, 1780) */ C. xanthostoma* (Charpentier, 1825). Conversely, the warning cases (and species) of intraspecific delimitation increased with *Coenagrion mercuriale* and *Lestes dryas* Kirby, 1890 showing three and two lineages respectively, whereas the same two lineages of *Erythromma lindenii* found in DS1 were confirmed. Finally, apart from the already cited case of *C. puella* and close congenerics, *Chalcolestes viridis* and *C. parvidens* showed a mixed delimitation pattern, similarly to DS1. In Anisoptera, the groups of species not delimited remained the same of DS1 (i.e., *Anax imperator* Leach, 1815 / *A. parthenope* (Sélys, 1839) and *Somatochlora metallica* (Vander Linden, 1825) / *S. meridionalis*), whereas the number of warnings due to high intraspecific delimitation increased from two (one species) to 11 (five species). Concerning these latter, two lineages were delimited in *Aeshna juncea* (Linnaeus, 1758), *Onychogomphus forcipatus*, *Stylurus flavipes* (Charpentier, 1825) and *Sympetrum danae* (Sulzer, 1776), while three groups were delimited from *Orthetrum albistylum* (Sélys, 1848) DNA barcode sequences.

[TABLE 2]

Finally, the "BIN Discordance" analysis conducted on BOLD was used to verify the warning cases mentioned above. Based on a similarity clustering strategy, COI barcode sequences are grouped into clusters which are assigned to a unique identifier called BIN

(Ratnasingham & Hebert, 2013). This tool confirmed almost all the warning cases revealed by

our multi-approach species delimitation. In particular, of the 88 barcoded Italian odonate

morphospecies, 68 were assigned to unique BINs, 13 shared the same BINs and 7 were assigned

to two or more BINs. The only case that did not match with our species delimitation results was

O. forcipatus, for which all the sequenced specimens clustered with the public deposited DNA

barcodes into a single BIN (see Appendix S1 for BINs assignment).

Warnings investigation

Some warning cases, resulting from the species delimitation of DS2, involved odonate

species represented by a high number of publicly available DNA barcode sequences from

different geographic regions or by the availability of multi-locus data deposited in GenBank.

These data allowed to better investigate the geographic distribution of genetic diversity at the

COI barcode region for at least four Italian species, also distributed at the Palearctic or Holarctic

scale (two Zygoptera, *Coenagrion mercuriale* and *Lestes dryas* and two Anisoptera, *Aeshna*

juncea and *Sympetrum danae*, see Figure 4). The three groups of *C. mercuriale* DNA barcodes

delimited in DS2 correspond precisely to three distinct haplogroups that are exclusive to Europe

(except for Italy), North Africa and Italy, showing no shared haplotypes and high values of K2P

and uncorrected *p*-distance among them (Figure 4, Appendix S7). Such isolation and genetic

distinctiveness are also supported by uncorrected genetic *p*-distance values at another

mitochondrial (16s rRNA) and three nuclear (AgT, PRMT, MLC) loci sequences retrieved in

GenBank (see Appendix S7).

In the other three species depicted in Figure 4, the European (and Italian) haplotypes were

clearly separated from the North American ones, supporting the delimitation inferred from DS2.

The number of substitutions characterizing this Holarctic disjunction was more than the double

in *S. danae* with respect to *A. juncea* and *L. dryas* delimited groups, with the only exception of

one *S. danae* haplotype from Canada (A388, see Appendix S2) that is much closer to the

Western Palearctic haplogroup.

Additional nuclear DNA sequences retrieved in GenBank allowed to better investigate the case

of absent interspecific delimitation among *Coenagrion puella*, *C. pulchellum* and *C. ornatum*.

However, these species were found to be clearly distinct at the three nuclear loci with no shared

haplotypes, thus indicating a mito-nuclear discordance pattern of genetic divergence (Appendix

S7).

454   **[FIGURE 4]**

Discussion

DNA barcoding performance and reference dataset

To the best of our knowledge, this study is one of the most comprehensive DNA

barcoding-based surveys on Odonata conducted on a national scale in the Northern Hemisphere.

Eighty-eight of the 89 Italian odonate species were successfully characterized by DNA

barcoding, thus increasing the available reference molecular data for many taxa distributed at

the European, Palearctic, or Holarctic scales (Figure 4, Table 2). *Ischnura fountaineae* Morton,

1905, was the only species which has not been collected, as just one isolated population is

known to occur on Pantelleria island (Corso et al., 2012).

The multi-approach species delimitation showed a moderately high congruence

between molecular groups and the Linnaean taxa with COI DNA barcodes, delimiting 85% of

species in the reference Italian database as separate MOTUs (71% and 93% in Zygoptera and

Anisoptera, respectively). Although most of the morphospecies are diagnosable, the

identification performance of this study is lower compared to other odonate DNA barcoding

surveys, which have reported up to a 95% success rate (Bergmann et al., 2013). However, all

the previous DNA barcoding studies regarding odonates were based on a lower number of

specimens and morphospecies (e.g., Bergmann et al., 2013; Kim et al., 2014; Koroiva et al.,

2017). Concerning sequence variation within and among morphospecies, the average values

obtained in this study are in line with those obtained by Koroiva and Kvist (2018) in their global

overview of DNA barcoding of odonates, with the only exception of the mean intraspecific

genetic variation in Italian Anisoptera (i.e., 0.33 DS1, 0.59 DS2), which is 4.8 (DS1) - 2.7 (DS2)

times lower than the global average (i.e., 1.60). No more than a slight variation was observed

in the average genetic distance values between DS1 and DS2. This was likely due to the higher

completeness of DS2, where intraspecific variation was the result of the inclusion of samples

from the whole Palearctic or Holarctic regions (Bergsten et al., 2012).

In the remaining cases (Table 1 and Table 2), the multi-approach species delimitation lumped

the morphospecies or split them into additional molecular groups.

Species delimitation and warnings

In this study, the multi-approach species delimitation conducted on DS1 showed that

32.3% of Zygoptera and 8.8% of Anisoptera species were highlighted by warnings (Table 1),

and these percentages increased in DS2 (38.7% and 15.8%). While the discrepancy between

the two datasets was due to the wider sampling area considered by the DS2 (see also Bergsten

et al., 2012), the higher warning rate in Zygoptera is more challenging to explain. In both

datasets, damselflies were involved in all the three possible groups of warnings, that even occur

simultaneously within the genus *Coenagrion* (Table 1, Figure 2, Appendix S6). The situation

of Italian Zygoptera could confirm previous findings that recorded a wide variety of global

biogeographical patterns among damselflies (e.g., Sánchez-Guillén et al., 2014a; Swaegers et

al., 2015; Ferreira et al., 2016; Wellenreuther & Sánchez-Guillén, 2016). Such a situation could

be partly due to the weaker dispersal ability of damselflies as compared to dragonflies (Heiser

& Schmitt 2010; Pinkert et al., 2018), which could have been resulted in more frequent isolation

and diversification events. However, other studies (reviewed in Wellenreuther & Sánchez-

Guillén 2016) point to sexual selection, and not ecological constraints, in playing the most

important role in damselflies diversification over time and space.

Warnings identified by species delimitation of DS1 and DS2 provide evidence for

different scenarios regarding Italian odonates taxonomy and conservation, with possible

influence also at the Holarctic scale.

No interspecific delimitation

This category of warning encompasses four cases in DS2 that were already confirmed

in the reference Italian sampling of DS1. The missing mitochondrial diversification in the

damselflies species pairs *Ischnura elegans/I. genei* and *Calopteryx splendens/C. xanthostoma*,

even though poorly investigated so far for Italian populations, was expected. The reproductive

isolation of both species pairs in Mediterranean areas is indeed highly debated by entomologists

with morphological and genetic evidence of introgressive hybridization, largely documented in

Ischnura (Sánchez-Guillén et al., 2014a; 2014b) and less clear in *Calopteryx* (Dumont,

Mertens, & De Coster, 1993; Weekers, De Jonckheere, & Dumont, 2001). Conversely, the

missing delimitation in *Somatochlora metallica/S. meridionalis* and *Anax imperator*/*A.*

parthenope was unprecedented, also because no Italian samples and no COI barcode sequences

at all for *S. meridionalis* were available before this study. This taxon was originally described

as a subspecies of *S. metallica* (Nielsen, 1935) and elevated at species rank more recently

(Schmidt, 1957); however, this status is still debated as intermediate specimens have been

recorded in the contact areas (Fleck, Grand, & Boudot, 2007). *S. meridionalis* replaces *S.*

metallica in southern Europe; in Italy, both the morphospecies occur with the former distributed

in central and southern portions of the peninsula (and in two disjunct and isolated populations

in the North) and the latter mainly spread in northern regions. Both taxa are sometimes found

syntopically (Boudot & Kalkman, 2015; Dijkstra & Kalkman, 2012) and lack clear structural

differences apart from subtle, though stable, differences in colouration in the adults (Boudot &

Kalkman, 2015) and minor differences at the larval stage (Seidenbusch, 1996). Species

delimitation based on COI barcode sequences from specimens coming from all the three

disjunct Italian ranges of *S. meridionalis* suggests that the two morphospecies in Italy are not

distinguishable according to mitochondrial DNA. More challenging is the case of the two *Anax*

species, which, contrary to *Somatochlora*, are well separated morphologically (but also

according to their ecology and life-history traits), although they may share the same

mitochondrial haplotypes. This scenario is not exclusive to Italian populations, as the same

pattern was confirmed by DS2 European and Palearctic populations (see also Rewicz et al.,

2020).

Intraspecific delimitation

The first group of morphospecies showing marked intraspecific genetic divergence

encompasses the Italian samples (DS1) of *Erythromma lindenii* and *Onychogomphus*

forcipatus. Concerning the former taxon, it was initially described as *Agrion lindenii*, placed in

the new genus *Cercion* by Navas (1907) and recently assigned to its current genus by Weekers &

Dumont (2004). Across its vast range, it shows remarkably little morphological variation, but the

eastern populations were described as *E. l. zernyi* (Schmidt, 1938). This taxon occurs from Turkey

eastward in isolated populations, surrounded by, and/or overlapping with the nominal subspecies;

thus, a longitudinal introgression generating hybrid populations was hypothesized, although no

genetic data has ever confirmed this (Dumont et al., 1995). Unexpectedly, our Italian sampling, and

the comparison with the few available European COI barcode sequences, provide evidence for the

occurrence of two supported mitochondrial lineages within western Palearctic (Figure 2, Appendix

S6). One of these lineages is completely new, and, based on these preliminary data, it is mainly

occurring in Montenegro and north-eastern Italy sampling localities.

Onychogomphus forcipatus has three recognized subspecies showing structural differentiation

(both in adults and nymphs) and occurring in Europe, with discrete and mostly non-overlapping

ranges. Of these, two subspecies are recorded in Italy, *O. f. forcipatus* being distributed only in the

northeast and allegedly in Sicily, and *O. f. unguiculatus* (Vander Linden, 1823) being distributed in

north-western and peninsular Italy (Boudot, Jacquemin, & Dumont, 1990, Boudot & Kalkman,

2015). Our species delimitation approach identifies two distinct lineages as well (also supported in

DS2); however, these do not show clear geographic separation, with a group including Sicilian,

central, southern, and north-western Italy samples, but also several samples from north-eastern Italy

and the other one encompassing a few samples from north-eastern Italy where the other lineage also

occurs. In this sense, our study, as that by Ferreira et al. (2014b), is still not resolutive, and further

integrated studies are needed to possibly solve this long-lasting taxonomic enigma.

Other warning cases of intraspecific delimitation emerged from the analysis of DS2 and

encompass lineages diverging at the Palearctic scale. The most interesting case is *Coenagrion*

mercuriale that forms three delimited haplogroups, of which one exclusive of the Italian

peninsula. In 1948, *C. castellani* was described from central Italy (Roberts, 1948), and in 1949

it was reported that *C. mercuriale* male specimens from Italy invariably showed clear

morphological differences (i.e., male appendages) with respect to specimens from central

Europe, suggesting to treat the Italian ones as *C. m. castellani* (Conci, 1949). Notably, the

Italian populations show a geographic disjunction from the range of *C. mercuriale*, as also

supported by genetic distance values at other mitochondrial and nuclear loci (Figure 4,

Appendix S7). Similarly, the populations from the Maghreb were referred to as *C. m.*

hermeticum (Selys 1872), since they appeared slightly morphologically distinct from the

nominal subspecies (Ben Azzouz, Guemmouh, & Aguesse, 1989). The validity of *hermeticum*

was questioned by several authors (e.g. Lieftinck, 1966), although the necessity of further

comparative investigations was advocated (Jacquemin & Boudot 1990; 1999). This long-

awaited study did not come to light until a recent multi-locus characterization of North African

populations (Ferreira et al., 2014a) and the data here provided for *castellani*. Although Boudot

& Kalkman (2015) stated that at present no subspecies of *C. mercuriale* are recognized, our

results suggest that three clearly distinct lineages occur within its range, one of which possibly

endemic to Italy. Further research including morphometric investigations is definitely needed

to clarify the matter, also considering the high conservation measures involving *C. mercuriale*

in Europe, which is listed in the Annex II of Council Directive 92/43/EEC.

Other two intraspecific delimitation warnings emerging from DS2 concerned *Orthetrum*

albistylum and *Stylurus flavipes*. These showed, respectively, two and one delimited Asian

lineages, in addition to the Italian/European one. Both taxa have a huge Eurasian range and

while some (maybe isolated) subspecies have been described for *O. albistylum* (e.g., *O. a.*

speciosum (Uhler, 1858), see Schorr & Paulson, 2019), *S. flavipes* has no currently accepted

subspecies (Steinmann, 2013; Schorr & Paulson, 2019). Unfortunately, only a few DNA barcode

sequences are available for these taxa and this does not permit further clarification of the observed

warnings.

The last group of intraspecific delimitation warnings includes taxa having a Holarctic

distribution, namely *Lestes dryas* (Zygoptera), *Aeshna juncea*, and *Sympetrum danae* (Anisoptera).

The COI DS2 haplotypes of these three species showed a marked geographic structure with

Italian/European haplotypes diverging from the North American ones (Figure 4, Appendix S6).

Regarding *A. juncea*, Bartenev in 1929 described the subspecies *A. j. americana* based on

material from Canada (Steinmann, 2013). Subsequently, Boudot & Kalkman (2015) stated that

none of the subspecies described should be considered valid while admitting that the specimens

from North America are clearly morphologically different from those found in Europe. *Lestes*

dryas and *S. danae* were described based on European specimens and no intraspecific taxa are

currently recognized (Bridges 1994; Steinmann 2013), making the warnings here presented and

their possible taxonomic significance an element of primary importance to be clarified.

Mixed species delimitation

This warning category involves two morphospecies groups of damselflies which DNA

barcoded individuals showed either marked genetic divergence or complete sequence identity

with congenerics. The first case, also occurring in DS1, concerns *Chalcolestes viridis* and *C.*

parvidens. The latter was originally described as a subspecies of *C. viridis* and was

subsequently elevated to the species rank (Dell'Anna, Utzeri, & De Matthaeis, 1996; Gyulavari

et al., 2011). Both species can occur syntopically in the Italian peninsula and in other parts of

Europe, with hybridization events reported (Olias et al., 2007). Our DNA barcoding data

confirm the genetic differentiation between *C. viridis* and *C. parvidens*, even if one specimen

from Apulia, morphologically identified as *C. viridis*, was genetically assigned to *C. parvidens*.

Several *Chalcolestes* populations from southern Italy are often difficult to be morphologically

identified (authors pers. obs.), and intermediate morphological traits could be due to the close

relationship and relatively recent divergence between the two taxa (Gyulavari et al., 2011).

Hybridization and/or incomplete lineage sorting events could have played a role in causing this

warning. A multi-locus phyogeographic survey, based on an extensive sampling, especially in

the contact zones between the two taxa, is needed to clarify the matter. Surprisingly, a mixed

species delimitation warning was found among *Coenagrion puella*, *C. pulchellum* and *C.*

ornatum in DS2, with *C. puella* showing a delimited diverging lineage corresponding to North

African populations and the other Italian/western Palearctic representatives of the three species

delimited in a single group and even sharing the same haplotypes (as already found in DS1).

These taxa can be found in syntopic conditions but can be clearly distinguished through

morphology, ecological requirements, life-history traits, though sporadic hybridization is

known to occur among them. The marked divergence at the investigated nuclear markers

(Appendix S7) indicates that a mito-nuclear discordance pattern occurs throughout their overall

distribution with the only exception of northern African populations, as already described by

Ferreira et al. (2016). Also in this case, further investigation is needed to better address such

discordance, also considering the high conservation value of *C. ornatum*, which is included in

the Annex II of Council Directive 92/43/EEC.

On the whole, our comprehensive DNA barcoding characterization of Italian Odonata

sheds light on several novel and other long-time debated taxonomic aspects of this taxon, with

an unexpected geographical resonance well beyond Italy. The newly detected lineages may lead

to the discovery of separate cryptic taxonomic entities within the Linnaean species and may

provide fertile ground for future studies. Conversely, the reduced or missing delimitation

between some known species challenges the current taxonomic knowledge on this presumed

"well-explored" group of insects (at least in Europe). The warnings resulting from our species

delimitation approach could indicate Italy as an important biogeographical and evolutionary

centre for odonates diversification. A wide array of possible scenarios could explain the

warnings found by our DNA barcoding-based species delimitation, such as the occurrence of

still undiscovered cryptic species (e.g., the unexpected and most surprising recent description

of a novel species endemic to Spain, *Onychogomphus cazuma*; López-Estrada et al., 2020) and

different forms of mito-nuclear discordance due to more or less ancient introgression

phenomena (e.g., Solano et al., 2018; Ottenburghs, 2020).

Further sampling campaigns, in Italy and throughout the whole range of the warning species

groups (as recommended by Gaytán et al., 2020), the integration of multiple genetic markers

(e.g., the universal single-copy orthologs USCOs recently described by Eberle et al., 2020), and

accurate morphometrics analyses will be necessary to correctly address each possible case of

taxonomic (and conservation) interest.

Finally, this thorough DNA barcoding investigation of Italian Odonata and related

warnings will allow to better support and interpret HTS eDNA metabarcoding data from

European and even Holarctic freshwater environments and, at the same time, to design species-

specific (or lineage-specific) probes to reliably address conservation studies on the whole range

of the investigated species.

Acknowledgments: Stefano Aguzzi, Gianluca Ancarani, Paolo Biella, Giovanni Boano, Marco Bonifacino, Martina Cadin, Gianmaria Carchini, Giovanni Colombo, Andrea Cusmano, Davide De Rosa, Federica Ferrario, Tiziano Fiorenza, Ilaria Fozzi, Carlo Galliani, Roberto Garavaglia, Gabriele Gheza, Luca Giussani, Marco Guglielmi, Cristiano Liuzzi, Davide Magnani, Alexandro Minicò, Valerio Orioli, Francesco Ornaghi, Silvia Pelti, Prisca, Verena Penna, Pietro Ramellini, Gloria Ramello, Francesca Ricci, Laura Ricci, Giuseppe Rossi, Leonardo Siddi, Elena Ternelli, Nicola Tommasi, Parco Adda Nord, Parco Lombardo della valle del Ticino, Parco della Valle Lambro, Parco Regionale Marturanum. We thank the association Odonata.it for warmly welcoming this project. The "Servizio Sviluppo Sostenibile e Aree Protette of Provincia Autonoma di Trento" allowed specimen collection in full-protected areas in Trentino (Prot. S175/2018/382452/17.11.2/LS/57EI); and the Italian Ministry of the Environment, Land and Sea released a national permit for the collection of species included in European and Italian conservation directives (Prot. 0031783.20-11-2019).

References

Altermatt, F., Little, C. J., Mächler, E., Wang, S., Zhang, X., & Blackman, R. C. (2020). Uncovering the complete biodiversity structure in spatial networks–the example of riverine systems. *Oikos*. https://doi.org/10.1111/oik.06806

Bergmann, T., Rach, J., Damm, S., DeSalle, R., Schierwater, B., & Hadrys, H. (2013). The potential of distance-based thresholds and character-based DNA barcoding for defining problematic taxonomic entities by CO 1 and ND 1. *Molecular Ecology Resources, 13(6)*, 1069-1081. https://doi.org/10.1111/1755-0998.12125

Ben Azzouz, B., Guemmouh, R. & Aguesse, P. (1989). À propos des *Coenagrion* du groupe *mercuriale* (Charpentier, 1840) et de la présence de *C. castellani* Roberts, 1948 au Maroc (Zygoptera: Coenagrionidae). *Odonatologica, 18*, 279-283.

Bergsten, J., Bilton, D. T., Fujisawa, T., Elliott, M., Monaghan, M. T., Balke, M., ... & Ribera, I. (2012). The effect of geographical scale of sampling on DNA barcoding. *Systematic Biology, 61(5)*, 851-869. https://doi.org/10.1093/sysbio/sys037

Bernini, G., Bellati, A., Pellegrino, I., Negri, A., Ghia, D., Fea, G., ... & Galeotti, P. (2016). Complexity of biogeographic pattern in the endangered crayfish *Austropotamobius italicus* in northern Italy: molecular insights of conservation concern. Conservation genetics, 17(1), 141-154. https://doi.org/10.1007/s10592-015-0767-4

Berthier, K., Chapuis, M. P., Moosavi, S. M., Tohidi-Esfahani, D. & Sword, G. A. (2011) Nuclear insertions and heteroplasmy of mitochondrial DNA as two sources of intra-individual genomic variation in grasshoppers. *Systematic Entomology, 36,* 285–299. https://doi.org/10.1111/j.1365-3113.2010.00561.x

Boudot, J.-P., Jacquemin, G., & Dumont, H. J. (1990). Revision of the subspecies of *Onychogomphus forcipatus* (Linnaeus, 1758) in Europe and Asia Minor, and the true distribution of *Onychogomphus forcipatus unguiculatus* (Vander Linden, 1832) (Odonata, Gomphidae). *Bulletin et annales de la Société royale belge d'entomologie, 126*, 95–111.

Boudot, J.-P., & Kalkman, V. J. (Eds.) (2015). Atlas of the European dragonflies and damselflies. The Netherlands: KNNV Publishing.

Bridges, C. A. 1993. Catalogue of the family-group, genus-group and species-group names of the Odonata of the world (Second Edition). Urbana: C. A. Bridges.

Bried, J. T., D'Amico, F., & Samways, M. J. (2012). A critique of the dragonfly delusion hypothesis: Why sampling exuviae does not avoid bias. *Insect Conservation and Diversity, 5(5)*, 398–402. https://doi.org/10.1111/j.1752-4598.2011.00171.x

Bried, J. T., & Hinchliffe, R. P. (2019). Improving taxonomic resolution in large-scale freshwater biodiversity monitoring: an example using wetlands and Odonata. *Insect Conservation and Diversity, 12(1)*, 9–17. https://doi.org/10.1111/icad.12323

Brown, S. D., Collins, R. A., Boyer, S., Lefort, M. C., Malumbres-Olarte, J., Vink, C. J., & Cruickshank, R. H. (2012). Spider: an R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. *Molecular Ecology Resources, 12(3)*, 562. https://doi.org/10.1111/j.1755-0998.2011.03108.x

Bush, A., Compson, Z. G., Monk, W., Porter, T. M., Steeves, R., Emilson, E. J., ... & Baird, D. (2019). Studying ecosystems with DNA metabarcoding: lessons from biomonitoring of aquatic macroinvertebrates. *Frontiers in Ecology and Evolution, 7,* 434. https://doi.org/10.3389/fevo.2019.00434

Carstens, B. C., Pelletier, T. A., Reid, N. M., & Satler, J. D. (2013). How to fail at species delimitation. *Molecular ecology*, *22*(17), 4369-4383. https://doi.org/10.1111/mec.12413

Collen, B., Whitton, F., Dyer, E. E., Baillie, J. E., Cumberlidge, N., Darwall, W. R., ... & Böhm, M. (2014). Global patterns of freshwater species diversity, threat and endemism. *Global Ecology and Biogeography, 23(1)*, 40-51. https://doi.org/10.1111/geb.12096

Conci, C. (1949) L'*Agrion mercuriale castellanii* Roberts in Italia (Odonata: Agrionidae). *Bollettino della Società Entomologica Italiana, 79*, 62-64.

Corso, A., Janni, O., Pavesi, M., Sammut, M., Sciberras, A., & Viganò, M. (2012). Annotated checklist of the dragonflies (Insecta Odonata) of the islands of the Sicilian Channel, including the first records of *Sympetrum sinaiticum* Dumont, 1977 and *Pantala flavescens* (Fabricius, 1798) for Italy. *Biodiversity Journal, 3*, 459–478.

Creedy, T. J., Norman, H., Tang, C. Q., Qing Chin, K., Andujar, C., Arribas, P., ... & Vogler, A. P. (2020). A validated workflow for rapid taxonomic assignment and monitoring of a national fauna of bees (Apiformes) using high throughput DNA barcoding. *Molecular Ecology Resources, 20(1)*, 40-53. https://doi.org/10.1111/1755-0998.13056

Dapporto, L. (2010). Speciation in Mediterranean refugia and post-glacial expansion of Zerynthia polyxena (Lepidoptera, Papilionidae). Journal of Zoological Systematics and Evolutionary Research, 48(3), 229-237. https://doi.org/10.1111/j.1439-0469.2009.00550.x

Dapporto, L., Cini, A., Vodă, R., Dincă, V., Wiemers, M., Menchetti, M., ... & Casacci, L. P. (2019). Integrating three comprehensive data sets shows that mitochondrial DNA variation is linked to species traits and paleogeographic events in European butterflies. *Molecular Ecology Resources, 19(6)*, 1623-1636. https://doi.org/10.1111/1755-0998.13059

Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods, 9(8)*, 772-772. https://doi.org/10.1038/nmeth.2109

Darwall, W., Bremerich, V., De Wever, A., Dell, A. I., Freyhof, J., Gessner, M. O., … & Weyl, O. (2018). The Alliance for Freshwater Life: A global call to unite efforts for freshwater biodiversity science and conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems, 28*, 1015–1022. https://doi.org/10.1002/aqc.2958

757 Dell'Anna, L., Utzeri, C., & De Matthaeis, E. (1996). Biological differentiation and
758 reproductive isolation of syntopic central Italian populations of *Chalcolestes viridis* (Vander
759 L.) and *C. parvidens* (Artobol.). *Notulae Odonatologicae, 4(8)*, 135–136.
760
761 DeSalle, R., & Goldstein, P.Z. (2019). Review and Interpretation of trends in DNA barcoding.
762 *Frontiers in Ecology and Evolution, 7,* 302. https://doi.org/10.3389/fevo.2019.00302
763
764 Dijkstra, K.-D. B., & Lewington, R. (2006). Field Guide to the Dragonflies of Britain and
765 Europe. British Wildlife Publishing.
766
767 Dijkstra, K. D., & Kalkman, V. J. (2012). Phylogeny, classification and taxonomy of European
768 dragonflies and damselflies (Odonata): a review. *Organisms Diversity & Evolution, 12(3)*, 209-
769 227. https://doi.org/10.1007/s13127-012-0080-8
770
771 Dijkstra, K.D.B., Kalkman, V.J., Dow, R.A., Stokvis, F. R., & Van Tol, J. A. N. (2014).
772 Redefining the damselfly families: a comprehensive molecular phylogeny of Zygoptera
773 (Odonata). *Systematic Entomology, 39(1)*, 68-96. https://doi.org/10.1111/syen.12035
774
775 Drummond, A. J., Suchard, M. A., Xie, D., & Rambaut, A. (2012). Bayesian phylogenetics
776 with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution, 29(8)*, 1969-1973.
777 https://doi.org/10.1093/molbev/mss075
778
779 Dufresnes, C., Mazepa, G., Jablonski, D., Oliveira, R. C., Wenseleers, T., Shabanov, D. A., ...
780 & Mulder, K. P. (2019). Fifteen shades of green: the evolution of Bufotes toads revisited.
781 *Molecular Phylogenetics and Evolution, 141*, 106615.
782 https://doi.org/10.1016/j.ympev.2019.106615
783
784 Dufresnes, C., Nicieza, A. G., Litvinchuk, S. N., Rodrigues, N., Jeffries, D. L., Vences, M., ...
785 & Martínez-Solano, Í. (2020). Are glacial refugia hotspots of speciation and cyto-nuclear
786 discordances? Answers from the genomic phylogeography of Spanish common frogs.
787 *Molecular Ecology, 29*, 986– 1000. https://doi.org/10.1111/mec.15368
788
Dumont, H. J., J. Mertens, & W. De Coster 1993. The *Calopteryx splendens*-cline in
southwestern France, analyzed by quantitative wingspot analysis (Zygoptera: Calopterygidae).
Odonatologica 22, 345– 351.
Dumont, H. J., Schneider, W., Verschuren, D., & Pavesi, M. (1995). *Coenagrion lindenii zernyi*
(Schmidt): an east Mediterranean taxon threatened with extinction by introgression from the
west (Zygoptera: Coenagrionidae). *Odonatologica, 24(4)*, 451–459.
Eberle, J., Ahrens, D., Mayer, C., Niehuis, O., & Misof, B. (2020). A plea for standardized
nuclear markers in metazoan DNA taxonomy. *Trends in Ecology & Evolution*, *35*(4), 336-345.
https://doi.org/10.1016/j.tree.2019.12.003
798
799 Ezard, T., Tomochika F., & Barraclough, T. G (2009). Splits: species' limits by threshold
800 statistics. R package.
801

Ferreira, S., Lorenzo-Carballa, M. O., Torres-Cambas, Y., Cordero-Rivera, A., Thompson, D. J., & Watts, P. C. (2014a). New EPIC nuclear DNA sequence markers to improve the resolution of phylogeographic studies of Coenagrionids and other odonates. *International Journal of Odonatology, 17(2-3)*, 135-147. https://doi.org/10.1080/13887890.2014.950698

Ferreira, S., Velo-Antón, G., Brochard, C., Vieira, C., Alves, P. C., Thompson, D. J., ... & Brito, J. C. (2014b). A Critically Endangered new dragonfly species from Morocco: *Onychogomphus boudoti* sp. nov.(Odonata: Gomphidae). *Zootaxa*, 3856(3), 349-365. http://dx.doi.org/10.11646/zootaxa.3856.3.3

Ferreira, S., Boudot, J. P., El Haissoufi, M., Alves, P. C., Thompson, D. J., Brito, J. C., & Watts, P. C. (2016). Genetic distinctiveness of the damselfly *Coenagrion puella* in North Africa: an overlooked and endangered taxon. *Conservation Genetics, 17(4)*, 985-991. https://doi.org/10.1007/s10592-016-0826-5

Fleck, G., Grand, D., & Boudot, J. P. (2007). Description of the last stadium larva of *Somatochlora borisi*, with comparison to that of somatochlora metallica meridionalis (Odonata: Corduliidae). *International Journal of Odonatology, 10(1)*, 43–52. https://doi.org/10.1080/13887890.2007.9748287

Fontaneto, D., Flot, J. F., & Tang, C. Q. (2015). Guidelines for DNA taxonomy, with a focus on the meiofauna. *Marine Biodiversity, 45(3)*, 433-451. https://doi.org/10.1007/s12526-015-0319-7

Fujisawa, T., & Barraclough, T. G. (2013). Delimiting species using single-locus data and the Generalized Mixed Yule Coalescent approach: a revised method and evaluation on simulated data sets. *Systematic Biology, 62(5),* 707-724. https://doi.org/10.1093/sysbio/syt033

Galimberti, A., Spada, M., Russo, D., Mucedda, M., Agnelli, P., Crottini, A., ... & Casiraghi, M. (2012). Integrated operational taxonomic units (IOTUs) in echolocating bats: a bridge between molecular and traditional taxonomy. *PloS one, 7(6)*. https://doi.org/10.1371/journal.pone.0040122

Gaytán, Á., Bergsten, J., Canelo, T., Pérez-Izquierdo, C., Santoro, M., & Bonal, R. (2020). DNA Barcoding and geographical scale effect: the problems of undersampling genetic diversity hotspots. *Authorea.* March 17, 2020. https://doi.org/10.22541/au.158447942.26442860

Gyulavari, H. A., Felföldi, T., Benken, T., József, L., Miskolczi, M., Cserháti, C., … Dévai, G. (2011). Morphometric and molecular studies on the populations of the damselflies *Chalcolestes viridis* and *C . parvidens* (Odonata, Lestidae). *International Journal of Odonatology, 14*, 329–339. https://doi.org/10.1080/13887890.2011.651983

Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series, 41*, 95-98.

Hebert, P. D., Cywinska, A., Ball, S. L., & Dewaard, J. R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1512), 313-321. https://doi.org/10.1098/rspb.2002.2218

849
850 Heiser, M, & Schmitt, T. (2010). Do different dispersal capacities influence the biogeography
851 of the western Palearctic dragonflies (Odonata)? *Biological Journal of the Linnean Society*, 99,
852 177–195. https://doi.org/10.1111/j.1095-8312.2009.01349.x
853
854 Heiser, M., Dapporto, L., & Schmitt, T. (2014). Coupling impoverishment analysis and
855 partitioning of beta diversity allows a comprehensive description of Odonata biogeography in
856 the Western Mediterranean. *Organisms Diversity and Evolution*, 14(2), 203–214.
857 https://doi.org/10.1007/s13127-013-0161-3
858
859 Hewitt, G. M. (2004). Genetic consequences of climatic oscillations in the Quaternary.
860 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 359(1442), 183–195.
861 https://dx.doi.org/10.1098%2Frstb.2003.1388
862
863 Husemann, M., Schmitt, T., Zachos, F. E., Ulrich, W., & Habel, J. C. (2014). Palaearctic
864 biogeography revisited: evidence for the existence of a North African refugium for Western
865 Palaearctic biota. *Journal of Biogeography*, 41(1), 81–94. https://doi.org/10.1111/jbi.12180
866
867 Jacquemin, G., & Boudot, J. P. (1990) A propos de Coenagrion mercuriale (Charpentier, 1840)
868 au Maroc (Zygoptera: Coenagrionidae). Notulae Odonatologicae, *3*: 91-94.
869
870 Jacquemin, G., & Boudot, J. P. (1999) Les libellules (odonates) du Maroc. Société française
871 d'odonatologie.
872
873 Kajtoch, Ł., Montagna, M., & Wanat, M. (2018). Species delimitation within the
874 *Bothryorrhynchapion* weevils: multiple evidence from genetics, morphology and ecological
875 associations. *Molecular Phylogenetics and Evolution, 120*, 354-363.
876 https://doi.org/10.1016/j.ympev.2017.12.022
877
878 Kalkman, V. J., Boudot, J. P., Bernard, R., De Knijf, G., Suhling, F., & Termaat, T. (2018).
879 Diversity and conservation of European dragonflies and damselflies (Odonata). *Hydrobiologia,*
880 *811(1)*, 269-282. https://doi.org/10.1007/s10750-017-3495-6
881
882 Kapli, P., Lutteropp, S., Zhang, J., Kobert, K., Pavlidis, P., Stamatakis, A., & Flouri, T. (2017).
883 Multi-rate Poisson tree processes for single-locus species delimitation under maximum
884 likelihood and Markov chain Monte Carlo. *Bioinformatics, 33(11)*, 1630-1638.
885 https://doi.org/10.1093/bioinformatics/btx025
886
887 Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version
888 7: improvements in performance and usability. *Molecular Biology and Evolution, 30(4)*, 772-
889 780. https://doi.org/10.1093/molbev/mst010
890
891 Kim, M. J., Jung, K. S., Park, N. S., Wan, X., Kim, K. G., Jun, J., ... & Kim, I. (2014). Molecular
892 phylogeny of the higher taxa of Odonata (Insecta) inferred from COI, 16S r RNA, 28S r RNA,
893 and EF 1-α sequences. *Entomological Research*, 44(2), 65-79. https://doi.org/10.1111/1748-
894 5967.12051

Koroiva, R., Pepinelli, M., Rodrigues, M. E., de Oliveira Roque, F., Lorenz-Lemke, A. P., & Kvist, S. (2017). DNA barcoding of odonates from the Upper Plata basin: Database creation and genetic diversity estimation. *PloS one, 12(8)*. https://doi.org/10.1371/journal.pone.0182283

Koroiva, R., & Kvist, S. (2018). Estimating the barcoding gap in a global dataset of cox1 sequences for Odonata: close, but no cigar. *Mitochondrial DNA Part A, 29(5)*, 765-771. https://doi.org/10.1080/24701394.2017.1357709

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution, 35(6)*, 1547-1549. https://doi.org/10.1093/molbev/msy096

Leigh, J. W., & Bryant, D. (2015). popart: full-feature software for haplotype network construction. *Methods in Ecology and Evolution, 6(9)*, 1110-1116. https://doi.org/10.1111/2041-210X.12410

Lieftinck, M.A. (1966) A survey of the dragonfly fauna of Morocco (Odonata). *Bulletin et annales de la Société royale belge d'entomologie, 42*, 1-63.

Lohse, K. (2009). Can mtDNA barcodes be used to delimit species? A response to Pons et al.(2006). *Systematic Biology*, 58(4), 439-442. https://doi.org/10.1093/sysbio/syp039

López-Estrada, E. K., Fernández, J. B., Cardo-Maeso, N., Montejano, S. T., & Díaz-Martínez, C. (2020). *Onychogomphus cazuma* sp. nov. from Spain: Molecular and morphological evidence supports the discovery of a new European dragonfly species (Odonata: Gomphidae). *Odonatologica*, 49(1/2), 125-154..ì

Lunghi E., Corti C., Manenti R. & Ficetola G.F. (2019). Consider species specialism when publishing datasets. *Nature Ecology & Evolution 3*, 319. https://doi.org/10.1038/s41559-019-0803-8

Marinov, M. (2001). Somatochlora borisi spec. nov., a new European dragonfly species from Bulgaria (Anisoptera: Corduliidae). *Internation Dragonfly Found report, 3(1/2)*, 9-16.

Matos-Maraví, P., Wahlberg, N., Antonelli, A., & Penz, C. M. (2019). Species limits in butterflies (Lepidoptera: Nymphalidae): Reconciling classical taxonomy with the multispecies coalescent. *Systematic Entomology, 44(4)*, 745-756. https://doi.org/10.1111/syen.12352

Mazzamuto, M. V., Galimberti, A., Cremonesi, G., Pisanu, B., Chapuis, J. L., Stuyck, J., ... & Wauters, L. A. (2016). Preventing species invasion: A role for integrative taxonomy? *Integrative Zoology, 11(3)*, 214-228. https://doi.org/10.1111/1749-4877.12185

Meyer, C. & Paulay, G. (2005) DNA barcoding: error rates based on comprehensive sampling. *PLOS Biology, 3,* e422. https://doi.org/10.1371/journal.pbio.0030422

Miller, M. A., Pfeiffer, W., & Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In 2010 gateway computing environments workshop (GCE) (pp. 1-8). Ieee.

Morinière, J., Balke, M., Doczkal, D., Geiger, M. F., Hardulak, L. A., Haszprunar, G., ... &
Schmidt, S. (2019). A DNA barcode library for 5,200 German flies and midges (Insecta:
Diptera) and its implications for metabarcoding-based biomonitoring. *Molecular Ecology*
Resources, 19(4), 900-928. https://doi.org/10.1111/1755-0998.13022

Navas, L., 1907. Neuropteros de Espana y Portugal [Odonata]. *Broteria (Zool.), 6*, 43-58.

Nielsen, C. (1935). Note odonatologiche. *Bolletino Società Entomologica Italiana, 67*, 59-62.

Olias, M., Weihrauch, F., Bedjanič, M., Hacet, N., Marinov, M., & Salamun, A. (2007). *Lestes*
parvidens and *L. viridis* in southeastern Europe: a chorological analysis (Odonata: Lestidae).
Libellula, 26(3/4), 243–247.

Ottenburghs, J. (2020). Ghost Introgression: Spooky Gene Flow in the Distant Past. *BioEssays*,
42(6), 2000012. https://doi.org/10.1002/bies.202000012

Padial, J. M., Miralles, A., De la Riva, I., & Vences, M. (2010). The integrative future of
taxonomy. *Frontiers in zoology, 7(1)*, 16. https://doi.org/10.1186/1742-9994-7-16

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution
in R language. *Bioinformatics*, 20(2), 289-290. https://doi.org/10.1093/bioinformatics/btg412

Pawlowski, J., Kelly-Quinn, M., Altermatt, F., Apothéloz-Perret-Gentil, L., Beja, P., Boggero,
A., ... & Feio, M. J. (2018). The future of biotic indices in the ecogenomic era: Integrating (e)
DNA metabarcoding in biological assessment of aquatic ecosystems. *Science of the Total*
Environment, 637, 1295-1310. https://doi.org/10.1016/j.scitotenv.2018.05.002

Petit, R. J., & Excoffier, L. (2009). Gene flow and species delimitation. *Trends in Ecology &*
evolution, 24(7), 386-393. https://doi.org/10.1016/j.tree.2009.02.011

Pinkert, S., Dijkstra, K. D. B., Zeuss, D., Reudenbach, C., Brandl, R., & Hof, C. (2018).
Evolutionary processes, dispersal limitation and climatic history shape current diversity
patterns of European dragonflies. *Ecography, 41(5)*, 795-804.
https://doi.org/10.1111/ecog.03137

Piper, A. M., Batovska, J., Cogan, N. O., Weiss, J., Cunningham, J. P., Rodoni, B. C., &
Blacket, M. J. (2019). Prospects and challenges of implementing DNA metabarcoding for high-
throughput insect surveillance. *GigaScience, 8(8)*, giz092.
https://doi.org/10.1093/gigascience/giz092

Pons, J., Barraclough, T. G., Gomez-Zurita, J., Cardoso, A., Duran, D. P., Hazell, S., ... &
Vogler, A. P. (2006). Sequence-based species delimitation for the DNA taxonomy of
undescribed insects. *Systematic Biology, 55(4)*, 595-609.
https://doi.org/10.1080/10635150600852011

Porter, T. M., & Hajibabaei, M. (2018). Over 2.5 million COI sequences in GenBank and
growing. *PloS One, 13(9)*. https://doi.org/10.1371/journal.pone.0200177

Puillandre, N., Lambert, A., Brouillet, S., & Achaz, G. (2012). ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology, 21(8)*, 1864-1877. https://doi.org/10.1111/j.1365-294x.2011.05239.x

R Core Team (2019). R: A language and environment for statistical computing. Available from http://www.R-project.org.

Rambaut, A., 2012. FigTree v1.4.2 [online]. A Graphical Viewer of Phylogenetic Trees. Available from http://tree.bio.ed.ac.uk/software/figtree

Rambaut, A., Suchard MA, Drummond AJ. 2014. Tracer v1.6 [online]. Available from http://beast.bio.ed.ac.uk/Tracer

Ratnasingham, S., & Hebert, P. D. (2007). BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). *Molecular Ecology Notes, 7(3)*, 355-364. https://dx.doi.org/10.1111%2Fj.1471-8286.2007.01678.x

Ratnasingham, S., & Hebert, P.D. (2013). A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PloS One, 8(7).* https://doi.org/10.1371/journal.pone.0066213

Reid, N. M., & Carstens, B. C. (2012). Phylogenetic estimation error can decrease the accuracy of species delimitation: a Bayesian implementation of the general mixed Yule-coalescent model. *BMC Evolutionary Biology*, 12(1), 196. https://doi.org/10.1186/1471-2148-12-196

Rewicz, T., Móra, A., Tończyk, G., Szymczak, A., Grabowski, M., Calleja, E. J., & Csabai, Z. (2020). First records raise questions: DNA barcoding of Odonata in the middle of the Mediterranean. *Genome*. https://doi.org/10.1139/gen-2019-0226

Riservato, E., Festi, A., Fabbri, R., Grieco, C., Hardersen, S., La Porta, G., … Utzeri, C. (2014a). Odonata- Atlante delle libellule italiana -preliminare. Società per lo Studio e la Conservazione delle Libellule. Latina: Edizioni Belvedere.

Riservato, E., Fabbri, R., Festi, A., Grieco, C., Hardersen, S., Landi, F., … Utzeri, C. (2014b). Lista rossa IUCN delle Libellule Italiane. Roma: Comitato italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare.

Roberts, J. E. H. (1948). *Coenagrion castellani*, a new species of dragonfly in Europe (order Odonata, Zygoptera). *Proceedings of the Royal Entomological Society of London (B)*, 17: 63-66.

Sahlén, G., Kalkman, V. J., Boudot, J. P., Bernard, R., Conze, K. J., De Knijf, G., ... & Riservato, E. (2010). *European red list of dragonflies*. Publications Office of the European Union.

Sánchez-Guillén, R. A., Córdoba-Aguilar, A., Cordero-Rivera, A., & Wellenreuther, M. (2014a). Genetic divergence predicts reproductive isolation in damselflies. *Journal of Evolutionary Biology, 27(1)*, 76-87. https://doi.org/10.1111/jeb.12274

Sánchez-Guillén, R. A., Córdoba-Aguilar, A., Cordero-Rivera, A., & Wellenreuther, M. (2014b). Rapid evolution of prezygotic barriers in non-territorial damselflies. *Biological Journal of the Linnean Society, 113(2)*, 485-496. https://doi.org/10.1111/bij.12347

Scalercio, S., Cini, A., Menchetti, M., Vodă, R., Bonelli, S., Bordoni, A., ... & Dapporto, L. (2020). How long is 3 km for a butterfly? Ecological constraints and functional traits explain high mitochondrial genetic diversity between Sicily and the Italian Peninsula. *Journal of Animal Ecology*. https://doi.org/10.1111/1365-2656.13196

Schmidt, E. (1957). Was ist *Somatochlora sibirica* Trybom. *Beiträge zur naturkundlichen Forschung in Südwestdeutschland, 16*, 92-100.

Schmid-Egger, C., Straka, J., Ljubomirov, T., Blagoev, G. A., Morinière, J., & Schmidt, S. (2019). DNA barcodes identify 99 per cent of apoid wasp species (Hymenoptera: Ampulicidae, Crabronidae, Sphecidae) from the Western Palearctic. *Molecular Ecology Resources, 19(2)*, 476-484. https://doi.org/10.1111/1755-0998.12963

Schmitt, T., & Varga, Z. (2012). Extra-Mediterranean refugia: The rule and not the exception? *Frontiers in Zoology*, 9(1), 22. https://doi.org/10.1186/1742-9994-9-22

Schorr, M., & Paulson, D. (2019). World Odonata List. Electronic Database accessible at https://www.pugetsound.edu/academics/academic-resources/slater-museum/biodiversity-resources/dragonflies/world-odonata-list2/. Accessed, February 20, 2020.

Seidenbusch, R. (1996). Description of the last instar larva of *Somatochlora meridionalis* Nielsen, 1935 (Anisoptera: Corduliidae). *Odonatologica, 25(3)*, 303–306.

Solano, E., Hardersen, S., Audisio, P., Amorosi, V., Senczuk, G., & Antonini, G. (2018). Asymmetric hybridization in Cordulegaster (Odonata: Cordulegastridae): Secondary postglacial contact and the possible role of mechanical constraints. *Ecology and Evolution, 8(19)*, 9657-9671. https://dx.doi.org/10.1002%2Fece3.4368

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics, 30(9)*, 1312-1313. https://doi.org/10.1093/bioinformatics/btu033

Steinmann H., 2013. World Catalogue of Odonata II. Anisoptera. De Gruiter

Swaegers, J., Mergeay, J., Van Geystelen, A., Therry, L., Larmuseau, M. H. D., & Stoks, R. (2015). Neutral and adaptive genomic signatures of rapid poleward range expansion. *Molecular Ecology, 24(24)*, 6163-6176. https://doi.org/10.1111/mec.13462

Viganò, M., Janni, O. & Corso, A. (2017). *Tramea basilaris* on Linosa Island, Italy: A new species for Europe and the Western Palaeartic (Odonata: Libellulidae). *Odonatologica, 46(1/2)*, 55-56

1075 Wauters, L. A., Amori, G., Aloise, G., Gippoliti, S., Agnelli, P., Galimberti, A., ... & Martinoli,
1076 A. (2017). New endemic mammal species for Europe: *Sciurus meridionalis* (Rodentia,
1077 Sciuridae). *Hystrix, 28(1)*: 1-8. https://doi.org/10.4404/hystrix-28.1-12015
1078
Weekers, P. H., De Jonckheere, J. F., & Dumont, H. J. (2001). Phylogenetic relationships
inferred from ribosomal ITS sequences and biogeographic patterns in representatives of the
genus *Calopteryx* (Insecta: Odonata) of the West Mediterranean and adjacent West European
zone. *Molecular Phylogenetics and Evolution, 20(1)*, 89-99.
https://doi.org/10.1006/mpev.2001.0947
Weekers, P. H. H., & Dumont, H. J. (2004). A molecular study of the relationship between the
Coenagrionid genera Erythromma and Cercion, with the creation of Paracercion Gen. Nov. for
the East asiatic "*Cercion*" (Zygoptera: Coenagrionidae). *Odonatologica, 33(2)*, 181–188.
1087
1088 Weigand, H., Beermann, A. J., Ciampor, F., Costa, F. O., Csabai, Z., Duarte, S., ... & Strand,
1089 M. (2019). DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-
1090 analysis and recommendations for future work. *Science of the Total Environment, 678*, 499-
1091 524. https://doi.org/10.1016/j.scitotenv.2019.04.247
1092
1093 Wellenreuther, M., & Sánchez-Guillén, R. A. (2016). Nonadaptive radiation in damselflies.
1094 *Evolutionary Applications, 9(1)*, 103-118. https://doi.org/10.1111/eva.12269
1095
1096 Zhang, J., Kapli, P., Pavlidis, P., & Stamatakis, A. (2013). A general species delimitation
1097 method with applications to phylogenetic placements. *Bioinformatics, 29(22)*, 2869-2876.
1098 https://doi.org/10.1093/bioinformatics/btt499
1099
1100

Data Accessibility:
All specimen data are accessible on BOLD (www.boldsystems.org) under the project name 'ZPLOD - DNA barcoding Italian Odonata'. The data include collection locality, geographic coordinates, collector, identifier, voucher depository, BINs and other sampling details. Sequence data are available on BOLD and have also been deposited in GenBank (Accession numbers MT298232 - MT298679).

Author contributions
AG and GA conceived the idea and designed the sampling protocol, which was then developed by all authors. AG, GA, DB, GB, IC, AC, VF, MG, LI, GLP, LL, FL, FM, SR, RS, GS, SS conducted the field sampling. FR with the help of AG stored the biological tissues, performed the lab analyses, and mined data from BOLD and GenBank. AG, GA, and DM ran the statistical analyses and all authors contributed to analyses checking/development and check the final model outcomes. AG, GA and DM lead the writing of a first draft of the manuscript and all authors critically contributed revising the final version.

Tables

TABLE 1: Species delimitation results. The table provides the number of groups delimited by
each distance-based (ABGD and OT) and tree-based (PTP, MPTP, GMYC and bGMYC)
approach conducted on the DNA barcode sequences of Italian odonate morphospecies grouped
by suborder (Zygoptera and Anisoptera) and by DNA barcoding dataset (DS1 and DS2). The
number of overall delimited groups, groups for each warning category and involved species
(within brackets) are also reported.

1126

			Species delimitation approaches and n of delimited groups							Warnings: n of cases (n of involved species)			
		n morphospecies	ABGD	OT	PTP	MPTP	GMYC	bGMYC	Overall n of groups	no warnings	no interspecific delimitation	intraspecific delimitation	mixed species delimitation
Zygoptera	DS1	31	26	28	34	30	30	30	28	21	3(7)	2(1)	2(2)
	DS2	31	32	31	33	32	36	40	32	19	2(4)	7(3)	4(5)
Anisoptera	DS1	57	55	56	59	55	61	58	56	52	2(4)	2(1)	0
	DS2	57	57	62	63	55	59	63	61	48	2(4)	11(5)	0

1137 **TABLE 2:** List of odonate species showing warnings as a result of species delimitation on DS2.
1138 The warning category (as in Table 1), the number of delimited groups and the geographic
1139 overall provenance of DNA barcode sequences included in DS2 for each species group are also
1140 reported.

1141

Suborder	Involved morphospecies	warning category	n of delimited groups	Geographic scale	Also occurring in DS1?
Zygoptera	*Coenagrion mercuriale*	intraspecific delimitation	3	Palearctic	no
Zygoptera	*Coenagrion puella / Coenagrion pulchellum / Coenagrion ornatum*	mixed species delimitation	2	Palearctic	no
Zygoptera	*Erythromma lindenii*	intraspecific delimitation	2	Palearctic	yes
Zygoptera	*Ischnura genei/ Ischnura elegans*	no interspecific delimitation	1	Palearctic	yes
Zygoptera	*Calopteryx splendens / Calopteryx xanthostoma*	no interspecific delimitation	1	Palearctic	yes
Zygoptera	*Lestes dryas*	intraspecific delimitation	2	Holarctic	no
Zygoptera	*Chalcolestes viridis / Calcholestes parvidens*	mixed species delimitation	2	Palearctic	yes
Anisoptera	*Aeshna juncea*	intraspecific delimitation	2	Holarctic	no
Anisoptera	*Anax imperator / Anax parthenope*	no interspecific delimitation	1	Palearctic	yes
Anisoptera	*Onychogomphus forcipatus*	intraspecific delimitation	2	Palearctic	yes
Anisoptera	*Stylurus flavipes*	intraspecific delimitation	2	Palearctic	no
Anisoptera	*Orthetrum albistylum*	intraspecific delimitation	3	Palearctic	no
Anisoptera	*Sympetrum danae*	intraspecific delimitation	2	Holarctic	no
Anisoptera	*Somatochlora metallica / Somatochlora meridionalis*	no interspecific delimitation	1	Palearctic	yes

1142

 Figure captions

1144

FIGURE 1: Collection sites for the sampled 88 odonate species included in the present study.
The left map shows the number of specimens and the number of collected species per site
indicated by circle size and colour respectively. The right map shows the borders of the macro-
regions used for the analysis of genetic distances and structure of each species. Details of the
328 sites and collected and DNA barcoded specimens are provided in Appendix1. This map
was created in QGIS version 3.4.13 (https://www.qgis.org/it/site/).

1151

FIGURE 2: Multi-approach species delimitation of the 31 Italian Zygoptera morphospecies
analyzed in this study, based on COI DNA barcode sequences from DS1. A Bayesian tree is
used as a base to summarize the two threshold-based (OT and ABGD) and four character
explicit-based (PTP, MPTP, GMYC, bGMYC) approaches. Specimen haplotype identifiers are
reported on tips (see Appendix S1 for further details), the number of specimens sharing each
haplotype is reported within brackets and abbreviations NW, NE, C, S, SI and SA correspond
to the geographic haplotype origin as schematized in Figure 1 (ITA corresponds to unknown
locality in Italy). Numbers above nodes represent BPP and BS, respectively, whereas asterisks
correspond to maximal node support (BPP $\geq$ 0.99 and BS $\geq$ 95). Vertical colored solid boxes
delimit putative species identified by the different approaches. Black: delimitation congruent
with the identified morphospecies; Red: intraspecific delimitation; Light Blue: no interspecific
delimitation; Yellow: Mixed species delimitation. In the case of intraspecific delimitation, each
lineage has been numbered with Roman numerals. This tree was created with BEAST 1.8.2.

1165

FIGURE 3: Multi-approach species delimitation of the 57 Italian Anisoptera morphospecies
analyzed in this study, based on COI DNA barcode sequences from DS1. A Bayesian tree is
used as a base to summarize the two threshold-based (OT and ABGD) and four character
explicit-based (PTP, MPTP, GMYC, bGMYC) approaches. Specimen haplotype identifiers are
reported on tips (see Appendix S1 for further details), the number of specimens sharing each
haplotype is reported within brackets and abbreviations NW, NE, C, S, SI and SA correspond
to the geographic haplotype origin as schematized in Figure 1. Numbers above nodes represent
Bayesian posterior probabilities (BPP) and maximum likelihood bootstrap values (BS),
respectively, whereas asterisks correspond to maximal node support (BPP $\geq$ 0.99 and BS $\geq$ 95).
Vertical colored solid boxes delimit putative species identified by the different approaches.
Black: delimitation congruent with the identified morphospecies; Red: intraspecific
delimitation; Light Blue: no interspecific delimitation. In the case of intraspecific delimitation,
each lineage has been numbered with Roman numerals. This tree was created with BEAST
1.8.2.

1180

FIGURE 4: Median-joining network of COI DNA barcode haplotypes of four warning cases
found by species delimitation of DS2 (*Coenagrion mercuriale*, *Lestes dryas*, *Aeshna juncea*
and *Sympetrum danae*; see Appendix S2 and S6 for haplotypes and lineages subdivision). Each
circle represents a haplotype and circle size is proportional to haplotype frequency. Colours
indicate different sampling regions. Small black traits represent possible median vectors, while
dashes represent substitutions (also indicated within brackets when > 10). Networks have been
created with PopArt 1.7.

1188

1189

Supporting information

APPENDIX S1: dataset DS1, composed by the DNA barcoding sequences produced in this study on Italian samples (409) plus 69 GenBank sequences from other Italian specimens. Species taxonomy, sex, specimen IDs, depositories accessions, sampling geographic details and COI haplotype are reported for each DNA barcoded sample.

APPENDIX S2: dataset DS2, composed by the sequences mined from BOLD and GenBank plus dataset DS1 and other COI barcode sequences obtained from additional samples retrieved by authors in other western Palearctic countries. Species taxonomy, DS1 specimen IDs, depositories accessions, are reported for each COI haplotype.

APPENDIX S3: Support values of the species delimitation hypotheses obtained with the Bayesian implementation of the PTP and GMYC methods, for both DS1 and DS2 (n.d.: not detected in DS1).

APPENDIX S4: Distributions of pairwise genetic distances (K2P) obtained with ABGD for a) Zygoptera DS1, b) Anisoptera DS1, c) Zygoptera DS2, and d) Anisoptera DS2.

APPENDIX S5: Barcode Gap Analysis of DS1 for Anisoptera and Zygoptera respectively, generated by BOLD. Three scatterplots are provided to confirm the existence and magnitude of the Barcode Gap. For each suborder, the first two scatterplots show the overlap of the max and mean intra-specific distances vs the inter-specific (nearest neighbour) distances. The third scatterplot plots the number of individuals in each species against their max intra-specific distances, as a test for sampling bias. Distance Model: Kimura 2 Parameter, Deletion Method: Pairwise Deletion, Alignment: BOLD Aligner (Amino Acid based HMM), Filters Applied: ≥ 500 bp only.

APPENDIX S6: Multi-approach species delimitation of the 31 Zygoptera and 57 Anisoptera species investigated in this study based on Holarctic COI DNA barcode sequences from DS2. A Bayesian tree is used as a base to summarize the two threshold-based (OT and ABGD) and four character explicit-based (PTP, MPTP, GMYC, bGMYC) approaches. Specimen haplotype identifiers are reported on tips (see Appendix S1 and S2 for further details) and the number of specimens sharing each haplotype is reported within brackets. Numbers above nodes represent BPP and BS, respectively, whereas asterisks correspond to maximal node support (BPP ≥ 0.99 and BS ≥ 95). Vertical colored solid boxes delimit putative species identified by the different approaches. Black: delimitation congruent with the identified morphospecies; Red: intraspecific delimitation; Light Blue: no interspecific delimitation; Yellow: Mixed species delimitation. In case of intraspecific delimitation, each lineage has been numbered with Roman numerals. This tree was created with BEAST 1.8.2.

APPENDIX S7: average genetic p-distance divergence and Standard Deviation values among the geographic population of *Coenagrion mercuriale* (ITA: Italy, EUR: Europe excluding Italy, NAF: North Africa) and among three species belonging to *Coenagrion* (ORN: *ornatum*, PUL: *pulchellum*, PUE: *puella*) calculated at two mitochondrial (16s rDNA and COI) and three nuclear (AgT, PRMT and MLC) markers. Sequence data were retrieved from GenBank. Concerning COI, also the sequences produced in this study were used to calculate the genetic distances. The exclusiveness of haplotypes belonging to the Italian populations of *C. mercuriale*

1238 and to the other three *Coenagrion* species is indicated for each genetic marker. Genetic distance
1239 values have been calculated using MEGA X (Kumar et al., 2018).

Coenagrion mercuriale

Lestes dryas

Aeshna juncea

Sympetrum danae

Italy   Europe (not Italy)   North Africa   North America