Differences in visually induced MEG oscillations reflect differences in deep cortical layer activity

D. A. Pinotsis[1,2,*], E. K. Miller[2]

[1]Centre for Mathematical Neuroscience and Psychology and Department of Psychology, City —University of London, London EC1V 0HB, United Kingdom

[2]The Picower Institute for Learning & Memory and Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

*pinotsis@mit.edu

Abstract

Neural activity is organized at multiple scales, ranging from the cellular to the whole brain level. Connecting neural dynamics at different scales is important for understanding brain pathology. Neurological diseases and disorders arise from interactions between factors that are expressed in multiple scales. Here, we suggest a new way to link microscopic and macroscopic dynamics through combinations of computational models. This exploits results from statistical decision theory and Bayesian inference. To validate our approach, we used two independent MEG datasets. In both, we found that variability in visually induced oscillations recorded from different people in simple visual perception tasks resulted from differences in the level of inhibition specific to deep cortical layers. This suggests differences in feedback to sensory areas and each subject's hypotheses about sensations due to differences in their prior experience. Our approach provides a new link between non-invasive brain imaging data, laminar dynamics and top-down control.

Introduction

A major challenge in treating neurological diseases and disorders is the their heterogeneity [1]. The same symptom can arise from multiple causes and the same cause can lead to various symptoms. Only a small fraction of patients might respond to a certain drug, which leads to frustration for both patients and clinicians. One way to address this challenge is to build computational models of brain dysfunction [2]. This approach has given rise to a new field, computational psychiatry [3]. Instead of focusing on symptomatology, computational models focus on describing brain neurobiology and its alterations in patient brains. However, computational models are not a *panacea:* they are limited by the spatial and temporal scale of the dynamics they describe. Spiking and compartmental models describe single neurons and firing rates while mean field models describe large brain networks and population activities. Thus, they are limited because neurological diseases and disorders involve interactions of many factors that are *simultaneously* expressed at multiple scales. They depend on both genetic variations and environmental factors spanning microscopic and macroscopic scales ranging from, for example, altered mitochondria and single neuron function to neuroinflammation of axons

connecting different brain areas [4]. Here, we present a new way to link multiple scales. We use combinations of computational models to describe both macro and micro scales.

We combine models of neural compartments, like dendrites and somata [5] with biophysical mean field models that predict non-invasive brain data [6]. We used statistical decision theory [7] to prove that the these two model classes can be combined to use non-invasive MEG data to infer neural dynamics at the level of cortical layers (laminar dynamics). In general, statistical decision theory prescribes the optimal way of using quantitative tools to make statistical decisions. It quantifies performance of these tools and what an optimal decision is. We found that in the context of statistical decision theory, compartmental and mean field models can also be thought of as quantitative tools (besides other examples that mathematicians have considered as tools so far). Then, estimating neurobiological parameters of both compartmental and mean field models is the same as making an optimal decision at different scales. After realizing this, we used insights from statistical decision theory to estimate biophysically accurate parameter sets that describe neural dynamics at both the macro and micro scales.

In [8] and [9], we used a similar combination of computational models to analyze invasive animal data. However, we did not provide a mathematical proof of why such a combination can be considered and focused on invasive electrophysiology. Here, we extend this work to non-invasive human data, consider brain activity measured with MEG, and give a proof that establishes the mathematical basis of our approach. We also consider between subject differences and quantify the neurobiological mechanisms that underlie variability in human MEG data. Together with [8] and [9], the results of the present paper suggest a general two step approach for understanding dynamics and neurobiology at the microscale using non-invasive human brain data. Step one: Construct a compartmental model that captures biophysical properties of single neurons (e.g., the geometry of the dendritic tree, kinetics and densities of ion channels, inputs from subcortical areas etc). Test it with intracranial recordings from non-human primates. Step two: Tune a mean field model to give the same predictions as the compartmental model. Test it with human M/EEG data.

Below, we demonstrate our approach using two independent human MEG datasets analyzed earlier in [6] and [10]. These datasets contain visually induced oscillations recorded during perception tasks. In earlier work, we had found that differences in these oscillations between different people were due to differences in the level of inhibition in the cortical source. To validate our new approach, we asked whether we could confirm those earlier results (obtained using different computational models). We also asked whether our new model of laminar dynamics could identify the cortical layer where these differences were more pronounced. Interestingly, both datasets led to the same result. Differences in visually induced oscillations between different people originated from differences in the level of inhibition in deep cortical layers. This provides new insights into the computations performed by different cortical layers.

Results

We used human MEG data to determine if we could explain individual differences in visually induced oscillations by localizing them to superficial versus deep cortical layers. We used statistical decision theory to establish that the compartmental model we used to describe laminar dynamics could make the same predictions as the mean field model fitted to human MEG data (Supplementary Materials and Methods). Then we used hierarchical Bayesian modelling to explain individual differences in single subject neurobiology and anatomy that underlie gamma oscillation variability.

Computational models of brain dynamics at the micro and macro scales predict similar laminar data

Following [8], we here used a macroscale (mean field[1]) model whose parameters had been tuned to predict similar[2] laminar data as a microscale (compartmental) model. The model predicts the same peak frequencies for superficial and deep pyramidal cells and similar power distribution across frequencies (Figure 1B). It describes a similar distribution of spatiotemporal dynamics across layers (see below). To construct this model, we adapted the macroscale model of [8] to deal with MEG as opposed to intracranial laminar data used in that earlier work. The new model predicts MEG oscillations from different cortical depths. Both macro- and microscale models describe the same microcircuit shown in Figure 1A. This includes two pairs of excitatory-inhibitory (E-I) populations, one in superficial and the other in deep layers. Each E-I population pair is connected with intralaminar connections. Superficial and deep populations are also connected with interlaminar connections. Arrows correspond to excitatory (black) and inhibitory (red) connections. Figure 1A also includes the mean field equations describing neural activity. A detailed description of the model can be found in [8].

Besides describing the same microcircuit, the macro- and microscale models predict similar laminar dynamics. The mathematical proof of the equivalence of their predictions is based on statistical decision theory and is included in Supplementary Materials and Methods. To make the two models predict similar dynamics, we developed an analysis pipeline that combines existing and validated methodologies (Figure 2). First, theoretical arguments from statistical decision theory required that the micro- scale model be adapted to have the same number of parameters as the macroscale model. We changed the compartmental model of [5], by reducing the number of its connection parameters (considering all synaptic weights equal) and called the resulting model the symmetric compartmental model [8], [9][3]. The remaining parameters are the weights of the connections between neural populations occupying different layers of the microcircuit

---

[1] The particular model used here is a so called neural mass model. This is one of the two main sub-classes of mean fields models. The other is neural fields considered in the Discussion. We here used the term "mean field" as this is associated in the literature with the macroscale.

[2] "Similar" has also a precise, mathematically rigorous, meaning. Examples of similar predictions are shown in Figure 1B and the mathematically rigourous meaning is explained in the Supplementary Materials and Methods. This is based on statistical decision theory. Briefly, the macro- and microscale models are reformulated in terms of Bayesian decision rules. Then it is shown that a Bayesian observer can not distinguish between the data predicted separately by each model. Thus, they predict "similar" data. Alternatively, this means that if one fits both models to the same data using Bayesian inference then the model fits are the same (same error or accuracy).

[3] In [7], we found that evoked responses from the original model of [5] and its symmetric variant were highly correlated, $r=0.9343$, $p < 0.001$, see the Supplementary Figure S1A.

shown in Figure 1A. These are the same parameters that the macroscale model has. Second, we simulated power spectra from the microscale (symmetric compartmental) model (green data in Figure 1B). Third, we fitted the macroscale model to these simulated power spectra using Dynamic Causal Modelling (DCM) for steady state responses [9], [11], [12]. This approach has been used to infer changes in synaptic plasticity in large cortical networks [11] in healthy and clinical populations [13] among other applications. It exploits the stationarity of variance of neuronal firing over the course of the task to efficiently fit mean field models to MEG data. After fitting, we obtained the parameter values of the macroscale model that fitted best the predictions of the microscale model. The predictions from the macroscale model for these parameter values are shown in Figure 1B in magenta. Predictions from both models are very similar. The magenta and green curves overlap for frequencies above 3 Hz for deep layers (Figure 1B bottom) and 8Hz for superficial layers (Figure 1B top). This difference in superficial layers is due to the fact that the macroscale model receives explicit white and pink noise input that propagates to all layers, while the microscopic models received Gaussian spike input targeting superficial layers.

To sum up we adapted the model of [5] and fine-tuned the parameters of a macroscale (mean field) model describing the same microcircuit so that the two models make the predictions of the same frequency peaks and similar power distribution across frequencies in superficial and deep layers. The parameter values of the two models for which this happens are included in Table 1. Parameters for the microscale model are shown in the left column while the parameters of the macroscale model in the right. These include the connection parameters between the populations shown in Figure 1. Each parameter corresponds to one arrow. There are in total ten connections in both the microscale and macroscale models.

To sum up, we fine tuned the macroscale model to predict similar laminar dynamics as the microscale model. Coupled with a mapping from laminar dynamics to MEG sensors, the microscale model has been used to *simulate* MEG oscillations in different frequency bands [5], [14]. Following our earlier work [6], [10], we then used the fine tuned macroscale model and a similar mapping[4] to *fit* MEG data and explain individual variability in human brain oscillations.

Variability of visually induced gamma oscillations from different datasets reveals differences in the level of inhibition across different people

We mapped differences in recorded power spectra from different people to differences in the macroscale model parameters and biophysics. We fitted the tuned macroscale model to oscillatory data from two different MEG datasets using DCM for steady state responses [12]. Both datasets contained visually induced gamma oscillations reported in [15], [16]. These oscillations where recorded in different subjects. The spatial extent of the V1 cortical source in each subject varied. In [16] this was the result of changing the size of the visual stimulus presented to the subjects. In [15], this was measured with retinotopic mapping. The tuned parameters described above and in Figure 2 were used as prior values for fitting the macroscale model to these MEG data.

We first fitted the macroscale model to the MEG data from [16]. These included visually induced oscillations while 12 subjects viewed grating stimuli of three different sizes— 2^0,

---

[4] This is given by $L_r(k, \varphi)$, see Methods.

$4^0 \& 8^0$. Thus the dataset from [16] included oscillations induced by three different stimulus sizes. Grand averages of model fits across all subjects for the smallest stimulus size are shown in Figure 3A. Dashed and solid lines correspond to data and model respectively. Shaded regions include 95% confidence intervals. These quantify variability across subjects and overlap for model predictions and data. Model fits to *individual* subjects and conditions are shown in Figure 3B[5].We fitted all three stimuli conditions simultaneously by modeling different sizes as condition-specific effects, see [10]. The three stimulus sizes correspond to red, blue and green lines. Dashed and solid lines correspond to data and model predictions. Differences in power spectra between conditions for the same subject are small and spectra corresponding to different stimulus sizes overlap. We fitted the data of each subject by changing the connection weights. These describe the strength of effective connections between different populations and describe the level of excitation and inhibition (E-I balance). Recall that priors for these parameters were chosen so that the macroscale model predicts the same data as the microscale model. After fitting individual subject data, the parameters of the macroscale model changed. We call the changes from priors, *connection changes*. Besides MEG oscillations, the two datasets from [15], [16] we analyse here, also included structural data, in particular V1 size, measured with MR spectroscopy. After obtaining subject specific parameters (connection changes), we tested if the variability across subjects we observed in them is linked to V1 size and frequency and amplitude of MEG oscillations.

In Figure 4A we included the correlations between connection changes and V1 size for the data reported in [16]. We found four correlations with connection changes, three of them corresponding to changes in connections in *deep* cortical layers. The second pair of statistics in parentheses are partial correlations obtained when controlling for the rest of the connection changes. The parameters that showed significant correlations with V1 size were: changes in the drive to deep pyramidal neurons from superficial interneurons, a_{31} (Figure 4Aiii, *R*=-0.664*, p*=0.022; *R*=-0.858, *p*=0.142). Changes in the local drive to deep pyramidal neurons from other pyramidal neurons, a_{33} (Figure 4Ax, *R*=-0.594 *p*=0.046; *R*=-0.008, *p*=0.992). Finally, changes in the drive to deep inhibitory interneurons, a_{34} (Figure 4Av, *R*=0.685, *p*=0.017; *R*=0.683, *p*=0.317). The only correlation involving changes in *superficial* as opposed to deep connections was the drive to inhibitory interneurons in superficial cells from pyramidal neurons, a_{14} (Figure 4Aii, *R*=-0.685, *p*=0.017; *R*=-0.864, *p*=0.136). All above results did not remain significant when considering partial correlations or using Bonferroni correction. This was not surprising as our macroscale model predicts temporal differences in cortical dynamics between layers (vertically). Correlations with horizontal V1 surface are weak as the model does not describe interactions between neurons within the same layer of V1 (horizontally). This is because it does not describe the spatial deployment of cortical sources in detail (like all mean field models of the same sort, known as neural masses). Spatial deployment can instead be described by an extension of neural masses known as a neural field, see our earlier work [10].

---

[5] Fits are obtained after convergence of the EM algorithm [17]. They are tight: variance explained is above 95%.

In Figure 4B, we included the correlations between connection changes and gamma peak frequency. We found that, in this case, changes to the drive to superficial pyramidal neurons from inhibitory interneurons across individuals significantly correlated with peak frequency, a_{41} (Figure 4Bviii, $R=0.782$, $p=0.003$). Interestingly, this correlation remained significant under a Bonferroni correction for multiple tests. It was also positive. An increase in the inhibitory drive parameter in superficial layers led to an increase in the frequency peak. This positive correlation has also been found in other studies with similar models [6], [18]. This is the reverse of the effect on the inhibitory time constant, whose increase leads to a peak decrease. This is thought to capture changes in IPSPs where an increase in inhibition leads to longer IPSPs and reduces the peak frequency [19]. Similarly to earlier results, this correlation did not remain significant when controlling for the remaining connection weights ($R=-0.758$, $p=0.452$). This was not surprising as the same change (increase or decrease) in gamma peak frequency (and amplitude) can result from increasing or decreasing many of the parameters of Table 1[6].

We also considered the correlations of connection changes with gamma amplitude (Supplementary Figure S2B) and relative gamma amplitude change between conditions (3 stimulus sizes; Supplementary Figure S3). Gamma frequency, amplitude and relative amplitude change between conditions were the three data features that [16] used to quantify the gamma oscillation and response curve shape (while stimulus size increased). Thus, we considered the same features here. We found that changes in recurrent connections in *deep* inhibitory interneurons, a_{22} ,correlated with changes in gamma amplitude across subjects (Figure S3Biv, $R=0.608$, $p=0.04$). No other connection changes were correlated. Similarly to the result above, a_{22} did not remain significant under a Bonferroni correction and when controlling for the remaining connections ($R=0.904$, $p=0.096$).

We then fitted the macroscale model to data from [15]. This was similar to our earlier work [6] where we had used a different macroscale model (neural field). The data from [15] included visually induced oscillations from the visual cortex of 16 subjects recorded while subjects fixated on the centre of a screen and viewed a static, high-contrast, square-wave, vertical grating. Individual gamma peak frequency correlated with V1 size recorded with MR spectroscopy [15] (Supplementary Figure S1B).

Connection changes are shown, for each subject, in Figure 5 and Supplementary Figure S2A. Figure 5 which shows correlations of connection changes with V1 size. The horizontal axes show V1 size in mm². We found that changes in the connections between the *deep* pair of excitatory and inhibitory populations, a_{23} and a_{32} significantly correlated with different V1 sizes[7] (Figure 5vi and ix, $R=0.626$, $p=0.017$). This did not remain significant with a Bonferroni correction for multiple tests at the $p<0.05$ level and when controlling for the remaining connections ($R=0.251$, $p=0.684$). We also considered the correlations of connection changes with gamma peak frequency (Supplementary Figure S2A). No changes were significantly correlated.

---

[6] See [6] for model parameter effects on gamma frequency and amplitude.

[7] Note that only changes from priors between different subjects were the same for the two parameters. The parameters themselves were *not* the same as their priors were different.

In brief, in both datasets, we found six parameters that were correlated with V1 size, five of them relating to connections targeting *deep* cortical layers. Besides V1 size, we also found significant correlations with gamma peak frequency and amplitude in one dataset. Taken together, we found correlations between eight connection changes and V1 size or gamma oscillations. Six of them described differences in the level of *inhibition* across subjects. In the visually induced oscillations from 12 subjects reported in [16] that we analysed, we found changes in the excitability of inhibitory interneurons. In the oscillations from 16 subjects reported in [15] that we also analysed, we found differences in the drive and input to inhibitory interneurons. In both cases, the interneurons involved are those appearing in deep cortical layers of the microcircuit of Figure 1.

To sum up, the above results suggest that variability in MEG oscillations across subjects originated from differences in the level of inhibition in deep cortical layers and correlated with V1 size. However, correlations did not remain significant when corrected for multiple comparisons using a Bonferroni correction or accounting for other sources of variance using partial correlations. To test our hypothesis about individual variability in MEG oscillations originating from deep cortical layers, we turned to hypothesis testing using Bayesian methods. This has the advantage that it allows us to test *all* alternative hypotheses in terms of which model parameters drive individual variability. This also addresses the above problem of correlations not remaining significant by expressing significance in terms of a (Bayesian) odds ratio (known as Bayes Factor (*BF*)). Specifically, we used a recent approach known as parametric empirical Bayes (PEB)(see Methods and Supplementary Methods for a summary of the theory behind this [20], [21]). This provided a BF that quantifies the relative probability of a general linear model (GLM) that included the connection change as a predictor of a data feature (e.g. V1 size) *vs.* a GLM that did *not* include this parameter. If *BF>20:1* we can conclude that this connection change predicts the chosen data feature with probability *p>0.95* [22] . This is discussed next.

Individual differences in MEG oscillations reflect differences in V1 size and deep layer activity

We considered local intracortical connectivity between E-I neurons and asked whether connection changes predict different data features: gamma peak frequency and V1 size. This provided an answer to the question which connections and cortical layer might drive differences in these data features. We used the PEB approach. It allowed us to test all alternative possible combinations of connection changes that could predict data features. These correspond to combinations of the ten connections shown in Figure 1A (arrows). The connection changes were predictors that entered a GLM. The PEB approach can be thought of as a numerically efficient way to use Bayesian statistics to score alternative GLMs. Because it is Bayesian, it downweighs the contribution of outliers (quantified in terms of posterior variance). Here we considered alternative GLMs that did not include one or more of the connections between populations shown in Figure 1.

We computed the BF between a GLM model that included a particular connection change as predictor *vs.* a GLM without changes. This quantifies the probability of a particular connection predicting the data features. We found that variability in V1 size reported in [15] could be best predicted by the recurrent connectivity of deep inhibitory interneurons, a_{22} (Figure 6A; very strong evidence, *p>0.95*); while variability in V1 size reported in [16] could be best predicted by the inhibitory drive to deep pyramidal cells, a_{31} (Figure

6B; weak evidence, $p>0.5$). Gamma peak frequency was not predicted by connection changes. Thus, our result suggests that variability in the level of inhibition in deep cortical layers obtained by fitting MEG spectra, predicts V1 size. To sum up, differences in oscillations across people seem to reflect macroscopic differences in V1 size and microscopic differences in neuromodulation specific to deep layers 5/6.

Discussion

We found that individual differences in visually induced gamma oscillations recorded with MEG might reflect differences in the level of inhibition. Our results fit with studies that found GABAergic channels to play a prominent role in generating gamma oscillations [23] and that gamma oscillation variability reflects differences resting in GABA concentration measured with MR spectroscopy [24]. Also, they confirm results we obtained earlier using different models and statistical analyses. In [6] we found that individual differences in gamma oscillations reflected the level of inhibition in the cortical source. They were driven by variations in the excitatory drive to GABAergic interneurons. In [10], we also found that these differences were driven by variations in not only the drive but also interneuron output.

In the visual cortex interneurons spread across layers [25]. Thus, the earlier work described above did not answer the question if there are any differences in the level of inhibition between cortical layers. Here, we used a combination of macro- and microscale computational models to address this question. Both models predicted the same laminar data, that is, responses from different cortical depths. By fitting the macroscale model to MEG data, we were able to locate the cortical layer from which differences between subjects' responses might originate. We found that the variability in the level of inhibition across subjects was specific to *deep* cortical layers. Thus, our results suggest that differences in gamma oscillations between subjects originate from deep cortical layers.

Interactions between deep cortical inhibition and gamma oscillations of the sort we found here using non-invasive *human* brain data have also been observed using laminar electrodes in *animals*. A recent study [26] found that during memory delay, activity in deep cortical layers releases inhibition that targets superficial cortical layers. This interaction is expressed via alpha/beta oscillations that are known to inhibit distracting stimuli [27]. Neural activity in layers 5/6 might modulate activity in layers 2/3 and from there to layer 4 input [28]. This suggests a control of superficial gamma oscillations from deep alpha/beta inhibition. During delay, inhibition release allows recurrent excitation in superficial layers that allows memory ensembles to persist [29]. The balance between gamma and lower (alpha/beta) frequencies has also been shown to reflect volitional control of working memory originating from frontal areas [30].

To sum up, animal studies found that deep layer activity in sensory areas modulates superficial gamma oscillations and is the result of feedback from higher areas [26], [30]. Here, we found that deep cortical inhibition is also related to gamma oscillation variability observed in human subjects. Thus, our results suggest that modulation of gamma oscillations by deep layer activity is different between different people. Predictive Coding (PC) offers a plausible explanation of this result. It suggests that deep layer activity corresponds to different hypotheses about sensory input that different people have [11], [31]. This in turn, might be due to their different prior experiences.

To sum up, PC offers a plausible explanation of deep layer activity underlying differences in gamma oscillations found in human MEG data here. PC suggests that deep cortical layer activity represents predictions about incoming sensory stimuli that interact with sensory driven superficial layer activity. A recent study using invasive, laminar electrodes in monkeys supports this hypothesis [32]. These authors found that changes in deep cortical inhibition correlated with stimulus predictability, or how confident the monkey was about the upcoming stimulus. This result can be reconciled with our current result in the context of the theory of PC: changes in deep inhibition can reflect both 1) different processing of the same sensory input by different people that we found here and 2) different sensory inputs (with different predictability) the same subject might have found in [32].

Our approach focuses on explaining differences in neural dynamics between different cortical layers. Because of constraints imposed upon the parameters of the microscale (compartmental) model, spatial effects within the same layer were neglected (e.g. interactions between cortical columns and the dependence of gamma peak frequency on the horizontal spread of the underlying cortical source). In future work, we will study the relationship between population activity predicted by the mean field (macroscale) model and the symmetric compartmental (microscale) model we used here. Spatial effects could be addressed by extending our approach to a different class of means field models besides the neural masses considered here, called neural fields [10]. In earlier work, we used neural fields to explain the relation between spatial and biophysical properties of cortical sources and observed brain dynamics [6]. We also revealed differences in brain structure and function between different people [10].

To sum up, we here used combinations of macroscale and microscale models, not isolated models. We estimated neural activity in different cortical depths of the visual cortex using non-invasive, human MEG data. We suggested that isolated models spanning a single spatial scale are not sufficient to fully understand such data and instead model combinations should be used. This will allow us to obtain information about human cortical layer activity and anatomy and exploit information at different spatial scales simultaneously.

More generally, our approach can help address the heterogeneity observed in neurological diseases and disorders. We claim that using isolated models is not enough to address this heterogeneity. Instead combinations of models spanning the macro- and micro- scales like the one introduced here should be considered. Spatial information at the macroscale is accessible via non-invasive human brain imaging. Spatial information at the microscale and the structure and function of local cortical circuits is accessible in vitro or through invasive recordings in animals. Developing hybrid models that combine both scales like we did here, exploits multiscale spatial information.

Our approach can help understand pathologies underlying neurological diseases and disorders. The link between deep layer activity, cortical inhibition and gamma oscillations found here and in [26], [32] is relevant to such disorders, where the excitation to inhibition (E-I) balance is disrupted [33]. Individual differences in E-I balance are one prominent feature of heterogeneity observed in neurological disorders [34]. Based on our results, we predict that E-I balance disruption will be more prominent in deep cortical layers. In future work, we will use the approach presented here to test this hypothesis. E-I differences are expressed at the local cortical circuit level. At the same time, they are reflected in

individual differences in human M/EEG data. Thus, our multiscale approach enables the study of E-I balance differences at the cortical circuit level using M/EEG data. It opens up the way to testing hypotheses about cortical circuit structure and function in human patient populations.

Materials and Methods

Data. We used two datasets to ask from similar but independent experiments to test whether our model will yield similar results. First, we used MEG beamformed data from [15]. These included visually induced oscillations from the visual cortex of 16 subjects. In that task, subjects fixated on the centre of a screen and viewed a static, high-contrast, square-wave, vertical grating. MEG data were recorded using a whole-head CTF axial gradiometer system with 275 channels, sampled at 600 Hz. Three electrical coils were placed at fiducial locations and used to monitor subject head movement. Data were analysed using SPM8 (http://www.fil.ion.ucl.ac.uk/spm). We used an LCMV beamformer algorithm implemented in SPM8 to quantify source power in the time window between 0.5 and 1.5 s after stimulus onset relative to baseline power over one second preceding stimulus onset. We located peak gamma activity in the medial occipital cortex, and at this peak location we used the beamformer weights to extract the time series of the virtual electrode. In the second dataset, also including visually induced oscillations taken from [16], whole-head MEG recordings were acquired using a similar CTF radial gradiometer system as above sampled at 1,200 Hz. In that task, a time series of the response at the location of the peak in each trial was generated by spatially filtering the sensor-level data through the corresponding beamformer weights at that location. These "virtual sensor" time series were then used to obtain power spectra from 12 healthy subjects while they viewed grating stimuli of three different sizes— 2^0, 4^0 & 8^0 . For more details on the above two datasets, see [15], [16].

Biophysical models. To construct the microscopic model (that we called symmetric compartmental model), we adapted NEURON code from [5]. This model can be found in https://github.com/pinotsislab/MicroMacro/ . The original model of [5] included a compartmental model of the local microcircuit shown in Figure 1. This comprised pyramidal neurons and interneurons [35]. This model was later extended to a network model of a cortical column in a key paper by [5]. The resulting network model provides detailed descriptions of intracellular (longitudinal) currents within the long apical dendrites of synchronized cortical pyramidal cells. It describes neuronal morphology and laminar structure of a cortical column and characterizes cellular and circuit level processes measured with multielectrode arrays or MEG. Driven with Gaussian input, the model accurately reproduced the S1 evoked response to a tap on the hand and described the intracellular currents that give rise to signal polarity.We here used a variant of this model where we increased the number of inhibitory units from three to 10 per layer, so that their number was equal to the number of the principal cells within each mini-column. To ensure that relative differences in interneuron densities were accommodated, we multiplied the maximum conductance values of the corresponding connections by a factor of 0.3. To reduce the number of connection parameters (synaptic strengths) to be equal to those of the macroscopic model, we assumed that the connection weights between different mini-columns comprising the microcircuit shown in Figure 1A were the same. We then simulated data from the symmetric compartmental model and made them amenable to a further DCM analysis, see *simData.mat* in the above Github repository.

We also implemented a macroscale (neural mass) model that describes the same local cortical circuit as the microscale model above. This assumes symmetry constraints on horizontal connectivity (within each cortical layer) of the sort assumed in mean field models that describe aggregate activity over hundreds of neurons. It is implemented as part of the DCM toolbox of SPM12, https://www.fil.ion.ucl.ac.uk/spm/software/spm12/, see *spm_fx_cmc_BS.m* in the above Github repository. Then we fitted this model to the simulated data above using DCM and estimated its parameters as described in the next section.

Parameter estimation. To fit the biophysical mean field model to simulated and MEG data we used DCM for steady state responses [12], that is implemented in the function *spm_dcm_csd.m* of the DCM toolbox. The inversion of mean field models above uses the standard DCM approach that we summarize below. First, it assumes that the mean field model is driven by endogenous neuronal fluctuations that produce observed power spectra according to the following likelihood model, see also [10]:

$$g(\omega) = g_Y(\omega, \theta^{(1)}) + g_N(\omega) + \varepsilon^{(1)}$$

$$g_Y(\omega, \theta^{(1)}) = \sum_k \left| Q L_r(k, \varphi) T(k, \omega, \theta^{(1)}) g_u(k, \omega) \right|^2$$

$$g_n(\omega) = \alpha_n + \beta_n / \omega \tag{1}$$

$$g_u(\omega) = \alpha_u + \beta_u / \omega$$

$$\mathrm{Re}(\varepsilon^{(1)}) \sim \mathcal{N}(0, \Sigma(\omega, \lambda)) \quad \mathrm{Im}(\varepsilon^{(1)}) \sim \mathcal{N}(0, \Sigma(\omega, \lambda))$$

Here, $g_u(\omega)$ is a spatiotemporal representation of fluctuations or inputs driving induced responses, which we assume to be spatially white and a mixture of white and pink temporal components, $L_r(k, \varphi)$ is the Fourier transform of the lead field of the MEG virtual electrode and $Q = [q_1, q_2, q_3, q_4]$ is a vector of coefficients that weights the contributions of each neuronal population (among the 4 shown in Figure 1A) to the observed MEG signal. These contributions are based on differences in anatomical properties and the lead field configuration of each population (e.g. inhibitory neurons do not generate a large dipole), where each electrode or sensor has its own sensitivity profile. The transfer function $T(k, \omega, \theta^{(1)})$ are the Fourier transform of the impulse response or first-order Volterra kernel associated with the ordinary differential equations shown in the boxes of Figure 1A. This transfer function describes how each of the four populations response to neuronal fluctuations, where the model parameters describe the connectivity architecture mediating responses, the observation function $\varphi \subset \theta^{(1)}$ and the spectra of the inputs and channel noise, $\{\alpha_n, \alpha_u, \beta_n, \beta_u\} \subset \theta^{(1)}$.

In summary, Equation (1) expresses the data features $g(\omega)$ as a mixture of predictions and sampling errors $\varepsilon^{(1)}$ with covariance $\Sigma(\omega, \lambda)$. Gaussian assumptions about these sampling errors provide the likelihood model at the first (within-subject) level: $p(g(\omega) | \theta^{(1)})$. The

predictions are themselves a mixture of predicted cross spectra and channel noise $g_N(\omega)$. This concludes the description of the likelihood model.

The optimization of this likelihood model is based on optimizing a free energy bound on the model log-evidence. The free energy bound is optimized with respect to a variational density $q(\theta) \sim \mathcal{N}(\mu, C)$ on the unknown model parameters. By construction, the free energy bound ensures that when the variational density maximizes free energy, it approximates the true posterior density over parameters, $q(\theta) \approx p(\theta \mid g_Y(\omega)_i, m)$. This bound is given by [6]:

$$\mathcal{F} = \mathcal{G}(\mu) + \tfrac{1}{2} \ln |\partial_{\mu\mu} \mathcal{G}|$$

$$\mathcal{G} = -\tfrac{1}{2} \mathrm{Re}(\varepsilon)^T \Sigma^{-1} \mathrm{Re}(\varepsilon) - \tfrac{1}{2} \mathrm{Im}(\varepsilon)^T \Pi \, \mathrm{Im}(\varepsilon) - \tfrac{1}{2} \rho^T \Omega^{-1} \rho - \tfrac{1}{2} \ln |\Sigma| - \tfrac{1}{2} \ln |\Omega|$$

$$\varepsilon = g_Y(\omega, \mu) + g_N(\omega, \mu) - \mathbf{g}_Y(\omega)$$

$$\rho = \mu - \phi \tag{2}$$

where $\varepsilon(\mu) \sim \mathcal{N}(I, \Sigma(\omega, \lambda))$ is the Gaussian observation noise and $p(\theta \mid m) = \mathcal{N}(\phi, \Omega)$ are the variational and prior densities. $g_Y(\omega, \mu) + g_N(\omega, \mu)$ are the predictions of the data features $\mathbf{g}_Y(\omega)$ and $\rho(\mu) \in \mathbb{R}$ are prediction errors on the parameters, in relation to their prior density and $\mathcal{G}$ is the Gibb's energy of the system.

Using DCM, we fitted the mean field model to power spectra between 30-80Hz from different subjects reported in [15], [16]. This frequency range contains gamma frequencies captured with MEG[15], [16]. This analysis was similar to our earlier work described in [6], [10]. To fit data from [16], we fitted all three stimuli conditions simultaneously by modeling different sizes as condition-specific effects (the B matrix that is used in DCM), see [10]. We next consider how this likelihood model is placed within a hierarchical model of responses from multiple subjects.

Between subject differences. To model between-subject effects we used a hierarchical Bayesian inference approach known as Parametric Empirical Bayes, see [10], [20], [21] and the function *spm_dcm_peb_bmc.m* in the DCM toolbox. At the first level, we used the mean field model described above and in Results. At the second level, we modelled individual differences in gamma frequency and amplitude. These were used as regressors in the design matrix entering the GLM at the second level of a hierarchical Bayesian model.

Below we summarize the PEB approach. For a detailed analysis, see [9], [10] and [16]. This approach is based on hierarchical model that accommodates both within and between subject effects given by the following Equations.

$$y_i = \Gamma_i(\theta^{(1)}) + \varepsilon_i^{(1)}$$
$$\theta^{(1)} = \Gamma(\theta^{(2)}) + \varepsilon^{(2)}$$
$$\theta^{(2)} = \eta + \varepsilon^{(3)} \tag{3}$$

where y_i is a matrix of i-th subject responses, $\Gamma_i(\theta^{(1)}) = g_Y(\omega, \mu, \theta^{(1)})$ represents the (differential equation or dynamic causal) model that generates these responses with parameters $\theta^{(1)}$, $\Gamma(\theta^{(2)})$ is the between subject (second level) model that describes intersubject variability in the parameters of the first level model. The second level maps second to first level parameters (e.g., group means to subject-specific parameters), where $\varepsilon^{(i)}$ represent random effects at each level (e.g., intersubject variability and observation noise).

In the context of non-invasive electrophysiology, the hierarchical model (3) poses the difficult inversion problem of finding neural source estimates in the context of intersubject variability. This involves (i) partitioning the covariance of observed data into observation error and components that can be explained in terms of neuronal responses, that themselves entail components due to second level (between-subject) level variability; (ii) exploiting differential equation models to provide anatomical and physiological constraints on the explanation for first level (within subject) responses – usually in terms of (synaptic) connectivity estimates. The second level covariance components specify whether the parameters of the dynamical model at the first level are random or fixed effects, while dynamical models provide predictions of the dynamics at source and sensor space, which depend upon cortical anatomy and physiology.

In summary, hierarchical or empirical Bayesian modelling of the sort implied by Equation (3) allows us to perform efficient source reconstruction and obtain connectivity estimates by replacing phenomenological constraints (e.g., based on autoregressive modelling and temporal smoothness considerations) by spatiotemporal constraints based on models of neuronal activity. This can be thought as an alternative to autoregressive models, which model statistical dependencies among measured signals – as opposed to the neuronal processes generating measurements. In DCM, one uses a forward or generative model of distributed processing to estimate the (coupling) parameters of that model. Inference then proceeds assuming nonlinear within-subject effects and linear between subject effects. This allows one to distinguish among competing hypotheses about the mechanisms and architectures generating the data and the nature of group effects in multiple subject studies

In this hierarchical model, constraints on the posterior density over model parameters are provided by the level above. In variational Bayesian inference, the approximate posterior over the second level parameters is obtained by optimising its sufficient statistics (i.e., mean and covariance) with respect to a (second level) free energy:

$$\hat{q}^{(1)*} = \arg\max_{\hat{q}^{(1)}} F^{(1)}(\hat{p}_F, \hat{q}^{(1)})$$

$$\hat{q}^{(2)*} = \arg\max_{\hat{q}^{(2)}} F^{(2)}(\hat{p}^{(2)}, \hat{q}^{(2)}, \hat{q}^{(1)*})$$

$$F^{(1)}(\hat{p}_R, \hat{q}^{(1)}) = E_{q^{(1)}}[\ln p(y \mid \theta^{(1)}, m)] - D_{KL}[q(\theta^{(1)} \mid \hat{q}^{(1)}) \| p(\theta^{(1)} \mid \hat{p}_R)]$$

$$F^{(2)}(\hat{p}^{(2)}, \hat{q}^{(2)}, \hat{q}^{(1)*}) = E_{q^{(2)}}[F^{(1)}(\hat{p}_R, \hat{q}^{(1)})] - D_{KL}[q(\theta^{(2)} \mid \hat{q}^{(2)}) \| p(\theta^{(2)} \mid \hat{p}^{(2)})]$$

$$\hat{p}_R = (\Gamma(\theta^{(2)}), \Sigma(\theta^{(2)})) \tag{4}$$

The key thing about this free energy is that it can be evaluated (using Bayesian Model Reduction;BMR [21]) without optimising the first level posterior. This means the second level parameters (e.g., group means) can be optimised or estimated, for any given model of priors, without reinventing the model at the first level. Technically, the inversion of the hierarchical or empirical Bayesian model only requires the posterior density from the inversion of each subject's DCM. In short, the use of BMR allows one to make inferences at the group level without having to re-estimate subject-specific parameters; see [16] for details and a study of robustness of this scheme.

References
There is only one reference list for all sources cited in the main text, figure and table legends, and Supplementary Materials, and this main list **should not exceed 40 citations**.

[1]   G. D. Pearlson, B. A. Clementz, J. A. Sweeney, M. S. Keshavan, and C. A. Tamminga, "Does biology transcend the symptom-based boundaries of psychosis?," *Psychiatric Clinics*, vol. 39, no. 2, pp. 165–174, 2016.
[2]   J.-M. Fellous, G. Sapiro, A. Rossi, H. S. Mayberg, and M. Ferrante, "Explainable Artificial Intelligence for Neuroscience: Behavioral Neurostimulation," *Frontiers in Neuroscience*, vol. 13, p. 1346, 2019.
[3]   P. R. Montague, R. J. Dolan, K. J. Friston, and P. Dayan, "Computational psychiatry," *Trends in cognitive sciences*, vol. 16, no. 1, pp. 72–80, 2012.
[4]   G. Lovas, N. Szilágyi, K. Majtényi, M. Palkovits, and S. Komoly, "Axonal changes in chronic demyelinated cervical spinal cord plaques," *Brain*, vol. 123, no. 2, pp. 308–317, 2000.
[5]   S. R. Jones, D. L. Pritchett, S. M. Stufflebeam, M. Hämäläinen, and C. I. Moore, "Neural correlates of tactile detection: a combined magnetoencephalography and biophysically based computational modeling study," *The Journal of Neuroscience*, vol. 27, no. 40, pp. 10751–10764, 2007.
[6]   Pinotsis, D. S. Schwarzkopf, V. Litvak, G. Rees, G. Barnes, and K. J. Friston, "Dynamic causal modelling of lateral interactions in the visual cortex," *NeuroImage*, vol. 66, pp. 563–576, 2013.
[7]   J. Berger, *Statistical decision theory: foundations, concepts, and methods*. Springer Science & Business Media, 2013.
[8]   D. A. Pinotsis *et al.*, "Linking canonical microcircuits and neuronal activity: Dynamic causal modelling of laminar recordings," *Neuroimage*, vol. 146, pp. 355–366, 2017.
[9]   D. A. Pinotsis, R. Loonis, A. M. Bastos, E. K. Miller, and K. J. Friston, "Bayesian modelling of induced responses and neuronal rhythms," *Brain topography*, pp. 1–14, 2016.
[10]   D. A. Pinotsis, G. Perry, V. Litvak, K. D. Singh, and K. J. Friston, "Intersubject variability and induced gamma in the visual cortex: DCM with empirical B ayes and neural fields," *Human brain mapping*, vol. 37, no. 12, pp. 4597–4614, 2016.
[11]   D. A. Pinotsis, T. J. Buschman, and E. K. Miller, "Working Memory Load Modulates Neuronal Coupling," *Cerebral Cortex*, 2018.

[12] R. J. Moran, K. E. Stephan, T. Seidenbecher, H. C. Pape, R. J. Dolan, and K. J. Friston, "Dynamic causal models of steady-state responses," *Neuroimage*, vol. 44, no. 3, pp. 796–811, 2009.

[13] R. E. Rosch *et al.*, "NMDA-receptor antibodies alter cortical microcircuit dynamics," *Proceedings of the National Academy of Sciences*, vol. 115, no. 42, pp. E9916–E9925, 2018.

[14] S. Lee and S. R. Jones, "Distinguishing mechanisms of gamma frequency oscillations in human current source signals using a computational model of a laminar neocortical network," *Frontiers in human neuroscience*, vol. 7, p. 869, 2013.

[15] D. S. Schwarzkopf, D. J. Robertson, C. Song, G. R. Barnes, and G. Rees, "The frequency of visually induced gamma-band oscillations depends on the size of early human visual cortex," *J Neurosci*, vol. 32, pp. 1507–1512, 2012.

[16] G. Perry, K. Hamandi, L. M. Brindley, S. D. Muthukumaraswamy, and K. D. Singh, "The properties of induced gamma oscillations in human visual cortex show individual variability in their dependence on stimulus size," *Neuroimage*, vol. 68, pp. 83–92, 2013.

[17] A. P. Dempster, N. M. Laird, and D. B. Rubin, "Maximum likelihood from incomplete data via the EM algorithm," *Journal of the Royal Statistical Society: Series B (Methodological)*, vol. 39, no. 1, pp. 1–22, 1977.

[18] A. D. Shaw *et al.*, "Neurophysiologically-informed markers of individual variability and pharmacological manipulation of human cortical gamma," *NeuroImage*, vol. 161, pp. 19–31, 2017.

[19] G. Buzsáki and X.-J. Wang, "Mechanisms of gamma oscillations," *Annual review of neuroscience*, vol. 35, p. 203, 2012.

[20] K. H. Preller, A. Razi, P. Zeidman, P. Stämpfli, K. J. Friston, and F. X. Vollenweider, "Effective connectivity changes in LSD-induced altered states of consciousness in humans," *Proceedings of the National Academy of Sciences*, vol. 116, no. 7, pp. 2743–2748, 2019.

[21] K. J. Friston *et al.*, "Bayesian model reduction and empirical Bayes for group (DCM) studies," *NeuroImage*, 2015.

[22] R. E. Kass and A. E. Raftery, "Bayes factors," *Journal of the American Statistical Association*, vol. 90, no. 430, pp. 773–795, 1995.

[23] N. Brunel and X.-J. Wang, "What determines the frequency of fast network oscillations with irregular neural discharges? I. Synaptic dynamics and excitation-inhibition balance," *Journal of Neurophysiology*, vol. 90, no. 1, pp. 415–430, 2003.

[24] S. D. Muthukumaraswamy, R. A. E. Edden, D. K. Jones, J. B. Swettenham, and K. D. Singh, "Resting GABA concentration predicts peak gamma frequency and fMRI amplitude in response to visual stimulation in humans," *Proceedings of the National Academy of Sciences*, vol. 106, no. 20, p. 8356, 2009.

[25] G. Ciceri, N. Dehorter, I. Sols, Z. J. Huang, M. Maravall, and O. Marín, "Lineage-specific laminar organization of cortical GABAergic interneurons," *Nature neuroscience*, vol. 16, no. 9, p. 1199, 2013.

[26] A. M. Bastos, R. Loonis, S. Kornblith, M. Lundqvist, and E. K. Miller, "Laminar recordings in frontal cortex suggest distinct layers for maintenance and control of working memory," *Proceedings of the National Academy of Sciences*, vol. 115, no. 5, pp. 1117–1122, 2018.

[27] T. J. Buschman and E. K. Miller, "Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices," *science*, vol. 315, no. 5820, pp. 1860–1862, 2007.

[28] K. D. Harris and T. D. Mrsic-Flogel, "Cortical connectivity and sensory coding," *Nature*, vol. 503, no. 7474, pp. 51–58, 2013.

[29] D. A. Pinotsis, S. L. Brincat, and E. K. Miller, "On memories, neural ensembles and mental flexibility," *NeuroImage*, vol. 157, pp. 297–313, 2017.

[30] M. Lundqvist, P. Herman, M. R. Warden, S. L. Brincat, and E. K. Miller, "Gamma and beta bursts during working memory readout suggest roles in its volitional control," *Nature communications*, vol. 9, no. 1, p. 394, 2018.

[31] A. M. Bastos *et al.*, "Visual areas exert feedforward and feedback influences through distinct frequency channels," *Neuron*, vol. 85, no. 2, pp. 390–401, 2015.

[32] A. M. Bastos, M. Lundqvist, A. S. Waite, N. Kopell, and E. K. Miller, "Layer and rhythm specificity for predictive routing," *NEURON-D-20-00146*, 2020.

[33] Y. Chen, K. Nakayama, D. Levy, S. Matthysse, and P. Holzman, "Processing of global, but not local, motion direction is deficient in schizophrenia," *Schizophrenia research*, vol. 61, no. 2–3, pp. 215–227, 2003.

[34] E. Milne, J. Swettenham, P. Hansen, R. Campbell, H. Jeffries, and K. Plaisted, "High motion coherence thresholds in children with autism," *Journal of Child Psychology and Psychiatry*, vol. 43, no. 2, pp. 255–263, 2002.

[35] P. C. Bush and T. J. Sejnowski, "Reduced compartmental models of neocortical pyramidal cells," *Journal of neuroscience methods*, vol. 46, no. 2, pp. 159–166, 1993.

[36] P. G. Bissiri, C. C. Holmes, and S. G. Walker, "A general framework for updating belief distributions," *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, vol. 78, no. 5, pp. 1103–1130, 2016.

[37] T. Zhang, "Information-theoretic upper and lower bounds for statistical estimation," *IEEE Transactions on Information Theory*, vol. 52, no. 4, pp. 1307–1321, 2006.

[38] V. Afraimovich, J. R. Chazottes, and A. Cordonet, "Synchronization in directionally coupled systems: Some rigorous results," *DISCRETE AND CONTINUOUS DYNAMICAL SYSTEMS SERIES B*, vol. 1, no. 4, pp. 421–442, 2001.

[39] M. Breakspear *et al.*, "A disturbance of nonlinear interdependence in scalp EEG of subjects with first episode schizophrenia," *Neuroimage*, vol. 20, no. 1, pp. 466–478, 2003.

[40] D. A. Pinotsis, R. J. Moran, and K. J. Friston, "Dynamic causal modeling with neural fields," *Neuroimage*, vol. 59, no. 2, pp. 1261–1274, 2012.

Acknowledgments

This work was supported by UKRI ES/T01279X/1, NIMH R37MH087027, ONR MURI N00014-16-1-2832, and The MIT Picower Institute Innovation Fund. We thank Dr Daniel Gibson for a careful reading of the manuscript and important suggestions, Dr Sam Schwarzkopf for illuminating discussions and providing data and Professor Krish Singh and Dr Gavin Perry for providing data.

Author contributions: DAP conceived the study and carried out the data analysis. EKM provided critical feedback and helped shape the research, analysis and manuscript.

Figures and Tables

Fig. 1. The Bush and Sejnowski microcircuit. (A) Two pairs of excitatory (black) and inhibitory (red) populations occupy superficial and deep cortical layers. Firing rates within each population provide inputs to other populations and convolution of presynaptic activity produces postsynaptic depolarization. Arrows denote

excitatory and inhibitory connections. All recurrent connections are inhibitory. The same microcircuit was implemented both as a mean field [8] and a compartmental model [5]. The equations describe the evolution of hidden states corresponding to activity in each of the four populations in the mean field model. **(B)** Both models predict LFPs and power spectra. The top plot shows predicted power spectra from both models generated by superficial pyramidal neurons. Predictions from the mean field model are shown in magenta, while predictions from the compartmental model are shown in green together with 95% confidence intervals. The bottom plot shows the same results for predicted spectra from deep pyramidal neurons. Insets in the top right corner of both plots show predicted LFPs. Note the peaked responses at 10Hz that are reminiscent of spiking burst input that are also captured by the mean field model responses.

Fig. 2. (A) Schematic of our analysis pipeline. A. We first establish a similarity

between the model of [5] and its symmetric variant. Here horizontal arrows of different widths in the left panel denote asymmetric connectivities and delays between mini-columns depicted as rectangles containing Superficial and Deep Pyramidal cells (SP and DP) and Inhibitory Interneurons (II). In the right panel, a symmetrisation of the model reduces the number of connectivity parameters to be the same as those in a homologous mean field model. **(B)** We then demonstrate the construct validity of the corresponding mean field model in relation to symmetric compartmental model above. This is achieved by fitting the model to synthetic (laminar) data obtained from its compartmental homologue. Red and green lines in the middle panel correspond to real data and model predictions. Solid and dashed lines to real and imaginary parts of cross spectra between deep and superficial pyramidal neurons.

Fig. 3. Model fits to data. (A) Empirical responses (power spectra) and model fits are shown in dashed and solid lines along with 95% confidence intervals across subjects for the data from [16] (smallest stimulus size; grand average across all subjects). These quantify variability across subjects and overlap for model predictions and data. **(B)** Example model fits to individual subjects in data from [16]. Fits are obtained for all three stimulus sizes simultaneously by modeling different sizes as condition-specific effects. The three stimulus sizes correspond to red, blue and green lines. Dashed and solid lines correspond to data and model predictions. Differences in power spectra between conditions for the same subject are small and spectra corresponding to different stimulus sizes overlap.

Fig. 4. (A) Correlations between connectivity parameters and V1 size. Parameters were obtained by fitting data from [16]. Changes in the drive to deep pyramidal neurons from both superficial interneurons, a_{31} (iii) and deep pyramidals, a_{33} (x) correlated with V1 size. Also changes in the drive to deep interneurons, a_{34} (v) and drive to inhibitory interneurons in superficial cells, a_{14} (ii).

Fig. 4. (B) Correlations between connectivity parameters and gamma peaks. Parameters were obtained by fitting data from [16]. Changes to the drive to superficial pyramidal neurons correlated with peak frequency, a_{41} (viii). Least squares fitted line shown in red.

Fig. 5. Connection changes in data from [15]. Correlations between connectivity parameters and V1 size using data from [15]. Changes in recurrent connections in deep inhibitory interneurons, a_{22} ,correlated with changes in gamma amplitude. Changes in the connections between the *deep* pair of excitatory and inhibitory populations, a_{23} and a_{32} significantly correlated with different V1 sizes (iv,vi and ix). Least squares fitted line shown in red.

Fig. 6. V1 size predictors. (A) We scored alternative GLMs where predictors of variability in V1 included any combination of the connections (arrows) in Fig.1A. We found that for the data from [15] V1 size could be best predicted by the recurrent connectivity of deep inhibitory interneurons, a_{22} (orange arrow). Evidence was very strong $p>0.95$. **(B)** Same as in (A) for data from [16] . V1 size variability reported in [16] could be best predicted by the inhibitory drive to deep pyramidal cells, a_{31} (orange arrow). Evidence was weak $p>0.5$.

Table 1 *Synaptic connectivity parameters for the microscale and macroscale models.*

	Max conductance (µS) Microscale model	Intrinsic connectivity Macroscale model (a.u)
a_{44}	0.001/0.0005	4.4
a_{14}	0.003	4.8
a_{34}	0.00025	23.3
a_{41}	0.015/0.015	3.8
a_{31}	0.0003	5.9
a_{11}	0.0006	4.2
a_{33}	0.005/0.0005	2.2
a_{23}	0.0003	4.6
a_{32}	0.0075/0.0075	6.9
a_{22}	0.0006	4.16