Thiocillin and Micrococcin Exploit the Ferrioxamine Receptor of *Pseudomonas aeruginosa* for Uptake

Derek C. K. Chan and Lori L. Burrows*

Michael G. DeGroote Institute for Infectious Disease Research, Department of Biochemistry and
Biomedical Sciences, McMaster University, 1200 Main Street West, Hamilton, Ontario L8N 3Z5, Canada

KEYWORDS: drug discovery, mechanism of uptake, thiopeptide, antimicrobial activity, siderophore, trojan
horse,

ABSTRACT

Thiopeptides are a class of Gram-positive antibiotics that inhibit protein synthesis. They have been
underutilized as therapeutics due to solubility issues, poor bioavailability, and lack of activity against
Gram-negative pathogens. We discovered recently that a member of this family, thiostrepton, has activity
against *Pseudomonas aeruginosa and Acinetobacter baumannii* under iron-limiting conditions.
Thiostrepton uses pyoverdine siderophore receptors to cross the outer membrane, and combining
thiostrepton with an iron chelator yielded remarkable synergy, significantly reducing the minimal
inhibitory concentration. These results led to the hypothesis that other thiopeptides could also inhibit
growth by using siderophore receptors to gain access to the cell. Here, we screened six thiopeptides for
synergy with the iron chelator deferasirox against *P. aeruginosa* and a mutant lacking the pyoverdine
receptors FpvA and FpvB. Our findings suggest that thiopeptides such as thiocillin cross the outer
membrane using FoxA, the ferrioxamine siderophore receptor. Other structurally related thiopeptides did
not inhibit growth of *P. aeruginosa*, but had greater potency against methicillin-resistant *Staphylococcus*
aureus than thiostrepton and related thiopeptides*.* These results suggest that thiopeptide structures have
evolved with considerations for target affinity and entry into cells.

INTRODUCTION

Thiopeptides are a class of natural products with potent antibacterial activity against Gram-
positive bacteria, including methicillin-resistant *Staphylococcus aureus* (MRSA)[1]. They are characterized
chemically by the oxidation state of a central pyridine ring and by the size of a core macrocyclic ring
comprised of thiazole moieties[2]. Thiopeptides act on the ribosome to inhibit protein translation but the
exact mechanism depends on the number of members in the core macrocyclic ring: 26-membered
macrocycles inhibit elongation factor G whereas 29-membered macrocycles block elongation factor Tu
binding[3–5]. The exact mechanism of action for 35-membered macrocycles remains unknown, although
they are reported to target the 50S ribosome, similar to thiostrepton (TS)[6]. TS is among the best-studied

26-membered thiopeptides and has multiple biological properties, including anticancer and antimalarial

activities[7,8].

In 1948, Su isolated a strain of *Micrococcus* from sewage samples that produced a compound with

activity against Gram positive but not Gram negative bacteria[9]. The compound, micrococcin, was the first

thiopeptide to be discovered. Since then, the thiopeptide family has expanded to over 100 members of

diverse structure and origin[10]. Thiopeptides belong to a larger class of natural products called ribosomally

synthesized and post-translationally modified peptides (RiPPs)[11]. These natural products are produced

from a peptide precursor whose sequence varies depending on the thiopeptide, and undergo many post-

translational modifications to reach their final structure[12]. Most thiopeptides are produced by Gram-

positive soil and marine organisms including *Bacillus cereus, Streptomyces* spp*.*, and *Nocardopsis* spp*.,*

although some Gram negatives such as *Serratia marcesens* have biosynthetic gene clusters encoding

thiopeptide synthesis[13–17].

Due to their poor solubility, challenging synthesis, and limited potency against Gram-negative

bacteria, pharmaceutical companies ruled out thiopeptides for development as clinically useful

antibiotics. Instead, researchers focused on addressing these issues through total synthesis, discovery of

new thiopeptides and structure elucidation, identification of thiopeptide biosynthetic gene clusters, and

modification of existing thiopeptides to improve solubility and activity [12,17–25]. However, there is a gap in

knowledge regarding their utility against Gram-negative bacteria which likely stems from the assumption

that thiopeptides cannot cross the outer membrane.

We showed recently that TS hijacks pyoverdine receptors to cross the outer membranes of

Pseudomonas aeruginosa and *Acinetobacter baumannii,* among the World Health Organization's priority

pathogens for new antibiotic development[26] [27] *P. aeruginosa* and *A. baumannii* are opportunistic

pathogens with intrinsic resistance to many antibiotics and are causative agents of healthcare-acquired

infections[28]. Both express receptors for pyoverdine, a siderophore with high affinity for iron. TS uptake

through pyoverdine receptors occurred under iron-limited conditions, and deferasirox (DSX), an FDA-

approved oral iron chelator, strongly synergized with TS to inhibit the growth of clinical isolates of both

species*.* Combinations of TS with other iron chelators were also effective[29]. These findings showed that

thiopeptides can cross the outer membranes of specific Gram-negative bacteria. TS lacks activity against

Escherichia coli because it does not express pyoverdine receptors. The thiopeptide family has over 100

members and recent bioinformatic analyses suggest that there could be hundreds more thiopeptides[30,31].

We hypothesized that other thiopeptides may also have activity against Gram-negative pathogens.

Here, we screened six thiopeptides (Figure 1, Supplementary Figure 1) for activity against *P.*

aeruginosa and potential synergy with DSX. Combinatorial assays showed that two of these used the

ferrioxamine siderophore receptor, FoxA, to enter the cell, confirming our hypothesis that other

thiopeptides could cross the outer membrane. The 26-membered macrocycles are of particular interest

Figure 1. Structures of Thiopeptides Tested for Synergy with DSX. Differences between structural analogues are highlighted in yellow. The following are structural analogues: TS and SM, TC and MC, (26-membered macrocycles) and BER & GEN (35-membered macrocycles). NH is structurally similar to TC and MC, but has a unique indole moiety (blue). SM and TS have a unique quinaldic acid moiety (orange), but SM differs by two amino acids.

because modifications to the core macrocyclic ring altered siderophore specificity. Other modifications

abolished Gram-negative activity but improved Gram-positive activity. Thiopeptides are an underutilitzed

class of antibiotics with potential for development as clinically useful drugs, especially as iron limitation

at sites of infection could promote thiopeptide uptake.

RESULTS

Siomycin A Uses the Pyoverdine Receptors for Uptake

We screened the activity of six thiopeptides in combination with DSX against *P. aeruginosa* PA14 using

checkerboard assays. We looked for individual thiopeptide activity, synergy, and activity against a mutant

lacking pyoverdine receptors FpvA and FpvB, which results in TS resistance. If the combination was active

against the wild type but not the *fpvAB* mutant, the thiopeptide likely uses those receptors to cross the

membrane. The six thiopeptides in the screen included siomycin A (SM), thiocillin I (TC), micrococcin P1

(MC), nosiheptide (NH), berninamycin A (BER), and geninthiocin A (GEN) (Figure 1).

SM differs from TS in that the two amino acids adjacent to the quinaldic acid moiety are dehydroalanine

and isoleucine instead of alanine and leucine. TC and MC are nearly identical except that TC has a

hydroxyvaline and MC has a valine in its macrocyclic ring. BER and GEN differ by a single methyl group on

Figure 2. Thiopeptides synergize with DSX against PA14 and PA14 *ΔfpvA fpvB*::Mar2xT7. Checkerboard assays were conducted in 10:90 medium. Top row are checkerboards conducted with each thiopeptide against wild type PA14 whereas bottom row are checkerboards conducted against its pyoverdine receptor-null mutant. The concentration of the thiopeptides are indicated along the column (maximum concentration of 10 µM in µg/mL except for SM which was tested at 5 µM) and the concentration of DSX is indicated along the row (maximum concentration of 64 µg/mL). Growth is proportional to the intensity of blue whereas white indicates no growth. Results are averaged from three independent experiments.

one of the oxazole rings. NH resembles TC and MC but is structurally unique from the other thiopeptides,

with an indole moiety linked to the core macrocyclic ring to form a secondary ring. TS, SM, TC, MP, and

NH belong to the 26-membered macrocycles whereas BER and GEN have 35-membered macrocycles.

Based on checkerboard assays, BER, GEN, and NH did not synergize with DSX (Supplementary

Figure 1). SM synergized with DSX against wild type PA14 but the combination had no effect against the

fpvAB mutant (Figure 2). Unlike the wild type, growth of the mutant was inhibited by 16 µg/ml DSX alone.

These data are similar to those obtained previously with TS + DSX[26] and indicates that SM likely crosses

the outer membrane using the pyoverdine receptors. However, the synergistic profile of SM + DSX was

less pronounced compared to TS + DSX. There are two potential explanations for its weaker synergistic

profile: 1) siomycin has less affinity for the target or 2) siomycin has less affinity for the receptors. To

distinguish between these possibilities, we tested each thiopeptdide against the Gram-positive pathogen

MRSA USA 300 in a broth dilution assay (Supplementary Figure 2). MRSA lacks the outer membrane;

therefore, the main consideration would be differences in affinity for the highly-conserved ribosomal

target. No differences between the minimal inhibitory concentration (MIC) of cells treated with TS or SM

were observed, suggesting that the differences in susceptibility in *P. aeruginosa* were related to receptor

affinity. Further, this result provides some structure-activity relationship (SAR) information regarding

interactions between TS and the pyoverdine receptors, specifically that the alanine and leucine adjacent

to the quinaldic acid do not affect target activity but improve uptake; however, they are not essential

because SM can still use the pyoverdine receptors to cross the outer membrane.

Thiocillin I and Micrococcin P1 Exploit the Ferrioxamine Receptor

In checkerboard assays, TC and MC synergized with DSX against *P. aeruginosa* PA14 (Figure 2).

The concentration of DSX was kept constant at 64 µg/mL, while the highest concentration of TC and MC

tested due to solubility limitations was 12 µg/mL (10 µM). Remarkably, this combination inhibited growth

of the *fpvAB* mutant (Figure 2). This result indicated that that TC and MC likely use a different siderophore

receptor to enter the cell. To identify that receptor, we screened a panel of 16 siderophore receptor-

deficient transposon mutants of PA14 (Supplementary Table 2). Only a *foxA* transposon mutant, lacking

the ferrioxamine receptor, displayed resistance to TC + DSX (Figure 3AB). When the mutant was

complemented with *foxA* in trans, susceptibility to TC was restored, confirming that *foxA* is necessary for

TC uptake (Figure 3C). The susceptibility of the complemented strain was slightly reduced compared to

the wild type, likely due to the low copy number of the vector in *P. aeruginosa*.

TC and MC inhibit the growth of TS-resistant clinical isolates

      In our previous studies, we identified *P. aeruginosa* clinical isolates resistant to the TS + DSX combination. One isolate, C0379, was resistant due to a 5' truncation of the gene encoding the FpvB receptor[26,29]. We hypothesized that TC or MC might inhibit the growth of this isolate because the only differences in its *foxA* gene compared to that of PAO1 were five silent mutations and a single base pair mutation (D760G) in one beta-strand belonging to a structural component of the receptor (Supplementary Table 3). Consistent with this idea, C0379 was susceptible to combinations of TC + DSX or MC + DSX (Supplementary Figure 3A). Bioinformatic analysis showed that *A. baumannii* also encodes the FoxA receptor. We tested one multidrug-resistant clinical isolate, C0286, for susceptibility to TC + DSX and found that it was susceptible even at the lowest concentration of TC tested (0.17 µg/mL) (Supplementary Figure 3B). However, the lab strain ATCC 17978 was not susceptible to the combination, suggesting inter-strain differences. This is not unusual as it is well established that lab adapted strains may not behave similarly to clinical isolates[32]. Overall, we conclude that TC + DSX can inhibit the growth of *P. aeruginosa* and *A. baumannii* clinical isolates including those resistant to TS.

Figure 3. TC crosses the outer membrane through the FoxA siderophore receptor. Wild type PA14 (A) and PA14 foxA::Mar2xT7 (B) were challenged with TC (pink) and TC + DSX (blue). Dose response assays were conducted in 10:90. The transposon mutant was complemented with wild type PA14 foxA (C) and challenged with TC in VBMM. The vector control is in black whereas the complemented mutant is in red. Results were averaged from three independent replicates. n.s, not significant; **, $p<0.01$; ***, $p<0.0005$; ****, $p<0.0001$. Significance was determined using two-way ANOVA followed by Sidak's multiple-comparison test.

A cocktail of TS + TC + DSX has an additive effect

Our observation that thiopeptides synergize with DSX against *P. aeruginosa* demonstrated the potential to target gram-negatives expressing either pyoverdine or ferrioxamine receptors using thiopeptide-chelator cocktails. Sub-MIC combinations of TS (0.1 μg/mL) + 64 μg/mL DSX and TC (0.2 μg/mL) + 64 μg/mL DSX, as well as TS + TC + DSX were tested against PA14 in VBMM (Figure 4). Each double combination inhibited growth to ~30% of control; however, the triple combination decreased the growth to only 10% of control. A multiplicative rule approach was used to determine the effect of combining both thiopeptides where the expected growth is the product of the growth inhibition for the individual combinations (i.e. TS + DSX or TC + DSX). For TS + DSX, growth was reduced to 31% of control and for TC + DSX, growth was reduced to

Figure 4. A thiopeptide cocktail significantly reduces PA14 growth. Control growth less than 20% was considered as the MIC cut-off. Assays were conducted in VBMM. Results are averaged from three independent experiments. ****,p<0.0001. Significance was determined using one-way ANOVA followed by Tukey honest significant difference post test.

29% of control. Therefore, the expected growth for the combination of TC + TS + DSX would be 9% of control. DSX alone had no effect on growth. The experimental value yielded 8.8% of control, suggesting an additive effect of the thiopeptides. These results indicate that thiopeptide cocktails could offer an advantage compared over single thiopeptide treatment by using different siderophore receptors to enter the cell.

Thiopeptides target specific bacterial genera and species

Unlike the pyoverdine receptors, the ferrioxamine receptor is not unique to Pseudomonads. Therefore, we tested whether other Gram-negative bacteria were susceptible to TC. Bioinformatic analyses of other priority Gram-negative pathogens such as *Escherichia coli, Klebsiella pneumoniae,* and *Salmonella* Typhimurium showed that they carry *foxA* in their genome. However, they were not

susceptible to TC or the combination of TC + DSX

at the highest concentrations tested, suggesting

that the receptors are sufficiently different in

sequence to prevent recognition of TC as a

ligand. FoxA proteins from *S.* Typhimurium

SL1344 and *P. aeruginosa* PAO1 share 40% amino

acid sequence identity (Supplementary Figure 4).

Residues Y218, H374, and Q441 that interact

with the native ferroxamine ligand are conserved

in both species, indicating that TC and MC must

rely on interactions with other amino acids for

uptake into Pseudomonads[33].

To determine if there were differences

in susceptibilities between different species of

Pseudomonas, we conducted dose-response

assays with TC and TC + DSX against *P. putida*

KT2440, *P. marginalis* CVCOB 1152, *P. protegens*

Pf-5, and *P. fluorescens* PV5(Supplementary

Figure 5. Similar experiments were done with TS

and TS + DSX. We defined resistance if the

percent of control growth exceeded 80%,

susceptible if below 20%, and intermediate if

between 20-80%. Interestingly, there was

variability in the susceptibility to the

Figure 5. Susceptibility to TS and TC are different between species and strains. Susceptibility of *Pseudomonas* and *Acinetobacter* to TS and TC in combination with DSX. Blue boxes indicate resistance (>80% of control growth). Yellow boxes indicate intermedicate susceptibility (20-80% of control growth). Orange boxes indicate susceptible (<20% of control growth).

combinations between species. *P. putida* was resistant to TS but susceptible to TC (Figure 5,

Supplementary Figure 5). *P. protegens* was resistant to all combinations (Figure 5, Supplementary Figure

5). Alignments of the FoxA sequences for each species showed 65%, 65%, and 60% similarity for *P.*

fluorescens, P. protegens, and *P. putida* compared to *P. aeruginosa* PAO1. Despite the differences in

susceptibility to TC + DSX, the key residues that interact with the native ligand are conserved, similar to *S.*

Typhimurium, suggesting that TC interacts with other amino acids. To focus on possible sites of

interaction, we looked for residues shared between *P. aeruginosa*, *P. putida*, and *P. fluorescens*, but not

P. protegens, which was resistant to TC and TC + DSX (Figure 5, Supplementary Figure 5). A total of 17

residues were identified following this criterion and mapped onto a co-crystal structure of FoxA bound to

ferrioxamine (PDB ID: 6I96)[33]. Only three were found in proximity to the native ligand – A620(P), S612(A),

and D380(N) (Supplementary Table 4). A629(P) indicates that in the FoxA receptor of *P. aeruginosa*, *P.*

putida, and *P. fluorescens* at position 620, the amino acid is alanine. However, in *P. protegens* the amino

acid at the same position is proline. A620(P) and S612(A) are located in loop 7 of FoxA, which closes upon

interaction of the ligand with the active site[33]. Closure of loop 7 prevents the native ligand from

dissociation. D380(N) is close to conserved residue H374, which interacts with the native ligand.

A BLAST search using the PAO1 FoxA amino acid sequence against *P. aeruginosa, P. putida, P.*

fluorescens (all TC susceptible under low iron)*,* and *P. protegens* (TC resistant) was conducted to see if

A620(P), S612(A), and D380(N) are conserved (Supplementary Figure 6). The top 100 hits were aligned

(MUSCLE), which revealed that alanine at position 620 and aspartic acid at position 380 are well conserved

in *P. aeruginosa, P. fluorescens,* and *P. putida*. Alanine was the most common amino acid at position 612

in *P. fluorescens* rather than serine, suggesting that this residue may not be important for TC interaction

with FoxA since this species was susceptible to TC + DSX. In *P. protegens,* which is resistant to TC + DSX,

lysine was the most abundant at residue 620. At residue 380, glutamine occurred more frequently. These

differences suggest that alanine and aspartic acid at residues 620 and 380 respectively may be involved

in TC uptake. These data suggest that TC uptake requires specific amino acid interactions that differ

between species.

Nosiheptide is a potent inhibitor of MRSA USA 300

The core structure of NH is nearly identical to TC and MC; however, a single indole is linked to the

core 26-membered macrocyclic ring (Figure 1). This small change in structure results in loss of synergy

with DSX (Supplementary Figure 1). Although synergy was not observed, there was a reduction in growth

when NH was combined with DSX. When tested against the *fpvAB* mutant, potentiation was lost

(Supplementary Figure 1).

There are examples – such as psychotrimine and daptomycin – of natural product antibiotics

containing tryptophan or its side chain, indole [34]. Daptomycin is a Gram-positive antibiotic that disrupts

the cytoplasmic membrane[35]. Like NH, daptomycin is a bulky molecule whose size prevents it from

crossing the outer membrane, making it ineffective against Gram-negative bacteria. Based on its unique

indole content, we hypothesized that NH might have improved efficacy against Gram-positive bacteria

compared to the other thiopeptides tested. All seven thiopeptides were tested in a dose-response assay

against MRSA USA 300, with 260 ng/mL as the highest concentration (Supplementary Figure 2). Consistent

with our hypothesis, NH was the most potent against MRSA, with a MIC of <4 ng/mL, followed by TS and

SM (32 ng/mL), TC and MC (130 ng/mL), and BER and GEN (>260 ng/mL). These results show that the

indole moiety of NH may confer an advantage against Gram-positives compared to other thiopeptides.

DISCUSSION

We previously showed that TS hijacked the pyoverdine receptors of *P. aeruginosa* and *A.*

baumannii to cross the outer membrane. These data informed the hypothesis that other thiopeptides

might similarly act as Trojan Horse antibiotics. In this work, we explored the ability of six other

thiopeptides to exploit siderophore receptors under iron-limited conditions. We showed that SM – a

thiopeptide structurally similar to TS – also uses pyoverdine receptors to enter cells. The differences in

structure of the two compounds provided SAR information. The IC_{50} of TS and SM are similar (0.1 µM vs

0.25 µM) based on *in vitro* studies of *E. coli* ribosomal activity[36]. This suggests that SM has weaker synergy

with DSX because of reduced affinity for the receptor compared to TS. Combinatorial analyses by

checkerboard assays suggested that TC and MC used a different siderophore receptor. We screened

mutants lacking various receptors in low iron growth media, leading to the discovery that those

thiopeptides used FoxA, a xenosiderophore receptor in *P. aeruginosa*. TC was also effective against a TS-

resistant *P. aeruginosa* clinical isolate. However, despite widespread expression of FoxA homologues,

susceptibility to TC was not universal among Gram-negative bacteria. Even within the genus

Pseudomonas, there was variability in susceptibility between species. Differences in the levels of

siderophore receptor expression could be a factor in the susceptibility profiles as a higher abundance

would allow more thiopeptide uptake.

This work suggests links between thiopeptide structure and activity against specific Gram-

negatives. Thiopeptides were thought to have little or no activity against Gram-negative pathogens due

to their inability to cross the outer membrane. However, thiopeptide producers are mainly marine and

soil bacteria, which inhabit environments shared by Pseudomonads. We propose that thiopeptide

structures evolved to target the siderophore receptors of cohabiting Gram-negatives. This is not

unprecedented as many other natural products can use such receptors to cross the outer membrane. For

example, *P. aeruginosa* produces pyocins that target the pyoverdine receptors of competing strains[37]..

NH failed to synergize with DSX, although the combination reduced growth. Potentiation was lost against

the *fpvAB* mutant. This result suggests that NH could have poor affinity for pyoverdine receptors, but

further work needs to be done to demonstrate this. We also showed that of the thiopeptides tested, NH

had the greatest activity against MRSA USA 300. The exact mechanism remains unknown, but while the

core of NH is nearly identical to TC and MC, the indole moiety makes it unique compared to the other

thiopeptides. NH does not synergize with DSX which suggests that the indole moiety could interfere with

its recognition by the FoxA receptor. Tryptophan and indole increase interactions of peptides with

membranes and can act as signaling molecules[38–41]. We speculate that the indole moiety may provide

enhanced peptide-membrane interactions, facilitating the ability of NH to cross the membrane and target

Gram-positive pathogens like MRSA. We observed a lower MIC with NH (<4 ng/mL) compared to TS (32

ng/mL) against MRSA in 10:90 medium, consistent with results from other groups using Mueller-Hinton

Broth[1,42]. In contrast, assays using purified ribosomes showed that TS had 40% greater inhibitory activity

compared to NH[43]. While NH has weaker inhibitory activity against purified ribosomes, it may compensate

with improved membrane permeability. NH has been tested against MRSA in an intraperitoneal mouse

infection model[1]. Mice treated with 20 mg/kg NH showed 90% survival over 5 days compared to 40% in

the vehicle control group.

Comparing TC and TS, it is apparent that the core macrocycle is nearly identical. However, TS has

a second macrocyclic ring with a structurally-distinct quinaldic acid moiety. It is likely that this second ring

allows TS to interact with the pyoverdine receptors rather than FoxA. In contrast, BER and GEN had no

activity against *P. aeruginosa* under iron-limitation. Even against MRSA, these 35-membered macrocycle

thiopeptides had no activity at the highest concentration tested (260 ng/mL). This is unsurprising as it was

previously reported that BER has weaker inhibitory activity on purified ribosomes compared to TS[6]. Our

results do not rule out the potential of 35-membered thiopeptides having other advantages. Furthermore,

in this study we did not test 29-membered thiopeptides such as GE2270A, thiomuracin A, and GE37468[44–

46]. These molecules could also have potential activity against *P. aeruginosa* and *A. baumannii* under iron-

limited conditions.

Schwalen et al. recently used a bioinformatics approach to search for enzymes that catalyze non-

spontaneous [4+2]-cycloadditions characteristic of thiopeptide biosynthesis. They identified 508 [4+2]-

cycloaddition enzymes, 372 of which were unique at the amino acid level[31]. These results suggest that

there are many more structurally unique thiopeptides than previously estimated. Diverse thiopeptides

may reveal interesting new activities with antimicrobial applications. One example is cyclothioazomycin,

a structurally distinct compound with activity against *Bacillus* spp., although weaker relative to other thiopeptides[47]. Interestingly, this thiopeptide has potent antifungal activity via its interactions with chitin[48]. It also was reported to be a human plasma renin inhibitor[49].

Trojan horse antibiotics are an interesting avenue for drug development. They have advantages over antibiotics that must simply diffuse through the outer membrane because of their greater specificity and uptake. Moreover, sites of infection are frequently iron-limited, which naturally potentiates such molecules. Hosts produce iron sequestration proteins such as transferrin and lactoferrin that restrict iron, a process called nutritional immunity[50–52].

While thiopeptides been overlooked for use in the clinic, we show the potential for these antibiotics to be developed as selective Gram-negative antimicrobials that can target high priority pathogens including *P. aeruginosa* and *A. baumannii*. Our work also supports the future development of NH as an anti-MRSA therapeutic.

METHODS

Chemicals

Thiopeptides were purchased from Cayman Chemicals and stored at -20°C. Thiopeptide purity was at least ≥95%. All thiopeptides were dissolved in dimethylsulfoxide (DMSO) with no solubility issues ranging from 1mg/mL to 20mg/mL. Deferasirox was purchased from AK Scientific (98% purity). A stock solution of 20 mg/mL in DMSO was made. Stock solutions were stored at -20°C until use and all serial dilutions were prepared using fresh DMSO at 75x the final concentration.

Bacterial Strains and Culture Conditions

All bacterial strains used in this study are listed in Supplementary Table 1. Bacteria were inoculated from glycerol stocks stored at -80°C and grown overnight in LB at 37°C with shaking (200 rpm). Bacteria were subcultured (1/500 dilution) into fresh 10:90 media or Vogel-Bonner Minimal Media (VBMM) and grown for 16 hours. Optical densities were standardized to OD_{600} 0.1 and diluted 1/500 into fresh 10:90 or VBMM, which was used in microbroth dilution MIC assays or checkerboard assays.

Generation of PA14 f*oxA*::Mar2xT7 pUCP20-*foxA*

foxA was PCR amplified from wild type PA14 genomic DNA prepared using Qiagen genomic DNA isolation

kit. The following forward and reverse primers were used to obtain wild type *foxA* with SacI and Xbal

restriction sites respectively (underlined): forward 5'GAAG<u>GAGCTC</u>GTGGACGCTTGCTTTCGT and reverse

5'GCGT<u>TCTAGA</u>GACGCCGGCGAATGCCC. The PCR product and pUCP20 were digested with SacI and Xbal

(Thermo Scientific) using Thermo Scientific Fast Digest. The products were run on a 1% agarose gel for

30mins at 120V, cut out, and purified using GeneJET Extraction Kit (Thermo Scientific). The purified

linearized vector and digested *foxA* gene were ligated (1:3 molar ratio) overnight at 4°C using T4 DNA

ligase (Thermo Scientific). The construct was transformed into competent *E. coli* DH5α by heat shock at

42°C and plated on LB plates with 100 µg/mL ampicillin and 5-bromo-4-chloro-3-indolyl-β-D-

galactopyranoside (IPTG) for blue-white screening (BioShop). Plates were incubated at 37°C overnight.

White single colonies were streaked on fresh LB plates with ampicillin to confirm that colonies contained

the construct. Minipreps were done using GeneJET Plasmid Miniprep Kit (Thermo Scientific) and

quantified by NanoDrop. The *foxA* transposon mutant was grown overnight in LB. 1 mL was centrifuged

and washed twice with nuclease-free water (Qiagen). The cell pellet was resuspended in 500 µL of

nuclease-free water. 100 ng of the construct was added to 100 µL of resuspended cells and electroporated

(2.50 V). After a recovery period of 2 hours, cells were plated onto LB agar with 200 µg/mL carbenicillin

(Bioshop). Plates were incubated at 37°C overnight. Single colonies were streaked on fresh plates to

confirm successful electroporation. A single colony was picked and grown in liquid for MIC assays. The

susceptibility of the *foxA* transposon mutant harbouring pUCP20-*foxA* to TC indicated successful

complementation. Similarly, the same *foxA* transposon mutant was electroporated with empty pUCP20

as vector control.

Microbroth Dilution and Checkerboard Assays

Stock solutions were diluted in DMSO and 2µL of the resulting solutions plus 148µL of a bacterial

suspension standardized to an OD600 of ~ 0.1 and diluted 1:500 in 10:90 was added to a 96 well plate

(Nunc) in triplicate. Control wells contained 148µL of 10:90 + 2µL DMSO (sterility control) or standardized

bacterial suspension + 2µL DMSO (growth control). After incubating at 37°C for 16h, with shaking at 200

rpm, the OD600 was read using a plate reader (Thermo Scientific). Assays were performed in triplicate,

calculations were done in Microsoft Excel, and results were graphed using Prism (GraphPad) as a

percentage of the DMSO control. For wells with 2 compounds, such as increasing TS concentrations with

64µg/mL of DSX, 4µL of DMSO was used in the sterility and growth controls.

Checkerboard assays were set-up using Nunc 96-well plates in an 8-well by 8-well format. Two columns

were allocated for vehicle controls and two columns for sterility controls. Vehicle controls contained 4 µL

DMSO, in 146 µL of 10:90 inoculated with PA14 as described in Growth Curves. Sterile controls contained

the same components in 10:90, without cells. Serial dilutions of each thiopeptide – with 10 µM being the

highest final concentration – was added along the ordinate of the checkerboard (increasing concentration

from bottom to top) whereas serial dilutions of DSX – with 64 µg/mL being the highest final concentration

– was added along the abscissa (increasing concentration from left to right). The final volume of each well

was 150 µL and each checkerboard was repeated at least three times. Plates were incubated and the final

OD600 determined as detailed above. Checkerboards were graphed as heatmaps in Prism (GraphPad).

Bioinformatic Analysis

MUSCLE alignments were conducted using Unipro UGENE and Geneious Prime. Accession numbers of *P.*

aeruginosa, P. fluorescens, P. protegens, P. putida, and *S.* Typimurium FoxA are as follows: AAG05854.1,

WP_122305244.1, WP_003218248.1, AAY95537.2, NP_742329.1, and CBW16459.1. Genomic analyses for

mutations in WCC C0379 were done with breseq by Kara Tsang using *P. aeruginosa* PAO1 as the reference

genome.

ASSOCIATED CONTENT

Supporting Information

Table S1, list of strains; Table S2, mutants tested to show that TC uses the FoxA receptor to cross the outer

membrane; Table S3, breseq genomic annotations of the FoxA receptor for WCC C0379; Table S4, FoxA

amino acid residues that are different in *P. protegens* compared to other *Pseudomonas* spp; Figure S1,

checkerboard assays for thiopeptides that did not synergize with DSX; Figure S2, thiopeptide activity

against MRSA USA300; Figure S3, thiopeptide activity against *P. aeruginosa* and *A. baumannii* TS-resistant

clinical isolates; Figure S4, MUSCLE alignment of *S.* Typhimurium and *P. aeruginosa* FoxA receptors; Figure

S5, dose-response assays of thiopeptides against different *Pseudomonas* spp. Figure S6, Sequence logos

of FoxA amino acid sequences of *Pseudomonas* spp.

AUTHOR INFORMATION

Corresponding Author

* lori.burrows@mcmaster.ca

ORCID

Lori L. Burrows: 0000-0003-0838-5040

Author Contributions

DCKC performed all experiments. DCKC and LLB wrote the paper.

Funding Sources

This work was funded by grants to LLB from the Natural Sciences and Engineering Research Council (NSERC) RGPIN-2016-06521 and the Ontario Research Fund RE07-048. DCKC holds an NSERC Canadian Graduate Scholarship – Master's Award.

ACKNOWLEDGMENTS

We thank Gerry Wright for strains from the Wright Clinical Collection, as well as Eric Brown and Brian Coombes for the following strains: MRSA USA 300 and *K. pneumoniae*, and S. Typhimurium SL1344. We thank John Whitney for the following strains: *P. protegens* Pf-5, and *P. putida* KT2440 and David Guttman for *P. fluorescens*. We thank Kara Tsang for her help with breseq analyses. We thank Christy Groves for assistance with the graphical abstract.

ABBREVIATIONS

TS, thiostrepton; TC, thiocillin; MC, micrococcin; NH, nosiheptide; MIC, minimal inhibitory concentration; SM, siomycin; BER, berninamycin; GEN, geninthiocin; DSX, deferasirox; FpvA, ferripyoverdine receptor A; FpvB, ferripyoverdine receptor B; FoxA, ferrioxamine receptor; WCC, Wright Clinical Collection; VBMM, Vogel-Bonner Minimal Media; MRSA, methicillin-resistant *Staphylococcus aureus*; *E. coli*, *Escherichia coli*; *K. pneumoniae*, *Klebsiella pneumoniae*; *S.* Typhimurium, *Salmonella* Typhimurium; *P. aeruginosa*, *Pseudomonas aeruginosa*; *P. fluorescens*, *Pseudomonas fluorescens*; *P. putida*, *Pseudomonas putida*; *P. protegens*, *Pseudomonas protegens*; *A. baumannii*, *Acinetobacter baumannii*.

REFERENCES

(1)   Haste, N. M.; Thienphrapa, W.; Tran, D. N.; Loesgen, S.; Sun, P.; Nam, S. J.; Jensen, P. R.; Fenical, W.; Sakoulas, G.; Nizet, V.; Hensler, M. E. Activity of the Thiopeptide Antibiotic Nosiheptide against Contemporary Strains of Methicillin-Resistant Staphylococcus Aureus. *J. Antibiot. (Tokyo).* **2012**, *65* (12), 593–598. https://doi.org/10.1038/ja.2012.77.

(2)   Bagley, M. C.; Dale, J. W.; Merritt, E. A.; Xiong, X. Thiopeptide Antibiotics. *Chemical Reviews*. 2005, pp 685–714. https://doi.org/10.1021/cr0300441.

(3)   Walter, J. D.; Hunter, M.; Cobb, M.; Traeger, G.; Spiegel, P. C. Thiostrepton Inhibits Stable 70S Ribosome Binding and Ribosome-Dependent GTPase Activation of Elongation Factor G and Elongation Factor 4. *Nucleic Acids Res.* **2012**, *40* (1), 360–370. https://doi.org/10.1093/nar/gkr623.

(4)   Bowen, W. S.; Van Dyke, N.; Murgola, E. J.; Lodmell, J. S.; Hill, W. E. Interaction of Thiostrepton and Elongation Factor-G with the Ribosomal Protein L11-Binding Domain. *J. Biol. Chem.* **2005**, *280* (4), 2934–2943. https://doi.org/10.1074/jbc.M407008200.

(5)   Heffron, S. E.; Jurnak, F. Structure of an EF-Tu Complex with a Thiazolyl Peptide Antibiotic Determined at 2.35 Å Resolution: Atomic Basis for GE2270A Inhibition of EF- Tu. *Biochemistry* **2000**, *39* (1), 37–45. https://doi.org/10.1021/bi9913597.

(6)   Thompson, J.; Cundliffe, E.; Stark, M. J. R. The Mode of Action of Berninamycin and the Mechanism of Resistance in the Producing Organism, Streptomyces Bernensis. *J. Gen. Microbiol.* **1982**, *128* (4), 875–884. https://doi.org/10.1099/00221287-128-4-875.

(7)   Hegde, N. S.; Sanders, D. A.; Rodriguez, R.; Balasubramanian, S. The Transcription Factor FOXM1 Is a Cellular Target of the Natural Product Thiostrepton. *Nat. Chem.* **2011**, *3* (9), 725–731. https://doi.org/10.1038/nchem.1114.

(8)   Aminake, M. N.; Schoof, S.; Sologub, L.; Leubner, M.; Kirschner, M.; Arndt, H. D.; Pradel, G. Thiostrepton and Derivatives Exhibit Antimalarial and Gametocytocidal Activity by Dually Targeting Parasite Proteasome and Apicoplast. *Antimicrob. Agents Chemother.* **2011**, *55* (4), 1338–1348. https://doi.org/10.1128/AAC.01096-10.

(9)   SU, T. L. Micrococcin, an Antibacterial Substance Formed by a Strain Of. *Br. J. Exp. Pathol.* **1948**, *29* (5), 473–481.

(10)   Li, J.; Qu, X.; He, X.; Duan, L.; Wu, G.; Bi, D.; Deng, Z.; Liu, W.; Ou, H. Y. ThioFinder: A Web-Based
Tool for the Identification of Thiopeptide Gene Clusters in DNA Sequences. *PLoS One* **2012**, *7* (9).
https://doi.org/10.1371/journal.pone.0045878.

(11)   Arnison, P. G.; Bibb, M. J.; Bierbaum, G.; Bowers, A. A.; Bugni, T. S.; Bulaj, G.; Camarero, J. A.;
Campopiano, D. J.; Challis, G. L.; Clardy, J.; Cotter, P. D.; Craik, D. J.; Dawson, M.; Dittmann, E.; Donadio,
S.; Dorrestein, P. C.; Entian, K. D.; Fischbach, M. A.; Garavelli, J. S.; Göransson, U.; Gruber, C. W.; Haft, D.
H.; Hemscheidt, T. K.; Hertweck, C.; Hill, C.; Horswill, A. R.; Jaspars, M.; Kelly, W. L.; Klinman, J. P.; Kuipers,
O. P.; Link, A. J.; Liu, W.; Marahiel, M. A.; Mitchell, D. A.; Moll, G. N.; Moore, B. S.; Müller, R.; Nair, S. K.;
Nes, I. F.; Norris, G. E.; Olivera, B. M.; Onaka, H.; Patchett, M. L.; Piel, J.; Reaney, M. J. T.; Rebuffat, S.; Ross,
R. P.; Sahl, H. G.; Schmidt, E. W.; Selsted, M. E.; Severinov, K.; Shen, B.; Sivonen, K.; Smith, L.; Stein, T.;
Süssmuth, R. D.; Tagg, J. R.; Tang, G. L.; Truman, A. W.; Vederas, J. C.; Walsh, C. T.; Walton, J. D.; Wenzel,
S. C.; Willey, J. M.; Van Der Donk, W. A. Ribosomally Synthesized and Post-Translationally Modified
Peptide Natural Products: Overview and Recommendations for a Universal Nomenclature. *Natural*
Product Reports. Royal Society of Chemistry 2013, pp 108–160. https://doi.org/10.1039/c2np20085f.

(12)   Brown, L. C. W.; Acker, M. G.; Clardy, J.; Walsh, C. T.; Fischbach, M. A. Thirteen Posttranslational
Modifications Convert a 14-Residue Peptide into the Antibiotic Thiocillin. *Proc. Natl. Acad. Sci. U. S. A.*
2009, *106* (8), 2549–2553. https://doi.org/10.1073/pnas.0900008106.

(13)   Khilyas, I. V.; Tursunov, K. A.; Shirshikova, T. V.; Kamaletdinova, L. K.; Matrosova, L. E.; Desai, P.
T.; McClelland, M.; Bogomolnaya, L. M. Genome Sequence of Pigmented Siderophore-Producing Strain
Serratia Marcescens SM6. *Microbiol. Resour. Announc.* **2019**, *8* (18). https://doi.org/10.1128/mra.00247-
19.

(14)   Cheng, T. H.; Saidin, J.; Danish-Daniel, M.; Gan, H. M.; Isa, M. N. M.; Bakar, M. F. A.; Ismail, N.
Genome Sequence of Serratia Marcescens Subsp. Sakuensis Strain K27, a Marine Bacterium Isolated from
Sponge   (Haliclona   Amboinensis).   *Genome   Announc.*   **2018**,   *6*   (6).
https://doi.org/10.1128/genomeA.00022-18.

(15)   Engelhardt, K.; Degnes, K. F.; Kemmler, M.; Bredholt, H.; Fjaervik, E.; Klinkenberg, G.; Sletta, H.;
Ellingsen, T. E.; Zotchev, S. B. Production of a New Thiopeptide Antibiotic, TP-1161, by a Marine
Nocardiopsis   Species.   *Appl.   Environ.   Microbiol.*   **2010**,   *76*   (15),   4969–4976.
https://doi.org/10.1128/AEM.00741-10.

(16)   Trejo, W. H.; Dean, L. D.; Pluscec, J.; Meyers, E.; Brown, W. E. Streptomyces Laurentii, a New

Species   Producing   Thiostrepton. *J. Antibiot. (Tokyo).* **1977**, *30* (8), 639–643.

https://doi.org/10.7164/antibiotics.30.639.

(17)   Acker, M. G.; Bowers, A. A.; Walsh, C. T. Generation of Thiocillin Variants by Prepeptide Gene

Replacement and in Vivo Processing by Bacillus Cereus. *J. Am. Chem. Soc.* **2009**, *131* (48), 17563–17565.

https://doi.org/10.1021/ja908777t.

(18)   Akasapu, S.; Hinds, A. B.; Powell, W. C.; Walczak, M. A. Total Synthesis of Micrococcin P1 and

Thiocillin   I   Enabled   by   Mo(vi)   Catalyst. *Chem. Sci.* **2019**, *10* (7), 1971–1975.

https://doi.org/10.1039/C8SC04885A.

(19)   Luo, X.; Zambaldo, C.; Liu, T.; Zhang, Y.; Xuan, W.; Wang, C.; Reed, S. A.; Yang, P. Y.; Wang, R. E.;

Javahishvili, T.; Schultz, P. G.; Young, T. S. Recombinant Thiopeptides Containing Noncanonical Amino

Acids.   *Proc.   Natl.   Acad.   Sci.   U.   S.   A.* **2016**, *113* (13), 3615–3620.

https://doi.org/10.1073/pnas.1602733113.

(20)   Wever, W. J.; Bogart, J. W.; Baccile, J. A.; Chan, A. N.; Schroeder, F. C.; Bowers, A. A.

Chemoenzymatic Synthesis of Thiazolyl Peptide Natural Products Featuring an Enzyme-Catalyzed Formal

[4 + 2] Cycloaddition. *J. Am. Chem. Soc.* **2015**, *137* (10), 3494–3497. https://doi.org/10.1021/jacs.5b00940.

(21)   Bogart, J. W.; Bowers, A. A. Thiopeptide Pyridine Synthase TbtD Catalyzes an Intermolecular

Formal   Aza-Diels-Alder   Reaction. *J.   Am.   Chem.   Soc.* **2019**, *141* (5), 1842–1846.

https://doi.org/10.1021/jacs.8b11852.

(22)   Liao, R.; Duan, L.; Lei, C.; Pan, H.; Ding, Y.; Zhang, Q.; Chen, D.; Shen, B.; Yu, Y.; Liu, W. Thiopeptide

Biosynthesis Featuring Ribosomally Synthesized Precursor Peptides and Conserved Posttranslational

Modifications. *Chem. Biol.* **2009**, *16* (2), 141–147. https://doi.org/10.1016/j.chembiol.2009.01.007.

(23)   Tran, H. L.; Lexa, K. W.; Julien, O.; Young, T. S.; Walsh, C. T.; Jacobson, M. P.; Wells, J. A. Structure-

Activity Relationship and Molecular Mechanics Reveal the Importance of Ring Entropy in the Biosynthesis

and   Activity   of   a   Natural   Product. *J.   Am.   Chem.   Soc.* **2017**, *139* (7), 2541–2544.

https://doi.org/10.1021/jacs.6b10792.

(24)   Wang, J.; Lin, Z.; Bai, X.; Tao, J.; Liu, W. Optimal Design of Thiostrepton-Derived Thiopeptide

Antibiotics and Their Potential Application against Oral Pathogens. *Org. Chem. Front.* **2019**, *6* (8), 1194–

1199. https://doi.org/10.1039/c9qo00219g.

(25)   Kelly, W. L.; Pan, L.; Li, C. Thiostrepton Biosynthesis: Prototype for a New Family of Bacteriocins.
J. Am. Chem. Soc. **2009**, *131* (12), 4327–4334. https://doi.org/10.1021/ja807890a.

(26)   Ranieri, M. R. M.; Chan, D. C. K.; Yaeger, L. N.; Rudolph, M.; Karabelas-Pittman, S.; Abdo, H.; Chee,
J.; Harvey, H.; Nguyen, U.; Burrows, L. L. Thiostrepton Hijacks Pyoverdine Receptors to Inhibit Growth of
Pseudomonas   Aeruginosa.   *Antimicrob.   Agents   Chemother.*   **2019**,   *63*   (9).
https://doi.org/10.1128/AAC.00472-19.

(27)   Tacconelli, E., Magrini, N., Kahlmeter, G. & Singh, N. Global Priority List of Antibiotic-Resistant
Batceria to Guide Research, Discovery, and Development of New Antibiotics. *WHO Publ.* **2017**, 7.

(28)   National Nosocomial Infection Surveillance System. National Nosocomial Infections Surveillance
(NNIS) System Report, Data Summary from January 1992 through June 2004, Issued October 2004.
American Journal of Infection Control. Mosby Inc. December 1, 2004, pp 470–485.
https://doi.org/10.1016/j.ajic.2004.10.001.

(29)   Chan, D. C. K.; Guo, I.; Burrows, L. L. Forging New Antibiotic Combinations under Iron-Limiting
Conditions. *Antimicrob. Agents Chemother.* **2020**, *64* (3). https://doi.org/10.1128/AAC.01909-19.

(30)   Merwin, N. J.; Mousa, W. K.; Dejong, C. A.; Skinnider, M. A.; Cannon, M. J.; Li, H.; Dial, K.;
Gunabalasingam, M.; Johnston, C.; Magarvey, N. A. DeepRiPP Integrates Multiomics Data to Automate
Discovery of Novel Ribosomally Synthesized Natural Products. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117* (1),
371–380. https://doi.org/10.1073/pnas.1901493116.

(31)   Schwalen, C. J.; Hudson, G. A.; Kille, B.; Mitchell, D. A. Bioinformatic Expansion and Discovery of
Thiopeptide   Antibiotics.   *J.   Am.   Chem.   Soc.*   **2018**,   *140*   (30),   9494–9501.
https://doi.org/10.1021/jacs.8b03896.

(32)   Di Venanzio, G.; Flores-Mireles, A. L.; Calix, J. J.; Haurat, M. F.; Scott, N. E.; Palmer, L. D.; Potter, R.
F.; Hibbing, M. E.; Friedman, L.; Wang, B.; Dantas, G.; Skaar, E. P.; Hultgren, S. J.; Feldman, M. F. Urinary
Tract Colonization Is Enhanced by a Plasmid That Regulates Uropathogenic Acinetobacter Baumannii
Chromosomal Genes. *Nat. Commun.* **2019**, *10* (1). https://doi.org/10.1038/s41467-019-10706-y.

(33)   Josts, I.; Veith, K.; Tidow, H. Ternary Structure of the Outer Membrane Transporter FoxA with
Resolved Signaling Domain Provides Insights into TonB-Mediated Siderophore Uptake. *Elife* **2019**, *8*.
https://doi.org/10.7554/eLife.48528.

(34)　Schallenberger, M. A.; Newhouse, T.; Baran, P. S.; Romesberg, F. E. The Psychotrimine Natural

Products Have Antibacterial Activity against Gram-Positive Bacteria and Act via Membrane Disruption. *J.*

Antibiot. (Tokyo). **2010**, *63* (11), 685–687. https://doi.org/10.1038/ja.2010.107.

(35)　Randall, C. P.; Mariner, K. R.; Chopra, I.; O'Neill, A. J. The Target of Daptomycin Is Absent from

Escherichia Coli and Other Gram-Negative Pathogens. *Antimicrob. Agents Chemother.* **2013**, *57* (1), 637–

639. https://doi.org/10.1128/AAC.02005-12.

(36)　Lentzen, G.; Klinck, R.; Matassova, N.; Aboul-Ela, F.; Murchie, A. I. H. Structural Basis for

Contrasting Activities of Ribosome Binding Thiazole Antibiotics. *Chem. Biol.* **2003**, *10* (8), 769–778.

https://doi.org/10.1016/S1074-5521(03)00173-X.

(37)　White, P.; Joshi, A.; Rassam, P.; Housden, N. G.; Kaminska, R.; Goult, J. D.; Redfield, C.; McCaughey,

L. C.; Walker, D.; Mohammed, S.; Kleanthous, C. Exploitation of an Iron Transporter for Bacterial Protein

Antibiotic Import. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114* (45), 12051–12056.

https://doi.org/10.1073/pnas.1713741114.

(38)　Chan, D. I.; Prenner, E. J.; Vogel, H. J. Tryptophan- and Arginine-Rich Antimicrobial Peptides:

Structures and Mechanisms of Action. *Biochimica et Biophysica Acta - Biomembranes*. Elsevier September

1, 2006, pp 1184–1202. https://doi.org/10.1016/j.bbamem.2006.04.006.

(39)　Gaede, H. C.; Yau, W. M.; Gawrisch, K. Electrostatic Contributions to Indole-Lipid Interactions. *J.*

Phys. Chem. B **2005**, *109* (26), 13014–13023. https://doi.org/10.1021/jp0511000.

(40)　Lee, J. H.; Lee, J. Indole as an Intercellular Signal in Microbial Communities. *FEMS Microbiology*

Reviews. Blackwell Publishing Ltd July 1, 2010, pp 426–444. https://doi.org/10.1111/j.1574-

6976.2009.00204.x.

(41)　Zhu, X.; Ma, Z.; Wang, J.; Chou, S.; Shan, A. Importance of Tryptophan in Transforming an

Amphipathic Peptide into a Pseudomonas Aeruginosa-Targeted Antimicrobial Peptide. *PLoS One* **2014**, *9*

(12), e114605. https://doi.org/10.1371/journal.pone.0114605.

(42)　Niu, H.; Yee, R.; Cui, P.; Tian, L.; Zhang, S.; Shi, W.; Sullivan, D.; Zhu, B.; Zhang, W.; Zhang, Y.

Identification of Agents Active against Methicillin- Resistant Staphylococcus Aureus Usa300 from a Clinical

Compound Library. *Pathogens* **2017**, *6* (3). https://doi.org/10.3390/pathogens6030044.

(43)　Cundliffe, E.; Thompson, J. The Mode of Action of Nosiheptide (Multhiomycin) and the Mechanism

of Resistance in the Producing Organism. *J. Gen. Microbiol.* **1981**, *126* (1), 185–192.

https://doi.org/10.1099/00221287-126-1-185.

(44)   Stella, S.; Montanini, N.; Le Monnier, F.; Ferrari, P.; Colombo, L.; Landini, P.; Ciciliato, I.; Goldstein,

B. P.; Selva, E.; Denaro, M. Antibiotic GE37468 A: A New Inhibitor of Bacterial Protein Synthesis: I. Isolation

and Characterization. *J. Antibiot. (Tokyo).* **1995**, *48* (8), 780–786.

https://doi.org/10.7164/antibiotics.48.780.

(45)   Fabbretti, A.; He, C. G.; Gaspari, E.; Maffioli, S.; Brandi, L.; Spurio, R.; Sosio, M.; Jabes, D.; Donadio,

S. A Derivative of the Thiopeptide GE2270A Highly Selective against Propionibacterium Acnes. *Antimicrob.*

Agents Chemother. **2015**, *59* (8), 4560–4568. https://doi.org/10.1128/AAC.05155-14.

(46)   LaMarche, M. J.; Leeds, J. A.; Brewer, J.; Dean, K.; Ding, J.; Dzink-Fox, J.; Gamber, G.; Jain, A.;

Kerrigan, R.; Krastel, P.; Lee, K.; Lombardo, F.; McKenney, D.; Neckermann, G.; Osborne, C.; Palestrant, D.;

Patane, M. A.; Rann, E. M.; Robinson, Z.; Schmitt, E.; Stams, T.; Tiamfook, S.; Yu, D.; Whitehead, L.

Antibacterial and Solubility Optimization of Thiomuracin A. *J. Med. Chem.* **2016**, *59* (14), 6920–6928.

https://doi.org/10.1021/acs.jmedchem.6b00726.

(47)   Cox, C. L.; Tietz, J. I.; Sokolowski, K.; Melby, J. O.; Doroghazi, J. R.; Mitchell, D. A. Nucleophilic 1,4-

Additions for Natural Product Discovery. *ACS Chem. Biol.* **2014**, *9* (9), 2014–2022.

https://doi.org/10.1021/cb500324n.

(48)   Mizuhara, N.; Kuroda, M.; Ogita, A.; Tanaka, T.; Usuki, Y.; Fujita, K. I. Antifungal Thiopeptide

Cyclothiazomycin B1 Exhibits Growth Inhibition Accompanying Morphological Changes via Binding to

Fungal Cell Wall Chitin. *Bioorganic Med. Chem.* **2011**, *19* (18), 5300–5310.

https://doi.org/10.1016/j.bmc.2011.08.010.

(49)   Aoki, M.; Ohtsuka, T.; Yamada, M.; Ohba, Y.; Yoshizaki, H.; Yasuno, H.; Sano, T.; Watanabe, J.;

Yokose, K.; Seto, H. Cyclothiazomycin, a Novel Polythiazole-Containing Peptide with Renin Inhibitory

Activity. Taxonomy, Fermentation, Isolation and Physico-Chemical Characterization. *J. Antibiot. (Tokyo).*

1991, *44* (6), 582–588. https://doi.org/10.7164/antibiotics.44.582.

(50)   Hood, M. I.; Skaar, E. P. Nutritional Immunity: Transition Metals at the Pathogen-Host Interface.

Nature Reviews Microbiology. NIH Public Access August 2012, pp 525–537.

https://doi.org/10.1038/nrmicro2836.

(51)   Iatsenko, I.; Marra, A.; Boquete, J.-P.; Peña, J.; Lemaitre, B. Iron Sequestration by Transferrin 1

Mediates Nutritional Immunity in Drosophila Melanogaster. *Proc. Natl. Acad. Sci.* **2020**, *117* (13),

201914830. https://doi.org/10.1073/pnas.1914830117.

(52)   Kortman, G. A. M.; Raffatellu, M.; Swinkels, D. W.; Tjalsma, H. Nutritional Iron Turned inside out:

Intestinal Stress from a Gut Microbial Perspective. *FEMS Microbiology Reviews*. 2014, pp 1202–1234.

https://doi.org/10.1111/1574-6976.12086.

For Table of Contents Only