Mining zebrafish microbiota reveals key community-level

resistance against fish pathogen infection.

Franziska A. STRESSMANN[1,2†], Joaquin BERNAL-BAYARD[1,3†], David PEREZ-

PASCUAL[1], Bianca AUDRAIN[1], Olaya RENDUELES[1,4], Valérie BRIOLAT[5],

Sebastian BRUCHMANN[6,7], Stevenn VOLANT[8], Amine GHOZLANE[8], Susanne

HAUSSLER[6,9], Eric DUCHAUD[10] Jean-Pierre LEVRAUD[5] and Jean-Marc GHIGO[1]*

9
10 [1] *Genetics of Biofilms Laboratory, Institut Pasteur,* UMR CNRS2001, *Paris 75015, France*

11 [2] *current address: Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Department of Chemical*
12
13 *Analytics and Biogeochemistry, Müggelseedamm 310, 12587, Berlin, Germany*

14 [3] *Departamento de Genética, Facultad de Biología, Universidad de Sevilla, Apartado 1095, 41080*
15
16 *Sevilla, Spain*

17 [4] *current address: Microbial Evolutionary Genomics Laboratory, Institut Pasteur, UMR3525, Paris*
18
19 *75015, France*

20 [5] *Macrophages and Development of Immunity Laboratory, Institut Pasteur, UMR3738 CNRS, Paris*
21
22 *75015, France*

23 [6] *Department of Molecular Bacteriology, Helmholtz Centre for Infection Research, Braunschweig,*
24
25 *Germany.*

26 [7] *current address: University of Cambridge, Department of Veterinary Medicine, Madingley Road*
27
28 *Cambridge CB3 0ES, UK*

29 [8] *Hub de Bioinformatique et Biostatistique – Département Biologie Computationnelle, Institut Pasteur,*
30
31 *USR 3756 CNRS, Paris, France*

32 [9] *current address:Department of Clinical Microbiology, Rigshospitalet; Copenhagen, 2100, Denmark*
33

34 [10] *Unité VIM, INRAE, Université Paris-Saclay, 78350 Jouy-en-Josas, France.*

36 † FS and JBB equally contributed to this work.
37

38 *** Corresponding author:** Jean-Marc Ghigo (jmghigo@pasteur.fr).

40 **Running title:** Microbiota-based infection resistance in zebrafish

42 **Keywords:** Zebrafish; germ-free; probiotic; microbiota; colonization resistance,

43 bacterial infection.

ABSTRACT

The long-known resistance to pathogens provided by host-associated microbiota fostered the notion that adding protective bacteria could prevent or attenuate infection. However, the identification of endogenous or exogenous bacteria conferring such protection is often hindered by the complexity of host microbial communities. Here, we used zebrafish and the fish pathogen *Flavobacterium columnare* as a model system to study the determinants of microbiota-associated colonization resistance. We compared infection susceptibility in germ-free, conventional and re-conventionalized larvae and showed that a consortium of 10 culturable bacterial species are sufficient to protect zebrafish. Whereas survival to *F. columnare* infection does not rely on host innate immunity, we used antibiotic dysbiosis to alter zebrafish microbiota composition, leading to the identification of two different protection strategies. We first identified that the bacterium *Chryseobacterium massiliae* individually protects both larvae and adult zebrafish. We also showed that an assembly of 9 endogenous zebrafish species that do not otherwise protect individually confer a community-level resistance to infection. Our study therefore provides a rational approach to identify key endogenous protecting bacteria and promising candidates to engineer resilient microbial communities. It also shows how direct experimental analysis of colonization resistance in low-complexity *in vivo* models can reveal unsuspected ecological strategies at play in microbiota-based protection against pathogens.

INTRODUCTION

Animal resident microbial consortia form complex and long-term associations with

important community-level functions essential for host development and physiology (1,

2). Microbial ecosystems also provide protection against exogenous pathogens by

inhibition of pathogen settlement and growth and/or stimulation of the host immune

system (3-8). From the perspective of microbial community composition, a shift or

reduction in resident microbial diversity, a phenomenon generally referred to as

dysbiosis, is often associated with increased susceptibility to infection due to the loss or

change in abundance of key microbial community members (3, 9). These observations

early supported the notion that addition or promotion of individually or communally

protective bacteria (such as probiotics) could minimize microbiota dysbiosis or directly

prevent infection to restore host health (10-12).

Although the efficacy of probiotics has been shown in animal and humans, their

mechanisms of action are poorly understood and low throughput experimental models

often offer limited information on the individual contribution of probiotic species to

community functions (1, 6, 7, 13, 14). Moreover, characterization of bacterial strains

improving colonization resistance is still hindered by the complexity of host-commensal

ecosystems. Zebrafish have recently emerged as a powerful tool to study microbe-

microbe and host-microbe interactions (15-19). Zebrafish can be easily reared germ-free

or gnotobiotically in association with specific bacterial species (15, 20). Moreover,

zebrafish bacterial communities are increasingly well characterized and a number of

phylogenetically distinct zebrafish gut bacteria can be cultured, making this model system

directly amenable to microbiota manipulation and assessment of probiotic effect on host

infection resistance (21-24). Several studies have used zebrafish to evaluate the effect of

exogenous addition of potential probiotics on host resistance to infection by various

pathogens (22-29). However, the role of the endogenous microbial community in

protecting against invasive pathogen was rarely assessed and the reported protections

were often partial, illustrating the difficulty in identifying fully protective exogenous

probiotics.

Here we used germ-free and conventional zebrafish larvae to mine the indigenous

commensal microbiota for bacterial species protecting against *Flavobacterium*

columnare, a bacterial pathogen affecting wild and cultured fish species. We identified

two distinct infection resistance strategies preventing mortality caused by *F. columnare*,

mediated either by an individual member of the microbiota, the *Bacteroidetes*

Chryseobacterium massiliae or by an assembly of 9 individually non-protecting bacterial

species that formed a protective community. Our results demonstrated that mining host

microbiota constitutes a powerful approach to identify key mediators of intrinsic

colonization resistance, providing insight into how to engineer ecologically resilient and

protective microbial communities.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Bacterial strains isolated from zebrafish microbiota are listed in Table 1. *F. columnare*

strains (Suppl. Table S3) were grown at 28°C in tryptone yeast extract salts (TYES) broth

[0.4 % (w/v) tryptone, 0.04 % yeast extract, 0.05 % (w/v) $MgSO_4$ $7H_2O$, 0.02 % (w/v)

$CaCl_2$ $2H_2O$, 0.05 % (w/v) D-glucose, pH 7.2]. *F. columnare* strains were assigned into

four genomovar groups using 16S rRNA restriction fragment length polymorphism

analysis, including genomovar I, I/II, II, and III (30). All 10 strains of the core zebrafish

microbiota species were grown in TYES or LB at 28°C.

Ethics statement

All animal experiments described in the present study were conducted at the Institut

Pasteur (larvae) or at INRA Jouy-en-Josas (adults) according to European Union

guidelines for handling of laboratory animals

(http://ec.europa.eu/environment/chemicals/lab_animals/home_en.htm) and were

approved by the relevant institutional Animal Health and Care Committees.

General handling of zebrafish.

Wild-type AB fish, originally purchased from the Zebrafish International Resource

Center (Eugene, OR, USA), or *myd88*-null mutants (*myd88^{hu3568/hu3568}*) (31), kindly

provided by A.H Meijer, (Leiden University, the Netherlands), were raised in our facility.

A few hours after spawning, eggs were collected, rinsed, and sorted under a dissecting

scope to remove faeces and unfertilized eggs. All following procedures were performed

in a laminar microbiological cabinet with single-use disposable plasticware. Fish were

kept in sterile 25 cm^3 vented cap culture flasks containing 20 mL of water (0-6 dpf- 15

fish per flasks) or 24-well microtiter plates (6-15 dpf-1 fish per 2 mL well) in autoclaved

mineral water (Volvic) at 28°C. Fish were fed 3 times a week from 4 dpf with germ-free

Tetrahymena thermophila protozoans (22). Germ-free zebrafish were produced after

sterilizing the egg chorion protecting the otherwise sterile egg, with antibiotic and

chemical treatments (see below), whereas conventional larvae (with facility-innate

microbiota) were directly reared from non-sterilized eggs and then handled exactly as the

germ-free larvae.

Sterilization of zebrafish eggs.

Egg sterilization was performed as previously described with some modifications (22).

Freshly fertilized zebrafish eggs were first bleached (0.003%) for 5 min, washed 3 times

in sterile water under gentle agitation and maintained overnight in groups of 100 eggs per

75 cm^3 culture flasks with vented caps containing 100 mL of autoclaved Volvic mineral

water supplemented with methylene blue solution (0.3 μg/mL). Afterwards, eggs were

transferred into 50 mL Falcon tubes (100 eggs per tube) and treated with a mixture of

antibiotics (500 μL of penicillin G: streptomycin, 10,000 U/ml: 10 mg/mL GIBCO

#P4333), 200 μL of filtered kanamycin sulfate (100 mg/mL, SERVA Electrophoresis

#26899) and antifungal drug (50 μL of amphotericin B solution Sigma-Aldrich (250

μg/mL) #A2942) for 2 h under agitation at 28°C. Eggs were then washed 3 times in sterile

water under gentle agitation and bleached (0.003%) for 15 min, resuspending the eggs

every 3 min by inversion. Eggs were washed again 3 times in water and incubated 10 min

with 0.01% Romeiod (COFA, Coopérative Française de l'Aquaculture). Finally, eggs

were washed 3 times in water and transferred into 25 cm^3 culture flasks with vented caps

containing 20 mL of water. After sterilization, eggs were transferred with approximately

30 to 35 eggs / flasks and were transferred into new flasks at 4 dpf before re-

conventionalization with 10 to 15 fish / flask. We monitored sterility at several points

during the experiment by spotting 50 µL of water from each flask on LB, TYES and on

YPD agar plates, all incubated at 28°C under aerobic conditions. Plates were left for at

least 3 days to allow slow-growing organisms to multiply. Spot checks for bacterial

contamination were also carried out by PCR amplification of water samples with the 16S

rRNA gene primers and procedure detailed further below. If a particular flask was

contaminated, those fish were removed from the experiment.

Procedure for raising germ-free zebrafish.

After hatching, fish were fed with germ-free *T. thermophila* 3 times per week from 4 dpf

onwards. *(i) T. thermophila stock.* A germ-free line of *T. thermophila* was maintained at

28°C in 20 mL of PPYE (0.25% proteose peptone BD Bacto #211684, 0.25% yeast

extract BD Bacto #212750) supplemented with penicillin G (10 unit/mL) and

streptomycin (10µg/mL). Medium was inoculated with 100 µL of the preceding *T.*

thermophila stock. After one week of growth, samples were taken, tested for sterility on

LB, TYES and YPD plates and restocked again. *(ii) Growth. T. thermophila* were

incubated at 28°C in MYE broth (1% milk powder, 1% yeast extract) inoculated from

stock suspension at a 1:50 ratio. After 24 h of growth, *T. thermophila* were transferred to

Falcon tubes and washed (4400 rpm, 3 min at 25°C) 3 times in 50 mL of autoclaved

Volvic water. Finally, *T. thermophila* were resuspended in sterile water and added to

culture flasks (500 µL in 20 mL) or 24-well plates (50 µL / well). Sterility of *T.*

thermophila was tested by plating and 16S rRNA PCR as described in the section above.

(iii) *Fine-powder feeding*. When indicated, fish were fed with previously γ-ray-sterilized

fine-powdered food suitable for an early first feeding gape size (ZM-000 fish feed, ZM

Ltd) every 48 hours (32).

Re-conventionalization of germ-free zebrafish.

At 4 dpf, just after hatching, zebrafish larvae were re-conventionalized with a single

bacterial population or a mix of several. The 10 bacterial strains constituting the core

protective microbiota were grown for 24h in suitable media (TYES or LB) at 28°C.

Bacteria were then pelleted and washed twice in sterile water, and all adjusted to the same

cell density (OD_{600} = 1 or 5.10^7 cfu/mL) (i) *Re-conventionalization with individual*

species. Bacteria were resuspended and transferred to culture flasks containing germ-free

fish at a final concentration of 5.10^5 cfu/mL. (ii) *Re-conventionalization with bacterial*

mixtures. For the preparation of Mix10, Mix9, Mix8 and all other mixes used, equimolar

mixtures were prepared by adding each bacterial species at initial concentration to 5.10^7

cfu/mL. Each bacterial mixture suspension was added to culture flasks containing germ-

free fish at a final concentration of 5.10^5 cfu/mL.

Infection challenges.

F. columnare strains (Suppl. Table S3) were grown overnight in TYES broth at 28°C.

Then, 2 mL of culture were pelleted (10,000 rpm for 5 min) and washed once in sterile

water. GF zebrafish were brought in contact with the tested pathogens at 6 dpf for 3h by

immersion in culture flasks with bacterial doses ranging from 5.10^2 to 5.10^7 cfu/mL. Fish

were then transferred to individual wells of 24-well plates, containing 2 mL of water and

50 μL of freshly prepared GF *T. thermophila* per well. Mortality was monitored daily as

described in (22) and as few as 54±9 cfu/larva of *F. columnare* were recovered from

infected larvae. All zebrafish experiments were stopped at day 9 post-infection and

zebrafish were euthanized with tricaine (MS-222) (Sigma-Aldrich #E10521). Each

experiment was repeated at least 3 times and between 10 and 15 larvae were used per

condition and per experiment.

Collection of eggs from other zebrafish facilities.

Conventional zebrafish eggs were collected in 50 mL Falcon tubes from the following

facilities: Facility 1 - zebrafish facility in Hospital Robert Debré, Paris; Facility 2 - Jussieu

zebrafish facility A2, University Paris 6; Facility 3 - Jussieu – zebrafish facility C8

(UMR7622), University Paris 6; Facility 4: AMAGEN commercial facility, Gif sur

Yvette; Larvae were treated with the same rearing conditions, sterilization and infection

procedures used in the Institut Pasteur facility.

Determination of fish bacterial load using cfu count. Zebrafish were euthanized with

tricaine (MS-222) (Sigma-Aldrich #E10521) at 0.3 mg/mL for 10 minutes. Then they

were washed in 3 different baths of sterile PBS-0.1% Tween to remove bacteria loosely

attached to the skin. Finally, they were transferred to tubes containing calibrated glass

beads (acid-washed, 425 μm to 600 μm, SIGMA-ALDRICH #G8772) and 500 μL of

autoclaved PBS. They were homogenized using FastPrep Cell Disrupter (BIO101/FP120

QBioGene) for 45 s at maximum speed (6.5 m/s). Finally, serial dilutions of recovered

suspension were spotted on TYES agar and cfu were counted after 48h of incubation at

28°C.

Characterization of zebrafish microbial content.

Over 3 months, the experiment was run independently 3 times and 3 different batches of

eggs were collected from different fish couples in different tanks. Larvae were reared as

described above. GF and Conv larvae were collected at 6 dpf and 11 dpf for each batch.

Infected Conv larvae were exposed to *F. columnare*[ALG] for 3h by immersion as described

above. For each experimental group, triplicate pools of 10 larvae (one in each

experimental batch) were euthanized, washed and lysed as above. Lysates were split into

3 aliquots, one for culture followed by 16S rRNA gene sequencing (A), one for 16S rRNA

gene clone library generation and Sanger sequencing (B), and one for Illumina

metabarcoding-based sequencing (C).

A) Bacterial culture followed by 16S rRNA gene-based identification

Lysates were serially diluted and immediately plated on R2A, TYES, LB, MacConkey,

BHI, BCYE, TCBS and TSB agars and incubated at 28°C for 24-72h. For each agar,

colony morphotypes were documented, and colonies were picked and re-streaked on the

same agar in duplicate. In order to identify the individual morphotypes, individual

colonies were picked for each identified morphotype from each agar, vortexed in 200 μL

DNA-free water and boiled for 20 min at 90°C. Five μL of this bacterial suspension were

used as template for colony PCR to amplify the 16S rRNA gene with the universal primer

pair for the Domain bacteria 8f (5'-AGA GTT TGA TCC TGG CTC AG-3') and 1492r

(5'-GGT TAC CTT GTT ACG ACT T-3'). Each primer was used at a final concentration

of 0.2 μM in 50 μL reactions. PCR cycling conditions were - initial denaturation at 94°C

for 2 min, followed by 32 cycles of denaturation at 94 °C for 1 min, annealing at 56°C for

1 min, and extension at 72°C for 2 min, with a final extension step at 72°C for 10 min.

16S rRNA gene PCR products were verified on 1% agarose gels, purified with the

QIAquick® PCR purification kit and two PCR products for each morphotype were sent

for sequencing (Eurofins, Ebersberg, Germany). 16S rRNA sequences were manually

proofread, and sequences of low quality were removed from the analysis. Primer

sequences were trimmed, and sequences were compared to GenBank (NCBI) with

BLAST, and to the Ribosomal Database Project with SeqMatch. For genus determination

a 95% similarity cut-off was used, for Operational Taxonomic Unit determination, a 98%

cut-off was used.

B) 16S rRNA gene clone library generation

Total DNA was extracted from the lysates with the Mobio PowerLyzer® Ultraclean® kit

according to manufacturer's instructions. Germ-free larvae and DNA-free water were

also extracted as control samples. Extracted genomic DNA was verified by Tris-acetate-

EDTA-agarose gel electrophoresis (1%) stained with GelRed and quantified by applying

2.5 μL directly to a NanoDrop® ND-1000 Spectrophotometer. The 16S rRNA gene was

amplified by PCR with the primers 8f and 1492r, and products checked and purified as

described in section A. Here, we added 25-50 ng of DNA as template to 50 μL reactions.

Clone libraries were generated with the pGEM®-T Easy Vector system (Promega)

according to manufacturer's instructions. Presence of the cloned insert was confirmed by

colony PCR with vector primers gemsp6 (5'-GCT GCG ACT TCA CTA GTG AT-3')

and gemt7 (5'-GTG GCA GCG GGA ATT CGA T-3'). Clones with an insert of the

correct size were purified as above and sent for sequencing (Eurofins, Ebersberg,

Germany). Blanks using DNA-free water as template were run for all procedures as

controls. For the three independent runs of the experiment, 10 Conv fish per condition (6

and 11 dpf, exposed or not to *F. columnare*) and per repeat were pooled. Each pool of 10

fish was sequenced separately, generating 3 replicates for each condition (n=12), resulting

in a total of 857 clones. Clone library coverage was calculated with the following formula

$[1-(n_1/N_2)] \times 100$, where n_1 is the number of singletons detected in the clone library, and

N_2 is the total number of clones generated for this sample. Clone libraries were generated

to a minimum coverage of 95%. Sequence analysis and identification was carried out as

in section A.

C) by 16S rRNA V3V4 Amplicon Illumina sequencing

To identify the 16S rRNA gene diversity in our facility and fish collected from 4 other

zebrafish facilities, fish were reared as described above. GF fish were sterilised as above,

and uninfected germ-free and conventional fish were collected at 6 dpf and 11 dpf.

Infection was carried out as above with *F. columnare*[ALG] for 3h by bath immersion,

followed by transfer to clean water. Infected conventional fish were collected at 6 dpf 6h

after infection with *F. columnare* and at 11 dpf, the same as uninfected fish. GF infected

larvae were only collected at 6 dpf 6h post infection, as at 11 dpf all larvae had succumbed

to infection. Triplicate pools of 10 larvae were euthanized, washed and lysed as above.

Total DNA was extracted with the Mobio PowerLyzer® Ultraclean® kit as described

above and quantified with a NanoDrop® ND-1000 Spectrophotometer and sent to IMGM

Laboratories GmbH (Germany) for Illumina sequencing. Primers Bakt_341F (5'-

CCTACGGGNGGCWGCAG-3') and Bakt_805R (5'-

GACTACHVGGGTATCTAATCC-3'), amplifying variable regions 3 and 4 of the 16S

gene were used for amplification (33). Each amplicon was purified with solid phase

reversible immobilization (SPRI) paramagnetic bead-based technology (AMPure XP

beads, Beckman Coulter) with a Bead:DNA ratio of 0.7:1 (v/v) following manufacturer´s

instructions. Amplicons were normalized with the Sequal-Prep Kit (Life Technologies),

so each sample contained approximately 1 ng/µl DNA. Samples, positive and negative

controls were generated in one library. The High Sensitivity DNA LabChip Kit (was used

on the 2100 Bioanalyzer system (both Agilent Technologies) to check the quality of the

purified amplicon library. For cluster generation and sequencing, MiSeq® reagents kit

500 cycles Nano v2 (Illumina Inc.) was used. Before sequencing, cluster generation by

two-dimensional bridge amplification was performed, followed by bidirectional

sequencing, producing 2 x 250 bp paired-end (PE) reads.

MiSeq® Reporter 2.5.1.3 software was used for primary data analysis (signal processing,

de-multiplexing, trimming of adapter sequences). CLC Genomics Workbench 8.5.1

(Qiagen) was used for read-merging, quality trimming and QC reports and OTU

definition were carried out in the CLC plugin Microbial Genomics module.

Comparison of whole larvae vs intestinal bacterial content.

Larvae re-conventionalized with Mix10 and infected with *F. columnare*[ALG] at 6 dpf for

3h were euthanized and washed. DNA was extracted from pools of 10 whole larvae or of

pools of 10 intestinal tubes dissected with sterile surgical tweezer and subjected to

Illumina 16S rRNA gene sequencing. GF larvae and dissected GF intestines were

sampled as controls. As dissection of the larval guts involved high animal loss and was a

potential important contamination source, we proceeded with using entire larvae for the

rest of the study.

Whole genome sequencing.

Chromosomal DNA of the ten species composing the core of zebrafish larvae microbiota

was extracted using the DNeasy Blood & Tissue kit (QIAGEN) including RNase

treatment. DNA quality and quantity were assessed on a NanoDrop ND-1000

spectrophotometer (Thermo Scientific).

DNA sequencing libraries were made using the Nextera DNA Library Preparation Kit

(Illumina Inc.) and library quality was checked using the High Sensitivity DNA LabChip

Kit on the Bioanalyzer 2100 (Agilent Technologies). Sequencing clusters were generated

using the MiSeq reagents kit v2 500 cycles (Illumina Inc.) according to manufacturer's

instructions. DNA was sequenced at the Helmholtz Centre for Infection Research by

bidirectional sequencing, producing 2 x 250 bp paired-end (PE) reads. Between 1,108,578

and 2,914,480 reads per sample were retrieved with a median of 1,528,402. Reads were

quality filtered, trimmed and adapters removed with trimmomatic 0.39 (34) and genomes

assembled using SPAdes 3.14 (35).

Bacterial species identification.

Whole genome-based bacterial species identification was performed by the TrueBac ID

system (v1.92, DB:20190603) [https://www.truebacid.com/; (36). Species-level

identification was performed based on the algorithmic cut-off set at 95% ANI when

possible or when the 16S rRNA gene sequence similarity was >99 %.

Monitoring of bacterial dynamics. T

hree independent experiments were run over 6 weeks with eggs collected from different

fish couples from different tanks to monitor establishment and recovery. Larvae were

reared, sterilized and infected as above with the only difference that 75 cm^3 culture flasks

with vented caps (filled with 50 mL of sterile Volvic) were used to accommodate the

larger number of larvae required, as in each experiment. Larvae for time course Illumina

sequencing were removed sequentially from the experiment that monitored the survival

of the larvae. Animals were pooled (10 larvae for each time point/condition), euthanized,

washed and lysed as described above and stored at -20ºC until the end of the survival

monitoring, and until all triplicates had been collected.

A) Community establishment

In order to follow the establishment of the 10 core strains in the larvae, GF larvae were

re-conventionalized with an equiratio Mix10 as above. Re-con^{Mix10} larvae were sampled

at 4 dpf immediately after addition of the 10 core species and then 20 min, 2h, 4h and 8h

after. Germ-free, conventional larvae and the inoculum were also sampled as controls.

B) Induction of dysbiosis

Different doses of kanamycin (dose 1= 200 µg/mL; dose 2= 50 µg/mL; dose 3= 25

µg/mL) and a penicillin/streptomycin antibiotic mix (dose 1= 250 µg/mL; dose 2= 15.6

µg/mL were tested on re-con^{Mix10} 4 dpf zebrafish larvae by adding them to the flask water

to identify antibiotic treatments that were non-toxic to larvae but that caused dysbiosis.

After 16 hours of treatment, antibiotics were extensively washed off with sterile water

and larvae were challenged with *F. columnare*^{ALG}, leading to the death of all larvae – e.g.

successful abolition of colonization resistance with best results in all repeats with 250

µg/mL penicillin/streptomycin and 50 µg/mL kanamycin as antibiotic treatment.

C) Community recovery

As in B) after 8h of incubation, 4 dpf recon^{Mix10} larvae were treated with 250 µg/mL

penicillin/streptomycin and 50 µg/mL kanamycin for 16h. Antibiotics were extensively

washed off and larvae were now left to recover in sterile water for 24h to assess resilience

of the bacterial community. Samples (pools of 10 larvae) were taken at 3h, 6h, 12h, 18h

and 24h during recovery and sent for 16S rRNA Illumina sequencing. Larvae were then

challenged at 6 dpf with *F. columnare*^{ALG} for 3h and survival was monitored daily for 9

days post-infection. All time course samples were sequenced by IMGM Laboratories

GmbH, as described above.

Statistical analysis of metataxonomic data. 16S RNA analysis was performed with

SHAMAN (37)]. Library adapters, primer sequences, and base pairs occurring at 5' and

3'ends with a Phred quality score <20 were trimmed off by using Alientrimmer (v0.4.0).

Reads with a positive match against zebrafish genome (mm10) were removed. Filtered

high-quality reads were merged into amplicons with Flash (v1.2.11). Resulting amplicons

were clustered into operational taxonomic units (OTU) with VSEARCH (v2.3.4) (38).

The process includes several steps for de-replication, singletons removal, and chimera

detection. The clustering was performed at 97% sequence identity threshold, producing

459 OTUs. The OTU taxonomic annotation was performed against the SILVA SSU

(v132) database (39) completed with 16S sequence of 10 bacterial communities using

VSEARCH and filtered according to their identity with the reference (40). Annotations

were kept when the identity between the OTU sequence and reference sequence is $\geq$

78.5% for taxonomic Classes, $\geq$ 82% for Orders, $\geq$ 86.5% for Families, $\geq$ 94.5% for

Genera and $\geq$ 98% for species. Here, 73.2% of the OTUs set was annotated and 91.69 %

of them were annotated at genus level.

The input amplicons were then aligned against the OTU set to get an OTU contingency

table containing the number of amplicon associated with each OTU using VSEARCH

global alignment. The matrix of OTU count data was normalized for library size at the

OTU level using a weighted non-null count normalization. Normalized counts were then

summed within genera. The generalized linear model (GLM) implemented in the DESeq2

R package (41) was then applied to detect differences in abundance of genera between

each group. We defined a GLM that included the treatment (condition) and the time

(variable) as main effects and an interaction between the treatment and the time. Resulting

P values were adjusted according to the Benjamini and Hochberg procedure.

The statistical analysis can be reproduced on shaman by loading the count table, the

taxonomic results with the target and contrast files that are available on figshare

https://doi.org/10.6084/m9.figshare.11417082.v2.

Determination of cytokine levels. Total RNAs from individual zebrafish larvae were

extracted using RNeasy kit (Qiagen), 18h post pathogen exposure (12h post-wash).

Oligo(dT17)-primed reverse transcriptions were carried out using M-MLV H- reverse-

transcriptase (Promega). Quantitative PCRs were performed using Takyon SYBR Green

PCR Mastermix (Eurogentec) on a StepOne thermocycler (Applied Biosystems). Primers

for *ef1a* (housekeeping gene, used for cDNA amount normalization), *il1b*, *il10* and *il22*

are described in (22). Data were analysed using the ΔΔCt method. Four larvae were

analysed per condition. Zebrafish genes and proteins mentioned in the text: *ef1a*

NM_131263*; il1b* BC098597*; il22* NM_001020792*; il10* NM_001020785*; myd88*

NM_212814.

Histological comparisons of GF, Conv and Re-Conv fish GF infected or not with *F.

*columnare***

Fish were collected 24h after infection (7 dpf) and were fixed for 24h at 4°C in Trump

fixative (4% methanol-free formaldehyde, 1% glutaraldehyde in 0.1 M PBS, pH 7.2) and

sent to the PIBiSA Microscopy facility services (https://microscopies.med.univ-tours.fr/)

in the Faculté de Médecine de Tours (France), where whole fixed animals were processed,

embedded in Epon. Semi-thin sections (1μm) and cut using a X ultra-microtome and then

either dyed with toluidine blue for observation by light microscopy and imaging or

processed for Transmission electron microscopy.

Adult zebrafish pre-treatment with *C. massiliae*

The zebrafish line AB was used. Fish were reared at 28°C in dechlorinated recirculated

water, then transferred into continuous flow aquaria when aging 3-4 months for infection

experiments. *C. massiliae* was grown in TYES broth at 150 rpm and 28°C until stationary

phase. This bacterial culture was washed twice in sterile water and adjusted to $OD_{600nm} =$

1. Adult fish re-conventionalization was performed by adding *C. massiliae* bacterial

suspension directly into the fish water (1L) at a final concentration of 2.10^6 cfu/mL.

Bacteria were maintained in contact with fish for 24 h by stopping the water flow then

subsequently removed by restoring the water flow. *C. massiliae* administration was

performed twice after water renewal. In the control group, the same volume of sterile

water was added.

Adult zebrafish infection challenge

F. columnare infection was performed just after fish re-conventionalization with *C.*

massiliae. The infection was performed as previously described by Li and co-workers

with few modifications [Li et al., 2017]. Briefly, *F. columnare* strain ALG-0530 was

grown in TYES broth at 150 rpm and 28 °C until late-exponential phase. Then, bacterial

cultures were diluted directly into the water of aquaria (200 mL) at a final concentration

of 5.10^6 cfu/mL. Bacteria were maintained in contact with fish for 1 h by stopping the

water flow then subsequently removed by restoring the water flow. Sterile TYES broth

was used for the control group. Bacterial counts were determined at the beginning of the

immersion challenge by plating serial dilutions of water samples on TYES agar. Water

was maintained at 28°C and under continuous oxygenation for the duration of the

immersion. Groups were composed of 10 fish. Virulence was evaluated according to fish

mortality 10 days post-infection.

Statistical methods.

Statistical analyses were performed using unpaired, non-parametric Mann-Whitney test

or unpaired t-tests. Analyses were performed using Prism v8.2 (GraphPad Software).

Evenness: The Shannon diversity index was calculated with the formula (H_S = -

$\Sigma[P(\ln(P))]$) where P is the relative species abundance. Total evenness was calculated for

the Shannon index as $E = H_S/H_{max.}$ The less evenness in communities between the species

(and the presence of a dominant species), the lower this index is.

Data availability

Bacterial genome sequences obtained in the present study are available at the European

Nucleotide Archive with the project number PRJEB36872, under accession numbers

numbers ERS4385993 (*Aeromonas veronii 1*); ERS4386000 (*Aeromonas veronii 2*);

ERS4385996 (*Aeromonas caviae*); ERS4385998 (*Chryseobacterium massiliae*);

ERS4385999 (*Phyllobacterium myrsinacearum*); ERS4406247 (*Pseudomonas*

sediminis); ERS4385994 (*Pseudomonas mossellii*) ERS4386001 (*Pseudomonas*

nitroreducens); ERS4385997 (*Pseudomonas peli*); ERS4385995 (*Stenotrophomas*

maltophilia);

471
472
473

RESULTS

***Flavobacterium columnare* kills germ-free but not conventional zebrafish**

To investigate microbiota-based resistance to infection in zebrafish, we compared the

sensitivity of germ-free (GF) and conventional (Conv) zebrafish larvae to *F. columnare*,

an important fish pathogen affecting carp, channel catfish, goldfish, eel, salmonids and

tilapia and previously shown to infect and kill adult zebrafish (12, 42-45). We used bath

immersion to expose GF and Conv zebrafish larvae at 6 days post fertilization (dpf), to a

collection of 28 *F. columnare* strains, belonging to four different genomovars for 3h at

5.10^5 colony forming units (cfu)/mL. Daily monitoring showed that 16 out of 28 *F.*

columnare strains killed GF larvae in less than 48h (Supplementary Fig. S1A), whereas

Conv larvae survived exposure to all tested virulent *F. columnare* strains (Supplementary

Fig. S1B). Exposure to the highly virulent strain ALG-00-530 (hereafter *F.*

columnare[ALG]) also showed that GF mortality was fast (1 day) and dose-dependent and

that Conv zebrafish survived all but the highest dose (10^7 cfu/mL) (Fig. 1). Similar

survival of infected Conv larvae was obtained with zebrafish AB strain eggs obtained

from 4 different zebrafish facilities (Supplementary Fig. S2), suggesting that

conventional zebrafish microbiota could provide protection against *F. columnare*

infection.

A community of 10 culturable bacterial strains protect against *F. columnare*

infection

In our rearing conditions, the conventional larval microbiota is acquired after hatching

from microorganisms present on the egg chorion and in fish facility water. To test the

hypothesis that microorganisms associated with conventional eggs provided protection

against *F. columnare*[ALG], we exposed sterilized eggs to either fish facility tank water or

to non-sterilized conventional eggs at 0 or 4 dpf (before or after hatching, respectively).

In both cases, these re-conventionalized (re-Conv) zebrafish survived *F. columnare*[ALG]

infection as well as Conv zebrafish (Supplementary Fig. S3). To determine the

composition of conventional zebrafish microbiota, we generated 16S rRNA gene clone

libraries from homogenate pools of Conv larvae aged 6 and 11 dpf exposed or not to *F.*

columnare[ALG], sampled over 3 months from 3 different batches of larvae (n=10). A total

of 857 clones were generated for all samples. We identified 15 operational taxonomical

units (OTUs), 10 of which were identified in all experiments (Table 1, Suppl. Table S1).

Two OTUs (belonging to an *Ensifer* sp. and a *Hydrogenophaga* sp.) were only detected

once, and a *Delftia* sp., a *Limnobacter* sp. and a *Novosphingobium* sp. were detected more

than once (2, 3 and 2 times, respectively), but not consistently in all batches of fish (Table

1, Suppl. Table S1). Moreover, deep-sequencing of the 16S rRNA V3-V4 region of

gDNA retrieved from larvae originating from the other four zebrafish facilities described

above, revealed that OTUs for all of these 10 strains were also detected in Conv larvae,

with the exception of *A. veronii* strain 2 that was not detected in all samples (Suppl. Table

S2).

To isolate culturable zebrafish microbiota bacteria, we plated dilutions of homogenized

6 dpf and 11 dpf larvae pools on various growth media and we identified 10 different

bacterial morphotypes. 16S-based analysis followed by full genome sequencing

identified 10 bacteria corresponding to 10 strains of 9 different species that were also

consistently detected by culture-free approaches (Table 1). To assess the potential

protective role of these 10 strains, we re-conventionalized GF zebrafish at 4 dpf with a

mix of all 10 identified culturable bacterial species (hereafter called Mix10), each at a

concentration of 5.10^5 cfu/mL and we monitored zebrafish survival after exposure to *F.*

columnare[ALG] at 6 dpf. We showed that zebrafish re-conventionalized with the Mix10

(Re-Conv[Mix10]) displayed a strong level of protection against all identified highly virulent

F. columnare strains (Supplementary Fig. S4). These results demonstrated that the Mix10

constitutes a core protective bacterial community providing full protection of zebrafish

larvae against *F. columnare* infection.

Community dynamics under antibiotic-induced dysbiosis reveal a key contributor

to resistance to *F. columnare* infection

To further analyze the determinants of Mix10 protection against *F. columnare*[ALG]

infection, we inoculated 4 dpf larvae with an equal-ratio mix of the 10 bacteria (at 5.10^5

cfu/mL each) and monitored their establishment over 8 hours. We first verified that whole

larvae bacterial content (OTU abundance) was not significantly different from content of

dissected intestinal tubes (p=0.99, two-tailed t-test) (Supplementary Fig. S5) and

proceeded to use entire larvae to monitor bacterial establishment and recovery in the rest

of the study. We then collected pools of 10 larvae immediately after re-

conventionalization (t_0), and then at 20 min, 2 hours, 4 hours and 8 hours in three

independent experiments. Illumina sequencing of the 16S rRNA gene was used to follow

bacterial relative abundance. At t_0, all species were present at > 4% in the zebrafish, apart

from *A. veronii* strains 1 (0.2%) and 2 (not detected) (Supplementary Fig. S6). *Aeromonas*

caviae was detected as the most abundant species (33%), followed by *Stenotrophomonas*

maltophilia (23%) and *Chryseobacterium massiliae* (12%), altogether composing 68% of

the community (Supplementary Fig. S6). The relative species abundance, possibly

reflecting initial colonization ability, was relatively stable for most species during

community establishment, with similar species evenness at t_0 (E = 0.84) and t_{8h} (E = 0.85).

Whereas both Conv and Re-Conv^{Mix10} larvae were protected against *F. columnare*^{ALG}

infection, the global structure of the reconstituted Mix10 population was different from

the conventional one at 4 dpf (Supplementary Fig. S6).

To test the sensitivity to disturbance and the resilience of the protection provided by

Mix10 bacterial community, we subjected Re-Conv^{Mix10} zebrafish to non-toxic antibiotic

treatment at 4dpf using either 250 µg/mL penicillin/streptomycin combination (all

members of the Mix10 bacteria are sensitive to penicillin/streptomycin) or 50 µg/mL

kanamycin (affecting all members of the Mix10 bacteria except *C. massiliae*, *P.*

myrsinacearum and *S. maltophilia*) (Suppl. Fig. S7). At 5 dpf, after 16 hours of exposure,

antibiotics were washed off and zebrafish were immediately exposed to *F. columnare*^{ALG}.

Both antibiotic treatments resulted in complete loss of the protection against *F.*

columnare^{ALG} infection observed in Re-Conv^{Mix10} (Fig. 2A). We then used the same

antibiotic treatments but followed by a 24h recovery period after washing off the

antibiotics at 5 dpf, therefore only performing the infection at 6 dpf (Fig. 2B). Whilst Re-

Conv^{Mix10} larvae treated with penicillin/streptomycin showed similar survival to infected

GF larvae, kanamycin-treated Re-Conv^{Mix10} zebrafish displayed restored protection after

24h recovery and survived similarly to untreated conventionalized fish (Fig. 2B).

Sampling and 16S gene analysis during recovery experiments at different time points

showed that bacterial community evenness decreased after antibiotic administration for

both treatments (E = 0.85 for 4 dpf control, E = 0.72 for t_0 kanamycin and E = 0.7 for t_0

penicillin/streptomycin), and continued to decrease during recovery (E = 0.6 and 0.64 for

kanamycin and penicillin/streptomycin treatment after 24h recovery, respectively).

Although *C. massiliae* remained detectable immediately after both antibiotic treatments,

penicillin/streptomycin treatment led a significant reduction in its relative abundance

(0.21%) (Fig. 2C). By contrast, *C. massiliae* relative abundance rebounded 6h after

cessation of kanamycin treatment and was the dominant member (52%) of the

reconstituted microbiota after 24h recovery period (Fig. 2D), suggesting that the

protective effect observed in the kanamycin-treated larvae might be due to the recovery

of *C. massiliae*.

Resistance to *F. columnare* infection is provided by both individual and

community-level protection

To test the potential key role played by *C. massiliae* in protection against *F. columnare*[ALG]

infection, we exposed 4 dpf GF zebrafish to *C. massiliae* only and showed that it

conferred individual protection at doses as low as 5.10^2 cfu/mL (Fig. 3). Whereas none

of the 9 other species composing the Mix10 were protective individually (Fig. 3A), their

equiratio combination (designated as Mix9) conferred protection to zebrafish, although

not at doses lower than 5.10^4 cfu/mL and not as reproducibly as with *C. massiliae* (Fig.

3B). To identify which association of species protected Re-Conv[Mix9] zebrafish against *F.*

columnare[ALG] infection, we tested all 9 combinations of 8 species (Mix8), as well as

several combinations of 7, 6, 4 or 3 species and showed no protection (Suppl. Fig. S8A

and Suppl. Table S4). We then tested whether lack of protection of Mix8 compared to

Mix9 could rely on a density effect by doubling the concentration of any of the species

within the non-protective Mix8a (Suppl. Fig. S8B) and showed no protection. These

results therefore indicated that microbiota-based protection against *F. columnare*[ALG]

infection relied on either *C. massiliae*-dependent membership effect or on a community-

dependent effect mediated by the Mix9 consortium.

Pro- and anti- inflammatory cytokine production does not contribute to microbiota-

mediated protection against *F. columnare*[ALG] infection.

To test the contribution of the immune response of zebrafish larvae to resistance to *F.*

columnare infection, we used qRT-PCR to measure cytokine mRNA expression in GF

and Conv zebrafish exposed or not to *F. columnare*[ALG]. We also tested the impact of re-

conventionalization with *C. massiliae* (re-Conv[Cm]), Mix10 (re-Conv[Mix10]) or with Mix4

(*A. caviae*, both *A. veronii* spp., *P. mossellii*) as a non-protective control (Suppl. Table

S4). We tested genes encoding IL1β (pro-inflammatory), IL22 (promoting gut repair),

and IL10 (anti-inflammatory) cytokines. While we observed some variation in *il10*

expression among non-infected re-conventionalized larvae, this did not correlate with

protection. Furthermore, *il10* expression was not modulated by infection in any of the

tested conditions (Fig. 4A). By contrast, we observed a strong induction of *il1b* and *il22*

in GF zebrafish exposed to *F. columnare*[ALG] (Fig. 4BC). However, although this

induction was reduced in protected Conv, Re-Conv[Cm] and Re-Conv[Mix10], it was also

observed in non-protective Re-Conv[Mix4] larvae, indicating that down-modulation of the

inflammatory response induced by *F. columnare* does not correlate with resistance to

infection. Consistently, the use of a *myd88* mutant, a key adaptor of IL-1 and toll-like

receptor signalling deficient in innate immunity[31, 46], showed that Conv or Re-

Conv^{Mix10}, but not GF *myd88* mutants survived *F. columnare* as well as wild-type

zebrafish (Figure 4D). Moreover, *il1b* induction by *F. columnare* infection was observed

only in GF larvae and was *myd88*-independent (Suppl. Figure S9). These results therefore

indicated that the tested cytokine responses do not play a significant role in the

microbiota-mediated protection against *F. columnare* infection.

C. massiliae and Mix9 protect zebrafish from intestinal damages upon *F.*

***columnare*^{ALG} infection**

Histological analysis of GF larvae fixed 24h after exposure to *F. columnare*^{ALG} revealed

extensive intestinal damage (Fig. 5A) prior to noticeable signs in other potential target

organs such as gills or skin. To test the requirement for gut access in *F. columnare*^{ALG}

infection process, we modified our standard rearing protocol of GF fish, which involves

feeding with live germ-free *T. thermophila*. We found that, if left unfed, GF zebrafish did

not die after *F. columnare*^{ALG} exposure, while feeding with either *T. thermophila* or

another food source such as sterile fish food powder, restored sensitivity to *F.*

columnare^{ALG} infection (Suppl. Fig S10), suggesting that successful infection requires

feeding and ingestion.

Histological sections consistently showed severe disorganization of the intestine with

blebbing in the microvilli and vacuole formation in *F. columnare*^{ALG}-infected GF larvae

(Fig. 5). In contrast, zebrafish pre-incubated with either *C. massiliae* or Mix9 consortium

at 4 dpf, and then exposed to *F. columnare*^{ALG} at 6 dpf showed no difference compared

to non-infected larvae or conventional infected larvae (Fig. 5), confirming full protection

against *F. columnare*^{ALG} at the intestinal level.

C. massiliae* protects larvae and adult zebrafish against *F. columnare

The clear protection provided by *C. massiliae* against *F. columnare*[ALG] infection

prompted us to test whether exogenous addition of this bacterium could improve

microbiota-based protection towards this widespread fish pathogen. We first showed that

zebrafish larvae colonized with *C. massiliae* were fully protected against all virulent *F.*

columnare strains identified in this study (Fig. 6A). To test whether *C. massiliae* could

also protect adult zebrafish from *F. columnare* infection, we pre-treated conventional 3-

4-month-old Conv adult zebrafish with *C. massiliae* for 48h before challenging them with

a high dose (5.10^6 cfu/mL) *F. columnare*[ALG]. Monitoring of mortality rate showed that

pre-treatment with *C. massiliae* significantly increased the survival rate of adult zebrafish

upon *F. columnare*[ALG] infection compared to non-treated conventional fish (p=0.0084

Mann-Whitney test, Figure 6B). Taken together, these results show that *C. massiliae* is a

putative broad-spectrum probiotic protecting zebrafish against columnaris disease caused

by *F. columnare*.

DISCUSSION

In this study, we used re-conventionalization of otherwise germ-free zebrafish larvae to show that conventional-level protection against infection by a broad range of highly virulent *F. columnare* strains is provided by a set of 10 culturable bacterial strains, belonging to 9 different species, isolated from the indigenous standard laboratory zebrafish microbiota. With the exception of the Bacteroidetes *C. massiliae*, this protective consortium was dominated by Proteobacteria such as *Pseudomonas* and *Aeromonas* spp., bacteria commonly found in aquatic environments (47, 48). Despite the relative permissiveness of zebrafish larvae microbiota to environmental variations and inherent variability between samples (49), we showed that these ten bacteria were consistently identified in four different zebrafish facilities, suggesting the existence of a core microbiota with important colonization resistance functionality. Use of controlled combinations of these 10 bacterial species enabled us to show a very robust species-specific protection effect in larvae mono-associated with *C. massiliae.* We also identified a community-level protection provided by the combination of the 9 other species that were otherwise unable to protect against *F. columnare* when provided individually. This protection was however less reproducible and required a minimum inoculum of 5.10^4 cfu/mL, compared to 5.10^2 cfu/mL with *C. massiliae*. These results therefore suggest the existence of two distinct microbiota-based protection scenarios: one based on a membership effect provided by *C. massiliae*, and the other mediated by the higher-order activity of the Mix9 bacterial community.

Although protection against *F. columnare* infection does not seem to rely on microbiota-

based immuno-modulation, we cannot exclude that, individually, some members of the

studied core zebrafish microbiota could induce pro- or anti-inflammatory responses

masked in presence of the full Mix10 consortium (1). We also acknowledge that the

mechanistic aspects of the two identified colonization resistance scenarios are still unclear

and further studies are required to determine whether they involve nutrient depletion,

adhesion inhibition, direct or indirect growth inhibition leading to niche exclusion or any

other process (6, 13). In the case of the protection provided by *C. massiliae* against *F.*

columnare, these two phylogenetically close bacteria could compete for similar resources

and directly antagonize each other, as shown to occur between *Bacteroidetes* species (50-

53). Interestingly, infected larvae re-conventionalized with either C. *massiliae* or Mix9

showed no signs of the intestinal damage displayed by germ-free larvae, suggesting that

both *C. massiliae* and Mix9 provide similar intestinal resistance to *F. columnare*

infection. Whereas microbial colonization contributes to gut maturation and stimulates

the production of epithelial passive defences such as mucus (54, 55), lack of intestinal

maturation is unlikely to be contributing to *F. columnare*-induced mortality, as mono-

colonized larvae or larvae re-conventionalized with non-protective mixes died as rapidly

as germ-free larvae.

Several studies have monitored the long-term assembly and development of the zebrafish

microbiota from larvae to sexually mature adults, however little is known about the initial

colonization establishment of the larvae after hatching (56, 57). Neutral (stochastic) and

deterministic (host niche-based) processes (58-60) lead to microbial communities that are

mostly represented by a limited number of highly abundant species with highly diverse

low-abundant populations. In our experiments, the Mix10 species inoculum corresponded

to an equiratio bacterial mix, thus starting from an engineered and assumed total evenness

(E = 1) (61, 62). Evenness was still relatively high (0.84) and remained similar up until

8h in our study, indicating that most of the ten species were able to colonize the larvae.

From the perspective of community composition, a loss of diversity is often associated

with decreased colonization resistance, but it remains unclear whether this increased

susceptibility is due to the loss of certain key member species of the microbial community

and/or a change in their prevalence (3, 9).

We further investigated resistance to infection by exposing established bacterial

communities to different antibiotic perturbations, followed by direct challenge with *F.*

columnare (to study core microbiota sensitivity to disturbance) or after recovery (to study

its resilience) (12, 63). Antibiotics are known to shift the composition and relative

abundances of the microbiota according to their spectrum (13, 64). We observed that

penicillin/streptomycin treatment that would affect most of the core species, reduced the

abundance of all but two species (*A. veronii* 1 and *P. myrsinacearum)* that became

relatively dominant during recovery, but failed to provide protection against *F.*

columnare. With the kanamycin treatment, colonization resistance was fully restored at

the end of the 24h recovery period, indicative of a resilience that could result from species

recovering quickly to their pre-perturbation levels due to fast growth rates, physiological

flexibility or mutations (65). Interestingly, even taking into account potential biases

associated with the use of the 16S rRNA as a proxy index to determine relative abundance

(66, 67), evenness was similarly reduced during recovery for both treatments, but

abundance at phylum level changed to 48% for Proteobacteria, and 52% for Bacteroidetes

compared to the >98% of Proteobacteria with the penicillin/streptomycin treatment.

Furthermore, *C. massiliae* was detected as rare (<1%) in conventional larvae, suggesting

that it could have a disproportionate effect on the community or that community-level

protection provided by the nine other bacteria was also responsible for the protection of

conventional larvae to *F. columnare* infection.

We showed that germ-free zebrafish larvae are highly susceptible to a variety of different

F. columnare genomovars isolated from different hosts, demonstrating that they are a

robust animal model for the study of its pathogenicity. Recently, *F. columnare* mutants

in Type 9 secretion system (T9SS) were shown to be avirulent in adult zebrafish,

suggesting that proteins secreted by the T9SS are likely to be key, but still largely

unidentified, *F. columnare* virulence determinants (45). Body skin, gills, fins and tail are

also frequently damaged in salmonid fish, whereas severe infection cases are associated

with septicemia (68). We could not identify such clear *F. columnare* infection sites in

zebrafish larvae, perhaps due to the very low dose of infection used, with less than 100

cfu recovered from infected moribund larvae. However, several lines of evidence suggest

that the gut is the main target of *F. columnare* infection in our model: (i) unfed germ-free

larvae survived exposure, (ii) histology analysis showing severe disruption of the

intestinal region just hours after infection in germ-free larvae, and (iii) induction of *il22*

in germ-free larvae exposed to *F. columnare*, since a major function of IL-22 is to

promote gut repair (69). This induction appears to be a consequence of the pathogen-

mediated damage, as there was no observed induction in conventional or re-

conventionalized larvae. The very rapid death of larvae likely caused by this severe

intestinal damage may explain why other common target organs of columnaris disease

showed little damage.

In this study, we showed that *C. massiliae* is a promising probiotic candidate that could

contribute to reduce the use of antibiotic to prevent columnaris diseases in research and

aquaculture settings. Whereas *C. massiliae* provided full and robust protection against all

tested virulent *F. columnare* genomovars and significantly increased survival of exposed

adult conventional zebrafish, further studies are needed to elucidate *C. massiliae*

protection potential in other teleost fish. However, the endogenous nature of *C. massiliae*

suggests that it could establish itself as a long-term resident of the zebrafish larval and

adult microbiota, an advantageous trait when seeking a stable modulation of the bacterial

community over long periods (33).

In conclusion, the use of a simple and tractable zebrafish larval model to mine indigenous

host microbial communities allowed us to identify two independent protection pathways

against the same pathogen. Whereas further study will determine how these pathways

may contribute to protection against a wider range of pathogens, this work also provides

insights into how to the engineering of stable protective microbial communities with

controlled colonization resistance functions.

ACKNOWLEDGEMENTS

We thank Mark McBride, Pierre Boudinot and Rebecca Stevick for critical reading of the manuscript. We are grateful to the late Covadonga Arias for the gift of *F. columnare* ALG 00-530, to Mark McBride for *F. columnare* C#2 strain and to Jean-François Bernardet for all other *F. columnare* strains. Prof. Annemarie Meijer (Leiden University) kindly provided the *myd88* mutant zebrafish line. We thank Julien Burlaud-Gaillard and Rustem Uzbekov the IBiSA Microscopy facility, Tours University, France and the following zebrafish facility teams for providing eggs: José Perez and Yohann Rolin(Institut Pasteur), Nadia Soussi-Yanicostas (INSERM Robert Debré), Sylvie Schneider-Manoury and Isabelle Anselme (UMR7622, University Paris 6) and Frédéric Sohm (AMAGEN Gif-sur-Yvette).

FUNDING

This work was supported by the Institut Pasteur, the French Government's *Investissement d'Avenir* program: *Laboratoire d'Excellence* 'Integrative Biology of Emerging Infectious Diseases' (grant no. ANR-10-LABX-62-IBEID to J.M.G.), the Fondation pour la Recherche Médicale (grant no. DEQ20180339185 to J.M.G.). F.S. was the recipient of a post-doctoral Marie Curie fellowship from the EU-FP7 program, J.B.B. was the recipient of a long-term post-doctoral fellowship from the Federation of European Biochemical Societies (FEBS) and by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 842629. D.P-P was supported by an Institut Carnot MS Postdoctoral fellowship.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CONFLICT OF INTEREST

The authors of this manuscript have the following conflict of interest: a provisional patent application has been filed: "*bacterial strains for use as probiotics, compositions thereof, deposited strains and method to identify probiotic bacterial strains*" by J.-M.G, F. A. S., D.P.-P. and J. B. B. The other authors declare no conflict of interest in relation to the submitted work.

AUTHOR CONTRIBUTIONS:

F. A. S., J. B. B., D.P.-P., J.-P. L. and J.-M.G. designed the experiments. O.R. contributed

to the initial experiments. V.B. and J.-P. L. provided zebrafish material and advice. F. A.

S, J. B. B., D.P.-P., B.A., V.B. and J.-P. L. performed the experiments. S.B., S.H.

performed bacterial genome sequencing and analysis, A.G., S.V., F. A. S and D.P.-P.

performed the bioinformatic and sequence analyses. F. A. S, J. B. B., D.P.-P., J.-P. L. and

J.-M.G. analysed the data and wrote the paper with significant contribution from O.R.

and E.D..

REFERENCES

1.   Rolig AS, Parthasarathy R, Burns AR, Bohannan BJ, Guillemin K. Individual Members of the Microbiota Disproportionately Modulate Host Innate Immune Responses. *Cell host & microbe*. 2015;18(5):613-20.

2.   McFall-Ngai MJ. Unseen forces: the influence of bacteria on animal development. *Dev Biol*. 2002;242(1):1-14.

3.   van der Waaij D, Berghuis-de Vries JM, Lekkerkerk-van der Wees JEC. Colonization resistance of the digestive tract and the spread of bacteria to the lymphatic organs in mice. *The Journal of Hygiene*. 1972;70:335-42.

4.   Cani PD, Delzenne NM. The role of the gut microbiota in energy metabolism and metabolic disease. *Curr Pharm Des*. 2009;15(13):1546-58.

5.   Stecher B, Hardt WD. The role of microbiota in infectious disease. *Trends Microbiol*. 2008;16(3):107-14.

6.   Stecher B, Hardt WD. Mechanisms controlling pathogen colonization of the gut. *Current opinion in microbiology*. 2011;14(1):82-91.

7.   Falcinelli S, Rodiles A, Unniappan S, Picchietti S, Gioacchini G, Merrifield DL, et al. Probiotic treatment reduces appetite and glucose level in the zebrafish model. *Sci Rep*. 2016;6:18061.

8.   Olsan EE, Byndloss MX, Faber F, Rivera-Chavez F, Tsolis RM, Baumler AJ. Colonization resistance: The deconvolution of a complex trait. *The Journal of biological chemistry*. 2017;292(21):8577-81.

9.   Littman DR, Pamer EG. Role of the commensal microbiota in normal and pathogenic host immune responses. *Cell host & microbe*. 2011;10(4):311-23.

10.   Heselmans M, Reid G, Akkermans LM, Savelkoul H, Timmerman H, Rombouts FM. Gut flora in health and disease: potential role of probiotics. *Curr Issues Intest Microbiol*. 2005;6(1):1-7.

11.   Boirivant M, Strober W. The mechanism of action of probiotics. *Curr Opin Gastroenterol*. 2007;23(6):679-92.

12.   Robinson CJ, Bohannan BJ, Young VB. From structure to function: the ecology of host-associated microbial communities. *Microbiol Mol Biol Rev*. 2010;74(3):453-76.

13.   Vollaard EJ, Clasener HA. Colonization resistance. *Antimicrob Agents Chemother*. 1994;38(3):409-14.

14.   Gill HS. Probiotics to enhance anti-infective defences in the gastrointestinal tract. *Best Pract Res Clin Gastroenterol*. 2003;17(5):755-73.

15.   Rawls JF, Samuel BS, Gordon JI. Gnotobiotic zebrafish reveal evolutionarily
conserved responses to the gut microbiota. *Proceedings of the National Academy of*
Sciences of the United States of America. 2004;101(13):4596-601.

16.   Milligan-Myhre K, Charette JR, Phennicie RT, Stephens WZ, Rawls JF,
Guillemin K, et al. Study of host-microbe interactions in zebrafish. *Methods in cell*
biology. 2011;105:87-116.

17.   Burns AR, Guillemin K. The scales of the zebrafish: host-microbiota
interactions from proteins to populations. *Current opinion in microbiology*.
2017;38:137-41.

18.   Douglas AE. Simple animal models for microbiome research. *Nature reviews*
Microbiology. 2019;17(12):764-75.

19.   Flores EM, Nguyen AT, Odem MA, Eisenhoffer GT, Krachler AM. The
zebrafish as a model for gastrointestinal tract-microbe interactions. *Cellular*
microbiology. 2020;22(3):e13152.

20.   Melancon E, Gomez De La Torre Canny S, Sichel S, Kelly M, Wiles TJ, Rawls
JF, et al. Best practices for germ-free derivation and gnotobiotic zebrafish husbandry.
Methods in cell biology. 2017;138:61-100.

21.   Cantas L, Sorby JR, Alestrom P, Sorum H. Culturable gut microbiota diversity
in zebrafish. *Zebrafish*. 2012;9(1):26-37.

22.   Rendueles O, Ferrieres L, Fretaud M, Begaud E, Herbomel P, Levraud JP, et al.
A new zebrafish model of oro-intestinal pathogen colonization reveals a key role for
adhesion in protection by probiotic bacteria. *PLoS pathogens*. 2012;8(7):e1002815.

23.   Caruffo M, Navarrete NC, Salgado OA, Faundez NB, Gajardo MC, Feijoo CG,
et al. Protective Yeasts Control V. anguillarum Pathogenicity and Modulate the Innate
Immune Response of Challenged Zebrafish (Danio rerio) Larvae. *Front Cell Infect*
Microbiol. 2016;6:127.

24.   Perez-Ramos A, Mohedano ML, Pardo MA, Lopez P. beta-Glucan-Producing
Pediococcus parvulus 2.6: Test of Probiotic and Immunomodulatory Properties in
Zebrafish Models. *Frontiers in microbiology*. 2018;9:1684.

25.   Chu W, Zhou S, Zhu W, Zhuang X. Quorum quenching bacteria Bacillus sp.
QSI-1 protect zebrafish (Danio rerio) from Aeromonas hydrophila infection. *Sci Rep*.
2014;4:5446.

26.   Wang Y, Ren Z, Fu L, Su X. Two highly adhesive lactic acid bacteria strains are
protective in zebrafish infected with Aeromonas hydrophila by evocation of gut
mucosal immunity. *Journal of applied microbiology*. 2016;120(2):441-51.

27.   Qin C, Zhang Z, Wang Y, Li S, Ran C, Hu J, et al. EPSP of L. casei BL23
Protected against the Infection Caused by Aeromonas veronii via Enhancement of
Immune Response in Zebrafish. *Frontiers in microbiology*. 2017;8:2406.

28.   Girija V, Malaikozhundan B, Vaseeharan B, Vijayakumar S, Gobi N, Del Valle
Herrera M, et al. In vitro antagonistic activity and the protective effect of probiotic
Bacillus licheniformis Dahb1 in zebrafish challenged with GFP tagged Vibrio
parahaemolyticus Dahv2. *Microbial pathogenesis*. 2018;114:274-80.

29.   Lin YS, Saputra F, Chen YC, Hu SY. Dietary administration of Bacillus
amyloliquefaciens R8 reduces hepatic oxidative stress and enhances nutrient
metabolism and immunity against Aeromonas hydrophila and Streptococcus agalactiae
in zebrafish (Danio rerio). *Fish Shellfish Immunol*. 2019;86:410-9.

30.   Garcia JC, LaFrentz BR, Waldbieser GC, Wong FS, Chang SF. Characterization
of atypical Flavobacterium columnare and identification of a new genomovar. *Journal
of fish diseases*. 2018;41(7):1159-64.

31.   van der Vaart M, van Soest JJ, Spaink HP, Meijer AH. Functional analysis of a
zebrafish myd88 mutant identifies key transcriptional components of the innate immune
system. *Disease models & mechanisms*. 2013;6(3):841-54.

32.   Pham LN, Kanther M, Semova I, Rawls JF. Methods for generating and
colonizing gnotobiotic zebrafish. *Nature protocols*. 2008;3(12):1862-75.

33.   Lemon KP, Armitage GC, Relman DA, Fischbach MA. Microbiota-targeted
therapies: an ecological perspective. *Sci Transl Med*. 2012;4(137):137rv5.

34.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics (Oxford, England)*. 2014;30(15):2114-20.

35.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.
SPAdes: a new genome assembly algorithm and its applications to single-cell
sequencing. *Journal of computational biology : a journal of computational molecular
cell biology*. 2012;19(5):455-77.

36.   Ha SM, Kim CK, Roh J, Byun JH, Yang SJ, Choi SB, et al. Application of the
Whole Genome-Based Bacterial Identification System, TrueBac ID, Using Clinical
Isolates That Were Not Identified With Three Matrix-Assisted Laser
Desorption/Ionization Time-of-Flight Mass Spectrometry (MALDI-TOF MS) Systems.
Annals of laboratory medicine. 2019;39(6):530-6.

37.   Volant S, Lechat P, Woringer P, Motreff L, Malabat C, Kennedy S, et al.
SHAMAN: a user-friendly website for metataxonomic analysis from raw reads to
statistical analysis. *bioRxiv*. 2019:2019.12.18.880773.

38.   Rognes T, Flouri T, Nichols B, Quince C, Mahe F. VSEARCH: a versatile open
source tool for metagenomics. *PeerJ*. 2016;4:e2584.

39.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Research. 2012;41(D1):D590-D6.

40.   Yarza P, Yilmaz P, Pruesse E, Glockner FO, Ludwig W, Schleifer KH, et al.
Uniting the classification of cultured and uncultured bacteria and archaea using 16S
rRNA gene sequences. *Nature reviews Microbiology*. 2014;12(9):635-45.

41.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*. 2014;15(12):550.

42.   Soto E, Mauel MJ, Karsi A, Lawrence ML. Genetic and virulence
characterization of Flavobacterium columnare from channel catfish (Ictalurus
punctatus). *Journal of applied microbiology*. 2008;104(5):1302-10.

43.   Olivares-Fuster O, Bullard SA, McElwain A, Llosa MJ, Arias CR. Adhesion
dynamics of Flavobacterium columnare to channel catfish Ictalurus punctatus and
zebrafish Danio rerio after immersion challenge. *Diseases of aquatic organisms*.
2011;96(3):221-7.

44.   Declercq AM, Haesebrouck F, Van den Broeck W, Bossier P, Decostere A.
Columnaris disease in fish: a review with emphasis on bacterium-host interactions. *Vet
Res*. 2013;44:27.

45.   Li N, Zhu Y, LaFrentz BR, Evenhuis JP, Hunnicutt DW, Conrad RA, et al. The
Type IX Secretion System Is Required for Virulence of the Fish Pathogen
Flavobacterium columnare. *Applied and environmental microbiology*. 2017;83(23).

46.   Cheesman SE, Neal JT, Mittge E, Seredick BM, Guillemin K. Epithelial cell
proliferation in the developing zebrafish intestine is regulated by the Wnt pathway and
microbial signaling via Myd88. *Proceedings of the National Academy of Sciences of the
United States of America*. 2011;108 Suppl 1:4570-7.

47.   Mena KD, Gerba CP. Risk assessment of Pseudomonas aeruginosa in water.
Reviews of environmental contamination and toxicology. 2009;201:71-115.

48.   Goncalves Pessoa RB, de Oliveira WF, Marques DSC, Dos Santos Correia MT,
de Carvalho E, Coelho L. The genus Aeromonas: A general approach. *Microbial
pathogenesis*. 2019;130:81-94.

49.   Stephens WZ, Burns AR, Stagaman K, Wong S, Rawls JF, Guillemin K, et al.
The composition of the zebrafish intestinal microbial community varies across
development. *ISME J*. 2016;10(3):644-54.

50.   Chatzidaki-Livanis M, Coyne MJ, Comstock LE. An antimicrobial protein of the
gut symbiont Bacteroides fragilis with a MACPF domain of host immune proteins.
Molecular microbiology. 2014;94(6):1361-74.

51.   Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, et al. A
type VI secretion-related pathway in Bacteroidetes mediates interbacterial antagonism.
Cell host & microbe. 2014;16(2):227-36.

52.   Roelofs KG, Coyne MJ, Gentyala RR, Chatzidaki-Livanis M, Comstock LE.
Bacteroidales Secreted Antimicrobial Proteins Target Surface Molecules Necessary for
Gut Colonization and Mediate Competition In Vivo. *mBio*. 2016;7(4).

53.   Jacobson A, Lam L, Rajendram M, Tamburini F, Honeycutt J, Pham T, et al. A
Gut Commensal-Produced Metabolite Mediates Colonization Resistance to Salmonella
Infection. *Cell host & microbe*. 2018;24(2):296-307.e7.

54.   Kamada N, Seo SU, Chen GY, Nunez G. Role of the gut microbiota in immunity
and inflammatory disease. *Nat Rev Immunol*. 2013;13(5):321-35.

55.   Ganz J, Melancon E, Eisen JS. Zebrafish as a model for understanding enteric
nervous system interactions in the developing intestinal tract. *Methods in cell biology*.
2016;134:139-64.

56.   Bates JM, Mittge E, Kuhlman J, Baden KN, Cheesman SE, Guillemin K.
Distinct signals from the microbiota promote different aspects of zebrafish gut
differentiation. *Dev Biol*. 2006;297(2):374-86.

57.   Yan Q, van der Gast CJ, Yu Y. Bacterial community assembly and turnover
within the intestines of developing zebrafish. *PLoS One*. 2012;7(1):e30603.

58.   Costello EK, Stagaman K, Dethlefsen L, Bohannan BJ, Relman DA. The
application of ecological theory toward an understanding of the human microbiome.
Science. 2012;336(6086):1255-62.

59.   Rogers GB, Hoffman LR, Carroll MP, Bruce KD. Interpreting infective
microbiota: the importance of an ecological perspective. *Trends Microbiol*.
2013;21(6):271-6.

60.   Burns AR, Stephens WZ, Stagaman K, Wong S, Rawls JF, Guillemin K, et al.
Contribution of neutral processes to the assembly of gut microbial communities in the
zebrafish over host development. *ISME J*. 2016;10(3):655-64.

61.   Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition:
surviving and thriving in the microbial jungle. *Nature reviews Microbiology*.
2010;8(1):15-25.

62.   Shafquat A, Joice R, Simmons SL, Huttenhower C. Functional and phylogenetic
assembly of microbial communities in the human microbiome. *Trends Microbiol*.
2014;22(5):261-6.

63.   Shade A, Peter H, Allison SD, Baho DL, Berga M, Burgmann H, et al.
Fundamentals of microbial community resistance and resilience. *Frontiers in
microbiology*. 2012;3:417.

64.   Willing BP, Russell SL, Finlay BB. Shifting the balance: antibiotic effects on
host-microbiota mutualism. *Nature reviews Microbiology*. 2011;9(4):233-43.

65.   Brugman S, Liu KY, Lindenbergh-Kortleve D, Samsom JN, Furuta GT,
Renshaw SA, et al. Oxazolone-induced enterocolitis in zebrafish depends on the
composition of the intestinal microbiota. *Gastroenterology*. 2009;137(5):1757-67 e1.

66.   Salonen A, Nikkila J, Jalanka-Tuovinen J, Immonen O, Rajilic-Stojanovic M,
Kekkonen RA, et al. Comparative analysis of fecal DNA extraction methods with
phylogenetic microarray: effective recovery of bacterial and archaeal DNA using
mechanical cell lysis. *J Microbiol Methods*. 2010;81(2):127-34.

67.   Kunin V, Engelbrektson A, Ochman H, Hugenholtz P. Wrinkles in the rare
biosphere: pyrosequencing errors can lead to artificial inflation of diversity estimates.
Environ Microbiol. 2010;12(1):118-23.

68.   Bernardet J-F, Bowman JP. The Genus Flavobacterium. *Prokaryotes*.
2006;7:481-531.

69.   Shabgah AG, Navashenaq JG, Shabgah OG, Mohammadi H, Sahebkar A.
Interleukin-22 in human inflammatory diseases and viral infections. *Autoimmunity*
reviews. 2017;16(12):1209-18.
1009
1010

1011 **TABLES**

1012

1013 **Table 1. The 10 strains composing the core of zebrafish larvae microbiota**

1014 Bacterial strains consistently detected at all time points (6 and 11 dpf) in all experiment

1015 runs and constituting the core of conventional zebrafish larval microbiota and their

1016 taxonomic affiliation.

1017

Bacterial species of the core zebrafish microbiota	ANI %[a]	16S rRNA (%)[b]	*recA* (%)[c]	*rplC* (%)[d]
Aeromonas veronii 1	96.52	98.27	97.00	99.84
Aeromonas veronii 2	96.58	99.53	98.31	99.68
Aeromonas caviae	97.97	99.94	98.78	99.84
Chryseobacterium massiliae	95.85	99.86	96.61	99.84
Phyllobacterium myrsinacearum	98.58	99.86	99.72	100
Pseudomonas sediminis	96.12	99.73	97.70	99.84
Pseudomonas mossellii	99.39	98.27	100	99.84
*Pseudomonas nitroreducens**	92.14	99.80	94.95	99.06
*Pseudomonas peli**	88.84	99.20	91.51	95.44
*Stenotrophomas maltophilia**	90.94	97.85	95.38	99.08

1018 [a] Average Nucleotide Identity value

1019 [b] 16S rRNA gene sequence similarity

1020 [c] *recA* gene sequence similarity

1021 [d] *rplC* gene sequence similarity

1022 *Species ambiguously identified

1023

1024

1025 **FIGURES**

1026

1027

Figure 1. *Flavobacterium columnare* **kills germ-free but not conventional zebrafish.**
6 dpf (corresponding to 0 dpi) GF or Conv zebrafish larvae were exposed to different
doses of *F. columnare*[ALG] by bath immersion and transferred after 3h into sterile water.
Mean survival is represented by a thick horizontal bar with standard deviation. For each
condition, n = 12 zebrafish larvae. Larvae mortality rate was monitored daily and
surviving fish were euthanized at day 9 post infection. Statistics correspond to unpaired,
non-parametric Mann-Whitney test comparing all conditions to non-infected GF (left) or
Conv (right). ****: $p<0.0001$; ***: $p<0.001$; *: $p<0.05$, absence of *: non-significant.
Blue mean bars correspond to non-exposed larvae and red mean bars correspond to larvae
exposed to *F. columnare*.

1038

1039

1040

1041

Figure 2 Analysis of protection against *F. columnare* infection after antibiotic dysbiosis. A: Response of zebrafish larvae to exposure to *F. columnare*[ALG] after antibiotic-induced dysbiosis with a diagram showing timing and treatments of the experiment. **B:** A 24h period after antibiotic treatment allows the recovery of protection in kanamycin-treated zebrafish larvae with a diagram showing timing and treatments. Mean survival is represented by a thick horizontal bar with standard deviation. For each condition, n = 12 zebrafish larvae. Blue mean bars correspond to larvae not exposed to the pathogen and red mean bars correspond to exposed larvae. Larvae mortality rate was monitored daily and surviving fish were euthanized at day 9 post exposition to the pathogen. Indicated statistics correspond to unpaired, non-parametric Mann-Whitney test. ****: p<0.0001; absence of *: non-significant. **C:** Community recovery profile with streptomycin/penicillin treatment. **D:** Community recovery profile with kanamycin treatment. Pools of 10 larvae were collected for 16S rRNA sequencing.

Figure 3. Protection against *F. columnare* in zebrafish re-conventionalized with individual or mixed bacterial strains isolated from zebrafish. A: Determination of the level of protection provided by each of the 10 bacterial strains composing the core protective zebrafish microbiota. Bacteria were added individually to the water on hatching day (dose 5.10^5 cfu/mL). **B**: Level of protection provided by different amount of *C. massiliae* and Mix9. Mix9 only protected at the highest inoculum doses. Mean survival is represented by a thick horizontal bar with standard deviation. For each condition, n = 12 zebrafish larvae. Blue mean bars correspond to larvae not exposed to the pathogen and red mean bars correspond to exposed larvae. Larvae mortality rate was monitored daily and surviving fish were euthanized at day 9 post exposition to the pathogen. Indicated statistics correspond to unpaired, non-parametric Mann-Whitney test. ****: $p<0.0001$; absence of *: non-significant.

1072

Figure 4. Zebrafish immune response to *F.columnare* infection. A-C: qRT-PCR
analysis of host gene expression, 18 hours after exposure to *F. columnare,* in larvae re-
conventionalized with indicated bacteria or bacterial mixes; each point corresponds to an
individual larva. Expression of *il10* (**A**), *il1b* (**B**), and *il22* (**C**), by wild-type AB zebrafish;
D: Comparison of the survival of *myd88-/-* and background-matched *myd88+/+* zebrafish
after re-conventionalization and exposure to *F. columnare*[ALG]. Mean survival is
represented by a thick horizontal bar with standard deviation. For each condition, n = 12
zebrafish larvae. Larvae mortality rate was monitored daily and surviving fish were
euthanized at day 9 post exposition to the pathogen. A-D: Blue bars correspond to larvae
not exposed to the pathogen and red mean bars correspond to exposed larvae. Indicated
statistics correspond to unpaired, non-parametric Mann-Whitney test. ****: p<0.0001;
**: p<0.005*: p<0.05, absence of *: non-significant.
1085

Figure 5. Intestine of infected germ-free zebrafish displays severe disorganization.
Germ-free, conventional and re-conventionalized zebrafish larvae. Re-conventionalized
zebrafish were inoculated at 4 dpf with Mix9 or *C. massiliae.* **A**: Representative picture
of intestines of non-infected larvae. Fish were fixed for histology analysis or electron
microscopy at 7 dpf. **B**: Representative picture of intestines of infected larvae exposed
at 7 dpf to *F.columnare*[ALG]. In **A** and **B**: *Left column:* Toluidine blue staining of Epon-
embedded zebrafish larvae for Light microscopy. *Right column*: Transmission electron
microscopy. at 7 dpf (*right*). L= intestinal lumen.

1098

1099

Figure 6. Pre-exposure to *C. massiliae* protects larval and adult zebrafish against *F. columnare* infection

A: Zebrafish larvae were inoculated at 4 dpf with 5.10^5 cfu/mL of *C. massiliae* for 48h before infection at 6 dpf with virulent *F. columnare* strains. **B:** Survival of adult zebrafish with or without pre-exposure to *C. massiliae* (2.10^6 cfu/mL for 48h) followed by exposure to *F. columnare*[ALG] (5.10^6 cfu/mL for 1h) Mean survival is represented by a thick horizontal bar with standard deviation. For each condition, n = 12 zebrafish larvae or 10 adult. Zebrafish mortality rate was monitored daily and surviving fish were euthanized at day 9 post exposition to the pathogens. Blue bars correspond to larvae not exposed to the pathogen and red mean bars correspond to exposed larvae. Indicated statistics correspond to unpaired, non-parametric Mann-Whitney test. ****: $p<0.0001$; **: $p<0.005$; absence of *: non-significant.

1112

1113

1114