Synchronizing vigilance or taking turns as sentinels? The importance of testing

coordination

Running title: Synchronized not sentinel high-quality vigilance

Annemarie van der Marel [a], Jane M. Waterman [a], Marta López-Darias [b]

[a] *Department of Biological Sciences, University of Manitoba, Winnipeg, Canada*

[b] *Island Ecology and Evolution Research Group, Instituto de Productos Naturales y Agrobiología*

(IPNA-CSIC), La Laguna, Tenerife, Canary Islands, Spain

Corresponding author: Annemarie van der Marel, email: avdmarel@outlook.com, phone: +1 204

330 7251, *current address*: Department of Biological Sciences, University of Cincinnati, 614

Rieveschl Hall, Cincinnati, OH, USA, 45221-0006

Abstract

Diurnal species can reduce the cost of the trade-off between feeding and predation risk by 1)

performing low-quality (vigilance while performing another behavior) instead of high-quality

vigilance (only performing vigilance) or 2) by vigilance coordination either by taking turns

acting as sentinels or by synchronizing vigilance bouts. A sentinel system assumes that sentinels

are located at raised positions, alarm signal, and alternate vigilance bouts. However, many

species with a described sentinel system have not been tested yet for coordination. We set out to

study coordinated behavior and the reasons for this behavior in the invasive Barbary ground

squirrel, *Atlantoxerus getulus*, using behavioral observations and genetic analyses. This species

performs a type of vigilance ('perch behavior') seemingly similar to sentinel behavior as

individuals performed high-quality vigilance at raised locations for over 30 s, but alternating

coordination is unknown. Perch behavior was coordinated but synchronized instead of taking

turns (sentinel). Both sexes performed perch behavior and individuals performed perch behavior

in the absence and presence of kin and offspring. We found that survival or time spent foraging

did not decrease for perched individuals, nor that individual survival increased. Perch behavior in

the invasive population of *A. getulus* may be synchronized 1) to perform an optimal activity

when satiated (low costs), and, 2) may be an adaptation to habitat structure (large benefits). Our

study demonstrates that the cost of the vigilance/foraging trade-off may not be high for invasive

species and argued the importance of testing for coordination in species with described sentinel

systems.

Keywords: anti-predator behavior, collective behavior, coordinated vigilance, invasive species,

ground squirrel

Introduction

Environmental conditions impose a notable investment in vigilance for diurnal species, as vigilance is required to avoid predation resulting in a trade-off with foraging(Bednekoff and Lima 1998). Thus, an individual will balance the trade-off to increase its survival chances. Different vigilance behaviors evolved due to the cost associated with this trade-off between vigilance and foraging. First, some species can reduce the cost of this trade-off by performing nonexclusive or low-quality vigilance, i.e., individuals are vigilant while performing another behavior (e.g. Cape ground squirrels, *Xerus inauris,* Unck et al. (2009), or eastern grey kangaroos, *Macropus giganteus,* Favreau et al. (2015)). Low-quality vigilance is often seen in habitats where the risk imposed by predators is relatively low. In riskier habitats, individuals increase their time spent in exclusive or high-quality vigilance –vigilance without performing any other behavior– while reducing the time spent foraging (Unck et al. 2009; Périquet et al. 2012).

The cost of the trade-off between feeding activity and predator avoidance can additionally lead to the evolution of dependent vigilance within groups –individuals within the group coordinate the vigilance bouts– either by synchronizing their vigilance bouts or by taking turns acting as sentinels (Beauchamp 2015). Synchronization of vigilance bouts arises when individuals are either vigilant or non-vigilant at the same time as other group members, resulting in periods that either no group members or multiple group members are simultaneously vigilant (Beauchamp 2015). This behaviour, which is frequent in both birds and mammals (Ebensperger et al. 2006; Pays, Jarman, et al. 2007; Pays, Renaud, et al. 2007; Pays et al. 2009, 2012; Öst and Tierala 2011; Podgórski et al. 2016; McDougall and Ruckstuhl 2018), could occur through different mechanisms. First, vigilance levels of the group may convey the group's fear level

('contagion-of-fear' hypothesis, Sirot and Touzalin 2009; Beauchamp 2015), i.e., the level of fear

is heightened when other group members experience many vigilant neighbors as an indication of

potential predation risk (Beauchamp 2015). Second, synchronization can arise when individuals

benefit by detecting predators and by reaching safety before group members, as predators target

individuals that are last to find cover ('predators-targeting-behavior' hypothesis, Sirot and

Touzalin 2009; Beauchamp 2015). Third, multiple guards may be necessary to increase the

collective view of the surroundings (Shackleton et al. 2018). Additionally, synchronization may

occur when individuals respond to a) the same external stimulus (e.g., a predator, termed induced

vigilance, Blanchard and Fritz 2007), b) group members that ceased chewing (McDougall and

Ruckstuhl 2018), or c) group members that started feeding (Beauchamp 2015). Thus, group

members may simply be copying the behavior of nearest neighbors (Evans et al. 2018).

The foraging-vigilance trade-off can also lead to the evolution of sentinel behavior. In

species with a sentinel system, one or two group members are always vigilant because

individuals forego feeding and take turns performing high-quality vigilance from raised positions

(Rasa 1987; Bednekoff 1997; Beauchamp 2015). Some highly social bird and mammal species

living or foraging in open arid habitats perform sentinel behavior (Rasa 1977; Ferguson 1987;

McGowan and Woolfenden 1989; Clutton-Brock 1999; Manser 1999; Wright, Berg, et al. 2001;

Newbold et al. 2008). A sentinel system assumes that sentinels warn group members of danger

(Bednekoff 1997) and that there is a continuous turn-over of who is sentinel, but there is no

change in sentinel number (McGowan and Woolfenden 1989; Bednekoff 2015). Since

non-sentinels can be located at raised positions or can alarm signal, Bednekoff (2015) argued that

the main criterion of a sentinel system is the alternation of the high-quality vigilance bouts. As

alternating reduces the cost of the foraging and vigilance trade-off for group members. However,

out of 40 species with a described sentinel system, coordination was tested for only 3 species,

although suggested for another 14 species (Bednekoff 2015). A sentinel system may evolve

because sentinels appear to be more effective at detecting predators due to the more focused

vigilance and the elevated positions they adopt (Ridley et al. 2013). Besides, sentinel behavior

may be a cooperative behavior (Santema and Clutton-Brock 2013), where individuals gain

fitness benefits through kin selection (sentinels only keep watch when close kin are present) or

reciprocal altruism (Rasa 1987; McGowan and Woolfenden 1989; Brandl and Bellwood 2015).

Nevertheless, sentinel behavior can be costly, as it is more likely to attract the attention of

predators (Ridley et al. 2013). Consequently, coordination of guarding bouts may be a necessity

to divide the costs of being on guard (increased predation risk and/or less time available for

foraging) over group members (Ridley et al. 2013). Yet, sentinels may be considered selfish

when they can detect a predator earlier and reach safety before group members or when the

decision to become sentinel is state-dependent (Bednekoff 1997; Clutton-Brock 1999; Bednekoff

2001; Wright, Berg, et al. 2001; Wright, Maklakov, et al. 2001; Bednekoff and Woolfenden

2003). Most species with a sentinel system present stable social groups with high-relatedness

among group members due to low levels of promiscuity (Fernández et al. 2003), suggesting that

kin selection is an important mechanism for sentinel behavior to occur. However, other species

are also sentinel when nonkin are nearby (Bednekoff and Woolfenden 2003). Thus, the role of

kin selection in coordinated vigilance is not necessarily a prerequisite.

Besides environmental conditions, social conditions –sex, age and possibly kinship- may

influence vigilance. For instance, due to intraspecific competition (Reboreda and Fernandez

1997) or mate protection (Horrocks and Hunte 1986) males are often the sex performing the

most vigilance (Li et al. 2012; Pays et al. 2012; Whiteside et al. 2016). As well, adults normally

spend more time vigilant than juveniles (newborn sexually immature individuals that need more

time feeding to maintain a high growth rate; Lung and Childress 2007; Pays et al. 2012). In some

species kinship positively affects vigilance (Black and Owen 1989; Griesser 2003), although this

pattern is not universal (Fairbanks and Dobson 2010; Quirici et al. 2013).

In the invasive population of Barbary ground squirrels, *Atlantoxerus getulus*, on

Fuerteventura, Canary Islands, Spain, both males and females perform high-quality vigilance

(actively scan the environment > 30 s without performing any other behavior for approximately

40% of their daily activity budget) while positioned on top of, mainly, man-made rock walls (van

der Marel et al. 2019). We termed this particular vigilance behavior 'perch behavior' because it is

different from high-quality vigilance in that individuals move away from foraging areas to

perform exclusive vigilance in a raised position (van der Marel et al. 2019). As well, perched

individuals alarm call to inform group members of danger (van der Marel et al. 2019). Hence,

perch behavior seems similar to sentinel behavior, although following Bednekoff (2015) such

behavior cannot be considered as sentinel behavior unless it is coordinated by individuals taking

turns. Thus, our main aim in this contribution is to deepen the understanding of factors

explaining vigilance behavior in a social ground squirrel, which will aid in our comprehension of

the evolution of coordinated vigilance in general. More specifically, our objectives are to

investigate whether 1) Barbary ground squirrels coordinate perch behavior either by taking turns

acting as sentinels or by synchronization, and are situated in higher positions when perched than

when they are performing other behaviors; 2) perch behavior in this species is an induced

vigilance; and, 3) social and environmental conditions -sex, age, kinship, resource availability or

predation pressure- influence perch behavior using the hypotheses and predictions presented in

Table 1.

Materials and methods

Study sites, species, and trapping protocol

We studied an invasive population of Barbary ground squirrels in the northwest of

Fuerteventura, the Canary Islands, Spain (28°34'60" N, 13°58'0" W) from March through July

2014, January through July 2015, and January through June 2016. The study area consisted of

three study sites that differed in the number of squirrels and size, but that did not represent

distinct squirrel' populations (see van der Marel et al. 2020 for more details). Fuerteventura has

an arid climate and semi-desert habitats characterized by xerophytic scrubland and ravines

caused by erosion (del Arco Aguilar et al. 2010). Dams and rock walls, built to make terraces for

land cultivation and to fence properties, characterize island landscapes and also our field sites,

and function as shelters and perches for the Barbary ground squirrels to watch for predators

(López-Darias and Lobo 2008; van der Marel et al. 2019). The main predator of Barbary ground

squirrels on Fuerteventura is the Eurasian buzzard, *Buteo buteo* (Gangoso et al. 2006). Other

aerial predators are the common raven, *Corvus corax* (Gangoso et al. 2006) and the common

kestrel, *Falco tinnunculus* (López-Darias and Lobo 2008), whereas the invasive feral cat, *Felis*

catus, is the only terrestrial predator (Medina et al. 2008).

Barbary ground squirrel is a social species because individuals share sleeping burrows and

show spatiotemporal overlap and cohesiveness (van der Marel et al. 2020). Females share

sleeping burrows with related adult females, whereas males share sleeping burrows with

unrelated males and subadults of either sex when they are present. Throughout the day, males

and females of different sleeping burrow associations can be active in the same area in a

site (van der Marel et al. 2020). Adult and subadult males and females are distinguished by the

state of their primary sexual characters (scrotum or vulva and nipples, van der Marel et al. 2020).

Subadults are over six months old, reached adult body size, but males do not have descended

testes and females do not have swollen vulva and nipples during the mating season (i.e. days

when females are in estrus). The mating season is distinct as males regress their scrotum once all

females mated (van der Marel 2019).

As part of a broader research project, we continuously monitored the population of the study

area by trapping Barbary ground squirrels weekly throughout the three field seasons. We

followed the trapping and marking procedures described in van der Marel et al. (2019, 2020)

and Piquet et al. (2018). For this study, we additionally measured squirrel body mass using a

spring scale (± 5 g; Pesola AG, Baar, Switzerland) and hindfoot length (mm) using a Digimatic

Plastic Caliper (Mitutoyo Corporation, Kawasaki, Japan), both as proxy measurements for body

size (Schulte-Hostedde et al. 2001).

Behavioral observations

To characterize perch behavior of Barbary ground squirrels and test the predictions

presented in Table 1, we used 10 min scan samples and all-occurrence sampling of perch bouts

throughout each behavioral observation period in addition to performing 10 min focal follows

(Altmann 1974). Table 1 also provides the behavioral observation method used to test our

predictions.

In our scan- and all-occurrence sampling, we observed the squirrels within each site from

elevated areas and roads at a distance of approximately 50 m, which did not affect the squirrels'

behavior. We defined a group as a set of individuals that were in each other's visible range during

an observation period (Stankowich 2003; van der Marel et al. 2019). We conducted behavioral

observations during all three field seasons between 10:00 h GMT and 2 h before sunset, i.e.,

when the squirrels are active above ground (Machado 1979; van der Marel et al. 2019). In 2014,

we first established sites 1 and 2 at the beginning of the field season, while site 3 was set at the

end, resulting in an unequal number of behavioral observations among sites this year (for a

breakdown of the total observation periods, the number of scans, and hours observed per site and

year see SM1). Primarily, we recorded observed perch behavior, low-quality vigilance (vigilance

at any location that lasted < 30 s or was performed while doing other behaviors), and feeding

(actively searching for, manipulating, and/or ingesting a food item). We provided an ethogram of

all recorded behaviors in SM2). We also recorded whether any predators or other external stimuli

were observed in a scan, as animals may respond to the same external stimulus (Beauchamp et

al. 2012; McDougall and Ruckstuhl 2018). Finally, we included the height from which the

squirrels performed the behaviors. To enter behavioral data, we used the spreadsheet program

Numbers (Apple, Cupertino, CA, USA) on an iPod (Apple, Cupertino, CA, USA) from 2014

until June 2015, and thereafter Prim8 Software (McDonald and Johnson 2014) on an Android

phone (Motorola Droid A850). Prim8 software measured the exact duration of perch behavior

instead of the number of times the behavior was observed, which we used to analyse

coordination of perch bouts by taking the proportion of time individuals were perched

(predictions 2 and 3; Table 1). In our analyses, we selected individuals that were observed for at

least 50 min over a minimum of five observation periods (Edwards and Waterman 2011).

To study perch behavior in more detail (predictions 1, 5, 6, 7, and 8, Table 1), we also

performed 10 min focal follows in 2015 (Altmann 1974), recording focal follows for 81

individuals (ranging from 1 to 7 focal follows per individual). Before sampling, we recorded

which of the three sites we sampled, focal ID, and the number and ID's of the individuals in sight

during an observation period (i.e. group size, see van der Marel et al. 2019). During the focal

follow, we recorded the behavior (following the behaviors noted in the ethogram, SM2),

location, the height of position (using a tape measure), the number of other individuals perching,

and the distance from the focal to cover. We randomized the order of the site in which we

performed focal follows using a random number generator (MS Excel 2013) and the order of

focal follows among individuals observed in the site directly in the field using a randomizer

application on an Android phone (Randomizer v. 0.2.6-beta, Mahé 2016). We only performed

focal follows when no predators were observed, removing the possibility that individuals were

responding to the same external stimulus (Beauchamp et al. 2012; McDougall and Ruckstuhl

2018).

Perch behavior general description

Using both the scan- and all-occurrence sampling to investigate whether all individuals

perform perch behavior, we first calculated the percentage of individuals that perch from all

marked individuals during the three field seasons. We then calculated the number of scans where

one, two, or more than two individuals were perched, and the proportion of group members

perched per scan excluding observations performed during the mating season. We analyzed the

relationship between the proportion of perched group members and group size with a beta

regression from the 'betareg' package version 3.1-1 (Cribari-Neto and Zeileis 2010) and the

likelihood ratio test from the 'lmtest' version 0.9-36 (Zeileis 2002). To study whether perch

behavior functioned as induced vigilance, we compared the proportion of perched individuals

when a predator was observed or when we observed an alarm call (external stimulus present) *vs.*

when no predator was observed or when no alarm call was heard (external stimulus absent) using

a Mann-Whitney U test due to the non-normality of the data (prediction 0; Table 1). To test

whether individuals perch from elevated positions (prediction 1; Table 1), we used a Mann-

Whitney U test to examine differences in height between perched squirrels and squirrels that

performed any other behavior.

To measure coordination of perch bouts (predictions 2 and 3; Table 1), we calculated the

difference between the observed and expected proportions of time when at least one individual

was perched per group (Pays, Jarman, et al. 2007; Öst and Tierala 2011; Beauchamp 2015). We

measured the expected time as $1 - \prod_{i=1}^{n}(1 - p_i)$, where n is the sample size and p_i is the

proportion of time individual i is perched, i.e., total time spent perched divided by the total time

observed for that individual (Beauchamp 2015; Bednekoff 2015). We used the number of

observation periods as our sample size if the individuals seen in an observation period belonged

to the same group. We measured the observed time as the sum of all perch bouts in the group

divided by the sum that each individual in the group was observed. If the difference in

proportions (p_{obs}-p_{exp}) did not differ from 0, then individual perch bouts were independent and

not coordinated; if the difference in proportions was greater than 0, then perch bouts were

sentinel; and if the difference was smaller than 0, then perch bouts were synchronized

(Beauchamp 2015). We calculated the duration of observed and expected time when at least one

individual was perched for all individuals, only adult individuals, and only adult females. We

excluded days that females were in estrus because male behavior changed as they search

competitively for estrous females, similar to what has been reported for the Cape ground squirrel

(Waterman 1998). We excluded unidentified individuals because a different unidentified squirrel

could have switched with the initial unidentified squirrel during a scan. Thus, start and endpoints

of perch bouts could have corresponded to different unidentified squirrels. We used a non-

parametric paired test (Wilcoxon signed-rank test with continuity correction) to test for

differences between the expected and observed values per observation period.

Perch behavior and social conditions analyses

To test whether adults and subadults spent more time perched than juveniles (prediction 4;

Table 1), we calculated activity budgets for perch behavior, low-quality vigilance, and foraging

of adults, subadults, and juveniles. We divided the number of scans in which each individual was

observed performing each behavior by the total scans in which they were observed. In the

absence of any sex differences in time allocation of low-quality vigilance and perch behavior

(van der Marel et al. 2019), we tested for differences among adults, subadults, and juveniles for

males and females combined. Since our data were not normally distributed, we used Kruskal-

Wallis tests to account for differences in proportion time spent in low-quality vigilance, perch

behavior and foraging for adults, subadults, and juveniles. If the results were significant, we

performed post-hoc tests using a "Holm" correction (Holm 1979).

To examine whether individuals only perch when close kin are nearby (prediction 5; Table

1), we extracted information from our focal follows ($N = 191$) and analyzed relationships

between individuals using genetics data extracted from van der Marel et al. (2020). Relatedness

ranged from non-related (r = 0) to highly-related (r = 0.69), but the average within-group

relatedness was 0.19 ± 0.01. We used generalized linear mixed models (GLMMs) to analyze the

proportion of time spent perched in relation with multiple measures for both our independent

variables (alone *vs.* in-group, kin-selection, and parental behavior), with ID as the random factor.

We then compared the model against a null model using the likelihood ratio test from the 'lmtest'

version 0.9-36 (Zeileis 2002). We used a 'glmmADMB' with the beta distribution (Bolker et al.

2012; Fournier et al. 2012) on the proportion of time spent perched for kin (parent-offspring to

first cousin; $r > 0.125$) *vs.* nonkin (unrelated to second cousin; $r < 0.125$), and a Spearman's rank

correlation to test for a correlation between proportional time spent perched and relatedness of

the group. To test whether perch behavior was a parental behavior (prediction 6; Table 1), we

analyzed the time adult females spent perched in presence and absence of vulnerable offspring

using a glmmADMB (Bolker et al. 2012; Fournier et al. 2012) with ID as a random factor.

Perch behavior and survival analyses

To study whether survival increased for perched individuals, we recorded whether perched

individuals were closer to cover than individuals performing another behavior and whether

individuals were perched when alone and, in a group (prediction 7; Table 1). We only used data

from individuals that were both feeding and foraging within the 10 min focal follow. Since the

average proportion of time spent perched was not normally distributed, we used a paired

Wilcoxon signed-rank test to test for a difference in distance to cover for perched *vs.* feeding

individuals. We then tested for differences in perch behavior when the individual was alone

compared to when group members were present using a glmmADMB.

We considered individuals as survivors if the focal squirrels were trapped in the following

field season (2016) to study whether survival decreased for perched individuals (prediction 8;

Table 1). We measured survival as the last Julian day observed (number of days counted from

our first fieldwork day, 12 February 2014 = Julian day 0) and age at disappearance. As we

performed multiple focal follows on the same individual, we averaged the proportional time

spent perched for analyses where we only had one measure per individual (survival, Julian date

lived until, and age at disappearance). We tested whether perched individuals (proportional time

spent perched > 0.1) experienced higher predation risk than individuals that were not perched

(proportional time spent perched < 0.1) using a Mann-Whitney U test. We performed a Spearman

rank correlation on the proportion of time spent perched and the last Julian date observed and a

Kruskal-Wallis test on age at disappearance. We calculated the proportion of time that each

squirrel allocated to foraging and perch behavior, which we analyzed using a Spearman's rank

correlation. We measured body condition by taking the residuals from a linear regression of log

body mass (g) on log hind foot length (mm) (Labocha et al. 2014; Piquet et al. 2018) and tested

the relationship between body condition and proportional time spent perched using a

glmmADMB with ID as a random factor.

We performed all analyses in R version 3.4.1 (R Core Team 2017) and we made the figures

using "ggplot2" package version 3.0.0 (Wickham 2009). We set statistical significance to the

level of 0.05. Unless otherwise indicated, we denote averages as mean ± standard error (SE).

Results

Perch behavior general description

Overall, 99.3% of the squirrels (adults, subadults, and juveniles, $N = 139$) were observed

perching. In 31.9% of a total of 3134 scans just one individual perched, while 2 or over 2

squirrels perched in 21.1 and 19.8% of scans, respectively. On average 38.4 ± 0.6 % of a group

was perched per scan ($N = 522$ scans), but we found no relation between number of perched

squirrels and group size (LR test: $\Lambda = 0.04$, $P = 0.85$). Individuals perched both in the presence

and absence of predators (prediction 0 Table 1). Although, the percentage of individuals perched

per scan when a predator was observed during the observation period ($53.3 \pm 1.3\%$, $N = 417$

scans) was significantly higher to that when no external stimulus was observed ($48.9 \pm 0.6\%$, N

= 2192 scans; Mann-Whitney U test: $U = 503630$, $P < 0.001$, effect size r = 0.07; prediction 0

Table 1). Barbary ground squirrels were located at more elevated positions when perched ($N =$

2203) compared to locations where the squirrels performed any other behavior (prediction 1

Table 1; $N = 5289$; Mann-Whitney U test: $U = 3958800$, $Z = -22.01$, $P < 0.001$, r = 0.25; Fig. 1).

We found that the proportion of observed time that at least one individual in the group was

perched (p_{obs}) was smaller than the proportion of expected time (p_{exp}, predictions 2 and 3; Table

2). Furthermore, p_{obs}-p_{exp} was smaller than 0 (prediction 2 and 3 Table 1, Table 2).

Perch behavior and social conditions analyses

We found differences in time spent foraging (Kruskal-Wallis test: $H_2 = 8.18$, $P = 0.02$, post-

hoc adults-juveniles $P = 0.04$), low-quality vigilance (Kruskal-Wallis test: $H_2 = 6.64$, $P = 0.04$,

post hoc subadults-juveniles $P = 0.03$) and perch behavior (Kruskal-Wallis test: $H_2 = 52.6$, $P <$

0.001, post-hoc adults-juveniles and subadults-juveniles $P < 0.001$) among age classes

(prediction 4 Table 1, Fig. 2). Time spent perched did not differ when individuals performed

perch behavior in the absence ($33.5 \pm 3.6\%$, $N = 77$ focal follows) or presence of kin ($26.7 \pm$

2.4%, $N = 114$ focal follows; LR test: $\Lambda = 2.93$, $P = 0.09$; GLMM: 0.35 ± 0.20, random factor:

ID = 0.25; prediction 5 Table 1). We found neither any correlation between perch behavior and

relatedness (Spearman's rank correlation: $r_s = -0.05$, $P = 0.49$). Adult females ($N = 86$) spent a

similar amount of time perched when vulnerable juveniles were absent ($29.3 \pm 3.3\%$, $N = 90$

focal follows) or present ($30.7 \pm 2.7\%$, $N = 80$ focal follows; LR test: $\Lambda = 0.39$, $P = 0.53$;

GLMM: 0.12 ± 0.20, random factor: ID = 0.16; prediction 6 Table 1).

Perch behavior and survival analyses

In our 2016 field season, 50 of our focal animals from 2015 survived and 32 disappeared.

We found that individuals were closer to cover when perched (0.58 ± 0.09 m) than when

foraging (3.20 ± 0.32 m; paired Wilcoxon signed-rank test: $V = 41$, $N = 94$, $P < 0.001$; prediction

7 Table 1). Individuals that were alone or in groups spent similar amounts of time perched (37.3

$\pm 5.2\%$, $N = 38$, and $29.5 \pm 2.0\%$, $N = 193$, respectively; LR test: $\Lambda = 0.61$, $P = 0.44$; GLMM: -

0.19 ± 0.24, random factor: ID = 0.23; prediction 7 Table 1). We found no difference in survival,

last Julian day observed, and age at disappearance with the proportion of time spent perched ($N =$

82 individuals, Mann-Whitney U test: $U = 803.5$, $P = 0.98$; Spearman's rank correlation: $r_s=$

0.04, $P = 0.73$; Kruskal-Wallis test: $H_5 = 1.46$, $P = 0.92$; respectively; prediction 8 Table 1).

Foraging was negatively correlated to time spent perched (prediction 8 Table 1, Fig. 3), and

perch behavior was not affected by body condition (LR test: $\Lambda = 0.24$, $P = 0.62$; GLMM: $0.54 \pm$

1.10, random factor: ID = 0.09).

Discussion

Barbary ground squirrels met two criteria of a sentinel system (van der Marel et al. 2019),

but coordination of perch bouts was unknown. Here, we examined the importance of

quantitatively testing coordination in species with a potential sentinel system. Barbary ground

squirrels' perch behavior was coordinated; however, individuals did not take turns acting as

sentinels as they never decreased the time between different perch bouts in a group. Instead,

perch bouts were synchronized because the observed time without a perched individual was

smaller than the expected time resulting in a difference in proportions smaller than zero (Table

2). We detected no relationship between the numbers of perched individuals and group size.

Furthermore, perch behavior cannot be considered induced vigilance (Blanchard and Fritz 2007)

because perch behavior was performed in the absence and presence of an external stimulus (a

predator). Finally, we detected no apparent costs or benefits of perch behavior at the individual

or group level. Thus, our result may impact the number of species with a true sentinel system

because coordination of sentinel bouts has not been quantitatively tested for 37 out of 40 species

with a described sentinel system (Bednekoff 2015). When taking turns in a potential sentinel

system is tested, coordination may be absent altogether (Hing et al. 2019) or synchronized, as in

our study.

Since it seems counterintuitive to be on the lookout at the same time as your group

members, the question is why group members synchronize high-quality vigilance bouts? First,

Barbary ground squirrels perch behavior may be an individual's optimal activity, as is seen in

suricates, *Suricata suricatta*, or Arabian babblers, *Argya squamiceps*, where individuals become

sentinels when satiated (Clutton-Brock 1999; Wright, Maklakov, et al. 2001). We did not find a

relationship between perch behavior and body condition, suggesting that Barbary ground

squirrels on Fuerteventura may be able to afford a reduction in foraging behavior perhaps due to

the available resources or the release of enemies (parasites and predators) on the island compared

to the native range (López-Darias 2007; López-Darias and Nogales 2008; Piquet et al. 2018).

Normally, herbivores spend a great proportion of the time foraging to meet daily energy

requirements (Fortin et al. 2004) and often rarely stop chewing (McDougall and Ruckstuhl

2018). But the mainly herbivorous Barbary ground squirrels spend a surprisingly small

proportion of time foraging (12.8 ± 1.1%, van der Marel et al. 2019) in comparison to other

species (52.0 ± 3.0% in Cape ground squirrels, Waterman, 1995; Unck et al. 2009; 87% in

eastern grey kangaroos, Gélin et al. 2013; 64% in chestnut-crowned babblers, *Pomatostomus*

ruficeps, Sorato et al. 2012; and 40% in Florida scrub-jays, *Aphelocoma coerulescens*, Fleischer

Jr et al. 2003). Hence, satiated squirrels may perform perch behavior as an optimal activity and

may imply that the trade-off between vigilance and feeding might be absent in the invasive

population of Barbary ground squirrels.

A second mechanism to explain synchronized vigilance is the structure of the habitat

because the presence of rock walls prevents the squirrels from looking on either side of the wall

when on the ground. Visual cues are essential for predator detection in diurnal species and, thus,

for ground-dwelling sciurids, whereby predator detection is easier in an open-structured habitat

(Ylönen and Brown 2007), which in turn can be considered a constraint on sociality. Hence, to

adequately detect predators, two Barbary ground squirrels may be necessary to look in both

directions of the rock wall. Therefore, perch behavior may simply be a form of extremely

high-quality vigilance where individuals improve their level of detection by synchronizing

vigilance bouts. In the stingless bee, *Tetragonisca angustula,* guards are present to defend the

nest entrance and these guards are evenly distributed on either side of the entrance to increase the

view of their surroundings (Shackleton et al. 2018). Therefore, similar to synchronized vigilance

in bees, squirrels may be vigilant at the same time on top of rock walls to watch 'each other's

back'. As such, vigilance could mean that at least two or more individuals have to be on the

lookout to watch at either side of the rock walls, instead of being a synchronized function as a

form of collective sentinel behavior. The possibility of collective sentinel behavior requires

further attention. Thus, the structure of the habitat (plains interspersed with ravines and rock

walls), and that they are group-living, could influence the synchronization of vigilance in the

invasive population of Barbary ground squirrels.

As we used an observational study in the field, overall vigilance levels, distance between

neighbors, and group size could have affected levels of synchronization. For example, when

vigilance levels are high in a group, there is a greater chance that multiple individuals are

vigilant at the same time, whereas for groups where vigilance is overall low, there is less chance

of detecting synchronization (Beauchamp 2015). Furthermore, monitoring of neighbors may

prove too difficult when the distance between neighbors is too large (Pays et al. 2009), or when

group sizes are too large (Beauchamp 2015), which may skew our results to synchronization

instead of taking turns.

Although social conditions, including sex, kinship, age, or being a parent, may influence

perch behavior, male and female Barbary ground squirrels spent the same amount of time

perched (van der Marel et al. 2019). This lack of sex-biased benefit on perching suggests that

perch behavior does not serve a function in intrasexual competition (Walker et al. 2016) or mate

protection (Horrocks and Hunte 1986). Group members of species with a sentinel system are

often closely-related (Fernández et al. 2003); however, the role of kin selection is not a

prerequisite to the evolution of such a system (Bednekoff and Woolfenden 2003). We found that

kin selection was not a mechanism explaining coordinated vigilance in Barbary ground squirrel –

with varying relatedness levels within groups due to promiscuity– because individuals were

perched independently of whether close kin was in their visible range or not. Adults and

subadults performed more perch behavior than juveniles, but juveniles spent more time foraging

than adults or subadults, which is similar to other bird and mammalian species (Rasa 1977;

Ferguson 1987). In contrast to other species, e.g., common eiders, *Somateria mollissima* (Öst and

Tierala 2011), we did not find that perch behavior increased for mothers when vulnerable young

were present, refuting the notion that perch behavior in Barbary ground squirrels was a form of

parental care. In species where mothers show maternal care, such as ungulates, mothers provide

extra vigilance so that their offspring can increase time spent feeding (Pays et al. 2012), which

could explain our finding that juveniles do not spend perched a similar amount of time as adults.

To summarise, environmental conditions may affect perch behavior in the Barbary ground

squirrels on Fuerteventura more than social conditions.

Our results also suggest that perch behavior did not impose survival costs or benefits to

the Barbary ground squirrel population on Fuerteventura because 1) distance to cover was

shorter for perched individuals than foragers, and, 2) we found no relation between the

proportion of time spent perched on survival. However, we were unable to measure predation

risk directly for perched individuals compared to foragers and we may not have had enough

variance in the time spent perched to detect an effect on survival. Individuals were as likely to

perform perch behavior alone as in groups, suggesting a selfish benefit to the perched individual,

such as increased detection of predators (Bednekoff 2001; Bednekoff and Woolfenden 2003).

The absence of an effect on survival could be explained by two mechanisms: 1) all individuals

that were observed in at least five scans in the population perform perch behavior, and, 2) there is

an ecological release of predators and parasites in the invasive range of Barbary ground squirrels

(López-Darias 2007; López-Darias and Nogales 2008; Piquet et al. 2018). In summary, we found

no apparent cost or benefit of perch behavior in the invasive population of the Barbary ground

squirrel.

Conclusions

Our results demonstrated the importance of testing for coordination in species with a

suggested sentinel system and showed that the invasive population of Barbary ground squirrels

synchronize perch behavior. Two non-mutually exclusive hypotheses explaining synchronization

of perch bouts are that: 1) squirrels are satiated and have the time and energy to perform

synchronized perch behavior (low cost), and, 2) the benefits are large in terms of evading

ambush predators and scanning effectiveness (watch each other's back). Future studies could test

for other possible hypotheses explaining synchrony of perch bouts, including contagion-of-fear,

induced synchrony, and mimicking of nearest neighbor hypotheses (Sirot and Touzalin 2009;

Beauchamp 2015). This study shows that quantitative testing of coordination is important in

species with a described sentinel system and that the cost of the trade-off between vigilance and

foraging may not be high for an invasive species.

Acknowledgements

We acknowledge the landowners of our study sites, the IPNA-CSIC by providing a research

vehicle, and the Cabildo of Fuerteventura for the use of the facilities and the help of its staff at

the Estación Biológica de La Oliva. AM was supported by a University of Manitoba Faculty of

Science graduate studentship and a Faculty of Graduate Studies Graduate Enhancement of the

Tri-Council Stipend (GETS). MLD was funded by the Program Agustín de Betancourt of the

Cabildo Insular de Tenerife under the identification mark "Tenerife 2030" (P. INNOVA 2016-

2021). JMW was funded by the National Research Council of Canada (#04362), the Canadian

Foundation for Innovation, and the University of Manitoba University Research Grant Program.

All methods in this study were approved by the University of Manitoba animal ethics committee

(protocol #F14-032), were permitted by the government of Fuerteventura (Cabildo Insular de

Fuerteventura #14885) and adhered to the ABS/ASAB guidelines for ethical animal research.

References

Altmann J. 1974. Observational study of behavior: Sampling methods. Behaviour. 49:227–267.

del Arco Aguilar M-J, González-González R, Garzón-Machado V, Pizarro-Hernández B. 2010.

Actual and potential natural vegetation on the Canary Islands and its conservation status.

Biodivers Conserv. 19(11):3089–3140. doi:10.1007/s10531-010-9881-2.

Beauchamp G. 2015. Synchronization and coordination of animal vigilance. In: Animal

Vigilance: Monitoring Predators and Competitors. Elsevier Science. p. 155–172.

Beauchamp G, Alexander P, Jovani R. 2012. Consistent waves of collective vigilance in groups

using public information about predation risk. Behav Ecol. 23(2):368–374.

doi:10.1093/beheco/arr194.

Bednekoff PA. 1997. Mutualism among safe, selfish sentinels: a dynamic game. Am Nat.

150(3):373–392.

Bednekoff PA. 2001. Coordination of safe, selfish sentinels based on mutual benefits. Ann Zool

Fenn. 38:5–14.

Bednekoff PA. 2015. Sentinel behavior: a review and prospectus. In: Advances in the study of

behavior. p. 115–145.

Bednekoff PA, Lima SL. 1998. Randomness, chaos and confusion in the study of antipredator

vigilance. Trends Ecol Evol. 13(7):284–7. http://www.ncbi.nlm.nih.gov/pubmed/21238305.

Bednekoff PA, Woolfenden GE. 2003. Florida scrub-jays (*Aphelocoma coerulescens*) are

sentinels more when well-fed (even with no kin nearby). Ethology. 109(11):895–903.

doi:10.1046/j.0179-1613.2003.00926.x.

Black JM, Owen M. 1989. Parent-offspring relationships in wintering barnacle geese. Anim

Behav. 37:187–198. doi:10.1016/0003-3472(89)90109-7.

Blanchard P, Fritz H. 2007. Induced or routine vigilance while foraging. Oikos. 116(10):1603–

1608. doi:10.1111/j.2007.0030-1299.15799.x.

Bolker B, Skaug H, Magnusson A, Nielsen A. 2012. Getting started with the glmmADMB

package. R Packag version 20–8.:12.

Brandl SJ, Bellwood DR. 2015. Coordinated vigilance provides evidence for direct reciprocity in

coral reef fishes. Sci Rep. 5:14556. doi:10.1038/srep14556.

Clutton-Brock TH. 1999. Selfish sentinels in cooperative mammals. Science. 284:1640–1644.

doi:10.1126/science.284.5420.1640.

Ebensperger LA, Hurtado MJ, Ramos-Jiliberto R. 2006. Vigilance and collective detection of

predators in degus (*Octodon degus*). Ethology. 112(9):879–887. doi:10.1111/j.1439-

0310.2006.01242.x.

Edwards S, Waterman JM. 2011. Vigilance and grouping in the southern African ground squirrel

(*Xerus inauris*). Afr J Ecol. 49(3):286–291. doi:10.1111/j.1365-2028.2011.01262.x.

Evans MHR, Lihou KL, Rands SA. 2018. Black-headed gulls synchronise their activity with

their nearest neighbours. Sci Rep. 8:9978. doi:10.1038/s41598-018-28378-x.

Fairbanks BM, Dobson FS. 2010. Kinship does not affect vigilance in Columbian ground

squirrels (*Urocitellus columbianus*). Can J Zool. 88:266–270. doi:10.1139/Z10-001.

Favreau F-R, Pays O, Fritz H, Goulard M, Best EC, Goldizen AW. 2015. Predators, food and

social context shape the types of vigilance exhibited by kangaroos. Anim Behav. 99:109–121.

doi:10.1016/j.anbehav.2014.11.001.

Ferguson JWH. 1987. Vigilance behaviour in white-browed sparrow-weavers *Plocepasser*

mahali. Ethology. 76(3):223–235. doi:10.1111/j.1439-0310.1987.tb00685.x.

Fernández GJ, Capurro AF, Reboreda JC. 2003. Effect of group size on individual and collective

vigilance in greater rheas. Ethology. 109(5):413–425. doi:10.1046/j.1439-0310.2003.00887.x.

Fleischer Jr AL, Bowman R, Woolfenden GE. 2003. Variation in foraging behavior, diet, and

time of breeding of Florida scrub-jays in suburban and wildland habitats. Condor. 105:515–527.

Fortin D, Boyce MS, Merrill EH. 2004. Multi-tasking by mammalian herbivores: Overlapping

processes during foraging. Ecology. 85(8):2312–2322. doi:10.1890/03-0485.

Fournier DA, Skaug HJ, Ancheta J, Ianelli J, Magnusson A, Maunder MN, Nielsen A, Sibert J.

2012. AD Model Builder: Using automatic differentiation for statistical inference of highly

parameterized complex nonlinear models. Optim Methods Softw. 27(2):233–249.

doi:10.1080/10556788.2011.597854.

Gangoso L, Donázar JA, Scholz S, Palacios CJ, Hiraldo F. 2006. Contradiction in conservation

of island ecosystems: plants, introduced herbivores and avian scavengers in the Canary Islands.

Biodivers Conserv. 15:2231–2248. doi:10.1007/s10531-004-7181-4.

Gélin U, Wilson ME, Coulson GM, Festa-Bianchet M. 2013. Offspring sex, current and previous

reproduction affect feeding behaviour in wild eastern grey kangaroos. Anim Behav. 86:885–891.

doi:10.1016/j.anbehav.2013.08.016.

Griesser M. 2003. Nepotistic vigilance behavior in Siberian jay parents. Behav Ecol. 14(2):246–

250. doi:10.1093/beheco/14.2.246.

Hing JS, Schmaltz G, Quinn JS. 2019. Testing for sentinel coordination in Smooth-billed Anis

(*Crotophaga ani*). J Ornithol. 160:165–172. doi:10.1007/s10336-018-1584-6.

Horrocks JA, Hunte W. 1986. Sentinel behaviour in vervet monkeys: Who sees whom first?

Anim Behav. 34(5):1566–1568. doi:10.1016/S0003-3472(86)80226-3.

Labocha MK, Schutz H, Hayes JP. 2014. Which body condition index is best? Oikos. 123:111–

119. doi:10.1111/j.1600-0706.2013.00755.x.

Li Chunlin, Jiang Z, Li L, Li Z, Fang H, Li Chunwang, Beauchamp G. 2012. Effects of

reproductive status, social rank, sex and group size on vigilance patterns in Przewalski's gazelle.

PLoS One. 7(2):e32607. doi:10.1371/journal.pone.0032607.

López-Darias M. 2007. Ecology of invasions: the case of the Barbary ground squirrel

(*Atlantoxerus getulus*) in Fuerteventura Island (Canary Islands). PhD thesis, University of La

Laguna. San Cristóbal de La Laguna, Spain.

López-Darias M, Lobo JM. 2008. Factors affecting invasive species abundance: the Barbary

Ground Squirrel on Fuerteventura Island, Spain. Zool Stud. 47(3):268–281.

López-Darias M, Nogales M. 2008. Effects of the invasive Barbary ground squirrel (*Atlantoxerus*

getulus) on seed dispersal systems of insular xeric environments. J Arid Environ. 72:926–939.

doi:10.1016/j.jaridenv.2007.12.006.

Lung MA, Childress MJ. 2007. The influence of conspecifics and predation risk on the vigilance

of elk (*Cervus elaphus*) in Yellowstone National Park. Behav Ecol. 18:12–20.

doi:10.1093/beheco/arl066.

Machado A. 1979. The introduction of the Getulian squirrel (*Atlantoxerus getulus L.*, 1758) in

Fuerteventura, Canary Islands. Egypt J Wildl Nat Resour. 2:182–202.

Manser MB. 1999. Response of foraging group members to sentinel calls in suricates, *Suricata*

suricatta. Proc R Soc B Biol Sci. 266:1013–1019. doi:10.1098/rspb.1999.0737.

van der Marel A. 2019. Life history traits, social organisation and the drivers of sociality in an

invasive ground squirrel. PhD thesis, University of Manitoba. Winnipeg, Canada.

http://hdl.handle.net/1993/34401.

van der Marel A, López-Darias M, Waterman JM. 2019. Group-enhanced predator detection and

quality of vigilance in a social ground squirrel. Anim Behav. 151:43–52.

doi:10.1016/j.anbehav.2019.02.017.

van der Marel A, Waterman JM, López-Darias M. 2020. Social organization in a North African

ground squirrel. J Mammal. XX:1–14. doi:10.1093/jmammal/gyaa031.

McDonald M, Johnson S. 2014. "There's an app for that": a new program for the collection of

behavioural field data. Anim Behav. 95:81–87. doi:10.1016/j.anbehav.2014.06.009.

McDougall PL, Ruckstuhl KE. 2018. Vigilance behaviour is more contagious when chewing

stops: examining the characteristics of contagious vigilance in bighorn sheep. Behav Ecol

Sociobiol. 72:143. doi:10.1007/s00265-018-2536-0.

McGowan K, Woolfenden G. 1989. A sentinel system in the Florida scrub jay. Anim Behav.

37:1000–1006.

Medina FM, López-Darias M, Nogales M, García R. 2008. Food habits of feral cats (*Felis*

silvestris catus L.) in insular semiarid environments (Fuerteventura, Canary Islands). Wildl Res.

35(2):162–169. doi:10.1071/WR07108.

Newbold T, Collins S, Behnke J, Eales J, El-Geznawy A, El-Tohamy T, Ezzat R, Farag D,

Gilbert F, Jobling S, et al. 2008. Sentinel behaviour and the watchman's call in the Chukar at St

Katherine Protectorate, Sinai, Egypt. Egypt J Biol. 10:42–53.

Öst M, Tierala T. 2011. Synchronized vigilance while feeding in common eider brood-rearing

coalitions. Behav Ecol. 22(2):378–384. doi:10.1093/beheco/arq223.

Pays O, Dubot AL, Jarman PJ, Loisel P, Goldizen AW. 2009. Vigilance and its complex

synchrony in the red-necked pademelon, *Thylogale thetis*. Behav Ecol. 20(1):22–29.

doi:10.1093/beheco/arn110.

Pays O, Jarman PJ, Loisel P, Gerard JF. 2007. Coordination, independence or synchronization of

individual vigilance in the eastern grey kangaroo? Anim Behav. 73:595–604.

doi:10.1016/j.anbehav.2006.06.007.

Pays O, Renaud PC, Loisel P, Petit M, Gerard JF, Jarman PJ. 2007. Prey synchronize their

vigilant behaviour with other group members. Proc R Soc B Biol Sci. 274:1287–1291.

doi:10.1098/rspb.2006.0204.

Pays O, Sirot E, Fritz H. 2012. Collective vigilance in the greater kudu: Towards a better

understanding of synchronization patterns. Ethology. 118:1–9. doi:10.1111/j.1439-

0310.2011.01974.x.

Périquet S, Todd-Jones L, Valeix M, Stapelkamp B, Elliot N, Wijers M, Pays O, Fortin D,

Madzikanda H, Fritz H, et al. 2012. Influence of immediate predation risk by lions on the

vigilance of prey of different body size. Behav Ecol. 23(5):970–976. doi:10.1093/beheco/ars060.

Piquet JC, López-Darias M, van der Marel A, Nogales M, Waterman J. 2018. Unraveling

behavioral and pace-of-life syndromes in a reduced parasite and predation pressure context:

personality and survival of the Barbary ground squirrel. Behav Ecol Sociobiol. 72:147.

doi:10.1007/s00265-018-2549-8.

Podgórski T, de Jong S, Bubnicki JW, Kuijper DPJ, Churski M, Jędrzejewska B. 2016. Drivers of

synchronized vigilance in wild boar groups. Behav Ecol. 27(4):1097–1103.

doi:10.1093/beheco/arw016.

Quirici V, Palma M, Sobrero R, Faugeron S, Ebensperger LA. 2013. Relatedness does not predict

vigilance in a population of the social rodent *Octodon degus*. Acta Ethol. 16(1):1–8.

doi:10.1007/s10211-012-0131-3.

Rasa OAE. 1977. The ethology and sociology of the dwarf mongoose (*Helogale undulata*

rufula). Z Tierpsychol. 43:337–406. doi:10.1111/j.1439-0310.1977.tb00487.x.

Rasa OAE. 1987. Vigilance behaviour in dwarf mongooses: selfish or altruistic? Suid-Afrikaanse

Tydskr vir Wet. 83:587–590.

Reboreda JC, Fernandez GJ. 1997. Sexual, seasonal and group size differences in the allocation

of time between vigilance and feeding in the greater rhea, *Rhea americana*. Ethology. 103:198–

207. doi:10.1111/j.1439-0310.1997.tb00116.x.

Ridley AR, Nelson-Flower MJ, Thompson AM. 2013. Is sentinel behaviour safe? An

experimental investigation. Anim Behav. 85:137–142. doi:10.1016/j.anbehav.2012.10.017.

Santema P, Clutton-Brock T. 2013. Meerkat helpers increase sentinel behaviour and bipedal

vigilance in the presence of pups. Anim Behav. 85:655–661. doi:10.1016/j.anbehav.2012.12.029.

Schulte-Hostedde AI, Millar JS, Hickling GJ. 2001. Evaluating body condition in small

mammals. Can J Zool. 79:1021–1029. doi:10.1139/z01-073.

Shackleton K, Alves DA, Ratnieks FLW. 2018. Organization enhances collective vigilance in the

hovering guards of *Tetragonisca angustula* bees. Behav Ecol. 29(5):1105–1112.

doi:10.1093/beheco/ary086.

Sirot E, Touzalin F. 2009. Coordination and synchronization of vigilance in groups of prey: The

role of collective detection and predators' preference for stragglers. Am Nat. 173:47–59.

doi:10.1086/593358.

Sorato E, Gullett PR, Griffith SC, Russell AF. 2012. Effects of predation risk on foraging

behaviour and group size: adaptations in a social cooperative species. Anim Behav. 84:823–834.

doi:10.1016/j.anbehav.2012.07.003.

Stankowich T. 2003. Marginal predation methodologies and the importance of predator

preferences. Anim Behav. 66:589–599. doi:10.1006/anbe.2003.2232.

Unck CE, Waterman JM, Verburgt L, Bateman PW. 2009. Quantity versus quality: how does

level of predation threat affect Cape ground squirrel vigilance? Anim Behav. 78:625–632.

doi:10.1016/j.anbehav.2009.05.028.

Walker LA, York JE, Young AJ. 2016. Sexually selected sentinels? Evidence of a role for

intrasexual competition in sentinel behavior. Behav Ecol. 27(5):1461–1470.

doi:10.1093/beheco/arw064.

Waterman JM. 1995. The social organization of the Cape ground squirrel (*Xerus inauris*;

Rodentia: Sciuridae). Ethology. 101:130–147.

Waterman JM. 1998. Mating tactics of male Cape ground squirrels, *Xerus inauris*: consequences

of year-round breeding. Anim Behav. 56:459–466. doi:10.1006/anbe.1998.0780.

Whiteside MA, Langley EJG, Madden JR. 2016. Males and females differentially adjust

vigilance levels as group size increases: effect on optimal group size. Anim Behav. 118:11–18.

doi:10.1016/j.anbehav.2016.04.025.

Wright J, Berg E, de Kort SR, Khazin V, Maklakov AA. 2001. Safe selfish sentinels in a

cooperative bird. J Anim Ecol. 70:1070–1079.

Wright J, Maklakov AA, Khazin V. 2001. State-dependent sentinels: an experimental study in the

Arabian babbler. Proc R Soc London Ser B Biol Sci. 268:821–826. doi:10.1098/rspb.2000.1574.

Ylönen H, Brown JS. 2007. Fear and the foraging, breeding and sociality of rodents. In: Wolff

JO, Sherman PW, editors. Rodent societies: An ecological & evolutionary perspective. Chicago,

IL: The University of Chicago Press. p. 328–341.

Zeileis A. 2002. Diagnostic checking in regression relationships. R News. 2(3):7–10.

655
656

Figure legends

Figure 1. Boxplot of height (m) of Barbary ground squirrels when perching or performing any

other behavior. The dark line shows the median, the box edges the upper and lower quartiles, the

whiskers are 50% from the median, and the closed circles are the outliers, calculated as the

values smaller or larger than 1.5 times the box length (i.e., upper - lower quantile).

Figure 2. Mean $\pm$ SE of the proportion of time spent feeding, performing low-quality vigilance,

or perching in adult, subadult and juvenile Barbary ground squirrels. Significant differences are

denoted by asterisks (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$).

Figure 3. Relation between the proportion of time spent foraging and time spent perching

(Spearman's rank correlation: $r_s = -0.38$, $P < 0.001$) in Barbary ground squirrels.

Tables and table legends

Table 1. Hypotheses and predictions on perch behavior (vigilance without performing any other

behavior for over 30 s and at raised positions of at least 30 cm above the ground) in Barbary

ground squirrels on Fuerteventura, Canary Islands, Spain.

Hypothesis	Predictions	Method	Supported
Perch behavior			
Perch = induced vigilance	0) Individuals only perch in the presence of an external stimulus, e.g., predator	scan	no
Perch = sentinel behavior	1) perched from elevated positions	scan + ff	yes
	2) continuous turn-over without changing the number of perched individuals any more than expected by chance	all	no
Perch = synchronized behavior	3) either multiple individuals are perched or non-vigilant at the same time	all	yes
Perch behavior and social conditions			
Social conditions influence who are perched	4) adults and sub-adults spent more time perched than juveniles	scan	yes
	5) individuals only perch when close kin are nearby (kin-selected behavior)	ff	no
	6) adult females only perch when own vulnerable offspring have emerged (parental behavior)	ff	no
Perch behavior and survival			
Survival increases for perched individuals	7) perched individuals are closer to cover than individuals performing another behavior & individuals are perched when alone and in a group	ff	yes
Survival decreases for perched individuals	8) greater risk of predation when located at raised positions & when in worse body condition	scan + ff	no

674 [a] Behavioral observation method used to test predictions: 10 min scan sampling (scan), all-

675 occurrences during scan sampling (all), and separate 10 min focal follows (ff).

677   **Table 2.** The proportion of observed (p_{obs}) and expected (p_{exp}) time at least one individual was

678   perched (mean ± SE) per observation period in Barbary ground squirrels calculated as duration

679   of perch bouts.

Observation period	N	p_{obs}	p_{exp}	P_{obs}-P_{exp}	V	P
All individuals	125	0.19 ± 0.02	0.54 ± 0.03	-0.36 ± 0.03	496	<0.001
Adults	122	0.20 ± 0.02	0.52 ± 0.03	-0.33 ± 0.03	158	<0.001
Adult females	107	0.19 ± 0.02	0.52 ± 0.03	-0.35 ± 0.02	20	<0.001

681   Sample size (*N*) and the pairwise test results (*V* and *P*) are included.