Comprehensive evolution and molecular characteristics of a large number of

SARS-CoV-2 genomes revealed its epidemic trend and possible origins

Yunmeng Bai[1#], Dawei Jiang[1#], Jerome R Lon[1#], Xiaoshi Chen[1], Meiling Hu[1], Shudai Lin[1], Zixi Chen[1],

Xiaoning Wang[1,2], Yuhuan Meng[3*], Hongli Du[1*]

[1]*School of Biology and Biological Engineering, South China University of Technology, Guangzhou*

510006, China

[2]*State Clinic Center of Gearitic, Chinese PLA General Hospital, Beijing 100853, China*

[3]*Guangzhou KingMed Transformative Medicine Institute Co., Ltd, Guangzhou 510330, China*

[#] Equal contribution.

[*] Corresponding authors.

E-mail: hldu@scut.edu.cn, zb-mengyuhuan@kingmed.com.cn

Abstract

Objectives:

To reveal epidemic trend and possible origins of SARS-CoV-2 by exploring its evolution and

molecular characteristics based on a large number of genomes since it has infected millions of

people and spread quickly all over the world.

Methods:

Various evolution analysis methods were employed.

Results:

The estimated Ka/Ks ratio of SARS-CoV-2 is 1.008 or 1.094 based on 622 or 3624 SARS-CoV-2

genomes, and the time to the most recent common ancestor (tMRCA) was inferred in late

September 2019. Further 9 key specific sites of highly linkage and four major haplotypes H1, H2,

H3 and H4 were found. The Ka/Ks, detected population size and development trends of each

major haplotype showed H3 and H4 subgroups were going through a purify evolution and almost

disappeared after detection, indicating H3 and H4 might have existed for a long time, while H1

and H2 subgroups were going through a near neutral or neutral evolution and globally increased

with time. Notably the frequency of H1 was generally high in Europe and correlated to death rate

(r>0.37).

Conclusions:

In this study, the evolution and molecular characteristics of more than 16000 genomic sequences

provided a new perspective for revealing epidemiology of SARS-CoV-2.

KEYWORDS: SARS-CoV-2; evolution; classification; haplotype

Introduction

The global outbreak of SARS-CoV-2 is currently and increasingly recognized as a serious, public

health concern worldwide. Coronaviruses exist widely around the world, and to date, a total of seven

types of coronaviruses that can infect humans have been found, of which four coronaviruses including

hCoV-229E, hCoV-NL63, hCoV-OC43 and hCoV-HKU1 cause a cold, while the other three including

SARS-CoV, MERS-CoV and SARS-CoV-2 usually cause mild to severe respiratory diseases. The total

number of SARS-CoV and MERS-CoV infections are only 8,069 and 2,494 with reproduction number

(R0) fluctuates 2.5-3.9 and 0.3-0.8, respectively. However, SARS-CoV-2 has infected 2471136 people

in 212 countries up to April 22th(WHO, 2020), with the basic R0 ranging from 1.4 to 6.49(Liu, et al.,

2020). Among these three typical coronavirus, MERS-CoV has the highest death rate of 34.40%,

SARS-CoV has the modest death rate of 9.59%, SARS-CoV-2 is about 6.99% and 7.69% death rate of

the global and Wuhan, but is quite high death rate in some countries of Europe, such as Belgium, Italy,

United Kingdom, Netherlands, Spain and France, which even reaches 14.95%, 13.39%, 13.48%,

11.61%, 10.42% and 13.60% respectively according to the data of April 22th, 2020. Except for the

shortage of medical supplies, aging and other factors, it is not clear whether there is virus mutation

effect in these countries with such a significantly increased death rate.

It has been reported that SARS-CoV-2 belongs to beta-coronavirus and is mainly transmitted by the

respiratory tract, which belongs to the same subgenus (SarbeCoVirus) as SARS-CoV(Lu, et al., 2020).

Through analyzing the genome and structure of SARS-CoV-2, its receptor-binding domain (RBD) was

found to bind with angiotensin-converting enzyme 2 (ACE2), which is also one of the receptors for

binding SARS-CoV(Wrapp, et al., 2020). Some early genomic studies have shown that SARS-CoV-2 is

similar to certain bat viruses (RaTG13, with the whole genome homology of 96.2%(Zhou, et al., 2020))

and Malayan pangolins coronaviruses (GD/P1L and GDP2S, with the whole genome homology of

92.4%(Lam, et al., 2020)). They have speculated several possible origins of SARS-CoV-2 based on its

spike protein characteristics (cleavage sites or the RBD)(Lam, et al., 2020, Zhang and Holmes, 2020,

Zhou, et al., 2020), in particular, SARS-CoV-2 exhibits 97.4% amino acid similarity to the Guangdong

pangolin coronaviruses in RBD, even though it is most closely related to bat coronavirus RaTG13 at

the whole genome level. However, it is not enough to present genome-wide evolution by a single gene

or local evolution of RBD, whether bats or pangolins play an important role in the zoonotic origin of

SARS-CoV-2 remains uncertain(Andersen, et al., 2020). Furthermore, since there is little molecular

characteristics analysis of SARS-CoV-2 based on a large number of genomes, it is difficult to

determine whether the virus has significant variation that affects its phenotype. Thereby, it is necessary

to further reveal the phylogenetic evolution and molecular characteristics of the whole genome of

SARS-CoV-2 in order to develop a comprehensive understanding of the virus and provide a basis for

the prevention and treatment of SARS-CoV-2.

Results

Genome sequences

We got a total of 1053 genomic sequences up to March 22th, 2020. According to the filter criteria, 37

sequences with ambiguous time, 314 with low quality and 78 with similarity of 100% were removed. A

total of 624 sequences were obtained to perform multiple sequences alignment. Two highly divergent

sequences (EPI_ ISL_414690, EPI_ISL_415710) according to the firstly constructed phylogenetic tree

were also filtered out (Table S1). The remaining 622 sequences were used to reconstruct a phylogenetic

tree. In addition, total of 3624 and 16373 genomic sequences were redownloaded up to April 6th, 2020

and May 10th, 2020 respectively for further exploring the evolution and molecular characteristics of

SARS-CoV-2.

Estimate of evolution rate and the time to the most recent common ancestor for SARS-CoV,

MERS-CoV, and SARS-CoV-2

In this study, date for SARS-CoV-2 ranged from 2019/12/26 to 2020/03/18 was collected. The average

Ka/Ks of all the coding sequences was closer to 1 (1.008), which indicating the genome was going

through a neutral evolution. We also reevaluated the Ka/Ks of SARS-CoV and MERS-CoV through the

whole period, and found the ratio were smaller than SARS-CoV-2 (Table 1). To estimate more credible

Ka/Ks for SARS-CoV-2, we recalculated it using redownloaded 3624 genome sequences ranged from

2019/12/26 to 2020/04/06, and the average Ka/Ks of it was 1.094 (Table 1), which was almost same

with the above result.

We assessed the temporal signal using TempEst v1.5.3(Rambaut, et al., 2016). All three datasets

exhibit a positive correlation between root-to-tip divergence and sample collecting time (Figure S1),

hence they are suitable for molecular clock analysis in BEAST(Bouckaert, et al., 2019, Rambaut, et al.,

2016). The substitution rate of SARS-CoV-2 genome was estimated to be 1.601×10^{-3} (95% CI: 1.418 x

10^{-3} - 1.796×10^{-3}, Table 2, Figure S2A) substitution/site/year, which is as in the same order of

magnitude as SARS-CoV and MERS-CoV. The tMRCA was inferred on the late September, 2019 (95%

CI: 2019/08/28- 2019/10/26,Table 2, Figure S2B), about 2 months before the early cases of

SARS-CoV-2(Huang, et al., 2020).

Phylogenetic tree and clusters of SARS-CoV-2

The no-root phylogenetic trees constructed by the maximum likelihood method with PhyML

3.1(Guindon, et al., 2010) and MEGA(Kumar, et al., 2018) were showed in Figure 1, Figure S3A and

3B. According to the shape of phylogenetic trees, we divided 622 sequences into three clusters: Cluster

1 including 76 sequences mainly from North America, Cluster 2 including 367 sequences from all

regions of the world, and Cluster 3 including 179 sequences mainly from Europe (Table S2).

The specific sites of each Cluster

The *Fst* and population frequency of a total of 9 sites (NC_045512.2 as reference genome) were

detected (Table 3, Table S3). Thereinto, three (C17747T, A17858G and C18060T) are the specific sites

of Cluster 1, and four (C241T, C3037T, C14408T and A23403G) are the specific sites of Cluster 3.

Notably, C241T was located in the 5'-UTR region and the others were located in coding regions (6 in

ofr1ab gene, 1 in *S* gene and 1 in *ORF8* gene). Five of them were missense variant, including C14408T,

C17747T and A17858G in *ofr1ab* gene, A23403G in *S* gene, and T28144C in *ORF8* gene. The PCA

results showed that these 9 specific sites could clearly separate the three Clusters, while all SNV

dataset could not clearly separate Cluster 1 and Cluster 2 (Figure S4), which further suggested that

these 9 specific sites are the key sites for separating the three Clusters.

Linkage of specific sites

We found the 9 specific sites are highly linkage based on 622 genome sequences (Figure 2A), then we

carried out a further linkage analysis using the 3624 genome sequences (Figure 2B). As a result, for the

3624 genome sequences, 3 specific sites in the Cluster 1 were almost complete linkage, and haplotype

CAC and TGT accounted for 98.65% of all the 3 site haplotypes. The same phenomenon was also

found in 4 specific sites of Cluster 3, and haplotype CCCA and TTTG accounted for 97.68% of all the

4 site haplotypes. Intriguingly, the 9 specific sites were still highly linkage, and four haplotypes,

including TTCTCACGT (H1), CCCCCACAT (H2), CCTCTGTAC (H3) and CCTCCACAC (H4),

accounted for 95.89% of all the 9 site haplotypes. Thereinto, H1 and H3 had completely different bases

at the 9 specific sites. The frequencies of each site and major haplotype for each country were showed

in Figure 3 and Table S4. The data showed that the haplotype TTTG of the 4 specific sites in Cluster 3

had existed globally at present, and still exhibited high frequencies in most European countries but

quite low in Asian countries. While the haplotype TGT of the 3 specific sites in Cluster 1 existed

almost only in North America and Australia. For the 9 specific sites, most countries had only two or

three major haplotypes, except America and Australia had all of four major haplotypes and with

relative higher frequencies.

Characteristics and epidemic trends of major haplotype subgroups

All haplotypes of 9 specific sites for 3624 or 16373 genomes and the numbers of them were shown in

Table S5. Four major haplotypes H1, H2, H3 and H4, three minor haplotypes H5, H7 and H8 close to

H1 and one minor haplotype H6 close to H3 were found in both datasets. The numbers of these

haplotypes for 16373 genomes with clear collection data detected in each country in chronological

order were shown in Figure 4A. From these results, H2 and H4 haplotype subgroups had existed for a

long time (2019-12-24 to 2020-04-28), the detected population size of H2 subgroup was far greater

than that of H4 subgroup, and the H3 haplotype subgroup almost disappeared after detection

(2020-02-18 to 2020-04-28), while H1 haplotype subgroup was globally increasing with time

(2020-02-18 to 2020-05-05), which indicated H1 subgroup had adapted to the human hosts, and was

under an adaptive growth period worldwide. However, due to the nonrandom sampling on early phase

(only patients having a recent travel to Wuhan were detected), some earlier cases of H3 may be lost, the

high proportion of H3 subgroup during February 18 to March 10 could confirm this point.

H3 and H4 subgroups had the lowest Ka/Ks ratio in 3624 or 16373 genomes among the 4 major

subgroups (Table 4), suggesting that H3 and H4 subgroups might be going through a purifying

evolution and have existed for a long time, while H2 and H1 subgroups might be going through a near

neutral or neutral evolution according to their Ka/Ks ratios (Table 4), which were consistent with the

above phenomenon that only H1 and H2 subgroups were expanding with time around the world. From

the whole genome mutations in each major haplotype subgroup (Figure 4B, Table S6), we found that

except the 9 specific sites, there was no common mutations with frequencies more than 0.05 in these

haplotype subgroups but between H2 and H4 subgroups.

Phylogenetic network of haplotype subgroups

Phylogenetic networks were inferred with 697 mutations called from 3624 genomes dataset. For these

datasets, the network structures of TCS and MSN were similar. The major haplotype subgroups H4 and

H2 were in the middle of the network, while H1 and H3 were in the end nodes of the network (Figure

5). According to the phylogenetic networks, we proposed four hypothesis: (1) the ancestral haplotypes

evolved in four directions to obtain H1, H2, H3 and H4 respectively or evolved in two or more

directions to obtain two or more major haplotypes and then involved into the other major haplotype(s);

(2) H2 or H4 evolved in two directions respectively and finally generated H1 and H3; (3) H1 evolved

in one direction to generate H2 and H4, and then evolved into H3; (4) H3 evolved in one direction to

generate H4 and H2, and then evolved into H1. Although we cannot exclude the first hypothesis based

on the present data, but if there are evolutionary relationships among the four major haplotypes,

according to the Ka/Ks, detected population size and development trends in chronological order of each

major haplotype subgroup, we speculate that the most likely evolution hypothesis is that H3 and H4 are

the earliest haplotypes, which is gradually eliminated with selection, while H2 is the transitional

haplotype in the evolution process, and H1 may be the haplotype to be finally fixed.

Correlation analysis of specific sites with death rate and infectivity

To explore the relationship between death rate and the 9 specific sites, we used Pearson method to

calculate the correlation coefficient between death rate and frequency of each specific site or major

haplotype in 17 countries with 3624 genomes at early stage. As a result, all r values of 241T, 3037T,

14408T, 23403G and haplotype TTTG and H1 were more than 0.4 (Figure 6, Table S4). We also

evaluated the correlation coefficient with 16373 genomes in 30 countries, the r values of haplotype

TTTG and H1 were still more than 0.37 (Table S4), which might be increased when the death rates of

some countries were further stabilized with time. These finding integrating their high frequencies in

most European countries indicated that the 4 sites and haplotype TTTG and H1 might be related to the

pathogenicity of SARS-CoV-2.

To explore the relationship between infectivity and the 9 specific sites, we used population size of

major haplotypes to deduce the possible specific sites related infectivity. We assumed that these major

haplotype in each country were subject to similar virus transmission and control patterns, while the

population sizes of H1 and H2 subgroups were far greater than those of H3 and H4 subgroups (Figure

4A, Table S4), then the common different specific sites of H1 and H2 subgroups with H3 and H4

subgroups, C8782T and T28144C, might be related to the infectivity of SARS-CoV-2, and the virus

with C8782 and T28144 might be more infectious than those virus with 8782T and 28144C.

Discussion

SARS-CoV-2 poses a great threat to the production, living and even survival of human beings(Guo, et

al., 2020). With the further outbreak of SARS-CoV-2 in the world, further understanding on the

evolution and molecular characteristics of SARS-CoV-2 based on a large number of genome sequences

will help us cope better with the challenges brought by SARS-CoV-2.

Exploring evolution rate, tMRCA and phylogenetic tree of SARS-CoV-2 would help us better

understand the virus(Yuen, et al., 2020). The average Ka/Ks for all the coding sequences of 622 and

3624 SARS-CoV-2 genomes was 1.008 and 1.094, which was higher than those of SARS-CoV and

MERS-CoV, indicating that the SARS-CoV-2 was going through a neutral evolution. In general,

SARS-CoV would show purifying selection pressures and the Ka/Ks would go down at the middle or

end of the epidemic(He, et al., 2004). Therefore, it seemed to be reasonable considering that

SARS-CoV-2 was under an exponential growth period around the world. Interestingly, we also found

that the subgroups of different haplotypes (H1, H2, H3 and H4) seemed to experience different

evolutionary patterns according to their Ka/Ks. The H3 haplotype subgroup disappeared soon after

detection (2020-02-18 to 2020-04-28, Figure 4A), while H1 haplotype subgroup was globally

increasing with time, these evolution and changes should be considered in developing therapeutic drugs

and vaccines. The tMRCA of SARS-CoV-2 was inferred on the late September, 2019 (95% CI:

2019/08/28- 2019/10/26), about 2 months before the early cases of SARS-CoV-2(Huang, et al., 2020).

We also estimated tMRCA of SARS-CoV and MERS-CoV with the same methods, both were about 3

months later than the corresponding tMRCAs estimated by previous studied(Cotten, et al., 2014, Zhao,

et al., 2004). A recent study used TreeDater method to estimate tMRCA for more than 7000

SARS-CoV-2 genomes and indicated the tMRCA of SARS-CoV-2 was around 6 October 2019 to 11

December 2019, which were about one and a half months later than ours and in broad agreement with

six previous studies all performed on no more than 120 early SARS-CoV-2 genomes with BEAST

method(van Dorp, et al., 2020). While we used the most common method BEAST to estimate tMRCA

of SARS-CoV-2 based on 622 genomes and combined with the comparison of tMRCA of SARS-CoV

and MERS-CoV, indicating that the estimated tMRCA in the present study are more reliable.

A recent study clustered 160 SARS-CoV-2 whole-genome sequences into A, B and C groups by a

phylogenetic network analysis by taking bat RaTG13 as root (Forster, et al., 2020). The cluster result

was similar to our study: both the samples in Cluster A and our Cluster 1 were mainly from the United

States; the samples in Cluster C and our Cluster 3 were mainly from European countries, while Cluster

B and our Cluster 2 were mainly from China and the other regions. It was interesting that the markers

C8782T and T28144C, which were also discovered by Yu et al(Yu, et al., 2020) , in Cluster B were

also found in our study, but the other markers in Cluster A (T29095C) and Cluster C (G26144T) were

not significantly in our study. That may be caused by different sample sizes and different constructing

methods of phylogenetic tree. Based on the base substitution model, the ML method avoids the

possible "long-branch attraction" problem in the maximum parsimony method and is faster than the

Bayesian method(Holder and Lewis, 2003), hence it could be used as a reliable method for

phylogenetic analysis. Some studies used the genome of bat SARS-like-CoV(Zhang, et al., 2020),

RaTG13(Zhang, et al., 2020) or MT019529 (https://bigd.big.ac.cn/ncov/tree) as the root of

phylogenetic tree. Unfortunately, there was no obvious evidence showing that SARS-CoV-2 was from

the bat coronavirus even the identity between SARS-CoV-2 and RaTG13 was up to 96.2%(Zhou, et al.,

2020). In our study, the tMRCA of SARS-CoV-2 was inferred on the late September 2019, which

indicated there might exist an earlier SARS-CoV-2 strain we didn't find. Then in the case of unclear

source of SARS-CoV-2 and high homology of its genomes (> 99.9% homology mostly), it may be

inappropriate to identify the evolutionary characteristics inside the genomes by taking bat

SARS-like-CoV, RaTG13 or MT019529 as root. Therefore, the maximum likelihood method was used

in current study to construct a no-root tree to obtain the reliable clusters with different characteristics.

Based on the no-root tree, we identified 9 specific sites of highly linkage that play a decisive role in the

classification of clusters successfully. Among the four major haplotypes, H1 and H3 were in Cluster 3

and Cluster 1, respectively, while there were two haplotypes H2 and H4 in Cluster 2 (Figure 1, Figure

S3B).

Among the 9 specific sites, 8 of them were located in coding regions (6 in *ofr1ab* gene, 1 in *S* gene

and 1 in *ORF8* gene), and 5 of them were missense variant, including C14408T, C17747T and

A17858G in *ofr1ab* gene, A23403G in *S* gene, and C28144T in *ORF8* gene. *Orf1ab* is composed of

two partially overlapping open reading frames (orf), orf1a and 1b. It is proteolytic cleaved into 16

non-structural proteins (nsp), including nsp1 (suppress antiviral host response), nsp9 (RNA/DNA

binding activity), nsp12 (RNA-dependent RNA polymerase), nsp13 (helicase) and others(Chan, et al.,

2020), indicating the vital role of it in transcription, replication, innate immune response and

virulence(Graham, et al., 2008). C14408T with high frequencies of T in European countries were

located at nsp12 region, indicating that this missense variant might influence the role of RNA

polymerase. Spike glycoprotein, the largest structural protein on the surface of coronaviruses,

comprises of S1 and S2 subunits which mediating binding the receptor on the host cell surface and

fusing membranes, respectively(Li, 2016). It has been reported that S protein of SARS-CoV-2 can bind

angiotensin-converting enzyme 2 (ACE2) with higher affinity than that of SARS(Wrapp, et al., 2020,

Zhou, et al., 2020). Whether the missense variant of A23403G in *S* gene, which with high frequencies

of G in European countries, will change the affinity between S protein and ACE2 remains to be further

investigated.

It seems to take a long time to finally fix mutations according to the mutation frequency of each

subgroup. For example, H2 and H4 subgroups, which have been detected for more than four months

from December 24, 2019 to May 5, 2020 (Figure 4A), have more mutations with higher frequencies,

but the highest mutation frequency is only 0.486 at the position of 11083 (Figure 4B, Table S6). From

these phenomena, it can be inferred that it takes a long time for the specific sites of each major

subgroup to be fixed, but it may be faster if the early population is small. In addition, there is also the

possibility that an ancestor strain evolved in four directions by obtaining the specific mutations directly

and produced the four current major haplotypes, so the evolution time for obtaining four major

haplotypes may be shorter, which seems to be consistent with the phenomenon that four major

haplotypes can be detected in two months (Figure 4A). However, if there exist evolution relationship

among the major haplotypes of SARS-CoV-2, it is difficult to complete the evolution among the four

major haplotypes within two months (2019-12-24 to 2020-02-18, Figure 4A) at the current evolution

rate of each major haplotype population (Table 4). Therefore, we speculate that the transformation

among the four major haplotypes may have been completed for a long time, which have not been

detected. What's interesting is that only United States and Australia among 29 countries had all of four

major haplotypes and with relative higher frequency (Figure 4A, Table S4), which indicated that the

two countries are the most likely places where the virus appeared earlier based on all the present data.

Previous studies showed that the death rate of SARS-CoV-2 is affected by many factors, such as

medical supplies, aging, life style and other factors(Liu, et al., 2020, Malavolta, et al., 2020). However,

our study found that the 4 specific sites (C241T, C3037T, C14408T and A23403G) in the Cluster 3

were almost complete linkage, and the frequency of haplotype TTTG was generally high in European

countries and correlated to death rate (r>0.37) both based on 3624 or 16373 SARS-CoV-2 genomes,

which provides a new perspective to the reasons of relatively high death rate in Europe, and these

specific sites should be considered in designing new vaccine and drug development of SARS-CoV-2.

Two possible specific sites C8782T and T28144C related to the infectivity of SARS-CoV-2 were also

deduced in the present study, which would provide some basis for SARS-CoV-2 epidemiology.

Conclusion and prospective

The Ka/Ks ratio of SARS-CoV-2 and tMRCA of SARS-CoV-2 (95% CI: 2019/08/28- 2019/10/26)

indicated that SARS-CoV-2 might have completed the selection pressure of cross-host evolution in the

early stage and be going through a neutral evolution at present. The 9 specific sites with highly linkage

were found to play a decisive role in the classification of clusters. Thereinto, 3 of them are the specific

sites of Cluster 1 mainly from North America and Australia, 4 of them are the specific sites of Cluster 3

mainly from Europe in the early stage, both of these 3 and 4 specific sites are almost complete linkage.

The frequencies of haplotype TTTG for the 4 specific sites and H1 for 9 specific sites were generally

high in European countries and correlated to death rate (r>0.37) based on 3624 or 16373 SARS-CoV-2

genomes, which suggested these haplotypes might relate to pathogenicity of SARS-CoV-2 and

provided a new perspective to the reasons of relatively high death rates in Europe. The relationship

between the haplotype TTTG or H1 and the pathogenicity of SARS-CoV-2 needs to be further verified

by clinical samples or virus virulence test experiment, and the 9 specific sites with highly linkage may

provide a starting point for the traceability research of SARS-CoV-2. Given that the different evolution

patterns of different haplotypes subgroups, we should consider these evolution and changes in the

development of therapeutic drugs and vaccines.

Materials and methods

Genome sequences

The complete genome sequences of SARS-CoV-2 were downloaded from China National Center for

Bioinformation (https://bigd.big.ac.cn/ncov/release_genome/) and GISAID (https://www.gisaid.org/)

up to March 22th, 2020. The sequences were filtered out according to the following criteria: (1)

sequences with ambiguous time; (2) sequences with low quality which contained the counts of

unknown bases > 15 and degenerate bases > 50 (https://bigd.big.ac.cn/ncov/release_genome); (3)

sequences with similarity of 100% were removed to unique one. Finally, 624 high quality genomes

with precise collection time were selected and aligned using MAFFT v7 with automatic parameters.

Besides, the genome sequences of 7 SARS-CoV and 475 MERS-CoV were also downloaded from

NCBI (https://www.ncbi.nlm.nih.gov/), and the MERS-CoV dataset including samples collected from

both human and camel. In addition, for further exploring evolution and molecular characteristics of

SARS-CoV-2 based on the larger amount of genomic data, we redownloaded validation datasets of the

genome sequences from GISAID up to April 6th, 2020 and May 10th, 2020 respectively.

312   **Estimate of evolution rate and the time to the most recent common ancestor for SARS-CoV,**

313   **MERS-CoV, and SARS-CoV-2**

314   The average Ka, Ks and Ka/Ks for all coding sequences were calculated using KaKs_Calculator

315   v1.2(Zhang, et al., 2006), and the substitution rate and tMRCA were estimated using BEAST

316   v2.6.2(Bouckaert, et al., 2019). The temporal signal with root-to-tip divergence was visualized in

317   TempEst v1.5.3(Rambaut, et al., 2016) using a ML whole genome tree with bootstrap value as input.

318   For SARS-CoV and SARS-CoV-2, we selected a strict molecular clock and Coalescent Exponential

319   Population Model. For MERS-CoV, we selected a relaxed molecular clock and Birth Death Skyline

320   Serial Cond Root Model. We used the tip dates and chose the HKY as the site substitution model in all

321   these analyses. Markov Chain Monte Carlo (MCMC) chain length was set to 10,000,000 steps

322   sampling after every 1000 steps. The output was examined in Tracer v1.6

323   (http://tree.bio.ed.ac.uk/software/tracer/).

324   **Variants calling of SARS-CoV-2 genome sequences**

325   Each genome sequence was aligned to the reference genome (NC_045512.2) using bowtie2 with

326   default parameters(Langmead and Salzberg, 2012), and variants were called by samtools (sort; mpileup

327   -gf) and bcftoots (call -vm). The merge VCF files were created by bgzip and bcftools (merge

328   --missing-to-ref)(Li, 2011, Li, et al., 2009).

329   **Phylogenetic tree construction and virus isolates clustering for SARS-CoV-2**

330   After alignment of 624 SARS-CoV-2 genomes with high quality and manually deleted 2 highly

331   divergent genomes (EPI_ISL_415710, EPI_ISL_414690) according to the firstly constructed

332   phylogenetic tree, the aligned dataset of 622 sequences was phylogenetically analyzed. The SMS

method was used to select GTR+G as the base substitution model(Lefort, et al., 2017), and the PhyML

3.1(Guindon, et al., 2010) and MEGA(Kumar, et al., 2018) were used to construct the no-root

phylogenetic tree by the maximum likelihood method with the bootstrap value of 100. The online tool

iTOL(Letunic and Bork, 2019) was used to visualized the phylogenetic tree. The clusters were defined

by the shape of phylogenetic tree.

Detection of specific sites from each Cluster

Information (ID, countries/regions and collection time) and variants (NC_045512.2 as reference

genome) of each genome from each Cluster were extracted. The allele frequency and nucleotide

divergency (pi) for each site in the virus population of each Cluster were measured by

vcftools(Danecek, et al., 2011). The Fst were also calculated by vcftools(Danecek, et al., 2011) to

assess the diversity between the Clusters. Sites with high level of Fst together with different major

allele in each Cluster were filtered as the specific sites. PCA were analyzed by the GCTA

v1.93.1beta(Yang, et al., 2011) with the specific sites and all SNV dataset respectively.

Linkage analysis of specific sites and characteristics of major haplotype subgroups

The linkage disequilibrium of the specific sites were analyzed by haploview(Barrett, et al., 2005), and

the statistics of the haplotype of the specific sites for each Cluster or country were used in-house perl

script.

Phylogenetic network of haplotype subgroups

The phylogenetic networks were inferred by PopART package v1.7.2(Leigh, et al., 2015) using TCS

and minimum spanning network (MSN) methods respectively.

Frequencies of specific sites or haplotypes and correlation with death rate

The frequencies of specific sites for each country were calculated. The death rate was estimated with

Total Deaths /Confirmed Cases based on the data from Johns Hopkins resources on May 12th, 2020

(https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6).The

correlation coefficient between death rate and frequencies of specific site or haplotype in different

countries was calculated using Pearson method.

Authors' contributions

HD conceived the study. YM, YB, DJ, JL and XC carried out the data analysis and wrote the

manuscript. MH, SL, and ZC collected data and revised the manuscript. XW attended the discussions.

HD and YM supervised the whole work and revised the manuscript.

Conflict of Interest

The authors declare no conflict of interest.

Funding sources

This work was supported by the National Key R&D Program of China [2018YFC0910201], the Key

R&D Program of Guangdong Province [2019B020226001], the Science and the Technology Planning

Project of Guangzhou [201704020176 and 2020Q-P013].

Ethical Approval

Not required.

References

Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. (2020) The proximal origin of
SARS-CoV-2. *Nature Medicine*, 10.1038/s41591-020-0820-9

Barrett JC, Fry B, Maller J, Daly MJ. (2005) Haploview: analysis and visualization of LD and
haplotype maps. *Bioinformatics*, 21:263-5.

Bouckaert R, Vaughan TG, Barido-Sottani J, et al. (2019) BEAST 2.5: An advanced software platform
for Bayesian evolutionary analysis. *PLoS Comput Biol*, 15:e1006650.

Chan JF, Kok KH, Zhu Z, et al. (2020) Genomic characterization of the 2019 novel human-pathogenic
coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes*
Infect, 9:221-36.

Cotten M, Watson SJ, Zumla AI, et al. (2014) Spread, circulation, and evolution of the Middle East
respiratory syndrome coronavirus. *mBio*, 5:

Danecek P, Auton A, Abecasis G, et al. (2011) The variant call format and VCFtools. *Bioinformatics*
(Oxford, England), 27:2156-8.

Forster P, Forster L, Renfrew C, Forster M. (2020) Phylogenetic network analysis of SARS-CoV-2
genomes. *Proceedings of the National Academy of Sciences*, 10.1073/pnas.2004999117202004999.

Graham RL, Sparks JS, Eckerle LD, Sims AC, Denison MR. (2008) SARS coronavirus replicase
proteins in pathogenesis. *Virus Res*, 133:88-100.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. (2010) New algorithms and
methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst*
Biol, 59:307-21.

Guo YR, Cao QD, Hong ZS, et al. (2020) The origin, transmission and clinical therapies on
coronavirus disease 2019 (COVID-19) outbreak - an update on the status. *Mil Med Res*, 7:11.

He J, Peng G, Min J. (2004) Molecular Evolution of the SARS Coronavirus During the Course of the
SARS Epidemic in China. *Science*, 303:

Holder M, Lewis PO. (2003) Phylogeny estimation: traditional and Bayesian approaches. *Nat Rev*
Genet, 4:275-84.

Huang C, Wang Y, Li X, et al. (2020) Clinical features of patients infected with 2019 novel coronavirus
in Wuhan, China. *Lancet*, 395:497-506.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. (2018) MEGA X: Molecular Evolutionary Genetics
Analysis across Computing Platforms. *Mol Biol Evol*, 35:1547-9.

Lam TT, Shum MH, Zhu HC, et al. (2020) Identifying SARS-CoV-2 related coronaviruses in Malayan
pangolins. *Nature*, 10.1038/s41586-020-2169-0

Langmead B, Salzberg SL. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods*, 9:357-9.

Lefort V, Longueville JE, Gascuel O. (2017) SMS: Smart Model Selection in PhyML. *Mol Biol Evol*,
34:2422-4.

Leigh JW, Bryant D, Nakagawa S. (2015) popart: full‑feature software for haplotype network
construction. *Methods in Ecology and Evolution*, 6:1110-6.

Letunic I, Bork P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res, 47:W256-W9.
Li F. (2016) Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol*,
3:237-61.
Li H. (2011) A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics (Oxford, England)*,
27:2987-93.
Li H, Handsaker B, Wysoker A, et al. (2009) The Sequence Alignment/Map format and SAMtools.
Bioinformatics (Oxford, England), 25:2078-9.
Liu Y, Du X, Chen J, et al. (2020) Neutrophil-to-lymphocyte ratio as an independent risk factor for
mortality in hospitalized patients with COVID-19. *J Infect*, 10.1016/j.jinf.2020.04.002
Liu Y, Gayle AA, Wilder-Smith A, Rocklov J. (2020) The reproductive number of COVID-19 is higher
compared to SARS coronavirus. *J Travel Med*, 27:
Lu R, Zhao X, Li J, et al. (2020) Genomic characterisation and epidemiology of 2019 novel
coronavirus: implications for virus origins and receptor binding. *Lancet*, 395:565-74.
Malavolta M, Giacconi R, Brunetti D, Provinciali M, Maggi F. (2020) Exploring the Relevance of
Senotherapeutics for the Current SARS-CoV-2 Emergency and Similar Future Global Health Threats.
Cells, 9:
Rambaut A, Lam TT, Max Carvalho L, Pybus OG. (2016) Exploring the temporal structure of
heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol*, 2:vew007.
van Dorp L, Acman M, Richard D, et al. (2020) Emergence of genomic diversity and recurrent
mutations in SARS-CoV-2. *Infection, Genetics and Evolution*,
https://doi.org/10.1016/j.meegid.2020.104351104351.
WHO. (2020) Coronavirus disease 2019 (COVID-19) Situation Report – 93. *WHO*,
Wrapp D, Wang N, Corbett KS, et al. (2020) Cryo-EM structure of the 2019-nCoV spike in the
prefusion conformation. *Science*, 367:1260-3.
Yang J, Lee SH, Goddard ME, Visscher PM. (2011) GCTA: a tool for genome-wide complex trait
analysis. *Am J Hum Genet*, 88:76-82.
Yu WB, Tang GD, Zhang L, Corlett RT. (2020) Decoding the evolution and transmissions of the novel
pneumonia coronavirus (SARS-CoV-2 / HCoV-19) using whole genomic data. *Zool Res*, 41:247-57.
Yuen K-S, Ye Z-W, Fung S-Y, Chan C-P, Jin D-Y. (2020) SARS-CoV-2 and COVID-19: The most
important research questions. *Cell Biosci*, 10:40.
Zhang L, Shen F-m, Chen F, Lin Z. (2020) Origin and Evolution of the 2019 Novel Coronavirus.
Clinical Infectious Diseases, 10.1093/cid/ciaa112
Zhang YZ, Holmes EC. (2020) A Genomic Perspective on the Origin and Emergence of SARS-CoV-2.
Cell, 181:223-7.
Zhang Z, Li J, Zhao XQ, Wang J, Wong GK, Yu J. (2006) KaKs_Calculator: calculating Ka and Ks
through model selection and model averaging. *Genomics Proteomics Bioinformatics*, 4:259-63.
Zhao Z, Li H, Wu X, et al. (2004) Moderate mutation rate in the SARS coronavirus genome and its
implications. *BMC Evol Biol*, 4:21.
Zhou P, Yang XL, Wang XG, et al. (2020) A pneumonia outbreak associated with a new coronavirus of
probable bat origin. *Nature*, 579:270-3.

Figure 1   Phylogenetic tree and clusters of 622 SARS-CoV-2 genomes

The 622 sequences were clustered into three clusters: Cluster 1 were mainly from North America,

Cluster 2 were from regions all over the world, and Cluster 3 were mainly from Europe.

Figure 2   Linkage disequilibrium plot of haplotypes of the 9 specific sites

A. The plot for 622 genome sequences; **B**. The plot for 3624 genome sequences.

Figure 3   The frequencies of both the 9 specific sites and haplotypes

The frequencies of the 9 specific sites (**A**) and haplotypes (**B**) in each country for 3624 genomes.

Figure 4   The characteristics of haplotype subgroups

A. The numbers of haplotypes of the 9 specific sites for 16373 genomes with clear collection data

detected in each country in chronological order; **B**. The whole genome mutations in each major

haplotype subgroup.

Figure 5   Phylogenetic network of haplotype subgroups for 3624 genomes

The network was inferred by POPART using TCS method. Each colored vertex represents a haplotype,

with different colors indicating the different sampling areas. Hatch marks along edge indicated the

number of mutations. Small black circles within the network indicated unsampled haplotypes. H1-H5

subgroups were point out according haplotypes of the 9 specific sites, and other small subgroups were

not specially pointed out.

Figure 6 The correlation between death rate and frequencies of both the 9 specific sites and

haplotypes

Tables

Table 1   Statistics of Ka, Ks and Ka/Ks ratios for all coding regions of the SARS-CoV,

MERS-CoV and SARS-CoV-2 genome sequences

Table 2 Substitution rate and tMRCA estimated by BEAST v2.6.2

Table 3 The information of the 9 specific sites in each Cluster

Table 4 Statistics of Ka, Ks and Ka/Ks ratios for all coding regions of each 4 major haplotype

subgroup with 3624 and 16373 genomes respectively

Supplementary material

Figure S1   Regression analyses of the root-to-tip divergence with the maximum-likelihood

phylogenetic tree of SARS-CoV (A), MERS-CoV (B) and SARS-CoV-2 (C)

The root-to-tip genetic distances were inferred in TempEst v1.5.3. Three data sets exhibit a positive

correlation between root-to-tip divergence and sample collecting time, and they appear to be suitable

for molecular clock analysis.

Figure S2 Substitution rate and tMRCA estimate of SARS-CoV-2

A. Estimated substitution rate of SARS-CoV-2 using BEAST v2.6.2 under a strict molecular clock. The

mean rate was 1.601×10^{-3} (substitutions per site per year). **B**. The estimated tMRCA of SARS-CoV-2

using BEAST v2.6.2 under a strict molecular clock. The mean tMRCA was 2019.74 (date:

2019-09-27).

Figure S3   Phylogenetic tree of 622 SARS-CoV-2 genomes

A. Phylogenetic tree constructed by PhyML with bootstrap value of 100; **B**. Phylogenetic tree

constructed by MEGA with bootstrap value of 100.

Figure S4   The PCAs analysis result based on the 9 specific sites (A) and all SNVs (B)

Table S1   The information of the first downloaded complete genome sequences

Table S2   The sequence IDs in each Cluster

Table S3   The details information of the SNVs in all Clusters

Table S4   The information of death rate, frequencies of the 9 specific sites and major haplotypes

in each country for 3624 and 16373 genomes respectively

Table S5   All haplotypes of the 9 specific sites and numbers of them for 3624 and 16373 genomes

respectively

Table S6   Whole genome mutations and frequencies in each major haplotype subgroup of 3624

and 16373 genomes respectively

Tree scale: 0.0001

Inner Color: Continents
- Asia
- Europe
- NorthAmerica
- Oceania
- SouthAmerica

Shape Plot: Haplotype
- H1
- H2
- H3
- H4
- H5
- H6
- Other

Outer Clolors: Countries
- Australia
- Belgium
- Brazil
- Cambodia
- Canada
- Chile
- China
- Denmark
- Finland
- France
- Georgia
- Germany
- HongKong_China
- Hungary
- India
- Ireland
- Italy
- Japan
- Kuwait
- Luxembourg
- Mexico
- Nepal
- Netherlands
- NewZealand
- Poland
- Portugal
- SaudiArabia
- Singapore
- SouthKorea
- Sweden
- Switzerland
- Taiwan_China
- Thailand
- UnitedKingdom
- UnitedStates
- Vietnam

Cluster 1

Cluster 2

Cluster 3

Cluster 2

		Ka (mean) 10⁻³	s.e. (Ka) 10⁻³	Ks (mean) 10⁻³	s.e. (Ks) 10⁻³	Ka/Ks (mean)	s.e. (Ka/Ks)	Hypothesis	P-value* 622 genomes	P-value* 3624 genomes
	SARS-CoV	0.985	0.018	1.310	0.049	0.760	0.038	Ka/Ks (SARS-CoV < SARS-CoV-2)	0.139	0.084
	MERS-CoV	1.319	0.040	4.887	0.096	0.260	0.003	Ka/Ks (MERS-CoV < SARS-CoV-2)	0.000	0.000
SARS-CoV-2	622 genomes	0.231	0.004	0.265	0.005	1.008	0.020			
	3624 genomes	0.287	0.002	0.298	0.002	1.094	0.010			

Note: *One-sided Mann-Whitney U-test for the means of two independent samples.

	substitution rate (10^{-3})	95% CI (10^{-3})	tMRCA	95% CI	references
SARS-CoV	1.050	0.489, 1.654	2002/6/28	2002/1/19, 2002/11/3	this study
	-	0.800, 2.380	spring of 2002	-	[17]
MERS-CoV	1.516	1.392, 1.632	2012/6/7	2012/6/4, 2012/6/9	this study
	1.120	0.876, 1.370	2012/3	2011/12, 2012/6	[18]
SARS-CoV-2	1.601	1.418, 1.796	2019/9/27	2019/8/28, 2019/10/26	this study

17. Zhao Z, Li H, Wu X, Zhong Y, Zhang K, Zhang YP, et al. Moderate mutation rate in the SARS coronavirus genome and its implications. BMC Evol Biol. 2004;4:21.

18. Cotten M, Watson SJ, Zumla AI, Makhdoom HQ, Palser AL, Ong SH, et al. Spread, circulation, and evolution of the Middle East respiratory syndrome coronavirus. mBio. 2014;5(1).

Pos	Ref	Alt	Cluster 1			Cluster 2			Cluster 3			Fst	Gene region	Mutation type	Protein changed	codon changed	Impact
			pi	Major allele	frequency	pi	Major allele	frequency	pi	Major allele	frequency						
241	C	T	0.0000	C	1.0000	0.0109	C	0.9945	0.0000	T	1.0000	0.9912	5'UTR	upstream	NA	NA	MODIFIER
3037	C	T	0.0000	C	1.0000	0.0109	C	0.9945	0.0000	T	1.0000	0.9912	gene-orf1ab	synonymous	924F	2772ttC>ttT	LOW
8782	C	T	0.0000	T	1.0000	0.3863	C	0.7390	0.0000	C	1.0000	0.5821	gene-orf1ab	synonymous	2839S	8517agC>agT	LOW
14408	C	T	0.0000	C	1.0000	0.0055	C	0.9973	0.0000	T	1.0000	0.9956	gene-orf1ab	missense	4715P>L	14144cCt>cTt	MODERAT
17747	C	T	0.0735	T	0.9620	0.0000	C	1.0000	0.0000	C	1.0000	0.9752	gene-orf1ab	missense	5828P>L	17483cCt>cTt	MODERAT
17858	A	G	0.0497	G	0.9747	0.0000	A	1.0000	0.0000	A	1.0000	0.9836	gene-orf1ab	missense	5865Y>C	17594tAt>tGt	MODERAT
18060	C	T	0.0000	T	1.0000	0.0164	C	0.9918	0.0000	C	1.0000	0.9761	gene-orf1ab	synonymous	5932L	17796ctC>ctT	LOW
23403	A	G	0.0000	A	1.0000	0.0164	A	0.9918	0.0000	G	1.0000	0.9869	gene-S	missense	614D>G	1841gAt>gGt	MODERAT
28144	T	C	0.0000	C	1.0000	0.3915	T	0.7335	0.0000	T	1.0000	0.5785	gene-ORF8	missense	84L>S	251tTa>tCa	MODERAT

		Ka (mean) 10^{-3}	s.e. (Ka)10^{-3}	Ks (mean) 10^{-3}	s.e. (Ks)10^{-3}	Ka/Ks (mean)	s.e. (Ka/Ks)	Hypothesis	P-value*
	H1	0.510	0.057	0.498	0.056	1.099	0.010	Ka/Ks (H3 < H1)	0.000
	H2	0.347	0.042	0.334	0.044	1.268	0.023	Ka/Ks (H3 < H2)	0.000
3624 genomes	H3	0.298	0.007	0.397	0.009	0.795	0.010	-	-
	H4	0.334	0.023	0.414	0.019	0.796	0.019	-	-
	H1	0.416	0.011	0.384	0.011	1.178	0.005	Ka/Ks (H3 < H1)	0.000
	H2	0.332	0.014	0.316	0.014	1.224	0.012	Ka/Ks (H3 < H2)	0.000
16373 genomes	H3	0.305	0.004	0.401	0.005	0.809	0.007	-	-
	H4	0.354	0.010	0.477	0.009	0.749	0.010	-	-

Note: *One-sided Mann-Whitney U-test for the means of two independent samples.