Evolution and molecular characteristics of SARS-CoV-2

genome

Yunmeng Bai[1#], Dawei Jiang[1#], Jerome R Lon[1#], Xiaoshi Chen[1], Meiling Hu[1], Shudai

Lin[1], Zixi Chen[1], Xiaoning Wang[1,2], Yuhuan Meng[3*], Hongli Du[1*]

[1]*School of Biology and Biological Engineering, South China University of Technology,*

Guangzhou 510006, China

[2]*State Clinic Center of Gearitic, Chinese PLA General Hospital, Beijing 100853,*

China

[3]*Guangzhou KingMed Transformative Medicine Institute Co., Ltd, Guangzhou*

510330, China

[#] Equal contribution.

[*] Corresponding authors.

E-mail: hldu@scut.edu.cn(Du HL), zb-mengyuhuan@kingmed.com.cn(Meng YH)

There are 6 figures, 4 tables, 4 supplementary figures and 6 supplementary tables.

Abstract

In the evolution analysis of 622 or 3624 complete SARS-CoV-2 genomes, the

estimated Ka/Ks ratio is 1.008 or 1.094, which is higher than that of SARS-CoV and

MERS-CoV, and the time to the most recent common ancestor (tMRCA) is inferred in

late September 2019, indicating that SARS-CoV-2 may have completed positive

selection pressure of the cross-host evolution in the early stage. Further we find 9 key

specific sites of highly linkage which play a decisive role in the classification of

SARS-CoV-2. Notably the frequencies of haplotype TTTG and H1 are generally high

in European countries and correlated to death rate (r>0.37) based on 3624 or 16373

SARS-CoV-2 genomes, which indicated that the haplotypes might be related to

pathogenicity of SARS-CoV-2. The development trends in chronological order and

Ka/Ks ratio of each major haplotype subgroup showed the four major haplotype

subgroups were going through different evolution patterns, the H3 haplotype

subgroup was going through a purifying evolution and almost disappeared after

detection, while H1 haplotype subgroup was going through a neutral evolution and

globally increased with time. The evolution and molecular characteristics of more

than 16000 genomic sequences provided a new perspective for revealing

epidemiology of SARS-CoV-2 and coping with SARS-CoV-2 effectively.

KEYWORDS: SARS-CoV-2; evolution; classification; haplotype

Introduction

The global outbreak of SARS-CoV-2 is currently and increasingly recognized as a

serious, public health concern worldwide. To date, a total of seven types of

coronaviruses that can infect humans have been found, among which the more typical

ones are SARS-CoV, MERS-CoV and SARS-CoV-2. They spread across species and

usually cause mild to severe respiratory diseases. The total number of SARS-CoV and

MERS-CoV infections are only 8,069 and 2,494 with reproduction number (R0)

fluctuates 2.5-3.9 and 0.3-0.8, respectively. However, SARS-CoV-2 has infected

2471136 people in 212 countries up to April 22th(1), with the basic R0 ranging from

1.4 to 6.49(2). Among these three typical coronavirus, MERS-CoV has the highest

death rate of 34.40%, SARS-CoV has the modest death rate of 9.59%, SARS-CoV-2

is about 6.99% and 7.69% death rate of the global and Wuhan, but is quite high death

rate in some countries of Europe, such as Belgium, Italy, United Kingdom,

Netherlands, Spain and France, which even reaches 14.95%, 13.39%, 13.48%,

11.61%, 10.42% and 13.60% respectively according to the data of April 22th, 2020.

Except for the shortage of medical supplies, aging and other factors, it is not clear

whether there is virus mutation effect in these countries with such a significantly

increased death rate.

It has been reported that SARS-CoV-2 belongs to beta-coronavirus and is mainly

transmitted by the respiratory tract, which belongs to the same subgenus

(SarbeCoVirus) as SARS-CoV(3). Through analyzing the genome and structure of

SARS-CoV-2, its receptor-binding domain (RBD) was found to bind with

angiotensin-converting enzyme 2 (ACE2), which is also one of the receptors for

binding SARS-CoV(4). Some early genomic studies have shown that SARS-CoV-2 is

similar to certain bat viruses (RaTG13, with the whole genome homology of 96.2%(5))

and Malayan pangolins coronaviruses (GD/P1L and GDP2S, with the whole genome

homology of 92.4%(6)). They have speculated several possible origins of

SARS-CoV-2 based on its spike protein characteristics (cleavage sites or the

RBD)(5-7), in particular, SARS-CoV-2 exhibits 97.4% amino acid similarity to the

Guangdong pangolin coronaviruses in RBD, even though it is most closely related to

bat coronavirus RaTG13 at the whole genome level. However, it is not enough to

present genome-wide evolution by a single gene or local evolution of RBD, whether

bats or pangolins play an important role in the zoonotic origin of SARS-CoV-2

remains uncertain(8). Furthermore, since there is little molecular characteristics

analysis of SARS-CoV-2 based on a large number of genomes, it is difficult to

determine whether the virus has significant variation that affects its phenotype.

Thereby, it is necessary to further reveal the phylogenetic evolution and molecular

characteristics of the whole genome of SARS-CoV-2 in order to develop a

comprehensive understanding of the virus and provide a basis for designing vaccines

and antiviral treatment strategies. Fortunately, with the increase of SARS-CoV-2

genome data, we have an opportunity to reveal phylogenetic evolution and molecular

characteristics of SARS-CoV-2 more comprehensively. In this study, the Ka/Ks ratio,

substitution rate and the time to the most recent common ancestor (tMRCA) of

SARS-CoV-2, SARS-CoV and MERS-CoV were analyzed using KaKs_Calculator

and BEAST, and then 622 SARS-CoV-2 genome sequences with high quality were

clustered using maximum likelihood (ML) with bootstrap of 100 to ensure the

reliability of evolutionary trees. According to the clusters of phylogenetic trees, the

characteristic mutations of SARS-CoV-2 were screened and their potential

significance were explored. The results of this study will provide a landscape for us to

further understand SARS-CoV-2, and supply a possible basis for the prevention and

treatment of SARS-CoV-2 in different areas.

Results

Genome sequences

We got a total of 1053 genomic sequences up to March 22th, 2020. According to the

filter criteria, 37 sequences with ambiguous time, 314 with low quality and 78 with

similarity of 100% were removed. A total of 624 sequences were obtained to perform

multiple sequences alignment. Two highly divergent sequences (EPI_ ISL_414690,

EPI_ISL_415710) according to the firstly constructed phylogenetic tree were also

filtered out (Table S1). The remaining 622 sequences were used to reconstruct a

phylogenetic tree. In addition, total of 3624 and 16373 genomic sequences were

redownloaded up to April 6th, 2020 and May 10th, 2020 respectively for further

exploring the evolution and molecular characteristics of SARS-CoV-2.

Estimate of evolution rate and the time to the most recent common ancestor for

SARS-CoV, MERS-CoV, and SARS-CoV-2

In this study, date for SARS-CoV-2 ranged from 2019/12/26 to 2020/03/18 was

collected. The average Ka/Ks of all the coding sequences was closer to 1 (1.008),

which indicating the genome was going through a neutral evolution. We also

reevaluated the Ka/Ks of SARS-CoV and MERS-CoV through the whole period, and

found the ratio were smaller than SARS-CoV-2 (**Table 1**). To estimate more credible

Ka/Ks for SARS-CoV-2, we recalculated it using redownloaded 3624 genome

sequences ranged from 2019/12/26 to 2020/04/06, and the average Ka/Ks of it was

1.094 (Table 1), which was almost same with the above result.

We assessed the temporal signal using TempEst v1.5.3(9). All three datasets

exhibit a positive correlation between root-to-tip divergence and sample collecting

time (Figure S1), hence they are suitable for molecular clock analysis in BEAST(9,

10). The substitution rate of SARS-CoV-2 genome was estimated to be 1.601×10^{-3}

(95% CI: 1.418 x 10^{-3} - 1.796 x 10^{-3}, **Table 2**, Figure S2A) substitution/site/year,

which is as in the same order of magnitude as SARS-CoV and MERS-CoV. The

tMRCA was inferred on the late September, 2019 (95% CI: 2019/08/28-

2019/10/26,Table 2, Figure S2B), about 2 months before the early cases of

SARS-CoV-2(11).

Phylogenetic tree and clusters of SARS-CoV-2

The no-root phylogenetic trees constructed by the maximum likelihood method with

PhyML 3.1(12) and MEGA(13) were showed in **Figure 1**, Figure S3A and 3B.

According to the shape of phylogenetic trees, we divided 622 sequences into three

clusters: Cluster 1 including 76 sequences mainly from North America, Cluster 2

including 367 sequences from all regions of the world, and Cluster 3 including 179

sequences mainly from Europe (Table S2).

The specific sites of each Cluster

The *Fst* and population frequency of a total of 9 sites (NC_045512.2 as reference

genome) were detected (**Table 3**, Table S3). Thereinto, three (C17747T, A17858G and

C18060T) are the specific sites of Cluster 1, and four (C241T, C3037T, C14408T and

A23403G) are the specific sites of Cluster 3. Notably, C241T was located in the

5'-UTR region and the others were located in coding regions (6 in *ofr1ab* gene, 1 in *S*

gene and 1 in *ORF8* gene). Five of them were missense variant, including C14408T,

C17747T and A17858G in *ofr1ab* gene, A23403G in *S* gene, and C28144T in *ORF8*

gene. The PCA results showed that these 9 specific sites could clearly separate the

three Clusters, while all SNV dataset could not clearly separate Cluster 1 and Cluster

2 (Figure S4), which further suggested that these 9 specific sites are the key sites for

separating the three Clusters.

Linkage of specific sites

We found the 9 specific sites are highly linkage based on 622 genome sequences

(**Figure 2A**), then we carried out a further linkage analysis using the 3624 genome

sequences (Figure 2B). As a result, for the 3624 genome sequences, 3 specific sites in

the Cluster 1 were almost complete linkage, and haplotype CAC and TGT accounted

for 98.65% of all the 3 site haplotypes. The same phenomenon was also found in 4

specific sites of Cluster 3, and haplotype CCCA and TTTG accounted for 97.68% of

all the 4 site haplotypes. Intriguingly, the 9 specific sites were still highly linkage, and

four haplotypes, including TTCTCACGT (H1), CCCCCACAT (H2), CCTCTGTAC

(H3) and CCTCCACAC (H4), accounted for 95.89% of all the 9 site haplotypes.

Thereinto, H1 and H3 had completely different bases at the 9 specific sites. The

frequencies of each site and major haplotype for each country were showed in **Figure**

3 and Table S4. The data showed that the haplotype TTTG of the 4 specific sites in

Cluster 3 had existed globally at present, and still exhibited high frequencies in most

European countries but quite low in Asian countries. While the haplotype TGT of the

3 specific sites in Cluster 1 existed almost only in North America and Australia. For

the 9 specific sites, most countries had only two or three major haplotypes, except

America and Australia had all of four major haplotypes with higher frequency.

Characteristics of major haplotype subgroups

All haplotypes of 9 specific sites for 3624 or 16373 genomes and the numbers of them

were shown in Table S5. Four major haplotypes H1, H2, H3 and H4, three minor

haplotypes H5, H7 and H8 close to H1 and one minor haplotype H6 close to H3 were

found in both datasets. The numbers of these haplotypes for 16373 genomes with

clear collection data detected in each country in chronological order were shown in

Figure 4A. From these results, H2 and H4 haplotype subgroups had existed for a long

time (2019-12-24 to 2020-04-28), and the H3 haplotype subgroup almost disappeared

after detection (2020-02-18 to 2020-04-28), while H1 haplotype subgroup was

globally increasing with time (2020-02-18 to 2020-05-05), which indicated H1

subgroup had adapted to the human hosts, and was under an adaptive growth period

worldwide. However, due to the nonrandom sampling on early phase (only patients

having a recent travel to Wuhan were detected), some earlier cases of H3 may be lost,

the high proportion of H3 subgroup during February 18 to March 10 could confirm

this point.

H3 and H4 subgroups had the lowest Ka/Ks ratio in 3624 or 16373 genomes among

the 4 major subgroups (**Table 4**), suggesting that H3 and H4 subgroup might be going

through a purifying evolution and have existed for a long time, while H2 and H1

subgroup might be going through a positive selection and a neutral evolution

respectively according to their Ka/Ks ratios (Table 4), which were consistent with the

above phenomenon that only H1 subgroup was expanding with time around the world.

From the whole genome mutations in each major haplotype subgroup (Figure 4B,

Table S6), we found that except the 9 specific sites, there was no common mutations

with frequencies more than 0.05 in these haplotype subgroups but between H2 and H4

subgroups.

Phylogenetic network of haplotype subgroups

Phylogenetic networks were inferred with 697 mutations called from 3624 genomes

dataset. For these datasets, the network structures of TCS and MSN were similar. The

major haplotype subgroups H4 and H2 were in the middle of the network, while H1

and H3 were in the end nodes of the network (**Figure 5**). According to the

phylogenetic networks, we proposed four hypothesis: (1) the ancestral haplotypes

evolved in four directions to obtain H1, H2, H3 and H4 respectively or evolved in two

or more directions to obtain two or more major haplotypes and then involved into the

other major haplotype(s); (2) H2 or H4 evolved in two directions respectively and

finally generated H1 and H3; (3) H1 evolved in one direction to generate H2 and H4,

and then evolved into H3; (4) H3 evolved in one direction to generate H4 and H2, and

then evolved into H1. Although we cannot exclude the first hypothesis based on the

present data, but if there are evolutionary transformation relationships between the

four major haplotypes, according to the Ka/Ks, population size and development trend

in chronological order of each major haplotype subgroup, we speculate that the most

likely transformation hypothesis is that H3 is the ancestral haplotype, which is

gradually eliminated with selection, while H4 and H2 is the transitional haplotype in

the evolution process, and H1 may be the haplotype to be finally fixed.

Correlation analysis of specific sites with death rate

To explore the relationship between death rate and the 9 specific sites, we used

Pearson method to calculate the correlation coefficient between death rate and

frequency of each specific site or major haplotype in 17 countries with 3624 genomes

at early stage. As a result, all r values of 241T, 3037T, 14408T, 23403G and haplotype

TTTG and H1 were more than 0.4 (**Figure 6**, Table S4). We also evaluated the

correlation coefficient with 16373 genomes in 30 countries, the r values of haplotype

TTTG and H1 were still more than 0.37 (Table S4), which might be increased when

the death rates of some countries were further stabilized with time. These finding

integrating their high frequencies in most European countries indicated that the 4 sites

and haplotype TTTG and H1 might be related to the pathogenicity of SARS-CoV-2.

Discussion

SARS-CoV-2 poses a great threat to the production, living and even survival of

human beings(14). With the further outbreak of SARS-CoV-2 in the world, further

understanding on the evolution and molecular characteristics of SARS-CoV-2 based

on a large number of genome sequences will help us cope better with the challenges

brought by SARS-CoV-2. In the present study, we estimated the Ka/Ks ratio and

tMRCA of SARS-CoV-2, SARS-CoV and MERS-CoV, and constructed the

phylogenetic tree of SARS-CoV-2 genomes. As a result, we found that 9 specific sites

of highly linkage could play a decisive role in the classification of SARS-CoV-2, and

the haplotype TTTG might be related to pathogenicity of SARS-CoV-2 based on more

than 16000 genomes. All these results could accelerate our further understanding of

SARS-CoV-2.

Exploring evolution rate, tMRCA and phylogenetic tree of SARS-CoV-2 would

help us better understand the virus(15). The average Ka/Ks for all the coding

sequences of 622 and 3624 SARS-CoV-2 genomes was 1.008 and 1.094, which was

higher than those of SARS-CoV and MERS-CoV, indicating that the SARS-CoV-2

was going through a neutral evolution. In general, SARS-CoV would show purifying

selection pressures and the Ka/Ks would go down at the middle or end of the

epidemic(16). Therefore, it seemed to be reasonable considering that SARS-CoV-2

was under an exponential growth period around the world. Interestingly, we also

found that the subgroups of different haplotypes (H1, H2, H3 and H4) seemed to

experience different evolutionary patterns according to their Ka/Ks. The H3 haplotype

subgroup disappeared soon after detection (2020-02-18 to 2020-04-07, Figure 4A),

while H1 haplotype subgroup was globally increasing with time, these evolution and

changes should be considered in developing therapeutic drugs and vaccines as soon as

possible. The tMRCA of SARS-CoV-2 was inferred on the late September, 2019 (95%

CI: 2019/08/28- 2019/10/26), about 2 months before the early cases of

SARS-CoV-2(11). We also estimated tMRCA of SARS-CoV and MERS-CoV with

the same methods, both were about 3 months later than the corresponding tMRCAs

estimated by previous studied(17, 18), which indicated the tMRCA of SARS-CoV-2

we estimated was reasonable. A recent study used TreeDater method to estimate

tMRCA for more than 7000 SARS-CoV-2 genomes and indicated the tMRCA of

SARS-CoV-2 was around 6 October 2019 to 11 December 2019, which were about

one and a half months later than ours and in broad agreement with six previous

studies all performed on smaller subsets of the SARS-CoV-2 genomes with BEAST

method(19). While we used the most common method BEAST to estimate tMRCA of

SARS-CoV-2 based on 622 genomes, indicating that the estimated tMRCA are

reliable.

A recent study clustered 160 SARS-CoV-2 whole-genome sequences into A, B

and C groups by a phylogenetic network analysis(20). The cluster result was similar

to our study: both the samples in Cluster A and our Cluster 1 were mainly from the

United States; the samples in Cluster C and our Cluster 3 were mainly from European

countries, while Cluster B and our Cluster 2 were mainly from China and the other

regions. It was interesting that the markers in Cluster B (T8782C and C28144T) were

also found in our study, but the other markers in Cluster A (T29095C) and Cluster C

(G26144T) were not significantly in our study. That may be caused by different

sample sizes and different constructing methods of phylogenetic tree. Based on the

base substitution model, the ML method avoids the possible "long-branch attraction"

problem in the maximum parsimony method and is faster than the Bayesian

method(21), hence it could be used as a reliable method for phylogenetic analysis.

Nowadays, there is still a shortage of studies on phylogenetic analysis of

SARS-CoV-2. Some studies used the genome of bat SARS-like-CoV(22), RaTG13(22)

or MT019529 from Wuhan (https://bigd.big.ac.cn/ncov/tree) as the root of

phylogenetic tree. Unfortunately, there was no obvious evidence showing that

SARS-CoV-2 was from the bat coronavirus even the identity between SARS-CoV-2

and RaTG13 was up to 96.2%(5). In our study, the tMRCA of SARS-CoV-2 was

inferred on the late September 2019, which indicated there might exist an earlier

SARS-CoV-2 strain we didn't find. Then in the case of unclear source of

SARS-CoV-2 and high homology of its genomes (> 99.9% homology mostly), it may

be inappropriate to identify the evolutionary characteristics inside the genomes by

taking bat SARS-like-CoV, RaTG13 or MT019529 as root. Therefore, the maximum

likelihood method was used in current study to construct a no-root tree to obtain the

reliable clusters with different characteristics. Based on the no-root tree, we identified

9 specific sites of highly linkage that play a decisive role in the classification of

clusters successfully. Among the four major haplotypes, H1 and H3 were in Cluster 3

and Cluster 1, respectively, while there were two haplotypes H2 and H4 in Cluster 2

(Figure 1, Figure S3B).

Among the 9 specific sites, 8 of them were located in coding regions (6 in *ofr1ab*

gene, 1 in *S* gene and 1 in *ORF8* gene), and 5 of them were missense variant,

including C14408T, C17747T and A17858G in *ofr1ab* gene, A23403G in *S* gene, and

C28144T in *ORF8* gene. *Orf1ab* is composed of two partially overlapping open

reading frames (orf), orf1a and 1b. It is proteolytic cleaved into 16 non-structural

proteins (nsp), including nsp1 (suppress antiviral host response), nsp9 (RNA/DNA

binding activity), nsp12 (RNA-dependent RNA polymerase), nsp13 (helicase) and

others(23), indicating the vital role of it in transcription, replication, innate immune

response and virulence(24). C14408T with high frequencies of T in European

countries were located at nsp12 region, indicating that this missense variant might

influence the role of RNA polymerase. Spike glycoprotein, the largest structural

protein on the surface of coronaviruses, comprises of S1 and S2 subunits which

mediating binding the receptor on the host cell surface and fusing membranes,

respectively(25). It has been reported that S protein of SARS-CoV-2 can bind

angiotensin-converting enzyme 2 (ACE2) with higher affinity than that of SARS(4, 5).

Whether the missense variant of A23403G in *S* gene, which with high frequencies of

G in European countries, will change the affinity between S protein and ACE2

remains to be further investigated.

It seems to take a long time to fix mutations finally according to the mutation

frequency of each subgroup. For example, H2 and H4 subgroups, which have been

detected for more than four months from December 24, 2019 to May 5, 2020 (Figure

4A), have more mutations with higher frequencies, but the highest mutation frequency

is only 0.486 at the position of 11083 (Figure 4B, Table S6). From these phenomena,

it can be inferred that it takes a long time for the specific sites of each major subgroup

to be fixed, but it may be faster if the early population is small. In addition, there is

also the possibility that an ancestor strain evolved in four directions by obtaining the

specific mutations directly and produced the four current major haplotypes, so the

evolution time may be shorter, which seems to be consistent with the phenomenon

that four major haplotypes can be detected in two months (Figure 4A). However, if

there exist transformation evolution pattern among the major haplotypes of

SARS-CoV-2, it is difficult to complete the transformation among the four major

haplotypes within two months at the current evolution rate of each major haplotype

population. Therefore, we speculate that the transformation between the four major

haplotypes may have been completed for a long time, which have not been detected.

Previous studies showed that the death rate of SARS-CoV-2 is affected by many

factors, such as medical supplies, aging, life style and other factors(26, 27). However,

our study found that the 4 specific sites (C241T, C3037T, C14408T and A23403G) in

the Cluster 3 were almost complete linkage, and the frequency of haplotype TTTG

was generally high in European countries and correlated to death rate (r>0.37) based

on 3624 or 16373 SARS-CoV-2 genomes, which provides a new perspective to the

reasons of relatively high death rate in Europe. In order to make frequency relatively

credible, we selected countries with more than 10 genomes (except South Korea with

16 genomes, all the other countries with more than 20 genomes) to analyze the

correlation between frequencies of haplotypes and death rate. Therefore, the

correlation between the frequencies of haplotypes and death rate obtained in this study

is quite reliable, and these specific sites should be considered in designing new

vaccine and drug development of SARS-CoV-2.

Conclusion and prospective

The Ka/Ks ratio of SARS-CoV-2 and tMRCA of SARS-CoV-2 (95% CI: 2019/08/28-

2019/10/26) indicated that SARS-CoV-2 might have completed the positive selection

pressure of cross-host evolution in the early stage and be going through a neutral

evolution at present. The 9 specific sites with highly linkage were found to play a

decisive role in the classification of clusters. Thereinto, 3 of them are the specific sites

of Cluster 1 mainly from North America and Australia, 4 of them are the specific sites

of Cluster 3 mainly from Europe in the early stage, both of these 3 and 4 specific sites

are almost complete linkage. The frequencies of haplotype TTTG for the 4 specific

sites and H1 for 9 specific sites were generally high in European countries and

correlated to death rate (r>0.37) based on 3624 or 16373 SARS-CoV-2 genomes,

which suggested these haplotypes might relate to pathogenicity of SARS-CoV-2 and

provided a new perspective to the reasons of relatively high death rates in Europe.

The relationship between the haplotype TTTG or H1 and the pathogenicity of

SARS-CoV-2 needs to be further verified by clinical samples or virus virulence test

experiment, and the 9 specific sites with highly linkage may provide a starting point

for the traceability research of SARS-CoV-2. Given that the different evolution

patterns of different haplotypes subgroups, we should consider these evolution and

changes in the development of therapeutic drugs and vaccines as soon as possible.

Materials and methods

Genome sequences

The complete genome sequences of SARS-CoV-2 were downloaded from China

National Center for Bioinformation (https://bigd.big.ac.cn/ncov/release_genome/) and

GISAID (https://www.gisaid.org/) up to March 22th, 2020. The sequences were

filtered out according to the following criteria: (1) sequences with ambiguous time; (2)

sequences with low quality which contained the counts of unknown bases > 15 and

degenerate bases > 50 (https://bigd.big.ac.cn/ncov/release_genome); (3) sequences

with similarity of 100% were removed to unique one. Finally, 624 high quality

genomes with precise collection time were selected and aligned using MAFFT v7

with automatic parameters. Besides, the genome sequences of 7 SARS-CoV and 475

MERS-CoV were also downloaded from NCBI (https://www.ncbi.nlm.nih.gov/), and

the MERS-CoV dataset including samples collected from both human and camel. In

addition, for further exploring evolution and molecular characteristics of

SARS-CoV-2 based on the larger amount of genomic data, we redownloaded

validation datasets of the genome sequences from GISAID up to April 6th, 2020 and

May 10th, 2020 respectively.

Estimate of evolution rate and the time to the most recent common ancestor for

SARS-CoV, MERS-CoV, and SARS-CoV-2

The average Ka, Ks and Ka/Ks for all coding sequences were calculated using

KaKs_Calculator v1.2(28), and the substitution rate and tMRCA were estimated using

BEAST v2.6.2(10). The temporal signal with root-to-tip divergence was visualized in

TempEst v1.5.3(9) using a ML whole genome tree with bootstrap value as input. For

SARS-CoV and SARS-CoV-2, we selected a strict molecular clock and Coalescent

Exponential Population Model. For MERS-CoV, we selected a relaxed molecular

clock and Birth Death Skyline Serial Cond Root Model. We used the tip dates and

chose the HKY as the site substitution model in all these analyses. Markov Chain

Monte Carlo (MCMC) chain length was set to 10,000,000 steps sampling after every

1000 steps. The output was examined in Tracer v1.6

(http://tree.bio.ed.ac.uk/software/tracer/).

Variants calling of SARS-CoV-2 genome sequences

Each genome sequence was aligned to the Wuhan-Hu-1 reference genome

(NC_045512.2) using bowtie2 with default parameters(29), and variants were called

by samtools (sort; mpileup -gf) and bcftoots (call -vm). The merge VCF files were

created by bgzip and bcftools (merge --missing-to-ref)(30, 31).

Phylogenetic tree construction and virus isolates clustering for SARS-CoV-2

After alignment of 624 SARS-CoV-2 genomes with high quality and manually deleted

2 highly divergent genomes (EPI_ISL_415710, EPI_ISL_414690) according to the

firstly constructed phylogenetic tree, the aligned dataset of 622 sequences was

phylogenetically analyzed. The SMS method was used to select GTR+G as the base

substitution model(32), and the PhyML 3.1(12) and MEGA(13) were used to

construct the no-root phylogenetic tree by the maximum likelihood method with the

bootstrap value of 100. The online tool iTOL(33) was used to visualized the

phylogenetic tree. The clusters were defined by the shape of phylogenetic tree.

Detection of specific sites from each Cluster

Information (ID, countries/regions and collection time) and variants (NC_045512.2 as

reference genome) of each genome from each Cluster were extracted. The allele

frequency and nucleotide divergency (pi) for each site in the virus population of each

Cluster were measured by vcftools(34). The *Fst* were also calculated by vcftools(34)

to assess the diversity between the Clusters. Sites with high level of Fst together with

different major allele in each Cluster were filtered as the specific sites. PCA were

analyzed by the GCTA v1.93.1beta(35) with the specific sites and all SNV dataset

respectively.

Linkage analysis of specific sites and characteristics of major haplotype

subgroups

The linkage disequilibrium of the specific sites were analyzed by haploview(36), and

the statistics of the haplotype of the specific sites for each Cluster or country were

used in-house perl script.

Phylogenetic network of haplotype subgroups

The phylogenetic networks were inferred by PopART package v1.7.2(37) using TCS

and minimum spanning network (MSN) methods respectively.

Frequencies of specific sites or haplotypes and correlation with death rate

The frequencies of specific sites for each country were calculated. The death rate was

estimated with Total Deaths /Confirmed Cases based on the data from Johns Hopkins

resources   on   May   12th,   2020

(https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467

b48e9ecf6).The correlation coefficient between death rate and frequencies of specific

site or haplotype in different countries was calculated using Pearson method.

Authors' contributions

HD conceived the study. YM, YB, DJ, JL and XC carried out the data analysis and

wrote the manuscript. MH, SL, and ZC collected data and revised the manuscript.

XW attended the discussions. HD and YM supervised the whole work and revised the

manuscript.

Competing interests

The authors declare no competing interests.

Acknowledgements

This work was supported by the National Key R&D Program of China (2018YFC0910201), the Key R&D Program of Guangdong Province (2019B020226001), the Science and the Technology Planning Project of Guangzhou (201704020176 and 2020Q-P013).

References

1. Coronavirus disease 2019 (COVID-19) Situation Report – 93. WHO. 2020.

2. Liu Y, Gayle AA, Wilder-Smith A, Rocklov J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. J Travel Med. 2020;27(2).

3. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. Lancet. 2020;395(10224):565-74.

4. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science. 2020;367(6483):1260-3.

5. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3.

6. Lam TT, Shum MH, Zhu HC, Tong YG, Ni XB, Liao YS, et al. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. Nature. 2020.

7. Zhang YZ, Holmes EC. A Genomic Perspective on the Origin and Emergence of SARS-CoV-2. Cell. 2020.

8. Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2. Nature Medicine. 2020.

9. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the temporal structure of heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol. 2016;2(1):vew007.

10. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchene S, Fourment M, Gavryushkina A, et al. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. PLoS Comput Biol. 2019;15(4):e1006650.

11. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506.

12. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol. 2010;59(3):307-21.

13. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol. 2018;35(6):1547-9.

14. Guo YR, Cao QD, Hong ZS, Tan YY, Chen SD, Jin HJ, et al. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak - an update on the status. Mil Med Res.

2020;7(1):11.

15.   Yuen K-S, Ye Z-W, Fung S-Y, Chan C-P, Jin D-Y. SARS-CoV-2 and COVID-19: The most
important research questions. Cell Biosci. 2020;10:40.

16.   Chinese SMEC. Molecular evolution of the SARS coronavirus during the course of the SARS
epidemic in China. Science. 2004;303(5664):1666-9.

17.   Zhao Z, Li H, Wu X, Zhong Y, Zhang K, Zhang YP, et al. Moderate mutation rate in the SARS
coronavirus genome and its implications. BMC Evol Biol. 2004;4:21.

18.   Cotten M, Watson SJ, Zumla AI, Makhdoom HQ, Palser AL, Ong SH, et al. Spread, circulation,
and evolution of the Middle East respiratory syndrome coronavirus. mBio. 2014;5(1).

19.   van Dorp L, Acman M, Richard D, Shaw LP, Ford CE, Ormond L, et al. Emergence of genomic
diversity and recurrent mutations in SARS-CoV-2. Infection, Genetics and Evolution. 2020:104351.

20.   Forster P, Forster L, Renfrew C, Forster M. Phylogenetic network analysis of SARS-CoV-2
genomes. Proceedings of the National Academy of Sciences. 2020:202004999.

21.   Holder M, Lewis PO. Phylogeny estimation: traditional and Bayesian approaches. Nat Rev Genet.
2003;4(4):275-84.

22.   Zhang L, Shen F-m, Chen F, Lin Z. Origin and Evolution of the 2019 Novel Coronavirus. Clinical
Infectious Diseases. 2020.

23.   Chan JF, Kok KH, Zhu Z, Chu H, To KK, Yuan S, et al. Genomic characterization of the 2019
novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting
Wuhan. Emerg Microbes Infect. 2020;9(1):221-36.

24.   Graham RL, Sparks JS, Eckerle LD, Sims AC, Denison MR. SARS coronavirus replicase proteins
in pathogenesis. Virus Res. 2008;133(1):88-100.

25.   Li F. Structure, Function, and Evolution of Coronavirus Spike Proteins. Annu Rev Virol.
2016;3(1):237-61.

26.   Malavolta M, Giacconi R, Brunetti D, Provinciali M, Maggi F. Exploring the Relevance of
Senotherapeutics for the Current SARS-CoV-2 Emergency and Similar Future Global Health Threats.
Cells. 2020;9(4).

27.   Liu Y, Du X, Chen J, Jin Y, Peng L, Wang HHX, et al. Neutrophil-to-lymphocyte ratio as an
independent risk factor for mortality in hospitalized patients with COVID-19. J Infect. 2020.

28.   Zhang Z, Li J, Zhao XQ, Wang J, Wong GK, Yu J. KaKs_Calculator: calculating Ka and Ks
through   model   selection   and   model   averaging.   Genomics   Proteomics   Bioinformatics.
2006;4(4):259-63.

29.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9(4):357-9.

30.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map
format and SAMtools. Bioinformatics (Oxford, England). 2009;25(16):2078-9.

31.   Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics (Oxford, England).
2011;27(21):2987-93.

32.   Lefort V, Longueville JE, Gascuel O. SMS: Smart Model Selection in PhyML. Mol Biol Evol.
2017;34(9):2422-4.

33.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments.

Nucleic Acids Res. 2019;47(W1):W256-W9.

34.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format

and VCFtools. Bioinformatics (Oxford, England). 2011;27(15):2156-8.

35.   Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait

analysis. Am J Hum Genet. 2011;88(1):76-82.

36.   Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype

maps. Bioinformatics. 2005;21(2):263-5.

37. Leigh JW, Bryant D, Nakagawa S. popart: full‐feature software for haplotype network

construction. Methods in Ecology and Evolution. 2015;6(9):1110-6.

Figure Legends

Figure 1   Phylogenetic tree and clusters of 622 SARS-CoV-2 genomes

The 622 sequences were clustered into three clusters: Cluster 1 were mainly from North America, Cluster 2 were from regions all over the world, and Cluster 3 were mainly from Europe.

Figure 2   Linkage disequilibrium plot of haplotypes of the 9 specific sites

A. The plot for 622 genome sequences; **B**. The plot for 3624 genome sequences.

Figure 3   The frequencies of both the 9 specific sites and haplotypes

The frequencies of the 9 specific sites (**A**) and haplotypes (**B**) in each country for 3624 genomes.

Figure 4   The characteristics of haplotype subgroups

A. The numbers of haplotypes of the 9 specific sites for 16373 genomes with clear collection data detected in each country in chronological order; **B**. The whole genome mutations in each major haplotype subgroup.

Figure 5   Phylogenetic network of haplotype subgroups for 3624 genomes

The network was inferred by POPART using TCS method. Each colored vertex represents a haplotype, with different colors indicating the different sampling areas. Hatch marks along edge indicated the number of mutations. Small black circles within the network indicated unsampled haplotypes. H1-H5 subgroups were point out according haplotypes of the 9 specific sites, and other small subgroups were not specially pointed out.

Figure 6   The correlation between death rate and frequencies of both the 9

specific sites and haplotypes

Tables

Table 1   Statistics of Ka, Ks and Ka/Ks ratios for all coding regions of the

SARS-CoV, MERS-CoV and SARS-CoV-2 genome sequences

Table 2   Substitution rate and tMRCA estimated by BEAST v2.6.2

Table 3   The information of the 9 specific sites in each Cluster

Table 4   Statistics of Ka, Ks and Ka/Ks ratios for all coding regions of each 4

major haplotype subgroup with 3624 and 16373 genomes respectively

Supplementary material

Figure S1   Regression analyses of the root-to-tip divergence with the

maximum-likelihood phylogenetic tree of SARS-CoV (A), MERS-CoV (B) and

SARS-CoV-2 (C)

The root-to-tip genetic distances were inferred in TempEst v1.5.3. Three data sets

exhibit a positive correlation between root-to-tip divergence and sample collecting

time, and they appear to be suitable for molecular clock analysis.

Figure S2   Substitution rate and tMRCA estimate of SARS-CoV-2

A. Estimated substitution rate of SARS-CoV-2 using BEAST v2.6.2 under a strict

molecular clock. The mean rate was 1.601×10^{-3} (substitutions per site per year). **B**.

The estimated tMRCA of SARS-CoV-2 using BEAST v2.6.2 under a strict molecular

clock. The mean tMRCA was 2019.74 (date: 2019-09-27).

Figure S3   Phylogenetic tree of 622 SARS-CoV-2 genomes

A. Phylogenetic tree constructed by PhyML with bootstrap value of 100; **B**. Phylogenetic tree

constructed by MEGA with bootstrap value of 100.

Figure S4   The PCAs analysis result based on the 9 specific sites (A) and all SNVs

(B)

Table S1   The information of the first downloaded complete genome sequences

Table S2   The sequence IDs in each Cluster

Table S3   The details information of the SNVs in all Clusters

Table S4   The information of death rate, frequencies of the 9 specific sites and

major haplotypes in each country for 3624 and 16373 genomes respectively

Table S5   All haplotypes of the 9 specific sites and numbers of them for 3624 and

16373 genomes respectively

Table S6   Whole genome mutations and frequencies in each major haplotype

subgroup of 3624 and 16373 genomes respectively

Tree scale: 0.0001

Inner Color: Continents
- Asia
- Europe
- NorthAmerica
- Oceania
- SouthAmerica

Shape Plot: Haplotype
- H1
- H2
- H3
- H4
- H5
- H6
- Other

Outer Clolors: Countries
- Australia
- Belgium
- Brazil
- Cambodia
- Canada
- Chile
- China
- Denmark
- Finland
- France
- Georgia
- Germany
- HongKong_China
- Hungary
- India
- Ireland
- Italy
- Japan
- Kuwait
- Luxembourg
- Mexico
- Nepal
- Netherlands
- NewZealand
- Poland
- Portugal
- SaudiArabia
- Singapore
- SouthKorea
- Sweden
- Switzerland
- Taiwan_China
- Thailand
- UnitedKingdom
- UnitedStates
- Vietnam

Cluster 1

Cluster 2

Cluster 2

Cluster 3

		Ka (mean) 10^{-3}	s.e. (Ka) 10^{-3}	Ks (mean) 10^{-3}	s.e. (Ks) 10^{-3}	Ka/Ks (mean)	s.e. (Ka/Ks)	Hypothesis	P-value* 622 genomes	P-value* 3624 genomes
SARS-CoV		0.985	0.018	1.310	0.049	0.760	0.038	Ka/Ks (SARS-CoV < SARS-CoV-2)	0.139	0.084
MERS-CoV		1.319	0.040	4.887	0.096	0.260	0.003	Ka/Ks (MERS-CoV < SARS-CoV-2)	0.000	0.000
SARS-CoV-2	622 genomes	0.231	0.004	0.265	0.005	1.008	0.020			
	3624 genomes	0.287	0.002	0.298	0.002	1.094	0.010			

Note: *One-sided Mann-Whitney U-test for the means of two independent samples.

	substitution rate (10^{-3})	95% CI (10^{-3})	tMRCA	95% CI	references
SARS-CoV	1.050	0.489, 1.654	2002/6/28	2002/1/19, 2002/11/3	this study
	-	0.800, 2.380	spring of 2002	-	[17]
MERS-CoV	1.516	1.392, 1.632	2012/6/7	2012/6/4, 2012/6/9	this study
	1.120	0.876, 1.370	2012/3	2011/12, 2012/6	[18]
SARS-CoV-2	1.601	1.418, 1.796	2019/9/27	2019/8/28, 2019/10/26	this study

17. Zhao Z, Li H, Wu X, Zhong Y, Zhang K, Zhang YP, et al. Moderate mutation rate in the SARS coronavirus genome and its implications. BMC Evol Biol. 2004;4:21.

18. Cotten M, Watson SJ, Zumla AI, Makhdoom HQ, Palser AL, Ong SH, et al. Spread, circulation, and evolution of the Middle East respiratory syndrome coronavirus. mBio. 2014;5(1).

os	Ref	Alt	Cluster 1			Cluster 2			Cluster 3			Fst	Gene region	Mutation type	Protein changed	codon changed	Impact
			pi	Major allele	frequency	pi	Major allele	frequency	pi	Major allele	frequency						
1	C	T	0.0000	C	1.0000	0.0109	C	0.9945	0.0000	T	1.0000	0.991240	5'UTR	upstream	NA	NA	MODIFIE
)37	C	T	0.0000	C	1.0000	0.0109	C	0.9945	0.0000	T	1.0000	0.991240	gene-orf1ab	synonymous	924F	2772ttC>ttT	LOW
'82	C	T	0.0000	T	1.0000	0.3863	C	0.7390	0.0000	C	1.0000	0.582113	gene-orf1ab	synonymous	2839S	8517agC>agT	LOW
408	C	T	0.0000	C	1.0000	0.0055	C	0.9973	0.0000	T	1.0000	0.995609	gene-orf1ab	missense	4715P>L	14144cCt>cTt	MODERA
'747	C	T	0.0735	T	0.9620	0.0000	C	1.0000	0.0000	C	1.0000	0.975245	gene-orf1ab	missense	5828P>L	17483cCt>cTt	MODERA
'858	A	G	0.0497	G	0.9747	0.0000	A	1.0000	0.0000	A	1.0000	0.983573	gene-orf1ab	missense	5865Y>C	17594tAt>tGt	MODERA
060	C	T	0.0000	T	1.0000	0.0164	C	0.9918	0.0000	C	1.0000	0.976100	gene-orf1ab	synonymous	5932L	17796ctC>ctT	LOW
403	A	G	0.0000	A	1.0000	0.0164	A	0.9918	0.0000	G	1.0000	0.986895	gene-S	missense	614D>G	1841gAt>gGt	MODERA
144	T	C	0.0000	C	1.0000	0.3915	T	0.7335	0.0000	T	1.0000	0.578530	gene-ORF8	missense	84L>S	251tTa>tCa	MODERA

		Ka (mean) 10^{-3}	s.e. (Ka)10^{-3}	Ks (mean) 10^{-3}	s.e. (Ks)10^{-3}	Ka/Ks (mean)	s.e. (Ka/Ks)	Hypothesis	P-value*
	H1	0.510	0.057	0.498	0.056	1.099	0.010	Ka/Ks (H3 < H1)	0.000
	H2	0.347	0.042	0.334	0.044	1.268	0.023	Ka/Ks (H3 < H2)	0.000
3624 genomes	H3	0.298	0.007	0.397	0.009	0.795	0.010	-	-
	H4	0.334	0.023	0.414	0.019	0.796	0.019	-	-
	H1	0.416	0.011	0.384	0.011	1.178	0.005	Ka/Ks (H3 < H1)	0.000
	H2	0.332	0.014	0.316	0.014	1.224	0.012	Ka/Ks (H3 < H2)	0.000
16373 genomes	H3	0.305	0.004	0.401	0.005	0.809	0.007	-	-
	H4	0.354	0.010	0.477	0.009	0.749	0.010	-	-

Note: *One-sided Mann-Whitney U-test for the means of two independent samples.