Overcoming bias in gene-set enrichment analyses of brain-wide transcriptomic data.

Ben D. Fulcher[1,2,✉], **Aurina Arnatkevičiūtė**[2], and **Alex Fornito**[2]

[1]School of Physics, The University of Sydney, Camperdown, NSW 2006, Australia.
[2]The Turner Institute for Brain and Mental Health, School of Psychological Sciences, and Monash Biomedical Imaging, Monash University, Victoria, Australia.

The recent availability of whole-brain atlases of gene expression, which quantify the transcriptional activity of thousands of genes across many different brain regions, has opened new opportunities to understand how gene-expression patterns relate to spatially varying properties of brain structure and function. To aid interpretation of a given neural phenotype, gene-set enrichment analysis (GSEA) has become a standard statistical methodology to identify functionally related groups of genes, annotated using systems such as the Gene Ontology (GO), that are associated with a given phenotype. While GSEA has identified functional groups of genes related to diverse aspects of brain structure and function in mouse and human, here we show that these results are affected by substantial statistical biases. Quantifying the false-positive rates of individual GO categories across an ensemble of completely random phenotypic spatial maps, we found an average 875-fold inflation of significant findings relative to expectation in mouse, and a 582-fold inflation in human, with some categories being judged as significant for over 20% of random phenotypes. Concerningly, the probability of a GO category being reported as significant in the extant literature increases with its estimated false-positive rate, suggesting that published reports are strongly affected by the reporting of false-positive bias. We show that the bias is primarily driven by gene–gene coexpression and spatial autocorrelation in transcriptional data, which are not accounted for in conventional GSEA nulls, and we introduce flexible ensemble-based null models that properly account for these effects. Using case studies of structural connectivity degree in mouse and human, we demonstrate that many GO categories that would conventionally be judged as highly significant are in fact consistent with ensembles of random phenotypes. Our results highlight major pitfalls with applying standard GSEA to brain-wide transcriptomic data and outline a solution to this pervasive problem, which is made available as a toolbox.

transcriptome | gene ontology | gene enrichment | genome | MRI
Correspondence: *ben.fulcher@sydney.edu.au*

Introduction

The brain's multi-scale organization spans at least five orders of magnitude in space and time (1, 2). Understanding how these distinct scales relate to each other has proven challenging, largely because typical assays face a trade-off between spatial resolution and anatomical coverage. For example, *in vivo* brain-imaging techniques like magnetic resonance imaging (MRI) afford an unparalleled capacity to measure diverse aspects of structure and function across the entire brain, but have a limited spatial resolution that rarely surpasses $1\,\mathrm{mm}^3$. By contrast, invasive physiological recordings enable the measurement of neural structure and dynamics with cellular resolution, but traditionally only within small subregions of the brain. In recent years, our capacity to understand variations in molecular function across the entire brain has been greatly enhanced by industrialized, high-throughput transcriptome profiling, which has yielded genome-wide quantification of expression levels across the entire brain (3, 4). Two of the most influential genome-wide resources are the Allen Human Brain Atlas (AHBA), which encompasses microarray measurements of expression in $> 20\,000$ genes across 3702 tissue samples from six postmortem brains (4), and the Allen Mouse Brain Atlas (AMBA), which comprises *in situ* hybridization measures of expression for $> 17\,000$ genes at cellular resolution (3).

The availability of these genome-wide expression atlases has generated new opportunities to bridge spatial scales and uncover the microscale molecular correlates of macroscale brain organization. This correspondence has commonly been assessed by comparing regional variations in molecular function (from gene-expression maps) to independently measured macroscale structural and functional properties, e.g., from imaging techniques like magnetic resonance imaging (MRI) (5, 6). Prior work has characterized a relationship between gene expression and structural connectivity in *Caenorhabditis elegans* (7–10), mouse (11–18), and human (19, 20). Human research has further characterized links between gene-expression patterns and correlations of neural dynamics (functional connectivity) estimated from functional MRI (fMRI) (21–23) and Electrocorticography (ECoG) (24); brain morphometry and microstructure (25–30); neurotransmitter receptor densities (31); and disease-related changes in brain structure and function (32–35). In these analyses, genes are typically scored according to the correspondence between their spatial ex-

pression map and anatomical variations of the independently measured phenotype, most commonly quantified as a simple correlation across space. Instead of performing inference on thousands of genes directly, statistical tests can be performed at the level of groups of functionally annotated sets of genes, such as those involved in different types of biological processes. This process of gene-set enrichment analysis (GSEA) allows one to investigate which gene categories are preferentially related to a spatial brain phenotype of interest, reducing the genome-wide multiple comparison burden and sometimes facilitating biological interpretation. GSEA uses a statistical hypothesis-testing framework to assess which categories of genes are most strongly related to a given phenotype, leveraging annotations of genes to categories from open ontologies like the Gene Ontology (GO) (36) and KEGG (37). GSEA has thus been applied extensively across species, scales, and diverse aspects of brain structure and function to gain insight into the biological processes that are mediated by genes with similar spatial expression profiles.

In the brain-imaging literature, a wide variety of publicly available tools have been used to perform GSEA (38–47). While software packages make enrichment analysis easy to run, there are numerous challenges to correctly interpreting the results of GSEA. This requires careful consideration of: (i) the reliability of gene annotations (including the rate of false-positive annotations (48)); (ii) the reproducibility of results (different software packages use different methods to perform enrichment with respect to the GO hierarchy; both annotations and GO terms are updated daily (49)); and (iii) the statistical inference procedure (the design of multiple-hypothesis testing across many non-independent and often hierarchically structured tests (48, 50)). Methodological variability can lead to vast discrepancies in results obtained using different software packages. For example, one test using identical inputs to different software packages, demonstrated a variation in p-values for some GO categories that spanned several orders of magnitude (48). Methodological development of GSEA is ongoing (51, 52).

Applications of GSEA, from the interpretation of genome-wide association study (GWAS) findings (53) to case–control comparisons of microarray data, are subject to the general statistical challenges outlined above. But the application of GSEA to whole-brain transcriptional atlas data is associated with unique challenges due to the spatial embedding of the atlas data. First, spatial embedding introduces coexpression between genes with similar spatial patterning of expression (4, 54, 55). This induces dependences between genes, violating the null assumption of independence in a way that is a generic characteristic of the expression dataset and not specific to the particular phenotype being studied. Second, transcriptional maps are strongly spatially autocorrelated, such that nearby anatomical regions have more similar

patterns of gene expression than distant regions, as has been observed in head neurons of *C. elegans* (10), the mouse brain (12, 18), and human cortex (5, 6, 54, 56–58). The neural phenotypes that are matched to gene-expression atlas data are also commonly spatially autocorrelated (10, 18, 59–63). Two spatial maps with similar spatial autocorrelation structure have a greater chance of exhibiting a high correlation to each other than two random spatial maps. This issue has been highlighted previously in neuroimaging, with researchers developing methods to better estimate null distributions in the presence of spatial autocorrelation, e.g., using spatial permutation methods like spatial-lag models (30, 64) and spin tests (65) to test against an ensemble of surrogate spatial maps, or by removing the effect of physical distance through regression (5, 12, 14, 16–18, 24, 66). These prominent factors affecting correlational analyses involving spatially embedded phenotypes are yet to be characterized in the context of GSEA.

Here we characterize the statistical biases involved in applying standard GSEA methodologies to brain-wide transcriptomic data. We demonstrate that the rate at which a GO category is judged as significantly correlated to a random phenotype (category false-positive rate, CFPR) is far higher than statistical expectation, exceeding 20% for some GO categories. We show that the literature is compatible with the reporting of this false-positive bias, finding a progressive increase in the reporting of GO categories with their false-positive rate under random phenotypes. These biases can be overcome using new ensemble-based null models that assess statistical significance relative to ensembles of randomized phenotypes (rather than the conventional randomized-gene null). Using case studies applying GSEA to structural connectivity degree (the number of inter-regional connections attached to a brain region) in mouse and human brains, we show that highly significant categories under conventional GSEA are often consistent with ensembles of randomized phenotypes. Our ensemble-based approach to GSEA overcomes biases in investigating the transcriptomic correlates of spatially varying neural phenotypes, and is made available as a Matlab toolbox.

Results

We first describe a typical GSEA pipeline applied to a spatial brain phenotype (SBP) of interest, depicted in Fig. 1A. The SBP is a spatial map of some measurement taken across brain areas, such as gray-matter volume, functional connectivity strength, or case–control differences in some neural property. Each gene in the atlas is scored using a measure of similarity between its expression pattern and the SBP (e.g., as a correlation coefficient), yielding high scores for genes with a spatial patterning resembling the SBP. GSEA then tests whether high-scoring genes are concentrated in particular types of GO categories by comparing the GO-category scores obtained from the data to those obtained from a null

model. Current GSEA analysis uses a 'random-gene null' to assess whether a GO category's score is higher than would be expected if genes were annotated to GO categories randomly (48). This is similar to shuffling gene identities, as depicted in Figs 1B and C. This general procedure captures the essence of applying GSEA to expression-atlas datasets. While here we focus just on this type of spatial comparison, we note that some studies instead focus on pairwise inter-regional phenotypes, such as the presence of a connection between brain areas and a measure of correlated gene expression (6, 18, 21). Throughout this work, we perform GSEA (67) on biological process GO categories (36) using brain-wide gene-expression data from the AMBA (3) for the mouse brain, and the AHBA (4) for the human cortex. As described in Methods, enrichment is performed using gene-score resampling (38) using open Matlab software that we developed. Given the sensitivity of GSEA results to specific methodological pipelines (48), our aim here is to highlight general issues with applying any GSEA pipeline to spatially embedded transcriptional-atlas data. Thus, while the quantitative results presented here vary across different GSEA packages and parameters, the statistical biases we characterize will apply to all current GSEA pipelines that assess significance relative to random-gene null models.

Diverse phenotypes in the literature are enriched for similar GO categories We were motivated to undertake this study because we observed a strong similarity in the results of GSEA analyses in the literature, despite these analyses involving very different phenotypes, species, measurement modalities, gene-expression processing pipelines, and software implementations of GSEA. To evaluate this consistency, we surveyed 16 mouse GSEA analyses (taken from eight different studies) and 60 human GSEA analyses (from 23 different studies). Despite investigating very different phenomena, these studies reported similar sets of significant GO categories, most frequently implicating categories related to metabolic, neuronal, and generic biological and behavioral processes. The most reported GO category in our survey was 'chemical synaptic transmission' (GO:000726), which has been reported in 15 human analyses from 10 different studies (21, 26, 32, 70, 73–78) and four different mouse analyses from three different studies (14, 17, 79). These studies implicate genes involved in chemical synaptic transmission in the organization of human resting-state functional connectivity (21), human adolescent cortical shrinkage and myelination (26), and tract-traced structural connectivity in the mouse brain (17). Some other of the most commonly reported categories include 'potassium ion transmembrane transport' (22, 26, 32, 68–70), 'learning or memory' (15, 70–73), and 'electron transport chain' (18, 19, 23, 26, 70) (see Supplementary File `LiteratureAnnotations_p005.csv` for a sorted list of GO categories, annotated by studies). All of these bi-

A Scoring genes and GO categories

Fig. 1. Pipeline for applying gene set enrichment analysis (GSEA) to brain-wide expression atlas data. **A** Given a phenotype map, we first compute the spatial correlation coefficient between that map and each gene. These gene scores are then agglomerated at the level of categories using an annotation system like the Gene Ontology (GO). For continuous scores, agglomeration is typically performed as the mean score of genes annotated to a category. **B** Statistical significance of a GO category is assessed relative to the random-gene null, which estimates a null distribution for each GO category by annotating genes to GO categories at random. **C** For every GO category, a p-value is estimated using a permutation test, by comparing the GO category score obtained from the real data to the null distribution.

ological processes are sufficiently broad to be plausibly linked to any type of brain-related phenotype, but we aimed to investigate whether the consistency of these findings might instead be driven by common statistical biases in GSEA that favor the selection of some GO categories over others.

The GO enrichment signature of randomized spatial maps We first tested for false-positive bias in GSEA by characterizing its results when applied to purely random SBPs, generated by assigning a random number to each brain area independently. We performed GSEA separately for each of 10 000 random SBPs, noting which GO categories were significantly related to each SBP (false discovery rate, FDR < 0.05). We then computed the category false-positive rate (CFPR) for each GO category as the proportion of random SBPs for which a statistically significant spatial correlation was identified. This method of computing CFPRs under random SBPs is depicted schematically in Fig. 2A(ii), labeled 'SBP-random'. As random SBPs are uninformative, an unbiased GSEA procedure should produce CFPRs consistent with the expected statistical false-positive level,

Fig. 2. Some GO categories have far higher false-positive rates under randomized spatial phenotypes than statistical expectation, and these GO categories are more likely to be reported as significant in existing literature studies. A A schematic of three GSEA analyses involving correlations between: (i) 'reference'—an ensemble of random phenotypes and randomized gene-expression data (green), (ii) 'SBP-random'—an ensemble of random phenotypes and real gene-expression data (red), and (iii) 'SBP-spatial'—an ensemble of spatially autocorrelated phenotypes and real gene-expression data (blue). For the human cortex, examples of spatial maps in each ensemble are plotted; relative to the SBP-random maps that have no spatial correlation structure, the SBP-spatial maps are more likely to have similar values in nearby locations. **B** Distributions of the category false-positive rate (CFPR) across all GO categories are shown as violin plots in mouse. Across an ensemble of 10 000 SBP-random or SBP-spatial maps, the CFPR is computed for each GO category as the proportion of phenotypes for which that GO category was found to be significant. Results are shown for the three analyses depicted in A: (i) 'reference' (green); (ii) 'SBP-random' (red); and (iii) 'SBP-spatial' (blue). Note the logarithmic vertical scale (and therefore exclusion of GO categories with CFPR = 0). **C** The proportion of literature-reported GO categories increases with the false-positive rate expected under ensembles of random phenotypes. GO categories were labeled from a literature surveys of GSEA analyses using atlas-based transcriptional data in human and mouse (see Supplementary Information for survey details). Across eight equiprobable bins of CFPR (i.e., each bin contains the same number of GO categories), we plot the proportion of all literature-reported GO categories that are contained in that bin. Results are shown for the SBP-random (red) and SBP-spatial (blue) ensembles in mouse (dotted) and human (solid). The position of each bin is shown as the mean of its extremities.

and all GO categories should have similar CFPRs. We estimated this statistically expected category-wise false-positive level numerically using a 'reference' case, shown in Fig. 2A(i), in which each gene's expression data is independently randomized. As this randomization destroys gene–gene coexpression structure in the gene-expression matrix, it allows us to isolate the contribution of non-random gene-expression structure to CFPRs through comparison to the 'SBP-random' results. Motivated by the strong spatial autocorrelation observed in many real SBPs (6), we also tested whether the use of spatially autocorrelated random phenotypes would affect CFPRs. These spatially autocorrelated SBPs, labeled 'SBP-spatial' [Fig. 2A(iii)], were generated using a spatial-lag model (80), with parameters determined from the spatial autocorrelation properties of the gene expression data (following Burt et al. (30), see Methods for details).

Distributions of non-zero CFPR across GO categories are shown for the three experiments in Fig. 2B for mouse (note the logarithmic scale); extremely similar quanti-

tative results were obtained in human (Fig. S1). In the 'reference' case, consistent with the lack of signal from randomized gene expression and random phenotypes, we observed very low CFPR across our ensemble of 10 000 random SBPs. The vast majority of GO categories (84% in mouse and human) received zero significant hits across the entire ensemble and the average CFPR was 0.002% (in both mouse and human). Furthermore, in both mouse and human, no individual GO category received more than three significant hits (maximal CFPR = 0.03%).

When repeating the same procedure with the same random SBPs but now using real gene-expression data ('SBP-random', Fig. 2A), 95% of GO categories increased their CFPR (in mouse; 92% in human), and the GO category with maximal CFPR jumped dramatically from the reference level of 0.03% to 23% (mouse) and 25% (human). The mean CFPR across all GO categories increased 875-fold in mouse (from 0.002% to 1.5%), and 582-fold in human (from 0.002% to 1.0%). When repeating the same procedure but now using random phe-

GO Category	Mouse CFPR [SBP-spatial] (%)	Human CFPR [SBP-spatial] (%)	References
Regulation of synaptic plasticity	23 [37]	10 [7]	(18, 69–71, 73)
Regulation of postsynaptic membrane potential	19 [35]	10 [17]	(18, 22, 70)
Glutamate receptor signaling pathway	16 [34]	6 [14]	(70–72, 79)
Respiratory electron transport chain	12 [4]	20 [30]	(18, 19, 23, 26, 70)
Learning	20 [34]	5 [2]	(70, 71, 73, 78)

Table 1. GO categories with high category false-positive rates (CFPRs) are predominantly related to neuronal and metabolic biological function in mouse and human. We list selected GO categories with amongst the highest CFPRs across mouse and human, across ensembles of spatially independent random phenotypes (SBP-random) and spatially autocorrelated random phenotypes (SBP-spatial). CFPR is listed for SBP-random [and for SBP-spatial in brackets]. A full ordered list is in Supplementary File `CFPRTable.csv`.

notypes with a preserved spatial autocorrelation structure ('SBP-spatial'), most GO categories decreased their CFPR (60% in mouse and 68% in human) but CFPRs increased on average, from 1.5% to 2.8% in mouse (maximal CFPR increased from 23% to 37%) and from 1.0% to 2.2% in human (maximal CFPR increased from 25% to 36%). The differences in results between these ensembles will be explained later through a more detailed investigation of particular GO categories.

Performing conventional GSEA on transcriptomic data thus yields significant false-positive relationships to SBPs made up of random numbers far beyond statistical expectation, indicating a striking methodological bias. This false-positive bias does not affect all categories equally: some GO categories exhibit reference-consistent CFPR ($\leq 0.03\%$) while others are flagged as significantly correlated to random spatial maps more than 20% of the time. GO categories with the highest CFPRs are dominated by brain-related functions, including neuronal, metabolic, and behavioral categories. A selection is listed in Table 1 along with literature studies that have reported their significant enrichment (see Supplementary File `CFPRTable.csv` for full list).

To assess whether GO categories commonly reported as significant in the published literature might be affected by this false-positive bias, we binned GO categories by their CFPR, placing an equal number of categories in each bin. We then analyzed the distribution of literature-reported GO categories across these bins by computing the proportion of all literature-significant categories in each bin. As shown in Fig. 2C, for both ensembles of null phenotypes and in both mouse and human, we find a progressive increase in the literature reporting of a GO category's significance with its CFPR. Another analysis of frequently reported GO categories, that incorporated information about the number of literature studies that have reported a given GO category, confirmed a similar increase with CFPR (Fig. S2). The increase in the reporting of GO categories with their false-positive rate under ensembles of randomized phenotypes suggests that the existing literature is heavily affected by the false-positive biases of conventional GSEA methodology.

Within-category coexpression drives false-positive bias
Since randomizing the gene coexpression structure ('ref-

erence') dramatically reduces CFPRs, we reasoned that dependencies between genes within a GO category must drive the vast differences in CFPR between categories. To investigate this possibility, we selected two similarly-sized GO categories in mouse that have different levels of within-category coexpression: 'regulation of dendritic spine morphogenesis' (RDSM, 40 genes), and 'zymogen activation' (ZA, 42 genes). We aimed to understand why, under the SBP-random ensemble, RDSM (CFPR = 13%) has a higher CFPR than ZA (0.07%). Region × gene expression matrices and gene × gene coexpression matrices are plotted for RDSM and ZA in Figs 3A and B, respectively. Compared to a representative random set of 40 genes, shown in Fig. 3C, we find a much more consistent spatial patterning of gene expression in the RDSM category, resulting in increased within-category gene × gene coexpression. This spatial coherency of expression patterning is consistent with genes associated with regulating dendritic-spine morphogenesis (RDSM) having a coordinated brain function that varies characteristically across brain regions. By contrast, GO categories that play a minimal or less-specific role in brain function, such as zymogen activation (ZA), exhibit noisier expression patterns and lower within-category coexpression (that is visually similar to that of a random set of genes).

To understand how these differences in within-category coexpression affect category false-positive rates, we investigated the distribution of mean correlation coefficients between the expression profiles of genes in each category and the ensembles of randomized SBPs analyzed above. These category-score distributions are plotted for SBP-random and SBP-spatial ensembles in Figs 3D and E, respectively. A category with wider distributions than a random set of genes (the null comparison used in GSEA) will have a greater probability of obtaining a significant correlation to that ensemble of phenotypes. First we note that, consistent with both SBP-random and SBP-spatial ensembles containing no information about gene expression, all category-score distributions are symmetric about zero. Under random phenotypes (Fig. 3D), RDSM has a wider distribution of category scores than ZA; it is more likely to exhibit a higher correlation to a random SBP. This widening is driven by the increased coexpression of RDMS genes,

Fig. 3. GO categories with high within-category gene–gene coexpression are most prone to false-positive enrichment. We plot expression (brain area × gene) and coexpression (gene × gene) heat maps for two example GO categories in the mouse brain: **A** 'regulation of dendritic spine morphogenesis' (40 genes); and **B** 'zymogen activation' (42 genes); as well as **C** a random set of 40 genes, for comparison. Each gene's expression is normalized (low to high) for visualization purposes. Genes annotated to 'regulation of dendritic spine morphogenesis' display a more characteristic spatial patterning and hence have higher coexpression. Distributions of each category's score (mean correlation between the genes in that category and a phenotype) across an ensemble of null phenotypes, are plotted as violin plots for: **D** the SBP-random ensemble of random-number phenotypes (cf. Fig. 2A(ii)), and **E** the SBP-spatial ensemble of random spatially autocorrelated phenotypes (cf. Fig. 2A(iii)). The mean of each distribution is annotated with a horizontal line.

such that a chance correlation between a random SBP and any single RDMS gene is amplified through a similar correlation with many other genes in the category. By contrast, categories like ZA contain genes with lower gene–gene coexpression, such that a chance correlation between an SBP and a given ZA gene is more likely to be cancelled out by a chance correlation in the opposite direction for another gene, driving category scores towards zero and resulting in a narrower category-score distribution.

In estimating category p-values, conventional GSEA compares the category score to that of a random set of the same number of genes, shown gray in Fig. 3D (where we have performed many randomizations of sets of 40 genes so as not to bias towards any particular random set). Due to the high coexpression of RDSM genes, this occurs for approximately for 13% of SBP-random phenotypes, but for ZA, which has a similar coexpression structure as random genes, this occurs for just 0.07% of random SBPs.

Having developed some intuition for the mechanism through which within-category coexpression can bias towards more extreme null category scores (and hence greater CFPRs), we next tested whether this effect holds across all GO categories. We defined a simple measure of within-category coexpression, $\langle r \rangle$, as the mean coexpression across all pairs of genes in a GO category (the mean of the upper diagonal of coexpression matrices plotted in Figs 3A and B). GO categories that contain genes with very similar spatial patterning across the brain (like RDSM) have higher $\langle r \rangle$ than categories with more diverse gene-expression patterns (like ZA). We found that GO categories with the highest within-category coexpression, $\langle r \rangle$, are strongly related to metabolic and neuronal functioning, in both mouse and human. For example, amongst the GO categories

with the highest $\langle r \rangle$ are: 'regulation of short-term neuronal synaptic plasticity' ($\langle r \rangle_{\text{mouse}} = 0.38$, $\langle r \rangle_{\text{human}} = 0.29$) and 'ATP synthesis coupled proton transport' ($\langle r \rangle_{\text{mouse}} = 0.34$, $\langle r \rangle_{\text{human}} = 0.29$) (see Supplementary File `WithinCategoryCoexp.csv` for a full list). As shown in Fig. 4A, $\langle r \rangle$ and CFPR (using the SBP-random ensemble) are positively correlated in both mouse and human. This relationship clearly demonstrates that gene–gene dependencies play a role in driving false-positive bias when applying GSEA to transcriptomic data. Brain-related categories of genes exhibit high within-category coexpression and thus high CFPRs, due to their characteristic spatial patterning across the brain. This relationship is a key factor behind the deceptively sensible brain-related enrichment results of published GSEAs across diverse brain phenotypes.

The role of spatial autocorrelation We next turn to the role of spatial autocorrelation in SBPs by investigating the SBP-spatial ensemble of spatially autocorrelated phenotypes. Relative to the SBP-random ensemble (of purely random phenotypes), the CFPR of RDSM increased under the SBP-spatial ensemble: $13\% \rightarrow 28\%$, whereas the CFPR for ZA decreased: $0.07\% \rightarrow 0.02\%$. Examining the category-score distributions in Fig. 3E, we first note that they are all much wider than for the SBP-random ensemble (Fig. 3D), including for random sets of genes. This is due to the predominance of spatial autocorrelation in the expression patterns of individual genes: a gene that exhibits a strongly autocorrelated expression map is more likely to be strongly correlated to a spatially autocorrelated SBP than a random spatial map. Relative to random genes, we see a wider category-score distribution for RDSM, and a narrower distribution for ZA, consistent with their CFPRs. Imposing the constraint of spatial autocorrelation (i.e., SBP-rand $\rightarrow$ SBP-spatial) can either increase a category's CFPR (if

Fig. 4. Category false-positive significance rates (CFPRs) vary with within-category gene–gene coexpression and spatial autocorrelation in mouse and human. **A** CFPR (%) computed from the SBP-rand ensemble of random spatial maps, increases with a measure of mean within-category coexpression, $\langle r \rangle$, across ten equiprobable bins in mouse (blue) and human (green). The extent of each bin is displayed as a horizontal line. **B** The percentage of GO categories that exhibited an increase in CFPR when using the SBP-spatial ensemble relative to the SBP-random ensemble, across ten equiprobable bins of the spatial autocorrelation score, $R^2_{\exp}$. This score captures the goodness of fit of each GO category's correlated gene expression to an exponential function with distance. More spatially autocorrelated GO categories are more likely to exhibit an increase in CFPR for spatially autocorrelated phenotypes (the SBP-spatial ensemble). The average value across all GO categories is shown as a horizontal dotted line.

it exhibits a more similar spatial autocorrelation structure to the SBP-spatial ensemble) or decrease it (if it exhibits a less similar spatial autocorrelation structure). To verify this reasoning, we investigated whether GO categories containing genes with high average spatial autocorrelation are more likely to increase their CFPR in the SBP-spatial ensemble relative to the SBP-random ensemble. We computed a simple measure of spatial autocorrelation for each GO category, $R^2_{\exp}$, that measures how well each category's correlated gene expression is fit to an exponential decay with distance (see Methods). Higher values of $R^2_{\exp}$ are given to GO categories containing genes that exhibit a stronger exponential distance-dependence of their gene-expression patterns. We then tested whether categories with high $R^2_{\exp}$ were more likely to exhibit an increase in CFPR when adding a spatial autocorrelation constraint to the null ensemble (i.e., using SBP-spatial rather than SBP-random). As shown in Fig. 4B, categories containing genes with stronger spatial autocorrelation scores, $R^2_{\exp}$, are more likely to increase their CFPR under the SBP-spatial ensemble. This effect is strongest for categories with an autocorrelation length scale close to that of the SBP-spatial ensemble (Fig. S3), confirming the intuition that a high correlation between a GO category and phenotype is more likely for categories of genes with a similar spatial autocorrelation structure to the phenotype. As this effect is not taken into account by conventional GSEA, it can drive increased CFPR.

Ensemble-based null models for spatial expression data. The above results reveal clear biases in the application of conventional GSEA to transcriptional-atlas data, with a strong false-positive bias in categories of genes that exhibit high within-category correlation, and those which display a similar spatial autocorrelation structure to any given phenotype. Since brain-related GO categories are characterized by high within-category coexpression and strong spatial autocorrelation, this false-

positive bias misleadingly favors brain-related GO categories. These methodological issues highlight the need for new null models to enable valid inference and interpretation of GSEA for spatially embedded transcriptional data.

In introducing new null models, we imagine a researcher who wishes to investigate the transcriptional correlates of their spatial brain phenotype, asking: *'Which GO categories contain genes that are significantly correlated to my phenotype?'* The conventional GSEA methodology evaluates the significance of a GO category against a null of random genes, depicted in Fig. 5A (cf. Fig. 1B). This generates null samples by randomizing the annotations of genes to GO categories, and asks the question of each GO category: *'Are genes in this category more strongly correlated to my phenotype than a random set of genes?'* Through the mechanisms described above, this randomization of gene-to-category annotations destroys within-category gene–gene coexpression structure, thereby driving high CFPRs (particularly for brain-related categories) characterized above.

To overcome this bias, we introduce a new way of performing GSEA that, instead of estimating statistical significance relative to randomized gene-to-category assignments, estimates the significance of a given SBP relative to ensembles of randomized SBPs. Two such ensembles of randomized are the the SBP-random and SBP-spatial ensembles defined above (Fig. 2A). By estimating each GO category's null distribution relative to an ensemble of randomized phenotypes, ensemble-based nulls test whether a given SBP is significantly related to a GO category beyond what would be expected from the null phenotype ensemble. For example, in testing a measured SBP relative to the SBP-random ensemble, the researcher can ask the question of each GO category: *'Are genes in this GO category more correlated to my phenotype than they would be to a random phenotype?'*, shown in Fig. 5B. The major biases with GSEA characterized above, are associated with the random-gene null

A Random-gene
'Are genes in this category more correlated to my phenotype than a random set of genes?'

B SBP-random
'Are genes in this GO category more correlated to my phenotype than they would be to a random phenotype?'

C SBP-spatial
'Are genes in this GO category more correlated to my phenotype than they would be to a random spatially autocorrelated phenotype?'

Fig. 5. The statistical significance of a GO category can be quantified relative to conventional random-gene nulls, or ensemble-based null models introduced here. For a given spatial brain phenotype (SBP) of interest, we depict the process through which null samples are generated for estimating statistical significance for GSEA across three different null models. **A** The conventional random-gene null tests whether the observed result is more extreme than if genes were assigned to GO categories at random (similar to the illustrated shuffling of gene identities). As this destroys within-category gene–gene correlation structure, it leads to high category false-positive rates for random phenotypes. An alternative is to compute null distributions for each category based on an ensemble of null phenotypes: **B** The SBP-random null tests whether the observed result is more extreme than if the phenotype of interest was a random spatial map. **C** The SBP-spatial null tests whether the observed result is more extreme than if the phenotype of interest was a random spatially autocorrelated map.

destroying the within-category coexpression structure, but are preserved in ensemble-based nulls. This ensures that enrichment results stemming from ensemble-based nulls cannot be explained by variations in gene–gene coexpression structure between GO categories (which are a generic property of the transcriptional atlas, independent of the phenotype of interest). Ensemble-based nulls also allow for additional constraints (e.g., spatial autocorrelation) to be incorporated straightforwardly. For example, testing against the SBP-spatial ensemble allows the researcher to ask: *'Are genes in this GO category more correlated to my phenotype than they would be to a random spatially autocorrelated phenotype?'*, shown in Fig. 5C. While here we use the generic SBP-spatial ensemble, the spatial autocorrelation properties of the ensemble could be fitted to match the phenotype of interest, to ensure that the spatially autocorrelated ensemble exhibits a similar spatial autocorrelation structure as the phenotype of interest (e.g., using the *BrainSMASH* package (64)).

Case study: Transcriptional correlates of structural connectivity degree. To demonstrate how the different null models, including the new ensemble-based nulls, can be applied in real settings, we present a demonstrative case study in mouse and human. We aimed to uncover whether particular classes of genes have an expression pattern that is significantly correlated to that of structural-connectivity degree, k, defined as the number of regions a given region is structurally connected to. Brain-network hubs, which have the highest k, are metabolically expensive (81–83) and characterized by long-range connectivity (84). However, the types of metabolic and neuronal GO categories that have been implicated in previous analyses of structural connectivity-related properties (18, 23) are also amongst those with the highest false-positive rates under conventional GSEA methodology. We therefore wished to verify whether these results stand up to the ensemble-based

null models that estimate significance relative to ensembles of null phenotypes (SBP-random and SBP-spatial). We computed the connectivity degree, k, of each brain region as the number of other regions to which it is connected, using diffusion-weighted imaging in human (85) and tract-tracing estimates in mouse (86) (see Methods for details).

Human cortex We performed GSEA for enrichment relative to structural connectivity degree, k. Conventional (random-gene null) GSEA identifies 365 significant GO categories, including many related to metabolism (e.g., 'ATP synthesis coupled electron transport', $q_{FDR} = 1 \times 10^{-12}$; and 'respiratory electron transport chain', $q_{FDR} = 2 \times 10^{-9}$) and neuronal function (e.g., 'chemical synaptic transmission, postsynaptic', $q_{FDR} = 4 \times 10^{-8}$; and 'excitatory postsynaptic potential', $q_{FDR} = 4 \times 10^{-7}$). FDR-corrected p-values q_{FDR}, for selected GO categories are listed in Table 2 (see Supplementary File `EnrichmentThreeWays_human_cortex.csv` for full list). These functions are sensibly related to what a researcher might expect of regions with high structural connectivity but, as characterized above, they are also amongst the categories with the highest false-positive significance rates under random (and spatially autocorrelated) SBPs. Indeed, relative to the SBP-random and SBP-spatial null ensembles, no GO categories were statistically significant in their relationship to k. These results suggest that within-category coexpression (and perhaps also spatial autocorrelation) of gene categories related to metabolic and neuronal function drive the enrichment findings under the random-gene null model.

Mouse Performing GSEA for k in the mouse brain yielded very different enrichment results across the three null models: random-gene (820 significant), SBP-random (1252 significant), and SBP-spatial (0 significant). Many of the significant GO categories under conventional random-gene and ensemble-based SBP-random nulls are sensibly related to metabolism and

GO Category	ID	q_{FDR} (random gene)	q_{FDR} (SBP-random)	q_{FDR} (SBP-spatial)
Human cortex				
ATP synthesis coupled electron transport	GO:0042773	$\mathbf{1 \times 10^{-12}}$	1.0	0.9
Chemical synaptic transmission, postsynaptic	GO:0099565	$\mathbf{4 \times 10^{-8}}$	1.0	0.9
Regulation of synaptic plasticity	GO:0048167	$\mathbf{4 \times 10^{-7}}$	1.0	0.9
Learning	GO:0007612	$\mathbf{3 \times 10^{-6}}$	1.0	0.9
Mouse brain				
ATP metabolic process	GO:0046034	$\mathbf{< 10^{-13}}$	$\mathbf{7 \times 10^{-3}}$	0.5
Learning	GO:0007612	$\mathbf{4 \times 10^{-11}}$	**0.01**	0.5
Dendritic spine development	GO:0060996	$\mathbf{5 \times 10^{-10}}$	**0.01**	0.5
Regulation of synaptic transmission, GABAergic	GO:0032228	$\mathbf{4 \times 10^{-4}}$	**0.004**	0.5
Mouse cortex				
ATP metabolic process	GO:0046034	$\mathbf{2 \times 10^{-4}}$	0.05	0.3
dendritic spine development	GO:0060996	**0.02**	0.06	0.3
Protein localization to synapse	GO:0035418	**0.04**	0.05	0.3
Synaptic vesicle priming	GO:0016082	0.04	0.06	0.3

Table 2. Testing against multiple null models allows more subtle interpretation of GSEA results, applied here to structural connectivity degree, k. We show selected results from biological process GO enrichment for the spatial variation of k, in: (i) mouse brain, (ii) mouse cortex, and (iii) human cortex. FDR-corrected p-values, q_{FDR}, are provided for: (i) the conventional random-gene null, (ii) the ensemble-based random phenotype null (SBP-random), and (iii) the ensemble-based spatially autocorrelated null (SBP-spatial). Significant $q_{\mathrm{FDR}} < 0.05$ are highlighted bold. Full tables containing results for all analyses summarized here are in Supplementary Files `EnrichmentThreeWays_human_cortex.csv`, `EnrichmentThreeWays_mouse_all.csv`, and `EnrichmentThreeWays_mouse_cortex.csv`.

neuronal function (Table 2; full results are in Supplementary File `EnrichmentThreeWays_mouse_all.csv`). The high number of significant categories under the SBP-random null is driven by a dominant expression signature that differentiates cortical areas from non-cortical areas (but is not specific to k). In the mouse brain, the vast majority of genes (73%: $14\,137/19\,417$) are differentially expressed in isocortical areas relative to all other areas ($q_{\mathrm{FDR}} < 0.05$, Wilcoxon rank-sum test). Connectivity degree, k, is also higher in isocortex than other areas ($p = 2 \times 10^{-3}$, Wilcoxon rank-sum test). Comparing all genes on: (i) their expression difference between cortical and non-cortical areas (Wilcoxon rank-sum test: $-\log_{10}(p)$), and (ii) their correlation to k (Spearman's ρ), we found a strong correlation, Spearman's $\rho = 0.67$. The predominance of genes that display differential expression in cortical areas thus drives an apparent GO enrichment signature for k that is actually dominated by the enrichment signatures of the isocortex. This result looks sensible because genes that are most differentially expressed in the isocortex are strongly related to neuronal, metabolic, and behavioral GO categories (cf. results of a conventional GSEA performed on differential expression in isocortex in Supplementary File `MouseIsocortexRanksumGSEA.csv`). This is another form of confound in the interpretation of GSEA analyses (that affects all null models) that, independent of statistical issues with null significance rates in GSEA, highlights the need to critically interpret apparently sensible GSEA results relative to alternative explanations.

As a simple way to examine the results without the dominant expression difference between cortical and non-cortical areas, we focused our next analysis on the 38 isocortical areas of the mouse brain, and computed intracortical degree, $k^{\mathrm{isocortex}}$ (using only cortico-cortical connections). Again, the number of significant GO categories varied across the null models: the conventional random-gene GSEA flagged 50 significant categories, whereas zero categories were significant relative to the SBP-random and SBP-spatial ensembles. Consistent with existing results relating hubs to genes involved in metabolic and neuronal function, significant GO categories under the random-gene null include 'ATP metabolic process' and 'protein localization to synapse', as listed in Table 2 (the full table is in Supplementary File `EnrichmentThreeWays_mouse_cortex.csv`). No GO categories are more correlated to k than would be expected from an ensemble of random maps (all $q_{\mathrm{FDR}} \geq 0.05$, but with many GO categories slightly above the 0.05 threshold), or a representative ensemble of spatially autocorrelated maps (all $q_{\mathrm{FDR}} > 0.3$).

This result indicates that while the spatial map of k is significantly correlated to the expression of some neuronal and metabolic gene categories in human cortex, mouse brain, and mouse cortex, this level of correlation is consistent with ensembles of non-specific expression patterns. Our results reveal how different null comparisons can drive significance in different GO categories, emphasizing the importance of proper comparison and careful consideration of the brain's spatial embedding when interpreting GSEA results.

Discussion

The ability to perform GSEA to identify groups of genes putatively related to spatially varying neural phenotypes has been facilitated by open neuroinformatics tools for accessing genome- and brain-wide transcriptional atlases in mouse and human (3–5, 87), and software tools for performing GSEA (38–42, 46, 47). While previous work has emphasized the need for care when applying GSEA for general applications—including substantial inter-software methodological variance, the reliably of gene-to-category annotations, and ongoing statistical improvements (48–50)—here we present a detailed anal-

ysis of methodological issues specifically associated with applications involving transcriptional atlas data, which are a new application for GSEA. We show that gene–gene coexpression, a generic property of the transcriptional atlas, makes some GO categories far more prone to false-positive bias than others. These GO categories with high false-positive bias are more likely to be reported as significant in the literature, suggesting that published results are heavily affected by statistical bias. We show that there is potential for further bias when analyzing phenotypes with similar spatial autocorrelation structure as GO categories. The flexible new ensemble-based framework that we introduce for generating GO-category null distributions enables more accurate inference of how spatially embedded brain phenotypes covary with the expression of specific functional categories of genes.

Our tests applying conventional GSEA (which relies on a random-gene null) to transcriptional atlas data using randomized phenotypes indicate heavily inflated false-positive rates for individual GO categories, at an average level of 875-fold in mouse and 630-fold in human (relative to the reference level). GO categories varied widely in their CFPRs, with some brain-related categories exhibiting CFPRs over 20% (relative to a maximum CFPR of 0.03% in the reference case). We show that this false-positive bias is primarily driven by within-category gene–gene coexpression: a generic property of the expression atlas, not of the specific spatial brain phenotype being analyzed. Categories of genes involved in brain function tend to exhibit spatially coordinated expression patterns (compared to the less correlated expression patterns of non-brain genes), yielding high gene–gene coexpression and thus high CFPR. This leads to the dangerous consequence that applying conventional GSEA to transcriptional-atlas data can misleadingly yield significant brain-related GO categories that are feasibly connected to the neural phenotype of interest. For example, GO categories related to oxidative metabolism have been linked to hub connectivity (18, 23), a result that can be plausibly interpreted in light of the known metabolic expense of brain-network hubs inferred from other modalities (81–83). While results of conventional GSEA reported the literature are not necessarily spurious, we must take care to validate these results using more rigorous statistical methodology, such as the ensemble-based nulls introduced here.

Ensemble-based hypothesis testing for GSEA is a flexible framework with which to perform valid inference of the enrichment of spatial phenotypes. In the SBP-random null model, random numbers are assigned to all brain regions, and in the SBP-spatial null model we instead generate an ensemble of phenotypes with a given spatial autocorrelation structure. The framework is flexible to other types of generative null models to test more specific questions. For example, the SBP-spatial ensemble used here takes parameters from the average spatial correlation properties of transcriptional atlas data, but they could instead be fitted to match the spatial autocorrelation properties of the phenotype of interest using, e.g., the *brainSMASH* toolbox (64). Another generative null model could also impose an anatomical constraint which generates an ensemble of phenotypes by randomizing the target phenotype separately within specific anatomical divisions—e.g., the ensemble could preserve differences in expression between cortical and non-cortical areas by randomizing cortical areas separately from subcortical areas. This flexibility to test against different null ensembles allows inference, and subsequent interpretation, to be tailored to the specific scientific question at hand. Ensemble-based nulls are expensive to compute, as GSEA often requires estimating very small *p*-values, and new SBPs must be generated for each null sample. To ease the computational burden, here we used a Gaussian approximation for the 40 000 sample null distribution computed for each GO category, allowing us to estimate small *p*-values from the fitted Gaussian to a greater precision than a direct permutation test. However, while the initial computation of null distributions for each GO category is expensive, it is a one-off computation for ensembles that are independent of the phenotype being tested (like the SBP-random and SBP-spatial ensembles studied here). That is, once the initial generic null computation has been computed, ensemble-based GSEA can then be applied efficiently for any new phenotype.

Much of the brain's structural organization can be accounted for by its physical embedding. Spatial autocorrelation means that samples are not independent, which requires corrections to any resulting statistical inference, as has been noted in neuroimaging applications (30, 64, 65, 88), in other fields like geography (89), and for data types like time series that also exhibit autocorrelation (90–92). The effect of spatial autocorrelation on GSEA is not straightforward: relative to independent random numbers (SBP-rand), applying an autocorrelation constraint (SBP-spatial) on average increased CFPRs modestly (1.3% in mouse and 1.2% in human), but the majority of GO categories exhibited a decrease in CFPR (60% in mouse and 68% in human). We showed that GO categories with a more similar spatial autocorrelation strength and length scale as the SBP-spatial ensemble have the greatest increase in CFPR relative to the SBP-rand ensemble (Fig. 4B, Fig. S3). Differences in CFPRs determined by different null ensembles can be understood by viewing each generative null model as defining a probability distribution in the space of all possible SBPs. For categories that have similar spatial autocorrelation properties as the SBP-spatial ensemble, the SBP-spatial ensemble will define a higher probability density around SBPs that are correlated to its genes, yielding CFPR(SBP-spatial) > CFPR(SBP-rand). Because here we use an SBP-spatial ensemble with a fixed strength and length

scale, only a minority of categories match this quite restricted set of phenotypes better than SBP-rand (40% in mouse and 32% in human), but for categories that match closely, CFPR can increase substantially. For example, in mouse, the CFPR of 'ionotropic glutamate receptor signaling pathway' increases from 8.8% (SBP-rand) to 29.2% (SBP-spatial) and 'dopamine receptor signaling pathway' increases from 6.5% to 26.2%. Interestingly, maps with strong spatial autocorrelation, but at a different length scales, can be less correlated to each other than to random maps; this effect could explain the large drops in CFPR under SBP-spatial of some highly spatially autocorrelated GO categories such as 'ATP metabolic process' (CFPR in mouse decreased from 20.1% to 10.7% under the SBP-spatial ensemble, despite high $R^2 = 0.37$). Thus, whether the SBP-spatial ensemble is more or less conservative than the SBP-rand ensemble depends on the specific spatial autocorrelation properties of each given GO category and how they relate to the properties of the SBP-spatial ensemble.

Our aim here was to characterize false-positive bias in applying GSEA to transcriptional data and uncover its key underpinnings, but there is much scope for future improvements. For example, our specific quantitative results depend on many parameters, including generic methodological settings of running GSEA. For demonstration, here we used a basic GSEA pipeline, but it will continue to be refined in future work, including better accounting for the hierarchical organization of GO terms and annotations (e.g., setting a level to test at, or defining rules for terminating hypothesis testing while ascending the hierarchy) (48, 52). Additionally, while the standard baseline gene set in GSEA includes all genes, future work may also investigate refining the reference set of genes (and thus GO categories) to, for example, those with feasible brain-related functions. This would circumvent the need to statistically correct across > 100s of GO categories that are not plausibly related to brain function (e.g., related to lung or heart function), and may facilitate the detection of biological processes relevant to neuronal systems. Finally, we note that while here we have focused on spatial correlations between phenotype maps and expression maps, there is important work to be done in extending this approach to connection-based (pairwise) analyses that score genes according to their contribution to correlated gene expression between pairs of brain areas (6, 10, 18). While an advantage of working in this pairwise domain is that the spatial dependence of correlated gene-expression can be modeled and removed, such analyses do not account for variations in within-category coexpression between GO categories. Extending the applicability of ensemble-based nulls to pairwise analyses will be an important extension of the current work.

Finally, we make some recommendations to improve the transparency and reproducibility of applying GSEA to transcriptomic atlas data. We first recommend generally

that researchers follow in the spirit of the open resources that enable their analysis (e.g., GO (36), KEGG (37), AHBA (4), and AMBA (3)) by providing code and data to reproduce their results. In reporting on GSEA analyses, we note that GO annotations are updated daily, such that GSEA tools can produce different results to identical inputs when run at different times. We thus recommend that researchers share the gene-score or gene-list inputs they used as input to the GSEA pipeline with a detailed methodological description including all settings used in performing enrichment. This would allow future studies to test for reproducibility through time, as gene annotations, categories, and category structures are updated (49), and would enable testing for robustness to other methodological implementations of GSEA (e.g., from different software packages). In reporting the results of GSEA, we discourage the practice of only listing manually selected GO categories. Rather, the full outputs of the enrichment procedure should be provided as supplementary files that include raw and corrected *p*-values. Finally, we note that data-processing choices for gene-expression data can have a large effect on GSEA results. It is therefore advisable to demonstrate robustness of a given set of enrichment results to a range of appropriate processing choices, including thoughtful justification of normalization procedures, distance-correction, and so on (see (5) for a review).

The open availability of gene transcriptional atlases provides an incredible opportunity to bridge scales of analysis in the brain to understand how spatial variations in molecular function relate to large-scale properties of brain structure and function (6, 93). GSEA plays a crucial methodological role in leveraging gene ontologies to interpret these multi-scale relationships. We highlight major statistical issues in applying conventional implementations of GSEA to assess the correlation between spatial brain phenotypes and the expression patterns of functionally annotated categories of genes, and introduce new ensemble-based null models that allow researchers to exploit whole-brain molecular maps in addressing diverse scientific questions. Extensive data tables for all enrichment analyses, code for reproducing all presented results, and a toolbox for performing ensemble-based enrichment accompany this paper.

Data and Methods

Documented code for reproducing all analyses presented here is available at https://github.com/benfulcher/GSEA_FalsePositives. All raw and processed data are available from an associated figshare repository. A toolbox for performing conventional (random-gene null) and ensemble-based GSEA in Matlab is available at https://github.com/benfulcher/GeneSetEnrichmentAnalysis.

Data.

Mouse   We used gene-expression data from the Allen Mouse Brain Atlas (AMBA) (3), obtained using the Allen Software Development Kit (SDK)[1] using the 213-region parcellation of Oh et al. (86). Gene transcription in a given brain area was summarized as the 'expression energy' (the mean ISH intensity across voxels of that brain area) (3, 94). Where multiple section datasets were available for the same gene, they were combined by first z-scoring expression data taken across areas, and then computing the expression value of a cortical area as the mean across these z-scored section datasets. For each of 19 419 genes, we applied a 50% quality threshold, first on genes (keeping genes with expression data for at least 50% of brain areas), and then on brain areas (keeping brain areas with expression data for at least 50% of genes). This resulted in a region × gene expression matrix of size $213 \times 19\,417$ (whole brain), and $38 \times 19\,417$ (isocortical areas).

For the case study involving structural connectivity degree, we used axonal-connectivity data based on 469 anterograde viral microinjection experiments in C57BL/6J male mice at age P56, obtained from the Allen Mouse Brain Connectivity Atlas (AMBCA) (86). We computed degree, k, from binarized ipsilateral axonal connectivity in the right-hemisphere, including edges with $p < 0.05$, as inferred from the whole-brain linear-regression model presented in Oh et al. (86) (see Fulcher and Fornito (18)).

Human   We use gene-expression data from the Allen Human Brain Atlas (AHBA) that consists of microarray expression measures across 3702 spatially distinct tissue samples collected from six neurotypical postmortem adult brains (4). Considering that two of the six brains were sampled from both left and right hemispheres while the other four brains had samples collected only from the left hemisphere, we focus our analyses on the left hemisphere only. The data was processed as outlined in Arnatkeviciūtė et al. (5). Specifically: i) probe-to-gene annotations were updated using the Re-Annotator toolbox (95); ii) intensity-based filtering was applied in order to exclude probes that do not exceed background noise in more than 50% of samples; iii) if more than one probe was available for a gene, a probe with the highest differential stability score (55) was selected; iv) gene expression samples from cortical regions were assigned to the regions-of-interest by generating donor-specific gray-matter parcellations (180 regions per hemisphere) and assigning samples located within 2 mm of the parcellation voxels; v) gene expression measures within a given brain were normalised first by applying a scaled robust sigmoid normalization (96) for every sample across genes and then for every gene across samples. This allowed us to evaluate the relative expression of each gene across regions, while controlling for donor-specific differences in gene expression. Normalized expression

measures in samples assigned to the same region were averaged within each donor brain and then aggregated into a $180 \times 15\,744$ region × gene matrix consisting of expression measures for 15 744 genes across 180 cortical regions in the left cortex. Three of the 180 regions had no gene expression samples assigned; all analyses shown here are of the remaining 177 regions.

Structural connectivity was estimated based on the minimally processed diffusion weighted imaging and structural data from the Human Connectome Project (85) for 972 participants ($\text{age}_{\text{mean}} = 28.7 \pm 3.7$, 522 females) (97, 98). Data were acquired on a customised Siemens 3 T 'Connectome Skyra' scanner (Washington University in St Louis, Missouri, USA) using a multi-shell protocol for the DWI: $1.25\,\text{mm}^3$ isotropic voxels, repetition time (TR) = 5.52 s, echo time (TE) = 89.5 ms, field-of-view (FOV) of 210 × 180 mm, 270 directions with $b = 1000$, 2000, $3000\,\text{s/mm}^2$ (90 per b value), and 18 $b = 0$ volumes. Structural T1-weighted data were collected using $0.7\,\text{mm}^3$ isotropic voxels, TR = 2.4 s, TE = 2.14 ms, FOV of 224 × 224 mm. The full details regarding data acquisition can be found elsewhere (97). For each individual network, nodes were defined using a recently-developed, data-driven group average parcellation of the cortex into 360 regions (180 per hemisphere) (99) using Freesurfer-extracted surfaces and subsequently projected to volumetric space.

Processing of the DWI data was performed using the MRtrix3 (100) and FMRIB Software Library (101). Tractography was performed in each participant's T1 space using second order integration over fibre orientation distributions (iFOD2) – a probabilistic algorithm that improves the quality of tract reconstruction in the presence of crossing fibers and high degree of fiber curvature (102). To further improve the biological accuracy of the structural networks we also applied Anatomically Constrained Tractography (ACT), which delineates the brain into different tissue types (cortical grey matter, subcortical grey matter, white matter, cerebrospinal fluid). This information is then used to ensure that streamlines are beginning, traversing, and terminating in anatomically plausible locations (103). Tissue types were determined using FSL software (101). A total of 10^7 streamlines were generated using a dynamic seeding approach. By evaluating the relative difference between the estimated and current reconstruction fibre density, it preferentially initiates seeding from areas of insufficient density (104).

The resulting tractogram was then combined with the cortical parcellation for each subject assigning streamline termination points to the closest region within 5 mm radius. Connection weights were quantified using streamline count (number of streamlines connecting two regions). Connectomes derived from probabilistic tractography algorithms are often thresholded due to the high probability of false-positive connections (105, 106), therefore a single group-average connectome was aggre-

---

[1]©2010 Allen Institute for Brain Science. Allen SDK. Available from: http://alleninstitute.github.io/AllenSDK/

gated by selecting connections that are present in at least 30% of subjects and retaining the strongest edges (based on the median streamline count across subjects) to achieve a connectome density of 20%.

Gene Set Enrichment Analysis (GSEA). We performed GSEA using gene-score resampling, where continuous scores are assigned to each gene, aggregated as the mean across individual GO categories, and then compared to a null distribution of category-level scores to perform statistical inference on GO categories [36].

We first describe our implementation of conventional GSEA using gene-score resampling against random-gene nulls (ensemble-based nulls are described later). We performed GSEA using Matlab-based software that we developed, available at https://github.com/benfulcher/GeneSetEnrichmentAnalysis. We used GO Term hierarchy and annotation files were downloaded from GO on 17th April 2019. We first matched annotations to GO Terms (excluding NOT and ND annotations [48]). To match genes listed in the GO annotation file (MGI identifiers) to expression data (NCBI identifiers), we used MouseMine [107]. In human, identification was made directly from the gene symbol. Annotations to genes that could not be mapped in this way were ignored. Direct gene-to-category annotations were propagated up to parents within the GO-Term hierarchy by taking the `is_a` keyword to indicate parent-child relationships. This process yielded a fully propagated set of GO categories and their gene annotations. We restricted our analyses to GO categories related to biological processes with between 10 and 200 annotations. We performed enrichment across all GO categories meeting these criteria (we did not perform a hierarchical tree-like statistical search that terminates the search at significant leaf nodes [48]).

Null distributions for a GO category of a given size, s, were generated through $40\,000$ random samples of s gene scores. Achieving stable p-value estimates from the empirical null distribution generated through permutation testing can require millions of null samples due to the small p-values involved in GSEA. As null distributions of category scores tend to be approximately Gaussian, we chose to estimate p-values from a Gaussian distribution fitted to a given set of null samples. That is, p-values were estimated numerically after fitting a Gaussian distribution to the estimated null distribution. Correction for multiple-hypothesis testing was achieved as a false-discovery rate [108], denoted as q_{FDR} and computed using the `mafdr` function in Matlab.

Analysis and Testing.

Generating spatially autocorrelated maps We generated SBP-spatial ensembles of spatially autocorrelated SBPs using the spatial-lag model [30, 80]. Code was written in Matlab, adapted from the Murray Lab's `surrogates` package by Joshua Burt, available

at https://bitbucket.org/murraylab/surrogates/src/master/ [30]. Note that a more recent *BrainSMASH* package has since been developed: https://github.com/murraylab/brainsmash [64]. In the spatial-lag model, for a Euclidean distance matrix constructed for all areas, two parameters determine the generated maps: a parameter controlling the strength of the spatial autocorrelation (relative to noise), ρ, and the characteristic spatial scale, d_0. We set $\rho = 0.8$ and estimated d_0 from an exponential fit to (Pearson) correlated gene expression (across all genes) as a function of distance [18], yielding $d_0 = 1.46\,\text{mm}$ in mouse brain, $d_0 = 1.84\,\text{mm}$ in mouse cortex, and $d_0 = 102.2\,\text{mm}$ in human cortex.

The within-category coexpression score, $\langle r \rangle$ To investigate the role of within-category coexpression in driving false-positive significance, we computed a simple measure of within-category coexpression, $\langle r \rangle$. For each GO category, we computed the gene $\times$ gene coexpression matrix, C, for all genes in that category as Spearman correlation coefficients between the spatial expression pattern of each pair of genes. The within-class coexpression metric, $\langle r \rangle$, was then computed as the mean of the unique (upper diagonal) entries of C.

The spatial autocorrelation score, R^2_{exp} To investigate the role of spatially autocorrelated SBPs in driving differential significance between GO categories, we computed a spatial autocorrelation score, R^2_{exp}, for each GO category. For each category, we computed a correlated gene expression (CGE) value for each pair of brain regions as a Pearson correlation between the expression values of genes in that category. We then fitted a three-parameter exponential function to the variation of CGE with distance, d, as $\text{CGE}(d) = A\exp(-d/\lambda) + B$. The goodness of this fit, measured as R^2_{exp}, provides a measure of how well a given category of genes exhibits a decaying distance-dependent expression similarity. R^2_{exp} will be high for strongly spatially autocorrelated categories of genes. For spatially autocorrelated GO categories, the length scale, λ, provides a measure of the autocorrelation length scale (used in Fig. S3). Note that we obtained similar results when computing a Spearman correlation between d and CGE, which does not assume the exponential functional form. Values of these fitted parameters for each GO category are in Supplementary Files `CategorySpatialScoring_human-cortex.csv` and `CategorySpatialScoring_mouse-all.csv`.

ACKNOWLEDGEMENTS
The authors would like to thank Stuart Oldham for assistance in processing human structural connectivity data and Oliver Cliff for thoughtful discussion and feedback on the manuscript. We thank Colorgorical [109] for assisting with the generation of color palettes. AF was supported by the Sylvia and Charles Viertel Charitable Foundation.

Bibliography

1. G. Buzsáki and A. Draguhn. Neuronal Oscillations in Cortical Networks. *Science* **304**, 1926 (2004).

2. J. W. Lichtman and W. Denk. The Big and the Small: Challenges of Imaging the Brain's Circuits. *Science* **334**, 618 (2011).

3. E. Lein, M. J. Hawrylycz, N. Ao, et al. Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168 (2007).

4. M. J. Hawrylycz, E. Lein, A. L. Guillozet-Bongaarts, et al. An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* **489**, 391 (2012).

5. A. Arnatkevičiūtė, B. D. Fulcher, and A. Fornito. A practical guide to linking brain-wide gene expression and neuroimaging data. *NeuroImage* **189**, 353 (2019).

6. A. Fornito, A. Arnatkevičiūtė, and B. D. Fulcher. Bridging the Gap between Connectome and Transcriptome. *Trends in Cognitive Sciences* **23**, 34 (2019).

7. A. Kaufman, G. Dror, I. Meilijson, and E. Ruppin. Gene Expression of Caenorhabditis elegans Neurons Carries Information on Their Synaptic Connectivity. *PLoS Comp. Biol.* **2**, e167 (2006).

8. V. Varadan, D. M. Miller III, and D. Anastassiou. Computational inference of the molecular logic for synaptic connectivity in C. elegans. *Bioinformatics* **22**, e497 (2006).

9. L. Baruch, S. Itzkovitz, M. Golan-Mashiach, E. Shapiro, and E. Segal. Using Expression Profiles of Caenorhabditis elegans Neurons To Identify Genes That Mediate Synaptic Connectivity. *PLoS Comp. Biol.* **4**, e1000120 (2008).

10. A. Arnatkevičiūtė, B. D. Fulcher, R. Pocock, and A. Fornito. Hub connectivity, neuronal diversity, and gene expression in the Caenorhabditis elegans connectome. *PLoS Comp. Biol.* **14**, e1005989 (2018).

11. L. Wolf, C. Goldberg, N. Manor, R. Sharan, and E. Ruppin. Gene Expression in the Rodent Brain is Associated with Its Regional Connectivity. *PLoS Comp. Biol.* **7**, e1002040 (2011).

12. L. French and P. Pavlidis. Relationships between gene expression and brain wiring in the adult rodent brain. *PLoS Comp. Biol.* **7**, e1001049 (2011).

13. L. French, P. P. C. Tan, and P. Pavlidis. Large-scale analysis of gene expression and connectivity in the rodent brain: insights through data integration. *Front. Neuroinf.* **5**, 12 (2011).

14. S. Ji, A. Fakhry, and H. Deng. Integrative analysis of the connectivity and gene expression atlases in the mouse brain. *NeuroImage* **84**, 245 (2014).

15. M. Rubinov, R. J. F. Ypma, C. Watson, and E. T. Bullmore. Wiring cost and topological participation of the mouse brain connectome. *Proc. Natl. Acad. Sci. USA* **112**, 10032 (2015).

16. A. Fakhry and S. Ji. High-resolution prediction of mouse brain connectivity using gene expression patterns. *Methods* **73**, 71 (2015).

17. A. Fakhry, T. Zeng, H. Peng, and S. Ji. Global analysis of gene expression and projection target correlations in the mouse brain. *Brain Inform.* **2**, 107 (2015).

18. B. D. Fulcher and A. Fornito. A transcriptional signature of hub connectivity in the mouse connectome. *Proc. Natl. Acad. Sci. USA* **113**, 1435 (2016).

19. L. Parkes, B. D. Fulcher, M. Yücel, and A. Fornito. Transcriptional signatures of connectomic subregions of the human striatum. *Genes, Brain and Behavior* **25**, 1176 (2017).

20. P. Goel, A. Kuceyeski, E. LoCastro, and A. Raj. Spatial patterns of genome-wide expression profiles reflect anatomic and fiber connectivity architecture of healthy human brain. *Hum. Brain Map.* **35**, 4204 (2014).

21. J. Richiardi, A. Altmann, A.-C. Milazzo, et al. Correlated gene expression supports synchronous activity in brain networks. *Science* **348**, 1241 (2015).

22. I. Diez and J. Sepulcre. Neurogenetic profiles delineate large-scale connectivity dynamics of the human brain. *Nat. Comm.* **9**, 1 (2018).

23. P. E. Vértes, T. Rittman, K. J. Whitaker, et al. Gene transcription profiles associated with inter-modular hubs and connection distance in human functional magnetic resonance imaging networks. *Phil. Trans. Roy. Soc. B* **371**, 20150362 (2016).

24. R. F. Betzel, J. D. Medaglia, A. E. Kahn, et al. Structural, geometric and genetic factors predict interregional brain connectivity patterns probed by electrocorticography. *Nature Biomedical Engineering* **106**, 1 (2019).

25. J. Ritchie, S. P. Pantazatos, and L. French. Transcriptomic characterization of MRI contrast, focused on the T1-w/T2-w ratio in the cerebral cortex. *NeuroImage* **174**, 504 (2018).

26. K. J. Whitaker, R. Romero-Garcia, F. Váša, et al. Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. *Proc. Natl. Acad. Sci. USA* **113**, 201601745 (2016).

27. J. Seidlitz, F. Váša, M. Shinn, et al. Morphometric Similarity Networks Detect Microscale Cortical Organization and Predict Inter- Individual Cognitive Variation. *Neuron* pp. 1–25 (2018).

28. R. Romero-Garcia, K. J. Whitaker, F. Váša, et al. Structural covariance networks are coupled to expression of genes enriched in supragranular layers of the human cortex. *NeuroImage* **171**, 256 (2018).

29. P. K. Reardon, J. Seidlitz, S. Vandekar, et al. Normative brain size variation and brain shape diversity in humans. *Science* **360**, eaar2578 (2018).

30. J. B. Burt, M. Demirtas, W. J. Eckner, et al. Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nat. Neurosci.* **27**, 889 (2018).

31. K. H. Preller, J. B. Burt, J. L. Ji, et al. Changes in global and thalamic brain connectivity in lsd-induced altered states of consciousness are attributable to the 5-ht2a receptor. *Elife* **7**, e35082 (2018).

32. I. A. C. Romme, M. A. de Reus, R. A. Ophoff, R. S. Kahn, and M. P. van den Heuvel. Connectome Disconnectivity and Cortical Gene Expression in Patients With Schizophrenia. *Biol. Psychiatry* **81**, 495 (2017).

33. T. Rittman, M. Rubinov, P. E. Vértes, and A. X. Patel. Regional expression of the MAPT gene is associated with loss of hubs in brain networks and cognitive impairment in Parkinson disease and progressive . . . . *Neurobiology of . . .* **48**, 153 (2016).

34. R. Romero-Garcia, V. Warrier, E. T. Bullmore, S. Baron-Cohen, and R. A. I. Bethlehem. Synaptic and transcriptionally downregulated genes are associated with cortical thickness differences in autism. *Molecular Psychiatry* **24**, 1053 (2019).

35. S. E. Morgan, J. Seidlitz, K. J. Whitaker, et al. Cortical patterning of abnormal morphometric similarity in psychosis is associated with brain expression of schizophrenia-related genes. *Proc. Natl. Acad. Sci. USA* **97**, 201820754 (2019).

36. M. Ashburner, C. A. Ball, J. A. Blake, et al. Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25 (2000).

37. M. Kanehisa and S. Goto. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucl. Acid. Res.* **28**, 27 (2000).

38. J. Gillis, M. Mistry, and P. Pavlidis. Gene function analysis in complex data sets using ErmineJ. *Nat. Protoc.* **5**, 1148 (2010).

39. G. Dennis Jr, B. T. Sherman, D. A. Hosack, and J. Yang. DAVID: database for annotation, visualization, and integrated discovery. *Genome Biol.* (2003).

40. P. Shannon, A. Markiel, O. Ozier, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* **13**, 2498 (2003).

41. E. Eden, R. Navon, I. Steinfeld, D. Lipson, and Z. Yakhini. Gorilla: a tool for discovery and visualization of enriched go terms in ranked gene lists. *BMC bioinformatics* **10**, 48 (2009).

42. H. Mi, A. Muruganujan, and P. D. Thomas. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucl. Acid. Res.* **41**, D377 (2013).

43. M. V. Kuleshov, M. R. Jones, A. D. Rouillard, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic acids research* **44**, W90 (2016).

44. D. Klopfenstein, L. Zhang, B. S. Pedersen, et al. Goatools: A python library for gene ontology analyses. *Scientific reports* **8**, 10872 (2018).

45. B. Zhang, S. Kirov, and J. Snoddy. Webgestalt: an integrated system for exploring gene sets in various biological contexts. *Nucleic acids research* **33**, W741 (2005).

46. S. Carbon, A. Ireland, C. J. Mungall, et al. Amigo: online access to ontology and annotation data. *Bioinformatics* **25**, 288 (2008).

47. J. Chen, E. E. Bardes, B. J. Aronow, and A. G. Jegga. Toppgene suite for gene list enrichment analysis and candidate gene prioritization. *Nucleic acids research* **37**, W305 (2009).

48. S. Y. Rhee, V. Wood, and K. Dolinski. Use and misuse of the gene ontology annotations. *Nat. Rev. Genet.* (2008).

49. A. Tomczak, J. M. Mortensen, R. Winnenburg, et al. Interpretation of biological experiments changes with evolution of Gene Ontology and its annotations. *bioRxiv* p. 228080 (2017).

50. S. Ballouz, P. Pavlidis, and J. Gillis. Using predictive specificity to determine when gene set analysis is biologically meaningful. *Nucl. Acid. Res.* **45**, e20 (2017).

51. X. Dong, Y. Hao, X. Wang, and W. Tian. LEGO: a novel method for gene set over-representation analysis by incorporating network-based gene weights. *Sci. Rep.* **6**, 18871 (2016).

52. A. Hukku, C. Quick, F. Luca, R. Pique-Regi, and X. Wen. BAGSE: a Bayesian hierarchical model approach for gene set enrichment analysis. *bioRxiv* **9**, 662171 (2019).

53. L. Wang, P. Jia, R. D. Wolfinger, X. Chen, and Z. Zhao. Gene set analysis of genome-wide association studies: Methodological issues and perspectives. *Genomics* **98**, 1 8 (2011).

54. M. J. Hawrylycz, A. Bernard, C. Lau, et al. Areal and laminar differentiation in the mouse neocortex using large scale gene expression data. *Methods* **50**, 113 (2010).

55. M. J. Hawrylycz, J. A. Miller, V. Menon, et al. Canonical genetic signatures of the adult human brain. *Nat. Neurosci.* **18**, 1832 (2015).

56. F. M. Krienen, B. T. T. Yeo, T. Ge, R. L. Buckner, and C. C. Sherwood. Transcriptional profiles of supragranular-enriched genes associate with corticocortical network architecture in the human brain. *Proc. Natl. Acad. Sci. USA* **113**, E469 (2016).

57. S. P. Pantazatos and X. Li. Commentary: BRAIN NETWORKS. Correlated gene expression supports synchronous activity in brain networks. Science 348, 1241-4. *Front. Neurosci.* **11**, 412 (2017).

58. J. Richiardi, A. Altmann, and M. Greicius. Distance Is Not Everything In Imaging Genomics Of Functional Networks: Reply To A Commentary On Correlated Gene Expression Supports Synchronous Activity In Brain Networks. *bioRxiv* p. 132746 (2017).

59. J. A. Henderson and P. A. Robinson. Geometric effects on complex network structure in the cortex. *Phys. Rev. Lett.* **107**, 018102 (2011).

60. J. A. Henderson and P. A. Robinson. Relations Between the Geometry of Cortical Gyrification and White-Matter Network Architecture. *Brain Conn.* **4**, 112 (2014).

61. J. A. Roberts, A. Perry, A. R. Lord, et al. The contribution of geometry to the human connectome. *NeuroImage* **124**, 379 (2016).

62. J. Stiso and D. S. Bassett. Spatial Embedding Imposes Constraints on Neuronal Network Architectures. *Trends in Cognitive Sciences* (2018).

63. P. A. Robinson. Physical brain connectomics. *Phys. Rev. E* **99**, 012421 (2019).

64. J. B. Burt, M. Helmer, M. Shinn, A. Anticevic, and J. D. Murray. Generative modeling of brain maps with spatial autocorrelation. *bioRxiv* (2020).

65. A. F. Alexander-Bloch, H. Shou, S. Liu, et al. On testing for spatial correspondence between maps of human brain structure and function. *NeuroImage* **178**, 540 (2018).

66. P. G. Reddy, R. F. Betzel, A. N. Khambhati, et al. Genetic and Neuroanatomical Support for Functional Brain Network Dynamics in Epilepsy. *arXiv* p. 1809.03934 (2018).

67. A. Subramanian, P. Tamayo, V. K. Mootha, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. USA* **102**, 15545 (2005).

68. K. M. Anderson, M. A. Collins, R. Chin, et al. The transcriptional landscape of cortical interneurons underlies in-vivo brain function and schizophrenia risk. *bioRxiv* p. 481036 (2018).

69. J. Liu, M. Xia, X. Wang, X. Liao, and Y. He. Transcriptional Signatures of the Hierarchical Chronnectome Organization in the Human Brain. *bioRxiv* **39**, 637942 (2019).

70. L. French and T. Paus. A FreeSurfer view of the cortical transcriptome generated from the Allen Human Brain Atlas. *Front. Psychiat.* **9**, 25 (2015).

71. Z. Liu, E. T. Rolls, J. Zhang, et al. The functional and genetic associations of neuroimaging data: a toolbox. *bioRxiv* p. 178640 (2017).

72. X. Yao, J. Yan, S. Kim, et al. Two-dimensional enrichment analysis for mining high-level imaging genetic associations. *Brain Informatics* **4**, 27 (2017).

73. M. Meijer, A. Keo, J. M. C. van Leeuwen, et al. Molecular characterization of the stress network in the human brain. *bioRxiv* **526**, 661587 (2019).

74. K. M. Anderson, F. M. Krienen, E. Y. Choi, et al. Gene expression links functional networks across cortex and striatum. *Nat. Comm.* **9**, 1428 (2017).

75. Z. Kuncheva, M. L. Krishnan, and G. Montana. Exploring brain transcriptomic patterns: a topological analysis using spatial expression networks. In *Pac. Soc. Biocomput.* (2017).

76. W. Liu, N. Peeters, G. Fernandez, and N. Kohn. Common neural and transcriptional correlates of inhibitory control across emotion, memory, and response inhibition. *bioRxiv* p. 708552 (2019).

77. D. Vidal-Pineiro, N. Parker, J. Shin, et al. Cellular correlates of cortical thinning throughout the lifespan. *bioRxiv* p. 585786 (2019).

78. P. P. C. Tan, L. French, and P. Pavlidis. Neuron-Enriched Gene Expression Patterns are Regionally Anti-Correlated with Oligodendrocyte-Enriched Patterns in the Adult Mouse and Human Brain. *Front. Psychiat.* **7** (2013).

79. B. D. Mills, D. S. Grayson, A. Shunmugavel, et al. Correlated gene expression and anatomical communication support synchronized brain activity in the mouse functional connectome. *J. Neurosci.* **38**, 2910 (2018).

80. L. Anselin. *Spatial econometrics: methods and models*, volume 4. Springer Science & Business Media (2013).

81. E. T. Bullmore and O. Sporns. The economy of brain network organization. *Nat. Rev. Neurosci.* **13**, 337 (2012).

82. G. Collin, O. Sporns, R. C. W. Mandl, and M. P. van den Heuvel. Structural and Functional Aspects Relating to Cost and Benefit of Rich Club Organization in the Human Cerebral Cortex. *Cereb. Cortex* **24**, 2258 (2014).

83. D. Tomasi, G. J. Wang, and N. D. Volkow. Energetic cost of brain functional connectivity. *Proc. Natl. Acad. Sci. USA* **110**, 13642 (2013).

84. M. P. van den Heuvel, R. S. Kahn, J. Goñi, and O. Sporns. High-cost, high-capacity backbone for global brain communication. *Proc. Natl. Acad. Sci. USA* **109**, 11372 (2012).

85. D. C. Van Essen, S. M. Smith, D. M. Barch, et al. The WU-Minn Human Connectome Project: An overview. *NeuroImage* **80**, 62 (2013).

86. S. W. Oh, J. A. Harris, L. Ng, et al. A mesoscale connectome of the mouse brain. *Nature* **508**, 207 (2014).

87. L. Ng, S. D. Pathak, C. Kuan, et al. Neuroinformatics for Genome-Wide 3-D Gene Expression Mapping in the Mouse Brain. *IEEE/ACM Trans. Comput. Biol. Bioinf.* **4**, 382 (2007).

88. M. Breakspear, M. J. Brammer, E. T. Bullmore, P. Das, and L. M. Williams. Spatiotemporal wavelet resampling for functional neuroimaging data. *Human brain mapping* **23**, 1 (2004).

89. M. Cardillo, L. Bromham, and S. J. Greenhill. Links between language diversity and species richness can be confounded by spatial autocorrelation. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20142986 (2015).

90. S. Afyouni, S. M. Smith, and T. E. Nichols. Effective degrees of freedom of the pearson's correlation coefficient under autocorrelation. *NeuroImage* (2019).

91. O. James, H. Park, and S.-G. Kim. Impact of sampling rate on statistical significance for single subject fmri connectivity analysis. *Human brain mapping* (2019).

92. O. M. Cliff, L. Novelli, B. D. Fulcher, J. M. Shine, and J. T. Lizier. Exact inference of linear dependence between multiple autocorrelated time series. *arXiv preprint arXiv:2003.03887* (2020).

93. B. D. Fulcher. Discovering Conserved Properties of Brain Organization Through Multimodal Integration and Interspecies Comparison:. *Journal of Experimental Neuroscience* **13**, 117906951986204 (2019).

94. L. Ng, A. Bernard, C. Lau, et al. An anatomic gene expression atlas of the adult mouse brain. *Nat. Neurosci.* **12**, 356 (2009).

95. J. Arloth, D. M. Bader, S. Röh, and A. Altmann. Re-Annotator: annotation pipeline for microarray probe sequences. *PLOS One* **10**, e0139516 (2015).

96. B. D. Fulcher, M. A. Little, and N. S. Jones. Highly comparative time-series analysis: the empirical structure of time series and their methods. *J. Roy. Soc. Interface* **10**, 20130048 (2013).

97. M. F. Glasser, S. N. Sotiropoulos, J. A. Wilson, et al. The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage* **80**, 105 (2013).

98. S. N. Sotiropoulos, S. Jbabdi, J. Xu, et al. Advances in diffusion MRI acquisition and processing in the Human Connectome Project. *Neuroimage* **80**, 125 (2013).

99. M. F. Glasser, T. S. Coalson, E. C. Robinson, et al. A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171 (2016).

100. J.-D. Tournier, F. Calamante, and A. Connelly. MRtrix: diffusion tractography in crossing fiber regions. *Int. J. Imaging Syst. Technol.* **22**, 53 (2012).

101. M. Jenkinson, C. F. Beckmann, T. E. Behrens, M. W. Woolrich, and S. M. Smith. FSL. *Neuroimage* **62**, 782 (2012).

102. J.-D. Tournier, F. Calamante, and A. Connelly. Improved probabilistic streamlines tractography by 2 nd order integration over fibre orientation distributions. *Ismrm* **88**, 2010 (2010).

103. R. E. Smith, J.-D. Tournier, F. Calamante, and A. Connelly. Anatomically-constrained tractography: improved diffusion MRI streamlines tractography through effective use of anatomical information. *Neuroimage* **62**, 1924 (2012).

104. R. E. Smith, J.-D. Tournier, F. Calamante, and A. Connelly. SIFT2: Enabling dense quantitative assessment of brain white matter connectivity using streamlines tractography. *Neuroimage* **119**, 338 (2015).

105. T. Sarwar, K. Ramamohanarao, and A. Zalesky. Mapping connectomes with diffusion MRI: deterministic or probabilistic tractography? *Magn. Reson. Med.* **81**, 1368 (2019).

106. S. N. Sotiropoulos and A. Zalesky. Building connectomes using diffusion MRI: why, how and but. *NMR Biomed.* p. e3752 (2017).

107. H. Motenko, S. B. Neuhauser, M. O'keefe, and J. E. Richardson. Mousemine: a new data warehouse for mgi. *Mammalian Genome* **26**, 325 (2015).

108. Y. Benjamini and Y. Hochberg. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. Roy. Stat. Soc. B* **57**, 289 (1995).

109. C. C. Gramazio, D. H. Laidlaw, and K. B. Schloss. Colorgorical: creating discriminable and preferable color palettes for information visualization. *IEEE Transactions on Visualization and Computer Graphics* (2017).