Characterising a stable five-species microbial community for use in experimental evolution and ecology.

Author Affiliations

Meaghan Castledine[1], Joseph Pennycook[2], Arthur Newbury[1], Luke Lear[1], Zoltan Erdos[1], Rai Lewis[1], Suzanne Kay[1], Dirk Sanders[1], David Sünderhauf[1], Angus Buckling[1], Elze Hesse[1], & Daniel Padfield[1].

[1]Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter, Penryn, Cornwall, TR10 9FE, U.K.

[2]University College Cork, Cork T12 K8AF

Corresponding author: Daniel Padfield (d.padfield@exeter.ac.uk)

Author contributions: J.P, E.H. and A.B. conceived the study and designed the experimental work. J.P, E.H., D.P and M.C conducted the experiments. R.L., and S.K. did DNA extractions and QC. D.P., and M.C. analysed the data. D.P. did the bioinformatic analyses. D.P. and M.C. wrote the first version of the manuscript and all authors contributed to editing subsequent versions.

Acknowledgements: This work was funded by NERC. D.P. was also supported by a NERC Independent Research Fellowship (NE/W008890/1). E.H. was supported by a UKRI Future Leaders Fellowship (MR/V022482/1). D.S. was supported in part by grant MR/N0137941/1 for the GW4 BIOMED MRC DTP, awarded from the Medical Research Council.

Conflict of interests: There are no conflicts of interest.

Abstract

Model microbial communities are regularly used to test ecological and evolutionary theory as

they are easy to manipulate and have fast generation times, allowing for large-scale, high

throughput experiments. A key assumption for most model microbial communities is that

they stably coexist, but this is rarely tested experimentally. Here we report the (dis)assembly

of a five-species microbial community from a metacommunity of soil microbes that can be

used for future experiments. Using reciprocal invasion from rare experiments we show that

all species can coexist and we demonstrate that the community is stable for a long time (~600

generations). Crucially for future work, we show that each species can be identified by their

plate morphologies, even after >1 year in co-culture. We characterise pairwise species

interactions and produce high-quality reference genomes for each species. This stable five-

species community can be used to test key questions in microbial ecology and evolution.

Introduction

Controlled experiments on synthetic microbial communities have been used to test general

ecological and evolutionary theories about - for instance - species coexistence [1], adaptive

radiations [2–4], and ecosystem functioning [5, 6]. The short generation times and large

population sizes of bacteria make them ideal for testing ideas that are experimentally

intractable in other model systems. Culturable bacteria are also easy to manipulate, allowing

us to change species richness [5, 7, 8], abundances [9, 10], and even interaction type [11, 12]

and strength [13] in a systematic and increasingly high throughput way.

The majority of previous experimental work has been done on monocultures or in

synthetic communities consisting of only a few species. However, an increasing number of

studies have used more diverse microbial communities to study how species rich

communities assemble and their stability, to understand the maintenance of the high levels of

microbial diversity seen in nature [8, 14–17]. Experiments on both simple and complex model

communities are key to understanding the processes underpinning community assembly and

coexistence, and there is an inherent trade-off between more detailed mechanistic

understanding and complexity [18].

A key assumption of both approaches is that the community members stably coexist,

meaning that the densities of species in the system do not show long-term trends [19].

Modern coexistence theory presents the ability to invade from rare (negative frequency

dependence) which tests that each species is able to increase from low density when in the

presence of the rest of the community, as a key test of coexistence [19, 20]. Experimentally

establishing that model communities are stable and that species coexist is key to increasing

the likelihood that the findings are relevant to natural communities (where high levels of

diversity are maintained, composition is relatively stable through time across space [21–23],

and where patterns of diversity cannot be explained solely through neutral processes [24–

26]). Second, when a model community is not stable, it risks biasing our understanding of

species interactions. Specifically, species will be driven locally extinct in unstable

communities, which will necessarily bias interactions towards being negative. Third,

demonstrating stable coexistence and understanding the mechanisms of coexistence (e.g.

equalising or stabilising [19]) would increase the generalisability of findings and allow future

work to understand under what circumstances stability breaks down.

There are several proposed methods to measure invasion growth rate [27], but these

become more complicated to test as species diversity increases, where you have to test

whether each species has a positive invasion growth rate while it is absent and the resident

community is at equilibrium [28]. For example, following the removal of a species, the

resident community may destabilise or convert to an alternative stable state [29]. Explicit tests

of coexistence in model microbial communities have been done when communities are

simple and contain only strains of one species [30–32], two [33], or three species [7], but they

remain scarce when species diversity goes beyond this ([5, 8, 13, 15, 17], but see [34]). Most

model microbial communities are performed in batch culture, where populations are regularly

transferred into fresh microcosms [2], meaning that "equilibrium" is never reached in a

measurable sense. Consequently, microbiologists have used invasion-from-rare assays to test

for negative frequency dependence and stable coexistence. In these assays, the community is

inoculated into fresh media, but one species (the invader) is at 100-fold lower density than the

other species (the resident community). If each species has a higher growth rate when rare,

relative to the resident community (a relative invader fitness greater than 1) then the

community shows negative frequency dependence and is likely to be stable [31, 35, 36].

Here, we report the characterisation of a stable 5 species community that can be used

for future experiments in ecology and evolution. We cultured a community pool of 46

isolates retrieved from soil over several weeks in lab media and the final community

composition was determined. As outcomes were highly repeatable, we took a single replicate

community and conducted reciprocal invasion experiments to check coexistence. In addition,

we measured species interactions using spent media and co-culture, tracked long-term

community dynamics, and generated high quality reference genomes. Finally, we discuss

potential questions that could be answered with this - or any - stable community when so

much information about its constituent species is known.

Methods

Community disassembly experiment

We created a model microbial community consisting of 5 species that (dis)assembled for

approximately 85 generations (Figure 1a). Briefly, this community was formed by

inoculating 46 bacterial strains – belonging to 27 genera (Table S1) – that were isolated from

an experimental compost community [37] into 6mL growth medium (1/64 Tryptic Soy Broth

(TSB) diluted with demineralised H$_2$O) in 25mL glass vials with loosened plastic lids. Forty

eight cultures were set up, incubated statically for 13 weeks at 28ºC. For the first five weeks

no transfers were made, after which weekly transfers were done, passaging 1% (60µl) of each

community into fresh 6mL microcosms every week. After this time, the 48 communities

converged on a similar composition of ~5 dominant species. We selected a single replicate of

these to explore its stability.

This community consisted of five species, *Achromobacter* sp., *Ochrobactrum* sp.,

Pseudomonas sp., *Stenotrophomonas* sp., and *Variovorax* sp.. Crucially, these species could

be reliably identified by their colony morphology (Figure 1b), allowing us to track each

species in the community and re-isolate them for phenotyping and clone re-sequencing. A

single clone of each member of this community was picked and grown in monoculture for

two days in 6mL of 1/64 TSB, shaking, in 25mL glass vials with loosened plastic lids at 28ºC

to achieve high cell densities. These samples were then cryogenically frozen at -70ºC in

glycerol (final concentration: 25%). These clones were used to recreate the community for

subsequent experiments.

Sanger sequencing of 16S rRNA of each morphotype

To understand how well our characterisation of different morphotypes represent different

species, we sequenced the 16S rRNA gene of multiple replicates of each morphotype. Of the

5 distinct morphotypes, we picked at least 8 colonies across all 48 microcosms. Polymerase

chain reactions (PCRs) were performed in 20µL reactions containing 10µL of DreamTaq

Green PCR MasterMix (2X) (Thermo Scientific), 0.4µL each of the 338F and 518R primers

(10µM) and 2µL of 1:100 diluted culture. In total 51 samples were Sanger sequenced using

the 515F primer (Eurofins Genomics).

The Sanger sequencing files were analysed in *R*, with *R* functions specified as

Rpackage::function. Chromatograms for each sample were created using

sangeranalyseR::secondary.peaks [38]. Low-quality bases were trimmed from the start and

end of each file using *sangeranalyseR::trimm.mott* which uses Mott's modified trimming

algorithm, with a cut-off of 40 to define a bad quality score. Files were further filtered to

retain only sequences that were longer than 100 base pairs long, had five or less secondary

peaks (where there is more than one peak at a given position; a secondary peak was defined

when the lower peak was at least a third as high as the higher peak), and had a mean quality

score of >30. This left 43 trimmed and cleaned sequences that went forward to further

analysis.

For each of the 5 identified morphs, we calculated a pairwise distance matrix from the

DNA sequences using *ape::dist.dna* [39], with a phylogenetic tree being constructed and

plotted using *ape::njs* and *ape::plot.phylo*. If all replicates of the same morph were attributed

to the same level on the tree, a single consensus sequence for the morph was constructed

using *msa::msaConsensusSequence* [40] after aligning multiple sequences of the same morph

using *msa::msa*. In instances where there was more than a single branch of the tree, samples

within a morph were split along branches to create multiple consensus sequences. Consensus

sequences were aligned using *msa::msa* to look at similarities between morphs. Taxonomy

was assigned to genus level using *dada2::assignTaxonomy* [41], which implements the RDP

Naive Bayesian Classifier to assign taxonomy to the RDP database (v16) [42].

Invasion-from-rare assays

To explicitly test for coexistence, we conducted invasion-from-rare assays [4]. To assess

whether all species were necessary for coexistence, we did assays where each species was

inoculated from rare in every species by diversity combination. In total, this yielded six

replicates of 75 treatments. Species were grown from frozen to achieve high cell densities as

described above. Cell density (colony-forming units [CFU]) was estimated from optical

density (600 nm wavelength; OD_{600}) using a species-specific calibration curve (Table S3 for

equations) and normalised to 10^5 CFU μL^{-1}. Focal species were invaded at a 100-fold lower

density (1.6μL) than that of the resident community (160μL in total). When there was more

than one resident species, resident communities were created from equal ratios of each

species. Cultures were incubated for one week, after which samples were cryogenically

frozen at -70ºC in glycerol. Species abundances (CFU mL^{-1}) were estimated by plating

cultures onto KB (King's medium B) agar and incubating for two days at 28°C.

Relative invader growth rate of the focal species was calculated as the ratio of

estimated Malthusian parameters (m), m_{focal}: $m_{community}$ where $m_{community}$ is the total density

change of all the other populations combined. $m = ln(N_1 / N_0)/t$ where N_1 is the final density,

N_0 is starting density, and t is the assay time (7 days) [4]. We performed independent one-

sample t-tests on each combination of focal species by resident community combination, with

the null-hypothesis being that the mean relative invader growth rate equals 1. A relative

invader growth rate above 1 would indicate that the invader had a faster growth rate than the

resident community, demonstrating negative frequency dependence. This resulted in 75

different statistical tests; P values were adjusted using the false discovery rate (*fdr*) method

[43].

Medium-term persistence assays

To test whether diversity impacts coexistence and persistence, and to quantify pairwise and

indirect (higher order) interactions, each species was also inoculated at equal density in every

possible community combination, from monoculture to five-species. Species monocultures

were prepared with 20µL of each species inoculated (total inoculum varied across diversity

levels) into fresh microcosms for each community combination. Twelve replicates of each

unique community combination were then incubated for either 1, 2, 3 or 4 weeks (three

replicates each) without resource replenishment. These assays allowed time for extinctions to

occur, to measure the effect of resource scarcity and secondary metabolites for cultures left

for 3 or 4 weeks, and for us to calculate species interactions of the community. At the end of

each incubation period, individual species abundance (CFU mL^{-1}) was estimated using

methods described above. Abundances from week 1 were removed as it was visually clear

that equilibrium density had not been reached for most species.

We calculated pairwise species interactions (e.g. effect of species *2* on species *1*, $w_{1/2}$)

as the abundance of species *1* in the presence of species *2* relative to that of species *1* grown

in monoculture (no interspecific interactions): $w_{1/2} = N_{1/2}/N_1$. If w is greater than 1, then the

interaction is positive, whereas a $w < 1$ demonstrates competitive interactions, with a value of

0 demonstrating competitive exclusion. To get a broad perspective of the nature of species

interactions, we calculated the average (the mean of ~9 replicates) for each pairwise species

interaction.

We estimated indirect (higher order) interactions from the persistence assays using a

modified version of a framework originally developed to help quantify higher order antibiotic

interactions on *Escherichia coli* [44]. As our assays cannot determine whether the presence of

a third, fourth, or fifth species changes competitor density (an interaction chain) or changes

per capita competitive effects (higher order interaction), our method calculates indirect

interactions generally. This method overcomes the issues of the additive model when single

species have large impacts on fitness, such that the additive expectation can never be

achieved (i.e. a population size < 0)[45, 46].

For each community combination with more than two species (e.g. species $1, 2, 3$),

the predicted effect on the focal species, $\hat{w}_{1/23}$, is the product of the independent pairwise

interactions: $\hat{w}_{1/23} = w_{1/2} \times w_{1/3}$. The indirect interaction (*ii*) of any community combination

on focal species 1, $ii_{1|23}$, was calculated as the deviation of the observed abundance relative to

monoculture, $w_{1|23}$, from that predicted from pairwise interactions: $ii_{1/23} = \hat{w}_{1/23} - w_{1/23}$. Using

this definition, if an *ii* is the same sign as the original prediction ($\hat{w}_{1/23}$), then it indicates

synergism: the combined presence of species 2 and 3 on species 1 is greater than either on

their own. If an *ii* is the opposite sign of the original prediction, then it indicates non-

additivity and potential buffering: the combined presence of species 2 and 3 on species 1 is

less than either on their own. To make all synergistic indirect interactions positive (and

buffering interactions negative), we multiplied indirect interactions of predicted negative

interactions by -1. We looked at whether the type of indirect interaction (buffering or

synergism) was related to the strength of the expected pairwise interaction using a linear

model with expected pairwise interaction as the response and type of indirect interaction as

the response. This was only done on predicted negative interactions as there were only seven

predicted positive interactions (all belonging to *Variovorax* sp.) in the dataset.

The original Tekin method rescales interaction estimates and quantifies indirect

interactions after accounting for all interactions at lower species combinations (for example,

in four species communities the indirect interaction of $ii_{1/234}$ includes taking into account the

estimate of $ii_{1/23}$, $ii_{1/24}$, and $ii_{1/34}$). Recent work has discussed whether pairwise interactions

alone can predict species abundances, so we chose to calculate all indirect interactions

relative to the prediction of only the pairwise interaction estimates.

Supernatant assays

Cell-free supernatants were used to establish cross-feeding assays in 96-well plates. To obtain

the supernatants, three replicate populations per species were established by inoculating a

single clone in 6mL of 1/64 TSB in 30mL glass vials. Following 2 days of growth at 28ºC,

we centrifuged individual cultures at 3000 rpm for 10 minutes, and filtered these through

0.22mm syringe filters (Merck, Millex PES). We confirmed supernatants were cell free by

plating these onto KB agar. For each focal strain we inoculated ~5 x 10^4 cells from an

overnight culture in triplicate wells, each containing either 200µL of fresh 1/64 TSB or cell-

free supernatant of either the same or a different species (n=3 per pairwise combination).

Following 48 hours of growth at 28℃, serial-diluted cultures were plated onto KB agar to

obtain final cell densities.

Growth rate per day was calculated as $ln(N_1 / N_0)/t$, where N_1 is final density, N_0 is the

initial density, and t is two (the number of days of growth). If the growth rate was negative,

we replaced the value with 0.001 to prevent negative values of relative growth required for

comparisons between species. Relative growth rate was calculated by dividing the growth

rate in supernatant by the growth rate in fresh media. In this way, if the value is above 1, the

supernatant has a positive effect on growth, and if it is below 1 it has a negative effect on

growth. This method gives a measure of the intraspecific effect and interspecific effect of

each species by comparing the relative growth in each species' own supernatant to those in

other species'. No formal statistics were done on the interactions of individual species,

choosing instead to only describe the types and strengths of interactions. These measurements

may help explain why we see coexistence, and can act as a benchmark against which to

understand how interactions change after future experiments.

We looked at whether estimates of species interactions from supernatant assays and

co-culture experiments matched both qualitatively and quantitatively. Quantitatively, we used

standardised major axis (SMA) regression to look at the relationship between the two

estimates for each pairwise combination. SMA was useful here as we have no expectation

about one variable being the response or the predictor, and allows us to test for an expected

slope of 1. Qualitatively, we looked at whether the same sign of interaction (positive or

negative) occurred when calculating pairwise interactions using co-culture and supernatant

assays.

Long-term stability

To understand long-term stability of the five-species community, 12 cultures were transferred

weekly for 60 weeks. To start the cultures, monocultures of species were grown from the

freezer stocks to achieve high cell densities as described above. Populations were set-up

using 20µL inoculum of each species' culture normalised using the species-specific

calibration curves and incubated at 28℃ [47]. Serial 100-fold dilutions (60µL culture into

6mL 1/64 TSB as described earlier) took place every week for over a year. Cultures were

plated every two weeks to check for contamination. After 60 weeks (at least 400 generations),

our plate counts and identification of morphotypes indicated that all 5 species remained

present in all cultures. To confirm this, we sequenced the 16S rRNA gene of multiple

replicates of each morphotype (species) across multiple replicates. We picked at least 30

colonies of each morphotype across all 12 populations, with 169 samples in total. DNA was

extracted by following Qiagen's standard DNeasy UltraClean Microbial protocol, and the

concentration of gDNA in each sample was determined using QuBit. A conserved fragment

from the hypervariable region of the 16s rRNA gene was targeted using N501F and N806R

primers and amplicon sequencing was undertaken by the Centre of Genomic Research

(Liverpool, UK) using an Illumina MiSeq to create 2x250bp paired end reads.

We processed and analysed the sequence data in R using the packages *dada2* and

phyloseq [48]. Following the standard full stack workflow, we estimated error rates, inferred

and merged sequences, constructed a sequence table, removed chimeric sequences and

assigned taxonomy. During processing, forward and reverse reads were truncated between 25

and 250 nucleotide positions due to poor quality scores. Assembled amplicon sequence

variants (ASVs) were assigned taxonomy using the RDP database. We then aggregated OTUs

at the genus level as we knew each of the five species belonged to a different genus. One

sample was removed as it only had 135 reads, and one sample was removed because the

dominant ASV (the identity of the clone) was <0.95, indicating that the colony likely

consisted of multiple species.

Whole genome sequencing of each morphotype

To better understand the genomic features of each species, we did *de novo* hybrid assembly

using both short- and long-read sequencing. DNA was using a NEB monarch genomic DNA

kit following the manufacturer's instructions, apart from for *Ochrobactrum* spp. samples,

which were vortexed with beads and had an extra proteinase-K treatment to aid lysis. Short

read Illumina sequencing was done by the Exeter Sequencing Service on a HiSeq 2500 ran in

standard mode. Long read sequencing was done using the PacBio HiFi platform on a Sequell

II using one SMRT cell by the Centre of Genomic Research (Liverpool, UK) and quality

processed by them.

Hybrid assembly was done using *HybridSPAdes* [49] with the option "*-isolate*"

specified. The assemblies were polished with the trimmed Illumina data using *Pilon* [50]

under default settings. *Circlator* [51] was used to circularise the assemblies, which utilised the

PacBio reads which were corrected using *Canu* [52] with the option "*genomeSize*" set to the

approximate size for each isolate (estimated from the BLAST results of each isolates 16S

sanger sequence). All contigs <500bp were removed, and a contig that was identified as the

phage *phiX* - used as spike in for the short-read sequencing - was removed. Assemblies were

annotated using *prokka* [53] with default settings. We ran *breseq* [54] to map the short reads

back to each assembly to identify potential errors in the assembly and to estimate depth.

Taxonomy was estimated using *gtdbtk* [55] which uses FastANI [56] to do whole genome

alignment, and by using *dada2* to assign taxonomy using 16S genes in the assemblies. The

quality of the assemblies was assessed using *CheckM2* [57].

Results

Final (dis)assembled community

From plating all 48 communities, we identified 5 distinct morphotypes that were dominant in

almost all communities. Based on Sanger sequencing, the five colony morphotypes were

identified as *Achromobacter* sp., *Ochrobactrum* sp., *Pseudomonas* sp., *Stenotrophomonas*

sp., and *Variovorax* sp. (Figure 1b). Overall we have high confidence that these morphs

represent different species. For *Pseudomonas* sp., *Stenotrophomonas* sp., and *Variovorax* sp.,

all morphotype replicates were assigned to the same consensus sequence. For *Ochrobactrum*

sp., 11/12 morphotype replicates were aligned to a single consensus sequence, with the

remaining replicate assigned to the closely related *Rhizobium* sp.. As both sequences are on

the same branch of the global tree, this morph is unlikely to have been misidentified to any

other of the four morphs. For isolate 1 (likely *Achromobacter* sp.), 4/8 replicates were

assigned to *Achromobacter* sp., 3/8 were assigned to *Stenotrophomonas* sp. and 1 replicate

was assigned to *Pseudomonas* sp. (Figure 1C). This may lead to some underreporting of

Achromobacter sp. and overreporting of *Stenotrophomonas* sp., but it is unlikely to

qualitatively alter results of experiments that use this community. We chose a single replicate

community and a single isolate of each colony morphology from which to do all future

experiments.

Reciprocal invasion-from-rare experiments

We determined whether all species were key for coexistence by invading each species from

rare in all species combinations. Species could invade, irrespective of resident community

composition or diversity, in 71 of the 75 combinations (one-sample t-test: $P_{adj} < 0.05$, Figure

2). Moreover, there was considerable variation in relative invader growth rate, ranging from a

minimum of 1.12 to a maximum of 4.34. Of the non-significant combinations, three were

when *Achromobacter* sp. was the invading species (including when it was invading into the

other 4 species together; Figure 2a), the other when *Stenotrophomonas* sp. was invading into

a community of *Achromobacter* sp., *Variovorax* sp. and *Ochrobactrum* sp. However,

although not significant, mean estimates of relative invader growth rate were still above 1.

Interactions based on co-culture and spent media

From doing co-culture assays in all combinations of species diversity, we were able to

explore the nature of species interactions between species pairs in the model community and

also how these changed at higher levels of diversity, allowing us to estimate indirect

interactions. Four of the five species (*Achromobacter* sp., *Ochrobactrum* sp., *Pseudomonas*

sp., *Stenotrophomonas* sp.) had competitive interactions, growing worse in co-culture with

other species compared to monoculture (Figure 3b), with three of the species negatively

impacted by all other species. Although *Variovorax* sp. negatively impacted the growth of all

other species, it grew better in the presence of three of the other species, demonstrating

asymmetric positive interactions (Figure 3b). The only species which had a symmetrically

competitive interaction with *Variovorax* sp. was *Stenotrophomas* sp..

In combinations of more than 2 species, we saw large variation in estimates of

indirect interactions (Figure 4a). Overall, *Ochrobactrum* sp., *Pseudomonas* sp. and

Stenotrophomas sp. had on average negative indirect interactions, which given they had

negative pairwise species interactions, indicates interactions in multispecies culture are more

synergistic than would be expected by pairwise interactions alone. The opposite is true for

Achromobacter sp., where we see on average buffering indirect interactions where

interactions in multispecies culture are less than would be expected by pairwise interactions

alone. *Variovorax* sp. has huge variation in indirect effects, indicating that interactions from

coculture are bad at estimating how it will do in multispecies culture.

When looking at the indirect interactions of all species together (Figure 4), we found

that 41% were synergistic (23 out of 55), with 59% being buffering, when compared to the

expected effect of pairwise interactions. For three- and four-species communities, buffering

was found 40% of the time, and synergistic 60% of the time, with this pattern switching when

all 5 species were present (albeit only with 5 data points). It is important to note that the

definition of synergism and buffering of indirect interactions is sensitive to the null model we

are comparing against, which we set here simply as the impact expected from pairwise

interactions alone [46]. If we included the indirect interactions of three species combinations

into the estimated indirect interactions of four species, we may have seen more buffering [44].

We chose not to do this for simplicity in explanation and interpretation.

No focal species consistently had buffering or synergistic indirect interactions

irrespective of species combination they were co-cultured with (Figure 4a). Across all

indirect interactions, we found that as pairwise interactions intensify, indirect interactions are

more likely to be buffering (lower effect than expected), with synergistic indirect interactions

more likely when the expected pairwise interaction is weaker (Figure 4b). This pattern is

statistically significant for expected negative interactions (ANOVA between models with and

without indirect interaction type as a predictor: $F_{1,47} = 4.51$, $p = 0.039$), and the pattern is the

same for expected positive interactions, albeit a lack of expected positive interactions

prevents us from doing formal statistical analysis.

The supernatant assays allowed us to get an estimate of intraspecific competition, as

well as interspecific competition, with coexistence expected when intraspecific competition is

greater than interspecific competition. In four of the five species (*Pseudomonas* sp. being the

exception, where no difference was detected) intraspecific competition was stronger than the

interspecific effect (Figure 3a). In 3 of the species the strength of intraspecific competition

was at least twice that of the estimate of the lower 95% confidence interval of interspecific

competition, and in *Variovorax* sp. intraspecific competition was twelve times stronger than

interspecific competition.

Both coculture and supernatant assays are commonly used to estimate species

interactions, but little work has compared the two methods. We find a positive relationship

between estimates of interactions from supernatant and coculture (slope = 1.46, 95%CI =

0.86-2.45, Figure 3c), but the model only explains 15% of the variation in the data, meaning

that the fit between the supernatant and coculture estimates is poor. However, qualitatively,

both approaches characterised the interaction as the same sign (positive or negative) in 18 of

the 20 interaction estimates (Figure 3c, inset). In the two instances where they disagreed, $w_{o/a}$

and $w_{s/o}$, one of the estimates was more or less neutral and therefore difficult to define as

either positive or negative.

Long-term coexistence

To examine whether the five-species community was stable long-term, we re-sequenced and

visually identified multiple clones of each morphotype across six communities that had been

cultured for 60 weeks. Crucially, this will also tell us whether morphotype differences still

exist after long-term culturing which is key for the use of model communities in

experimental-evolution studies.

Of the 167 samples that passed quality control, 160 (~96%) were correctly identified

from colony morphology (the genus assigned by the 16S sequencing was the same as that

assigned by the assessor). Of the seven that were mis-assigned, 2 *Achromobacter* sp. were

identified as *Ochrobactrum* sp., 2 *Pseudomonas* sp. were identified as *Variovorax* sp., 2

Variovorax sp. were identified as *Pseudomonas* sp., and 1 *Achromobacter* sp. was identified

as *Pseudomonas* sp.. These mis-assignments do not match reliably with those from the initial

Sanger sequencing but remain extremely rare so are again unlikely to qualitatively alter

results of experiments that use this community. This confirms that the community has long-

term stability and that the individual species can still be identified by their morphotype after

this time.

Genomic composition

Using short- and long-read sequencing, we created high quality reference genomes for each

of the five species. All of the genomes have very high completion (>99.9%) and very low

contamination (<2%), with the worst assembly having only 3 contigs (Table 1). Reassuringly,

taxonomic assignment from GTDBtk corresponds to the genus-level taxon assignment from

the Sanger sequencing, but for 3 of the 5 isolates a species name is now available (Table 1).

Using GTDBtk to assign taxonomy using genome similarity, 4 of the species show >98%

similarity to previously sequenced genomes (Table 1). The isolate genomes have now been

submitted to NCBI and have been given strain names: *Achrombacter veterisilvae* AB1,

Ochrobactrum teleogrylli AB1, *Pseudomonas fluorescens* AB1, *Stenotrophomonas sp.* AB1,

and *Variovorax sp.* AB1. Genomic elements associated with plasmids, prophages, secondary

metabolites, defence systems, and antibiotic resistance are present in all species. These

genomes can be used as high quality references to look at genetic changes in future

experimental evolution experiments.

Discussion

We characterised a model five species community natural community that can be used in

experiments in microbial ecology and evolution. The community contains species with

mostly competitive interactions, but *Variovorax* sp. demonstrates exploitative (potentially

cross-feeding) interactions with 3 of the 4 other species. We found that estimates of

interactions from supernatant assays and coculture assays agreed qualitatively, but not

quantitatively, with estimates of interactions in coculture being stronger on average than

those in supernatant. This suggests that contact dependent interactions (e.g. T6SS and biofilm

formation) may also be important, but it also may just reflect methodological differences in

the approaches. We found evidence of indirect interactions, but they became weaker (and

more likely to be buffering) as expected pairwise interaction increased. This meant that in the

full five species community, indirect interactions were relatively weak.

This community has numerous advantages as a model system for ecological and

evolutionary studies. Sequencing has confirmed that each species has a distinct colony

morphology that means it can be tracked and re-isolated from the other members of the

community, and that these colony morphologies remain distinct through a year of long-term

culture. This makes this community ideal for experimental evolution. Furthermore, we used

invasion-from-rare assays to experimentally demonstrate species coexistence, and sequencing

showed that the community is stable through long-term (>1 year) batch culture. We

encourage - where possible - more explicit tests of coexistence in microbial community

experiments.

Our model community is composed of diverse bacterial species all isolated from the

same soil sample. Even so, this community is not designed to mimic a soil community.

Instead the aim was to create a community that we could show empirically to be stable that

could then be used to investigate general questions about ecology and evolution. The long-

term stability of the community and the unique ability to re-isolate each species to test for

evolutionary responses in phenotype make the community perfect for doing long-term

experimental evolution experiments, a multi-species version of Lenski's pioneering LTEE.

The genome assemblies provide a reference database to track genetic changes through

evolutionary time.

The utility of this 5 species community is demonstrated in the current and previous

work by us, and its potential to test future ideas. For example, we have demonstrated how

species become (mal)adapted after coevolutionary time [47] and how invaders and

disturbances interact synergistically in their effect on resident diversity [58]. While the 5

species community as described only contains bacteria, we have also discovered three phages

that each can infect one of the five species [59] and combined the model community with a

broad host range plasmid which can be carried by all 5 species [60]. Moreover, we have used

this community to demonstrate that fitness effects of plasmids shape the structure of bacteria–

plasmid interaction networks [60]. Given its long-term stability and the distinct morphotypes

of each species, this community is ideal to understand how microbial communities evolve

and how this alters stability in the face of environmental stressors. Furthermore, its stability

means this community is an ideal one to use to measure invasibility, or as a way to validate

and improve Genome Scale Models (GSMs) that use genomic information to predict

metabolic functioning and species interactions [61]. Environmental conditions such as

primary substrate, pH, or temperature could be changed to see how well GSMs do under

different scenarios.

Data availability

All data and code for recreating the analyses and plots from the processed datasets is

available on GitHub: https://github.com/padpadpadpad/five_species_community_paper. The

assembled genomes have been submitted to NCBI and are available under the following

accessions: *Achromobacter veterisilvae* AB1 (CP148753.1), *Ochrobactrum teleogrylli* AB1

(JBBHKQ000000000.1), *Pseudomonas fluorescens* AB1 (CP148752), *Stenotrophomonas sp.*

AB1(2024)(JBBHKO000000000.1), and *Variovorax sp.* AB1(2024)(JBBHKP000000000.1).

The raw sequencing of the original colony identification (Sanger sequencing) is available on

the GitHub repository, and long-term colony identification (16s amplicon sequencing) will be

made available on ENA. The raw short- and long-read whole genome sequencing data can be

made immediately available on request, as can the code that creates the processed datasets.

References

1. **Barber JN, Nicholson LC, Woods LC, Judd LM, Sezmis AL, *et al.*** Species
interactions constrain adaptation and preserve ecological stability in an experimental
microbial community. *ISME J* 2022;16:1442–1452.

2. **Buckling A, Craig Maclean R, Brockhurst MA, Colegrave N**. The Beagle in a bottle.
Nature 2009;457:824–829.

3. **Jousset A, Eisenhauer N, Merker M, Mouquet N, Scheu S**. High functional diversity
stimulates diversification in experimental microbial communities. *Sci Adv*
2016;2:e1600124.

4. **Lenski RE, Rose MR, Simpson SC, Tadler SC**. Long-term experimental evolution in
Escherichia coli. I. Adaptation and divergence during 2,000 generations. *Am Nat*
1991;138:1315–1341.

5. **García FC, Bestion E, Warfield R, Yvon-Durocher G**. Changes in temperature alter
the relationship between biodiversity and ecosystem functioning. *Proc Natl Acad Sci*
2018;115:10989–10994.

6. **Sanchez-Gorostiaga A, Bajić D, Osborne ML, Poyatos JF, Sanchez A**. High-order
interactions distort the functional landscape of microbial consortia. *PLOS Biol*
2019;17:e3000550.

7. **Friedman J, Higgins LM, Gore J**. Community structure follows simple assembly rules in
microbial microcosms. *Nat Ecol Evol* 2017;1:1–7.

8. **Hu J, Amor DR, Barbier M, Bunin G, Gore J**. Emergent phases of ecological diversity
and dynamics mapped in microcosms. *Science* 2022;378:85–89.

9. **Acosta F, Zamor RM, Najar FZ, Roe BA, Hambright KD**. Dynamics of an experimental
microbial invasion. *Proc Natl Acad Sci* 2015;112:11594–11599.

10. **Jones ML, Ramoneda J, Rivett DW, Bell T**. Biotic resistance shapes the influence of
propagule pressure on invasion success in bacterial communities. *Ecology*
2017;98:1743–1749.

11. **Hammarlund SP, Gedeon T, Carlson RP, Harcombe WR**. Limitation by a shared
mutualist promotes coexistence of multiple competing partners. *Nat Commun*
2021;12:619.

12. **Harcombe W**. NOVEL COOPERATION EXPERIMENTALLY EVOLVED BETWEEN
SPECIES. *Evolution* 2010;64:2166–2172.

13. **Ratzke C, Barrere J, Gore J**. Strength of species interactions determines biodiversity
and stability in microbial communities. *Nat Ecol Evol* 2020;4:376–383.

14. **Bittleston LS, Gralka M, Leventhal GE, Mizrahi I, Cordero OX**. Context-dependent
dynamics lead to the assembly of functionally distinct microbial communities. *Nat*
Commun 2020;11:1440.

15. **Estrela S, Sanchez-Gorostiaga A, Vila JC, Sanchez A**. Nutrient dominance governs
the assembly of microbial communities in mixed nutrient environments. *eLife*
2021;10:e65948.

16. **Estrela S, Sánchez Á, Rebolleda-Gómez M**. Multi-replicated enrichment communities
as a model system in microbial ecology. *Front Microbiol* 2021;12:657467.

17. **Goldford JE, Lu N, Bajić D, Estrela S, Tikhonov M, *et al.*** Emergent simplicity in
microbial community assembly. *Science* 2018;361:469–474.

18. **Blasche S, Kim Y, Oliveira AP, Patil KR**. Model microbial communities for ecosystems
biology. *Curr Opin Syst Biol* 2017;6:51–57.

19. **Chesson P**. Mechanisms of maintenance of species diversity. *Annu Rev Ecol Syst*
2000;31:343–366.

20. **Grainger TN, Levine JM, Gilbert B**. The invasion criterion: A common currency for
ecological research. *Trends Ecol Evol*.

21. **Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, *et al.*** Bacterial Community
Variation in Human Body Habitats Across Space and Time. *Science* 2009;326:1694–
1697.

22. **Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A, Eldridge DJ,**
et al. A global atlas of the dominant bacteria found in soil. *Science* 2018;359:320–325.

23. **Fierer N, Jackson RB**. The diversity and biogeography of soil bacterial communities.
Proc Natl Acad Sci 2006;103:626–631.

24. **Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH**. Relative roles of niche and
neutral processes in structuring a soil microbial community. *ISME J* 2010;4:337–345.

25. **Liao J, Cao X, Zhao L, Wang J, Gao Z, *et al.*** The importance of neutral and niche
processes for bacterial community assembly differs between habitat generalists and
specialists. *FEMS Microbiol Ecol* 2016;92:fiw174.

26. **Ofiţeru ID, Lunn M, Curtis TP, Wells GF, Criddle CS, *et al.*** Combined niche and
neutral effects in a microbial wastewater treatment community. *Proc Natl Acad Sci*
2010;107:15345–15350.

27. **Godwin CM, Chang F, Cardinale BJ**. An empiricist's guide to modern coexistence
theory for competitive communities. *Oikos* 2020;129:1109–1127.

28. **Spaak JW, Schreiber SJ**. Building modern coexistence theory from the ground up: The
role of community assembly. *Ecol Lett* 2023;26:1840–1861.

29. **Chesson P**. Updates on mechanisms of maintenance of species diversity. *J Ecol*
2018;106:1773–1794.

30. **Pande S, Merker H, Bohl K, Reichelt M, Schuster S, *et al.*** Fitness and stability of
obligate cross-feeding interactions that emerge upon gene loss in bacteria. *ISME J*
2014;8:953–962.

31. **Preussger D, Giri S, Muhsal LK, Oña L, Kost C**. Reciprocal Fitness Feedbacks
Promote the Evolution of Mutualistic Cooperation. *Curr Biol* 2020;30:3580-3590.e7.

32. **Zhao L, Zhang Q-G, Zhang D-Y**. Evolution alters ecological mechanisms of coexistence
in experimental microcosms. *Funct Ecol* 2016;30:1440–1446.

33. **Hammarlund SP, Chacón JM, Harcombe WR**. A shared limiting resource leads to
competitive exclusion in a cross-feeding system. *Environ Microbiol* 2019;21:759–771.

34. **Chang C-Y, Bajić D, Vila JCC, Estrela S, Sanchez A**. Emergent coexistence in
multispecies microbial communities. *Science* 2023;381:343–348.

35. **Le Gac M, Plucain J, Hindré T, Lenski RE, Schneider D**. Ecological and evolutionary
dynamics of coexisting lineages during a long-term experiment with Escherichia coli.
Proc Natl Acad Sci 2012;109:9487–9492.

36. **Turner PE, Souza V, Lenski RE**. Tests of Ecological Mechanisms Promoting the Stable
Coexistence of Two Bacterial Genotypes. *Ecology* 1996;77:2119–2129.

37. **Hesse E, O'Brien S, Tromas N, Bayer F, Luján AM, *et al.*** Ecological selection of
siderophore-producing microbial taxa in response to heavy metal contamination. *Ecol*
Lett 2018;21:117–127.

38. **Chao K-H, Barton K, Palmer S, Lanfear R**. sangeranalyseR: Simple and Interactive
Processing of Sanger Sequencing Data in R. *Genome Biol Evol* 2021;13:evab028.

39. **Paradis E, Schliep K**. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* 2019;35:526–528.

40. **Bodenhofer U, Bonatesta E, Horejš-Kainrath C, Hochreiter S**. msa: an R package for
multiple sequence alignment. *Bioinformatics* 2015;31:3997–3999.

41. **Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, *et al.*** DADA2: high-
resolution sample inference from Illumina amplicon data. *Nat Methods* 2016;13:581.

42. **Maidak BL, Cole JR, Lilburn TG, Parker Jr CT, Saxman PR, *et al.*** The RDP
(ribosomal database project) continues. *Nucleic Acids Res* 2000;28:173–174.

43. **Benjamini Y, Hochberg Y**. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. *J R Stat Soc Ser B Methodol* 1995;57:289–300.

44. **Tekin E, White C, Kang TM, Singh N, Cruz-Loya M, *et al.*** Prevalence and patterns of
higher-order drug interactions in Escherichia coli. *Npj Syst Biol Appl* 2018;4:1–10.

45. **Tekin E, Savage VM, Yeh PJ**. Measuring higher-order drug interactions: a review of
recent approaches. *Curr Opin Syst Biol* 2017;4:16–23.

46. **Tekin E, Yeh PJ, Savage VM**. General form for interaction measures and framework for
deriving higher-order emergent effects. *Front Ecol Evol* 2018;6:166.

47. **Castledine M, Padfield D, Buckling A**. Experimental multi-species microbial (co)
evolution results in local maladaptation. *Ecol Lett*.

48. **McMurdie PJ, Holmes S**. phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. *PloS One* 2013;8:e61217.

49. **Antipov D, Korobeynikov A, McLean JS, Pevzner PA**. hybridSPAdes: an algorithm for
hybrid assembly of short and long reads. *Bioinformatics* 2016;32:1009–1015.

50. **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, *et al.*** Pilon: an integrated tool for
comprehensive microbial variant detection and genome assembly improvement. *PloS*
One 2014;9:e112963.

51. **Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, *et al.*** Circlator: automated
circularization of genome assemblies using long sequencing reads. *Genome Biol*
2015;16:294.

52. **Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, *et al.*** Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Res 2017;27:722–736.

53. **Seemann T**. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*
2014;30:2068–2069.

54. **Deatherage DE, Barrick JE**. Identification of Mutations in Laboratory-Evolved Microbes
from Next-Generation Sequencing Data Using breseq. In: Sun L, Shou W (editors).
Engineering and Analyzing Multicellular Systems. New York, NY: Springer New York.
pp. 165–188.

55. **Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH**. GTDB-Tk v2: memory friendly
classification with the genome taxonomy database. *Bioinformatics* 2022;38:5315–5316.

56. **Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S**. High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*
Commun 2018;9:5114.

57. **Chklovski A, Parks DH, Woodcroft BJ, Tyson GW**. CheckM2: a rapid, scalable and
accurate tool for assessing microbial genome quality using machine learning. *Nat*
Methods 2023;20:1203–1212.

58. **Lear L, Inamine H, Shea K, Buckling A**. Diversity loss from multiple interacting
disturbances is regime-dependent. *Ecol Lett* 2023;26:2056–2065.

59. **Castledine M, Newbury A, Lewis R, Hacker C, Meaden S, *et al.*** Antagonistic Mobile
Genetic Elements Can Counteract Each Other's Effects on Microbial Community
Composition. *mBio* 2023;14:e00460-23.

60. **Newbury A, Dawson B, Klümper U, Hesse E, Castledine M, *et al.*** Fitness effects of
plasmids shape the structure of bacteria–plasmid interaction networks. *Proc Natl Acad*
Sci 2022;119:e2118361119.

61. **Frioux C, Singh D, Korcsmaros T, Hildebrand F**. From bag-of-genes to bag-of-
genomes: metabolic modelling of communities in the era of metagenome-assembled
genomes. *Comput Struct Biotechnol J* 2020;18:1722–1734.

627
628

Figures and tables

Figure 1. Community (dis)assembly experiment to form a stable community. (a)

Schematic of the experimental design for the formation of the stable microbial community.

(b) Five morphs dominated the community. (1) *Achromobacter* sp., the smallest colony,

circular and uniform in size. (2) *Ochrobactrum* sp. white/grey in colour and opaque. (3)

Pseudomonas sp. is the largest colony, cream in colour and opaque. (4) *Stenotrophomonas*

sp. is typically larger than *Achromobacter* sp. although variable in size and has a stronger

colour. (5) *Variovorax* sp. has a spreading mat surrounding the colony and is variable in size.

(c) Reliability of morphotypes for identifying the different species.

Figure 2. Ability of species to invade-from-rare in all combinations of species and diversity. Panels represent the invader species. Colours of pie points reflect the mix of the resident community. Black points are observed values of relative invader growth rate (n~6), pie points are the estimate of relative invader growth rate from each one-sample t-test, and error bars represent lower and upper 95% confidence intervals. Shaded regions separate different levels of resident community diversity; dashed, vertical line drawn where relative invader growth rate = 1, values above which indicate the invading species can successfully re-establish from rare.

Figure 3. Estimated pairwise interactions of the model community from supernatant and coculture assays. (a) Growth of each species was in the supernatant of each species (including itself) and fresh media. (b) Growth of each species was measured in coculture and in monoculture. (c) Relationship between estimates of interactions estimated from supernatant assays and coculture. In all plots, values below one indicate negative interactions. In (a) and (b) large coloured points are the mean value of all interspecific interactions with 95% confidence intervals, and small coloured points represent individual interspecific interactions (the median value of three technical replicates). In (c) the solid fitted line is the best fit of a standardised major axis regression, the dashed lines demonstrate where interactions go from being negative to positive, and the inset plot demonstrates how often the interactions are qualitatively the same type using coculture and supernatant assays.

Figure 4. Estimated indirect interactions of the model community from coculture
assays. (a) Total indirect interactions across levels of diversity show lots of variation across
all focal species. (b) Impact of the strength of negative and positive pairwise interactions on
whether the indirect interaction is synergistic (stronger than expected) or buffering (less than
expected). As negative and positive pairwise interactions get stronger, there is a tendency for
the indirect interaction to be buffering. In (a) and (b) large, coloured points are the mean
value of all interspecific interactions with 95% confidence intervals, and small coloured
points represent individual indirect interactions (the median value of three technical
replicates).

671   **Table 1. Genome characteristics and taxonomic assignment of the 5 species genomes.**

Isolate	NCBI submission name	Sanger sequence assignment	GTDBtk assignment	16s copy number	Genome size (bp)	Number of contigs	GC content	Total coding sequences	Completeness	Contamination
1	*Achromobacter veterisilvae* AB1	Achromobacter	*Achromobacter veterisilvae*	4	6,549,044	1	0.67	5,991	100.00	0.98
2	*Ochrobactrum teleogrylli* AB1	Ochrobactrum	*Ochrobactrum_B teleogrylli*	1	5,064,753	3	0.57	4,714	99.97	1.41
3	*Pseudomonas fluorescens* AB1	Pseudomonas	*Pseudomonas_E fluorescens_BH*	2	6,759,727	1	0.60	6,053	100.00	0.15
4	*Stenotrophomonas* sp. AB1 (2024)	Variovorax	*Variovorax sp003019815*	1	6,611,336	2	0.68	6,041	99.99	0.37
5	*Variovorax* sp. AB1 (2024)	Stenotrophomonas	*Stenotrophomonas spp.*	1	3,926,254	2	0.66	3,384	100.00	0.04

673