Biochemical, structural insights of newly isolated AA16 family of Lytic Polysaccharide Monooxygenase (LPMO) from *Aspergillus fumigatus* and investigation of its synergistic effect using biomass.

Musaddique Hossain, Subba Reddy Dodda, Bishwajit Singh Kapoor, Kaustav Aikat, and Sudit S. Mukhopadhyay*

Department of Biotechnology, National Institute of Technology Durgapur-713209, West Bengal, India

Running title: Biochemical, structural insights, and investigation of the synergistic effect of newly isolated AA16 family of Lytic Polysaccharide Monooxygenase (LPMO) from *Aspergillus fumigatus*.

* To whom the corresponding author should be addressed.

E-mail: suditmukhopadhy@yahoo.com

Phone: +919434788139

Abstract

The efficient conversion of lignocellulosic biomass into fermentable sugar is a bottleneck for the cheap production of bio-ethanol. The recently identified enzyme Lytic Polysaccharide Monooxygenase (LPMO) family has brought new hope because of its boosting capabilities of cellulose hydrolysis. In this report, we have identified and characterized a new class of auxiliary (AA16) oxidative enzyme LPMO from the genome of a locally isolated thermophilic fungus *Aspergillus fumigatus* (NITDGPKA3) and evaluated its boosting capacity of biomass hydrolysis. The *Af*LPMO16 is an intronless gene and encodes the 29kDa protein. While Sequence-wise, it is close to the C1 type of *Aa*AA16 and cellulose-active AA10 family of LPMOs, but the predicted three-dimensional structure shows the resemblance with the AA11 family of LPMO (PDB Id: 4MAH). The gene was expressed under an inducible promoter (AOX1) with C-terminal His tag in the *Pichia pastoris*. The protein was purified using Ni-NTA affinity chromatography, and we studied the enzyme kinetics with 2,6-dimethoxyphenol. We observed polysaccharides depolymerization activity with Carboxymethyl cellulose (CMC) and Phosphoric acid swollen cellulose (PASC). Moreover, the simultaneous use of cellulase cocktail (commercial) and *Af*LPMO16 enhances lignocellulosic biomass hydrolysis by 2-fold, which is highest so far reported in the LPMO family.

Importance

The auxiliary enzymes, such as LPMOs, have industrial importance. These enzymes are used in cellulolytic enzyme cocktail due to their synergistic effect along with cellulases. In our study, we have biochemically and functionally characterized the new AA16 family of LPMO from *Aspergillus fumigatus* (NITDGPKA3). The biochemical characterization is the fundamental scientific elucidation of the newly isolated enzyme. The functional characterization, biomass degradation activity of *Af*LPMO16, and cellulase cocktail (commercial) combination enhancing the activity by 2-fold. This enhancement is the highest reported so far, which gives the enzyme *Af*LPMO16 enormous potential for industrial use.

Keywords: *A.fumigatus*, Auxiliary activity, Cloning, Kinetics, LPMO, Lignocelluloses, Molecular docking

Introduction

The diminution of fossil fuels and the growing concern of environmental consequences, particularly climate changes, have steered our fast-growing economy for clean and renewable energy production [1]. Among different renewable energy sources, bioethanol is one of the promising alternatives to fossil fuel because of its low CO_2 emission [2, 3] and its manufacturing reliance on lignocellulosic biomass, which is bio-renewable and abundance on earth. However, the structural complexity and the recalcitrance of this renewable carbon source [4] have hindered its optimal use. The current process of saccharification of lignocellulosic biomass is time-consuming and costly. Therefore, the requirement of cost-effective and fast controlled destruction of lignocellulose has driven the bioethanol industry to explore the accessory enzymes to achieve a better and efficient enzyme cocktail for the commercial production of lignocellulose-derived ethanol.

A breakthrough in such exploration came into existence when a mono-copper redox enzyme, known as Lytic polysaccharide monooxygenase (LPMO), was first reported in 2010 [5-8]. LPMO increases lignocellulosic biomass conversion efficiency[9,10] by catalyzing the hydroxylation of C1 and/or C4 carbon involved in glycosidic bonds that connect glucose unit in cellulose and allow cellulase enzymes to process the destabilized complex polysaccharides [11-15]. Harris et al., in their study, used LPMO from *T reesei* along with classical cellulases and showed that the degradation of polysaccharide substrates was increased by a factor of two when compared with the activity of classical cellulases alone [16]. A CBM33 domain-containing enzyme identified from *Serratia marcescens* with boosting chitinase activity, later classified as LPMO. A study by Nakagawa et al. showed that an AA10 family of LPMO from *Streptomyces griseus* could increase the efficiency of chitinase enzymes by 30- and 20-fold on both α and β forms of chitin, respectively [17]. Along with this work, there are some recent reports of the synergistic effect of LPMOs with glycoside hydrolases on polysaccharide substrates [18-20].

LPMOs are classified as AA9, AA10, AA11, AA13, AA14, and AA15 in the CAZy database (http://www.cazy.org/), based on their amino acid sequence similarity. Recently Filiatrault-Chastel et al. identified the AA16, a new family of LPMO from the secretome of a fungi *Aspergillus aculeatus* (*Aa*AA16). The *Aa*AA16 was initially isolated as X273 protein (unnamed domain) and later identified as C1-oxidizing LPMO active on cellulose [21]. *Aa*AA16, the only AA16 family of LPMO so far, has been identified, and it lacks complete

biochemical characterization. The biochemical characterization, structural characterization,

and the assessment of biomass conversion efficiency are required to understand better the

action of members of this new family on plant biomass and their possible biological roles.

While we were analyzing the cellulose hydrolyzing genes from the genome of *A. fumigatus*

(*Aspergillus* genome database)*,* we identified five LPMOs, one belonging to AA16 family

because of its X273 domain. Further, we cloned the *Af*LPMO16 gene from the genome of our

locally isolated strain of *A. fumigatus* (NITDGPKA3) [22] (GenBank accession No.

JQ046374) by designing the primers based on the *A. fumigatus* LPMO sequence

(CAF32158.1)(NCBI). The cloned *A. fumigatus* (NITDGPKA3) LPMO (after cloning and

sequencing the sequence submitted to GenBank; accession No. MT462230) is expressed in

Pichia pastoris X33. The heterologous protein (*Af*LPMO16) purified and used for

biochemical and functional characterization. The saccharification rate assessment suggests

that *Af*LPMO16 has fast and effective glucose releasing ability from lignocellulose and

cellulose when used with a commercial cellulase cocktail. Enzyme kinetics using 2,6-

dimethoxyphenol as a substrate [23] confirmed the oxidative activity. The lignocellulosic

biomass (alkaline pre-treated raw rice straw) conversion efficiency along with cellulases

suggests that *Af*LPMO16 could be an essential member of the cellulase cocktail for industrial

use.

Results

Cloning, expression, and purification of AfLPMO16

*Af*LPMO16 (GenBank accession No. MT462230) is an intronless 870 nucleotide long gene

that encodes 290 amino acids. The theoretical molecular mass is 29KDa (including signal

peptide). The gene sequence of AA16 from our isolated strain of *A.fumigatus* (NITDGPKA3)

has shown almost 99.6% homology with the gene sequence of AA16 present in the genome

database of *A.fumigatus* (CAF32158.1) (NCBI database).

The protein of *Af*LPMO16 (GenBank accession No. MT462230) was produced in *Pichia*

pastoris X33 without its C-terminal extension. After the optimization of the expression

procedure, we achieved approximately 0.8 mg/ml of purified protein. The SDS-PAGE

analysis (Fig 1) confirmed the single band of the purified protein (Fig. 1: lanes 5 and 6). We

further confirmed the purified recombinant protein bearing the 6X His-tag by Western blot

using an anti-His antibody (Fig. 1: Lane W1 & W2); the purified protein (lane 5 & 6 of SDS-

PAGE) used for western blot. The expressed recombinant *Af*LPMO16 band appeared at approximately 32kDa position in SDS-PAGE (Fig. 1), which is slightly higher than the expected size. It is probably due to glycosylation [24], or recombinant protein has *c-myc* epitope and 6x His tag in its c-terminal that can increase the molecular mass by 2.7KDa. For further confirmation of N-glycosylation, we checked the *Af*LPMO16 sequence glycosylation site using NetNGlyc 1.0 server (DTU Bioinformatics, Technical University of Denmark, http://www.cbs.dtu.dk/services/NetNGlyc/) [36]. There were two N-glycosylation sites present above the 0.5 threshold value at 114 & 149 amino acid sequence positions with 0.76 and 0.56 potential values, respectively.

Enzyme assay and Kinetics

LPMO converts 2,6-dimethoxyphenol (2,6-DMP) into 1-Coerulignone (Fig. 2a) due to its oxidative property, and 1-Coerulignone has an extinction coefficient of $53200 M^{-1} cm^{-1}$. 1-Coerulignone gives absorbance at 469nm wavelength; therefore, we can easily quantify it using a spectrophotometer [21]. The OD at 469nm wavelength steadily increases with time that clearly indicates the steady conversion of 2,6-dimethoxyphenol to 1-Coerulignone (Fig. 2a). It also suggests the sufficient activity of the enzyme *Af*LPMO16. Temperature and pH influence the activity of LPMO. Thus, during the kinetic study, we used optimum temperature 30°C and pH 6.0, as described by [21]. *Af*LPMO16 showed proper activity for the chemical substrate 2,6-dimethoxyphenol; there was a steady release of 1-Coerulignone when incubated 2,6-dimethoxyphenol with *Af*LPMO16. The enzyme kinetics was performed with different concentrations of 2,6-dimethoxyphenol. We obtained the Kinetics parameters such as Michaelis Menten constant (K_m) and maximum velocity (V_{max}) from the Line-weaver-Burk plot (Fig. 2b) as 5.4mM, and 0.153 U/mg, respectively. The calculated catalytic activity K_{cat} was 277.67 min^{-1} (Table 1). These kinetics parameters suggest that the oxidative property of *Af*LPMO16.

In-silico analysis for substrate specificity

The *Af*LPMO16 contains 19 amino acids long N-terminal signal peptide before His1 catalytic domain (1-169aa), and C terminal Serine rich region (170-271aa) (Fig. 3a). This N-terminal sequence is one of the marker features of fungal LPMOs, but this serine-rich C-terminal or linker is a feature of AA16 family. It also lacks the CBM1 module or glycosylphosphatidylinositol (GPI) anchor, like other AA16 LPMOs [19]. *Af*LPMO16 also

has conserved Histidine at 1^{st} and 109^{th} positions, which are mainly involved in copper binding, the signature characteristic of LPMOs. There are other conserved sequences like Gly, Pro, Asn, Cys, Try, Tyr, Leu, and Asp, including GNV(I)QGELQ motif (Fig. 3b) The fully conserved sequences (highlighted with red background) are the marker amino acids represent the LPMOs. The partially conserved sequences (within the blue boxes) are the marker of different auxiliary families (Fig. 3b). The sequence alignment studies of AA16 family (including *Af*LPMO16) with other families (AA9, AA10, and AA11) of LPMOs suggested (Fig. S1) a co-relation between AA10 family and AA16 LPMOs. The substrate-binding motif in the L2 loop of cellulose active LPMO10 has some similarities with AA16 L2 loop motif (marked with black box) and cellulose active motif (Fig. 3b). In AA16 LPMOs the conserved motif in L2 loop GNI(V)QGEL the region is replaced by YNWFG(A)NL for C1 oxidizing AA10 LPMOs, which are also cellulose active. The previous study suggests that the amino acids (Y79, N80, F82, Y111, and W141) in loop L2 take part in substrate specificity for LPMO 10, and mutations (Y79, N80D, F82A, Y111F, W141Q) alter the specificity of the substrate from chitin to cellulose [37]. In *Af*LPMO16, the corresponding amino acids GNQYR (Fig. 3b) (marked with black arrows), some amino acids from these positions (N & Y) are also present in cellulose-active AA10 LPMOs. Hopefully, the polar amino acids (Q & R) are charged and may interact with chitin due to electrostatic interaction. Alternatively, there are high chances that few mutations in these amino acids may help *Af*LPMO16 to interact with chitin. Further, in chitin active LPMOs, more than 70% residues of the motif (Y(W)EPQSVE) are polar, including two negatively charged Glu (E). In cellulose active LPMOs, 70% residues of the motif (Y(W)NWFGVL) are hydrophobic [38]. In contrast, in *Af*LPMO16, 70% residues are polar, including one negatively charged Glu (E), one hydrophobic Tyr (Y), and others are neutral. The presence of polar residue and negative charged Glu (E) suggests that *Af*LPMO16 may bind to chitin. Electrostatics interaction between the substrate and enzyme active site plays a pivotal role in substrate binding. The electrostatic potential surface at the catalytic site of the *Af*LPMO16 was found unchanged or slightly positive-charged at pH 6.0 (Fig. 3c) (Marked in the figure). The electrostatic interaction study suggests that the *Af*LPMO16 may also bind to cellulose [52].

Regioselectivity of AfLPMO16

Amino acids on the substrate-binding surface determine the oxidative regioselectivity of LPMOs [29]. Sequence comparison and mutation studies revealed that the conserved amino

acids near the catalytic center in C1 and C1/C4 oxidizing AA10 and AA9 LPMOs are

responsible for regioselectivity. In the case of C1/C4 oxidizing AA10, the amino acid Asn85

near the catalytic center is responsible for C4 oxidizing activity. Alteration of this amino acid

(N85F) diminished the C4 activity and produced only C1 oxidized product [39]. In C1

oxidizing AA9 LPMOs, hydrophobic amino acids Phe and Tyr are conserved in addition to

Asn. While in C1 oxidizing AA10 LPMOs, the Phe amino acid has replaced the

corresponding Asn site (Fig. 3b)(marked with red arrow). The Phe is also parallel to the

substrate-binding surface [47]. In AA16, the corresponding Gln (Q) may be parallel to the

substrate-binding region (Fig. 3b). The function of conserved Gln (Q) is not clear. However,

this polar amino acid has a similar side chain with polar Asn (N). The axial distance between

the conserved amino acid and copper catalytic center is another crucial factor for

regioselectivity. The C1/C4 oxidizing AA10 LPMOs have more open or wider axial gaps

than C1 oxidizing AA10 LPMOs [39]. Here the distance between Gln56 and His20 is 7.7Å,

and the distance between Gln56 and Cu catalytic center is 11.1Å. In the absence of the AA16

structure (crystal or model), we cannot compare the lengths; nevertheless, this distance may

play a key role in regioselectivity.

Phylogenetic tree construction and analysis

The sequential and functional relationship of AA10 and AA16 LPMOs has been discussed,

but phylogenetic studies based on the sequence similarity give an evolutionary origin. Based

on sequence comparison, *Af*LPMO16 is evolutionarily closer to the LPMO of *Aspergillus*

fisheri (91% sequence homology). The constructed phylogenetic tree contains two main

clades and two subclades (Fig. 4). The first clade contains all AA10 LPMOs from bacterial

species such as *Bacillus thuringiensis*, *Bacillus amyloliquefaciens, Streptomyces lividans,* and

Enterococcus faecalis. The second clade includes all fungal AA10 and AA16LPMOs, mainly

belongs to *Aspergillus,* and *Penicillium* species in which AA16 LPMOs are mostly from

A.niger, *A.fumigatus*, *A.fisheri*, *Aspergillus kawachii* (Fig. 4).

Model structure prediction and molecular docking analysis

I-TASSER was used to predict the three-dimensional structure of the *Af*LPMO16. Most of the

LPMOs have immunoglobulin-like distorted β-sandwich fold like structures, in which loops

connect seven antiparallel β-strands with a different number of α-helix insertions (Fig. 5a).

The final model has a β-sandwich structure connected by loops with two α-helices. The

superimposition of the *Af*LPMO16 with other LPMO families like AA9, AA10, AA11, and AA13 showed that they share common antiparallel β-strands and helices with more loops, which indicate higher flexibility. Moreover, *Af*LPMO16 showed 1.2Å RMSD with AA11 (PDB Id: 4MAH) LPMO lower than the other LPMOs. So the 3D structure of *Af*LPMO16 suggests that it has more structural resemblance with AA11 LPMO. We also found one disulfide bond in *Af*LPMO16 between the Cys78-Cys186 amino acids, signature of thermo-stability (Fig. S2). The histidine brace amino acids, such as His20 and His109, participate in coordination bond with Cu ions. The surface of *Af*LPMO16 has an active site (Fig. 5b). The interaction studies with cellohexose suggest amino acids like Gln48, Gln181, Ser178, His109, His20, Asn54, Asp50, Tyr52, and Glu58 (Active enzyme starts with His1; so His20 will His1 and corresponding amino acids can be numbered accordingly) are in the active site and are involved in the interaction with the substrate (Fig. 5c). Molecular docking suggests that *Af*LPMO16 has a cellulose-binding surface (Fig. 5b & 5c). This study also suggests that the binding energy between *Af*LPMO16 and cellulose is -7.0 kcal/mol, which is highest compared to chitin (-5.5kcal/mol) and other polysaccharides.

Polysaccharides depolymerization by AfLPMO16

*Af*LPMO16 showed efficient depolymerization activity on both CMC and PASC (Fig. 6a & 6b). We quantified the amount of reducing sugar released by enzymatic degradation. When incubated CMC with increasing concentrations of the enzyme, the amount of product (reducing sugar) increased with the increase of *Af*LPMO16 concentration (Fig. 6a). When we added 50µg of the enzyme, nearly 0.05mg/ml of reducing sugar was released. For 100µg of the enzyme, the product was nearly 0.136mg/ml, and for 200µg of the enzyme, the amount of product released was approximately 0.356mg/ml (Fig. 6a). This result indicates the polysaccharide (CMC) depolymerization activity of *Af*LPMO16.

Further, we used insoluble PASC as a substrate and incubated with an increasing concentration of *Af*LPMO16, and determined the relative absorbance of PASC with the growing amount of enzyme. The enzyme degrades the polysaccharide (substrate) into smaller polysaccharide units (monosaccharides, disaccharides, etc.), which are soluble and make the reaction mixture clearer. Therefore, it leads to a decrease in the absorbance resulting increment in relative absorbance [40]. Ultimately we will find a graph where relative absorbance increase with increasing concentration of *Af*LPMO16. Hence In this experiment, we found a rise in relative absorbance concerning the untreated substrate with a high

concentration of enzyme *Af*LPMO16 (Fig. 6b). The graph (Fig. 6b) showed that 0.17
absorbance difference concerning untreated substrate when we used 50µl (concentration
0.8µg/µl) of the enzyme. The difference in absorbance steadily increased with the escalation
of enzyme concentration (200µl of the enzyme at the concentration of 0.8µg/µl the relative
absorbance reached nearly 0.36). Hence these experiments confirmed the intrinsic
polysaccharide degradation property of the *Af*LPMO16 like other LPMOs. In these
experiments, we used the heat-inactivated *Af*LPMO16 and ascorbic acid-deficient set to
verify these results (data not shown).

Pre-treated lignocellulosic biomass and cellulose hydrolysis with simultaneous treatment of
AfLPMO16 and commercial cellulase

There are two modes of action to show the synergy or boosting effect of LPMO while using
with cellulase- sequential assay and simultaneous assay. In the sequential assay, LPMO
should add a prior time limit to cellulase. And in the simultaneous assay, both the enzymes
LPMO and cellulase are being used together to the substrate. In this study, we chose to
perform a simultaneous assay for two reasons; simultaneous assay shows better synergy or
boosting in crystalline cellulose [41] than sequential one. Furthermore, we aimed to check the
synergy or stimulating activity of commercial cellulase by *Af*LPMO16 so that it may include
in the cocktail for better depolymerizing action. Here the boosting effect of *Af*LPMO16 was
studied with a commercial cellulase cocktail on both cellulose (Avicel) and lignocellulosic
biomass (alkaline pre-treated rice straw). The alkaline pre-treatment has a beneficiary over
acid pre-treatment in terms of hydrolysis yield [48]. The reason is that alkaline pre-treatment
sufficiently removes the lignin [42], but it preserves hemicelluloses [43]. When incubating
Avicel with *Af*LPMO16 and cellulase, the amount of reducing sugar released was almost
double compared to Avicel incubated with either cellulase alone or cellulase along with heat-
inactivated *Af*LPMO16 (Fig. 7b). A similar kind of boosting effect we observed in every
time point from 5 hrs to 72 hrs. We also found the synergistic impact of *Af*LPMO16 in
lignocellulosic biomass transformation to fermentable sugar (Fig. 7a). When we incubated
the alkaline pre-treated rice straw with 100 µg and 200µg of *Af*LPMO16 along with cellulase,
almost 1.7 fold and slightly above 2-fold of reducing sugar were released respectively
compared to lignocellulose incubated with either cellulase alone or cellulase along with heat-
inactivated *Af*LPMO16 (Fig. 7a) suggests the enhancement is dependent on the amount of
auxiliary enzyme *Af*LPMO16. For further elaboration of the synergistic effect of *Af*LPMO16,

another set of reactions prepared where the biomass was treated with an increasing concentration of only *Af*LPMO16. A minimal amount of hydrolysis activity was there, nearly 0.04 mg/ml to 0.06 mg/ml, reducing sugar quantified for *Af*LPMO16 treated biomass (Fig. 7c). This hydrolysis activity of *Af*LPMO16 alone is negligible compare to only cellulase treated biomass.

Nevertheless, the simultaneous use of *Af*LPMO16 and cellulase enhances the hydrolysis activity two-fold compared to the only cellulase treated biomass (Fig. 7c). This result strongly indicates the synergistic effect of *Af*LPMO16 with cellulase. All these results confirmed the boosting effect or synergistic effect of *Af*LPMO16 on the hydrolytic activity of cellulase for both cellulosic and lignocellulosic biomass degradation. So far highest synergistic effect was reported by AA9 (Table 2), which is less than two-fold [44, 45].

Discussion

The gene was cloned in pPICZαA vector under the control of *AOX1* promoter by following the same strategy developed for *Aa*AA16 and PMO9A_MLACI [19, 26]. The nucleotide sequence of *Af*LPMO16 was codon-optimized for *Pichia pastoris*. The recombinant protein containing a C-terminal polyhistidine tag was produced in flasks in the presence of trace metals, including copper, and purified from the culture supernatant by immobilized metal ion affinity chromatography (IMAC: Ni-NTA affinity chromatography), following the same protocol used for *Aa*AA16 [19]. We were successful in producing the active *Af*LPMO16 in *P.pastoris* X33 (Fig. 1) in a shake flask. Despite the chance of N-terminal modification in shake flask culture instead of bioreactor culture [19], the amount of active enzyme obtained in shake flask was sufficient for characterization. The enzyme activity determined by 2,6-dimethoxyphenol concerning the heat-inactivated enzyme and without ascorbic acid as negative controls (data not shown). The enzyme activity suggests the successful production of active protein (Fig. 2a), and interestingly, the initial reaction rate is faster compared to later time span. Lytic polysaccharide monooxygenase (LPMO) releases a spectrum of cleavage products from their polymeric substrates cellulose, hemicellulose, or chitin. The correct identification and quantitation of these released products is the basis of MS/HPLC-based detection methods for LPMO activity, which is time taking and is required specialized laboratories to measure LPMO activity in day-to-day work. A spectrophotometric assay based on the 2,6-dimethoxyphenol can accurately measure the enzymatic action and can be used for enzyme screening, production, and purification, and can also be applied to study

enzyme Kinetics [21]. Thus it is swift, robust for biochemical characterization, and also
accurately determines the active enzyme.

Sequence analyses indicating that the *Af*LPMO16 has some signature characteristics for both
cellulose and chitin-binding and both C1 and C1/C4 oxidizing activity. However,
experimental confirmation is required to establish the presence or absence of any chitin-
binding nature and C1/C4 oxidizing capability of *Af*LPMO16. The constructed phylogenetic
tree (Fig. 4) suggests that the fungal AA10 and AA16 LPMOs are more likely to come from a
common ancestor. Molecular docking study suggests that *Af*LPMO16 has the highest affinity
towards cellulose among the known substrates, based on the binding energy. The binding
energy between cellulose and *Af*LPMO16 is -7.0 Kcal/mol, which makes thermodynamically
strong binding between enzyme and substrate (Fig. 5b & 5c) compared to other substrates.
The LPMOs are essential for their auxiliary activity and polysaccharide degrading property.
We observed polysaccharide depolymerizing activity on carboxymethyl cellulose (CMC) and
phosphoric acid swollen cellulose (PASC) (Fig. 6a & 6b). Due to its auxiliary activity, it
enhances the action of the cellulase enzyme for the degradation of cellulose and
lignocelluloses [49]. The only identified AA16 family, the *Aa*AA16, showed a sequential
boosting effect with *T. reesei* CBHI on nano-fibrillated cellulose (NFC) and PASC. The
*Aa*AA16, the recent addition of the AA16 family of LPMO in the CAZY database, showed
synergism with the CBH1 for the degradation of cellulose [19]. However, *Aa*AA16 study did
not deal with the biomass hydrolysis boosting effect of the AA16 family. The boosting result
is most important in the technical aspect for enhancing the activity of the cellulase cocktail.
LPMO enzyme has earned much research interest due to their synergistic effect or boosting
effect on cellulase enzyme [45]. *Af*LPMO16 showed a boosting impact on cellulose and
lignocellulose hydrolysis (Fig. 7a & 7b). The synergism of *Af*LPMO16 has shown in (Fig.
7c), where the only *Af*LPMO16 and only cellulase treated biomass hydrolysis activity is low
compare to the combined effect of these two enzymes. The simultaneous use of *Af*LPMO16
and cellulase enhances nearly two-fold biomass hydrolysis compare to the only cellulase
treated biomass hydrolysis. This enhancement of two-fold biomass hydrolysis is higher than
that of other LPMO families [50]. However, the synergy or boosting effect depends on many
factors such as pre-treatment [51], the lignin content of lignocelluloses and acting cellulase
[46]. Still, over 50% enhancement suggests intense demands on inclusion on cellulase
cocktail. However, the mechanism of synergism with the cellulase enzyme complex is poorly
understood. The probable explanation of such a boosting effect could be that the cellulosic

biomass is partially depolymerized by the LPMO, which gives further access to the cellulase
enzymes.

Conclusion

In concluding remark, *Af*LPMO16 is the second report of the AA16 family of LPMO, but for
the first time, we have characterized the AA16 family biochemically and structurally. *In-*
silico sequence analysis, structure analysis, and molecular docking studies suggest some
unique characteristics of the *Af*LPMO16, like cellulose-binding ability, chances of chitin-
binding, and C1 and C4 oxidizing property. Further studies, including the engineering
approach, are required to confirm these characteristics. Nevertheless, the most crucial aspect
of *Af*LPMO16 is the significant boosting effect on commercial cellulase cocktail in
lignocellulosic biomass conversion, and that suggests its importance in the bioethanol
industry.

Materials and Methods

Sequence analysis and Phylogenetic analysis:

*Af*LPMO16 sequence (CAF32158.1) was obtained from NCBI, and the sequence was further
confirmed from the *Aspergillus* genome database (http://www.aspgd.org/). To avoid
interference from the presence or the absence of additional residues or domains, the signal
peptides, and C-terminal extensions were removed before the alignment. Homology sequence
alignment was performed by the BLAST [22]. Clustal Omega [23] was used for multiple
sequence alignment. The sequence alignment was edited with Espript for better visualization.
Pymol [24] and MEGA7 [25] were used to construct a phylogenetic tree after sequence
alignment. To build the phylogenetic tree, the sequences of twenty-seven (27) LPMO genes
(edited to remove N-terminal signal sequence, C-terminal extension or GPI anchor, CBM1
module) were taken from different species belong to AA10 and AA16 family of LPMOs. The
neighbor-joining tree was constructed with 1000 bootstrap replications.

Cloning of AfLPMO16

Aspergillus fumigatus NITDGPKA3 was grown on CMC agar media containing 2% CMC,
0.2% peptone, 2% agar in basal medium (0.2% $NaNO_3$, 0.05%KCl, 0.05%$MgSO_4$,
0.001%$FeSO_4$, 0.1%K_2HPO_4). The fungal biomass was then milled in a pestle and mortar
followed by rapid overtaxing in solution with an appropriate lysis buffer for proper lysis of

the cell. Genomic DNA was isolated from the fungal biomass using the DNA extraction

buffer (400mM Tris-HCl, 150mM NaCl, 0.5M EDTA, 1%SDS) and followed by Phenol,

chloroform and isoamyl alcohol (25:24:1) extraction. The final pellet was washed with 70%

alcohol, air-dried, and dissolved in sterile water. *Af*LPMO16 gene was amplified by

polymerase chain reaction (PCR). The codon-optimized gene for *Pichia pastoris* was inserted

into the pPICZαA vector (Invitrogen Carlsbad, California, USA). The gene was cloned with

the native signal sequence and 6x His-tag at the C-terminal [26]. The cloning was done by

following the same protocol as *Aa*AA16 and PMO9A_MLACI [19, 26]. The vector

(pPICZαA) containing the *Af*LPMO16 gene was linearized by Pme1 (New England BioLabs)

and transformed to *Pichia pastoris* X33 competent cells. The Zeocin resistant transformants

were picked and screened for protein production. The cloned gene was further confirmed by

sequencing and the sequence submitted to GenBank (GenBank accession No. MT462230).

Expression and purification of AfLPMO16

The positive colonies were selected on YPDS (Zeocin: 100µg/ml) plates. The positive

transformants were further screened by the colony PCR and expression studies. Protein

expression was carried out initially in BMGY media containing 1ml/L *Pichia* trace minerals

4 (PTM4) salt (2g/L $CuSO_4·5H_2O$, 3g/L $MnSO4·H_2O$, 0.2g/L $Na_2MoO_4·2H_2O$, 0.02g/L

H_3BO_3, 0.5g/L $CaSO_4·2H_2O$, 0.5g/L $CoCl_2$, 12.5g/L $ZnSO_4·7H_2O$, 22g/L $FeSO_4·7H_2O$, NaI

0.08g/L, H_2SO_4 1mL/L) and 0.1 g/L of biotin. Then after 16 hours, *Pichia* cells were

transferred into BMMY medium (PTM4 salt) with continuous induction by the addition of

1% methanol (optimized) every day (after every 24 hours) for three days. After three days,

the culture media was spun down (8,000rpm for 10mins) at 4^0C. The pellet was discarded,

and the media was collected. The protein was precipitated from the media by ammonium

sulfate precipitation (90% saturation). The pellet was redissolved in Tris buffer (Tris-HCl

50mM pH-7.8, NaCl-400mM, Imidazole-10mM). The recombinant protein was purified by

immobilized ion affinity chromatography (Ni-NTA affinity chromatography)[27], followed

by dialysis with 50mM phosphate buffer, pH 6.0. We followed the expression and

purification procedure, same as *Aa*AA16 [19]. The yield of the purified protein was almost

0.8 mg/ml. The concentration was measured by Bradford assay, and BSA was used for

standard concentration. The protein was separated by SDS-PAGE using 12% acrylamide in

resolving gel(dH$_2$O-3.6 ml, Acrylamide+Bisacrylamide – 4.0 ml, 1.5M Tris-2.6 ml,

10%SDS-0.1 ml, 10% APS-0.1 ml, TEMED- 0.01 ml; for 10 ml), stained with coomassie

blue, and the purified protein band was also confirmed by Western blot analysis by using an

anti-His antibody (Abcam).

Biochemical assays of AfLPMO16

Biochemical characterization of AfLPMO16

2,6 DMP (2,6-dimethoxyphenol) was used as a substrate for *Af*LPMO16 in this study. The

reaction was done in phosphate buffer (100mM pH 6.0) containing 10mM 2,6-

dimethoxyphenol, 5μM hydrogen peroxide, and 50μg of purified *Af*LPMO16 at 30□C. The

amount of product 1-coerulignone was measured by spectrophotometer using the standard

extinction coefficient ($53200M^{-1}cm^{-1}$) and Lambert-Beer law. For kinetic assay different 2,6-

dimethoxyphenol concentrations (1mM, 5mM, 10mM, 20mM, 25mM, 30mM, 40mM,

50mM, 70mM and 100mM) were used. The kinetic parameters were calculated based on the

Line-weaver-Burk plot (LB plot). One unit of enzyme activity is defined as the amount of

enzyme which releases 1μM of 1-coerulignone (product) per minute in standard reaction

condition.

Polysaccharides depolymerization by AfLPMO16

Different cellulosic compounds such as PASC, avicel®PH-101 (SIGMA), and carboxyl

methylcellulose (CMC) was used. We used 1% Avicel®PH-101 (SIGMA) (crystalline

cellulose) and 1% CMC (Carboxyl methylcellulose sodium salt) with different concentrations

of purified *Af*LPMO16 for different incubation time. Reducing sugar was determined by

Dinitro salicylic acid (DNS) assay. For PASC assay, we used 0.25% PASC and incubated

with increasing concentration of *Af*LPMO16 for 6 hours and measured the OD after 6hrs of

incubation and plot the relative absorbance ([OD of *Af*LPMO16 treated PASC]-[OD of

untreated substrate]) with enzyme concentration [28].

Biomass and cellulose hydrolysis by cellulase and AfLPMO16

Cellulose and lignocellulose (alkaline pre-treated raw rice straw) [29] was used to determine

the cellulose hydrolysis enhancing capacity. Rice straw was pre-treated with 5% NaOH (1:10

W/V ratio) at 120□C at 15Psi pressure for 1 hour, and sodium azide (20%) 10μl (per 10ml)

was added at the reaction mixture to prevent any microbial contamination. The reaction was

performed at 50□C, and the amount of reducing sugar was quantified after 5hours, 24hours,

48hours, and 72 hours by Dinitro salicylic acid (DNS) assay. 20μl of cellulase (commercial)

(MP Biomedicals LLC) (5mg/ml) was used along with two different concentrations of

*Af*LPMO16 125μl (100μg) and 250μl (200μg) [concentration 0.8mg/ml]. Reaction sets were

prepared using the only cellulase, only *Af*LPMO16 with different concentrations, combined

*Af*LPMO16 and cellulase and lastly, cellulase with inactivated *Af*LPMO16. *Af*LPMO16 was

heat-inactivated by keeping at 100□C temperature for 30 minutes. Reducing sugar from each

triplicate sets were quantified. In the case of cellulose degradation, 400μl (1%) of avicel

(SIGMA) was incubated with 10μl of cellulase (commercial) (MP Biomedicals LLC)

(5mg/ml). Reducing sugar was quantified after 5 hours of incubation. For these biochemical

assays, we used 100mM phosphate buffer (pH-6.0), and heat-inactivated *Af*LPMO16 was

taken as a negative control.

Molecular modeling and Molecular docking

I-TASSER [30] server was used to model the *Af*LPMO16. The final model was energy

minimized by Gromacs software [31]. The Ramachandran plot [32] and Procheck [33] was

used to evaluate the final model. For Metal Ion-Binding site prediction and docking server or

MIB server (http://bioinfo.cmu.edu.tw/MIB/) were used to identify the copper (Cu) ion

position. A molecular docking study was performed by the Autodock Vina [34] using MGL

tools (Molecular graphics laboratory). The optimized substrate structures were prepared by

Autodock vina and saved in PDBQT format. The grid size parameters used in this docking

were 44, 46, 46, and grid center parameters used in this study were 49, 45, and 55. The

genetic algorithm was also used for docking. Molecular interactions between enzyme and

substrate were analyzed by the MGL tools [35]. The electrostatic potential surface of the

*Af*LPMO16 is calculated by the APBS plugin available in Pymol at pH 6.0.

Acknowledgments

MH is thankful to DBT, and SRD is grateful to DST Inspire for their fellowship. The authors

are also thankful to DST-FIST grant of the Department of Biotechnology, NIT Durgapur.

Funding

This study is financially supported by the DBT, Govt. of India (Grant No. BT/PR13127/

PBD/26/447/2015).

Authors' contribution

MH and SM designed the research work. MH, BSK, and SM wrote the manuscript. MH performed biochemical assays. SRD performed *In-Silico* analysis. MH and KA analyzed the results. All authors read and approved the manuscript.

Conflict of interest

Authors have no competing interests. The manuscript has been spell-checked, grammar checked and plagiarism-checked by "Grammarly."

Ethical approval

No human participants or animal is being used during the study.

References

1. Dias De Oliveira Me, Vaughan Be, Rykiel EJ (2005) Ethanol as Fuel: Energy, Carbon Dioxide Balances, and Ecological Footprint. Bioscience. https://doi.org/10.1641/0006-3568(2005)055[0593:eafecd]2.0.co;2

2. Saricks C, Santini D, Wang M (1999) Effects of Fuel Ethanol Use on Fuel-Cycle Energy and Greenhouse Gas Emissions

3. X. Lang, D. G. MacDonald, G. A. Hil (2002) Recycle Bioreactor for Bioethanol Production from Wheat Starch II. Fermentation and Economics. Energy Sources. https://doi.org/10.1080/009083101300058426

4. Somerville C, Bauer S, Brininstool G, et al (2004) Toward a systems approach to understanding plant cell walls. Science (80-.).

5. Forsberg Z, Vaaje-kolstad G, Westereng B, et al (2011) Cleavage of cellulose by a cbm33 protein. Protein Sci. https://doi.org/10.1002/pro.689

6. Phillips CM, Beeson WT, Cate JH, Marletta MA (2011) Cellobiose dehydrogenase and a copper-dependent polysaccharide monooxygenase potentiate cellulose degradation by Neurospora crassa. ACS Chem Biol. https://doi.org/10.1021/cb200351

7. Quinlan RJ, Sweeney MD, Lo Leggio L, et al (2011) Insights into the oxidative degradation of cellulose by a copper metalloenzyme that exploits biomass components. Proc Natl Acad Sci. https://doi.org/10.1073/pnas.1105776108

8. Johansen KS (2016) Discovery and industrial applications of lytic polysaccharide monooxygenases. Biochem Soc Trans. https://doi.org/10.1042/bst20150204

9.

10. Beeson WT, Vu V V., Span EA, et al. (2015) Cellulose Degradation by Polysaccharide Monooxygenases. Annu Rev Biochem. https://doi.org/10.1146/annurev-biochem-060614-034439

11. Vermaas J V., Crowley MF, Beckham GT, Payne CM (2015) Effects of lytic polysaccharide monooxygenase oxidation on cellulose structure and binding of oxidized cellulose oligomers to cellulases. J Phys Chem B. https://doi.org/10.1021/acs.jpcb.5b00778

12. Forsberg Z, Vaaje-kolstad G, Westereng B, et al (2011) Cleavage of cellulose by a cbm33 protein. Protein Sci. https://doi.org/10.1002/pro.689

13. Vermaas J V., Crowley MF, Beckham GT, Payne CM (2015) Effects of lytic polysaccharide monooxygenase oxidation on cellulose structure and binding of oxidized cellulose oligomers to cellulases. J Phys Chem B. https://doi.org/10.1021/acs.jpcb.5b00778

14. Harris P V., Welner D, McFarland KC, et al (2010) Stimulation of lignocellulosic biomass hydrolysis by proteins of glycoside hydrolase family 61: Structure and function of a large, enigmatic family. Biochemistry. https://doi.org/10.1021/bi100009p

15. Nakagawa YS, Kudo M, Loose JSM, et al (2015) A small lytic polysaccharide monooxygenase from Streptomyces griseus targeting α- And β-chitin. FEBS J. https://doi.org/10.1111/febs.13203

16. Crouch LI, Labourel A, Walton PH, et al (2016) The contribution of non-catalytic carbohydrate-binding modules to the activity of lytic polysaccharide monooxygenases. J Biol Chem. https://doi.org/10.1074/jbc.M115.702365

17. Chabbert B, Habrant A, Herbaut M, et al (2017) Action of lytic polysaccharide monooxygenase on plant tissue is governed by cellular type. Sci Rep. https://doi.org/10.1038/s41598-017-17938-2

18. Liu B, Krishnaswamyreddy S, Muraleedharan MN, et al (2018b) Side-by-side biochemical comparison of two lytic polysaccharide monooxygenases from the white-rot fungus Heterobasidion irregulare on their activity against crystalline cellulose and glucomannan. PLoS One. https://doi.org/10.1371/journal.pone.0203430

19. Filiatrault-Chastel C, Navarro D, Haon M, et al (2019) AA16, a new lytic polysaccharide monooxygenase family identified in fungal secretomes. Biotechnol Biofuels. https://doi.org/10.1186/s13068-019-1394-y

20. Sarkar N, Aikat K (2014) Aspergillus fumigatus NITDGPKA3 provides for increased
cellulase production. Int J Chem Eng 2014:. https://doi.org/10.1155/2014/959845

21. Breslmayr E, Hanžek M, Hanrahan A, et al (2018) A fast and sensitive activity assay for
lytic   polysaccharide   monooxygenase.   Biotechnol   Biofuels.
https://doi.org/10.1186/s13068-018-1063-6

22. Altschul SF, Gish W, Miller W, et al (1990) Basic local alignment search tool. J Mol Biol.
https://doi.org/10.1016/S0022-2836(05)80360-2

23. Sievers F, Higgins DG (2014) Clustal Omega, accurate alignment of very large numbers
of sequences. Methods Mol Biol. https://doi.org/10.1007/978-1-62703-646-7_6

24. DeLano W. . (2002) Pymol: An open-source molecular graphics tool. CCP4 Newsl
Protein Crystallogr

25. Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics
Analysis   Version   7.0   for   Bigger   Datasets.   Mol   Biol   Evol.
https://doi.org/10.1093/molbev/msw054

26. Basotra N, Dhiman SS, Agrawal D, et al (2019) Characterization of a novel Lytic
Polysaccharide Monooxygenase from Malbranchea cinnamomea exhibiting dual
catalytic behavior. Carbohydr Res. https://doi.org/10.1016/j.carres.2019.04.006

27. Bennati-granier C, Garajova S, Champion C, et al (2015) Substrate specificity and
regioselectivity of fungal AA9 lytic polysaccharide monooxygenases secreted by
Podospora anserina To cite this version□: Substrate specificity and regioselectivity of
fungal AA9 lytic polysaccharide monooxygenases secreted by Pod. Biotechnol Biofuels.
https://doi.org/10.1186/s13068-015-0274-3

28. Hansson H, Karkehabadi S, Mikkelsen N, et al (2017) High-resolution structure of a lytic
polysaccharide monooxygenase from Hypocrea jecorina reveals a predicted linker as an
integral part of the catalytic domain. J Biol Chem 292:19099–19109.
https://doi.org/10.1074/jbc.M117.799767

29. Yoswathana (2010) Bioethanol Production from Rice Straw. Energy Res J 1:26–31.
https://doi.org/10.3844/erjsp.2010.26.31

30. Zhang R, Liu Y, Zhang Y, et al (2019) Identification of a thermostable fungal lytic
polysaccharide monooxygenase and evaluation of its effect on lignocellulosic
degradation.   Appl   Microbiol   Biotechnol   103:5739–5750.
https://doi.org/10.1007/s00253-019-09928-3

31. Pronk S, Páll S, Schulz R, et al (2013) GROMACS 4.5: A high-throughput and highly

parallel open source molecular simulation toolkit. Bioinformatics. https://doi.org/10.1093/bioinformatics/btt055

32. Gopalakrishnan K, Sowmiya G, Sheik SS, Sekar K (2007) Ramachandran plot on the web (2.0). Protein Pept Lett

33. Laskowski RA, MacArthur MW, Moss DS, Thornton JM (2002) PROCHECK: a program to check the stereochemical quality of protein structures. J Appl Crystallogr. https://doi.org/10.1107/s0021889892009944

34. Trott O, Olson AJ (2010) Software news and update AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. J Comput Chem. https://doi.org/10.1002/jcc.21334

35. Morris GM, Huey R, Lindstrom W, et al (2009) AutoDock4 and AutoDockTools4: Automated Docking with Selective Receptor Flexibility. J Comp Chem. https://doi.org/10.1002/jcc.21256

36. Agrawal, D., Kaur, B., Kaur Brar, K., Chadha, B.S., 2020. An innovative approach of priming lignocellulosics with lytic polysaccharide monooxygenases prior to saccharification with glycosyl hydrolases can economize the second-generation ethanol process.Bioresour. Technol.308, 123257. https://doi.org/10.1016/j.biortech.2020.123257

37. Jensen MS, Klinkenberg G, Bissaro B, et al (2019) Engineering chitinolytic activity into a cellulose-active lytic polysaccharide monooxygenase provides insights into substrate specificity. J Biol Chem. https://doi.org/10.1074/jbc.RA119.010056

38. Zhou X, Zhu H (2020) Current understanding of substrate specificity and regioselectivity of LPMOs. Bioresour Bioprocess 7:. https://doi.org/10.1186/s40643-020-0300-6

39. Forsberg Z, Mackenzie AK, Sørlie M, et al (2014) Structural and functional characterization of a conserved pair of bacterial cellulose-oxidizing lytic polysaccharide monooxygenases. Proc Natl Acad Sci U S A. https://doi.org/10.1073/pnas.1402771111

40. Hansson H, Karkehabadi S, Mikkelsen N, et al (2017) High-resolution structure of a lytic polysaccharide monooxygenase from Hypocrea jecorina reveals a predicted linker as an integral part of the catalytic domain. J Biol Chem 292:19099–19109. https://doi.org/10.1074/jbc.M117.799767

41. Eibinger, M., Ganner, T., Bubner, P., Rošker, S., Kracher, D., Haltrich, D., Ludwig, R., Plank, H., Nidetzky, B., 2014. Cellulose surface degradation by a lytic polysaccharide monooxygenase and its effect on cellulase hydrolytic efficiency. J. Biol. Chem. https://doi.org/10.1074/jbc.M114.602227

42. Kim, I.J., Youn, H.J., Kim, K.H., 2016. Synergism of an auxiliary activity 9 (AA9) from
Chaetomium globosum with xylanase on the hydrolysis of xylan and lignocellulose.
Process Biochem. 51, 1445–1451. https://doi.org/10.1016/j.procbio.2016.06.017

43. Kim, I.J., Jung, J.Y., Lee, H.J., Park, H.S., Jung, Y.H., Park, K., Kim, K.H., 2015.
Customized optimization of cellulase mixtures for differently pre-treated rice straw.
Bioprocess Biosyst. Eng. 38, 929–937. https://doi.org/10.1007/s00449-014-1338-7

44. Zhang R, Liu Y, Zhang Y, et al (2019) Identification of a thermostable fungal lytic
polysaccharide monooxygenase and evaluation of its effect on lignocellulosic
degradation.   Appl   Microbiol   Biotechnol   103:5739–5750.
https://doi.org/10.1007/s00253-019-09928-3

45. Hemsworth, G.R., Johnston, E.M., Davies, G.J., Walton, P.H., 2015. Lytic Polysaccharide
Monooxygenases   in   Biomass   Conversion.   Trends   Biotechnol.   xx,   1–15.
https://doi.org/10.1016/j.tibtech.2015.09.006

46. Dimarogona, M., Topakas, E., Olsson, L., Christakopoulos, P., 2012. Bioresource
Technology Lignin boosts the cellulase performance of a GH-61 enzyme from
Sporotrichum   thermophile.   Bioresour.   Technol.   110,   480–487.
https://doi.org/10.1016/j.biortech.2012.01.116

47.   Liu B, Kognole AA, Wu M, et al (2018a) Structural and molecular dynamics
studies of a C1-oxidizing lytic polysaccharide monooxygenase from Heterobasidion
irregulare reveal amino acids important for substrate recognition. 285:2225–2242.
https://doi.org/10.1111/febs.14472

48.   Kim, I.J., Youn, H.J., Kim, K.H., 2016. Synergism of an auxiliary activity 9
(AA9) from Chaetomium globosum with xylanase on the hydrolysis of xylan and
lignocellulose.   Process   Biochem.   51,   1445–1451.
https://doi.org/10.1016/j.procbio.2016.06.017

49.   Corrêa TLR, Júnior AT, Wolf LD, et al (2019) An actinobacteria lytic
polysaccharide monooxygenase acts on both cellulose and xylan to boost biomass
saccharification. Biotechnol Biofuels 12:1–14. https://doi.org/10.1186/s13068-019-
1449-0

50.   Hu J, Tian D, Renneckar S, Saddler JN (2018) Enzyme mediated
nanofibrillation of cellulose by the synergistic actions of an endoglucanase, lytic
polysaccharide monooxygenase (LPMO) and xylanase. Sci Rep 8:4–11.
https://doi.org/10.1038/s41598-018-21016-6

51.   Müller G, Várnai A, Johansen KS, et al (2015) Harnessing the potential of
LPMO-containing cellulase cocktails poses new demands on processing conditions.
Biotechnol Biofuels. https://doi.org/10.1186/s13068-015-0376-y

52. Pantoom S, Songsiriritthigul C, Suginta W (2008) The effects of the surface-exposed
residues on the binding and hydrolytic activities of Vibrio carchariae chitinase A. BMC
Biochem 9:1–11. https://doi.org/10.1186/1471-2091-9-2

Figure legends

Figure 1 Expression and purification of *Af*LPMO16 (marked with red arrow). SDS PAGE analysis; lane1, flow-through, lane2,3&4 wash, lane 5 & 6. Purified *Af*LPMO16: Western blot analysis using purified protein presented in lane 5 & 6 of SDS page marked as lane W1 and W2

Figure 2 Enzyme kinetics studies of *Af*LPMO16 with 2,6-DMP (mean values are plotted). **(a)** Chemical reaction to convert 2,6DMP to 1-coerulignone; OD at 469 nm vs. time plot. **(b)** LB plot or 1/v vs 1/[s] plot.

Figure 3 *In silico* analysis of *Af*LPMO16. **(a)** Schematic diagram of *Af*LPMO16; signal peptide: 19 amino acids, catalytic domain: 1-169 amino acids, and a serine-rich domain: 169-271 amino acids. **(b)** Multiple sequence alignment of AA16 LPMOs, C1 oxidizing, and C1/C4 oxidizing AA10 LPMOs: Conserved sequences are highlighted. The red arrow indicates the amino acid responsible for regioselectivity; the Black arrow represents the amino acid responsible for substrate specificity, the black box represents the AA16 conserved motif. **(c)** The electrostatic surface potential of *Af*LPMO16 model structure at pH6.0, blue and red color represents positive and negative potential surface respectively. The area surrounded by the ring represents the catalytic site.

Figure 4 Phylogenetic relationship of *Af*LPMO16 with AA10 LPMOs. A neighbor-joining tree from MEGA showing C1(Bacterial) & C2(Fungal) clades and C2 clade further divided into C2.1 (*Penicillium* & other) & C2.2 (*Aspergillus*) subclades.

Figure 5 Model structure and molecular docking of *Af*LPMO16. **(a)** Predicted three-dimensional models of the *Af*LPMO16 showing functional loops LS(orange), L2(blue),

L3(green), LC(magenta) loops surrounding the copper active site. **(b)** Histidine brace (His20,

His109) of *Af*LPMO16 surrounding the copper metal. **(c)** Amino acids involved in substrate

binding: Gln48, Gln181, Ser178, His109, His20, Asn54, Asp50, Tyr52, Glu58

Figure 6 Polysaccharides degradation activity of *Af*LPMO16. **(a)** CMC depolymerization:

estimation of reducing sugar with the increasing amount of *Af*LPMO16. **(b)** PASC

hydrolysis: relative absorbance at 405nm vs. *Af*LPMO16 quantity plot. Results are the mean

value of the minimum three experiments. The bar represents the standard deviation (SD)

Figure 7 Boosting effect of *Af*LPMO16. **(a)** Hydrolysis of alkali pre-treated rice straw: light-

grey bar indicates only cellulase and deep-grey indicates heat inactive *Af*LPMO16 with

cellulase, dark-grey and black bar indicates cellulase along with two different quantity of

*Af*LPMO16. **(b)** Avicel hydrolysis: reducing sugar estimation. Light-grey bar indicates only

cellulase and deep-grey indicates heat inactive *Af*LPMO16 with cellulase, dark-grey and

black bar indicates cellulase along with two different quantities of *Af*LPMO16. **(c)**

Synergistic effect: light-grey bars indicate biomass hydrolysis by two different concentrations

of *Af*LPMO16; dark-grey bar indicating the only cellulase treated biomass and black bar

indicating combined treated biomass with *Af*LPMO16 & cellulase. Error bars represent the

standard deviation of experiments ran in triplicate. The different number of asterisks (*)

indicate a significant difference between glucose release in the presence of *Af*LPMO16 by

one-way ANOVA followed by Student's t-test ($P \ll 0.05$).

Enzyme Kinetics Parameter	Values
V_{max} in U/mg	0.153
K_m in mM	5.4
K_{cat} in min^{-1}	277.67

Table 1: Enzyme kinetics of *Af*AA16 with 2,6, DMP as a substrate.

Substrates (Biomass)	Cellulases	LPMOs	Fold increase	% increase	References
Wheat straw	Celluclast (Novozymes)	*St*Cel61a (AA9)	-	20%	[46]
Corn stover	Celluclast (Novozymes)	*Ta*AA9		25%	[50]
Raw rice straw	Celluclast (Novozymes)	*Cg*AA9	1.1-1.2	-	[48]
Raw rice straw	**Cellulase (MP Biomedicals)**	***Af*LPMO16**	**2**	**~100%**	**-**

683　　　　　**Table 2: Lignocellulosic biomass hydrolysis enhancement by LPMOs**

Figure: 1

(a)

(b)

Figure: 2

Figure: 3

Figure: 4

(a)

L2 loop

L3 loop

LS loop

LC loop

(b)

Met16

cu504

His20

His109

(c)

ILE
A:179

LYS
A:82

LEU
A:59

SER
A:49

GLU
A:58

TYR
A:81

SER
A:178

GLN
A:48

HIS
A:109

ASN
A:46

ASN
A:54

GLN
A:181

ALA
A:107

HIS
A:20

ASP
A:50

GLY
A:53

TYR
A:52

LYS
A:79

ASN
A:51

Figure: 5

(a)

(b)

Figure: 6

Figure: 7