Consensus transcriptional regulatory networks of coronavirus-infected human cells

Scott A Ochsner[1], Rudolf T Pillich[2] and Neil J McKenna[1]

[1] The Signaling Pathways Project and Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX 77030; [2] Department of Medicine, University of California San Diego, La Jolla, CA 92093.

Address Correspondence To:

Neil J. McKenna

Department of Molecular and Cellular Biology

Baylor College of Medicine

Houston, TX 77030

USA

e: nmckenna@bcm.edu

t: 713-798-8568

Abstract

Establishing consensus around the transcriptional interface between coronavirus (CoV) infection and human cellular signaling pathways can catalyze the development of novel anti-CoV therapeutics. Here, we used publicly archived transcriptomic datasets to compute consensus regulatory signatures, or consensomes, that rank human genes based on their rates of differential expression in MERS-CoV (MERS), SARS-CoV-1 (SARS1) and SARS-CoV-2 (SARS2)-infected cells. Validating the CoV consensomes, we show that high confidence transcriptional targets (HCTs) of CoV infection intersect with HCTs of signaling pathway nodes with known roles in CoV infection. Among a series of novel use cases, we gather evidence for hypotheses that SARS2 infection efficiently represses E2F family target genes encoding key drivers of DNA replication and the cell cycle; that progesterone receptor signaling antagonizes SARS2-induced inflammatory signaling in the airway epithelium; and that SARS2 HCTs are enriched for genes involved in epithelial to mesenchymal transition. The CoV infection consensomes and HCT intersection analyses are freely accessible through the Signaling Pathways Project knowledgebase, and as Cytoscape-style networks in the Network Data Exchange repository.

Introduction

Infection of humans by coronaviruses (CoV) represents a major current global public health concern. Signaling within and between airway epithelial and immune cells in response to infections by CoV and other viruses is coordinated by a complex network of signaling pathway nodes. These include chemokine and cytokine-activated receptors, signaling enzymes and transcription factors, and the genomic targets encoding their downstream effectors[1–3]. Placing the transcriptional events resulting from CoV infection in context with those associated with host signaling paradigms has the potential to catalyze the development of novel therapeutic approaches. The CoV research community has been active in generating and archiving transcriptomic datasets documenting the transcriptional response of human cells to infection by the three major CoV strains, namely, Middle East respiratory syndrome coronavirus (MERS-CoV, or MERS) and severe acute respiratory syndrome coronaviruses 1 (SARS-CoV-1, or SARS1) and 2 (SARS-CoV-2, or SARS2)[4–9]. To date however the field has lacked a resource that fully capitalizes on these datasets by, firstly, using them to identify human genes that are most consistently transcriptionally responsive to CoV infection and secondly, contextualizing these transcriptional responses by integrating them with 'omics data points relevant to host cellular signaling pathways.

We recently described the Signaling Pathways Project (SPP)[10], an integrated 'omics knowledgebase designed to assist bench researchers in leveraging publically archived transcriptomic and ChIP-Seq datasets to generate research hypotheses. A unique aspect of SPP is its collection of consensus regulatory signatures, or consensomes, which rank genes based on the frequency of their significant differential expression

across transcriptomic experiments mapped to a specific signaling pathway node or

node family. By surveying across multiple independent datasets, we have shown that

consensomes recapitulate pathway node-genomic target regulatory relationships to a

high confidence level[10]. Here, as a service to the research community to catalyze the

development of novel CoV therapeutics, we generated consensomes for infection of

human cells by MERS, SARS1 and SARS2 CoVs. Computing the CoV consensomes

against those for a broad range of cellular signaling pathway nodes, we discovered

robust intersections between genes with high rankings in the CoV consensomes and

those of nodes with known roles in the response to CoV infection. Integration of the CoV

consensomes with the existing universes of SPP transcriptomic and ChIP-Seq data

points in a series of use cases illuminates previously uncharacterized interfaces

between CoV infection and human cellular signaling pathways. Moreover, while this

paper was under review and revision, numerous contemporaneous and independent

wet bench-based studies came to light that corroborate *in silico* predictions made using

our analysis pipeline. To reach the broadest possible audience of experimentalists, the

results of our analysis were made available in the SPP website, as well as in the

Network Data Exchange (NDEx) repository. Collectively, these networks constitute a

unique and freely accessible framework within which to generate mechanistic

hypotheses around the transcriptional interface between human signaling pathways and

CoV infection.

Results

Generation of the CoV consensomes

We first set out to generate a set of consensomes[10] ranking human genes based on

statistical measures of the frequency of their significant differential expression in

response to infection by MERS, SARS1 and SARS2 CoVs. To do this we searched the

Gene Expression Omnibus (GEO) and ArrayExpress databases to identify datasets

involving infection of human cells by these strains. Many of these datasets emerged

from a broad-scale systematic multi-omics Pacific Northwest National Library analysis of

the host cellular response to infection across a broad range of pathogens[11]. Since an

important question in the development of CoV therapeutics is the extent to which CoVs

have common transcriptional impacts on human cell signaling that are distinct from

those of other viruses, we also searched for transcriptomic datasets involving infection

by human influenza A virus (IAV). From this initial collection of datasets, we next carried

out a three step quality control check as previously described[10], yielding a total of 3.3

million data points in 156 experiments from 38 independent viral infection transcriptomic

datasets (figshare File F1, section 1). Using these curated datasets, we next used

consensome analysis (see Methods and previous SPP publication[10]) to generate

consensomes for each CoV strain. figshare File F1 contains the full human SARS1

(Section 2), SARS2 (Section 3), MERS (Section 4) and IAV (Section 5) infection

transcriptomic consensomes. To assist researchers in inferring CoV infection-

associated signaling networks, the consensomes are annotated using the previously

described SPP convention[10] to indicate the identity of a gene as encoding a receptor,

protein ligand, enzyme, transcription factor, ion channel or co-node (figshare File F1,

sections 2-5, columns A-C).

Ranking of interferon-stimulated genes (ISGs) in the CoV consensomes

As an initial benchmark for our CoV consensome analysis, we assembled a list of 20

canonical interferon-stimulated genes (ISGs), whose role in the anti-viral response is

best characterized in the context of IAV infection[12]. As shown in Figure 1, many ISGs

were assigned elevated rankings across the four viral consensomes. The mean

percentile of the ISGs was however appreciably higher in the IAV (98.7th percentile) and

SARS1 (98.5th percentile; $p = 6e\text{-}1$, t-test IAV vs SARS1) consensomes than in the

SARS2 (92nd percentile, $p = 5e\text{-}2$, t-test IAV v SARS2) and MERS (82nd percentile; $p =$

7e-5, t-test IAV v MERS) consensomes. This is consistent with previous reports of an

appreciable divergence between the IAV and SARS2 transcriptional responses with

respect to the interferon response[8]. Other genes with known critical roles in the

response to viral infection have high rankings in the CoV consensomes, including

NCOA7[13] (percentiles: SARS1, 98th; SARS2, 97th; MERS, 89th; IAV, 99th), *STAT1*[14]

(percentiles: SARS1, 99th; SARS2, 98th; MERS, 89th; IAV, 99th) and *TAP1*[15] (percentiles:

SARS1, 99th; SARS2, 94th; MERS, 83rd; IAV, 99th). In addition to the appropriate

elevated rankings for these known viral response effectors, the CoV consensomes

assign similarly elevated rankings to transcripts that are largely or completely

uncharacterized in the context of viral infection. Examples of such genes include

PSMB9, encoding a proteasome 20S subunit (percentiles: SARS1, 98th; SARS2, 97th;

MERS, 98th; IAV, 98th); *CSRNP1,* encoding a cysteine and serine rich nuclear protein

(percentiles: SARS1, 99th; SARS2, 94th; MERS, 98th; IAV, 94th); and *CCNL1,* encoding a

member of the cell cycle-regulatory cyclin family (percentiles: SARS1, 99th; SARS2,

94th; MERS, 99th; IAV, 97th). Finally, a CRISPR/Cas9 study posted as a preprint while

this manuscript was under review validated 27 human genes as critical modulators of

the host response to SARS2 infection of human cells[16]. Corroborating our analysis, 16

of these genes have significant ($q < 0.05$) rankings in the SARS2 consensome,

including *ACE2* and *DYRK1A* (both 97th percentile), *CTSL* (96th percentile), *KDM6A*,

ATRX, *PIAS1* (all 94th percentile), *RAD54L2* and *SMAD3* (90th percentile).

To illuminate human signaling pathways orchestrating the transcriptional response to

CoV infection, we next compared transcripts with elevated rankings in the CoV

consensomes with those that have predicted high confidence regulatory relationships

with cellular signaling pathway nodes. We generated four lists of genes corresponding

to the MERS, SARS1, SARS2 and IAV transcriptomic consensome 95th percentiles. We

then retrieved genes in the 95th percentiles of available SPP human transcriptomic (n =

25) consensomes and ChIP-Seq (n = 864) pathway node consensomes[10]. For

convenience we will refer from hereon to genes in the 95th percentile of a viral infection,

node (ChIP-Seq) or node family (transcriptomic) consensome as high confidence

transcriptional targets (HCTs). We then used the R GeneOverlap package[17] to compute

the extent and significance of intersections between CoV HCTs and those of the

pathway nodes or node families. We interpreted the extent and significance of

intersections between HCTs for CoVs and pathway node or node families as evidence

for a biological relationship between loss or gain of function of that node (or node family)

and the transcriptional response to infection by a specific virus.

Results of viral infection and signaling node HCT intersection analyses are shown in

Figure 2 (based on receptor and enzyme family transcriptomic consensomes), Figures 3

and 4 (based on ChIP-Seq consensomes for transcription factors and enzymes,

respectively) and figshare File F2 (based on ChIP-Seq consensomes for selected co-

nodes). figshare File F1, sections 6 (node family transcriptomic HCT intersection

analysis) and 7 (node ChIP-Seq HCT intersection analysis) contain the full underlying

numerical data. We surveyed $q < 0.05$ HCT intersections to identify (i) canonical

inflammatory signaling pathway nodes with characterized roles in the response to CoV

infection, thereby validating the consensome approach in this context; and (ii) evidence

for nodes whose role in the transcriptional biology of CoV infection is previously

uncharacterized, but consistent with their roles in the response to other viral infections.

In the following sections all q-values refer to those obtained using the GeneOverlap

analysis package in R[17].

Receptors Reflecting their well-documented roles in the response to CoV infection[18–21],

we observed appreciable significant intersections between CoV HCTs and those of the

toll-like (TLRs; q-values: SARS1, 3e-85; SARS2, 5e-49; MERS, 2e-33), interferon

(IFNR; q-values: SARS1, 1e-109; SARS2, 6e-53; MERS, 1e-24) and tumor necrosis

factor (TNFR; q-values: SARS1, 1e-48; SARS2, 1e-35; MERS, 5e-32) receptor families

(Fig. 2). HCT intersections between CoV infection and receptor systems with previously

uncharacterized connections to CoV infection, including epidermal growth factor

receptors (EGFR; q-values: SARS1, 4e-21; SARS2, 3e-48; MERS, 1e-35), and Notch

receptor signaling (q-values: SARS1, 6e-24; SARS2, 2e-33; MERS, 2e-29; Fig. 2), are

consistent with their known role in the context of other viral infections[22–26]. The Notch

receptor HCT intersection points to a possible mechanistic basis for the potential of

Notch pathway modulation in the treatment of SARS2[27]. The strong HCT intersection

between CoV infection and xenobiotic receptors (q-values: SARS1, 1e-30; SARS2, 1e-

44; MERS, 5e-32; Fig. 2) reflects work describing a role for pregnane X receptor in

innate immunity[28] and points to a potential role for members of this family in the

response to CoV infection. In addition, the robust intersection between HCTs for SARS2

infection and vitamin D receptor (q = 2e-35) is interesting in light of epidemiological

studies suggesting a link between risk of SARS2 infection and vitamin D deficiency[29,30].

Consistent with a robust signature for the glucocorticoid receptor across all CoVs (GR;

q-values: SARS1, 3e-35; SARS2, 1e-35; MERS, 7e-32), while this paper was under

review, studies were published showing the GR agonist dexamethasone was a

successful therapeutic for SARS2 infection[31]. Finally, and also while this paper was

under review, in vitro analyses confirmed our predictions of the modulation by SARS2

infection of ErbB/EGFR[20,32] and TGFBR[16,32] signaling systems (Fig. 2).

Transcription factors Not unexpectedly – and speaking again to validation of the

consensomes - the strongest and most significant CoV HCT intersections were

observed for HCTs for known transcription factor mediators of the transcriptional

response to CoV infection, including members of the NFκB (q-value ranges: SARS1,

1e-7-1e-9; SARS2, 9e-3-2e-3; MERS, 1e-3-1e-4)[33–35], IRF (q-value ranges: SARS1, 2e-

2-1e-31; SARS2, 2e-4-1e-17; MERS, 9e-4-7e-5)[36] and STAT (q-value ranges: SARS1,

1e-7-1e-55; SARS2, 2e-3-3e-29; MERS, 5e-2-3e-5)[37–39] transcription factor families

(Fig. 3). Consistent with the similarity between SARS1 and IAV consensomes with

respect to elevated rankings of ISGs (Fig. 2a & d), the IRF1 HCT intersection was

strongest with the SARS1 ($q = $2e-34) and IAV ($q = $3e-49) HCTs. Corroborating our

finding of a strong intersection between STAT2 and SARS2 infection HCTs ($q = $3e-29),

a study that appeared while this manuscript was under review showed that STAT2 plays

a prominent role in the response to SARS2 infection of Syrian hamsters[40]. HCT

intersections for nodes originally characterized as having a general role in RNA Pol II

transcription, including TBP (q-values: SARS1, 2e-10; SARS2, 6e-23; MERS, 3e-16),

GTF2B/TFIIB (q-values: SARS1, 7e-10; SARS2, 3e-23; MERS, 9e-14) and GTF2F1 (q-

values: SARS1, 2e-4; SARS2, 2e-13; MERS, 5e-5) were strong across all CoVs, and

particularly noteworthy in the case of SARS2. In the case of GTF2B, these data are

consistent with previous evidence identifying it as a specific target for orthomyxovirus[41],

and the herpes simplex[42] and hepatitis B[43] viruses. Moreover, a proteomic analysis that

appeared in BioRXiv while this paper was under review identified a high confidence

interaction between GTF2F2 and the SARS2 NSP9 replicase[32].

In general, intersections between viral infection and ChIP-Seq enrichments for

transcription factors and other nodes were more specific for individual CoV infection

HCTs (compare Fig. 2 with Figs. 3 & 4 and figshare File F1, sections 6 and 7). This is

likely due to the fact that ChIP-Seq consensomes are based on direct promoter binding

by a specific node antigen, whereas transcriptomic consensomes encompass both

direct and indirect targets of specific receptor and enzyme node families.

Enzymes Compared to the roles of receptors and transcription factors in the response

to viral infection, the roles of signaling enzymes are less well illuminated – indeed, in the

context of CoV infection, they are entirely unstudied. Through their regulation of cell

cycle transitions, cyclin-dependent kinases (CDKs) play important roles in the

orchestration of DNA replication and cell division, processes that are critical in the viral

life cycle. CDK6, which has been suggested to be a critical G1 phase kinase[44,45], has

been shown to be targeted by a number of viral infections, including Kaposi's sarcoma-

associated herpesvirus[46] and HIV-1[47]. Consistent with this common role across distinct

viral infections, we observed robust intersection between the CDK family HCTs (q-

values: SARS1, 8e-23; SARS2, 2e-31; MERS, 1e-30; Fig. 2) and the CDK6 HCTs (q-

values: SARS1, 1e-7; SARS2, 8e-8; MERS, 3e-4; Fig. 4) and those of all viral HCTs. As

with the TLRs, IFNRs and TNFRs, which are known to signal through CDK6[48–50],

intersection with the CDK6 HCTs was particularly strong in the case of the SARS2

HCTs (Fig. 4). Again, the subsequent proteomic analysis we alluded to earlier[32]

independently corroborated our prediction of a role for CDK6 in the response to SARS2

infection.

CCNT2 is another member of the cyclin family that, along with CDK9, is a component of

the viral-targeted p-TEFB complex[51]. Reflecting a potential general role in viral infection,

appreciable intersections were observed between the CCNT2 HCTs and all viral HCTs

(q-values: SARS1, 4e-4; SARS2, 6e-3; MERS, 7e-5; Fig. 4). Finally in the context of

enzymes, the DNA topoisomerases have been shown to be required for efficient

replication of simian virus 40[52] and Ebola[53] viruses. The prominent intersections

between DNA topoisomerase-dependent HCTs and the CoV HCTs (q-values: SARS1,

3e-15; SARS2, 6e-21; MERS, 1e-26; Fig. 4) suggest that it may play a similar role in

facilitating the replication of these CoVs.

Hypothesis generation use cases

We next wished to show how the CoV consensomes and HCT intersection networks,

supported by existing canonical literature knowledge, enable the user to generate novel

hypotheses around the transcriptional interface between CoV infection and human

cellular signaling pathways. Given the current interest in SARS2, we have focused our

use cases on that virus. In addition to these use cases, figshare File F2 contains a

number of additional use cases omitted from the main text due to space constraints.

Unless otherwise stated, all *q*-values below were obtained using the GeneOverlap

analysis package in R[17]. We stress that all use cases represent preliminary *in silico*

evidence only, and require rigorous pressure-testing at the bench for full validation.

Hypothesis generation use case 1: transcriptional regulation of the SARS2

receptor gene, *ACE2*

ACE2, encoding membrane-bound angiotensin converting enzyme 2, has gained

prominence as the target for cellular entry by SARS1[54] and SARS2[55]. An important

component in the development of ACE2-centric therapeutic responses is an

understanding of its transcriptional responsiveness to CoV infection. Interestingly,

based on our CoV consensome analysis, *ACE2* is more consistently transcriptionally

responsive to infection by SARS CoVs (SARS1: 98[th] percentile, consensome *q* value

(CQV)[10] = 1e-25; SARS2: 97[th] percentile, CQV = 4e-7) than by IAV (78[th] percentile,

CQV = 3e-8) or MERS (49[th] percentile, CQV = 2e-16; figshare File F1, sections 2-5).

The data points underlying the CoV consensomes indicate evidence for tissue-specific

differences in the nature of the regulatory relationship between *ACE2* and viral infection.

In response to SARS1 infection, for example, *ACE2* is induced in pulmonary cells but

repressed in kidney cells (Fig. 5). On the other hand, in response to SARS2 infection,

ACE2 is repressed in pulmonary cells - a finding corroborated by other studies [56,57] - but

inducible in gastrointestinal cells (Fig. 5). These data may relate to the selective

transcriptional response of *ACE2* to signaling by IFNRs (92[nd] percentile; figshare File

F1, section 8) rather than TLRs (48[th] percentile; figshare File F1, section 9) or TNFRs

(13[th] percentile, figshare File F1, section 10). While this manuscript was under review,

another study appeared confirming repression of induction of *ACE2* by interferon

stimulation and by IAV infection[58]. Our data reflect a complex transcriptional relationship

between *ACE2* and viral infection that may be illuminated in part by future single cell

RNA-Seq analysis in the context of clinical or animal models of SARS2 infection.

Hypothesis generation use case 2: evidence for antagonistic cross-talk between

progesterone receptor and interferon receptor signaling in the airway epithelium

A lack of clinical data has so far prevented a definitive evaluation of the connection

between pregnancy and susceptibility to SARS2 infection in CoVID-19. That said,

SARS2 infection is associated with an increased incidence of pre-term deliveries[59], and

pregnancy has been previously associated with the incidence of viral infectious

diseases, particularly respiratory infections[60,61]. We were therefore interested to observe

consistent intersections between the progesterone receptor (PGR) HCTs and CoV

infection HCTs (q-values: SARS1, 3e-35; SARS2, 5e-41; MERS 5e-28), with the

intersection being particularly evident in the case of the SARS2 HCTs (Fig. 2; figshare

File F1, section 6). To investigate the specific nature of the crosstalk implied by this

transcriptional intersection in the context of the airway epithelium, we first identified a

set of 12 genes that were HCTs for both SARS2 infection and PGR. Interestingly, many

of these genes encode members of the classic interferon-stimulated gene (ISG)

response pathway[12]. We then retrieved two SPP experiments involving treatment of

A549 airway epithelial cells with the PGR full antagonist RU486 (RU), alone or in

combination with the GR agonist dexamethasone (DEX). As shown in Figure 6, there

was unanimous correlation in the direction of regulation of all 12 genes in response to

CoV infection and PGR loss of function. These data are consistent with the reported

pro-inflammatory effects of RU486 in a mouse model of allergic pulmonary

inflammation[62]. Interestingly, SARS2-infected pregnant women are often

asymptomatic[63,64]. Based on our data, it can be reasonably hypothesized that

suppression of the interferon response to SARS2 infection by elevated circulating

progesterone during pregnancy may contribute to the asymptomatic clinical course.

Indeed, crosstalk between progesterone and inflammatory signaling is well

characterized in the reproductive system, most notably in the establishment of uterine

receptivity[65] as well as in ovulation[66]. Consistent with our hypothesis, while this paper

was under review, a clinical trial was launched to evaluate the potential of progesterone

for treatment of COVID-19 in hospitalized men[67]. Interestingly, and also while this paper

was under review, a paper appeared showing that progesterone inhibited SARS2

replication in African green monkey kidney Vero 6 cells[68]. These results indicate an

additional mechanism, distinct from its potential crosstalk with the interferon response,

by which progesterone signaling may impact SARS2 infection.

Hypothesis generation use case 3: association of an epithelial to mesenchymal

transition transcriptional signature with SARS2 infection

Epithelial to mesenchymal transition (EMT) is the process by which epithelial cells lose

their polarity and adhesive properties and acquire the migratory and invasive

characteristics of mesenchymal stem cells[69]. EMT is known to contribute to pulmonary

fibrosis[70], acute interstitial pneumonia[71] and acute respiratory distress syndrome

(ARDs)[72], all of which have been reported in connection with SARS2 infection in

COVID-19[73–75]. We were interested to note therefore that significant HCT intersections

for three well characterized EMT-promoting transcription factors were specific to SARS2

infection (q-values: SNAI2/Slug[76], 2e-2; EPAS1/HIF2α[77], 9e-9; LEF1[78], 1e-3; Fig. 3, bold

symbols; figshare File F1, section 7). Consistent with this, intersections between HCTs

for TGFBRs, SMAD2 and SMAD3, known regulators of EMT transcriptional programs[79]

– were stronger with HCTs for SARS2 (q-values: TGFBRs, 2e-31; SMAD2, 2e-7;

SMAD3, 5e-17) than with those of SARS1 (q-values: TGFBRs, 6e-29; SMAD2, 2e-2;

SMAD3, 3e-9) and MERS (q-values: TGFBRs, 1e-16; SMAD2, 3e-3; SMAD3, 2e-12) –

see also Figs. 2 and 3 and figshare File F1, sections 6 and 7). Moreover, a recent

CRISPR/Cas9 screen identified a requirement for both TGFBR signaling and *SMAD3* in

mediating SARS2 infection[16].

To investigate the connection between SARS2 infection and EMT implied by these HCT

intersections, we then computed intersections between the individual viral HCTs and a

list of 335 genes manually curated from the research literature as EMT markers[80]

(figshare File F1, section 11). In agreement with the HCT intersection analysis, we

observed significant enrichment of members of this gene set within the SARS2 HCTs (q

$= 4e\text{-}14$), but not the SARS1 or MERS (both $q = 2e\text{-}1$) HCTs (Fig. 7a). Consistent with

previous reports of a potential link between EMT and IAV infection[81], we observed

significant intersection between the EMT signature and the IAV HCTs ($q = 1e\text{-}04$).

One possible explanation for the selective intersection between the literature EMT

signature and the SARS2 HCTs relative to SARS1 and MERS was the fact that the

SARS2 consensome was exclusively comprised of epithelial cell lines, whereas the

SARS1 and MERS consensomes included non-epithelial cell biosamples (figshare File

F1, section 1). To exclude this possibility therefore, we next calculated airway epithelial

cell-specific consensomes for SARS1, SARS2 and MERS and computed intersections

between their HCTs and the EMT signature. We found that significant intersection of the

EMT signature with the CoV HCTs remained specific to SARS2 (q-values: SARS1, 2e-

1; SARS2, 1e-8; MERS, 2e-1) in the lung epithelium-specific CoV consensomes.

We next retrieved the canonical EMT genes in the SARS2 HCTs and compared their

percentile rankings with the other CoV consensomes. Although some EMT genes, such

as *CXCL2* and *IRF9*, had elevated rankings across all four viral consensomes, the

collective EMT gene signature had a significantly higher mean percentile value in the

SARS2 consensome than in each of the other viral consensomes (Fig. 7b; SARS2

mean percentile = 97.5; SARS1 mean percentile = 86, *p* = 1e-5, t-test; MERS mean

percentile = 63, *p* = 1e-9, t-test; IAV mean percentile = 76, p = 2e-7, t-test). A column

named "EMT" in figshare File F1, sections 2 (SARS1), 3 (SARS2), 4 (MERS) and 5

(IAV) identifies the ranking of the EMT genes in each of the viral consensomes.

Given that EMT has been linked to ARDs[72], we speculated that the evidence connecting

EMT and SARS2 acquired through our analysis might be reflected in the relatively

strong intersection between ARDs markers in SARS2 HCTs compared to other viral

HCTs. To test this hypothesis we carried out a PubMed search to identify a set of 88

expression biomarkers of ARDs or its associated pathology, acute lung injury (ALI). A

column named "ALI/ARDs" in figshare File F1, sections 2 (SARS1), 3 (SARS2) 4

(MERS) and 5 (IAV) identifies the expression biomarker genes using the PubMed

identifiers for the original studies in which they were identified. Consistent with our

hypothesis, we observed appreciable intersections between this gene set and the HCTs

of all four viruses (SARS1 odds ratio (OR) = 7, q = 5e-9; SARS2 OR = 10.4, q = 1e-9;

MERS, OR = 4.2, q = 2e-5; IAV OR = 6.8; q = 9e-8) with a particularly strong

intersection evident in the SARS2 HCTs.

Although EMT has been associated with infection by transmissible gastroenteritis virus[82]

and IAV[81], this is to our knowledge the first evidence connecting CoV infection, and

specifically SARS2 infection, to an EMT signature. Interestingly, lipotoxin A4 has been

shown to attenuate lipopolysaccharide-induced lung injury by reducing EMT[83].

Moreover, several members of the group of SARS2-induced EMT genes have been

associated with signature pulmonary comorbidities of CoV infection, including *ADAR*[84],

CLDN1[85] and *SOD2*[86]. Of note in the context of these data is the fact that signaling

through two SARS2 cellular receptors, ACE2/AT2 and CD147/basigin, has been linked

to EMT in the context of organ fibrosis[87–89]. Finally, while this manuscript was under

review, a preprint was posted that described EMT-like transcriptional and metabolic

changes in response to SARS2 infection[90]. Collectively, our data indicate that EMT

warrants further investigation as a SARS2-specific pathological mechanism.

Hypothesis generation use case 4: SARS2 repression of E2F family HCTs

encoding cell cycle regulators

Aside from EPAS1 and SNAI2, the only other transcription factors with significant HCT

intersections that were specific to the SARS2 HCTs were the E2F/FOX class members

E2F1 (q-values: SARS1, 5e-1; SARS2, 1e-2; MERS, 4e-1), E2F3 (q-values: SARS1,

6e-1; SARS2, 5e-2; MERS, 7e-1), E2F4 (q-values: SARS1, 1; SARS2, 9e-3; MERS, 1)

and TFDP1/Dp-1 (q-values: SARS1, 1; SARS2, 3e-4; MERS, 1; Fig. 3, bold symbols;

figshare File F1, section 7). These factors play well-documented interdependent roles in

the promotion (E2F1, E2F3, TFDP1) and repression (E2F4) of cell cycle genes[91,92].

Moreover, E2F family members are targets of signaling through EGFRs[93] and CDK6[94],

both of whose HCTs had SARS2 HCT intersections that were stronger those of the

other CoVs (EGFRs: q-values: SARS1, 4e-21; SARS2, 3e-48; MERS, 1e-35; CDK6: q-

values: SARS1, 1e-7; SARS2, 8e-8; MERS, 2e-4); Figs. 2 & 4). Based on these data,

we speculated that SARS2 infection might impact the expression of E2F-regulated cell

cycle genes more efficiently than other CoVs. To investigate this we retrieved a set of

SARS2 HCTs that were also HCTs for at least three of E2F1, E2F3, E2F4 and TFDP1

(figshare File F1, section 3, columns P-T). Consistent with the role of E2F/Dp-1 nodes in

the regulation of the cell cycle, many of these genes – notably *CDK1*, *PCNA*, *CDC6*,

CENPF and *NUSAP1* – are critical positive regulators of DNA replication and cell cycle

progression[95–99] and are known to be transcriptionally induced by E2Fs[100–103]. Strikingly,

with the exception of *E2F3*, all were consistently repressed in response to SARS2

infection (Fig. 8a). To gain insight into the relative efficiency with which the four viruses

impacted expression of the E2F/Dp-1 HCT signature, we compared their mean

percentile values across the viral consensomes. Consistent with efficient repression of

the E2F/Dp-1 HCTs by SARS2 infection relative to other viruses, their mean percentile

ranking was appreciably higher in the SARS2 consensome (97[th] percentile) than in the

SARS1 (76[th] percentile; *p* = 6e-12, t-test), MERS (71.2 percentile; p = 9e-6, t-test) and

IAV (71.2 percentile; p = 2e-5, t-test) consensomes (Fig. 8b). Although manipulation of

the host cell cycle and evasion of detection through deregulation of cell cycle

checkpoints has been described for other viruses[104–106], this represents the first

evidence for the profound impact of SARS2 infection on host cell cycle regulatory

genes, potentially through disruption of E2F mediated signaling pathways. The SARS2

infection-mediated induction of *E2F3* (Fig. 8a) may represent a compensatory response

to transcriptional repression of other E2F family members, as has been previously

observed for this family in other contexts[107,108]. Consistent with our prediction in this use

case, while this paper was in revision, a study appeared showing that infection by

SARS2 results in cell cycle arrest[109]. Our results represent evidence that efficient

modulation by SARS2 of E2F signaling, resulting in repression of cell cycle regulatory

genes, may contribute to its unique pathological impact.

Visualization of the CoV transcriptional regulatory networks in the Signaling

Pathways Project knowledgebase and Network Data Exchange repository

To enable researchers to routinely generate mechanistic hypotheses around the

interface between CoV infection human cell signaling, we next made the consensomes

and accompanying HCT intersection analyses freely available to the research

community in the SPP knowledgebase and the Network Data Exchange (NDEx)

repository. Table 1 contains digital object identifier (DOI)-driven links to the consensome

networks in SPP and NDEx, and to the HCT intersection networks in NDEx.

We have previously described the SPP biocuration pipeline, database and web

application interface[10]. Figure 9 shows the strategy for consensome data mining on the

SPP website. The individual CoV consensomes can be accessed by configuring the

SPP Ominer query form as shown, in this example for the SARS2 consensome (Fig.

9a). Figure 9b shows the layout of the consensomes, showing gene symbol, name,

percentile ranking and other essential information. Genes in the 90^{th} percentile of each

consensome are accessible via the user interface, with the full consensomes available

for download in a tab delimited text file. Target gene symbols in the consensome link to

the SPP Regulation Report, filtered to show only experimental data points that

contributed to that specific consensome (Fig. 9c). This view gives insights into the

influence of tissue and cell type context on the regulatory relationship. These filtered

reports can be readily converted to default Reports that show evidence for regulation of

a specific gene by other signaling pathway nodes. As previously described, pop-up

windows in the Report provide experimental details, in addition to links to the parent

dataset (Fig. 9d), curated accordingly to our previously described protocol[10]. Per FAIR

data best practice, CoV infection datasets – like all SPP datasets – are associated with

detailed descriptions, assigned a DOI, and linked to the associated article to place the

dataset in its original experimental context (Fig. 9d). The full list of datasets is available

for browsing in the SPP Dataset listing (https://www.signalingpathways.org/index.jsf).

The NDEx repository facilitates collaborative publication of biological networks, as well

as visualization of these networks in web or desktop versions of the popular and

intuitive Cytoscape platform[110–112]. Figure 10 shows examples of consensome and HCT

intersection network visualizations within the NDEx user interface. For ease of viewing,

the initial rendering of the full SARS2 (Fig. 10a) and other consensome networks shows

a sample (Fig. 10a, red arrow 1) containing only the top 5% of regulated transcripts; the

full data can be explored using the "Neighborhood Query" feature available at the

bottom of the page (red arrow 2). The integration in NDEx of the popular Cytoscape

desktop application enables any network to be seamlessly be imported in Cytoscape for

additional analysis (red arrow 3). Zooming in on a subset of the SARS2 consensome

(orange box) affords an appreciation of the diversity of molecular classes that are

transcriptionally regulated in response to SARS2 infection (Fig. 10b). Transcript size is

proportional to rank percentile, and edge weight is proportional to the transcript

geometric mean fold change (GMFC) value. Selecting a transcript allows the associated

consensome data, such as rank, GMFC and family, to be examined in detail using the

information panel (Fig. 10b, right panel). Highlighted to exemplify this feature is IL6, an

inflammatory ligand that has been previously linked to SARS2 pathology[8,113].

Consensome GMFCs are signless with respect to direction of regulation[10]. Researchers

can therefore follow the SPP link in the side panel (Fig. 10b, red arrow 4) to view the

individual underlying experimental data points on the SPP site (Fig. 9c shows the

example for *IFI27*). A network of the top 20 ranked transcripts in the SARS2

consensome (Fig. 10c) includes genes with known (*OAS1*, *MX1*[114]) and previously

uncharacterized (*PDZKIP1*, *SAT1*, *TM4SF4*) transcriptional responses to SARS2

infection. Finally, to afford insight into pathway nodes whose gain or loss of function

contributes to SARS2 infection-induced signaling, Figure 10d shows the top 5% ranked

nodes in the SARS2 node HCT ChIP-Seq intersection network (see figshare File F1,

section 7; see also Figs. 2 & 3 and accompanying discussion above). In this, as with all

HCT intersection networks, node size is proportional to the q-value, such that the larger

the circle, the lower the q-value, and the higher the confidence that a particular node or

node family is involved in the transcriptional response to viral infection.

The visual organization of the NDEx interface offers insights into the impact of CoV

infection on human cell signaling that are not readily appreciated in the current SPP

interface. For example, it is readily apparent from the NDEx SARS2 consensome

network (Fig. 10c; Table 1) that the single largest class of SARS2 HCTs encodes

immunomodulatory ligands (OR = 4.6, p = 3.8 e-24, hypergeometric test), many of

which are members of the cytokine and chemokine superfamilies. In contrast, although

still overabundant (OR = 1.58, p = 6.8e-4, hypergeometric test), inflammatory ligands

comprise a considerably smaller proportion of the SARS1 HCTs (Table 1). These data

represent evidence that SARS2 infection is relatively efficient in modulating a

transcriptional inflammatory response in host cells. Consistent with this hypothesis, and

while this manuscript was under review, a study appeared showing induction of

interferon-stimulated genes in COVID-19 patients was more robust than in response to

SARS1 infection[115].

Discussion

An effective research community response to the impact of CoV infection on human health demands systematic exploration of the transcriptional interface between CoV infection and human cell signaling systems. It also demands routine access to computational analysis of existing datasets that is unhindered either by paywalls or by lack of the informatics training required to manipulate archived datasets in their unprocessed state. Moreover, the substantial logistical obstacles to high containment laboratory certification emphasize the need for fullest possible access to, and re-usability of, existing CoV infection datasets to focus and refine hypotheses prior to carrying out *in vivo* CoV infection experiments. Meta-analysis of existing datasets represents a powerful approach to establishing consensus transcriptional signatures – consensomes – which identify those human genes whose expression is most consistently and reproducibly impacted by CoV infection. Moreover, integrating these consensus transcriptional signatures with existing consensomes for cellular signaling pathway nodes can illuminate transcriptional convergence between CoV infection and human cell signaling nodes.

To this end, we generated a set of CoV infection consensomes that rank human genes by the reproducibility of their differential expression ($p < 0.05$) in response to infection of human cells by CoVs. Using HCT intersection analysis, we then computed the CoV consensomes against high confidence transcriptional signatures for a broad range of cellular signaling pathway nodes, affording investigators with a broad range of signaling interests an entrez into the study of CoV infection of human cells. Although other enrichment based pathway analysis tools exist[116], HCT intersection analysis differs from

these by computing against only genes that have the closest predicted regulatory

relationships with upstream pathway nodes (i.e. HCTs). The use cases described here

represent illustrative examples of the types of analyses that users are empowered to

carry out in the CoV infection knowledgebase.

Previous network analyses across independent viral infection transcriptomic datasets,

although valuable, have been limited to stand-alone studies[117,118]. Here, to enable

access to the CoV consensomes and their >3,000,000 underlying data points by the

broadest possible audience, we have integrated them into the SPP knowledgebase and

NDEx repository to create a unique, federated environment for generating hypotheses

around the impact of CoV infection on human cell signaling. NDEx provides users with

the familiar look and feel of Cytoscape to reduce barriers of accessibility and provides

for intuitive click-and-drag data mining strategies. Incorporation of the CoV data points

into SPP places them in the context of millions more existing SPP data points

documenting transcriptional regulatory relationships between human pathway nodes

and their genomic targets. In doing so, we provide users with evidence for signaling

nodes whose gain or loss of function in response to CoV infection gives rise to these

transcriptional patterns. The transcriptional impact of viral infection is known to be an

amalgam of host antiviral responses and co-option by viruses of the host signaling

machinery in furtherance of its life cycle. It is hoped that dissection of these two distinct

modalities in the context of CoV infection will be facilitated by the availability of the CoV

consensomes in the SPP and NDEx knowledgebases.

The CoV consensomes have a number of limitations. Primarily, since they are

predicated specifically on transcriptional regulatory technologies, they will assign low

rankings to transcripts that may not be transcriptionally responsive to CoV infection, but

whose encoded proteins nevertheless play a role in the cellular response. For example,

MASP2, which encodes an important node in the response to CoV infection[119], has

either a very low consensome ranking (SARS1, MERS and IAV), or is absent entirely

(SARS2), indicating that it is transcriptionally unresponsive to viral infection and likely

activated at the protein level in response to upstream signals. This and similar instances

therefore represent "false negatives" in the context of the impact of CoV infection on

human cells. Another limitation of the transcriptional focus of the datasets is the

absence of information on specific protein interactions and post-translational

modifications, either viral-human or human-human, that give rise to the observed

transcriptional responses. Although these can be inferred to some extent, the availability

of existing[32,68,109] and future proteomic and kinomic datasets will facilitate modeling of

the specific signal transduction events giving rise to the downstream transcriptional

responses. Finally, although detailed metadata are readily available on the underlying

data points, the consensomes do not directly reflect the impact of variables such as

tissue context or duration of infection on differential gene expression. As additional

suitable archived datasets become available, we will be better positioned to generate

more specific consensomes of this nature.

The human CoV and IAV consensomes and their underlying datasets are intended as

"living" resources in SPP and NDEx that will be updated and versioned with appropriate

datasets as resources permit. This will be particularly important in the case of SARS2,

given the expanded budget that worldwide funding agencies are likely to allocate to

research into the impact of this virus on human health. Incorporation of future datasets

will allow for clarification of observations that are intriguing, but whose significance is

currently unclear, such as the intersection between the CoV HCTs and those of the

telomerase catalytic subunit (figshare File F2), as well as the enrichment of EMT genes

among those with elevated rankings in the SARS2 consensome (Fig. 7). Although they

are currently available on the SPP website, distribution of the CoV consensome data

points via the SPP RESTful API[10] will be essential for the research community to fully

capitalize on this work. For example, several co-morbidities of SARS2 infection,

including renal and gastrointestinal disorders, are within the mission of the National

Institute of Diabetes, Digestive and Kidney Diseases. In an ongoing collaboration with

the NIDDK Information Network (DKNET)[120], the SPP API will make the CoV

consensome data points available in a hypothesis generation environment that will

enable users to model intersections of CoV infection-modulated host signaling with their

own research areas of interest. We welcome feedback and suggestions from the

research community for the future development of the CoV infection consensomes and

HCT node intersection networks.

Methods

Consistent with emerging NIH mandates on rigor and reproducibility, we have used the

Research Resource Identifier (RRID) standard[121] to identify key research resources of

relevance to our study.

Dataset biocuration

Datasets from Gene Expression Omnibus (SCR_005012) and Array Express

(SCR_002964) were biocurated as previously described, with the incorporation of an

additional classification of peptide ligands[122] to supplement the existing mappings

derived from the International Union of Pharmacology Guide To Pharmacology

(SCR_013077).

Dataset processing and consensome analysis

Array data processing To process microarray expression data, we utilized the log2

summarized and normalized array feature expression intensities provided by the

investigator and housed in GEO. These data are available in the corresponding "Series

Matrix Files(s)". The full set of summarized and normalized sample expression values

were extracted and processed in the statistical program R. To calculate differential gene

expression for investigator-defined experimental contrasts, we used the linear modeling

functions from the Bioconductor limma analysis package [123]. Initially, a linear model was

fitted to a group-means parameterization design matrix defining each experimental

variable. Subsequently, we fitted a contrast matrix that recapitulated the sample

contrasts of interest, in this case viral infection vs mock infection, producing fold-change

and significance values for each array feature present on the array. The current

BioConductor array annotation library was used for annotation of array identifiers. P

values obtained from limma analysis were not corrected for multiple comparisons. _RNA-_

Seq data processing. To process RNA-Seq expression data, we utilized the aligned, un-

normalized, gene summarized read count data provided by the investigator and housed

in GEO. These data are available in the corresponding "Supplementary file" section of

the GEO record. The full set of raw aligned gene read count values were extracted and

processed in the statistical program R using the limma[123] and edgeR analysis[124]

packages. Read count values were initially filtered to remove genes with low read

counts. Gene read count values were passed to downstream analysis if all replicate

samples from at least one experimental condition had cpm > 1. Sequence library

normalization factors were calculated to apply scale normalization to the raw aligned

read counts using the TMM normalization method implemented in the edgeR package

followed by the voom function[125] to convert the gene read count values to log2-cpm.

The log2-cpm values were initially fit to a group-means parameterization design matrix

defining each experimental variable. This was subsequently fit to a contrast design

matrix that recapitulates the sample contrasts of interest, in this case viral infection vs

mock infection, producing fold-change and significance values for each aligned

sequenced gene. If necessary, the current BioConductor human organism annotation

library was used for annotation of investigator-provided gene identifiers. P values

obtained from limma analysis were not corrected for multiple comparisons.

Differential expression values were committed to the consensome analysis pipeline as

previously described[10]. Briefly, the consensome algorithm surveys each experiment

across all datasets and ranks genes according to the frequency with which they are

significantly differentially expressed. For each transcript, we counted the number of

experiments where the significance for differential expression was ≤0.05, and then

generated the binomial probability, referred to as the consensome p-value (CPV), of

observing that many or more nominally significant experiments out of the number of

experiments in which the transcript was assayed, given a true probability of 0.05. Genes

were ranked firstly by CPV, then by geometric mean fold change (GMFC). A more

detailed description of the transcriptomic consensome algorithm is available in a

previous publication[10]. The consensomes and underlying datasets were loaded into an

Oracle 13c database and made available on the SPP user interface as previously

described[10].

Statistical analysis

High confidence transcript intersection analysis was performed using the Bioconductor

GeneOverlap analysis package[17] (SCR_018419) implemented in R. Briefly, given a

whole set I of IDs and two sets A ∈ I and B ∈ I, and S = A ∩ B, GeneOverlap calculates

the significance of obtaining S. The problem is formulated as a hypergeometric

distribution or contingency table, which is solved by Fisher's exact test. *p*-values were

adjusted for multiple testing by using the method of Benjamini & Hochberg to control the

false discovery rate as implemented with the p.adjust function in R, to generate q-

values. The universe for the intersection was set at a conservative estimate of the total

number of transcribed (protein and non protein-coding) genes in the human genome

(25,000)[126]. R code for analyzing the intersection between an investigator gene set and

CoV consensome HCTs has been deposited in the SPP Github account. A two tailed

two sample t-test assuming equal variance was used to compare the mean percentile

ranking of the EMT (12 degrees of freedom) and E2F (14 degrees of freedom)

signatures in the MERS, SARS1, SARS2 and IAV consensomes using the PRISM

software package (SCR_005375).

Consensome generation

The procedure for generating transcriptomic consensomes has been previously

described[10]. To generate the ChIP-Seq consensomes, we first retrieved processed

gene lists from ChIP-Atlas[127] (SCR_015511), in which human genes are ranked based

upon their average MACS2 occupancy across all publically archived datasets in which a

given pathway node is the IP antigen. Of the three stringency levels available (10, 5 and

1 kb from the transcription start site), we selected the most stringent (1 kb). According to

SPP convention[10], we then mapped the IP antigen to its pathway node category, class

and family, and the experimental cell line to its appropriate biosample physiological

system and organ. We then organized the ranked lists into percentiles to generate the

node ChIP-Seq consensomes. The 95[th] percentiles of all consensomes (HCTs, high

confidence transcriptional targets) was used as the input for the HCT intersection

analysis.

SPP web application

The SPP knowledgebase (SCR_018412) is a gene-centric Java Enterprise Edition 6,

web-based application around which other gene, mRNA, protein and BSM data from

external databases such as NCBI are collected. After undergoing semiautomated

processing and biocuration as described above, the data and annotations are stored in

SPP's Oracle 13c database. RESTful web services exposing SPP data, which are

served to responsively designed views in the user interface, were created using a Flat

UI Toolkit with a combination of JavaScript, D3.JS, AJAX, HTML5, and CSS3.

JavaServer Faces and PrimeFaces are the primary technologies behind the user

interface. SPP has been optimized for Firefox 24+, Chrome 30+, Safari 5.1.9+, and

Internet Explorer 9+, with validations performed in BrowserStack and load testing in

LoadUIWeb. XML describing each dataset and experiment is generated and submitted

to CrossRef (SCR_003217) to mint DOIs[10].

Data availability

Important note on data availability: this paper refers to the first versions of the consensomes and HCT intersection networks based on the datasets available at the time of publication. As additional CoV infection datasets are archived over time, we will make updated versions of the consensomes and HCT intersection analyses accessible in future releases. The entire set of experimental metadata is available in figshare File F1, section 1. Consensome data points are in figshare File F1, sections 2-5.

SPP SPP MERS[137], SARS1[141], SARS2[145] and IAV[149] consensomes, their underlying data points and metadata, as well as original datasets, are freely accessible at https://ww.signalingpathways.org. Programmatic access to all underlying data points and their associated metadata are supported by a RESTful API at https://www.signalingpathways.org/docs/. All SPP datasets are biocurated versions of publically archived datasets, are formatted according to the recommendations of the FORCE11 Joint Declaration on Data Citation Principles, and are made available under a Creative Commons CC BY 4.0 license. The original datasets are available are linked to from the corresponding SPP datasets using the original repository accession identifiers. These identifiers are for transcriptomic datasets, the Gene Expression Omnibus (GEO) Series (GSE); and for cistromic/ChIP-Seq datasets, the NCBI Sequence Read Archive (SRA) study identifier (SRP). SPP consensomes are assigned DOIs as shown in Table 1.

NDEx NDEx versions of consensomes (MERS[138], SARS1[142], SARS2[146] and IAV[150]) and node family (MERS[139], SARS1[143], SARS2[147] and IAV[151]) and node (MERS[140],

SARS1[144], SARS2[148] and IAV[152]) HCT intersection networks are freely available in the

NDEx repository and assigned DOIs as shown in Table 1. NDEx is a recommended

repository for Scientific Data, Springer Nature and the PLOS family of journals and is

registered on FAIRsharing.org; for additional info and documentation, please visit

http://ndexbio.org. The official SPP account in NDEx is available at:

https://bit.ly/30nN129.

Code availability

698   SPP source code is available in the SPP GitHub account under a Creative Commons

699   CC BY 4.0 license at https://github.com/signaling-pathways-project.

Acknowledgements

This work was supported by the National Institute of Diabetes, Digestive and Kidney Diseases NIDDK Information Network (DK097748), the National Cancer Institute (CA125123, CA184427) and by the Brockman Medical Research Foundation. We acknowledge the assistance of Apollo McOwiti, Shijing Qu and Michael Dehart in making the datasets and consensomes available in the SPP knowledgebase. We thank all investigators who archive their datasets, without whom SPP would not be possible.

Author contributions

Dataset biocuration: SO
Data analysis: SO, RP, NM
Manuscript drafting: NM
Manuscript editing: NM, RP, SO

Competing interests
The authors declare no competing interests.

References

References

1.   Takeda, K., Kaisho, T. & Akira, S. Toll-like receptors. *Annu. Rev. Immunol.* **21**,

335–376 (2003).

2.   Stark, G. R., Kerr, I. M., Williams, B. R., Silverman, R. H. & Schreiber, R. D. How

cells respond to interferons. *Annu. Rev. Biochem.* **67**, 227–64 (1998).

3.   Darnell, J. E., Kerr, I. M. & Stark, G. R. Jak-STAT pathways and transcriptional

activation in response to IFNs and other extracellular signaling proteins. *Science*

(80-.). **264**, 1415–1421 (1994).

4.   DeDiego, M. L. *et al.* Severe acute respiratory syndrome coronavirus envelope

protein regulates cell stress response and apoptosis. *PLoS Pathog.* **7**, e1002315

(2011).

5.   Josset, L. *et al.* Cell host response to infection with novel human coronavirus

EMC predicts potential antivirals and important differences with SARS

coronavirus. *MBio* **4**, e00165-13 (2013).

6.   Sims, A. C. *et al.* Release of severe acute respiratory syndrome coronavirus

nuclear import block enhances host transcription in human lung cells. *J. Virol.* **87**,

3885–3902 (2013).

7.   Yoshikawa, T. *et al.* Dynamic innate immune responses of human bronchial

epithelial cells to severe acute respiratory syndrome-associated coronavirus

infection. *PLoS One* **5**, e8729 (2010).

8.   Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives

Development of COVID-19. *Cell* **181**, 1036-1045.e9 (2020).

9.   Lamers, M. M. *et al.* SARS-CoV-2 productively infects human gut enterocytes.

Science https://doi.org/10.1126/science.abc1669 (2020).

10.   Ochsner, S. A. *et al.* The Signaling Pathways Project, an integrated 'omics

knowledgebase for mammalian cellular signaling pathways. *Sci. data* **6**, 252

(2019).

11.   Aevermann, B. D. *et al.* A comprehensive collection of systems biology data

characterizing the host response to viral infection. *Sci. data* **1**, 140033 (2014).

12.   Schneider, W. M., Chevillotte, M. D. & Rice, C. M. Interferon-stimulated genes: a

complex web of host defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).

13.   Doyle, T. *et al.* The interferon-inducible isoform of NCOA7 inhibits endosome-

mediated viral entry. *Nat. Microbiol.* **3**, 1369–1376 (2018).

14.   Chapgier, A. *et al.* A partial form of recessive STAT1 deficiency in humans. *J.*

Clin. Invest. **119**, 1502–1514 (2009).

15.   Gruter, P. *et al.* TAP, the human homolog of Mex67p, mediates CTE-dependent

RNA export from the nucleus. *Mol. Cell* **1**, 649–659 (1998).

16.   Wei, J. *et al.* Genome-wide CRISPR screen reveals host genes that regulate

SARS-CoV-2 infection. *bioRxiv* 2020.06.16.155101

https://doi.org/10.1101/2020.06.16.155101 (2020) .

17.   Shen, L. & Sinai, M. GeneOverlap: GeneOverlap: Test and visualize gene

overlaps. R package version 1.24.0 http://shenlab-sinai.github.io/shenlab-sinai/

(2020).

18.   Totura, A. L. *et al.* Toll-Like Receptor 3 Signaling via TRIF Contributes to a

Protective Innate Immune Response to Severe Acute Respiratory Syndrome Coronavirus Infection. *MBio* **6**, e00638-15 (2015).

19. Hensley, L. E. *et al.* Interferon-beta 1a and SARS coronavirus replication. *Emerg. Infect. Dis.* **10**, 317–319 (2004).

20. Appelberg, S. *et al.* Dysregulation in mTOR/HIF-1 signaling identified by proteo-transcriptomics of SARS-CoV-2 infected cells. *bioRxiv* https://doi.org/10.1101/2020.04.30.070383 (2020).

21. Wang, W. *et al.* Up-regulation of IL-6 and TNF-alpha induced by SARS-coronavirus spike protein in murine macrophages via NF-kappaB pathway. *Virus Res.* **128**, 1–8 (2007).

22. Zheng, K., Kitazato, K. & Wang, Y. Viruses exploit the function of epidermal growth factor receptor. *Rev. Med. Virol.* **24**, 274–286 (2014).

23. Ng, S. S. M., Li, A., Pavlakis, G. N., Ozato, K. & Kino, T. Viral infection increases glucocorticoid-induced interleukin-10 production through ERK-mediated phosphorylation of the glucocorticoid receptor in dendritic cells: potential clinical implications. *PLoS One* **8**, e63587 (2013).

24. Ostler, J. B., Harrison, K. S., Schroeder, K., Thunuguntla, P. & Jones, C. The Glucocorticoid Receptor (GR) Stimulates Herpes Simplex Virus 1 Productive Infection, in Part Because the Infected Cell Protein 0 (ICP0) Promoter Is Cooperatively Transactivated by the GR and Kruppel-Like Transcription Factor 15. *J. Virol.* **93**, (2019).

25. Hayward, S. D. Viral interactions with the Notch pathway. *Semin. Cancer Biol.* **14**,

784  387–396 (2004).

785 26. Ito, T. *et al.* The critical role of Notch ligand Delta-like 1 in the pathogenesis of

786  influenza A virus (H1N1) infection. *PLoS Pathog.* **7**, e1002341 (2011).

787 27. Rizzo, P. *et al.* COVID-19 in the heart and the lungs: could we 'Notch' the

788  inflammatory storm? *Basic research in cardiology* vol. 115 31 (2020).

789 28. Wang, S. *et al.* Xenobiotic pregnane X receptor (PXR) regulates innate immunity

790  via activation of NLRP3 inflammasome in vascular endothelial cells. *J. Biol.*

791  *Chem.* **289**, 30075–30081 (2014).

792 29. Marik, P. E., Kory, P. & Varon, J. Does vitamin D status impact mortality from

793  SARS-CoV-2 infection? *Medicine in drug discovery*

794  https://doi.org/10.1016/j.medidd.2020.100041 (2020).

795 30. Rhodes, J. M., Subramanian, S., Laird, E. & Kenny, R. A. Editorial: low population

796  mortality from COVID-19 in countries south of latitude 35 degrees North supports

797  vitamin D as a factor determining severity. *Alimentary pharmacology &*

798  *therapeutics* https://doi.org/10.1111/apt.15777 (2020).

799 31. Ledford, H. Coronavirus breakthrough: dexamethasone is first drug shown to save

800  lives. *Nature* https://doi.org/10.1038/d41586-020-01824-5 (2020).

801 32. Stukalov, A. *et al.* Multi-level proteomics reveals host-perturbation strategies of

802  SARS-CoV-2 and SARS-CoV. *bioRxiv* 2020.06.17.156455

803  https://doi.org/10.1101/2020.06.17.156455 (2020).

804 33. Poppe, M. *et al.* The NF-kappaB-dependent and -independent transcriptome and

805  chromatin landscapes of human coronavirus 229E-infected cells. *PLoS Pathog.*

13, e1006286 (2017).

34. Ludwig, S. & Planz, O. Influenza viruses and the NF-kappaB signaling pathway -

towards a novel concept of antiviral therapy. *Biol. Chem.* **389**, 1307–1312 (2008).

35. Ruckle, A. *et al.* The NS1 protein of influenza A virus blocks RIG-I-mediated

activation of the noncanonical NF-kappaB pathway and p52/RelB-dependent

gene expression in lung epithelial cells. *J. Virol.* **86**, 10211–10217 (2012).

36. Chiang, H.-S. & Liu, H. M. The Molecular Basis of Viral Inhibition of IRF- and

STAT-Dependent Immune Responses. *Front. Immunol.* **9**, 3086 (2018).

37. Frieman, M. *et al.* Severe acute respiratory syndrome coronavirus ORF6

antagonizes STAT1 function by sequestering nuclear import factors on the rough

endoplasmic reticulum/Golgi membrane. *J. Virol.* **81**, 9812–9824 (2007).

38. Garcia-Sastre, A. *et al.* The role of interferon in influenza virus tissue tropism. *J.*

Virol. **72**, 8550–8558 (1998).

39. Blaszczyk, K. *et al.* The unique role of STAT2 in constitutive and IFN-induced

transcription and antiviral responses. *Cytokine Growth Factor Rev.* **29**, 71–81

(2016).

40. Boudewijns, R. *et al.* STAT2 signaling as double-edged sword restricting viral

dissemination but driving severe pneumonia in SARS-CoV-2 infected hamsters.

bioRxiv https://doi.org/10.1101/2020.04.23.056838 (2020).

41. Haas, D. A. *et al.* Viral targeting of TFIIB impairs de novo polymerase II

recruitment and affects antiviral immunity. *PLoS Pathog.* **14**, e1006980 (2018).

42. Gelev, V. *et al.* A new paradigm for transcription factor TFIIB functionality. *Sci.*

Rep. **4**, 3664 (2014).

43.   Haviv, I., Shamay, M., Doitsh, G. & Shaul, Y. Hepatitis B virus pX targets TFIIB in
830   transcription coactivation. *Mol. Cell. Biol.* **18**, 1562–1569 (1998).

831   44.   Bellail, A. C., Olson, J. J. & Hao, C. SUMO1 modification stabilizes CDK6 protein
832   and drives the cell cycle and glioblastoma progression. *Nat. Commun.* **5**, 4234
833   (2014).

834   45.   Grossel, M. J. & Hinds, P. W. Beyond the cell cycle: a new role for Cdk6 in
835   differentiation. *J. Cell. Biochem.* **97**, 485–493 (2006).

836   46.   Kaldis, P., Ojala, P. M., Tong, L., Makela, T. P. & Solomon, M. J. CAK-
837   independent activation of CDK6 by a viral cyclin. *Mol. Biol. Cell* **12**, 3987–3999
838   (2001).

839   47.   Pauls, E. *et al.* Cell cycle control and HIV-1 susceptibility are linked by CDK6-
840   dependent CDK2 phosphorylation of SAMHD1 in myeloid and lymphoid cells. *J.*
841   *Immunol.* **193**, 1988–1997 (2014).

842   48.   Cingoz, O. & Goff, S. P. Cyclin-dependent kinase activity is required for type I
843   interferon production. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E2950–E2959 (2018).

844   49.   Hennessy, E. J., Sheedy, F. J., Santamaria, D., Barbacid, M. & O'Neill, L. A. J.
845   Toll-like receptor-4 (TLR4) down-regulates microRNA-107, increasing
846   macrophage adhesion via cyclin-dependent kinase 6. *J. Biol. Chem.* **286**, 25531–
847   25539 (2011).

848   50.   Handschick, K. *et al.* Cyclin-dependent kinase 6 is a chromatin-bound cofactor for
849   NF-kappaB-dependent gene expression. *Mol. Cell* **53**, 193–208 (2014).

51. Zaborowska, J., Isa, N. F. & Murphy, S. P-TEFb goes viral. *Bioessays* **38 Suppl**

1, S75-85 (2016).

52. Wobbe, C. R. *et al.* In vitro replication of DNA containing either the SV40 or the

polyoma origin. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **317**, 439–453 (1987).

53. Takahashi, K. *et al.* DNA topoisomerase 1 facilitates the transcription and

replication of the Ebola virus genome. *J. Virol.* **87**, 8862–8869 (2013).

54. Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the

SARS coronavirus. *Nature* **426**, 450–454 (2003).

55. Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2

and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e8

(2020).

56. Verdecchia, P., Cavallini, C., Spanevello, A. & Angeli, F. The pivotal link between

ACE2 deficiency and SARS-CoV-2 infection. *Eur. J. Intern. Med.*

https://doi.org/10.1016/j.ejim.2020.04.037 (2020).

57. Fadason, T. *et al.* A transcription regulatory network within the ACE2 locus may

promote a pro-viral environment for SARS-CoV-2 by modulating expression of

host factors. *bioRxiv* https://doi.org/10.1101/2020.04.14.042002 (2020).

58. Ziegler, C. G. K. *et al.* SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated

Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets

across Tissues. *Cell* https://doi.org/10.1016/j.cell.2020.04.035 (2020).

59. Zhu, H. *et al.* Clinical analysis of 10 neonates born to mothers with 2019-nCoV

pneumonia. *Transl. Pediatr.* **9**, 51–60 (2020).

60.   Sappenfield, E., Jamieson, D. J. & Kourtis, A. P. Pregnancy and susceptibility to

infectious diseases. *Infect. Dis. Obstet. Gynecol.* **2013**, 752852 (2013).

61.   Siston, A. M. *et al.* Pandemic 2009 influenza A(H1N1) virus illness among

pregnant women in the United States. *JAMA* **303**, 1517–1525 (2010).

62.   Pastva, A., Estell, K., Schoeb, T. R. & Schwiebert, L. M. RU486 blocks the anti-

inflammatory effects of exercise in a murine model of allergen-induced pulmonary

inflammation. *Brain. Behav. Immun.* **19**, 413–422 (2005).

63.   Breslin, N. *et al.* COVID-19 infection among asymptomatic and symptomatic

pregnant women: Two weeks of confirmed presentations to an affiliated pair of

New York City hospitals. *Am. J. Obstet. Gynecol. MFM* 100118

https://doi.org/10.1016/j.ajogmf.2020.100118 (2020).

64.   Sutton, D., Fuchs, K., D'Alton, M. & Goffman, D. Universal Screening for SARS-

CoV-2 in Women Admitted for Delivery. *The New England journal of medicine*

https://doi.org/10.1056/NEJMc2009316 (2020).

65.   Bazer, F. W., Burghardt, R. C., Johnson, G. A., Spencer, T. E. & Wu, G.

Interferons and progesterone for establishment and maintenance of pregnancy:

interactions among novel cell signaling pathways. *Reprod. Biol.* **8**, 179–211

(2008).

66.   Richards, J. S., Russell, D. L., Ochsner, S. & Espey, L. L. *Ovulation: New*

dimensions and new regulators of the inflammatory-like response. Annual Review

of Physiology vol. 64 (2002).

67.   Ghandehari, S. Progesterone for the Treatment of COVID-19 in Hospitalized Men.

(2020).

68.   Gordon, D. E. *et al.* A SARS-CoV-2 protein interaction map reveals targets for

drug repurposing. *Nature* https://doi.org/10.1038/s41586-020-2286-9 (2020).

69.   Lamouille, S., Xu, J. & Derynck, R. Molecular mechanisms of epithelial-

mesenchymal transition. *Nat. Rev. Mol. Cell Biol.* **15**, 178–196 (2014).

70.   Hill, C., Jones, M. G., Davies, D. E. & Wang, Y. Epithelial-mesenchymal transition

contributes to pulmonary fibrosis via aberrant epithelial/fibroblastic cross-talk. *J.*

lung Heal. Dis. **3**, 31–35 (2019).

71.   Li, H. *et al.* Alveolar epithelial cells undergo epithelial-mesenchymal transition in

acute interstitial pneumonia: a case report. *BMC Pulm. Med.* **14**, 67 (2014).

72.   Gouda, M. M., Shaikh, S. B. & Bhandary, Y. P. Inflammatory and Fibrinolytic

System in Acute Respiratory Distress Syndrome. *Lung* **196**, 609–616 (2018).

73.   George, P., Wells, A. & Jenkins, G. Pulmonary fibrosis and COVID-19: the

potential role for antifibrotic therapy. *Lancet* https://doi.org/10.1016/S2213-

2600(20)30225-3 (2020).

74.   Adair, L. B. 2nd & Ledermann, E. J. Chest CT Findings of Early and Progressive

Phase COVID-19 Infection from a US Patient. *Radiology case reports*

https://doi.org/10.1016/j.radcr.2020.04.031 (2020).

75.   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of

probable bat origin. *Nature* **579**, 270–273 (2020).

76.   Bolós, V. *et al.* The transcription factor Slug represses E-cadherin expression and

induces epithelial to mesenchymal transitions: a comparison with Snail and E47

repressors. *J. Cell Sci.* **116**, 499–511 (2003).

77. Yang, J. *et al.* HIF-2α promotes epithelial-mesenchymal transition through

regulating Twist2 binding to the promoter of E-cadherin in pancreatic cancer. *J.*

Exp. Clin. Cancer Res. **35**, 26 (2016).

78. Kobayashi, W. & Ozawa, M. The transcription factor LEF-1 induces an epithelial-

mesenchymal transition in MDCK cells independent of β-catenin. *Biochem.*

Biophys. Res. Commun. **442**, 133–138 (2013).

79. Xu, J., Lamouille, S. & Derynck, R. TGF-beta-induced epithelial to mesenchymal

transition. *Cell Res.* **19**, 156–172 (2009).

80. Zhao, M., Kong, L., Liu, Y. & Qu, H. dbEMT: an epithelial-mesenchymal transition

associated gene resource. *Sci. Rep.* **5**, 11459 (2015).

81. Moheimani, F. *et al.* Influenza A virus infection dysregulates the expression of

microRNA-22 and its targets; CD147 and HDAC4, in epithelium of asthmatics.

Respir. Res. **19**, 145 (2018).

82. Xia, L., Dai, L., Yu, Q. & Yang, Q. Persistent Transmissible Gastroenteritis Virus

Infection Enhances Enterotoxigenic Escherichia coli K88 Adhesion by Promoting

Epithelial-Mesenchymal Transition in Intestinal Epithelial Cells. *J. Virol.* **91**,

(2017).

83. Yang, J.-X. *et al.* Lipoxin A(4) ameliorates lipopolysaccharide-induced lung injury

through stimulating epithelial proliferation, reducing epithelial cell apoptosis and

inhibits epithelial-mesenchymal transition. *Respir. Res.* **20**, 192 (2019).

84. Diaz-Pina, G. *et al.* The Role of ADAR1 and ADAR2 in the Regulation of miRNA-

21 in Idiopathic Pulmonary Fibrosis. *Lung* **196**, 393–400 (2018).

85.   Vukmirovic, M. *et al.* Identification and validation of differentially expressed transcripts by RNA-sequencing of formalin-fixed, paraffin-embedded (FFPE) lung tissue from patients with Idiopathic Pulmonary Fibrosis. *BMC Pulm. Med.* **17**, 15 (2017).

86.   Gao, F., Kinnula, V. L., Myllarniemi, M. & Oury, T. D. Extracellular superoxide dismutase in pulmonary fibrosis. *Antioxid. Redox Signal.* **10**, 343–354 (2008).

87.   Kato, N. *et al.* Basigin/CD147 promotes renal fibrosis after unilateral ureteral obstruction. *Am. J. Pathol.* **178**, 572–579 (2011).

88.   Ruster, C. & Wolf, G. Angiotensin II as a morphogenic cytokine stimulating renal fibrogenesis. *J. Am. Soc. Nephrol.* **22**, 1189–1199 (2011).

89.   Wang, C. *et al.* CD147 Induces Epithelial-to-Mesenchymal Transition by Disassembling Cellular Apoptosis Susceptibility Protein/E-Cadherin/beta-Catenin Complex in Human Endometriosis. *Am. J. Pathol.* **188**, 1597–1607 (2018).

90.   Stewart, C. A. *et al.* SARS-CoV-2 infection induces EMT-like molecular changes, including ZEB1-mediated repression of the viral receptor ACE2, in lung cancer models. *bioRxiv* 2020.05.28.122291 https://doi.org/10.1101/2020.05.28.122291 (2020).

91.   Shan, B., Farmer, A. A. & Lee, W. H. The molecular basis of E2F-1/DP-1-induced S-phase entry and apoptosis. *Cell growth Differ. Mol. Biol. J. Am. Assoc. Cancer Res.* **7**, 689–697 (1996).

92.   Harbour, J. W. & Dean, D. C. The Rb/E2F pathway: expanding roles and

emerging paradigms. *Genes Dev.* **14**, 2393–2409 (2000).

93. Reimer, D. *et al.* E2F3a is critically involved in epidermal growth factor receptor-
directed proliferation in ovarian cancer. *Cancer Res.* **70**, 4613–4623 (2010).

94. Classon, M. & Harlow, E. The retinoblastoma tumour suppressor in development
and cancer. *Nat. Rev. Cancer* **2**, 910–917 (2002).

95. Diril, M. K. *et al.* Cyclin-dependent kinase 1 (Cdk1) is essential for cell division
and suppression of DNA re-replication but not for liver regeneration. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 3826–3831 (2012).

96. Shivji, K. K., Kenny, M. K. & Wood, R. D. Proliferating cell nuclear antigen is
required for DNA excision repair. *Cell* **69**, 367–374 (1992).

97. Borlado, L. R. & Méndez, J. CDC6: from DNA replication to cell cycle checkpoints
and oncogenesis. *Carcinogenesis* **29**, 237–243 (2008).

98. Holt, S. V *et al.* Silencing Cenp-F weakens centromeric cohesion, prevents
chromosome alignment and activates the spindle checkpoint. *J. Cell Sci.* **118**,
4889–4900 (2005).

99. Vanden Bosch, A. *et al.* NuSAP is essential for chromatin-induced spindle
formation during early embryogenesis. *J. Cell Sci.* **123**, 3244–3255 (2010).

100. Engeland, K. Cell cycle arrest through indirect transcriptional repression by p53: I
have a DREAM. *Cell Death Differ.* **25**, 114–132 (2018).

101. Wang, N. *et al.* Increased expression of RRM2 by human papillomavirus E7
oncoprotein promotes angiogenesis in cervical cancer. *Br. J. Cancer* **110**, 1034–
1044 (2014).

102. Li, Y.-Y., Wang, L. & Lu, C.-D. An E2F site in the 5'-promoter region contributes to

serum-dependent up-regulation of the human proliferating cell nuclear antigen

gene. *FEBS Lett.* **544**, 112–118 (2003).

103. Furukawa, Y., Terui, Y., Sakoe, K., Ohta, M. & Saito, M. The role of cellular

transcription factor E2F in the regulation of cdc2 mRNA expression and cell cycle

control of human hematopoietic cells. *J. Biol. Chem.* **269**, 26249–26258 (1994).

104. Yuan, X. *et al.* SARS coronavirus 7a protein blocks cell cycle progression at

G0/G1 phase via the cyclin D3/pRb pathway. *Virology* **346**, 74–85 (2006).

105. Fan, Y., Sanyal, S. & Bruzzone, R. Breaking Bad: How Viruses Subvert the Cell

Cycle. *Front. Cell. Infect. Microbiol.* **8**, 396 (2018).

106. Gasser, S. & Raulet, D. H. The DNA damage response arouses the immune

system. *Cancer Res.* **66**, 3959–3962 (2006).

107. Kong, L.-J., Chang, J. T., Bild, A. H. & Nevins, J. R. Compensation and specificity

of function within the E2F family. *Oncogene* **26**, 321–327 (2007).

108. To, B. & Andrechek, E. R. Transcription factor compensation during mammary

gland development in E2F knockout mice. *PLoS One* **13**, e0194937 (2018).

109. M. Bouhaddou & Krogan, N. The Global Phosphorylation Landscape of SARS-

CoV-2 Infection. *Cell* https://doi.org/10.1016/j.cell.2020.06.034 (2020).

110. Pillich, R. T., Chen, J., Rynkov, V., Welker, D. & Pratt, D. NDEx: A Community

Resource for Sharing and Publishing of Biological Networks. *Methods Mol. Biol.*

1558, 271–301 (2017).

111. Pratt, D. *et al.* NDEx 2.0: A Clearinghouse for Research on Cancer Pathways.

Cancer Res. **77**, e58–e61 (2017).

112. Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

113. Chen, X. *et al.* Detectable serum SARS-CoV-2 viral load (RNAaemia) is closely correlated with drastically elevated interleukin 6 (IL-6) level in critically ill COVID-19 patients. *Clin. Infect. Dis.* https://doi.org/10.1093/cid/ciaa449 (2020).

114. Sungnak, W. *et al.* SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat. Med.* **26**, 681–687 (2020).

115. Zhou, Z. *et al.* Heightened Innate Immune Responses in the Respiratory Tract of COVID-19 Patients. *Cell Host Microbe* **27**, 883-890.e2 (2020).

116. Cirillo, E., Parnell, L. D. & Evelo, C. T. A Review of Pathway-Based Analysis Tools That Visualize Genetic Variants. *Front. Genet.* **8**, 174 (2017).

117. Mitchell, H. D. *et al.* A network integration approach to predict conserved regulators related to pathogenicity of influenza and SARS-CoV respiratory viruses. *PLoS One* **8**, e69374 (2013).

118. Ackerman, E. E., Alcorn, J. F., Hase, T. & Shoemaker, J. E. A dual controllability analysis of influenza virus-host protein-protein interaction networks for antiviral drug target discovery. *BMC Bioinformatics* **20**, 297 (2019).

119. Wilk, C. M. Coronaviruses hijack the complement system. *Nat. Rev. Immunol.* https://doi.org/10.1038/s41577-020-0314-5 (2020).

120. Whetzel, P. L., Grethe, J. S., Banks, D. E. & Martone, M. E. The NIDDK

Information Network: A Community Portal for Finding Data, Materials, and Tools

for Researchers Studying Diabetes, Digestive, and Kidney Diseases. *PLoS One*

10, e0136206 (2015).

121. Bandrowski, A. E. & Martone, M. E. RRIDs: A Simple Step toward Improving

Reproducibility through Rigor and Transparency of Experimental Methods.

Neuron **90**, 434–436 (2016).

122. Ramilowski, J. A. *et al.* A draft network of ligand-receptor-mediated multicellular

signalling in human. *Nat. Commun.* **6**, 7866 (2015).

123. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-

sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

124. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. *Bioinformatics*

26, 139–140 (2010).

125. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock

linear model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29

(2014).

126. Pertea, M. *et al.* CHESS: a new human gene catalog curated from thousands of

large-scale RNA sequencing experiments reveals extensive transcriptional noise.

Genome Biol. **19**, 208 (2018).

127. Oki, S. *et al.* ChIP-Atlas: a data-mining suite powered by full integration of public

ChIP-seq data. *EMBO Rep.* **19**, (2018).

128. Muzikar, K., Nickols, N. & Dervan, P. Analysis of the dexamethasone (Dex)-

dependent transcriptome in A549 lung adenocarcinoma cells. Signaling Pathways

Project Datasets. https://doi.org/10.1621/xigKzGn1se (2015).

129. Schulze-Gahmen, U. *et al.* The AFF4 scaffold binds human P-TEFb adjacent to

HIV Tat. *Elife* **2**, e00327 (2013).

130. Chalabi Hagkarim, N. *et al.* Degradation of a Novel DNA Damage Response

Protein, Tankyrase 1 Binding Protein 1, following Adenovirus Infection. *J. Virol.*

92, (2018).

131. Yang, J. *et al.* Telomerase activation by Epstein-Barr virus latent membrane

protein 1 is associated with c-Myc expression in human nasopharyngeal epithelial

cells. *J. Exp. Clin. Cancer Res.* **23**, 495–506 (2004).

132. Klingelhutz, A. J., Foster, S. A. & McDougall, J. K. Telomerase activation by the

E6 gene product of human papillomavirus type 16. *Nature* **380**, 79–82 (1996).

133. Gewin, L., Myers, H., Kiyono, T. & Galloway, D. A. Identification of a novel

telomerase repressor that interacts with the human papillomavirus type-16 E6/E6-

AP complex. *Genes Dev.* **18**, 2269–2282 (2004).

134. Yin, L., Hubbard, A. K. & Giardina, C. NF-kappa B regulates transcription of the

mouse telomerase catalytic subunit. *J. Biol. Chem.* **275**, 36671–36675 (2000).

135. Akiyama, M. *et al.* Nuclear factor-kappaB p65 mediates tumor necrosis factor

alpha-induced nuclear translocation of telomerase reverse transcriptase protein.

Cancer Res. **63**, 18–21 (2003).

136. Ghosh, A. *et al.* Telomerase directly regulates NF-kappaB-dependent

transcription. *Nat. Cell Biol.* **14**, 1270–1281 (2012).

137.   *Signaling Pathways Project Datasets.* The MERS-CoV transcriptomic

consensome. https://doi.org/10.1621/jgxM527b8s.1 (2020)

138.   *The Network Data Exchange.* The MERS-CoV transcriptomic consensome

network. https://doi.org/10.18119/N9QG7S (2020)

139.   *The Network Data Exchange.* MERS-CoV node family high confidence

transcriptional target intersection analysis network.

https://doi.org/10.18119/N9PG63 (2020)

140.   *The Network Data Exchange.* MERS-CoV node high confidence transcriptional

target intersection analysis network. https://doi.org/10.18119/N96G6R (2020)

141.   *Signaling Pathways Project Datasets.* The SARS-CoV-1 transcriptomic

consensome. https://doi.org/10.1621/jgxM527b8s.1 (2020)

142.   *The Network Data Exchange.* The SARS-CoV-1 transcriptomic consensome

network. https://doi.org/10.18119/N9QG7S (2020)

143.   *The Network Data Exchange.* SARS-CoV-1 node family high confidence

transcriptional target intersection analysis network.

https://doi.org/10.18119/N9PG63 (2020)

144.   *The Network Data Exchange.* SARS-CoV-1 node high confidence transcriptional

target intersection analysis network. https://doi.org/10.18119/N96G6R (2020)

145.   *Signaling Pathways Project Datasets.* The SARS-CoV-2 transcriptomic

consensome. https://doi.org/10.1621/jgxM527b8s.1 (2020)

146.   *The Network Data Exchange.* The SARS-CoV-2 transcriptomic consensome

1091   network. https://doi.org/10.18119/N9QG7S (2020)

1092   147.   *The Network Data Exchange.* SARS-CoV-2 node family high confidence

1093   transcriptional target intersection analysis network.

1094   https://doi.org/10.18119/N9PG63 (2020)

1095   148.   *The Network Data Exchange.* SARS-CoV-2 node high confidence transcriptional

1096   target intersection analysis network. https://doi.org/10.18119/N96G6R (2020)

1097   149.   *Signaling Pathways Project Datasets.* The IAV transcriptomic consensome.

1098   https://doi.org/10.1621/jgxM527b8s.1 (2020)

1099   150.   *The Network Data Exchange.* The IAV transcriptomic consensome network.

1100   https://doi.org/10.18119/N9QG7S (2020)

1101   151.   *The Network Data Exchange.* IAV node family high confidence transcriptional

1102   target intersection analysis network. https://doi.org/10.18119/N9PG63 (2020)

1103   152.   *The Network Data Exchange.* IAV node high confidence transcriptional target

1104   intersection analysis network. https://doi.org/10.18119/N96G6R (2020)

1105

1106

1107

Figure Titles and Legends

Figure 1. Rankings of canonical interferon-stimulated genes (ISGs) in the viral consensomes. Shown are the percentile rankings of 20 ISGS[12] in the SARS1 **(a)**, SARS2 **(b)**, MERS **(c)** and IAV **(d)** consensomes. Note that numerous genes have identical q-value and percentile values and are therefore superimposed in the plots. Full underlying data are provided in figshare File 1. Please refer to the Methods section for a full description of the consensome algorithm.

Figure 2. High confidence transcriptional target (HCT) intersection analysis of viral infection and human receptors or signaling enzymes based on transcriptomic consensomes. Full numerical data are provided in figshare File F1, section 6. Due to space constraints some class and family names may differ slightly from those in the SPP knowledgebase. All q-values refer to those obtained using the GeneOverlap analysis package in R[17].

Figure 3. High confidence transcriptional target (HCT) intersection analysis of viral infection and human transcription factors based on ChIP-Seq consensomes. White cells represent $q > 5e-2$ intersections. Full numerical data are provided in figshare File F1, section 7. Due to space constraints some class and family names may differ slightly from those in the SPP knowledgebase. All q-values refer to those obtained using the GeneOverlap analysis package in R[17].

Figure 4. High confidence transcriptional target (HCT) intersection analysis of viral infection and human signaling enzymes based on ChIP-Seq consensomes. White cells represent non-significant ($q > 5e-2$) intersections. Full numerical data are provided in figshare File F1, section 7. Due to space constraints some class and family

names may differ slightly from those in the SPP knowledgebase. All q-values refer to

those obtained using the GeneOverlap analysis package in R[17].

Figure 5. Hypothesis generation use case 1: strain- and tissue-specific regulation

of *ACE2* in response to CoV infection of human cells. All data points are p < 0.05.

Refer to figshare File F1, section 1 for full details on the underlying datasets.

Figure 6. Hypothesis generation use case 2: antagonism between PGR and

SARS2 inflammatory signaling in the regulation of viral response genes in the

airway epithelium. GMFC: geometric mean fold change. PGR loss of function

experiments were retrieved from the SPP knowledgebase[128].

Figure 7. Hypothesis generation use case 3: evidence for a SARS2 infection-

associated EMT transcriptional signature. a. CoV HCT intersection with the

literature-curated EMT signature for all-biosample and lung epithelium-specific

consensomes. The IAV consensome is comprised of lung epithelial cell lines and was

therefore omitted from the lung epithelium-only consensome analysis. Refer to the

column "EMT" in figshare File F1, section 3 for the list of EMT SARS2 HCTs. q-values

refer to those obtained using the GeneOverlap analysis package in R[17]. **b.** Comparison

of mean percentile ranking of the EMT-associated SARS2 HCTs across viral

consensomes. Note that SARS2 HCTs are all in the 97-99[th] percentile and are therefore

superimposed in the scatterplot. Indicated are the results of the two-tailed two sample t-

test assuming equal variance comparing the percentile rankings of the SARS2 EMT

HCTs across the four viral consensomes.

Figure 8. Hypothesis generation use case 4: efficient SARS2 repression of E2F

family HCTs encoding key cell cycle regulators. a. Relative abundance of E2F HCT

cell cycle regulators in response to SARS2 infection. **b.** Comparison of SARS2, SARS1,

MERS and IAV consensome percentiles of the E2F HCT cell cycle regulators. Indicated

are the results of the two-tailed two sample t-test assuming equal variance comparing

the percentile rankings of the SARS2 EMT HCTs across the four viral consensomes.

Figure 9. Mining of CoV consensomes and underlying data points in the SPP

knowledgebase. a. The Ominer query form can be configured as shown to access the

CoV infection consensomes. In the example shown, the user wishes to view the SARS2

consensome. **b**. Consensomes are displayed in a tabular format. Target transcript

symbols in the consensomes link to SPP transcriptomic Regulation Reports (**c**) **c.**

Regulation Reports for consensome transcripts are filtered to show only data points that

contributed to their consensome ranking. Clicking on a data point opens a Fold Change

Information window that links to the SPP curated version of the original archived dataset

(d). **d.** Like all SPP datasets, CoV infection datasets are comprehensively aligned with

FAIR data best practice and feature human-readable names and descriptions, a DOI,

one-click addition to citation managers, and machine-readable downloadable data files.

For a walk-through of CoV consensome data mining options in SPP, please refer to the

accompanying YouTube video (http://tiny.cc/2i56rz).

Figure 10. Visualization of viral consensomes and HCT intersection networks in

the NDEx repository. In all panels, transcripts (consensome networks; panels a, b & c)

and nodes (HCT intersection network; panel d) are color-coded according to their

category as follows: receptors (orange), enzymes (blue), transcription factors (green),

ion channels (mustard) and co-nodes (grey). Additional contextual information is

available in the description of each network on the NDEx site. Red arrows are explained

in the text. **a.** Sample view of SARS2 consensome showing top 5% of transcripts. White

rectangles represent classes to which transcripts have been mapped in the SPP

biocuration pipeline[10]. Orange rectangle refers to the view in panel b. **b.** Zoomed-in view

of orange rectangle in panel A. IL6 transcript is highlighted to show the contextual

information available in the side panel. **c.** Top 20 ranked transcripts in the SARS2

consensome. Edge widths are proportional to the GMFC. **d.** Selected classes

represented in the top 5% of nodes in the SARS2 ChIP-Seq HCT intersection network.

Node circle size is inversely proportional to the intersection q-value.

1185

 Tables and Table legends

 Table 1. DOI-driven links to consensomes and HCT intersection networks. SPP

 DOIs point to the web browser version of the consensome, which contains a

 downloadable version of the full consensome. For clarity of visualization, NDEx

 consensome DOIs point to networks containing transcripts in the top 5% of each

 consensome (i.e. HCTs for each viral infection); the full consensome network can be

 reached from this page. Similarly, NDEx HCT intersection DOIs point to networks

 containing nodes in the top 5% of each HCT intersection network; the full HCT

 intersection network can be reached from this page. TX, transcriptomic node family

 intersection; CX, ChIP-Seq node intersection.

1197

Table 1

Virus	Resource	Network type	DOI	Reference
MERS-CoV	SPP	Consensome	10.1621/jgxM527b8s.1	137
	NDEx	Consensome	10.18119/N9QG7S	138
		HCT intersection (TX)	10.18119/N9PG63	139
		HCT intersection (CX)	10.18119/N96G6R	140
SARS-CoV-1	SPP	Consensome	10.1621/jgxM527b8s.1	141
	NDEx	Consensome	10.18119/N9KP4G	142
		HCT intersection (TX)	10.18119/N9JS46	143
		HCT intersection (CX)	10.18119/N92P56	144
SARS-CoV-2	SPP	Consensome	10.1621/k9ygy4i49j.1	145
	NDEx	Consensome	10.18119/N9G02W	146
		HCT intersection (TX)	10.18119/N9F016	147
		HCT intersection (CX)	10.18119/N9Z01V	148
IAV	SPP	Consensome	10.1621/58AOyXDIAH.1	149
	NDEx	Consensome	10.18119/N9B60Z	150
		HCT intersection (TX)	10.18119/N9989R	151
		HCT intersection (CX)	10.18119/N9T609	152

1198

1199

1200

1201

1202

1203

1204

Pathway node family transcriptomic HCTs			Viral HCTs intersection q-value			
Category	Class	Family	SARS1	SARS2	MERS	IAV
Receptors	Catalytic receptors	Collagen receptors				
		Epidermal growth factor receptors				
		Fibroblast growth factor receptors				
		Insulin receptor family				
		Interferon receptor family				
		Notch receptors				
		Toll-like receptors				
		Transforming growth factor-β receptor family				
		Tumor necrosis factor receptors				
	G protein coupled receptors	Class Frizzled GPCRs				
	Nuclear receptors	Androgen receptor				
		Estrogen receptors				
		Estrogen-related receptors				
		Glucocorticoid receptor				
		Peroxisome proliferator-activated receptors				
		Progesterone receptor				
		Retinoic acid receptors				
		Retinoid X receptors				
		Vitamin D receptor				
		Xenobiotic receptors				
Enzymes	Kinases	Abl kinases (ABL)				
		Cyclin-dependent kinases (CDK)				
		Src kinases				
	Nucleotidyltransferases	Telomerase reverse transcriptase				
		DNA topoisomerases (TOP)				

q-value

color	range
	<1E-65
	1E-60-1E-65
	1E-55-1E-60
	1E-50-1E-55
	1E-45-1E-50
	1E-40-1E-45
	1E-35-1E-40
	1E-30-1E-35
	1E-25-1E-30
	1E-20-1E-25
	1E-15-1E-20
	1E-10-1E-15
	1E-5-1E-10
	5E-2-1E-5

Heat-map data table. Column headers for both panels:

Left panel — Pathway node ChIP-Seq consensome HCTs / Viral HCTs intersection q-value

Class	Family	Node	SARS1	SARS2	MERS	IAV
ARID domain	ARID1 family	ARID1A				
	ARID2 family	ARID2				
	ARID3 family	ARID3A				
BHLH factors	Ahr-like family	AHR				
		AHRR				
		EPAS1				
		HIF1A				
	AP-2 family	TFAP2A				
		TFAP2C				
	AP-4 family	TFAP4				
	Arnt-like	ARNT				
		ARNTL				
		CLOCK				
	E2A-related	TCF12				
		TCF3				
		TCF4				
	Hairy-related	BHLHE40				
		HEY1				
	Mad-like	MXI1				
	MESP	TCF21				
	Mondo-like	MLXIP				
	Myc / Max	MAX				
		MYC				
		MYCN				
	Myogenic TFs	MYF5				
		MYOD1				
	Neurogenin-Atonal like	NEUROD1				
	SREBP	SREBF1				
		SREBF2				
	Tal/HEN-like	TAL1				
	TFE3-like	MITF				
		TFEB				
	Twist-like	HAND2				
	USF	USF1				
		USF2				
BZIP factors	ATF-2-like	ATF2				
		ATF3				
		JDP2				
	ATF-4-related	ATF4				
	B-ATF-related	BATF				
		BATF3				
	C/EBP	CEBPA				
		CEBPB				
		CEBPD				
		DDIT3				
	CREB-like	CREB1				
	Fos	FOS				
		FOSL1				
		FOSL2				
	Jun	JUN				
		JUNB				
		JUND				
	Large Maf	MAF				
	NF-E2-like	BACH1				
		BACH2				
		NFE2L2				
	Small Maf	MAFF				
		MAFG				
		MAFK				
C2H2 Zn finger factors	B-cell lymphoma 13	BCL11B				
	B-cell lymphoma 2	BCL11A				
	BCL6	BCL6				
	CTCF-like	CTCF				
		CTCFL				
	Early growth response	EGR1				
	GFI1	GFI1				
	Gli-like	GLI2				
	Hypermethy. in Cancer	HIC1				
	Ikaros	IKZF1				
	Krüppel-like	KLF1				
		KLF11				
		KLF4				
		KLF5				
		KLF6				
	MAZ-like	MAZ				
	PLAG Zinc Finger	PLAG1				
	REST	REST				
	Sal-like	SALL3				
	Snail-like	**SNAI2**				
	Sp1-like	SP1				
		SP2				
		SP4				
	YY1-like	YY1				
	ZBTB17	ZBTB17				
	ZBTB7	ZBTB7A				
	ZFX/ZFY	ZFX				
	ZNF263	ZNF263				
	ZNF341	ZNF341				
	ZNF362-like	ZNF384				
	ZNF366-like	ZNF366				
	ZNF639-like	ZNF711				
	ZNF76-like	ZNF143				
	ZNF83	ZNF83				
	ZNF92	ZNF92				
	ZNF99-like	ZBTB48				
CXXC zinc finger	CpG-binding protein	CXXC1				
E2F/FOX	Dp-1	**TFDP1**				
	E2F	**E2F1**				
		E2F4				
		E2F6				
	FOXA	FOXA1				
		FOXA2				
	FOXF	FOXF1				
	FOXH	FOXH1				
	FOXK	FOXK1				

Right panel — Pathway node ChIP-Seq consensome / Viral HCTs intersection q-value

Class	Family	Node	SARS1	SARS2	MERS	IAV
E2F/FOX	FOXM	FOXM1				
	FOXO	FOXO1				
	FOXP	FOXP1				
		FOXP2				
Forkhead/winged helix	Regulatory X (RFX)	RFX5				
Grainyhead domain	GRH-like protein	GRHL2				
		GRHL3				
Heat shock factors	HSF	HSF1				
Heteromeric CCAAT bdg	Heteromeric CCAAT bdg	NFYA				
		NFYB				
		NFYC				
HMG domain	Canonical HMG protein	HMGB2				
	Group B	SOX2				
		SOX3				
	Group E	SOX9				
	TCF-7-related	**LEF1**				
		TCF7L2				
	UBF-related	UBTF				
	WHSC1-related	SSRP1				
Homeo domain	Caudal type homeobox (CDX)	CDX2				
	HNF1-like	HNF1B				
	HOX4	HOXA4				
	HOX5	HOXC5				
	HOX6-7	HOXA6				
	Nanog homeobox	NANOG				
	NK-2.1	NKX2-1				
	Oct-1/2-like (POU2)	POU2F2				
	Orthodenticle homeobox	OTX2				
	PBX	PBX1				
		PBX3				
		PBX4				
	SATB	SATB1				
	SIX1-like	SIX1				
		SIX2				
	TLX	TLX1				
	ZHX	ZHX2				
	Zn finger E-box bdg homeobox	ZEB1				
MADS box	Myocyte enhancer 2	MEF2A				
		MEF2B				
		MEF2C				
	Responders to external signals	SRF				
p53 domain	p53-related	TP53				
		TP63				
Paired box	PAX-2-like (partial homeobox)	PAX5				
Rel Homology Region	Early B-Cell-related	EBF1				
		EBF3				
	IkappaB-related	BCL3				
	M	RBPJ				
	NFAT-related	NFATC1				
	NF-kappaB p50 subunit-like	NFKB1				
		NFKB2				
		REL				
		RELA				
		RELB				
Runt domain	Core-binding subunit	CBFB				
		RUNX1				
		RUNX1T1				
		RUNX2				
		RUNX3				
SAND domain	Sp140-like	SP140				
SMAD/NF1 DBD	Co-activating (Co) Smads	SMAD4				
	Regulatory (R) Smads	SMAD1				
		SMAD2				
		SMAD3				
STAT domain	STAT	STAT1				
		STAT2				
		STAT3				
		STAT4				
		STAT5A				
		STAT5B				
T Box factors	TBrain-related	TBX21				
	TBX2-related (TBX)	TBX2				
TEA domain	TEF-1-related	TEAD1				
		TEAD4				
Tryptophan cluster	EHF-like	ELF3				
	Elf-1-like	ELF1				
	Elk-like	ELK1				
		ELK4				
		ETV1				
		ETV4				
		ETV5				
	Ets-like	ERG				
		ETS1				
		FLI1				
		GABPA				
	Interferon-regulatory	IRF1				
		IRF2				
		IRF4				
		IRF8				
	Myb-like	MYB				
		MYBL2				
	Nuclear receptor corepressor	NCOR1				
		NCOR2				
	REST corepressor	RCOR1				
	Spi-like	SPI1				
Others	Bromodomain PHD finger	BPTF				
	Nuclear I	NFIC				
	PR/SET domain	PRDM1				
	Single GATA-type zinc-finger	MTA3				
		TRPS1				
	Two zinc-finger GATA	GATA1				
		GATA2				
		GATA3				
		GATA4				
		GATA6				

q-value legend:

q-value
<1E-30
1E-25-1E-30
1E-20-1E-25
1E-15-1E-20
1E-10-1E-15
1E-5-1E-10
1E-2-1E-5
5E-2-1E-2

Pathway node ChIP-Seq consensome HCTs

Viral HCTs intersection q-value

Class	Family	Node	SARS1	SARS2	MERS	IAV
Acetyltransferases	CBP/p300	CREBBP				
		EP300				
	Lysine acetyltransferases (KAT)	KAT7				
	Nuclear receptor coactivator (NCOA)	NCOA1				
ATPases	EP400	EP400				
Deacetylases	Histone deacetylases (HDAC)	HDAC1				
		HDAC2				
		HDAC6				
Demethylases	Histone-H3-lysine-36 demethylases (KDM)	KDM1A				
		KDM2B				
		KDM4A				
		KDM4C				
		KDM5A				
		KDM5B				
		KDM5D				
		KDM6A				
		KDM6B				
	Jumonji domain containing	JMJD1C				
		JMJD6				
Dioxygenases	Ten-eleven translocation (TET)	TET2				
E3 ubiquitin ligases	BRCA1	BRCA1				
	Protein inhibitor of activated STAT (PIAS)	PIAS1				
	Tripartite motif-containing (TRIM)	TRIM24				
		TRIM25				
		TRIM28				
Helicases (DNA)	ATRX chromatin remodeler	ATRX				
	Chromodomain-helicases-DNA binding (CHD)	CHD1				
		CHD2				
	ERCC excision repair (ERCC)	ERCC3				
Kinases	Cyclin-dependent kinases (CDK)	CDK6				
		CDK7				
		CDK8				
		CDK9				
	Extracellular signal-regulated kinases (ERK)	MAPK1				
	Mammalian target of rapamycin (MTOR)	MTOR				
	Mitogen-activated protein kinases (MAPK)	MAPK14				
	Protein kinase C (PKC)	PRKCQ				
Methyltransferases	ASH like methyltransferase	ASH2L				
	DNA (cytosine-5-)-methyltransferases (DNMT)	DNMT3A				
	Euchromatic histone-lysine N-methyltransferases	EHMT2				
	Histone-lysine N-methyltransferases (KMT)	KMT2A				
		KMT2B				
		KMT2C				
		KMT2D				
	Protein arginine methyltransferases (PRMT)	PRMT1				
	SET domain-containing (SETD)	SETD1A				
Peptidases	Proteasome 20S subunit	PSMB5				
Regulatory factors	Cyclin-dependent kinase inhibitors (CDKN)	CDKN1B				
	Cyclins (CCN)	CCND2				
		CCNT2				
	Elongation factors for RNA polymerase II	ELL2				
	PAF1 homolog	PAF1				
	Protein phosphatase 1 regulatory subunits	NONO				
		SFPQ				
Ribonucleases	Argonaute	AGO2				
Topoisomerases	DNA topoisomerases (TOP)	TOP1				
Other enzymes	Recombination activating	RAG1				
		RAG2				

q-value

Color	Range
	<1E-9
	1E-8-1E-9
	1E-7-1E-8
	1E-6-1E-7
	1E-5-1E-6
	1E-4-1E-5
	1E-3-1E-4
	1E-2-1E-3
	5E-2-1E-2

Symbol	Name	log10 GMFC SARS2	log10 FC PGR LOF	
			RU/DEX vs V	RU vs V (DEX)
CXCL1	C-X-C motif chemokine ligand 1	0.36		0.18
CXCL2	C-X-C motif chemokine ligand 2	0.53	0.57	
IER3	immediate early response 3	0.21		0.53
IFIT3	interferon induced protein with tetratricopeptide repeats 3	0.96	0.16	0.26
IFITM3	interferon induced transmembrane protein 3	0.33	0.16	0.20
IL1B	interleukin 1 beta	0.38	0.30	0.29
ISG15	ISG15 ubiquitin like modifier	0.70	0.23	0.27
ISG20	interferon stimulated exonuclease gene 20	0.44	0.20	0.22
NFKBIA	NFKB inhibitor alpha	0.54	0.24	
OAS1	2'-5'-oligoadenylate synthetase 1	0.45	0.16	0.28
STAT1	signal transducer and activator of transcription 1	0.30	0.21	0.14
TNFAIP3	TNF alpha induced protein 3	0.65	0.27	

a.

	EMT signature intersection analysis					
	All tissues viral HCTs			Lung epithelium viral HCTs		
Virus	INT	OR	*q*	INT	OR	*q*
MERS	19	1.3	1.8e-1	13	1.3	1.9e-1
SARS1	15	1.2	2.3e-1	16	1.2	1.9e-1
SARS2	34	5.8	4.5e-14	32	3.7	1.1e-8
IAV	24	2.5	1.4e-4			

b.

The Signaling Pathways Project

A multi-omics knowledgebase for cellular signaling pathways

Home   Datasets   Ominer   Collaborations   About

A. Ominer query form

Target gene(s) of interest

Consensome

'Omics Category

Transcriptomics

Signaling Pathway Module Category

Models

Viral infection

Severe Acute Respiratory Syndrome coronavirus 2 (SARS-CoV-2)

B. Consensome

Calculated across **215, 000** data points from **17** experiments in **7** datasets.

Search: _______

Target	Gene Name	Discovery Rate	GMFC	CPV	Percentile
PDZK1IP1	PDZK1 interacting protein 1	0.882	1.943	3.77E-18	100
BAAT	bile acid-CoA:amino acid N-acyltransferase	0.824	1.646	3.60E-16	99
CFB	complement factor B	0.824	2.822	3.60E-16	99
IFI27	interferon alpha inducible protein 27	0.824	5.078	3.60E-16	99
OAS1	2'-5'-oligoadenylate synthetase 1	0.824	2.833	3.60E-16	99
PIGR	polymeric immunoglobulin receptor	0.824	1.777	3.60E-16	99

D. Dataset

Overview

Dataset Name :
Analysis of the SARS-CoV-2-, RSV-, and IAV-dependent transcriptomes in human primary bronchial epithelial cells and alveolar basal epithelial cells

Description :
Human NHBE primary bronchial epithelial cells and A549 alveolar basal epithelial cells were mock infected or infected at a MOI of 2 and 0.2 respectively with SARS-CoV-2 USA-WA1/2020.

Dataset Type : Transcriptomic   **Release Date :** May 08, 2020   **DOI :** 10.1621/AVmtFHml4t

Version : Version 1.0 of an annotated derivative of the original dataset, which can be found in GSE147507

Dataset Citation : 🌐
tenOever BR and Blanco-Melo D (2020) Analysis of the SARS-CoV-2-, RSV-, and IAV-dependent transcriptomes in human primary bronchial epithelial cells and alveolar basal epithelial cells, v 1.0 Signaling Pathways Project Datasets. 10.1621/AVmtFHml4t

Download Citation :

Associated Article : 🌐
Blanco-Melo D, Nilsson-Payant BE, Liu WC, Uhl S, Hoagland D, Møller R, Jordan TX, Oishi K, Panis M, Sachs D, Wang TT, Schwartz RE, Lim JK, Albrecht RA and tenOever BR (2020) Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19 Cell **181** 1036-1045 View Abstract | View PubMed

 Download Dataset

C. Regulation Report

Display by: Biosample ▾

Transcript Relative Abundance (Fo

Fold Change Information ✕

Symbol:IFI27
Fold Change:8.519
p value:1e-7
Biosample:Respiratory,Lung,epithelium, Calu-3 cells
Experiment:SARS-CoV-2 Wuhan/IVDC-HB-01/2019 vs Mock | 12 h
Species:Human

More Information

Gastrointestinal

Other

small intestine, enteroids

Respiratory

Lung

epithelium, bronchus, NHBE cells

epithelium, A549 cells

epithelium, Calu-3 cells