Identification and Characterization of Fbxl22, a novel skeletal muscle atrophy-promoting E3 ubiquitin ligase

David C. Hughes[1†], Leslie M. Baehr[1†], Julia R. Driscoll[2], Sarah A. Lynch[2], David S. Waddell[2*] & Sue C. Bodine[1*]

[1]Department of Internal Medicine, Division of Endocrinology and Metabolism, Carver College of Medicine, University of Iowa, Iowa City, Iowa, 52242 USA.

[2]University of North Florida, Department of Biology, 1 UNF Drive, Jacksonville, Florida, 32224, USA.

[†] These authors contributed equally

*Co-Corresponding Authors:

Sue C. Bodine, University of Iowa, Dept. of Internal Medicine, 1283D CBRB, 285 Newton Rd, Iowa City, IA 52242 (e-mail: sue-bodine@uiowa.edu)

David S. Waddell, University of North Florida, Department of Biology, 1 UNF Drive, Jacksonville, FL, 32224 (e-mail: d.s.waddell@unf.edu)

Abstract

Muscle-specific E3 ubiquitin ligases have been identified in muscle atrophy-inducing conditions. The purpose of the current study was to explore the functional role of Fbxl22, and a newly identified splice variant (Fbxl22-193), in skeletal muscle homeostasis and neurogenic muscle atrophy. In mouse C_2C_{12} muscle cells, promoter fragments of the Fbxl22 gene were cloned and fused with the secreted alkaline phosphatase reporter gene to assess the transcriptional regulation of Fbxl22. The

tibialis anterior muscles of male C57/BL6 mice (12-16 weeks old) were electroporated with expression plasmids containing the cDNA of two Fbxl22 splice variants and tissues collected after 7, 14 and 28 days. Gastrocnemius muscles of wild type and MuRF1 knockout mice were electroporated with an Fbxl22 RNAi or empty plasmid, denervated three days post-transfection, and tissues collected 7 days post-denervation. The full-length gene and novel splice variant are transcriptionally induced early (after 3 days) during neurogenic muscle atrophy. In vivo overexpression of Fbxl22 isoforms in mouse skeletal muscle lead to evidence of myopathy/atrophy suggesting that both are involved in the process of neurogenic muscle atrophy. Knockdown of Fbxl22 in MuRF1 KO muscles resulted in significant additive muscle sparing at 7 days of denervation. Targeting two E3 ubiquitin ligases appears to have a strong additive effect on protecting muscle mass loss with denervation and these findings have important implications in the development of therapeutic strategies to treat muscle atrophy.

Key Words: Denervation, electroporation, MuRF1, muscle atrophy, protein degradation

Running head: Fbxl22 and Muscle Atrophy

Abbreviations

AAV, adeno-associated viral vector; CSPD, 3-(1-chloro-3′-methoxyspiro-(adamantane-4,4′-dioxetane)-3′-yl) phenyl dihydrogen phosphate; DMEM, Dulbecco's Modified Eagle's Medium; Fbxl22, F-box and leucine-rich protein 22; GEO, Gene Expression Omnibus; GFP, Green Fluorescent Protein*;* H&E*,* Hematoxylin & Eosin*;* LGA, lateral gastrocnemius; MAFbx, muscle atrophy F-box; MGA, medial gastrocnemius; MuRF1, muscle-specific RING finger 1; MUSA1, muscle ubiquitin ligase of the SCF complex in

46   atrophy-1; MRF, myogenic regulatory factor; RLU, Relative Light Units; SEAP,

47   Secreted Alkaline Phosphatase; SMART, Specific of Muscle Atrophy and Regulated by

48   Transcription; TA, tibialis anterior; TGFβ1, transforming-growth factor-β1; UBR5,

49   ubiquitin protein ligase E3 component n-recognin 5; UPS, ubiquitin-proteasome system.

Introduction

Skeletal muscle wasting can occur as a result of both internal (e.g. genetic disorders) and external stimuli (e.g. disuse) (16, 23, 69). Central to the process of muscle wasting/atrophy is increased protein degradation through the ubiquitin-proteasome system (UPS) and autophagosome lysosome system (5, 8, 55).   The UPS targets proteins for degradation by the 26S proteasome. The ubiquitin ligase enzyme, or E3, is critical for catalyzing the movement of ubiquitin from the E2 to the protein substrate which can lead to altered function or degradation by the proteasome (5, 55). There are approximately 40 E2 enzymes and over 600 E3 ubiquitin ligases in the human genome making it an extremely complex and relatively understudied system (5, 35). In skeletal muscle, only a few E3 ubiquitin ligases have been characterized and associated with muscle atrophy, including MuRF1 (muscle-specific RING finger 1), MAFbx (muscle atrophy F-box; also known as Atrogin-1) (9, 28), MUSA1 (muscle ubiquitin ligase of the SCF complex in atrophy-1) (57) and SMART (Specific of Muscle Atrophy and Regulated by Transcription) (45). Recently, we identified UBR5 as an E3 ubiquitin ligase that may play a role in muscle growth and hypertrophy (59). Considerably more E3 ubiquitin ligases are expressed in skeletal muscle, most of which remain to be explored in the context of muscle physiology and disease.

Of all the E3 ubiquitin ligases, MuRF1 and MAFbx have been the most studied, especially during the progression of skeletal muscle atrophy induced by a wide range of stressors (such as neural inactivity, starvation, and aging) since their discovery in 2001 (9, 28). Under certain atrophy stimuli, increased MuRF1 and MAFbx expression occurs via activation of the glucocorticoid receptor and/or FoxO transcription factors (56, 66). A

null deletion in MuRF1 and MAFbx leads to significant muscle sparing after 14 days of

neural inactivity (9). Interestingly, after 7 days of neural inactivity, MuRF1 knockout (KO)

animals display muscle atrophy similar to wild type animals, suggesting other

unidentified factors contribute to the process of skeletal muscle wasting (9, 21). In our

previously published microarray data set, 400-600 genes showed significant up-

regulation in wild type and/or MuRF1 KO skeletal muscle exposed to neural inactivity for

3 days (25). Of the 400-600 genes identified, many were originally identified by the

RIKEN project and thus of unknown name and function at the time of the study (36).

One such RIKEN gene from the microarray data set was subsequently identified to be

an E3 ubiquitin ligase from the F-box family, named F-box and leucine-rich protein 22

(Fbxl22) (25). In cardiac muscle, Fbxl22 is enriched and, similar to MAFbx, a

component of the Skp1-Cullin-F-box (SCF) ubiquitin ligase complex through its N-

terminal F-box domain (9, 28, 62). Using a yeast 2-hybrid screen and cardiomyocytes,

Fbxl22 has been suggested to target filamin-C and α-actinin-2 for proteasomal

degradation (62). Filamin-C and α-actinin-2 are sarcomeric proteins localized to the Z-

line and are critical for maintaining the spatial relationship between myofilaments and

stabilizing the myofilament lattice (4, 34, 39). Little is known about the role of Fbxl22 in

skeletal muscle and its involvement in the atrophy process. Therefore, we sought to

characterize Fbxl22 expression and function in muscle cells and investigate the role of

Fbxl22 in skeletal muscle homeostasis and neural inactivity-induced atrophy.

Materials and methods

Cell Culture

Cell culture was performed as previously described (29, 30). Briefly, mouse C_2C_{12} myoblasts were obtained from the American Type Culture Collection (Manassas, VA) and maintained in Dulbecco's Modified Eagle's Medium (DMEM) (Thermo Fischer Scientific, Waltham, MA) supplemented with 10% Fetal Bovine Serum (FBS) (GE Healthcare HyClone Laboratories, Logan, UT), non-essential amino acids (GE Healthcare HyClone Laboratories, Logan, UT), 1X Penicillin/Streptomycin and Gentamicin (Thermo Fischer Scientific, Waltham, MA). C_2C_{12} cells were maintained in a humidified chamber at 37°C and 5% CO_2 and grown to 100% confluency and then switched to DMEM supplemented with 2% serum to induce myogenic differentiation.

Secreted Alkaline Phosphatase (SEAP) reporter gene assays

SEAP reporter assays were performed as previously described (29, 30, 42). All reporter assays were conducted in C_2C_{12} cells cultured in 12-well plates at a seeding density of 100,000 cells/well. Cells were transfected at 70-80% confluency with 1 µg of total DNA/well (250 ng reporter plasmid, 125 ng internal control plasmid, 125–250 ng of expression plasmids, and empty pBluescript vector as filler DNA to 1 µg/well). DNA was complexed with Turbofect transfection reagent (Thermo Fisher Scientific, Waltham, MA) for 20 minutes prior to overlaying of cells according to the manufacturer's protocol. The media was sampled at 24-hours post-transfection and then switched to DMEM supplemented with 2% serum to induce differentiation. Media was sampled at 2- and 7-days post-media change and reporter activity was assessed using the 3-(1-chloro-3′-

methoxyspiro-(adamantane-4,4′-dioxetane)-3′-yl) phenyl dihydrogen phosphate (CSPD)

chemiluminescent substrate and the Phospha-Light SEAP Reporter Gene Assay

System according to the manufacturer's protocol (Thermo Fisher Scientific, Waltham,

MA). Luminescence levels were detected using a Synergy 2 microplate reader set for

an endpoint read with a 2s integration time. At the end of the time course, cells were

lysed with Passive Lysis Buffer (Promega, Madison, WI) and homogenates were

cleared by centrifugation and analyzed for β-galactosidase activity. β-galactosidase

activity was determined by incubating cell lysates with ortho-nitrophenyl-β-D-

galactopyranoside (ONPG) dissolved in Z Buffer (60 mM $Na_2HPO_4 \cdot 7H_2O$, 40 mM

$NaH_2PO_4 \cdot H_2O$, 10 mM KCl, 1 mM $MgSO_4$, 50 mM β-mercaptoethanol, pH 7.0)

overnight at 37°C and subsequently read using a Synergy 2 plate reader at 420nm.

Reporter activity was expressed as relative light units of SEAP divided by β-

galactosidase activity.

RNA isolation and cDNA synthesis from C_2C_{12} cells

Total RNA was isolated from C_2C_{12} myoblasts and differentiated myotubes using the

RNeasy Mini kit (Qiagen, Valencia, CA) according the manufacturer's protocol. Purified

total RNA was reverse transcribed to cDNA using M-MLV Reverse Transcriptase and a

Poly T primer according to the manufacturer's instructions (Thermo Fisher Scientific,

Waltham, MA).

Cloning of a novel Fbxl22 cDNA and promoter fragments

A novel open reading frame of the Fbxl22 gene (GenBank Accession Number:

MT011389), termed Fbxl22-193 in this study, was amplified using cDNA generated from

RNA isolated from C_2C_{12} cells as described above (Thermo Fisher Scientific, Waltham,

MA). Using the cDNA primers listed in Table 1, the Fbxl22-193 cDNA was amplified and

subcloned into the HindIII and EcoRI sites of pcDNA3.1(+) (ThermoFisher Scientific,

Waltham, MA). A pCMV6-myc-DDK-Fbxl22 full-length cDNA (GenBank Accession

Number: NM_175206) plasmid was purchased from Origene (Rockville, MD). The full-

length Fbxl22 variant was termed Fbxl22-236 in this study. An untagged version of the

full-length mouse Fbxl22 gene was generated by cutting the full-length Fbxl22-myc-DDK

cDNA out of pCMV6 and cloning it into the EcoRI and NotI sites of pcDNA3.1 and then

performing site-directed mutagenesis to introduce a stop codon immediately upstream

of the myc and DDK tags. The untagged full-length Fbxl22 cDNA was then sequenced

to confirm correct orientation, the successful introduction of the correct stop codon, and

the absence of any other mutations (Eurofins Genomics, Louisville, KY).

The Fbxl22 proximal promoter region was amplified from genomic DNA isolated from

C_2C_{12} cells using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA) according to

the manufacturer's protocol. Fbxl22 promoter specific primers were designed using

genomic DNA sequences on deposit with the Ensembl database to amplify

approximately 500 and 1000bp fragments of the proximal 5' regulatory region (Table 1).

Promoter fragments were amplified using 1 µg of genomic DNA and Taq Polymerase

according to the manufacturer's protocol (Thermo Fisher Scientific, Waltham, MA).

Amplified fragments were subcloned into the MluI and EcoRI sites of the pSEAP2-Basic

reporter plasmid (Clontech, Mountain View, CA) and sequenced to confirm correct

orientation (Eurofins Genomics, Louisville, KY).

Site-directed mutagenesis

Site-directed mutagenesis was used to mutagenize conserved putative Ebox sequences

in the Fbxl22 proximal promoter and to introduce a stop codon in the pCMV6-myc-DDK-

Fbxl22 plasmid using the primers listed in Table 1. The QuikChange II Site-Directed

Mutagenesis protocol was performed using 200 ng of plasmid DNA and PfuTurbo DNA

Polymerase according to the manufacturer's protocol (Agilent Technologies, Santa

Clara, CA). Mutagenesis reactions were then digested with Dpn I (New England

Biolabs, Ipswich, MA) and transformed into sub-cloning efficiency competent cells (New

England Biolabs, Ipswich, MA). All constructs were sequenced to confirm successful

generation of the expected mutation (Eurofins Genomics, Louisville, KY).

174   <<INSERT TABLE 1. NEAR HERE>>

Quantitative RT-PCR analysis of Fbxl22 expression in cell culture

Total RNA was isolated from C_2C_{12} cells using the RNeasy minikit (Qiagen, Valencia,

CA) and reverse transcribed into cDNA using 1 μg of RNA and iScript Reverse

Transcriptase (Bio-Rad, Hercules, CA) according to the manufacturer's protocol.

Quantitative polymerase chain reaction was performed with 100 ng of cDNA, 100 nM of

each qPCR primer listed in Table 1, and iTaq Universal SYBR Green Supermix (Bio-

Rad, Hercules, CA) according to the manufacturer's protocol. Relative Fbxl22

expression levels were determined using a Bio-Rad CFX Connect machine and

normalized to GAPDH expression using the $-\Delta\Delta C_t$ method.

Study approval for animal experiments

The mice used in these studies were males from the C57BL/6 strain obtained from

Charles River laboratories at ages 3-4 months and used for experiments within 2 weeks

of their arrival. Male MuRF1 knockout (KO) mice were obtained from a breeding colony

maintained at the University of Iowa at 6 months of age and used for denervation

studies. The generation of mice with a null deletion of MuRF1 has been previously

described (9). Animals were housed in ventilated cages maintained in a room at 21 °C

with 12-h light/ dark cycles and had *ad libitum* access to standard chow (Harlan-Teklad

formula 7913) and water throughout the study. All animal procedures were approved by

the Institutional Animal Care and Use Committee at the University of Iowa.

Denervation Model

Targeted denervation of the lower limb muscles in the right leg was accomplished

through transection of the sciatic nerve as previously described (27). Under isoflurane

anesthesia (2-4% inhalation) and with the use of aseptic surgical techniques, the sciatic

nerve was isolated in the midthigh region and cut with sharp scissors. Mice were given

an analgesic (buprenorphine, 0.1 mg/kg) immediately following the surgery and returned

to their cage following recovery. A group of true control animals were used for the in-

depth time course analysis of muscle mass plus MuRF1 and Fbxl22 isoform mRNA

expression.

In vivo electroporation

Transfection of mouse skeletal muscle with plasmid DNA was performed in mice under

isoflurane (2-4% inhalation) anesthesia as previously described (22, 59). Briefly, after a

2hr pre-treatment with 0.4 units/µl of bovine placental hyaluronidase (Sigma)

resuspended in sterile 0.9% saline, 20µg of plasmid DNA was injected into either the

tibialis anterior (TA) or the lateral and medial gastrocnemius muscles (lateral

gastrocnemius, LGA; medial gastrocnemius, MGA). The hind limbs were placed

between two-paddle electrodes and subjected to 10 pulses (20 msec) of 175 V/cm

using an ECM-830 electroporator (BTX Harvard Apparatus, Holliston, MA). For in vivo

overexpression studies, an additional 2 µg of emerald-GFP plasmid was electroporated

for identification of GFP positive fibers. For RNAi plus denervation experiments,

plasmids were electroporated into hindlimb muscles three days prior to implementation

of sciatic nerve surgery (detailed above). Following transfection, mice were returned to

their cages to resume normal activities until tissue collection.

Tissue Collection

Following completion of the appropriate time period, mice were anesthetized with

isoflurane, and the TA and GA muscles were excised, weighed, frozen in liquid nitrogen,

and stored at –80°C for later analysis. A subset of muscles were collected for histology

and processed as described below. On completion of tissue removal, mice were

euthanized by exsanguination.

Histology

Harvested muscles were immediately fixed in 4% (w/v) paraformaldehyde for 16h at

4°C. Following a sucrose gradient incubation period, the TA muscles were embedded in

Tissue Freezing Medium (Triangle Biomedical Sciences), and a Thermo HM525

cryostat was used to prepare 10-µm serial sections from the muscle midbelly. All

sections were examined and photographed using a Nikon Eclipse Ti automated inverted

microscope equipped with NIS-Elements BR digital imaging software.

Laminin Stain: TA muscle sections were permeabilized in PBS with 1% triton for 10

minutes at room temperature. After washing with PBS, sections were blocked with 5%

goat serum for 15 minutes at room temperature. Sections were incubated with Anti-

Laminin (1:500, Sigma Aldrich, Cat no. L9393) in 5% goat serum for 2 hours at room

temperature, followed by two 5-minute washes with PBS. Goat-anti-rabbit AlexaFluor®

555 secondary (1:333) in 5% goat serum was then added for 1 hour at room

temperature. Slides were cover slipped using ProLong Gold Antifade reagent (Life

Technologies). Image analysis was performed using Myovision software (68). Skeletal

muscle fiber size was analyzed by measuring ≥ 350 transfected muscle fibers per

muscle (GFP-positive), per animal (10x magnification). In Fbxl22 overexpression

studies, we also measured the size of non-transfected fibers (GFP-negative) from the

transfected muscles at all time points. Comparison of the CSA distributions of GFP-

negative fibers in EV, -236 and 193 transfected muscles revealed no difference

between groups. Further in some muscles the transfection frequency was so high that

there were few GFP-negative fibers within the same region as the GFP-positive fibers.

Therefore, for Fbxl22 overexpression studies, fiber size comparisons were made

between the GFP-positive fibers in the EV controls and the -293 and 193 transected

muscles.

Hematoxylin & Eosin stain: A standard H&E protocol was performed (adapted from

Wang et al. (67)). Briefly, TA muscle sections were incubated with hematoxylin solution

for 10 minutes to stain the nuclei and then stained with Eosin solution for 3 min.

Sections were then immersed in 70% ethanol for 20 sec, 90% ethanol for 20 sec, 100%

ethanol for 1 min, and citrisolv for 3 min. Finally, slides were allowed to air dry and a

coverslip was added using Permount media (Thermo Fisher Scientific, Waltham, MA).

Percentage of centrally nucleated myofibers was calculated by dividing the number of

transfected fibers containing ≥1 centrally located nucleus by the total number of

transfected fibers (as previously described (24)). An average of 1500 myofibers were

counted per animal (N=4-5/group), and quantification was performed using ImageJ

software (National Institutes of Health, Bethesda, MD, USA).

RNAi Plasmids

For RNAi experiments, the negative control RNAi plasmid was described previously (22,

59) and encodes emerald green fluorescent protein (EmGFP) and a non-targeting

pre-miRNA under bicistronic control of the cytomegalovirus (CMV) promoter in the

pcDNA6.2GW/EmGFP-miR plasmid (Invitrogen, Carlsbad, CA). The Fbxl22 RNAi

plasmid encodes EmGFP and an artificial pre-miRNA targeting the full length gene of

mouse Fbxl22 under bicistronic control of the CMV promoter; it was generated by

ligating the Mmi576126 oligonucleotide duplex (Invitrogen, Carlsbad, CA) into the

pcDNA6.2GW/EmGFP-miR plasmid.

Immunoblotting

Frozen TA and GA muscles were homogenized in sucrose lysis buffer (50 mM Tris pH

7.5, 250 mM sucrose, 1 mM EDTA, 1 mM EGTA, 1% Triton X 100, 50 mM NaF). The

supernatant was collected following centrifugation at 8,000 g for 10 minutes and protein

concentrations were determined using the 660 protein assay (Thermo Fisher Scientific,

Waltham, MA). Twelve micrograms of protein were subjected to SDS-PAGE on 4-20%

Criterion TGX stain-free gels (Bio-Rad, Hercules, CA) and transferred to polyvinylidene

diflouride membranes (PVDF, Millipore, Burlington, MA). Membranes were blocked in

3% nonfat milk in Tris-buffered saline with 0.1% Tween-20 added for one hour and then

probed with primary antibody overnight at 4°C. Membranes were washed and incubated

with HRP-conjugated secondary antibodies at 1:10,000 for one hour at room

temperature. Immobilon Western Chemiluminescent HRP substrate was then applied to

the membranes prior to image acquisition. Image acquisition and band quantification

were performed using the Azure C400 System (Azure Biosystems, Dublin, CA, USA)

and Image Lab, version 6.0.1 (Bio-Rad), respectively. Total protein loading of the

membranes captured from images using stain-free gel technology was used as the

normalization control for all blots. The following antibodies were used: Ubiquitin (1:1000,

Life Sensors, VU-1), p62 (1:1000, Sigma, P0067), LC3B (1:1000, Sigma, L7543),

Desmin (1:500, Cell Signaling, 5332), Dystrophin (1:1000, Santa Cruz, sc-365954), α-

actinin (1:1000, Cell Signaling, 3134), Vimentin (1:1000, Santa Cruz, sc-373717).

Gene expression by Quantitative RT-PCR in skeletal muscle

Frozen muscle powder was homogenized using RNAzol RT reagent (Sigma-Aldrich, St

Louis, MO) in accordance with the manufacturer's instructions. cDNA was synthesized

using a reverse transcription kit (iScript cDNA synthesis kit; Bio-Rad, Hercules, CA)

from 1 µg of total RNA. PCR reactions (10 µL) were set up as: 2 µL of cDNA, 0.5 µL

(10 µM stock) forward and reverse primer, 5 µL of Power SYBR Green master mix

(Thermo Fisher Scientific, Waltham, MA) and 2 µL of RNA/DNA free water. Gene

expression analysis was then performed by quantitative PCR on a Quantstudio 6 Flex

Real-time PCR System (Applied Biosystems, Foster City, CA) using the mouse primers

shown in Table 1. PCR cycling comprised: hold at 50°C for 5 min, 10 min hold at 95°C,

before 40 PCR cycles of 95°C for 15 s followed by 60°C for 1 min (combined annealing

and extension). Melt curve analysis at the end of the PCR cycling protocol yielded a

single peak. As a result of reference gene instability, gene expression was normalized

to tissue weight and subsequently reported as the fold change relative to control

muscles, as described previously (31). This type of analysis has previously been used

extensively by our group (2, 3, 32).

Statistical Analysis

Results are presented as mean ± standard deviation (SD) for all in vitro experiments

and mean ± standard error of measure (SEM) for all in vivo experiments. Statistical

differences for all in vitro reporter assays were determined using a two-sample t-test

and performed using the data analysis add-in for Excel. Statistical differences for in vivo

overexpression studies were determined using a paired students' t-test. The

denervation time course experiment was assessed for differences in muscle mass and

mRNA expression using a one-way analysis of variance (ANOVA) with Dunnett's

multiple comparisons test. For the RNAi plasmid plus denervation experiments,

differences were determined using an unpaired students' t-test. All in vivo statistical

analyses were performed using GraphPad Prism (GraphPad Software, Inc., La Jolla,

CA). Results were considered significant when $P \leq 0.05$.

Results

Fbxl22 is induced during neurogenic skeletal muscle atrophy

An Illumina beadchip microarray was previously performed on RNA isolated from triceps

surae (TS) muscle of mice undergoing denervation-induced skeletal muscle atrophy and

described in Furlow *et al.* (25). Further analysis of the data SubSeries (GEO:

GSE44205) submitted to the NCBI Gene Expression Omnibus (GEO) revealed

numerous genes that previously have not been described in skeletal muscle, but show

differential expression following sciatic nerve transection. One of these previously

undescribed genes was originally identified from the microarray by its Riken

designation, 1110004B15RIK, and later determined to be Fbxl22, an E3 ubiquitin ligase

known to be enriched in cardiac muscle (18). Based on the microarray data, basal

expression of Fbxl22 is low in skeletal muscle of wild type (WT) and MuRF1 KO mice,

but is significantly elevated after three days of denervation (Fig 1A). Interestingly,

Fbxl22 expression returns to baseline expression by 14 days of denervation, in both WT

and MuRF1 KO skeletal muscle (Fig. 1B).

To confirm the identification of Fbxl22 from our previously published microarray data

(25) and to examine levels of endogenous expression for a novel Fbxl22 splice variant

(identification detailed below), we performed an in depth time course of denervation-

induced muscle atrophy. For the gastrocnemius complex (lateral and medial

gastrocnemius, plantaris, and soleus muscles), we observed significant reductions in

muscle mass at 7, 14 and 21 days post sciatic nerve transection (Fig. 1C). A significant

increase in MuRF1 mRNA expression occurred at 3 and 7 days post-denervation (Fig.

1D). Interestingly, Fbxl22 (both full length, Fbxl22-236 and novel splice variant, Fbxl22-

193) mRNA expression was significantly increased at 3 days post-denervation, but was

back to baseline levels by 7 days (Fig. 1E and 1F).

<<INSERT FIGURE 1. NEAR HERE>>

Identification and cloning of a novel Fbxl22 splice variant from skeletal muscle cells

Fbxl22 is located on chromosome 9 in mouse and encodes two alternative transcripts

that consist of either 2 coding exons (Fbxl22-Full Length) or 3 coding exons (Fbxl22-

Novel Transcript) (Fig. 2A). To demonstrate that this gene is expressed endogenously

at the transcriptional level in muscle cells, a cDNA comprising the open reading frame

(ORF) of Fbxl22 was amplified from C_2C_{12} myoblasts. The amplified cDNA was

sequenced in both directions and found to contain an in frame 129 nucleotide deletion in

the second exon based on comparisons to the predicted sequence available from the

Ensembl database (Fig. 2A). This deletion results in the removal of 43 amino acids in

the leucine-rich repeat region of Fbxl22 and essentially removes one leucine-rich

repeat. To further assess the endogenous Fbxl22 expression profile in muscle cells,

real-time RT-qPCR was performed using RNA isolated from proliferating myoblasts and

myotubes at early and late stages of differentiation. The results demonstrate that

Fbxl22-193 (novel) and Fbxl22-236 (full length) expression levels are highest in fully

differentiated myotubes (DD7) compared to proliferating myoblasts (PD2) (Fig. 2B and

2C). To determine if the Fbxl22-193 splice variant can be generated from the full-length

Fbxl22 transcript, C_2C_{12} cells were transfected with an expression plasmid containing

the Fbxl22-236 cDNA and quantitative RT-qPCR was performed using primers specific

for Fbxl22-236 or Fbxl22-193. The results show that overexpression of Fbxl22-236 in

muscle cells leads to significantly elevated levels of both the full-length Fbxl22 transcript

and the novel Fbxl22-193 splice variant (Fig. 2D and 2E). Furthermore, this observation

was confirmed in vivo with Fbxl22-236 overexpression in mouse skeletal TA muscle

leading to significantly increased levels of the novel Fbxl22-193 splice variant (Figure

S1).

373   <<INSERT FIGURE 2. NEAR HERE>>

Fbxl22 regulatory region is transcriptionally active in muscle cells and is induced by

myogenic regulatory factors

The predicted regulatory region of Fbxl22 was cloned using the primer pairs listed in

Table 1. The isolated promoter fragments spanned from +40 to -457 and +40 to -967 of

the mouse Fbxl22 gene (Fig. 3A). These fragments were fused with the secreted

alkaline phosphatase (SEAP) reporter gene to create the pSEAP-Fbxl22-Pro500 and

pSEAP-Fbxl22-Pro1000 reporter plasmids. C_2C_{12} mouse myoblast cells were transiently

transfected with the Fbxl22 reporter constructs and SEAP activity levels were analyzed

5-7 days following the switch to differentiation media and compared to SEAP activity

levels from cells transfected with the empty pSEAP2-Basic plasmid. The Fbxl22 reporter

plasmids showed significant transcriptional activity and had significantly higher

expression compared to the empty reporter plasmid alone (Fig. 3A). To determine if the

Fbxl22 reporter gene activity mirrors the transcriptional activity of the endogenous gene

(Fig. 2B and 2C), the pSEAP-Fbxl22-Pro500 reporter was transfected into C_2C_{12}

myoblasts and media was then analyzed for SEAP activity at 24 hours post-transfection

(PD2) and at 2 (DD2) and 7 (DD7) days post-switch to differentiation media. The data

shows that the Fbxl22 reporter gene has elevated transcriptional activity as C_2C_{12}

differentiate (Fig. 3B), mirroring the qPCR data.

Neurogenic atrophy results in a dramatic increase in myogenic regulatory factor (MRF)

expression, including significant increases in MyoD1 and myogenin expression (18, 25,

44, 48, 63). Therefore, since MyoD1 and myogenin are elevated during neurogenic

atrophy and are capable of binding canonical E-box elements in muscle-specific genes,

we evaluated the ability of the Fbxl22 reporters to respond to ectopic expression of

MyoD1 and myogenin in C_2C_{12} cells transiently transfected with the Fbxl22-Pro500 wild-

type reporter constructs. Cells transiently transfected with MyoD1 or myogenin in

combination with Tcf3 or Tcf12 expression plasmids showed significantly higher Fbxl22

reporter gene activity compared to cells that were not (Fig. 3C and 3D).

Identification of conserved, putative E-box elements in the Fbxl22 regulatory region

To better assess the level of conservation between the regulatory regions of mouse, rat,

and human Fbxl22 and to identify potential transcriptional regulatory elements, the

genomic sequences corresponding to the regulatory region of this gene for mouse, rat,

and human were aligned (Fig. 3E). Several putative enhancer elements were identified

in the proximal promoter region of Fbxl22, including one putative AP-2 consensus

sequence (37), one putative CCAAT box (43), and one putative FoxO binding element

(56, 66) (Fig. 3E). Furthermore, since MRFs are elevated during neurogenic atrophy,

are capable of binding E-box elements in muscle-specific genes, and appear to

positively regulate Fbxl22 reporter gene activity, we analyzed the Fbxl22 regulatory

region and identified two putative E-box elements located just upstream of the start of

transcription of the Fbxl22 gene. These two putative E-box elements lie between -145

and -70 of the proximal promoter region of Fbxl22 (Fig. 3E). To determine if the E-box

elements are necessary for proper Fbxl22 promoter activity, site-directed mutagenesis

was performed using the -500 bp Fbxl22 reporter plasmid, resulting in the generation of

the pSEAP-Fbxl22-Pro500 E-box2, pSEAP-Fbxl22-Pro500 E-box3, and the pSEAP-

Fbxl22-Pro500 E-box2/3 mutant constructs. Mutation of either E-box in the 500 bp

Fbxl22 promoter construct resulted in significantly lower reporter gene activity compared

to the wild-type reporter construct in transiently transfected C_2C_{12} cells (Fig. 3F),

suggesting that MRF activation of Fbxl22 expression may occur via these proximal E-

box elements.

424   <<INSERT FIGURE 3. NEAR HERE>>

Overexpression of the Fbxl22 isoforms in unchallenged TA muscles for 7, 14 and 28

days post electroporation

We sought to manipulate gene expression levels of both Fbxl22 isoforms (-236 and -

193) though an in vivo electroporation approach into unchallenged TA muscles for

periods of 7,14, and 28 days. Transfection of the full-length Fbxl22-236 construct in the

TA resulted in no changes in muscle mass at 7 and 28 days, but an increase after 14

days post electroporation (P = 0.01; Fig. 4C). Measurement of muscle fiber CSA

showed no differences between Fbxl22-236 transfected muscles and EV control

muscles after 7 days. A shift in the distribution of fiber CSAs towards smaller muscle

fibers was observed in Fbxl22-236 transfected muscles after 14 days but was no longer

evident after 28 days (Fig. 4D). On the other hand, transfection of the novel Fbxl22-193

isoform in the TA lead to a significant increase in muscle mass after 7 days, but no

differences in muscle mass at 14 or 28 days (Fig. 4E). In terms of muscle CSA, there

were increases in the percentage of smaller and larger fibers with the presence of

Fbxl22-193 overexpression compared to EV control muscles at 14 and 28 days (Fig.

4F).

At 14 days, Fbxl22-236 and Fbxl22-193 transfected muscles displayed increased cell

infiltration, necrosis and degeneration, and the large fibers were rounded in appearance

which is abnormal (Fig. 5A). With prolonged Fbxl22-193 overexpression for 28 days, we

observed large fibers with numerous centrally located nuclei and reduced amounts of

cell infiltration (Fig. 5A) suggestive of regeneration. Overexpression of -236 and -193

isoforms resulted in a significant amount of transfected fibers with centrally located

nuclei after 14 and 28 days (Fig 5B and 5C).

At the biochemical level, there was no change in total ubiquitin or p62 levels with

Fbxl22-236 overexpression at 14 days, but there was an increase in LC3B II content

(Fig. 6B). Further, a reduction in dystrophin protein and significant elevations in desmin

and vimentin were observed (Fig. 6C). Overexpression of Fbxl22-236 for 7 days

resulted in alterations in total ubiquitin, p62 and LC3B II protein content, but no

significant changes in dystrophin and intermediate filament protein levels (Fig. S2). The

only significant differences observed in sarcomeric protein levels after 28 days of

Fbxl22-236 overexpression was a decrease in α-actinin and an increase in vimentin

(Fig. S2). In comparison, there were significant increases in total ubiquitin, p62 and

LC3B II content in Fbxl22-193 transfected muscles compared to the EV control group at

14 days (Fig. 6E). In addition, we observed a significant reduction in dystrophin protein

and changes in α-actinin isoform levels (Fig. 6F). The alterations in α-actinin isoforms

were accompanied by 3- to 4-fold increases in desmin and vimentin (Fig. 6F). After 7

days of electroporation, Fbxl22-193 transfected muscles had elevation in p62 levels,

reduction in dystrophin, and alterations in α-actinin isoforms and vimentin (Fig. S3). At

28 days, total ubiquitin and LC3B II content remained elevated (1.5-fold), while no

significant changes in cytoskeleton protein content were observed (Fig. S3). The overall

changes in key sarcomeric proteins may be indicative of destabilization in the

myofilament lattice and the appearance of regenerating muscle fibers (13, 26, 34, 41).

469   <<INSERT FIGURE 4. NEAR HERE>>

470   <<INSERT FIGURE 5. NEAR HERE>>

471   <<INSERT FIGURE 6. NEAR HERE>>

Overexpression of Fbxl22 isoforms in unchallenged lateral and medial gastrocnemius

muscles for 14 days

The H&E staining of both GA muscles showed a similar pattern to the TA muscle after

14 days of overexpression, where GA muscles transfected with either Fbxl22 isoform

displayed increased cell infiltration, necrosis and degeneration of skeletal muscle fibers

(Fig. 7A and 7B). In the lateral GA muscle, the overexpression of Fbxl22-236 led to a

significant increase in muscle mass (Fig. 7C), while overexpression of the Fbxl22-193

isoform caused no change in muscle mass compared to EV transfected muscles (Fig.

7D). In Fbxl22-236 transfected muscles, there was a shift in the distribution of fiber CSA

primarily towards smaller fibers compared to the EV control muscle (Fig. 7C). Enlarged

fibers could be found in the Fbxl22-236 transfected muscles, however, they were

usually abnormally rounded in shape. The Fbxl22-193 transfected muscle displayed a

slight shift towards an increase in smaller fibers compared to the contralateral EV

control group (Fig. 7D). In the medial GA muscle, Fbxl22-236 transfected muscles

displayed a significant reduction in muscle mass and a leftward shift towards smaller

muscle fibers compared to the EV transfected muscles (Fig. 7E). The overexpression of

Fbxl22-193 isoform in the medial GA muscle caused no change in muscle mass but did

produce a shift in the distribution of fiber CSA primarily towards an increase in the

number of smaller muscle fibers (Fig. 7F). In addition, we observed a significant

increase in fibers with centrally located nuclei for -193 transfected GA muscles after 14

days (Fig 7D and 7F).

The histological changes observed with Fbxl22 overexpression in the GA muscle

corresponded with increased levels of markers of protein degradation (total ubiquitin,

p62 and LC3B II) and alterations in cytoskeleton proteins (dystrophin, α-actinin, desmin,

and vimentin) compared to EV transfected muscle (Fig. S4 and S5). These observations

were present in both the lateral and medial GA muscles after overexpression of the full-

length and novel Fbxl22 isoforms for 14 days. These data highlight the variation in the

morphological and biochemical response of different skeletal muscles (TA versus

gastrocnemius), depending on which Fbxl22 isoform is overexpressed.

503   <<INSERT FIGURE 7. NEAR HERE>>

Knockdown of Fbxl22 leads to partial muscle sparing in wild type muscle during denervation-induced muscle atrophy

Given the early upregulation of Fbxl22 expression following denervation, we studied whether knock down of Fbxl22 in muscle would have an effect on muscle atrophy following denervation. We first confirmed that our Fbxl22 RNAi construct could knock down Fbxl22 mRNA expression after 3 days of denervation. The Fbxl22 RNAi transfected muscles displayed a 60-70% reduction in endogenous expression of Fbxl22 mRNA (for both isoforms) in the LGA and MGA muscles compared to the EV transfected muscles (Fig. S6). After 7 days of denervation, total GA mass in WT mice significantly decreased in EV transfected and Fbxl22 RNAi transfected muscles, with no significant difference between groups (Fig. 8A). In relation to the contralateral control muscle, EV transfected muscles exhibited a 31% loss in muscle mass, whereas Fbxl22 RNAi transfected muscles displayed a 27% loss (P = 0.054; Fig. 8B). When we examined the two heads of GA separately, we found that in the LGA muscle there was no muscle sparing in the Fbxl22 RNAi transfected group (Fig. 8C), but in the MGA there was significant muscle sparing in the presence of the Fbxl22 RNAi (P = 0.03; Fig. 8D). The mean fiber CSA for GFP-positive fibers was significantly larger in the RNAi transfected muscles compared to the EV group (P = 0.04; Fig. 8E). Measurement of fiber CSA in both heads of the GA revealed that Fbxl22 RNAi resulted in partial preservation of muscle CSA size compared to non-transfected fibers, as signified by a reduced leftward shift in distribution towards smaller muscle fibers (Fig. 8F). In

526   comparison, the EV displayed a leftward shift towards smaller muscle fibers that was

527   similar to the non-transfected fibers (Fig. 8G).

529   <<INSERT FIGURE 8. NEAR HERE>>

Knockdown of Fbxl22 in MuRF1 knockout muscle results in greater muscle sparing

under denervation-induced muscle atrophy

As muscle sparing is evident in the MuRF1 KO mice at 14 days, but not 7 days (9), we

sought to determine the effect of knockdown of Fbxl22 expression in MuRF1 KO

muscles. After 7 days of denervation in MuRF1 KO mice, total GA mass was

significantly smaller in the EV transfected muscles compared to the Fbxl22 RNAi

transfected muscle (P = 0.03; Fig. 9A). MuRF1 KO muscles transfected with the EV

exhibited a 27% loss in muscle mass compared to the contralateral control (Fig. 9B),

while MuRF1 KO muscles transfected with the Fbxl22 RNAi construct showed

significant muscle sparing with only a 19% loss in muscle mass compared to the

contralateral control muscle (P = 0.01; Fig. 9B). Examination of the two heads

separately showed that in the LGA muscle there was partial (albeit non-significant)

sparing in the Fbxl22 RNAi transfected group (Fig. 9C) and significant muscle sparing in

the MGA muscle with the presence of the Fbxl22 RNAi (P = 0.01; Fig. 9D). The mean

fiber CSA for GFP-positive fibers was significantly larger in the RNAi transfected

muscles compared to the EV group (P = 0.01; Fig. 9E). Analysis of muscle fiber CSA

revealed Fbxl22 RNAi resulted in preservation of muscle CSA size compared to non-

transfected fibers, as signified by a lack of a leftward shift in distribution towards smaller

muscle fibers (Fig. 9F). In comparison, the EV transfected fibers displayed a leftward

shift towards smaller CSAs that was similar to the non-transfected fibers (Fig. 9G).

551   <<INSERT FIGURE 9. NEAR HERE>>

Discussion

Skeletal muscle has the ability to regulate its size in response to internal and external

cues (such as neural activity, nutritional status, etc.) throughout the lifespan (60).

Homeostasis of skeletal muscle size occurs through a balance between protein

synthesis and breakdown, with the balance shifting to net protein breakdown during

skeletal muscle atrophy events (7, 47, 58). The UPS is an integral part of the protein

breakdown process in skeletal muscle and E3 ubiquitin ligases are critical in targeting

substrates for ubiquitination (8, 55). Surprisingly, to date only a handful of E3 ubiquitin

ligases have been identified and characterized as important in the process of skeletal

muscle atrophy (9, 28, 45, 51, 57). In our present study, we have identified Fbxl22 and a

novel splice variant that code for E3 ubiquitin ligases that are significantly upregulated in

skeletal muscle exposed to neural inactivity. From our previously published microarray

data (25), we found the RIKEN gene 1110004B15RIK to be at the top of the list of

genes upregulated after 3 days of denervation in wild type and MuRF1 KO muscle. We

subsequently identified this RIKEN gene to be Fbxl22. Through an in-depth time course

analysis of denervation, Fbxl22 isoforms were found to significantly increase early in the

atrophy process and return to baseline levels by 7 days. In comparison, MuRF1 and

MAFbx expression remain elevated for a longer time period in neurogenic muscle

atrophy models, returning to baseline after 14 days (9, 25, 38, 48, 59). Thus, we

hypothesized that Fbxl22 could be involved in the early initiation of the muscle atrophy

process.

Fbxl22 is a member of the F-box family (which includes MAFbx, MUSA1, and SMART

(9, 45, 57)) and has previously been observed to play a role in sarcomeric turnover in

cardiac muscle (62). To our knowledge, the study by Spaich and colleagues (62) is the

only other published investigation into the role of Fbxl22 in striated muscle. In the study

by Spaich et al, the authors observed localization of Fbxl22 at the sarcomeric z-line and

in vitro overexpression studies identified filamin C and α-actinin to be Fbxl22 targeted

substrates for ubiquitination. Through analysis of Fbxl22 overexpression at multiple time

points and in multiple muscles, we observed significant alterations in α-actinin isoforms,

desmin, and vimentin which are all integral for maintaining muscle fiber integrity and

function (26, 34, 40, 50). Specifically, both Fbxl22 isoforms cause a reduction in one of

the α-actinin isoforms and significant elevations in desmin and vimentin. Interestingly,

expression of desmin and vimentin is suggested to be indicative of regenerating muscle

fibers (13, 26, 65). It is important to note that desmin has been observed to be targeted

for degradation in muscle atrophy models, specifically by the E3 ubiquitin ligase TRIM32

(17). We also observed significant reductions in dystrophin protein content in Fbxl22

transfected muscles. Dystrophin is essential for lateral force transmission and protecting

skeletal muscle from contraction-induced injury (32, 33, 41, 54). The alterations in these

sarcomeric proteins coincided with the development of a degenerative muscle fiber

phenotype in Fbxl22 transfected tissues. Other E3 ubiquitin ligases, such as MuRF1,

have been reported to target a distinct subset of sarcomeric proteins (such as titin and

myosin heavy chain) for degradation in skeletal muscle (14, 15, 17, 53), suggesting that

the protein turnover that accompanies muscle atrophy is likely orchestrated by several

different E3 ligases.

Upon neurogenic atrophy, there is a dramatic increase in MRF expression including

MyoD1 and myogenin (18, 44, 48). Within the Fbxl22 promoter, we identified two

putative E-box elements immediately upstream of the start of transcription, of which one

is fully conserved between rodents and humans. Furthermore, Fbxl22 reporter gene

activity appears to be positively regulated by MRF expression, and thus highlights the

potential role for Fbxl22 in the early initiation of the muscle atrophy process.

Interestingly, the development of a degenerative fiber phenotype in our overexpression

studies took a progressive period of time to develop (up to 14 days). The prolonged

expression of Fbxl22 alone may explain the appearance of the observed myopathy,

since Fbxl22 induction appears to be short-lived and returns rapidly to baseline after just

3 days of denervation. Furthermore, it is also possible that other genes that are

differentially expressed in response to denervation may work to direct the activity of

Fbxl22 during skeletal muscle atrophy and prevent the development of myopathy (25).

Although, we observed no dramatic effect on muscle mass with Fbxl22 isoform

overexpression, except in the MGA muscle, we did observe a distinct phenotype in

muscle fiber CSA with the appearance of small degenerating fibers and large abnormal

fibers which may have masked the lack of effect on overall muscle mass. In the MGA

muscle, the fiber CSA distribution shifted towards the left due to an increase in smaller

fibers with no large abnormal fibers present, which coincided with a significant loss of

muscle mass with Fbxl22 overexpression. The lack of uniform response on muscle

mass in Fbxl22 isoform overexpression studies highlights the complexity of studying a

single component of the muscle atrophy program in multiple hindlimb muscles (3, 6, 21).

The manifestation of the degenerative phenotype in Fbxl22 transfected muscles

appears to differ between muscles and is dependent on which Fbxl22 isoform is

overexpressed, this being reflected in the subtle differences observed in our

biochemical analyses. We have previously observed muscle-specific changes in levels

of atrophy upon implementation of a hindlimb unloading model (2, 3, 46). While

differences in muscle fiber type, neural input, and/or loading history may all contribute to

our previous findings, it is important to highlight that the changes in E3 ubiquitin ligase

expression (MuRF1 and MAFbx) varied between the rat soleus, medial gastrocnemius

and TA muscles during the unloading period (3). It would appear that different muscles

may atrophy in different ways at the molecular level depending on which genes are and

are not expressed (1, 3, 25). To note, we only implemented a neural inactivity stimulus

for inducing atrophy and thus future studies are needed to explore Fbxl22 expression in

other atrophy models where the motor unit remains intact (e.g. hindlimb suspension,

glucocorticoids). It is possible that the action of Fbxl22 may differ between hindlimb

muscles and this may explain why we observed muscle atrophy in MGA transfected

muscles, but a degenerative myopathy in LGA and TA transfected muscles. In addition,

differences in the ratio of E1 and E2 enzymes present in each manipulated muscle (TA,

LGA and MGA) and/or differences in Fbxl22 E2 binding partners could influence the

muscles' response to Fbxl22 overexpression. Furthermore, we observed different levels

of total ubiquitin between Fbxl22 isoforms and the muscle transfected, which may

indicate different substrates being targeted for degradation. Future studies will seek to

identify which E2 enzymes pair with Fbxl22 and the specific substrates that Fbxl22

targets for ubiquitination.

In the reported literature, Fbxl22 has been indirectly assessed in skeletal muscle

through transcriptomic analysis and genome-wide screens in various models. In human

skeletal muscle exposed to either bed rest or unilateral lower limb immobilization,

Fbxl22 does not appear to be transcriptionally active in vastus lateralis biopsy samples

between 2 and 14 days of intervention (52). Interestingly, Fbxl22 has been reported in

C_2C_{12} muscle cells to inhibit SMAD4 translocation and appears to modulate

transforming-growth factor-β1 (TGFβ1) signaling (20). To note, our previously published

microarray data and other studies have independently observed alterations in the TGF-

β signaling pathway in denervated muscle tissues (10, 25). As the network of TGF-β

ligands have been observed to be negative regulators of muscle mass (11, 12), future

studies may seek to explore the possible molecular interactions between Fbxl22 and

TGF-β signaling in muscle mass regulation.

Multiple E3 ubiquitin ligases have been targeted individually for knock down or knock

out in various skeletal muscle atrophy models, often resulting in sparing of muscle mass

(9, 28, 45, 51, 57). We report that RNAi knockdown of Fbxl22 mRNA expression leads

to partial muscle sparing in wild type muscle undergoing denervation-induced atrophy.

However, knock down of Fbxl22 in muscle with a deletion of MuRF1 resulted in

significantly greater sparing of muscle mass following 7 days of denervation. This result

is particularly interesting since the deletion of MuRF1 alone does not lead to sparing of

muscle mass at 7 days of denervation. Interestingly, MuRF1 KO muscles display

increased Fbxl22 expression after 3 days of denervation, and thus Fbxl22 could be an

important factor driving the onset of the early atrophy phase, prior to when the deletion

of MuRF1 results in muscle sparing (9, 25). In both wild type and MuRF1 KO muscle,

the medial gastrocnemius muscle displayed significant muscle sparing with the

presence of the Fbxl22 RNAi vector compared to the lateral muscle after 7 days of

denervation. In addition, the medial gastrocnemius muscle was the only muscle to

display muscle atrophy with Fbxl22 overexpression. MuscleDB database analysis

shows varying basal mRNA expression levels of Fbxl22 (64) in various hindlimb

muscles (plantaris, TA, gastrocnemius, etc.), which may explain the differing degrees of

sparing observed during neurogenic muscle atrophy. Another possible explanation is

the difference in transfection efficiency for the plasmid vectors between the lateral and

medial gastrocnemius, which differ in size and fiber type (61). Indeed, we observed

significant effects of muscle sparing with the Fbxl22 RNAi when single transfected

muscle fibers were analyzed for CSA verses whole muscle mass measures which

includes non-transfected fibers. The development of a muscle specific Fbxl22 knockout

animal or use of an adeno-associated viral vector (AAV) may be advantageous for

future studies (19, 49). Finally, subsequent studies are required to explore the possible

mechanisms for the strong effect in muscle sparing with combined inhibition of Fbxl22

and MuRF1 upon neurogenic muscle atrophy.

Conclusions

In the present study we extend our understanding of Fbxl22 by identifying a novel splice

variant of the gene, by demonstrating that in vivo overexpression causes muscle

atrophy and possibly a myopathy after 14 days, and that knockdown of Fbxl22 leads to

muscle sparing in a neurogenic muscle atrophy model. Fbxl22 overexpression resulted

in increased protein degradation and alterations in key sarcomeric proteins that

maintain muscle fiber structure and function. Also, by targeting two E3 ubiquitin ligases

(Fbxl22 and MuRF1) simultaneously, we observed enhanced muscle sparing after only

7 days of exposure to an atrophy stimulus. Future studies will seek to investigate the

potential role of Fbxl22 in various other atrophy stimuli such as disuse, nutritional

deprivation and glucocorticoid treatment.

Data Availability

693 The data that support the findings of this study are available from the corresponding

694 author upon reasonable request.

Online supplementary material

696 Additional supporting information may be found online via the link for Figshare data:

697 https://figshare.com/s/4bae1a93d95da33645d3.

Author Contributions

699 Conception and design of the experiments: DCH, LMB, DSW and SCB. Collection,

700 analysis and interpretation of data: DCH, LMB, JRD, SAL, DSW and SCB. Drafting the

701 article or revising it critically for important intellectual content: DCH, LMB, DSW and

702 SCB. All authors read and approved the final version of the manuscript and all authors

703 listed qualify for authorship.

Conflict of Interest

705   SCB is on the scientific advisory board for Emmyon Inc. All other authors declare that

706   they do not have a conflict of interest.

Figure Legends

Figure 1 Fbxl22 is induced during neurogenic skeletal muscle atrophy. Whole genome expression analysis was conducted on triceps surae muscle from wild-type (WT) and MuRF1-null (KO) mice after **(A)** 3 days and **(B)** 14 days of denervation (DEN). Fbxl22 expression increased significantly at 3 days of denervation but returned to basal expression levels by 14 days post-denervation. Each condition represents the average expression from three animals (n = 3) and error bars represent ± SD. Gray bars, controls; black bars, DEN. Significant difference between denervated mice and control mice in the same group, (*: $P \leq 0.05$, **: $P \leq 0.01$). Data in Panel A and B from a previously published study (25). (**C**) Muscle mass (milligrams) of Gastrocnemius Complex (GSTC) muscle exposed to denervation (DEN) for 3, 7, 14 and 21 days. The loss of mass as a percentage of control muscles is also given for each time point. (**D**) MuRF1, **(E)** Fbxl22-236 and **(F)** Fbxl22-193 mRNA expression during denervation time course in GSTC muscle. Data presented as Mean ± SEM (N= 5-6/group). Φ and Ψ depict $P \leq 0.05$ verses control group.

Figure 2 Fbxl22-236 and Fbxl22-193 are expressed in skeletal muscle and are upregulated during muscle cell differentiation. (**A**) Schematic of the full-length (Fbxl22-236) and novel splice variant (Fbxl22-193) transcripts in mouse. Darkened rectangles represent exons containing translated region, open rectangles represent exons containing the untranslated regions and the lines connecting the rectangles represent introns. (**B**) Fbxl22-193 and (**C**) Fbxl22-236 expression increases as C_2C_{12} cells differentiate. qPCR analysis of Fbxl22 expression in proliferating (PD2) C_2C_{12}

myoblasts and myotubes at early (DD2) and late (DD7) differentiation. RNA was

isolated from three biological replicates (n = 3) for each time point and the experiment

was repeated at least three times. Significant difference between proliferating and

differentiated cells, (*: $P \leq 0.01$). Ectopic expression of Fbxl22-236 cDNA in C_2C_{12} cells

results in increased levels of both the (**D**) Fbxl22-236 and (**E**) Fbxl2-193 transcripts.

Significant difference between controls cells and cells transfected with the Fbxl22-236

expression plasmid, (*: $P \leq 0.01$).

Figure 3 Fbxl22 reporter gene is transcriptionally upregulated during muscle cell

differentiation and is activated by myogenic regulatory factors. (**A**) C_2C_{12}

myoblasts were transfected with either an empty reporter plasmid or a reporter plasmid

containing ~500 bp or ~1000 bp of the Fbxl22 promoter fused to the Secreted Alkaline

Phosphatase (SEAP) reporter gene. Significant differences in activity between the

control empty reporter plasmid (pSEAP2-Basic) and the pSEAP-Fbxl22-Pro500 and

pSEAP-Fbxl22-Pro1000 reporter constructs (*: $P \leq 0.01$). C_2C_{12} myoblasts were

transfected with the (**B**) pSEAP-Fbxl22-Pro500 and cells were differentiated for seven

days and the media was analyzed at proliferation day 2 (PD2), differentiation day 2

(DD2) and differentiation day 7 (DD7) and SEAP numbers were normalized to β-

galactosidase activity to correct for variations in transfection efficiency. Significant

differences between the activities of the pSEAP-Fbxl22-Pro500 reporter construct at

PD2 compared to DD2 and DD7 (*: $P < 0.01$). C_2C_{12} myoblasts were transfected with

the pSEAP-Fbxl22-Pro500 reporter plasmid alone or in combination with (**C**) MyoD or

(**D**) myogenin with or without Tcf3 or Tcf12 expression plasmids. The media was then

assayed for SEAP activity at 48 hours following the switch to differentiation media.

Significant differences between the activities of the pSEAP-Fbxl22-Pro500 alone and in

combination with (**C**) MyoD alone or MyoD±Tcf3 or MyoD±Tcf12 or (**D**) myogenin alone

or myogenin±Tcf3 or myogenin±Tcf12 overexpression are indicated (*: P < 0.01). (**E**)

Analysis of the Fbxl22 promoter region revealed three conserved, putative consensus

E-box elements 5'-CANNTG-3' (red diamonds). N represents any nucleotide. (**F**) C_2C_{12}

myoblasts were transfected with the pSEAP-Fbxl22-Pro500-wild-type, pSEAP-Fbxl22-

Pro500-E-box2-mutant,   pSEAP-Fbxl22-Pro500-E-box3-mutant,   or   pSEAP-Fbxl22-

Pro500-E-box2/3-mutant reporter constructs with MyoD and Tcf3 expression plasmids.

The media was then assayed for SEAP activity at 48 hours following the switch to

differentiation media. Significant differences between the activities of the pSEAP-

Fbxl22-Pro500-wild-type   construct   and   the   pSEAP-Fbxl22-Pro500-E-box-mutant

reporter constructs (*: P < 0.01). For all Fbxl22 reporter gene experiments, each

condition was performed in triplicate and each experiment was repeated at least three

times. The graphs are of representative experiments (n = 3) and values correspond to

the mean relative light unit (RLU) values ± SD.

Figure 4 Overexpression of Fbxl22 isoforms in unchallenged tibialis anterior (TA)

muscle for 7, 14 and 28 days.   (**A**) Representative microscope images for GFP

fluorescence and laminin in TA mouse muscles transfected with empty vector (EV),

Fbxl22-236 and -193 plasmids (10x magnification; scale bar = 100μm) highlighting

transfection efficiency and muscle morphology after 7 (left), 14 (middle) and 28 (right)

days. Muscle mass measurements (milligrams) for muscles transfected with either the

EV or Fbxl22-236 (**C**) and -193 (**E**) plasmids (n=10/group) after 7 (left), 14 (middle) and

28 (right) days. Muscle cross-sectional area (CSA) distribution for EV and Fbxl22-236

(**D**) and -193 (**F**) transfected muscles (7 days = left; 14 days = middle; 28 days = right).

GFP-positive fibers were measured for CSA, with ≥ 350 transfected fibers analyzed per

animal, per muscle (n= 6/group) and data presented as a percentage of sizes between

0-5000µm. For 14- and 28-day muscle CSA data, fibers that were 5000µm-plus were

included in the analysis due to the phenotype observed with Fbxl22-236 and -193

overexpression at these time points. Data presented as Mean ± SEM.

Figure 5 Quantification of Centrally located nuclei in tibialis anterior (TA) muscle

fibers with Fbxl22 isoform overexpression for 7, 14 and 28 days. Representative

microscope images for hematoxylin and eosin (H & E) in TA mouse muscles transfected

with empty vector (EV), Fbxl22-236 and -193 plasmids (10x and 20x magnification;

scale bar = 100µm) highlighting muscle morphology after 7 (left), 14 (middle) and 28

(right) days. Examples of centrally located nuclei are identified on H & E images with

yellow arrows. Fibers with centrally located nuclei with quantified for -236 (B) and -193

(C) transfected TA muscles for 7 (left), 14 (middle) and 28 (right) days (n = 5/group).

Percentage of centrally nucleated myofibers was calculated by dividing the number of

transfected fibers containing ≥1 centrally located nuclei by the total number of

transfected fibers. Statistical significance is depicted where present. Data presented as

Mean ± SEM.

Figure 6 Alterations in ubiquitin, autophagy markers and cytoskeleton protein

levels in Fbxl22 transfected tibialis anterior (TA) muscles after 14 days. (**A and D**)

Representative immunoblot images for empty vector (EV), Fbxl22-236 and -193

transfected muscles after 14 days (n=4/group). (**B and E**) Total protein levels for

ubiquitin, p62 and LC3B II. (**C and F**) Total protein levels for dystrophin, α-actinin (lower

band), desmin, and vimentin. Total protein loading was used as the normalization

control for all blots. Statistical significance is depicted where present. Data presented as

Mean ± SEM.

Figure 7 Overexpression of Fbxl22 isoforms in unchallenged lateral and medial

gastrocnemius muscles for 14 days. Representative microscope images for GFP

fluorescence, laminin and H& E in lateral (LGA; Panel **A**) and medial (MGA; Panel **B**)

gastrocnemius mouse muscles transfected with empty vector (EV) or Fblx22-236 and

Fbxl22-193 constructs (i, 10x magnification and ii, 20x magnification; scale bar =

100µm) highlighting transfection efficiency and muscle morphology. Examples of

centrally located nuclei are depicted on H & E images with yellow arrows. Muscle mass

measurements (milligrams) for LGA (Panel **C** and **D**) and MGA (Panel **E** and **F**) muscles

transfected with either the EV, Fbxl22-236 and Fbxl22-193 constructs (n=10/group).

Muscle cross-sectional area (CSA) distribution in LGA (Panel **C** and **D**) and MGA (Panel

E and **F**) transfected muscles with either the EV, Fbxl22-236 or Fbxl22-193 constructs.

GFP-positive fibers were measured for CSA, with ≥ 350 transfected fibers analyzed per

animal, per muscle (n= 5/group) and data presented as a percentage of sizes between

0-5000µm. Fibers with centrally located nuclei (≥1) quantified in LGA (Panel **C** and **D**)

and MGA (Panel **E** and **F**) transfected muscles with either the EV, Fbxl22-236 and

Fbxl22-193 constructs (n=4/group). Statistical significance is depicted where present.

Data presented as Mean ± SEM.

Figure 8 Fbxl22 knockdown leads to partial muscle sparing after 7 days of

denervation in wild type mice. (**A**) Total muscle mass for gastrocnemius (GA; lateral

and medial) muscles transfected with either empty vector (EV) or Fblx22 RNAi

constructs. (**B**) Loss of GA muscle mass with either EV or Fbxl22 RNAi present

compared to contralateral control (innervated) muscle. Data expressed as percent lost

compared to the contralateral control. Muscle mass for **(C)** lateral and **(D)** medial

gastrocnemius muscles transfected with EV or Fbxl22 RNAi construct after 7 days of

denervation. (**E**) Mean Fiber cross-sectional area (CSA) for GFP-positive only fibers in

transfected GA muscles was determined.   Muscle cross-sectional area (CSA)

distribution in transfected muscles with the **(F)** Fbxl22 RNAi or **(G)** EV constructs. GFP-

positive fibers were measured for CSA, with ≥ 350 transfected fibers analyzed per

animal, per muscle (n= 5/group). In addition, the muscle CSA distributions for non-

transfected (red bars) and innervated (blue triangle) contralateral control muscles were

measured. Data presented as a percentage of sizes between 0-5000µm. (**H**)

Representative images for GFP fluorescence and laminin stain in Fbxl22 RNAi and EV

transfected lateral and medial gastrocnemius muscles after 7 days of denervation (10x

magnification; scale bar = 100µm). Statistical significance is depicted where present.

Data presented as Mean ± SEM (N= 5/group).

Figure 9 Knockdown of Fbxl22 in MuRF1 KO muscle results in improved muscle

sparing after 7 days of denervation. (**A**) Total muscle mass for gastrocnemius (GA;

lateral and medial) muscles transfected with either empty vector (EV) or Fblx22 RNAi

constructs. (**B**) Loss of GA muscle mass with either EV or Fbxl22 RNAi present

compared to contralateral control (innervated) muscle. Data expressed as percent loss

compared to contralateral control. Muscle mass for **(C)** lateral and **(D)** medial

gastrocnemius muscles transfected with EV or Fbxl22 RNAi construct after 7 days of

denervation. (**E**) Mean Fiber cross-sectional area (CSA) for GFP-positive only fibers in

transfected GA muscles was determined. Muscle CSA distribution in transfected

muscles with the **(F)** Fbxl22 RNAi or **(G)** EV constructs. GFP-positive fibers were

measured for CSA, with ≥ 350 transfected fibers analyzed per animal, per muscle (n=

5/group). In addition, the muscle CSA distributions for non-transfected (red bars) and

innervated (blue triangle) contralateral control muscles were measured. Data presented

as a percentage of sizes between 0-5000µm. (**H**) Representative images for GFP

fluorescence and laminin stain in Fbxl22 RNAi and EV transfected lateral and medial

gastrocnemius muscles after 7 days of denervation (10x magnification; scale bar =

100µm). Statistical significance is depicted where present. Data presented as Mean ±

SEM (N= 5-6/group).

References

1.   **Atherton PJ, Greenhaff PL, Phillips SM, Bodine SC, Adams CM, and Lang CH**. Control of skeletal muscle atrophy in response to disuse: clinical/preclinical contentions and fallacies of evidence. *American journal of physiology Endocrinology and metabolism* 311: E594-E604, 2016.

2.   **Baehr LM, West DWD, Marcotte G, Marshall AG, De Sousa LG, Baar K, and Bodine SC**. Age-related deficits in skeletal muscle recovery following disuse are associated with neuromuscular junction instability and ER stress, not impaired protein synthesis. *Aging* 8: 127-146, 2016.

3.   **Baehr LM, West DWD, Marshall AG, Marcotte GR, Baar K, and Bodine SC**. Muscle-specific and age-related changes in protein synthesis and protein degradation in response to hindlimb unloading in rats. *J Appl Physiol (1985)* 122: 1336-1350, 2017.

4.   **Beggs AH, Byers TJ, Knoll JH, Boyce FM, Bruns GA, and Kunkel LM**. Cloning and characterization of two human skeletal muscle alpha-actinin genes located on chromosomes 1 and 11. *J Biol Chem* 267: 9281-9288, 1992.

5.   **Bilodeau PA, Coyne ES, and Wing SS**. The ubiquitin proteasome system in atrophying skeletal muscle: roles and regulation. *American Journal of Physiology-Cell Physiology* 311: C392-C403, 2016.

6.   **Bodine SC**. Adolph Lecture: Skeletal Muscle Atrophy: Multiple Pathways Leading to a Common Outcome. *Journal of Applied Physiology*.

7.   **Bodine SC**. Disuse-induced muscle wasting. *The international journal of biochemistry & cell biology* 45: 2200-2208, 2013.

8.   **Bodine SC, and Baehr LM**. Skeletal muscle atrophy and the E3 ubiquitin ligases MuRF1 and MAFbx/atrogin-1. *American Journal of Physiology-Endocrinology and Metabolism* 307: E469-E484, 2014.

9.   **Bodine SC, Latres E, Baumhueter S, Lai VK-M, Nunez L, Clarke BA, Poueymirou WT, Panaro FJ, Na E, and Dharmarajan K**. Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science* 294: 1704-1708, 2001.

10.   **Calura E, Cagnin S, Raffaello A, Laveder P, Lanfranchi G, and Romualdi C**. Meta-analysis of expression signatures of muscle atrophy: gene interaction networks in early and late stages. *BMC Genomics* 9: 630, 2008.

11.   **Chen JL, Colgan TD, Walton KL, Gregorevic P, and Harrison CA**. The TGF-β Signalling Network in Muscle Development, Adaptation and Disease. *Advances in experimental medicine and biology* 900: 97-131, 2016.

12.   **Chen JL, Walton KL, Hagg A, Colgan TD, Johnson K, Qian H, Gregorevic P, and Harrison CA**. Specific targeting of TGF-β family ligands demonstrates distinct roles in the regulation of muscle mass in health and disease. *Proceedings of the National Academy of Sciences of the United States of America* 114: E5266-E5275, 2017.

13.   **Čížková D, Soukup T, and Mokrý J**. Expression of nestin, desmin and vimentin in intact and regenerating muscle spindles of rat hind limb skeletal muscles. *Histochemistry and cell biology* 131: 197-206, 2009.

14.   **Clarke BA, Drujan D, Willis MS, Murphy LO, Corpina RA, Burova E, Rakhilin SV, Stitt TN, Patterson C, and Latres E**. The E3 Ligase MuRF1 degrades myosin heavy chain protein in dexamethasone-treated skeletal muscle. *Cell metabolism* 6: 376-385, 2007.

15.   **Cohen S, Brault JJ, Gygi SP, Glass DJ, Valenzuela DM, Gartner C, Latres E, and Goldberg AL**. During muscle atrophy, thick, but not thin, filament components are degraded by MuRF1-dependent ubiquitylation. *The Journal of cell biology* 185: 1083-1095, 2009.

16.   **Cohen S, Nathan JA, and Goldberg AL**. Muscle wasting in disease: molecular mechanisms and promising therapies. *Nature reviews Drug discovery* 14: 58, 2015.

17.   **Cohen S, Zhai B, Gygi SP, and Goldberg AL**. Ubiquitylation by Trim32 causes coupled loss of desmin, Z-bands, and thin filaments in muscle atrophy. *J Cell Biol* 198: 575-589, 2012.

18.   **Cohen TJ, Waddell DS, Barrientos T, Lu Z, Feng G, Cox GA, Bodine SC, and Yao T-P**. The histone
deacetylase HDAC4 connects neural activity to muscle transcriptional reprogramming. *Journal of*
Biological Chemistry 282: 33752-33759, 2007.
19.   **Daya S, and Berns KI**. Gene therapy using adeno-associated virus vectors. *Clinical microbiology*
reviews 21: 583-593, 2008.
20.   **Deglincerti A, Haremaki T, Warmflash A, Sorre B, and Brivanlou AH**. Coco is a dual activity
modulator of TGFβ signaling. *Development* 142: 2678-2685, 2015.
21.   **Ebert SM, Al-Zougbi A, Bodine SC, and Adams CM**. Skeletal Muscle Atrophy: Discovery of
Mechanisms and Potential Therapies. *Physiology* 34: 232-239, 2019.
22.   **Ebert SM, Dyle MC, Kunkel SD, Bullard SA, Bongers KS, Fox DK, Dierdorff JM, Foster ED, and**
Adams CM. Stress-induced skeletal muscle Gadd45a expression reprograms myonuclei and causes
muscle atrophy. *J Biol Chem* 287: 27290-27301, 2012.
23.   **Evans WJ**. Skeletal muscle loss: cachexia, sarcopenia, and inactivity. *The American Journal of*
Clinical Nutrition 91: 1123S-1127S, 2010.
24.   **Fuqua JD, Mere CP, Kronemberger A, Blomme J, Bae D, Turner KD, Harris MP, Scudese E,**
Edwards M, Ebert SM, GO de Sousa L, Bodine SC, Yang L, Adams CM, and Lira VA. ULK2 is essential for
degradation of ubiquitinated protein aggregates and homeostasis in skeletal muscle. *The FASEB Journal*
33: 11735-12745, 2019.
25.   **Furlow JD, Watson ML, Waddell DS, Neff ES, Baehr LM, Ross AP, and Bodine SC**. Altered gene
expression patterns in muscle ring finger 1 null mice during denervation- and dexamethasone-induced
muscle atrophy. *Physiol Genomics* 45: 1168-1185, 2013.
26.   **Gallanti A, Prelle A, Moggio M, Ciscato P, Checcarelli N, Sciacco M, Comini A, and Scarlato G**.
Desmin and vimentin as markers of regeneration in muscle diseases. *Acta neuropathologica* 85: 88-92,
1992.
27.   **Gomes AV, Waddell DS, Siu R, Stein M, Dewey S, Furlow JD, and Bodine SC**. Upregulation of
proteasome activity in muscle RING finger 1-null mice following denervation. *The FASEB Journal* 26:
2986-2999, 2012.
28.   **Gomes MD, Lecker SH, Jagoe RT, Navon A, and Goldberg AL**. Atrogin-1, a muscle-specific F-box
protein highly expressed during muscle atrophy. *Proceedings of the National Academy of Sciences* 98:
14440-14445, 2001.
29.   **Haddock AN, Labuzan SA, Haynes AE, Hayes CS, Kakareka KM, and Waddell DS**. Dual-specificity
phosphatase 4 is upregulated during skeletal muscle atrophy and modulates extracellular signal-
regulated kinase activity. *American Journal of Physiology-Cell Physiology* 316: C567-C581, 2019.
30.   **Hayes CS, Labuzan SA, Menke JA, Haddock AN, and Waddell DS**. Ttc39c is upregulated during
skeletal muscle atrophy and modulates ERK1/2 MAP kinase and hedgehog signaling. *Journal of cellular*
physiology 2019.
31.   **Heinemeier KM, Olesen JL, Haddad F, Schjerling P, Baldwin KM, and Kjaer M**. Effect of
unloading followed by reloading on expression of collagen and related growth factors in rat tendon and
muscle. *Journal of applied physiology* 106: 178-186, 2009.
32.   **Hughes DC, Marcotte GR, Baehr LM, West DW, Marshall AG, Ebert SM, Davidyan A, Adams**
CM, Bodine SC, and Baar K. Alterations in the muscle force transfer apparatus in aged rats during
unloading and reloading: impact of microRNA-31. *The Journal of physiology* 596: 2883-2900, 2018.
33.   **Hughes DC, Marcotte GR, Marshall AG, West DW, Baehr LM, Wallace MA, Saleh PM, Bodine**
SC, and Baar K. Age-related differences in dystrophin: impact on force transfer proteins, membrane
integrity, and neuromuscular junction stability. *The Journals of Gerontology: Series A* 72: 640-648, 2017.
34.   **Hughes DC, Wallace MA, and Baar K**. Effects of aging, exercise, and disease on force transfer in
skeletal muscle. *American Journal of Physiology-Endocrinology and Metabolism* 309: E1-E10, 2015.

35.   **Hutchins AP, Liu S, Diez D, and Miranda-Saavedra D**. The repertoires of ubiquitinating and
deubiquitinating enzymes in eukaryotic genomes. *Molecular biology and evolution* 30: 1172-1187, 2013.
36.   **Kawai J, Shinagawa A, Shibata K, Yoshino M, Itoh M, Ishii Y, Arakawa T, Hara A, Fukunishi Y,**
Konno H, Adachi J, Fukuda S, Aizawa K, Izawa M, Nishi K, Kiyosawa H, Kondo S, Yamanaka I, Saito T,
Okazaki Y, Gojobori T, Bono H, Kasukawa T, Saito R, Kadota K, Matsuda H, Ashburner M, Batalov S,
Casavant T, Fleischmann W, Gaasterland T, Gissi C, King B, Kochiwa H, Kuehl P, Lewis S, Matsuo Y,
Nikaido I, Pesole G, Quackenbush J, Schriml LM, Staubli F, Suzuki R, Tomita M, Wagner L, Washio T,
Sakai K, Okido T, Furuno M, Aono H, Baldarelli R, Barsh G, Blake J, Boffelli D, Bojunga N, Carninci P, de
Bonaldo MF, Brownstein MJ, Bult C, Fletcher C, Fujita M, Gariboldi M, Gustincich S, Hill D, Hofmann M,
Hume DA, Kamiya M, Lee NH, Lyons P, Marchionni L, Mashima J, Mazzarelli J, Mombaerts P, Nordone
P, Ring B, Ringwald M, Rodriguez I, Sakamoto N, Sasaki H, Sato K, Schönbach C, Seya T, Shibata Y,
Storch KF, Suzuki H, Toyo-oka K, Wang KH, Weitz C, Whittaker C, Wilming L, Wynshaw-Boris A,
Yoshida K, Hasegawa Y, Kawaji H, Kohtsuki S, Hayashizaki Y, The RGERGPIIT, the FC, The RGERGPIIT,
Consortium F, and General o. Functional annotation of a full-length mouse cDNA collection. *Nature* 409:
685-690, 2001.
37.   **Kroeger KM, and Abraham LJ**. Identification of an AP-2 element in the-323 to-285 region of the
TNF-α gene. *IUBMB Life* 40: 43-51, 1996.
38.   **Lang F, Aravamudhan S, Nolte H, Türk C, Hölper S, Müller S, Günther S, Blaauw B, Braun T, and**
Krüger M. Dynamic changes in the mouse skeletal muscle proteome during denervation-induced
atrophy. *Disease models & mechanisms* 10: 881-896, 2017.
39.   **Leber Y, Ruparelia AA, Kirfel G, van der Ven PF, Hoffmann B, Merkel R, Bryson-Richardson RJ,**
and Furst DO. Filamin C is a highly dynamic protein associated with fast repair of myofibrillar
microdamage. *Hum Mol Genet* 25: 2776-2788, 2016.
40.   **Li Z, Mericskay M, Agbulut O, Butler-Browne G, Carlsson L, Thornell L-E, Babinet C, and Paulin**
D. Desmin is essential for the tensile strength and integrity of myofibrils but not for myogenic
commitment, differentiation, and fusion of skeletal muscle. *The Journal of cell biology* 139: 129-144,
1997.
41.   **Lovering RM, O'Neill A, Muriel JM, Prosser BL, Strong J, and Bloch RJ**. Physiology, structure, and
susceptibility to injury of skeletal muscle in mice lacking keratin 19-based and desmin-based
intermediate filaments. *American Journal of Physiology-Cell Physiology* 300: C803-C813, 2011.
42.   **Lynch S, McLeod M, Orsech H, Cirelli A, and Waddell D**. Zinc finger protein 593 is upregulated
during skeletal muscle atrophy and modulates muscle cell differentiation. *Experimental cell research*
383: 111563, 2019.
43.   **Mantovani R**. The molecular biology of the CCAAT-binding factor NF-Y. *Gene* 239: 15-27, 1999.
44.   **Merlie JP, Mudd J, Cheng T-C, and Olson EN**. Myogenin and acetylcholine receptor alpha gene
promoters mediate transcriptional regulation in response to motor innervation. *Journal of Biological*
Chemistry 269: 2461-2467, 1994.
45.   **Milan G, Romanello V, Pescatore F, Armani A, Paik J-H, Frasson L, Seydel A, Zhao J, Abraham R,**
and Goldberg AL. Regulation of autophagy and the ubiquitin–proteasome system by the FoxO
transcriptional network during muscle atrophy. *Nature communications* 6: 6670, 2015.
46.   **Miller BF, Baehr LM, Musci RV, Reid JJ, Peelor FF, 3rd, Hamilton KL, and Bodine SC**. Muscle-
specific changes in protein synthesis with aging and reloading after disuse atrophy. *J Cachexia*
Sarcopenia Muscle 2019.
47.   **Mitch WE, and Goldberg AL**. Mechanisms of muscle wasting—the role of the ubiquitin–
proteasome pathway. *New England Journal of Medicine* 335: 1897-1905, 1996.
48.   **Moresi V, Williams AH, Meadows E, Flynn JM, Potthoff MJ, McAnally J, Shelton JM, Backs J,**
Klein WH, and Richardson JA. Myogenin and class II HDACs control neurogenic muscle atrophy by
inducing E3 ubiquitin ligases. *Cell* 143: 35-45, 2010.

49.   **Naso MF, Tomkowicz B, Perry WL, and Strohl WR**. Adeno-associated virus (AAV) as a vector for
gene therapy. *BioDrugs* 31: 317-334, 2017.
50.   **Palmisano MG, Bremner SN, Hornberger TA, Meyer GA, Domenighetti AA, Shah SB, Kiss B,**
Kellermayer M, Ryan AF, and Lieber RL. Skeletal muscle intermediate filaments form a stress-
transmitting and stress-signaling network. *J Cell Sci* 128: 219-224, 2015.
51.   **Paul PK, Bhatnagar S, Mishra V, Srivastava S, Darnay BG, Choi Y, and Kumar A**. The E3 Ubiquitin
Ligase TRAF6 Intercedes in Starvation-Induced Skeletal Muscle Atrophy through Multiple Mechanisms.
Molecular and Cellular Biology 32: 1248-1259, 2012.
52.   **Pillon NJ, Gabriel BM, Dollet L, Smith JAB, Sardón Puig L, Botella J, Bishop DJ, Krook A, and**
Zierath JR. Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nature*
Communications 11: 470, 2020.
53.   **Polge C, Heng A-E, Jarzaguet M, Ventadour S, Claustre A, Combaret L, Béchet D, Matondo M,**
Uttenweiler-Joseph S, Monsarrat B, Attaix D, and Taillandier D. Muscle actin is polyubiquitinylated in
vitro and in vivo and targeted for breakdown by the E3 ligase MuRF1. *The FASEB Journal* 25: 3790-3802,
2011.
54.   **Ramaswamy KS, Palmer ML, Van Der Meulen JH, Renoux A, Kostrominova TY, Michele DE, and**
Faulkner JA. Lateral transmission of force is impaired in skeletal muscles of dystrophic mice and very old
rats. *The Journal of physiology* 589: 1195-1208, 2011.
55.   **Sandri M**. Protein breakdown in muscle wasting: role of autophagy-lysosome and ubiquitin-
proteasome. *The international journal of biochemistry & cell biology* 45: 2121-2129, 2013.
56.   **Sandri M, Sandri C, Gilbert A, Skurk C, Calabria E, Picard A, Walsh K, Schiaffino S, Lecker SH,**
and Goldberg AL. Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and
cause skeletal muscle atrophy. *Cell* 117: 399-412, 2004.
57.   **Sartori R, Schirwis E, Blaauw B, Bortolanza S, Zhao J, Enzo E, Stantzou A, Mouisel E, Toniolo L,**
and Ferry A. BMP signaling controls muscle mass. *Nature genetics* 45: 1309, 2013.
58.   **Schiaffino S, Dyar KA, Ciciliot S, Blaauw B, and Sandri M**. Mechanisms regulating skeletal
muscle growth and atrophy. *Febs j* 280: 4294-4314, 2013.
59.   **Seaborne RA, Hughes DC, Turner DC, Owens DJ, Baehr LM, Gorski P, Semenova EA, Borisov**
OV, Larin AK, Popov DV, Generozov EV, Sutherland H, Ahmetov II, Jarvis JC, Bodine SC, and Sharples
AP. UBR5 is a novel E3 ubiquitin ligase involved in skeletal muscle hypertrophy and recovery from
atrophy. *The Journal of Physiology* 597: 3727-3749, 2019.
60.   **Sharples AP, Hughes DC, Deane CS, Saini A, Selman C, and Stewart CE**. Longevity and skeletal
muscle mass: the role of IGF signalling, the sirtuins, dietary restriction and protein intake. *Aging cell* 14:
511-523, 2015.
61.   **Sokołowska E, and Błachnio-Zabielska AU**. A Critical Review of Electroporation as A Plasmid
Delivery System in Mouse Skeletal Muscle. *International journal of molecular sciences* 20: 2776, 2019.
62.   **Spaich S, Will RD, Just S, Spaich S, Kuhn C, Frank D, Berger IM, Wiemann S, Korn B, Koegl M,**
Backs J, Katus HA, Rottbauer W, and Frey N. F-Box and Leucine-Rich Repeat Protein 22 Is a Cardiac-
Enriched F-Box Protein That Regulates Sarcomeric Protein Turnover and Is Essential for Maintenance of
Contractile Function In Vivo. *Circulation Research* 111: 1504-1516, 2012.
63.   **Tang H, Macpherson P, Marvin M, Meadows E, Klein WH, Yang X-J, and Goldman D**. A histone
deacetylase 4/myogenin positive feedback loop coordinates denervation-dependent gene induction and
suppression. *Molecular biology of the cell* 20: 1120-1131, 2009.
64.   **Terry EE, Zhang X, Hoffmann C, Hughes LD, Lewis SA, Li J, Wallace MJ, Riley LA, Douglas CM,**
and Gutierrez-Monreal MA. Transcriptional profiling reveals extraordinary diversity among skeletal
muscle tissues. *Elife* 7: e34613, 2018.

65.   **Vaittinen S, Lukka R, Sahlgren C, Hurme T, Rantanen J, Lendahl U, Eriksson JE, and Kalimo H**.
The expression of intermediate filament protein nestin as related to vimentin and desmin in
regenerating skeletal muscle. *Journal of neuropathology & experimental neurology* 60: 588-597, 2001.
66.   **Waddell DS, Baehr LM, Van Den Brandt J, Johnsen SA, Reichardt HM, Furlow JD, and Bodine
SC**. The glucocorticoid receptor and FOXO1 synergistically activate the skeletal muscle atrophy-
associated MuRF1 gene. *American Journal of Physiology-Endocrinology and Metabolism* 295: E785-E797,
2008.
67.   **Wang C, Yue F, and Kuang S**. Muscle Histology Characterization Using H&E Staining and Muscle
Fiber Type Classification Using Immunofluorescence Staining. *Bio Protoc* 7: 2017.
68.   **Wen Y, Murach KA, Vechetti IJ, Jr., Fry CS, Vickery C, Peterson CA, McCarthy JJ, and Campbell
KS**. MyoVision: software for automated high-content analysis of skeletal muscle immunohistochemistry.
J Appl Physiol (1985) 124: 40-51, 2018.
69.   **Workeneh BT, and Mitch WE**. Review of muscle wasting associated with chronic kidney disease.
The American journal of clinical nutrition 91: 1128S-1132S, 2010.

1059

1060

1062   **Figure 1**

1063

gtggccgcgcacgggcgcgcgctgcagactctgcacgtggacttctgccgcaacgtgagcgccgccggcctgctccgcctccgcgccgcctgcccgaacctgcgcctcagcgccgaacgcagcgcggcc

1064

1065 **Figure 2**

1066

1067

Figure 3

1069

1071 **Figure 4**

1072

1073   **Figure 5**

1074

1075 **Figure 6**

1076

Figure 7

1078

1079

1080   **Figure 8**

1082   **Figure 9**