Gap junctions deliver malonyl-CoA from soma to germline to support

embryogenesis in *Caenorhabditis elegans*

Todd A. Starich[1,2] and David Greenstein[1]

[1]Department of Genetics, Cell Biology and Development, University of

Minnesota, Minneapolis, Minnesota 55455 USA

[2]Email: stari001@umn.edu

Abstract

Gap junctions are ubiquitous in metazoans and play critical roles in

important biological processes, including electrical conduction and development.

Yet, only a few defined molecules passing through gap junction channels have

been linked to specific functions. We isolated gap junction channel mutants that

reduce coupling between the soma and germ cells in the *C. elegans* gonad. We

provide evidence that malonyl-CoA, the rate-limiting substrate for fatty acid

synthesis (FAS), is produced in the soma and delivered through gap junctions to

the germline; there it is used by fatty acid synthase to critically support embryonic

development. Separation of malonyl-CoA production from its site of utilization

facilitates somatic control of germline development. Additionally, we demonstrate

that loss of malonyl-CoA production in the intestine negatively impacts germline

development independently of FAS. Our results suggest that metabolic

outsourcing of malonyl-CoA may be a strategy by which the soma communicates

nutritional status to the germline.

Impact Statement

Malonyl-CoA, the rate-limiting substrate for fatty acid synthesis, is produced in

the soma and delivered through gap junctions to the germline to promote

reproduction and coordinate it with nutritional status.

Introduction

Gap junctions are clusters of intercellular channels allowing direct passage of

small molecules between coupled cells. They are widespread in multicellular

organisms, though different gene families encode the protein subunits in

invertebrates (innexins) and vertebrates (primarily connexins) (*Skerrett et al.,*

2017). While the capability of small molecules to pass through gap junctions can

be shown readily in cultured cells, identification of essential molecules that must

pass through channels *in vivo* is challenging. Additionally, disruption of gap

junction coupling can affect cell adhesion (*Meyer et al., 1992*), and resultant

defects in intercellular communication may not necessarily relate to the passage

of molecules through junctional channels.

In *C. elegans*, germ cells are coupled throughout their development to the

somatic gonad by gap junctions (*Starich et al., 2014*). Germ cells progress in

assembly-line fashion through the early stages of meiosis (*Figure 1A*). Gonadal

gap junctions have been implicated in germ cell proliferation, oocyte growth,

regulation of oocyte meiotic maturation, early embryogenesis and sperm

guidance (*Govindan et al., 2006; Whitten et al., 2007; Nadarajan et al., 2009;*

Edmonds et al., 2011; Starich et al., 2014). Gap junction hemichannels (half-

channels) in the somatic gonad are comprised of the recently duplicated innexins

INX-8 and INX-9; apposing germ cell hemichannels are heteromeric and consist

of INX-14 in conjunction with either INX-21 or INX-22 (*Figure 1–figure*

supplement 1). Germ cells fail to proliferate in *inx-8(0) inx-9(0)* mutants, at a point

upstream of, or parallel to, GLP-1/Notch activation of germ cells by Delta class

ligands LAG-2 and APX-1. These ligands are produced in the distal tip cell (DTC)

that maintains the proliferative stem cell niche (*Austin and Kimble, 1987, 1989;*

Henderson et al., 1994; Nadarajan et al., 2009).

Germ cell proliferation can be rescued in *inx-8(0) inx-9(0)* mutants to half

the level of wild type by driving *inx-8* expression in the DTC with the *lag-2*

promoter (*Starich et al., 2014*). In these animals, gap junctions between soma

and germline are established in the distal gonad arm, but ostensibly not in the

proximal arm; many resultant embryos exhibit characteristics of the Pod

phenotype (Polarity and Osmotic sensitivity Defect) (*Tagawa et al., 2001*),

including eggshell and permeability barrier deficiencies, cytokinesis failures, and

polar body extrusion defects. Other genes with mutations producing Pod

embryos include those involved in eggshell synthesis, fatty acid synthesis (FAS),

the APC/separase pathway, and a coronin-like protein (*Rappleye et al., 1999,*

2003; Tagawa et al., 2001; Zhang et al., 2005; Johnston et al., 2006; Olson et al.,

2006; Benenati et al., 2009).

Molecules moving through soma-germ cell gap junctions necessary for

proliferation and early embryogenesis are unknown. We use genetic methods to

show that malonyl-CoA (mal-CoA) is produced by *pod-2* (acetyl-CoA

carboxylase, ACC) in the soma. Transfer of mal-CoA to the germline, for use by

fasn-1 (fatty acid synthase, FASN), is impacted by the integrity of INX-8-

containing channels under conditions in which gap junction formation appears

normal. Rescue of the Pod phenotype in *pod-2* genetic mosaics indicates a

somatic sheath focus of action. Our results support the model that mal-CoA

transits from soma to germline through gap junction channels.

76 **Results**

77 **Gametogenesis is delayed by the *inx-8(tn1513 tn1555)* reduction of**

78 **function (*rf*) mutation**

79 We previously isolated (in an *inx-9* null background) a strong loss-of-function *inx-*

80 *8* allele *(tn1513)* with sufficient residual function to contribute to channel

81 formation and cell adhesion but not to support germ cell proliferation (*Starich et*

82 *al., 2014*). *tn1513* encodes a T239I change in the second extracellular loop of the

83 innexin monomer, predicted to be involved in docking with octameric

84 hemichannels on apposing cells (*Figure 1–figure supplement 1*) (*Oshima et al.,*

85 *2016*). Using this allele in a genetic screen for restoration of germ cell

86 proliferation, six *inx-8* intragenic suppressor mutations were isolated (to be

87 described elsewhere). One of these, *tn1555*, encodes a D24N change in the

88 amino terminus (*Figure 1–figure supplement 1*), near a site that may associate

89 with a cytoplasmic dome structure surrounding the hemichannel outer pore

90 (*Oshima et al., 2016*). *inx-8(tn1513 tn1555rf)*—for brevity to be referred to as *inx-*

91 *8(rf)*—partially restored germ cell numbers and fertility to one-third the level of

92 wild type (*Figure 1–figure supplement table 1*). *inx-8(rf)* is rescued by an *inx-8(+)*

93 extrachromosomal array, and rescue of *inx-8(0) inx-9(0)* with an *inx-8(rf)::gfp*

94 array mimics the phenotype of the suppressor mutant; therefore observed

95 phenotypes can be attributed to *inx-8(rf)* and not to other possible background

96 mutations.

There is an ~18 hr delay to the onset of fertilization in *inx-8(rf)* compared

to wild type, as determined by relating the timing of first ovulation to the L4 molt

(L4-to-adult transition) (*Figure 1B*). This delay correlates with late expression of

gfp::lin41 in relation to L4 sub-stages of vulval development (*Mok et al., 2015*);

LIN-41 is the earliest known marker of oogenesis (*Spike et al., 2014*). In wild type

at 20^{o}C, *gfp::lin41* appears by sub-stage L4.7 (*Figure 1C*); in *inx-8(rf)*, *gfp::lin41*

is expressed after completion of vulval morphogenesis but prior to the L4 molt

(*Figure 1C; Figure 1–figure supplement table 2*). Spermatogenesis is also

delayed—wild-type sperm appear as early as the L4.7 stage, but *inx-8(rf)* sperm

first appear after completion of vulval morphogenesis (*Figure 1–figure*

supplement table 2). Once started, egg-laying in *inx-8(rf)* continues apace such

that ~99% of progeny are laid within a 96-hour window (3144/3179 larvae; n=25),

similar to wild type (5847/5929 larvae; n=21).

inx-8(rf) gap junction formation is more severely affected in proximal

gonad arms

To examine expression of INX-8(T239I D24N) mutant innexins encoded by *inx-*

8(rf), specific antibodies were used to label somatic (anti-INX-8) or the two

classes of germline (anti-INX-21 or anti-INX-22) hemichannels. In wild-type

hermaphrodites, clusters of gap junction puncta form between somatic DTC or

sheath cells and each underlying germ cell (*Figure 2A*). In distal arms of wild

type, loosely clustered sheath-germ cell gap junctions form more basally on germ

cells; in *inx-8(rf)* distal arms, both INX-21 and INX-22 puncta tend to localize

more laterally, where the sheath cells dip down between adjacent germ cells

(*Figure 2B*). In the loop region (transition from meiotic pachytene to diplotene),

large, higher-order clusters of gap junction puncta form in wild-type animals;

these are absent in *inx-8(rf)* (*Figure 2C*). In wild-type proximal arms (diakinesis),

developing oocytes exhibit extensive formation of INX-22 gap junction puncta,

while INX-21 puncta become faint; in *inx-8(rf)* proximal arms, INX-22 and INX-8

are strongly but diffusely expressed, with little co-localization evident (*Figure 2D*).

These results suggest there may be little gap junction coupling in *inx-8(rf)*

proximal arms. We showed previously that sheath-oocyte gap junctions are

endocytosed by the oocyte during ovulation (*Starich et al., 2014*). Embryos from

inx-8(0) inx-9(0) rescued with *inx-8(rf)::gfp* show internalized GFP puncta,

confirming that *inx-8(rf)* establishes at least some sheath-oocyte gap junctions in

the proximal arm (*Figure 2–figure supplement 1*).

Reduced proximal arm expression of gap junctions is seen in *inx-8(0) inx-*

9(0); Ex[lag-2p::inx-8::gfp] animals—simplified to *inx-8/9(0); Ex[inx-8(DTC+, Sh-*

)]—which are rescued for germ cell proliferation by driving expression of INX-

8::GFP in the DTC, but not sheath (Sh), using the *lag-2* promoter (*Starich et al.,*

2014). These animals produce primarily dead embryos, many with Pod

characteristics (*Table 1*). In contrast to *inx-8(rf)*, these embryos do not show

evidence of *inx-8::gfp* endocytosis, consistent with absence of proximal arm

sheath-germline gap junction coupling (*Figure 2–figure supplement 1*). *inx-8/9(0);*

Ex[inx-8(DTC+, Sh-)] animals also show delayed onset of *gfp::lin41* expression,

similar to *inx-8(rf)* (*Figure 1C; Figure 1–figure supplement table 2*).

We conclude that delayed gametogenesis in *inx-8(rf)* mutants arises from

compromised soma-germline gap junction communication, due to altered channel

properties that influence the formation and localization of gap junctions,

especially in the loop and proximal arm. These regions are additionally implicated

because DTC expression of wild-type INX-8 fails to rescue *inx-8(rf)* phenotypes

(n>20).

***inx-8(rf)* interacts with FAS mutants**

Because *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* produces Pod-like embryos, and *inx-*

8(rf) shows reduced proximal arm gap junction coupling similar to *inx-8/9(0);*

Ex[inx-8(DTC+, Sh-)], possible genetic interactions between *inx-8(rf)* and other

Pod mutants were explored (*Figure 1–figure supplement table 1*). Conditional

alleles of genes in the FAS pathway (*Figure 3A*) were used to construct double

mutants with *inx-8(rf)*. Alleles included: *pod-2(ye60)*, a cold-sensitive (*cs*) allele of

the *C. elegans* acetyl-CoA carboxylase ACC Type 1 (*Rappleye et al., 2003*); a

temperature-sensitive (*ts*) allele of fatty acid synthase *fasn-1(g43) (Jaramillo-*

Lambert et al., 2015); and *emb-8(hc69)*, a *ts* allele of an NADPH-cytochrome-

P450 reductase, likely necessary for reduction of CYP-450 enzymes involved in

fatty acid modifications (*Rappleye et al., 2003; Benenati et al., 2009*). Possible

interaction between *pod-2(ye60cs)* and *inx-8(rf)* was difficult to assess–*ye60cs*

grown at 15°C was quite sensitive to brief handling at room temperature. At a

partially restrictive temperature (22°C), *fasn-1(g43ts); inx-8(rf)* and *emb-*

8(hc69ts); inx-8(rf) double mutants exhibited markedly reduced brood sizes.

Similar reductions in brood size were also apparent in *emb-8(hc69ts); +/inx-8(rf)*,

suggesting *emb-8(hc69ts)* might be highly dose-sensitive to *inx-8(+)* levels

(*Figure 1–figure supplement table 1*).

FAS pathway shifts from soma to germline after *pod-2*

Genetic interactions between *inx-8* and FAS genes could represent activity in the

same or parallel pathways. If gap junctions are positioned in the FAS pathway, it

might be possible to find a step within this pathway where gene function shifts

from a somatic to a germline focus. *C. elegans* is particularly amenable to such

an investigation because simple (highly repetitive) extrachromosomal arrays

used to rescue gene function tend to express well in the soma, whereas arrays

with more sequence complexity are generally required to prevent silencing of

multi-copy gene expression in the germline (*Kelly et al., 1997*).

Previous characterization of *pod-2* implied rescue by a simple array

(*Rappleye et al., 2003*); we likewise found that *pod-2(ye60cs)* is rescued by

simple extrachromosomal arrays at 15°C (see Materials and methods). In

contrast, 2 fosmids encoding *emb-8*, used individually to generate simple arrays,

yielded 65 transformed lines, only one of which rescued *emb-8(hc69ts)* at 25°C;

the array in this line was eventually silenced. For *fasn-1*, two fosmids used to

generate 93 simple array lines and 20 complex array lines failed to rescue *fasn-*

1(g43ts). These results were consistent with a model in which FAS affecting

embryonic development requires *pod-2* in the soma, but shifts to a germline

requirement for *fasn-1* and *emb-8*.

To avoid misrepresentation of rescue results due to the special nature of a

conditional allele, null alleles of *pod-2(tn1691)* and *fasn-1(tn1762)*–simplified to

(0)–were generated using CRISPR-Cas9 methodology (*Figure 3B, C*) (Arribere *et*

al., 2014). *pod-2(0)* embryos from *pod-2(0)/+* parents arrest late in

embryogenesis (pretzel stage; *Figure 3D*); most *fasn-1(0)* embryos from *fasn-*

1(0)/+ parents shows signs of differentiation, and many arrest late in

embryogenesis; occasional hatchlings emerge with probable hypodermal defects

(*Figure 3E-G*). *pod-2(0)* was rescued by simple extrachromosomal arrays,

supporting probable somatic function. Using a PCR-amplified genomic fragment

we were able to rescue somatic but not germline function of *fasn-1(0)* with a

simple or complex array (see Materials and methods). A genetically-balanced line

with a complex array, *hT2/ fasn-1(0)*; *Ex[fasn-1(+); sur-5::gfp(+)]*, segregates

healthy homozygous *fasn-1(0); Ex[fasn-1(+); sur-5::gfp(+)]* adults that lay almost

exclusively Pod embryos (Table 1). (*sur-5::gfp* is a co-injection marker expressed

in somatic cell nuclei (*Yochem et al., 19*98); corresponding extrachromosomal

arrays can be lost mitotically to generate mosaics—see below). Gonad arms

appear normal, and animals lay wild-type numbers of defective embryos (Table

1); many of these display polarity defects (4/14 symmetric first divisions) and

exhibit extra nuclei in early embryos. Embryos typically arrest prior to

morphogenesis (see below). We conclude this *fasn-1(+)* PCR fragment rescues

somatic *fasn-1* function and is either silenced in the germline or lacks an element

necessary for germline expression; in either case a germline requirement is

indicated. In sum, rescue results are consistent with a fertility requirement for

pod-2 in the somatic gonad and *fasn-1* in the germline. This conclusion is

supported by genetic mosaic and gene expression data described below. If *inx-8*

gap junctions act in this pathway, it would position these junctions between *pod-2*

and *fasn-1* as possible mediators of mal-CoA transfer from soma to germline.

GFP::POD-2 and GFP::FASN-1 are tissue restricted

Endogenous, homozygous-viable, fertile GFP-labeled alleles of *pod-2* and *fasn-1*

were generated to determine if patterns of localization are consistent with

predicted foci of action. The GFP moiety in *gfp::pod-2(tn1765)* is expected to

label all 3 predicted isoforms (*Figure 3B*) (*WormBase*). GFP::POD-2 expression

is strongest in the intestine, and prominent in the hypodermis, gonadal sheath

(especially proximal sheath), CAN neuron, and excretory duct (*Figure 4A-C*).

gfp::fasn-1(tn1782) labeled at the amino terminus (*Figure 3C*) is

expressed in the hypodermis, gonad sheath (esp. proximal) and excretory duct;

intestinal and CAN neuron expression are not evident (*Figure 5A-C*). This

expression pattern partially overlaps with that of a *fasn-1* promoter::GFP reporter

fusion, which was expressed in the intestine (*Lee et al., 2010*). WormBase

predicts a single FASN-1 isoform, but a second in-frame ATG lies in Exon 2;

intestinal FASN-1 could be derived from translational initiation at this site, which

would not be revealed in our construct. Attempts to GFP-label FASN-1

downstream of this second ATG, or at the C-terminus, were unsuccessful.

Neither *gfp::pod-2* nor *gfp::fasn-1* is expressed in the DTC.

Germline expression for either GFP-tagged gene was not detected (*Figure*

4C; Figure 5B). Anti-GFP staining of dissected gonads failed to detect germline

GFP::POD-2 expression (*Figure 4C*). The earliest detectable GFP::FASN-1 in

embryos appears in developing hypodermis (*Figure 5D*); GFP::POD-2 is

prominent in developing hypodermis and intestine (*Figure 4D, E*). GFP::POD-2

expression decreases in response to starvation (*Figure 4F*).

Expression of both genes is consistent with primarily somatic functions.

Germline *fasn-1* activity may derive from low levels of maternal expression,

though its presence is confirmed by the late arrest phenotype of *fasn-1(0)*

embryos derived from *+/fasn-1(0)* parents (*Figure 3E-G*).

INX-8 channels influence effects of overexpressed malonyl-CoA on

downstream FAS mutations

If *inx-8* is required for transit of mal-CoA from soma to germline through gap

junctions, the fertility of mutants downstream of *pod-2* in the FAS pathway could

be compromised by inclusion of INX-8(T239I D24N) into somatic hemichannels.

Expression of an *Ex[pod-2(+); sur-5::gfp]* simple array increases the brood size of

emb-8(hc69ts) at 22°C. This increase is dependent on INX-8 hemichannel

integrity—*inx-8(+)* homozygotes facilitate the increase more effectively than

+/inx-8(rf) heterozygotes (*Figure 6A*). Importantly, anti-INX-8 and anti-INX-22 co-

localization in *emb-8(hc69ts); +/inx-8(rf)* dissected gonads is indistinguishable

from wild type, implying that cell adhesion is not affected (*Figure 2E*).

In contrast to *emb-8(hc69ts)*, expression of *Ex[pod-2(+); sur-5::gfp]* in

fasn-1(g43ts) significantly <u>reduced</u> brood size (p<0.01), and brood size reduction

is more severe in homozygous *inx-8(+)* than heterozygous *+/inx-8(rf)*

backgrounds (ρ<0.05) (*Figure 6B*). This result likely relates to the mechanism of

FASN production of fatty acids—acetyl-CoA (FA chain initiation) and mal-CoA

(chain elongation) load at a common, non-selective acyl-CoA binding site and are

regarded as competitive inhibitors (*Cox and Hammes, 1983; Chang and*

Hammes, 1990). Increasing the ratio of mal-CoA to acetyl-CoA may inhibit

FASN-1 enzyme by interfering with chain initiation. Both wild-type and enhanced

levels of mal-CoA delivered through wild-type INX-8 hemichannels appear to

inhibit *fasn-1(g43ts)*; this mutant form of FASN may be especially sensitive to

inhibition by higher levels of mal-CoA. Importantly, heterozygous *+/inx-8(rf)*

channels ameliorate this inhibition, likely by reducing mal-CoA transfer.

Enhancement by *inx-8(+)* hemichannels of both a positive *(emb-8)* and a

negative *(fasn-1)* effect on FAS mutations downstream of overexpressed *pod-*

2(+) strongly argues that *inx-8* acts in the same pathway as *pod-2,* and is

consistent with gap junctions acting as conveyors of mal-CoA from soma to

germline (*Figure 6C*).

Genetic mosaic analyses associate *pod-2* sheath cell loss with the Pod

phenotype

To verify a somatic *pod-2* focus of action, genetic mosaic animals derived from

pod-2(0); Ex[pod-2(+); sur-5::gfp] were sought. 17/833 adult hermaphrodites

were identified that produced dead eggs but no viable progeny; these adults were

examined for cell lineage loss of *Ex[pod-2(+); sur-5::gfp]*. Animals fell into two

classes: (1) eight animals lost *Ex[pod-2(+); sur-5::gfp]* from most or all gonad

sheath cells, and (2) nine animals lost *Ex[pod-2(+); sur-5::gfp]* from the intestine

(*Figure 7A*).

Within the first class of mosaics, five animals lost the array in all sheath

cells in both gonad arms. Three other animals exhibited a difference in

phenotypic severity between gonad arms—embryos derived from one arm

developed to later multicellular stages *in utero* (*Figure 7B-D*). Differences were

coincident with loss of the array from all sheath cells in the more severely

affected arm, versus retention of the array in a single sheath cell of the better

arm. In subsequent screening, five more mosaics were identified that displayed

phenotypic disparity between the two gonad arms; in one example, one arm

produced defective embryos while the better arm gave rise to wild-type embryos

(producing ≥40 viable progeny). These results indicate that the focus of *pod-2(0)*

rescue is autonomous to each gonad arm, consistent with the interpretation that

pod-2 expression in the somatic sheath is essential to embryonic viability.

In subsequent screening for germline (P_4 –) mosaics, four animals were

identified that maintained the array in the soma but lost the array in all embryos

(*Figure 7A*). The terminal stages of embryonic development resembled those of

pod-2(0) embryos derived from *+/pod-2(0)* parents, i.e. embryos developed to 3-

fold or pretzel stages but failed to hatch (*Figure 7E-G*). These stages are more

advanced than terminal stages of many dead embryos derived from parents with

somatic (but not germline) loss of the rescuing array; such embryos fail to

differentiate (*Figure 7H*). This early developmental arrest resembles terminal-

stage Pod embryos from *fasn-1(0); Ex[fasn-1(+))*] (compare panels H and I in

Figure 7). We conclude that somatic *pod-2(+)* supplies sufficient levels of mal-

CoA to the germline not only to establish a wild-type 1-cell embryo, but also to

support continued development, presumably until zygotic *pod-2* expression

becomes adequate for completion of embryogenesis. These studies establish

that metabolites essential for embryonic development can be produced in the

somatic gonad and provided to the oocyte through gap junctions.

Intestinal loss of *pod-2*, but not *fasn-1*, results in a germline starvation

response

The other class of *pod-2(0)* genetic mosaics producing dead eggs in our original

screen lost *Ex[pod-2(+); sur-5::gfp]* from the intestine (E–) (*Figure 7A*). E(–)

mosaic animals were smaller than wild type, and gonad arms were severely

reduced in size (*Figure 8A*). Only the most proximal oocytes showed appreciable

growth. E(–) mosaic gonad arms resemble those seen in gonads undergoing the

oocyte starvation response, or adult reproductive diapause (ARD), a process

induced by starvation during the L4 larval stage proposed to maintain germline

stem cells for repopulation of the gonad upon the resumption of feeding (*Angelo*

and Van Gilst, 2009; Seidel and Kimble, 2011). During ARD, germline stem cells

cease dividing, the gonad shrinks due to extensive apoptosis of differentiated

germ cells, and growth is largely restricted to the most proximal oocyte.

The *C. elegans* intestine is rich with lipid droplets and the source of yolk,

and regarded as a major site of FAS (*Watts and Ristow, 2017*). A genetic

pathway in *C. elegans* required for intestinal apical polarity establishment leading

to proper lumen formation has been described; genes in this pathway, including

pod-2, encode enzymes required for glycosphingolipid synthesis (*Zhang et al.,*

2012). Lipid biosynthesis is required throughout larval growth, coincident with

membrane expansion. The identification of adult mosaics lacking *pod-2(+)* in the

intestine was therefore surprising. We subsequently identified two presumptive

P_1(–) *pod-2* mosaics, predicted to have lost *pod-2(+)* function in the intestine,

somatic gonad, germline, many muscle and hypodermal cells, among others

(*Figure 7A*). The most striking phenotype distinguishing these animals from E(–)

mosaics was an extremely constipated intestine (*Figure 8B*). The absence of this

constipation in E(–) mosaic animals implies that loss of *pod-2(+)* from the

intestine does not prevent the processing of bacteria through the intestinal lumen.

(Both P$_1$ mosaics showed tissue extruded from the rectum, which may be the

cause of constipation.)

Although *gfp::fasn-1(tn1782)* is not detected in the intestine, we asked if

there is a functional role for *fasn-1* similar to *pod-2*. As described above,

homozygous *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]* progeny from heterozygous

parents produce Pod embryos but exhibit no other phenotype; within this class of

progeny we screened for E(−) mosaics. Ten (of 800) such animals were identified

(*Figure 7A*); in stark contrast to *pod-2*, their gonad arms appeared normal. These

animals displayed no overt defects outside of Pod embryo production (*Figure*

8C). Because maternal FASN-1 rescues *fasn-1(0)* homozygotes to later

embryonic stages, an early intestinal function for *fasn-1* equivalent to *pod-2* could

be rescued maternally. However, the surprising conclusion from this mosaic

analysis is that there is no significant intestinal requirement for zygotic *fasn-1* in

larval and adult stages. The implication is that the primary function of mal-CoA

produced in the intestine is not as substrate for intestinal FASN-1.

Is malonyl-CoA required in other germline processes?

inx-8(rf) and *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* share delayed oogenesis and a

reduction in gap junction coupling, germ cell proliferation, and brood size. *inx-*

21(0) mutants are sterile, and *inx-22(0)* mutants are fertile (*Starich et al., 2014*);

therefore mal-CoA required for embryos must transit through gap junction

channels comprised of INX-14/INX-21 hemichannels, which are expressed

throughout the distal arm (*Figure 2B*). We asked if mal-CoA levels might

influence phenotypes preceding embryogenesis, such as oocyte production.

Supplying mal-CoA transiently by microinjection (0.5-10 mM) into *inx-*

8/9(0); Ex[inx-8(DTC+, Sh-)] L4 gonad arms failed to ameliorate phenotypes such

as Pod-embryo production or low brood size. Ectopic expression of GFP::POD-2

in the germline in *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* or *inx-8(rf)* genetic

backgrounds also failed to improve phenotypes, including delayed

gametogenesis (*Figure 7–figure supplement 1*). However, whether endogenous

mal-CoA levels are altered under these conditions is uncertain.

Effects of global mal-CoA levels were indirectly examined by comparing

higher or lower *pod-2(+)* copy number rescue of *pod-2(0)*; numbers of dead eggs

and larvae were summed to reflect total embryo output (Table 1). Lower copy

number rescue coincided with reduced embryo output (84% of wild-type) and

lower proportion of viable larvae within a brood (~37% of total embryos) (Table

1). Embryo output may therefore by impacted by *pod-2* expression; however,

somatic cells outside the gonad could be responsible. We therefore asked

specifically if FAS is required within the germline. Somatically-rescued *fasn-1(0);*

Ex[fasn-1(+); sur-5::gfp] embryo output is comparable to wild-type (94%);

however, embryo output in the double mutant *fasn-1(0); inx-8(rf); Ex[fasn-1(+);*

sur-5::gfp] is only ~50% of *inx-8(rf)* alone (Table 1). The FAS pathway may

therefore play a germline role affecting embryo output. Surprisingly, *inx-8(rf)*

partially suppresses the embryonic lethality of *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]*

(Table 1). Possibly the smaller gonad arms of *inx-8(rf)* may concentrate levels of

FASN-1 or mal-CoA. Alternatively, *Ex[fasn-1(+); sur-5::gfp]* germline expression

may be de-silenced in this genetic background. Further study is needed to

address the nature of this suppression.

Discussion

The genetic analysis of mutant *inx-8* soma-germline gap junction channels has led us to three conclusions that have significance beyond the *C. elegans* model system: (1) Mal-CoA produced in the somatic sheath is delivered through gap junctions to the germline; this mal-CoA is necessary to establish a proper one-cell embryo and to support continued embryogenesis. To our knowledge, this segregation of the cellular source of mal-CoA from the eventual cell type in which it is used as substrate is novel. As the rate-limiting step in FAS, mal-CoA production is an attractive control point to tie aspects of germ cell developmental progression to somatic control. (2) *pod-2* expression (and presumably mal-CoA production) is exceptionally robust in the intestine and essential for normal germline development. In the absence of intestinal POD-2, the germline exhibits phenotypic hallmarks of germline ARD despite the presence of a replete food source. By contrast, intestinal *fasn-1* expression is not similarly required. (A possible caveat is that maternal *fasn-1* in the intestine may be sufficient to reach adulthood, which would be surprising.) In light of the abundant expression level of *gfp::pod-2* in the intestine, and our finding that mal-CoA can be directionally transferred intercellularly, the question arises as to whether the intestine provides mal-CoA to other cells. Alternatively, a deficit of mal-CoA production in the intestine might initiate a systemic starvation response. Recently, *Gerisch et al.,* (2020) have shown that the HLH-30/TEFB transcription factor functions as a master regulator of the ARD state. *hlh-30* mutants fail to effect the system-wide

changes promoting long-term survival and recovery associated with ARD. ARD

longevity does not appear to rely on fatty acid metabolism. However, entry of *hlh-*

30 mutants into the ARD state appears normal; therefore effects of decreased

mal-CoA expression in the intestine may lie upstream of *hlh-30* activity. (3) A

pronounced delay in gametogenesis along with reduction in brood size results

from reduced soma-germline gap junction communication; it is unknown if this

delay derives from generally slower mitotic or meiotic progression, or from a

developmental block at a particular meiotic stage. The nature of potential

molecules passing through gap junctions that are responsible for timely meiotic

progression is unknown, but mal-CoA has thus far not been implicated.

Our results are consistent with other findings in *C. elegans* related to fatty

acid metabolism in oocytes. *pod-2*/ACC requires post-translational covalent

attachment of biotin for enzymatic activity, and *bpl-1* (*b*iotin *p*rotein *l*igase-1) is

the *C. elegans* ortholog of the necessary holocarboxylase synthetase (*Watts et*

al., 2018). Mal-CoA provided in the bacterial diet enables *bpl-1* mutants to

develop to adulthood, but *de novo* synthesis of polyunsaturated fatty acids

(PUFAs) is required for embryonic viability (*Watts et al., 2018*). Our results

indicate that mal-CoA supplied for PUFA synthesis (and other FAS) in the early

embryo is derived from the somatic sheath via gap junctions. PUFA-dependent

prostaglandin signaling from oocytes influences sperm guidance, and reduction

of *inx-14* function disrupts this signaling (*Kubagawa et al., 2006; Edmonds et al.,*

2011). We suggest that reduced levels of mal-CoA delivered to the oocyte

through mutant INX-14-containing gap junctions may reduce prostaglandin

synthesis and its signaling function for sperm migration.

At least one route outside of gap junctions may deliver mal-CoA to the

germline early in its development. The primordial germ cells Z2 and Z3 associate

with a pair of intestinal cells during embryogenesis; contacts are lost before

hatching (*Sulston et al., 1983; Abdu et al., 2016*). Therefore an early germline

requirement for mal-CoA could be fulfilled directly from the intestine. Loss of such

a function would be encompassed within the ARD-like phenotype seen in *pod-2*

E(–) mosaics.

In addition to its metabolic function in FAS, mal-CoA has established roles

as a regulatory molecule. It inhibits beta-oxidation of fatty acids by mitochondria.

Inhibitory mal-CoA derives from a second acetyl-CoA carboxylase (ACC2),

associated with the mitochondrial membrane. Manipulation of mal-CoA levels in

the mouse hypothalamus regulates food intake (Wolfgang and Lane, 2006).

Regulation is effected through the brain-specific CPT1-c, via a mechanism

distinct from the enzymatic activity of other CPT1 enzymes (*Sierra et al., 2008;*

Reilly and Mak, 2012). Little information regarding the *C. elegans* ACC2 is

known; somatically-delivered mal-CoA could potentially influence germline

mitochondrial function.

Malonylation as a post-translational lysine modification has recently been

described (*Peng et al., 2011*). Malonylation may inhibit target proteins, such as

the glycolytic enzymes glyceraldehyde 3-phosphate dehydrogenase and

pyruvate kinase (*Kulkarni et al., 2017*). Malonylation of mTOR reduces mTORC1

complex activity, decreasing endothelial cell proliferation and reducing

angiogenesis in a human umbilical model (*Bruning et al., 2018*). Together these

studies and others suggest a potentially pervasive role for mal-CoA in sensing

and responding to nutritional status. One trait associated with rapid tumor growth

is an increase in FAS, and FASN has been explored as a cancer therapy target

(*Menendez et al., 2007*). Possibly the ability of mal-CoA to transit through gap

junctions might sustain FAS during tumor progression. The discovery that mal-

CoA transits through gap junctions from soma to germline may be indicative of

other unappreciated intercellular roles for this important molecule.

Materials and methods

Strains

C. elegans strains were grown on standard NGM media (plus Nystatin, 6.25 mg/l) with *E. coli* strain OP50 as food source. Temperatures used for conditional mutants are indicated in main text. In addition to the wild-type strain N2, the following alleles were used: Chr. *I*: *fasn-1(g43ts); fasn-1(tn1762null); fasn-1(tn1782[gfp::fasn-1]); lin-41(tn1541[gfp::tev::s-tag::lin-41]);* Chr. *II*: *pod-2(ye60cs); pod-2(tn1691null); pod-2(tn1765[gfp::tev::3xFLAG::pod-2]);* Chr. *III*: *emb-8(hc69ts); dpy-17(e1295);* Chr. *IV*: *inx-8(tn1474null) inx-9(ok1502null); inx-8(tn1513) inx-9(ok1502); inx-8(tn1513 tn1555) inx-9(ok1502)*—a.k.a. *inx-8(rf)* in main text. Balancers used included: *tmC18 [dpy-5(tmIs1236)] I; hT2[bli-4(e937) let-?(q782) qIs48] (I:III); mIs11 IV*. Extrachromosomal arrays used included: *tnEx195[inx-8(+) inx-9(+); sur-5::gfp]; tnEx205[lag-2p::inx-8::gfp; str-1::gfp]; tnEx212[W09B6-pod-2(+)—20 ng/µl—hi copy; sur-5::gfp]; tnEx218[fasn-1(+); sur-5::gfp]; tnEx219[W09B6-pod-2(+)—4 ng/µl—lo copy; sur-5::gfp];* and *tnEx221[inx-8(tn1513 tn1555); str-1::gfp].*

Microinjection rescue of FAS mutants with extrachromosomal arrays

 The cosmid W09B6 was used at 50 ng/µl (2 lines) or 20 ng/µl (1 line) with *sur-5::gfp* (75 ng/µl) to rescue *pod-2(ye60cs). sur-5::gfp* expression was faint for both lines obtained with 50 ng/µl and these were not further characterized. *pod-2(tn1691null)* was rescued by microinjection using W09B6 at 4 ng/µl (5 lines) or

by crossing in *tnEx212*. Fosmids WRM0613bD07(10 ng/µl, or 50 ng/µl—1 line

that silenced) and WRM067cB03 (25 ng/ul) were used to attempt rescue of *emb-*

8(hc69ts). Numerous conditions using fosmids WRM0612bD08 and

WRM0614aH05 in simple or complex arrays were used to try to rescue *fasn-*

1(g43ts) without success. However, FASN enzyme functions as a dimer;

formation of dimers between mutant (*g43*) and wild-type monomers might inhibit

phenotypic rescue. After *fasn-1(tn1762null)* was generated, somatic but not

germline rescue was obtained using a *fasn-1* genomic PCR product forming

simple arrays (1 ng/µl + *sur-5::gfp-75* ng/µl, at least 2/6 lines segregated *fasn-*

1(0); Ex[fasn-1(+); sur-5::gfp(+)] animals that laid only Pod embryos), or a

complex array (1 ng/µl + 20 ng/µl *sur-5::gfp* + 100 ng/µl salmon sperm,1 line

from 30 injected animals) as described in the main text.

Recombinant DNA constructs for isolation of new *pod-2* and *fasn-1* alleles

Standard CRISPR/Cas-9 methods were used to generate deletions and GFP

insertions in *pod-2* and *fasn-1.* For deletions, 100-nt repair templates were

synthesized (Ultramers; Integrated DNA Technologies, Coralville, IA, USA) and used

in a *dpy-10(cn64)* co-conversion strategy previously described (*Arribere et al., 2014*).

Injection mixes included: pDD162 (Cas9), 50 ng/µl; 25 ng/µl (*pod-2*) or 10 ng/µl (*fasn-*

1) each single guide; Ultramer repair template, 500 nM; *dpy-10* single guide (pJA58),

25 ng/µl; *dpy-10(cn64)* Ultramer A2-2F-827, 500nM. Single guides were made by

annealing oligo pairs and cloning into pRB1017 *(Arribere et al., 2014)*: Deletions were

verified by PCR. The PCR fragment used to rescue *fasn-1(tn1762null)* was amplified

from genomic DNA using primers 5'-tacgtcaaacatccgtttgtcaacgtcac-3' and 5'-

ttcttctcctcgtcctttgatacgaagatc-3'.

Generation of GFP insertions into *pod-2* using pDD282 followed established

protocols (*Dickinson et al., 2015*). Injection mix included the single guide (10 ng/µl),

repair template (10 ng/µl), pDD162 (50 ng/µl), and *sur-5::gfp* (80 ng/µl). The insertion

was verified by sequencing PCR products generated with primers positioned outside

the homology arms.

For *gfp::fasn-1*, we were unable to isolate 3' insertions using the Dickinson

method (*Dickinson et al., 2015*). This method relies on isolation of candidates based

on the Roller phenotype, and because *fasn-1* has been implicated in molting, we

turned to a co-conversion approach (*Arribere et al., 2014*). A repair template was

generated that included the GFP sequence from pDD282, and *fasn-1* homology arms

on either side. The injection mix contained pDD162, 50 ng/µl; *fasn-1* single guide 10

ng/µl; *gfp::fasn-1* repair template, 20 ng/µl; *dpy-10* single guide (pJA58), 25 ng/µl;

dpy-10(cn64) Ultramer A2-2F-827, 500nM; and pCFJ190, 2.5 ng/µl. Rol candidates

were screened directly for GFP expression. The insertion was verified by sequencing

PCR fragments generated using primers placed outside of the repair template

sequence.

mex-5p::gfp::pod-2::tbb-2-3'UTR was constructed by Gibson assembly of

component PCR products representing the *mex-5* promoter, the "a" isoform of

gfp::pod-2, and the tbb-2-3'UTR to prevent germline RNA degradation.

Immunofluorescence

Antibody staining of dissected gonads was as previously described (*Starich et*

al., 2014). DIC and fluorescent images were acquired on a Zeiss (Thornwood,

NY, USA) motorized Axioplan 2 microscope with either a 40x Plan-Neofluar

(numerical aperture 1.3), a 63x Plan-Apochromatic (numerical aperture 1.4), or

100x PlanApochromatic (N.A. 1.4) objective lens using a AxioCam MRm camera

and AxioVision software (Zeiss).

Genetic mosaic analysis

To associate the somatic focus of action of *pod-2* for germline phenotypes, array

losses were characterized in animals of genotype *pod-2(1691null);*

tnEx212[W09B6-pod-2(+)—20 ng/μl—hi copy; sur-5::gfp] that were isolated on

the basis of sterility. Subsequently, potential candidate mosaics of interest were

prescreened directly at the dissecting microscope level for asymmetry in the

phenotypic severity seen between anterior and posterior gonad arms, and then

examined at the compound microscope level for loss of *sur-5::gfp* expression.

Potential *pod-2(tn1691null); tnEx219[W09B6-pod-2(+)—4 ng/μl—lo copy; sur-*

5::gfp] genetic mosaics were also prescreened in this manner. Intestinal (E–)

mosaics were prescreened using low-power fluorescence SMZ18 Nikon (Melville,

NY, USA) microscope with a Lumencor (Beaverton, OR, USA) SOLA Light

Engine, but for *pod-2* many of these candidate mosaics were seen to express

sur-5::gfp at very low levels, or in just a few intestinal cells, when examined at

high power. This was not an issue when isolating *fasn-1(tn1762null); tnEx218[fasn-1(+); sur-5::gfp]* E(−) mosaics.

Oligonucleotides used

fasn-1(tn1762) isolation

Single guide pairs:

5'-tcttggtatcttcccactcactga-3' plus 5'-aaactcagtgagtgggaagatacc-3', and

5'-cttggatgactttccttgcacga-3' plus 5'-aaactcgtgcaaggaaagtcatcc-3'.

Repair template:5'-

cgtatgacgtgattttttaatgtgataattactctgaaatatctcaaaacgtgcaaggaaagtcatcttcagtaacagtt

gaacacatcaacaggattat-3'

Verification of *tn1762*:

5'- ctgacaagacgacggacaactcagg-3' and 5'- ctacgtaaccgtaacggcatggcac-3'

pod-2(tn1691) isolation

Single guide pairs:

5'-tcttggaaaactgtccatatacaa-3' plus 5'-aaacttgtatatggacagttttcc-3', and

5'-tcttgacagaacaggaaaaagtgg-3' plus 5'-aaacccagttttttcctgttctgtc-3'.

Repair template: 5'-

ccgattttttgtgcaatttcagagcaatataagtatataaacgttattttcagagccatcacttttttcctgttctgtgatata

caagaaacctattcagtgccttatttgattgacgactac-3'

Verification of *tn1691*employed oligos:

5'-cgacgaagtaagtgagcctaattgtac-3' with 5'-ttgtcaaccaccttacagtggcatg-3'

gfp::pod-2(tn1765) isolation

Single guide pair:

5'-tcttgcagggttacggatcttgga-3' and 5'-aaactccaagatccgtaaccctgc-3

Homology arm oligo pairs:

5'-acgttgtaaaacgacggccagtcgccggcatctacctgtgtacctgacctgaccagattc-3' plus

5'-aactccagtgaacaattcttctcctttactcattgtgaatgccagtccaagatccgtaac-3';

5'-cgtgattacaaggatgacgatgacaagagagttgtaaacgggcaaaaaccagatatcag-3' plus

5'-tcacacaggaaacagctatgatatgtgagagtgaatgaactgctccatgctctc-3'

gfp::fasn-1(tn1782) isolation

Single guide pair:

5'-tcttgaaggatcgtccgattcgag-3' plus 5'-aaaactcgaatcggacgatccttc-3'

Primer used to introduce silent mutations to recognition site:

5'-gataatactggagagggctcatcagactcaagtggaacttgggagcgaatttcggac-3'

Homology flanking GFP:

5'-gggataataagtaggttctgacctcctctcccg-3' to 5'-gatccattcgtgcttcttcttgtcccgtgag-3'

Acknowledgements

We are grateful to our colleagues Gabriela Huelgas-Morales, Caroline Spike, Tatsuya Tsukamoto, and Jocelyn Shaw for discussions regarding experiments and comments on the manuscript. We would also like to thank Ross Johnson and David Hall for many years of sharing their knowledge and enthusiasm in all things concerning gap junctions. This work was supported by National Institutes of Health (NIH) grant GM57173 to DG. Some strains were provided by the Caenorhabditis Genetics Center, which is funded by grant P40OD010440 from the NIH Office of Research Infrastructure Programs.

Additional information

Funding

Funder	Grant reference number	Author
National Institutes of Health	GM57173	David Greenstein

Author contributions

TS conducted the experiments, prepared the figures, and wrote the manuscript. TS and DG designed and conceptualized the research study. DG obtained funding, supervised the study, and edited the manuscript.

References

Abdu Y, Maniscalco C, Heddleston, JM, Chew, T-L, Nance J. 2016.
Developmentally programmed germ cell remodelling by endodermal cell
cannibalism. *Nature Cell Biology* **18**:1302–1310. DOI: 10.1038/ncb3439, PMID:
27842058

Angelo G, Van Gilst MR. 2009. Starvation protects germline stem cells and
extends reproductive longevity in *C. elegans*. *Science* **326**:954–958. DOI:
10.1126/science.1178343, PMID: 19713489

Arribere JA, Bell RT, Fu BX, Artiles KL, Hartman PS, Fire AZ. 2014. Efficient
marker-free recovery of custom genetic modifications with CRISPR/Cas9 in
Caenorhabditis elegans. *Genetics* 198:837–846. DOI:
10.1534/genetics.114.169730, PMID: 25161212

Austin J, Kimble J. 1987. *glp-1* is required in the germ line for regulation of the
decision between mitosis and meiosis in *C. elegans*. *Cell* **51**:589–599. DOI:
10.1016/0092-8674(87)90128-0, PMID: 3677168

Austin J, Kimble J. 1988. Transcript analysis of *glp-1* and *lin-12*, homologous
genes required for cell interactions during development of *C. elegans*. *Cell*
58:565–571. DOI: 10.1016/0092-8674(89)90437-6, PMID: 2758467

Benenati G, Penkov S, Müller-Reichert T, Entchev EV, Kurzchalia TV. 2009. Two
cytochrome P450s in *Caenorhabditis elegans* are essential for the organization
of eggshell, correct execution of meiosis and polarization of the embryo.

Mechanisms of Development **126**:382–393. DOI: 10.1016/j.mod.2009.02.001, PMID: 19368796

Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE. 2000. The protein data bank. *Nucleic Acids Research* **28**:235–242. DOI: 10.1107/s0907444902003451, PMID: 10592235

Bruning U, Morales-Rodriguez F, Kalucka J, Goveia J, Taverna F, Queiroz KCS, Dubois C, Cantelmo AR, Chen R, Loroch S, Timmerman E, Caixeta V, Bloch K, Conradi LC, Treps L, Staes A, Gevaert K, Tee A, Dewerchin M, Semenkovich CF, Impens F, Schilling B, Verdin E, Swinnen JV, Meier JL, Kulkarni RA, Sickmann A, Ghesquière B, Schoonjans L, Li X, Mazzone M, Carmeliet P. 2018. Impairment of angiogenesis by fatty acid synthase inhibition involves mTOR malonylation. *Cell Metabolism* **28**:866–880. DOI: 10.1016/j.cmet.2018.07.019, PMID: 30146486

Chang S-I, Hammes GG. 1990. Structure and mechanism of action of a multifunctional enzyme: fatty acid synthase. *Accounts of Chemical Research* **23**:363–369. DOI: https://doi.org/10.1021/ar00179a003

Cox BG, Hammes GG. 1983. Steady-state kinetic study of fatty acid synthase from chicken liver. *Proceedings of the National Academy of Sciences USA* **80**:4233–4237. DOI: 10.1073/pnas.80.14.4233, PMID: 6576333

Dickinson DJ, Pani AM, Heppert JK, Higgins CD, Goldstein B. 2015. Streamlined genome engineering with a self-excising drug selection cassette. *Genetics* **200**:1035–1049. DOI: 10.1534/genetics.115.178335, PMID: 26044593

Edmonds JW, McKinney SL, Prasain JK, Miller MA. 2011. The gap junctional

protein INX-14 functions in oocyte precursors to promote *C. elegans* sperm

guidance. *Developmental Biology* **359**:47–58. DOI:

10.1016/j.ydbio.2011.08.014, PMID: 21889935

Gerisch B, Tharyan RG, Mak J, Denzel SI, Popkes-van Oepen T, Henn N, Antebi

A. 2020. HLH-30/TFEB is a master regulator of reproductive quiescence.

Developmental Cell **53**:1–14. DOI: 10.1016/j.devcel.2020.03.014, PMID:

32302543

Govindan JA, Cheng H, Harris JE, Greenstein D. 2006. G$\alpha_{o/i}$ and Gα_s signaling

function in parallel with the MSP/Eph receptor to control meiotic diapause in *C.*

elegans. *Current Biology* **16**:1257–1268. DOI: 10.1016/j.cub.2006.05.020,

PMID: 16824915

Henderson, ST, Gao D, Lambie EJ, Kimble J. 1994. *lag-2* may encode a

signaling ligand for the GLP-1 and LIN-12 receptors of *C. elegans*.

Development **120**:2913–2924. PMID: 7607081

Jaramillo-Lambert A, Fuchsman AS, Fabritius AS, Smith HE, Golden A. 2015.

Rapid and efficient identification of *Caenorhabditis elegans* legacy mutations

using Hawaiian SNP-based mapping and whole-genome sequencing. *G3*

5:1007–1019. DOI: 10.1534/g3.115.017038, PMID: 25740937

Johnston WL, Krizus A, Dennis JW. 2006. The eggshell is required for meiotic

fidelity, polar-body extrusion and polarization of the *C. elegans* embryo. *BMC*

Biology **4**:35. DOI: 10.1186/1741-7007-4-35, PMID: 17042944

Kelly WG, Montgomery MK, Fire A. 1997. Distinct requirements for somatic and germline expression of a generally expressed *Caenorhabditis elegans* gene. *Genetics* **146**:227–238. PMID: 9136012

Kubagawa HM, Watts JL, Corrigan C, Edmonds JW, Sztul E, Browse J, Miller MA. 2006. Oocyte signals derived from polyunsaturated fatty acids control sperm recruitment *in vivo*. *Nature Cell Biology* **8**:1143–1148. DOI: 10.1038/ncb1476, PMID: 16998478

Kulkarni RA, Worth AJ, Zengeya TT, Shrimp JH, Garlick JM, Roberts AM, Montgomery DC, Sourbier C, Gibbs BK, Mesaros C, Tsai YC, Das S, Chan KC, Zhou M, Andresson T, Weissman AM, Linehan WM, Blair IA, Snyder NW, Meier JL. 2017. Discovering targets of non-enzymatic acylation by thioester reactivity profiling. *Cell Chemical Biology* **24**:231–242. DOI: 10.1016/j.chembiol.2017.01.002, PMID: 28163016.

Lee K-Z, Kniazeva M, Han M, Pujol N, Ewbank, J. 2010. The fatty-acid synthase *fasn-1* acts upstream of WNK and Ste20/GCK-VI kinases to modulate antimicrobial peptide expression in *C. elegans* epidermis. *Virulence* **1**:113–122. DOI: 10.4161/viru.1.3.10974, PMID: 21178429

McCarter J, Bartlett B, Dang T, Schedl T. 1997. Soma-germ cell interactions in *Caenorhabditis elegans*: multiple events of hermaphrodite germline development require the somatic sheath and spermathecal lineages. *Developmental. Biology* **181**:121–143. DOI: 10.1006/dbio.1996.8429, PMID: 9013925

Menendez JA, Lupu R. 2007. Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nature Reviews Cancer* **7**:763–777. DOI: 10.1038/nrc2222, PMID: 17882277

Meyer RA, Laird DW, Revel J-P, Johnson RG. 1992. Inhibition of gap junctions and adherens junction assembly by connexin and A-CAM antibodies. *Journal of Cell Biology* **119**:179–189. DOI: 10.1083/jcb.119.1.179, PMID: 1326565

Mok DZL, Sternberg PW, Inoue T. 2015. Morphologically defined sub-stages of *C. elegans* vulval development in the fourth larval stage. *BMC Developmental. Biology* **15**: 26. DOI: 0.1186/s12861-015-0076-7, PMID: 26066484

Nadarajan S, Govindan JA, McGovern M, Hubbard EJ, Greenstein D. 2009. MSP and GLP-1/Notch signaling coordinately regulate actomyosin-dependent cytoplasmic streaming and oocyte growth in *C. elegans*. *Development* **136**: 2223–2234. DOI: 10.1242/dev.034603, PMID: 19502484

Olson SK, Bishop JR, Yates JR, Oegema K, Esko JD. 2006. Identification of novel chondroitin proteoglycans in *Caenorhabditis elegans*: embryonic cell division depends on CPG-1 and CPG-2. *Journal of Cell Biology* **173**:985–994. DOI: 10.1083/jcb.200603003 PMID, 16785326

Oshima A, Tani K, Fujiyoshi Y. 2016. Atomic structure of the innexin-6 gap junction channel determined by cryo-EM. *Nature Communications* **7**:1368. DOI: 10.1038/ncomms13681, PMID: 27905396

Peng C, Lu Z, Xie Z, Cheng Z, Chen Y, Tan M, Luo H, Zhang Y, He W, Yang K, Zwaans BM, Tishkoff D, Ho L, Lombard D, He TC, Dai J, Verdin E, Ye Y, Zhao

Y. 2011. The first identification of lysine malonylation substrates and its regulatory enzyme. *Molecular and Cellular Proteomics* **10**:M111.012658. DOI: 10.1074/mcp.M111.012658, PMID: 21908771

Rappleye CA, Paredez AR, Smith CW, McDonald KL, Aroian RV. 1999. The coronin-like protein POD-1 is required for anterior-posterior axis formation and cellular architecture in the nematode *Caenorhabditis elegans*. *Genes and Development* **13**:2838–2851. DOI: 10.1101/gad.13.21.2838, PMID: 10557211

Rappleye CA, Tagawa A, Le Bot N, Ahringer J, Aroian RV. 2003. Involvement of fatty acid pathways and cortical interaction of the pronuclear complex in *Caenorhabditis elegans* polarity. *BMC Developmental Biology* **3**:8. DOI: 10.1186/1471-213X-3-8, PMID: 14527340

Rappleye CA, Tagawa A, Lyczak R, Bowerman B, Aroian RV. 2002. The anaphase-promoting complex and separin are required for embryonic anterior-posterior axis formation. *Developmental Cell* **2**:195–206. DOI: 10.1016/s1534-5807(02)00114-4, PMID: 11832245

Reilly PT, Mak TW. 2012. Molecular pathways: tumor cells co-opt the brain-specific metabolism gene *CPT1C* to promote survival. *Clinical Cancer Research* **18**:5850–5855. DOI: 10.1158/1078-0432.CCR-11-3281, PMID: 22952346

Rose AS, Bradley AR, Valasatava Y, Duarte JM, Prlic A, Rose PW. 2018. NGL viewer: web-based molecular graphics for large complexes. *Bioinformatics* 34:3755–3758. DOI: 10.1093/bioinformatics/bty419, PMID: 29850778

Seidel HS, Kimble, J. 2011. The oogenic germline starvation response in *C. elegans*. *PloS ONE* **6**:e28074. DOI: 10.1371/journal.pone.0028074, PMID: 22164230

Sierra AY, Gratacós E, Carrasco P, Clotet J, Ureña J, Serra D, Asins G, Hegardt FG, Casals N. 2008. CPT1c is localized in endoplasmic reticulum of neurons and has carnitine palmitoyltransferase activity. *Journal of Biological Chemistry* **283**:6878–6885. DOI: 10.1074/jbc.M707965200, PMID: 18192268

Skerrett IM, Williams JBA. 2017. A structural and functional comparison of gap junction channels composed of connexins and innexins. *Developmental Neurobiology* **77**:522–547. DOI: 10.1002/dneu.22447, PMID: 27582044

Spike CA, Coetzee D, Eichten C, Wang X, Hansen D, Greenstein D. 2014. The NHL-TRIM protein LIN-41 and the OMA RNA-binding proteins antagonistically control the prophase-to-metaphase transition and growth of *C. elegans* oocytes. *Genetics* **198**:1535–1558. DOI: 10.1534/genetics.114.168831, PMID: 25261698

Starich TA, Hall DH, Greenstein D. 2014. Two classes of gap junction channels mediate soma-germline interactions essential for germline proliferation and gametogenesis in *C. elegans*. *Genetics* **198**:1127–1153. DOI: 10.1534/genetics.114.168815, PMID: 25195067

Sulston JE, Schierenberg E, White JG, Thomson JN. 1983. The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Developmental Biology* **100**: 64–119. DOI: 10.1016/0012-1606(83)90201-4, PMID: 6684600

Tagawa A, Rappleye CA, Aroian RV. 2001. *pod-2*, along with *pod-1*, defines a

new class of genes required for polarity in the early *Caenorhabditis elegans*

embryo. *Developmental Biology* **233**:412–424. DOI: 10.1006/dbio.2001.0234,

PMID: 11336504

Watts JS, Morton DG, Kemphues KJ, Watts JL. 2018. The biotin-ligating protein

BPL-1 is critical for lipid biosynthesis and polarization of the *Caenorhabditis*

elegans embryo. *Journal of Biological Chemistry* **293**:610–622. DOI:

10.1074/jbc.M117.798553, PMID: 29158261

Watts JL, Ristow M. 2017. Lipid and carbohydrate metabolism in *Caenorhabditis*

elegans. *Genetics* **207**:413–446. DOI: 10.1534/genetics.117.300106, PMID:

28978773

Whitten SJ, Miller MA. 2007. The role of gap junctions in *Caenorhabditis elegans*

oocyte maturation and fertilization. *Developmental Biol*ogy **301**:432–446. DOI:

10.1016/j.ydbio.2006.08.038, PMID: 16982048

Wolfgang MJ, Lane MD. 2006. The role of hypothalamic malonyl-CoA in energy

homeostasis. *Journal of Biological Chemistry* **281**:37265–37269. DOI:

10.1074/jbc.R600016200, PMID: 17018521

WormBase (www.wormbase.org).

Yochem J, Gu T, Han, M. 1998. A new marker for mosaic analysis in

Caenorhabditis elegans indicates a fusion between hyp6 and hyp7, two major

components of the hypodermis. *Genetics* **149**:1323–1334. PMID: 9649523

Zhang H, Abraham N, Khan LA, Hall DH, Fleming JT, Göbel V. 2012. Apicobasal

domain identities of expanding tubular membranes depend on

glycosphingolipid biosynthesis. *Nature Cell Biology* **13**:1189–1201. DOI:

10.1038/ncb2328, PMID: 21926990

Zhang Y, Foster JM, Nelson LS, Ma D, Carlow, CKS. 2005. The chitin synthase

genes *chs-1* and *chs-2* are essential for *C. elegans* development and

responsible for chitin deposition in the eggshell and pharynx, respectively.

*Developmental Biol*ogy **285**:330–339. DOI: 10.1016/j.ydbio.2005.06.037, PMID:

16098962

Figure Legends

Figure 1. Ovulation and oocyte development are delayed in *inx-8(rf)*. (A)

Diagram of adult *C. elegans* gonad. Posterior arm shows the positions of the

somatic distal tip cell (DTC) and five pairs of sheath cells (nuclei are numbered).

Anterior arm shows the underlying germline, approximate regions of germ cell

proliferation (Prolif.), meiotic pachytene stage (Pach.), and developing oocytes.

Embryos typically undergo some cell division in the uterus before being laid. Sp,

spermatheca; vul, vulva. (B) Compared to wild type (N2), ovulation in *inx-*

8(tn1513 tn1555rf) mutants (a.k.a. *inx-8[rf]*) is delayed ~18 hrs. Bars

representing 1-3 hr windows indicate approximate time of the fourth larval stage

(L4 to adult) molt, followed by time of appearance of the first fertilized embryo

ovulated into the uterus. (C) Onset of *gfp::lin-41* expression as a marker of

oocyte development is delayed in *inx-8(rf)*. Panels show DIC images of vulval

development (left), loop region of gonad arm (middle), and expression of *gfp::lin-*

41 (right). Wild-type hermaphrodites express *gfp::lin-41* by the approximate L4.7

sub-stage of vulval development (vul); *inx-8(rf)* hermaphrodites do not express

gfp::lin-41 until after vulval development is complete but before the L4-to-adult

molt. Null *inx-8(tn1474) inx-9(ok1502)* hermaphrodites, rescued for *inx-8::gfp*

expression in the DTC with *Ex[lag-2p::inx-8::gfp]*—a.k.a. *inx-8/9(0); Ex[inx-*

8(DTC+, Sh-)]—show a similar delayed onset of *gfp::lin-41* expression as *inx-*

8(rf). Bar, 20 μm.

Figure supplement 1. Predicted topological sites of amino acids mutated in *inx-*

8(tn1513 tn1555) inx-9(0).

Figure supplement table 1. *inx-8(rf)* exhibits reduced brood size and interacts

with FA synthesis pathway genes.

Figure supplement table 2. Reduced soma-germline gap junction coupling

delays gametogenesis.

Figure 2. INX-8(T239I D24N) compromises gap junction formation between

soma and germline. (A) Somatic INX-8 in wild-type (WT) gonads localizes to

puncta representing soma-germline gap junctions where DTC or sheath cells

overlie germ cells. (B) In WT distal gonad arms, gap junction puncta aggregate in

small clusters associated with each germ cell; in *inx-8(rf)* distal arms, gap

junction puncta localize more apically, where sheath dips down between germ

cells, outlining the cell (honeycomb pattern). Localization of INX-21 and INX-22

define the two different classes of germline gap junction hemichannels, both of

which include INX-14 subunits. (C) At the loop region, WT gap junction puncta

form large higher-order aggregates (arrows) which are absent in *inx-8(rf)*. (D) In

WT proximal arms, gap junction formation with INX-22-containing hemichannels

is extensive; in *inx-8(rf)*, INX-22 and INX-8(T239I D24N) are strongly expressed

but show little evidence of gap junction formation. "–1" indicates the most

proximal oocyte; *inx-8(rf)* hermaphrodites frequently show undersized oocytes at

this position as well (<). (E) Heterozygous *+/inx-8(rf)* shows little effect on gap

junction localization in an *emb-8(hc69ts)* background. Antibodies specific to INX-

8, INX-21, or INX-22 were used as indicated. DAPI-stained DNA in blue. Bar, 50

μm.

Figure supplement 1. INX-8(rf)::GFP can be endocytosed by oocytes.

Figure 3. (A) Predicted assignment of FAS mutants in a simplified FAS pathway.

Targeted deletions and GFP insertions are shown for *pod-2* (B) and *fasn-1* (C).

Predicted isoforms for POD-2 are indicated. (D-G) Terminal phenotypes of (D)

pod-2(0) and (E-G) *fasn-1(0)* mutant embryos from heterozygous *+/(0)* parents.

Arrows in (D) and (F) indicate differentiated anterior pharynx; arrow in (E)

indicates differentiated intestinal cells that formed gut granules. Bar, 20 μm.

Figure 4. GFP::POD-2 expression. (A) Expression in the head includes

hypodermis (carets) and excretory duct cell (arrow). Int, intestine. (B) Primary

expression in the tail includes hypodermis (carets) and intestine (Int). Left panel

DIC image, right panel GFP fluorescence in live animals. (C) Dissected animal

stained with anti-GFP antibody shows abundant intestinal expression (Int) and

strong expression in proximal sheath of exposed gonad arm (prox) that

diminishes in the distal arm. No germline expression is seen in the distal arm

where germ cells are revealed due to the absence of sheath coverage. Arrow

indicates CAN neuron, one of the few neurons essential for viability in *C.*

elegans. DAPI-staining of DNA in blue. (D, E) Earliest expression in embryos is

evident in developing hypodermis and intestine. (A, B, D, E) DIC images in left

panels, live GFP fluorescence in right panels. Bars, 20 μm, except (C) 200 μm.

(F) GFP::POD-2 is responsive to starvation. Top row, animals briefly washed and

transferred to bacteria-free NGM plates for 4 hours; lower row, animals similarly-

treated but fed.

Figure 5. GFP::FASN-1 expression. (A) GFP::FASN-1 is expressed throughout

the hypodermis (small arrows). Excretory duct cell expression is also indicated

(larger arrow). (B) Arrow indicates proximal gonad sheath expression. (C) Arrow

indicates developing vulva. (D) Earliest expression in developing embryos is

seen in hypodermis. DIC images in upper panels, live GFP fluorescence in lower

panels. Bars, 20 μm.

Figure 6. Wild-type INX-8 facilitates effects of *Ex[pod-2(+)]* overexpression on

conditional mutations downstream in the fatty acid synthesis (FAS) pathway.

Brood size was measured as the number of viable progeny produced. The

balancer *mIs11* provided *inx-8(+)* for all strains listed. (A) Embryonic lethality in

emb-8(ts) mutants at 22^{o}C is more severe in heterozygous *+/inx-8(rf)* than *inx-*

8(+) backgrounds *(1)*. Expression of *Ex[pod-2(+)]* in *emb-8(ts)* significantly

rescues embryonic lethality *(2)*. Rescue of *emb-8(ts)* by *Ex[pod-2(+)]* is also

dependent on *inx-8(+)* copy number *(3)*. (B) Expression of *Ex[pod-2(+)]* in *fasn-*

1(ts) mutants at 20^{o}C reduces brood size *(1)*. *fasn-1(ts); inx-8(+)* homozygotes

show greater embryonic lethality than *fasn-1(ts); (+)/inx-8(rf)* heterozygotes in the

absence *(2)* or presence *(3)* of *Ex[pod-2(+)]*. Averages indicated with dotted

lines. Probability was calculated using Student's 2-tailed *t*-test. (C) Enzymes

(purple) implicated in germline FAS. Malonyl-CoA transits from soma sheath to

germline through gap junction channels comprised of INX-8/9 (soma) and INX-

14/INX-21 (germline) hemichannels. Germline FASN-1 uses malonyl-CoA to

synthesize palmitate, which may be further extended and modified to produce

other long-chain fatty acids. EMB-8 (NADPH-dependent CYP450 reductase) and

POD-7/8 (CYP450s) are necessary to synthesize the lipid-rich (inner) layer of the

eggshell and support polarity establishment (*Rappleye et al., 2003; Benenati et*

al., 2009); the composition of lipid layer fatty acids is unknown. *de novo* PUFA

synthesis is required in the germline for embryonic viability, and is implicated in

sperm guidance (*Kubagawa et al., 2006; Watts et al., 201*8). The identities of

other long-chain fatty acids that may be synthesized in the germline are

unknown. PUFAs, polyunsaturated fatty acids; LFAs, long-chain fatty acids;

MUFA, monounsaturated fatty acids; pm, plasma membrane.

Source data 1. Viable progeny counts for the Figure 6 panels A, B.

Figure 7. Genetic mosaic analyses of *pod-2* reveal a mal-CoA requirement in

somatic gonad sheath. (A) Abbreviated cell lineage of *C. elegans*. Mosaics

characterized correspond to loss of *sur-5::gfp* expression in the germline (P$_4$),

intestine (E), or somatic gonad. Five *pod-2(0); Ex[pod-2(+); sur-5::gfp]* animals

lost the array from all somatic gonad sheath cells and produced no viable

embryos. (B) Eight animals showed differences in array retention in each gonad

arm with phenotypic consequences. Arrays were generated by microinjecting

cosmid W09B6 *[pod-2(+)]* at 4 ng or 20 ng/µl (low/high copy). (C) Uterus of non-

mosaic *pod-2(0); Ex[pod-2(+)—high copy]* hermaphrodite; developmental

progression of embryos from each gonad arm is comparable. (D) Mosaic *pod-*

2(0); Ex[pod-2(+)—high copy] with array lost from posterior gonad arm sheath

but retained in a single Sh4 cell of anterior arm. Anterior arm embryos have more

distinct eggshells and develop to later stages. (E-H) Embryonic development

depends on somatic levels of mal-CoA. Shown are terminal-stage *pod-2(0)*

embryos derived from parents of designated genotypes. (E) Pretzel-stage

embryo from *+/pod-2(0)*. (F) 3-fold stage embryo from *pod-2(0); Ex[pod-2(+)—*

low copy], with array maintained in soma but lost in the germline (P4–). (G)

Pretzel-stage embryo from P4(–) mosaic *pod-2(0); Ex[pod-2(+)—high copy]*. (H)

Embryo from mosaic *pod-2(0); Ex[pod-2(+)—low copy]* with uncharacterized

somatic loss of array, but array retained in the germline (P4+). This parent

produced exclusively dead embryos. (I) Typical terminal undifferentiated *fasn-*

1(0) Pod embryo from *fasn-1(0); Ex[fasn-1(+)]*. ant ute, anterior uterus; post ute,

posterior uterus; v, vulva; –1, most proximal oocyte. Bar, 20 µm.

Figure supplement 1. Expression of *gfp::pod-2* in the germline fails to rescue

inx-8(rf) phenotypes.

Figure 8. Mosaic loss from the intestine of *pod-2*, but not *fasn-1,* leads to an

ARD-like gonad arm. (A) E(–) genetic mosaic *pod-2(0); Ex[pod-2(+); sur-5::gfp]*

gonad arms (from same animal). Gonad arm on right partially outlined. (B) Two

focal planes of a presumptive $P_1(-)$ *pod-2(0); Ex[pod-2(+); sur-5::gfp]* mosaic with

constipated intestine. Upper plane (left) with partially outlined intestine; lower

plane (right) shows underlying distal gonad arm. int, intestine; p, pachytene

region; v, vulva. (C) E(–) genetic mosaic *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]*

gonad arm. –1, most proximal oocyte. Bars, 20 μm.

Table 1

Brood sizes suggest FA synthesis influences embryo output

Genotype	Larvae	Dead embryos	Total (embryo output)
Wild type (N2)	291±29 (n=33)	2±2	293±29
inx-8(rf)	108±40 (n=90)	11±6 (n=30)	119±42
inx-8/9(0); Ex[inx-8(DTC+,Sh-)]	2±2 (n=32)	19±13	20±14
pod-2(0); Ex[pod-2(+)] high copy (20 ng/µl)	164±50 (n=31)	118±41	282±59
pod-2(0); Ex[pod-2(+)] low copy (4 ng/µl)	91±24 (n=41)	156±37	247±50
fasn-1(0); Ex[fasn-1(+)	0.05 (n=80)	276±18 (n=16)	276±18
fasn-1(0); inx-8(rf); Ex[fasn-1(+)]	17±15 (n=47)	43±27	60±37

Source data 1. Brood counts for genotypes shown in Table 1.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

intestine
9x E-
10x E-

germline
4x P₄-

uterus+spermatheca

DTC DTC

post.
sheath
(x5)

post.
sheath
(x5)

ant.
sheath
(x5)

ant.
sheath
(x5)

uterus+spermatheca

pod-2(+) losses
fasn-1(+) losses

5 complete, 8 partial sheath losses

B

	Embryo quality/arm	
	better	**worse**
pod-2(0);	Sh1+	Sh-
Ex[high copy]	Sh3+	Sh-
	Sh3+	Sh-
	Sh4+	Sh-
	Sh4+	Sh-
Ex[low copy]	Sh1+, 2x Sh4+	Sh-
	Sh1+, Sh5+	Sh5+
	Sh5+ (fertile)	Sh-

Figure 7

Figure 8

Figure 1–figure supplement table 1

inx-8(rf) exhibits reduced brood size and interacts with FA synthesis pathway genes

Genotype	Brood size	Germ cells per gonad arm
20°C		
N2 (wild type)	291 ± 29 (n=33)	≥1000[1]
inx-8(rf)[2]	108 ± 40 (n=90)	322±18 (n=3; day 2 adult)
22°C		
inx-8(+)[3]	266 ± 40 (n=32)	
+/inx-8(rf)	255 ± 36 (n=32)	
fasn-1(g43ts); inx-8(+)	115 ± 41 (n=11)	
fasn-1(g43ts); +/ inx-8(rf)	114 ± 47 (n=11)	
fasn-1(g43ts); inx-8(rf)	4.5 ± 2.3 (n=12)	
emb-8(hc69ts); inx-8(+)	95 ± 50 (n=15)	
emb-8(hc69ts); +/inx-8(rf)	35 ± 37 (n=15)	
emb-8(hc69ts); inx-8(rf)	11 ± 9 (n=15)	

[1]_McCarter et al., 1997._
[2]Full genotype _inx-8(tn1513 tn1555) inx-9(ok1502)._ Approximately 4% of animals are infertile in one or both gonad arms (n=168).
[3]_inx-8(+)_ provided on the balancer _mIs11_ IV.
Source data 1. Brood size data for Figure 1–figure supplement table 1.

Figure 1–figure supplement table 2

Reduced soma-germline gap junction coupling delays gametogenesis

(a) *gfp::lin-41* **expression at L4 vulval sub-stages**

Genotype	L4.4	L4.5	L4.6	L4.7	L4.8	L4.9	L4.9< < molt
gfp::lin-41; inx-8(+)	1/11		3/3	5/6	5/6	4/4	
gfp::lin-41; inx-8(rf)		0/2	0/3	0/6		0/2	16/37
gfp::lin-41; inx-8/9(0); Ex[inx-8(DTC+,Sh-)]	0/11	0/3		0/2	0/1	0/3	6/14

(b) sperm appearance at L4 sub-stages

Genotype	L4.4	L4.5	L4.6	L4.7	L4.8	L4.9	L4.9< < molt
N2	0/9	0/2	0/5	4/11	1/2	1/2	6/6
inx-8(rf)	0/3		0/1	0/2	0/3	0/1	1/10

Figure 1–figure supplement 1

Figure 2–figure supplement 1

Figure 7–figure supplement 1

 Legends to Figure Supplements

Figure 1–figure supplement 1. Predicted topological sites of amino acids

mutated in *inx-8(tn1513 tn1555) inx-9(0)*. (A) Atomic structure of the *C. elegans*

homomeric, homotypic INX-6 gap junction full channel, determined by *Oshima et*

al., (2016) (PDB I.D. 5H1R, rcsb.org (*Berman et al., 2000*), viewed with NGL

viewer (*Rose et al., 2018*). Octameric hemichannels associate via extracellular

loops to form an enclosed channel in the region classically identified by TEM as

the "gap" between closely aligned plasma membranes. cyto, cytoplasmic

membrane face; p.m., plasma membrane. (B-F) Model for INX-8 hemichannel

based on homology alignment to INX-6 (aquaria.ws). (B) Cytoplasmic view. (C)

Single subunit highlighted. In wild type this hemichannel is comprised of both

INX-8 and INX-9 subunits, and associates with germline hemichannels

comprised of INX-14+INX-21 or INX-14+INX-22. (D) Side view of hemichannel.

Beta-strands (yellow) in the extracellular loops contribute to walling off the pore.

(E) Single subunit highlighted. Kinked region (gray arrow) allows the cytoplasmic

structure of the subunits (cytoplasmic loop and C-terminus) to form a dome at the

pore entry. Amino termini (N-term) line the pore as a funnel; a loop in the amino

terminus allows for an interaction with residues in the dome (*Oshima et al.,*

2016). D24 in INX-8 (red arrow) lies near the D21 residue predicted by homology

with INX-6 (D25) to interact with the cytoplasmic dome. (F) INX-8 T239 is

predicted to lie adjacent to a beta-strand contributing to hemichannel-

hemichannel association (red arrow). (To expose the extracellular loop region in

relation to (E), the aspect of view is rotated towards the viewer and angled up to

the left.)

Figure 2–figure supplement 1. INX-8(rf)::GFP can be endocytosed by oocytes.

(A) *inx-8(+)::gfp* expressed in the somatic gonad sheath is endocytosed during

ovulation and appears as distinct fluorescent puncta in early embryos. (b) *inx-*

8(rf)::gfp expressed on an extrachromosomal array in *inx-8(0) inx-9(0)* animals is

endocytosed by ovulating oocytes. (c) *inx-8::gfp* expressed by the DTC in *inx-*

8/9(0); Ex[inx-8(DTC+,Sh-)] is not observed in early embryos. Bar, 20 µm.

Figure 7–figure supplement 1. Expression of *gfp::pod-2* in the germline fails to

rescue *inx-8(rf)* phenotypes. (**A**) Young (first-day) adult *inx-8(rf)* gonad arm. (B,

C) Young adult *inx-8(rf)* gonad arm expressing *mex-5p::gfp::pod-2::tbb-2-3'UTR*.

The *pod-2* isoform in this construct corresponds to *gfp::pod-2* isoform "a" in

Figure 3B. This isoform encompasses sequence homology to all of the

enzymatic sites in other ACC multi-functional enzymes. Sp, spermatheca; -1,

most proximal oocyte; bar, 50 µm.