Gap junctions deliver malonyl-CoA from soma to germline to support

embryogenesis in *Caenorhabditis elegans*

Todd A. Starich[1,2] and David Greenstein[1]

[1]Department of Genetics, Cell Biology and Development, University of

Minnesota, Minneapolis, Minnesota 55455 USA

[2]Email: stari001@umn.edu

Abstract

Gap junctions are virtually ubiquitous in metazoans and play critical roles

in many important biological processes, including electrical conduction and

development. Despite this ubiquity, only a small number of molecules passing

through gap junction channels have been definitively linked to specific functions.

We isolated gap junction channel mutants that reduce junctional coupling

between the soma and germ cells in the *C. elegans* gonad. We use genetic

methods to show that malonyl-CoA, the rate-limiting substrate for fatty acid

synthesis (FAS), is produced in the soma and delivered through gap junctions to

the germline; there it is used by fatty acid synthase to critically support embryonic

development. Tissue separation of malonyl-CoA production from its site of

utilization facilitates somatic control of germline development and reproduction.

Further, our results suggest that this metabolic outsourcing may be a strategy by

which the soma communicates nutritional status to the germline.

Gap junctions are clusters of intercellular channels allowing direct passage

of small molecules between coupled cells. They are widespread in multicellular

organisms, though different gene families encode the protein subunits in

invertebrates (innexins) and vertebrates (primarily connexins)[1]. While the

capability of small molecules to pass through gap junctions can be shown readily

in cultured cells, identification of essential molecules that must pass through

channels *in vivo* is challenging. Additionally, disruption of gap junction coupling

can affect cell adhesion[2], and resultant defects in intercellular communication

may not necessarily relate to the passage of molecules through junctional

channels.

In *C. elegans*, germ cells are coupled throughout their development to the

somatic gonad by gap junctions[3]. Germ cells progress in assembly-line fashion

through the early stages of meiosis (Fig. 1a). Gonadal gap junctions have been

implicated in germ cell proliferation, oocyte growth, regulation of oocyte meiotic

maturation, early embryogenesis and sperm guidance[3-7]. Gap junction

hemichannels (half-channels) in the somatic gonad are comprised of the recently

duplicated innexins INX-8 and INX-9; apposing germ cell hemichannels are

heteromeric and consist of INX-14 in conjunction with either INX-21 or INX-22

(Supplementary Fig. 1). Germ cells fail to proliferate in *inx-8(0) inx-9(0)* mutants,

at a point upstream of, or parallel to, GLP-1/Notch activation of germ cells by

Delta class ligands LAG-2 and APX-1. These ligands are produced in the distal

tip cell (DTC) that maintains the proliferative stem cell niche[4,8-10].

Germ cell proliferation can be rescued in *inx-8(0) inx-9(0)* mutants to half

the level of wild type by driving *inx-8* expression in the DTC with the *lag-2*

promoter[3]. In these animals, gap junctions between soma and germline are

established in the distal gonad arm, but ostensibly not in the proximal arm; many

resultant embryos exhibit characteristics of the Pod phenotype (Polarity and

Osmotic sensitivity Defect)[11], including eggshell and permeability barrier

deficiencies, cytokinesis failures, and polar body extrusion defects. Other genes

with mutations producing Pod embryos include those involved in eggshell

synthesis, fatty acid synthesis (FAS), the APC/separase pathway, and a coronin-

like protein[11-18].

Molecules moving through soma-germ cell gap junctions necessary for

proliferation and early embryogenesis are unknown. We use genetic methods to

show that malonyl-CoA (mal-CoA) is produced by *pod-2* (acetyl-CoA

carboxylase, ACC) in the soma. Transfer of mal-CoA to the germline, for use by

fasn-1 (fatty acid synthase, FASN), is impacted by the integrity of INX-8-

containing channels under conditions in which gap junction formation appears

normal. Rescue of the Pod phenotype in *pod-2* genetic mosaics indicates a

somatic sheath focus of action. Our results support the model that mal-CoA

transits from soma to germline through gap junction channels.

Results

Gametogenesis is delayed by the inx-8(tn1513 tn1555) reduction of function (rf)

mutation

We previously isolated (in an *inx-9* null background) a strong loss-of-

function *inx-8* allele *(tn1513)* with sufficient residual function to contribute to

channel formation and cell adhesion but not to support germ cell proliferation[3].

tn1513 encodes a T239I change in the second extracellular loop of the innexin

monomer, predicted to be involved in docking with octameric hemichannels on

apposing cells (Supplementary Fig. 1)[19]. Using this allele in a genetic screen for

restoration of germ cell proliferation, six *inx-8* intragenic suppressor mutations

were isolated (to be described elsewhere). One of these, *tn1555*, encodes a

D24N change in the amino terminus (Supplementary Fig. 1), near a site that may

associate with a cytoplasmic dome structure surrounding the hemichannel outer

pore[19]. *inx-8(tn1513 tn1555rf)*--for brevity to be referred to as *inx-8(rf)*--partially

restored germ cell numbers and fertility to one-third the level of wild type

(Supplementary Table 1). *inx-8(rf)* is rescued by an *inx-8(+)* extrachromosomal

array, and rescue of *inx-8(0) inx-9(0)* with an *inx-8(rf)::gfp* array mimics the

phenotype of the suppressor mutant; therefore observed phenotypes can be

attributed to *inx-8(rf)* and not to other possible background mutations.

There is an ~18 hr delay to the onset of fertilization in *inx-8(rf)* compared

to wild type, as determined by relating the timing of first ovulation to the L4 molt

(L4-to-adult transition) (Fig. 1b). This delay correlates with late expression of

gfp::lin41 in relation to L4 sub-stages of vulval development[20]; LIN-41 is the

earliest known marker of oogenesis[21]. In wild type at $20^{\circ}C$, *gfp::lin41* appears by

sub-stage L4.7 (Fig. 1c); in *inx-8(rf)*, *gfp::lin41* is expressed after completion of

vulval morphogenesis but prior to the L4 molt (Fig. 1c; Supplementary Table 2).

Spermatogenesis is also delayed—wild-type sperm appear as early as the L4.7

stage, but *inx-8(rf)* sperm first appear after completion of vulval morphogenesis

(Supplementary Table 2). Once started, egg-laying in *inx-8(rf)* continues apace

such that ~99% of progeny are laid within a 96-hour window (3144/3179 larvae;

n=25), similar to wild type (5847/5929 larvae; n=21).

inx-8(rf) gap junction formation is more severely affected in proximal gonad arms

99   To examine expression of INX-8(T239I D24N) mutant innexins encoded

by *inx-8(rf)*, specific antibodies were used to label somatic (anti-INX-8) or the two

classes of germline (anti-INX-21 or anti-INX-22) hemichannels. In wild-type

hermaphrodites, clusters of gap junction puncta form between somatic DTC or

sheath cells and each underlying germ cell (Fig. 2a). In distal arms of wild type,

loosely clustered sheath-germ cell gap junctions form more basally on germ cells;

in *inx-8(rf)* distal arms, both INX-21 and INX-22 puncta tend to localize more

laterally, where the sheath cells dip down between adjacent germ cells (Fig. 2b).

In the loop region (transition from meiotic pachytene to diplotene), large, higher-

order clusters of gap junction puncta form in wild-type animals; these are absent

in *inx-8(rf)* (Fig. 2c). In wild-type proximal arms (diakinesis), developing oocytes

exhibit extensive formation of INX-22 gap junction puncta, while INX-21 puncta

become faint; in *inx-8(rf)* proximal arms, INX-22 and INX-8 are strongly but

diffusely expressed, with little co-localization evident (Fig. 2d). These results

suggest there may be little gap junction coupling in *inx-8(rf)* proximal arms. We

showed previously that sheath-oocyte gap junctions are endocytosed by the

oocyte during ovulation[3]. Embryos from *inx-8(0) inx-9(0)* rescued with *inx-*

8(rf)::gfp show internalized GFP puncta, confirming that *inx-8(rf)* establishes at

least some sheath-oocyte gap junctions in the proximal arm (Supplementary Fig.

2).

Reduced proximal arm expression of gap junctions is seen in *inx-8(0) inx-*

9(0); Ex[lag-2p::inx-8::gfp] animals—simplified to *inx-8/9(0); Ex[inx-8(DTC+, Sh-*

)]–which are rescued for germ cell proliferation by driving expression of INX-

8::GFP in the DTC, but not sheath (Sh), using the *lag-2* promoter[3]. These

animals produce primarily dead embryos, many with Pod characteristics (Table

1). In contrast to *inx-8(rf)*, these embryos do not show evidence of *inx-8::gfp*

endocytosis, consistent with absence of proximal arm sheath-germline gap

junction coupling (Supplementary Fig. 2). *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]*

animals also show delayed onset of *gfp::lin41* expression, similar to *inx-8(rf)* (Fig.

1c, Supplementary Table 2).

We conclude that delayed gametogenesis in *inx-8(rf)* mutants arises from

compromised soma-germline gap junction communication, due to altered channel

properties that influence the formation and localization of gap junctions,

especially in the loop and proximal arm. These regions are additionally implicated

because DTC expression of wild-type INX-8 fails to rescue *inx-8(rf)* phenotypes

(n>20).

inx-8(rf) interacts with FAS mutants

Because *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* produces Pod-like embryos, and

inx-8(rf) shows reduced proximal arm gap junction coupling similar to *inx-8/9(0);*

Ex[inx-8(DTC+, Sh-)], possible genetic interactions between *inx-8(rf)* and other

Pod mutants were explored (Supplementary Table 1). Conditional alleles of

genes in the FAS pathway (Fig. 3a) were used to construct double mutants with

inx-8(rf). Alleles included: *pod-2(ye60)*, a cold-sensitive (*cs*) allele of the *C.*

elegans acetyl-CoA carboxylase ACC Type 1[16]; a temperature-sensitive (*ts*)

allele of fatty acid synthase *fasn-1(g43)*[22]; and *emb-8(hc69)*, a *ts* allele of an

NADPH-cytochrome-P450 reductase, likely necessary for reduction of CYP-450

enzymes involved in fatty acid modifications[14,16]. Possible interaction between

pod-2(ye60cs) and *inx-8(rf)* was difficult to assess–*ye60cs* grown at 15°C was

quite sensitive to brief handling at room temperature. At a partially restrictive

temperature (22°C), *fasn-1(g43ts); inx-8(rf)* and *emb-8(hc69ts); inx-8(rf)* double

mutants exhibited markedly reduced brood sizes. Similar reductions in brood size

were also apparent in *emb-8(hc69ts); +/inx-8(rf)*, suggesting *emb-8(hc69ts)* might

be highly dose-sensitive to *inx-8(+)* levels (Supplementary Table 1).

FAS pathway shifts from soma to germline after pod-2

Genetic interactions between *inx-8* and FAS genes could represent activity

in the same or parallel pathways. If gap junctions are positioned in the FAS

pathway, it might be possible to find a step within the pathway where gene

function shifts from a somatic to a germline focus. *C. elegans* is particularly

amenable to such an investigation because simple (highly repetitive)

extrachromosomal arrays used to rescue gene function tend to express well in

the soma, whereas arrays with more sequence complexity are generally required

to prevent silencing of multi-copy gene expression in the germline[23].

Previous characterization of *pod-2* implied rescue by a simple array[16]; we

likewise found that *pod-2(ye60cs)* is rescued by simple extrachromosomal arrays

at 15°C (see Methods). In contrast, 2 fosmids encoding *emb-8*, used individually

to generate simple arrays, yielded 65 transformed lines, only one of which

rescued *emb-8(hc69ts)* at 25°C; the array in this line was eventually silenced. For

fasn-1, two fosmids used to generate 93 simple array lines and 20 complex array

lines failed to rescue *fasn-1(g43ts)*. These results were consistent with a model in

which FAS affecting embryonic development requires *pod-2* in the soma, but

shifts to a germline requirement for *fasn-1* and *emb-8*.

To avoid misrepresentation of rescue results due to the special nature of a

conditional allele, null alleles of *pod-2(tn1691)* and *fasn-1(tn1762)*–simplified to

(0)–were generated using CRISPR-Cas9 methodology (Fig. 3b,c)[24]. *pod-2(0)*

embryos from *pod-2(0)/+* parents arrest late in embryogenesis (pretzel stage;

Fig. 3d); most *fasn-1(0)* embryos from *fasn-1(0)/+* parents shows signs of

differentiation, and many arrest late in embryogenesis; occasional hatchlings

emerge with probable hypodermal defects (see below). *pod-2(0)* was rescued by

simple extrachromosomal arrays, supporting probable somatic function. Using a

PCR-amplified genomic fragment we were able to rescue somatic but not

germline function of *fasn-1(0)* with a simple or complex array (see Methods). A

genetically-balanced line with a complex array, *hT2/ fasn-1(0)*; *Ex[fasn-1(+); sur-*

5::gfp(+)], segregates healthy homozygous *fasn-1(0); Ex[fasn-1(+); sur-5::gfp(+)]*

adults that lay almost exclusively Pod embryos (Table 1). (*sur-5::gfp* is a co-

injection marker expressed in somatic cell nuclei; corresponding

extrachromosomal arrays can be lost mitotically to generate mosaics—see

below[25].) Gonad arms appear normal, and animals lay wild-type numbers of

defective embryos (Table 1); many of these display polarity defects (4/14

symmetric first divisions) and exhibit extra nuclei in early embryos (Fig. 3e-h).

Embryos typically arrest prior to morphogenesis. We conclude this *fasn-1(+)* PCR

fragment rescues somatic *fasn-1* function and is either silenced in the germline or

lacks an element necessary for germline expression; in either case a germline

requirement is indicated. In sum, rescue results are consistent with a fertility

requirement for *pod-2* in the somatic gonad and *fasn-1* in the germline. This

conclusion is supported by genetic mosaic and gene expression data described

below. If *inx-8* gap junctions act in this pathway, it would position these junctions

between *pod-2* and *fasn-1* as possible mediators of mal-CoA transfer from soma

to germline.

GFP::POD-2 and GFP::FASN-1 are tissue restricted

Endogenous, homozygous-viable, fertile GFP-labeled alleles of *pod-2* and

fasn-1 were generated to determine if patterns of localization are consistent with

predicted foci of action. The GFP moiety in *gfp::pod-2(tn1765)* is expected to

label all 3 predicted isoforms (Fig. 3b)[26]. GFP::POD-2 expression is strongest in

the intestine, and prominent in the hypodermis, gonadal sheath (especially

proximal sheath), CAN neuron, and excretory duct (Supplementary Fig. 3).

GFP::POD-2 expression decreases in response to starvation (Supplementary

Fig. 3).

gfp::fasn-1(tn1782) labeled at the amino terminus (Fig. 3c) is expressed

in the hypodermis, gonad sheath (esp. proximal) and excretory duct; intestinal

and CAN neuron expression are not evident (Supplementary Fig. 4). This

expression pattern partially overlaps with that of a *fasn-1* promoter::GFP reporter

which was expressed in the intestine[27]. WormBase[26] predicts a single FASN-1

isoform, but a second in-frame ATG lies in Exon 2; intestinal FASN-1 could be

derived from translational initiation at this site, which would not be revealed in our

construct. Attempts to GFP-label FASN-1 downstream of this second ATG, or at

the C-terminus, were unsuccessful. Neither *gfp::pod-2* nor *gfp::fasn-1* is

expressed in the DTC.

Germline expression for either GFP-tagged gene was not detected. Anti-

GFP staining of dissected gonads failed to detect germline GFP expression

(Supplementary Fig. 3). The earliest detectable GFP::FASN-1 in embryos

appears in developing hypodermis; GFP::POD-2 is prominent in developing

hypodermis and intestine (Supplementary Fig. 3,4). Expression of both genes is

consistent with primarily somatic functions. Germline *fasn-1* activity may derive

from low levels of expression, though its presence is confirmed by the late arrest

phenotype of *fasn-1(0)* embryos derived from *+/fasn-1(0)* parents

(Supplementary Fig. 4, e-h).

inx-8 channels influence effects of overexpressed malonyl-CoA on downstream

FAS mutations

If *inx-8* is required for transit of mal-CoA from soma to germline through

gap junctions, the fertility of mutants downstream of *pod-2* in the FAS pathway

could be compromised by inclusion of INX-8(T239I D24N) into somatic

hemichannels. Expression of an *Ex[pod-2(+); sur-5::gfp]* simple array increases

the brood size of *emb-8(hc69ts)* at 22^{o}C. This increase is dependent on INX-8

hemichannel integrity—*inx-8(+)* homozygotes facilitate the increase more

effectively than *+/inx-8(rf)* heterozygotes (Fig. 4a). Importantly, anti-INX-8 and

anti-INX-22 co-localization in *emb-8(hc69ts); +/inx-8(rf)* dissected gonads is

indistinguishable from wild type, implying that cell adhesion is not affected (Fig.

2e).

In contrast to *emb-8(hc69ts)*, expression of *Ex[pod-2(+); sur-5::gfp]* in

fasn-1(g43ts) significantly <u>reduced</u> brood size ($p < 0.01$), and brood size reduction

is more severe in homozygous *inx-8(+)* than heterozygous *+/inx-8(rf)*

backgrounds ($\rho < 0.05$) (Fig. 4b). This result likely relates to the mechanism of

FASN production of fatty acids--acetyl-CoA (FA chain initiation) and mal-CoA

(chain elongation) load at a common, non-selective acyl-CoA binding site and are

regarded as competitive inhibitors[28,29]. Increasing the ratio of mal-CoA to acetyl-

CoA may inhibit FASN-1 enzyme by interfering with chain initiation. Both wild-

type and enhanced levels of mal-CoA delivered through wild-type INX-8

hemichannels appear to inhibit *fasn-1(g43ts)*; this mutant form of FASN may be

especially sensitive to inhibition by higher levels of mal-CoA. Importantly,

heterozygous *+/inx-8(rf)* channels ameliorate this inhibition, likely by reducing

mal-CoA transfer.

Enhancement by *inx-8(+)* hemichannels of both a positive *(emb-8)* and a

negative *(fasn-1)* effect on FAS mutations downstream of overexpressed *pod-*

2(+) strongly argues that *inx-8* acts in the same pathway as *pod-2*, and is

consistent with gap junctions acting as conveyors of mal-CoA from soma to

germline (Fig. 4c).

Genetic mosaic analyses associate pod-2 sheath cell loss with the Pod

phenotype

264   To verify a somatic *pod-2* focus of action, genetic mosaic animals derived

from *pod-2(0); Ex[pod-2(+); sur-5::gfp]* were sought. 17/833 adult hermaphrodites

were identified that produced dead eggs but no viable progeny; these adults were

examined for cell lineage loss of *Ex[pod-2(+); sur-5::gfp]*. Animals fell into two

classes: 1) eight animals lost *Ex[pod-2(+); sur-5::gfp]* from most or all gonad

sheath cells, and 2) nine animals lost *Ex[pod-2(+); sur-5::gfp]* from the intestine

(Fig. 5a).

271   Within the first class of mosaics, five animals lost the array in all sheath

cells in both gonad arms. Three other animals exhibited a difference in

phenotypic severity between gonad arms—embryos derived from one arm

developed to later multicellular stages *in utero* (Fig. 5b-d). Differences were

coincident with loss of the array from all sheath cells in the more severely

affected arm, versus retention of the array in a single sheath cell of the better

arm. In subsequent screening, five more mosaics were identified that displayed

phenotypic disparity between the two gonad arms; in one example, one arm

produced defective embryos while the better arm gave rise to wild-type embryos

(producing ≥40 viable progeny). These results indicate that the focus of *pod-2(0)*

rescue is autonomous to each gonad arm, consistent with the interpretation that

pod-2 expression in the somatic sheath is essential to embryonic viability.

In subsequent screening for germline (P_4 –) mosaics, four animals were

identified that maintained the array in the soma but lost the array in all embryos

(Fig. 5a). The terminal stages of embryonic development resembled those of

pod-2(0) embryos derived from *+/pod-2(0)* parents, i.e. embryos developed to 3-

fold or pretzel stages but failed to hatch (Fig. 5e-g). These stages are more

advanced than terminal stages of many dead embryos derived from parents with

somatic (but not germline) loss of the rescuing array; such embryos fail to

differentiate (Fig. 5h). This early developmental arrest resembles terminal-stage

Pod embryos from *fasn-1(0); Ex[fasn-1(+))]* (compare panels h and i in Fig. 5).

We conclude that somatic *pod-2(+)* supplies sufficient levels of mal-CoA to the

germline not only to establish a wild-type 1-cell embryo, but also to support

continued development, presumably until zygotic *pod-2* expression becomes

adequate for completion of embryogenesis. These studies establish that

metabolites essential for embryonic development can be produced in the somatic

gonad and provided to the oocyte through gap junctions.

Intestinal loss of pod-2, but not fasn-1, results in a germline starvation response

The other class of *pod-2(0)* genetic mosaics producing dead eggs in our

original screen lost *Ex[pod-2(+); sur-5::gfp]* from the intestine (E–) (Fig. 5a). E(–)

mosaic animals were smaller than wild type, and gonad arms were severely

reduced in size (Fig. 6a). Only the most proximal oocytes showed appreciable

growth. E(–) mosaic gonad arms resemble those seen in gonads undergoing the

oocyte starvation response, or adult reproductive diapause (ARD), a process

induced by starvation during the L4 larval stage proposed to maintain germline

stem cells for repopulation of the gonad upon the resumption of feeding[30,31].

During ARD, germline stem cells cease dividing, the gonad shrinks due to

extensive apoptosis of differentiated germ cells, and growth is largely restricted to

the most proximal oocyte.

The *C. elegans* intestine is rich with lipid droplets and the source of yolk,

and regarded as a major site of FAS[32]. A genetic pathway in *C. elegans* required

for intestinal apical polarity establishment leading to proper lumen formation has

been described; genes in this pathway, including *pod-2*, encode enzymes

required for glycosphingolipid synthesis[33]. Lipid biosynthesis is required

throughout larval growth, coincident with membrane expansion. The identification

of adult mosaics lacking *pod-2(+)* in the intestine was therefore surprising. We

subsequently identified two presumptive $P_1(-)$ *pod-2* mosaics, predicted to have

lost *pod-2(+)* function in the intestine, somatic gonad, germline, many muscle and

hypodermal cells, among others (Fig. 5a). The most striking phenotype

distinguishing these animals from E(−) mosaics was an extremely constipated

intestine (Fig. 6b). The absence of this constipation in E(−) mosaic animals

implies that loss of *pod-2(+)* from the intestine does not prevent the processing of

bacteria through the intestinal lumen. (Both P_1 mosaics showed tissue extruded

from the rectum, which may be the cause of constipation.)

Although *gfp::fasn-1(tn1782)* is not detected in the intestine, we asked if

there is a functional role for *fasn-1* similar to *pod-2*. As described above,

homozygous *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]* progeny from heterozygous

parents produce Pod embryos but exhibit no other phenotype; within this class of

progeny we screened for E(–) mosaics. Ten (of 800) such animals were identified

(Fig. 5a); in stark contrast to *pod-2*, their gonad arms appeared normal. These

animals displayed no overt defects outside of Pod embryo production (Fig. 6c).

Because maternal FASN-1 rescues *fasn-1(0)* homozygotes to later embryonic

stages, an early intestinal function for *fasn-1* equivalent to *pod-2* could be

rescued maternally. However, the surprising conclusion from this mosaic analysis

is that there is no significant intestinal requirement for zygotic *fasn-1* in larval and

adult stages. The implication is that the primary function of mal-CoA produced in

the intestine may not be as substrate for intestinal FASN-1.

Is malonyl-CoA required in other germline processes?

inx-8(rf) and *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* share delayed oogenesis

and a reduction in gap junction coupling, germ cell proliferation, and brood size.

inx-21(0) mutants are sterile, and *inx-22(0)* mutants are fertile[3]; therefore mal-

CoA required for embryos must transit through gap junction channels comprised

of INX-14/INX-21 hemichannels, which are expressed throughout the distal arm

(Fig. 2b). We asked if mal-CoA levels might influence phenotypes preceding

embryogenesis, such as oocyte production.

Microinjection of mal-CoA (0.5-10 mM) into *inx-8/9(0); Ex[inx-8(DTC+,*

Sh-)] L4 gonad arms failed to ameliorate phenotypes such as Pod-embryo

production or low brood size. Ectopic expression of GFP::POD-2 in the germline

in *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* or *inx-8(rf)* genetic backgrounds also failed to

improve phenotypes, including delayed gametogenesis (Supplementary Fig. 5).

However, whether endogenous mal-CoA levels are altered under these

conditions is uncertain.

Effects of global mal-CoA levels were indirectly examined by comparing

higher or lower *pod-2(+)* copy number rescue of *pod-2(0)*; numbers of dead eggs

and larvae were summed to reflect total embryo output (Table 1). Lower copy

number rescue coincided with reduced embryo output (84% of wild-type) and

lower proportion of viable larvae within a brood (~37% of total embryos) (Table

1). Embryo output may therefore by impacted by *pod-2* expression; however,

somatic cells outside the gonad could be responsible. We therefore asked

specifically if FA synthesis is required within the germline. Somatically-rescued

fasn-1(0); Ex[fasn-1(+); sur-5::gfp] embryo output is comparable to wild-type

(94%); however, embryo output in the double mutant *fasn-1(0); inx-8(rf); Ex[fasn-*

1(+); sur-5::gfp] is only ~50% of *inx-8(rf)* alone (Table 1). The FAS pathway may

therefore play a germline role affecting embryo output. Surprisingly, *inx-8(rf)*

partially suppresses the embryonic lethality of *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]*

(Table 1). Possibly the smaller gonad arms of *inx-8(rf)* may concentrate levels of

FASN-1 or mal-CoA. Alternatively, *fasn-1* array germline expression may be de-

silenced in this genetic background. Further study is needed to address the

nature of this suppression.

Discussion

The genetic analysis of mutant *inx-8* soma-germline gap junction channels

has led us to three conclusions that have significance beyond the *C. elegans*

model system: 1) Mal-CoA produced in the somatic sheath is delivered through

gap junctions to the germline; this mal-CoA is necessary to establish a proper

one-cell embryo and to support continued embryogenesis. To our knowledge,

this segregation of the cellular source of mal-CoA from the eventual cell type in

which it is used as substrate is novel. As the rate-limiting step in FAS, mal-CoA

production is an attractive control point to tie aspects of germ cell developmental

progression to somatic control. 2) *pod-2* expression (and presumably mal-CoA

production) is exceptionally robust in the intestine and essential for normal

germline development. In the absence of intestinal POD-2, the germline exhibits

phenotypic hallmarks of germline ARD despite the presence of a replete food

source. By contrast, intestinal *fasn-1* expression is not similarly required. (A

possible caveat is that maternal *fasn-1* in the intestine may be sufficient to reach

adulthood, which would be surprising.) In light of the abundant expression level of

gfp::pod-2 in the intestine, and our finding that mal-CoA can be directionally

transferred intercellularly, the question arises as to whether the intestine provides

mal-CoA to other cells. Alternatively, a deficit of mal-CoA production in the

intestine might initiate a systemic starvation response. Recently, Gerisch *et al.*

(2020)[34] have shown that the HLH-30/TEFB transcription factor functions as a

master regulator of the ARD state. *hlh-30* mutants fail to effect the system-wide

changes promoting long-term survival and recovery associated with ARD. ARD

longevity does not appear to rely on fatty acid metabolism. However, entry of *hlh-*

30 mutants into the ARD state appears normal; therefore effects of decreased

mal-CoA expression in the intestine may lie upstream of *hlh-30* activity. 3) A

pronounced delay in gametogenesis along with reduction in brood size results

from reduced soma-germline gap junction communication; it is unknown if this

delay derives from generally slower mitotic or meiotic progression, or from a

developmental block at a particular meiotic stage. The nature of potential

molecules passing through gap junctions that are responsible for timely meiotic

progression is unknown, but mal-CoA has thus far not been implicated.

Our results are consistent with other findings in *C. elegans* related to fatty

acid metabolism in oocytes. *pod-2*/ACC requires post-translational covalent

attachment of biotin for enzymatic activity, and *bpl-1* (*b*iotin *p*rotein *l*igase-1) is

the *C. elegans* ortholog of the necessary holocarboxylase synthetase[35]. Mal-CoA

provided in the bacterial diet enables *bpl-1* mutants to develop to adulthood, but

de novo synthesis of polyunsaturated fatty acids (PUFAs) is required for

embryonic viability[35]. Our results indicate that mal-CoA supplied for PUFA

synthesis (and other FAS) in the early embryo is derived from the somatic sheath

via gap junctions. PUFA-dependent prostaglandin signaling from oocytes

influences sperm guidance, and reduction of *inx-14* function disrupts this

signaling[7,36]. We suggest that reduced levels of mal-CoA delivered to the oocyte

through mutant *inx-14*-containing gap junctions may reduce prostaglandin

synthesis and its signaling function for sperm migration.

At least one route outside of gap junctions may deliver mal-CoA to the

germline early in its development. The primordial germ cells Z2 and Z3 associate

with a pair of intestinal cells during embryogenesis; contacts are lost before

hatching[37,38]. Therefore an early germline requirement for mal-CoA could be

fulfilled directly from the intestine. Loss of such a function would be

encompassed within the ARD-like phenotype seen in *pod-2* E(–) mosaics.

In addition to its metabolic function in FAS, mal-CoA has established roles

as a regulatory molecule. It inhibits beta-oxidation of fatty acids by mitochondria.

Inhibitory mal-CoA derives from a second acetyl-CoA carboxylase (ACC2),

associated with the mitochondrial membrane. Manipulation of mal-CoA levels in

the mouse hypothalamus regulates food intake[39]. Regulation is effected through

the brain-specific CPT1-c, via a mechanism distinct from the enzymatic activity of

other CPT1 enzymes[40,41]. Little information regarding the *C. elegans* ACC2 is

known; somatically-delivered mal-CoA could potentially influence germline

mitochondrial function.

Malonylation as a post-translational lysine modification has recently been

described [42]. Malonylation may inhibit target proteins, such as the glyocolytic

enzymes glyceraldehyde 3-phosphate dehydrogenase and pyruvate kinase[43].

Malonylation of mTOR reduces mTORC1 complex activity, decreasing

endothelial cell proliferation and reducing angiogenesis in a human umbilical

model[44]. Together these studies and others suggest a potentially pervasive role

for mal-CoA in sensing and responding to nutritional status. One trait associated

with rapid tumor growth is an increase in FAS, and FASN has been explored as a

cancer therapy target[45]. Possibly the ability of mal-CoA to transit through gap

junctions might sustain FAS during tumor progression. The discovery that mal-

CoA transits through gap junctions from soma to germline may be indicative of

other unappreciated intercellular roles for this important molecule.

References

1. Skerrett, I. M., and Williams, J. B. A structural and functional comparison of gap junction channels composed of connexins and innexins. *Develop. Neurobiol.* **77,** 522-547 (2017)

2. Meyer, R. A., Laird, D. W., Revel, J.-P., and Johnson, R. G. Inhibition of gap junctions and adherens junction assembly by connexin and A-CAM antibodies. *J. Cell Biol.* **119,** 179-189.

3. Starich, T. A., Hall, D. H., and Greenstein, D. Two classes of gap junction channels mediate soma-germline interactions essential for germline proliferation and gametogenesis in *C. elegans. Genetics* **198,** 1127–1153 (2014).

4. Nadarajan, S., Govindan, J. A., McGovern, M., Hubbard, E. J., and Greenstein, D. MSP and GLP-1/Notch signaling coordinately regulate actomyosin-dependent cytoplasmic streaming and oocyte growth in *C. elegans. Development* **136,** 2223–2234 (2009).

5. Govindan, J. A., Cheng, H., Harris, J. E., and Greenstein, D. $G\alpha_{o/i}$ and $G\alpha_s$ signaling function in parallel with the MSP/Eph receptor to control meiotic diapause in *C. elegans. Curr. Biol.* **16,** 257–1268 (2006).

6. Whitten, S. J., and Miller, M. A. The role of gap junctions in *Caenorhabditis elegans* oocyte maturation and fertilization. *Dev. Biol.* **301,** 432–446 (2007).

7. Edmonds, J. W., McKinney, S. L., Prasain, J. K., and Miller, M. A. The gap junctional protein INX-14 functions in oocyte precursors to promote *C. elegans* sperm guidance. *Dev. Biol.* **359,** 47–58 (2011).

8. Austin, J., and Kimble, J. *glp-1* is required in the germ line for regulation of the decision between mitosis and meiosis in *C. elegans*. *Cell* **51**, 589-599 (1987).

9. Austin, J., Kimble, J. Transcript analysis of *glp-1* and *lin-12,* homologous genes required for cell interactions during development of C. elegans. *Cell* **58,** 565-571 (1989).

10. Henderson, S. T., Gao, D., Lambie, E. J., and Kimble, J. *lag-2* may encode a signaling ligand for the GLP-1 and LIN-12 receptors of *C. elegans*. *Development* **120,** 2913-2924 (1994).

11. Tagawa, A., Rappleye, C. A., and Aroian, R. V. *pod-2*, along with *pod-1*, defines a new class of genes required for polarity in the early *Caenorhabditis elegans* embryo. *Dev. Biol.* **233,** 412–424 (2001).

12. Johnston, W. L., Krizus, A., and Dennis, J. W. The eggshell is required for meiotic fidelity, polar-body extrusion and polarization of the *C. elegans* embryo. *BMC Biol.* **4,** 35 (2006).

13. Olson, S. K., Bishop, J. R., Yates, J. R., Oegema, K., and Esko, J. D. Identification of novel chondroitin proteoglycans in *Caenorhabditis elegans*: embryonic cell division depends on CPG-1 and CPG-2. *J. Cell Biol.* **173,** 985–994 (2006).

14. Benenati G., Penkov, S., Müller-Reichert, T., Entchev, E. V., and Kurzchalia, T. V. Two cytochrome P450s in *Caenorhabditis elegans* are essential for the organization of eggshell, correct execution of meiosis and polarization of the embryo. *Mech. Dev.* **126,** 382–393 (2009).

15. Zhang, Y., Foster, J. M., Nelson, L. S., Ma, D., Carlow, C. K. S. The chitin synthase

genes *chs-1* and *chs-2* are essential for *C. elegans* development and responsible for chitin deposition in the eggshell and pharynx, respectively. *Dev. Biol.* **285,** 330–339 (2005).

16. Rappleye, C. A., Tagawa, A., Le Bot, N., Ahringer, J., and Aroian, R. V. Involvement of fatty acid pathways and cortical interaction of the pronuclear complex in *Caenorhabditis elegans* polarity. *BMC Dev. Biol.* **3,** 8 (2003).

17. Rappleye, C. A., Tagawa, A., Lyczak, R., Bowerman, B., and Aroian, R. V. The anaphase-promoting complex and separin are required for embryonic anterior-posterior axis formation. *Dev. Cell* **2,** 195–206 (2002).

18. Rappleye, C. A., Paredez, A. R., Smith, C. W., McDonald, K. L., and Aroian, R. V. The coronin-like protein POD-1 is required for anterior-posterior axis formation and cellular architecture in the nematode *Caenorhabditis elegans*. *Genes Dev.* **13,** 2838–2851 (1999).

19. Oshima, A., Tani, K., and Fujiyoshi, Y. Atomic structure of the innexin-6 gap junction channel determined by cryo-EM. *Nat. Commun* **7,** 13681 (2016). https://doi.org/10.1038/ncomms13681

20. Mok, D. Z. L., Sternberg, P. W., and Inoue, T. Morphologically defined sub-stages of *C. elegans* vulval development in the fourth larval stage. *BMC Dev. Biol.* **15,** 26 (2015).

21. Spike, C. A., Coetzee, D., Eichten, C., Wang, X., Hansen, D., and Greenstein, D. The NHL-TRIM protein LIN-41 and the OMA RNA-binding proteins antagonistically control the prophase-to-metaphase transition and growth of *C. elegans* oocytes.

Genetics **198,** 1535–1558 (2014).

22. Jaramillo-Lambert, A., Fuchsman, A. S., Fabritius, A. S., Smith, H. E., and Golden, A. Rapid and efficient identification of *Caenorhabditis elegans* legacy mutations using Hawaiian SNP-based mapping and whole-genome sequencing. *G3* **5,** 1007-1019 (2015).

23. Kelly, W. G., Montgomery, M. K., and Fire, A. Distinct requirements for somatic and germline expression of a generally expressed *Caenorhabditis elegans* gene. *Genetics* **146,** 227-238 (1997).

24. Arribere, J. A., *et al.* Efficient marker-free recovery of custom genetic modifications with CRISPR/Cas9 in *Caenorhabditis elegans*. *Genetics* **198,** 837-846 (2014)

25. Yochem, J., Gu. T., and Han, M. A new marker for mosaic analysis in *Caenorhabditis elegans* indicates a fusion between hyp6 and hyp7, two major components of the hypodermis. *Genetics* **149,** 1323-1334 (1998)

26. WormBase (www.wormbase.org)

27. Lee, K.-Z., Kniazeva, M., Han, M., Pujol, N., and Ewbank, J. The fatty-acid synthase *fasn-1* acts upstream of WNK and Ste20/GCK-VI kinases to modulate antimicrobial peptide expression in *C. elegans* epidermis. *Virulence* **1,** 113-122 (2010)

28. Cox, B. G., and Hammes, G. G. Steady-state kinetic study of fatty acid synthase from chicken liver. *Proc. Natl. Acad. Sci.* **80,** 4233-4237 (1983)

29. Chang, S.-I., and Hammes, G. G. Structure and mechanism of action of a multifunctional enzyme: fatty acid synthase. *Acc. Chem. Res.* **23,** 363-369 (1990)

30. Angelo, G., and Van Gilst, M. R. Starvation protects germline stem cells and extends

reproductive longevity in *C. elegans*. *Science* **326,** 954-958 (2009)

31. Seidel, H. S., and Kimble, J. The oogenic germline starvation response in *C. elegans*. *PloS ONE* **6**(12)**,** e28074. (2011)

32. Watts, J. L., and Ristow, M. Lipid and carbohydrate metabolism in *Caenorhabditis elegans*. *Genetics* **207,** 413-446 (2017)

33. Zhang, H., *et al*. Apicobasal domain identities of expanding tubular membranes depend on glycosphingolipid biosynthesis. *Nat. Cell Biol.* **13,** 1189-1201 (2012)

34. Gerisch, B., *et al*. HLH-30/TFEB is a master regulator of reproductive quiescence. *Develop. Cell* **53,** 1-14 (2020)

35. Watts, J. S., Morton, D. G., Kemphues, K. J., and Watts, J. L. The biotin-ligating protein BPL-1 is critical for lipid biosynthesis and polarization of the *Caenorhabditis elegans* embryo. *J. Biol. Chem.* **293,** 610-622 (2018)

36. Kubagawa, H. M., *et al*. Oocyte signals derived from polyunsaturated fatty acids control sperm recruitment *in vivo*. *Nat. Cell Biol.* **8**, 1143-1148 (2006)

37. Sulston, J. E., Schierenberg, E., White, J. G. & Thomson, J. N. The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* **100,** 64–119 (1983).

38. Abdu, Y., Maniscalco, C., Heddleston, J. M., Chew, T.-L., and Nance, J. Developmentally programmed germ cell remodelling by endodermal cell cannibalism. *Nat. Cell Biol.* **18,** 1302-1310 (2016)

39. Wolfgang, M. J., and Lane, M. D. The role of hypothalamic malonyl-CoA in energy homeostasis. *J. Biol. Chem.* **281,** 37265-37269 (2006)

40. Sierra, A. Y., et al. CPT1c is localized in endoplasmic reticulum of neurons and has

carnitine palmitoyltransferase activity. *J. Biol. Chem.* **283,** 6878-6885 (2008)

41. Reilly, P. T., and Mak, T. W. Molecular pathways: tumor cells co-opt the brain-specific metabolism gene *CPT1C* to promote survival. *Clin. Cancer Res.* **18(21)** (2012) doi: 10.1158/1078-0432.CCR-11-3281

42. Peng, C., *et al.* The first identification of lysine malonylation substrates and its regulatory enzyme. *Mol. Cell. Proteomics* **10(12)** doi:10.1074/mcp.M111.012658

43. Kulkami, R. A., *et al.* Discovering targets of non-enzymatic acylation by thioester reactivity profiling. *Cell Chem. Biol.* **24,** 231-242 (2017)

44. Bruning, U., *et al.* Impairment of angiogenesis by fatty acid synthase inhibition involves mTOR malonylation. *Cell Metabolism* **28,** 866-880 (2018)

45. Menendez, J. A., and Lupu, R. Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nature Rev.* **7,** 763-777 (2007)

46. McCarter, J., Bartlett, B., Dang, T., and Schedl, T. Soma-germ cell interactions in *Caenorhabditis elegans*: multiple events of hermaphrodite germline development require the somatic sheath and spermathecal lineages. *Dev. Biol.* **181,** 121-143 (1997).

47. Berman,H. M., *et al.* The protein data bank. *Nuc. Acids Res.* **28,** 235-242 (2000)

48. NGL viewer: web-based molecular graphics for large complexes. Bioinformatics doi:10.1093/bioinformatics/bty419 (2018)

49. Dickinson, D. J., Pani, A. M., Heppert, J. K., Higgins, C. D., and Goldstein, B. Streamlined genome engineering with a self-excising drug selection cassette. *Genetics* **200,** 1035-1049 (2015)

Methods

Strains

C. elegans strains were grown on standard NGM media (plus Nystatin, 6.25 mg/l)

with *E. coli* strain OP50 as food source. Temperatures used for conditional

mutants are indicated in main text. In addition to the wild-type strain N2, the

following alleles were used: Chr. *I*: *fasn-1(g43ts); fasn-1(tn1762null); fasn-*

1(tn1782[gfp::fasn-1]); lin-41(tn1541[gfp::tev::s-tag::lin-41]); Chr. *II*: *pod-*

2(ye60cs); pod-2(tn1691null); pod-2(tn1765[gfp::tev::3xFLAG::pod-2]); Chr. *III*:

emb-8(hc69ts); dpy-17(e1295); Chr. *IV*: *inx-8(tn1474null) inx-9(ok1502null); inx-*

8(tn1513) inx-9(ok1502); inx-8(tn1513 tn1555) inx-9(ok1502)—a.k.a. *inx-8(rf)* in

main text. Balancers used included: *tmC18 [dpy-5(tmIs1236)] I; hT2[bli-4(e937)*

let-?(q782) qIs48] (I:III); mIs11 IV. Extrachromosomal arrays used included:

tnEx195[inx-8(+) inx-9(+); sur-5::gfp]; tnEx205[lag-2p::inx-8::gfp; str-1::gfp];

tnEx212[W09B6-pod-2(+)--20 ng/µl—hi copy; sur-5::gfp]; tnEx218[fasn-1(+); sur-

5::gfp]; tnEx219[W09B6-pod-2(+)--4 ng/µl—lo copy; sur-5::gfp]; and *tnEx221[inx-*

8(tn1513 tn1555); str-1::gfp].

Microinjection rescue of FAS mutants with extrachromosomal arrays

The cosmid W09B6 was used at 50 ng/µl (2 lines) or 20 ng/µl (1 line) with *sur-*

5::gfp (75 ng/µl) to rescue *pod-2(ye60cs). sur-5::gfp* expression was faint for both

lines obtained with 50 ng/µl and these were not further characterized. *pod-*

2(tn1691null) was rescued by microinjection using W09B6 at 4 ng/µl (5 lines) or

by crossing in *tnEx212*. Fosmids WRM0613bD07(10 ng/µl, or 50 ng/µl—1 line

that silenced) and WRM067cB03 (25 ng/ul) were used to attempt rescue of *emb-*

8(hc69ts). Numerous conditions using fosmids WRM0612bD08 and

WRM0614aH05 in simple or complex arrays were used to try to rescue *fasn-*

1(g43ts) without success. However, FASN enzyme functions as a dimer;

formation of dimers between mutant (*g43*) and wild-type monomers might inhibit

phenotypic rescue. After *fasn-1(tn1762null)* was generated, somatic but not

germline rescue was obtained using a *fasn-1* genomic PCR product forming

simple arrays (1 ng/µl + *sur-5::gfp-75* ng/µl, at least 2/6 lines segregated *fasn-*

1(0); Ex[fasn-1(+); sur-5::gfp(+)] animals that laid only Pod embryos), or a

complex array (1 ng/µl + 20 ng/µl *sur-5::gfp* + 100 ng/µl salmon sperm,1 line

from 30 injected animals) as described in the main text.

Recombinant DNA constructs for isolation of new pod-2 and fasn-1 alleles

Standard CRISPR/Cas-9 methods were used to generate deletions and GFP

insertions in *pod-2* and *fasn-1.* For deletions, 100-nt repair templates were

synthesized (Ultramers; IDT) and used in a *dpy-10(cn64)* co-conversion strategy

previously described[24]. Injection mixes included: pDD162 (Cas9), 50 ng/µl; 25 ng/µl

(*pod-2*) or 10 ng/µl (*fasn-1*) each single guide; Ultramer repair template, 500 nM; *dpy-*

10 single guide (pJA58), 25 ng/µl; *dpy-10(cn64)* Ultramer A2-2F-827, 500nM. Single

guides were made by annealing oligo pairs and cloning into pRB1017[24]: Deletions

were verified by PCR. The PCR fragment used to rescue *fasn-1(tn1762null)* was

amplified from genomic DNA using primers 5'-tacgtcaaacatccgtttgtcaacgtcac-3' and

5'-ttcttctcctcgtcctttgatacgaagatc-3'.

Generation of GFP insertions into *pod-2* using pDD282 followed established

protocols[49]. Injection mix included the single guide (10 ng/μl), repair template (10

ng/μl), pDD162 (50 ng/μl), and *sur-5::gfp* (80 ng/μl). The insertion was verified by

sequencing PCR products generated with primers positioned outside the homology

arms.

For *gfp::fasn-1*, we were unable to isolate 3' insertions using the Dickinson

method[49]. This method relies on isolation of candidates based on the Roller

phenotype, and because *fasn-1* has been implicated in molting, we turned to a co-

conversion approach[24]. A repair template was generated that included the GFP

sequence from pDD282, and *fasn-1* homology arms on either side. The injection mix

contained pDD162, 50 ng/μl; *fasn-1* single guide 10 ng/μl; *gfp::fasn-1* repair template,

20 ng/μl; *dpy-10* single guide (pJA58), 25 ng/μl; *dpy-10(cn64)* ultramer A2-2F-827,

500nM; and pCFJ190, 2.5 ng/μl. Rol candidates were screened directly for GFP

expression. The insertion was verified by sequencing PCR fragments generated

using primers placed outside of the repair template sequence (5'-

ggaacacaaacattccacatcgccg-3' and 5'-gtgattcatctgaattcgctcactcc-3').

mex-5p::gfp::pod-2::tbb-2-3'UTR was constructed by Gibson assembly of

component PCR products representing the *mex-5* promoter, the "a" isoform of

gfp::pod-2, and the tbb-2-3'UTR to prevent germline RNA degradation. Sequences for

all oligos to generate single guides, repair templates, and recombinant plasmids are

available upon request.

Immunofluorescence

Antibody staining of dissected gonads was as previously described[3]. DIC and

fluorescent images were acquired on a Zeiss motorized Axioplan 2 microscope

with either a 40x Plan-Neofluar (numerical aperture 1.3), a 63x Plan-

Apochromatic (numerical aperture 1.4), or 100x PlanApochromatic (N.A. 1.4)

objective lens using a AxioCam MRm camera and AxioVision software (Zeiss).

Mosaic analysis

To associate the somatic focus of action of *pod-2* for germline phenotypes, array

losses were characterized in animals of genotype *pod-2(1691null); tnEx212[W09B6-*

pod-2(+)--20 ng/µl—hi copy; sur-5::gfp] that were isolated on the basis of sterility.

Subsequently, potential candidate mosaics of interest were prescreened directly at

the dissecting microscope level for asymmetry in the phenotypic severity seen

between anterior and posterior gonad arms, and then examined at the compound

microscope level for loss of *sur-5::gfp* expression. Potential *pod-2(tn1691null);*

tnEx219[W09B6-pod-2(+)–4 ng/µl—lo copy; sur-5::gfp] genetic mosaics were also

prescreened in this manner. Intestinal (E–) mosaics were prescreened using low-

power fluorescence (Nikon SMZ18, with Lumencor SOLA Light Engine), but for *pod-2*

many of these candidate mosaics were seen to express *sur-5::gfp* at very low levels,

or in just a few intestinal cells, when examined at high power. This was not an issue

when isolating *fasn-1(tn1762null); tnEx218[fasn-1(+); sur-5::gfp]* E(−) mosaics.

Data availability

All *C. elegans* strains are available from the Caenorhabditis Genetics Center or by request. DNA sequences for all oligonucleotides used to generate single guides, repair templates, and plasmids are available upon request. All other data and materials supporting the findings of this study are available upon request.

Acknowledgements

We are grateful to our colleagues Gabriela Huelgas-Morales, Caroline Spike, and Tatsuya Tsukamoto for suggestions regarding experiments and comments on the manuscript. This work was supported by National Institutes of Health (NIH) grant GM57173 to D.G. Some strains were provided by the Caenorhabditis Genetics Center, which is funded by grant P40OD010440 from the NIH Office of Research Infrastructure Programs.

Author contributions

T.S. conducted the experiments, prepared the figures, and wrote the manuscript. T.S. and D.G. designed and conceptualized the research study. D.G. obtained funding, supervised the study, and edited the manuscript.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. Ovulation and oocyte development are delayed in *inx-8(rf)*. **a**) Diagram of adult *C. elegans* gonad. Posterior arm shows the positions of the somatic distal tip cell (DTC) and five pairs of sheath cells (nuclei are numbered). Anterior arm shows the underlying germline, approximate regions of germ cell proliferation (Prolif.), meiotic pachytene stage (Pach.), and developing oocytes. Embryos typically undergo some cell division in the uterus before being laid. Sp, spermatheca; vul, vulva. **b**) Compared to wild type (N2), ovulation in *inx-8(tn1513 tn1555rf)* mutants (a.k.a. *inx-8[rf]*) is delayed ~18 hrs. Bars representing 1-3 hr windows indicate approximate time of the fourth larval stage (L4 to adult) molt, followed by time of appearance of the first fertilized embryo ovulated into the uterus. **c)** Onset of *gfp::lin-41* expression as a marker of oocyte development is delayed in *inx-8(rf)*. Panels show DIC images of vulval development (left), loop region of gonad arm (middle), and expression of *gfp::lin-41* (right). Wild-type hermaphrodites express *gfp::lin-41* by the approximate L4.7 sub-stage of vulval development (vul); *inx-8(rf)* hermaphrodites do not express *gfp::lin-41* until after vulval development is complete but before the L4-to-adult molt. Null *inx-8(tn1474) inx-9(ok1502)* hermaphrodites, rescued for *inx-8::gfp* expression in the DTC with *Ex[lag-2p::inx-8::gfp]*--a.k.a. *inx-8/9(0); Ex[inx-8(DTC+, Sh-)]* -- show a similar delayed onset of *gfp::lin-41* expression as *inx-8(rf)*. Bar, 20 μm.

Figure 2. INX-8(T239I D24N) compromises gap junction formation between soma and germline. **a**) Somatic INX-8 in wild-type (WT) gonads localizes to

puncta representing soma-germline gap junctions where DTC or sheath cells

overlie germ cells. **b)** In WT distal gonad arms, gap junction puncta aggregate in

small clusters associated with each germ cell; in *inx-8(rf)* distal arms, gap

junction puncta localize more apically, where sheath dips down between germ

cells, outlining the cell (honeycomb pattern). Localization of INX-21 and INX-22

define the two different classes of germline gap junction hemichannels, both of

which include INX-14 subunits. **c)** At the loop region, WT gap junction puncta

form large higher-order aggregates (arrows) which are absent in *inx-8(rf)*. **d)** In

WT proximal arms, gap junction formation with INX-22-containing hemichannels

is extensive; in *inx-8(rf)*, INX-22 and INX-8(T239I D24N) are strongly expressed

but show little evidence of gap junction formation. "–1" indicates the most

proximal oocyte; *inx-8(rf)* hermaphrodites frequently show undersized oocytes at

this position as well (<). **e)** Heterozygous *+/inx-8(rf)* shows little effect on gap

junction localization in an *emb-8(hc69ts)* background. Antibodies specific to INX-

8, INX-21, or INX-22 were used as indicated. DAPI-stained DNA in blue. Bar, 50

μm.

Figure 3. a**)** Predicted assignment of FAS mutants in a simplified FAS pathway.

Targeted deletions and GFP insertions are shown for *pod-2* (**b**) and *fasn-1* (**c**).

Predicted isoforms for POD-2 are indicated. **d-h**) Phenotypes of homozygous

mutant embryos from **d**) *+/pod-2(0)*; **e-h**) *fasn-1(0); Ex[fasn-1(+)]*. Arrow in (**d**)

indicates differentiated anterior pharynx; arrow in (**e**) indicates multiple ectopic

nuclei present at fertilization that persist during early stages of embryogenesis.

Figure 4. Wild-type INX-8 facilitates effects of *Ex[pod-2(+)]* overexpression on

conditional mutations downstream in the fatty acid synthesis (FAS) pathway.

Brood size was measured as the number of viable progeny produced. The

balancer *mIs11* provided *inx-8(+)* for all strains listed. **a**) Embryonic lethality in

emb-8(ts) mutants at 22°C is more severe in heterozygous *+/inx-8(rf)* than *inx-*

8(+) backgrounds *(1)*. Expression of *Ex[pod-2(+)]* in *emb-8(ts)* significantly

rescues embryonic lethality *(2)*. Rescue of *emb-8(ts)* by *Ex[pod-2(+)]* is also

dependent on *inx-8(+)* copy number *(3)*. **b**) Expression of *Ex[pod-2(+)]* in *fasn-*

1(ts) mutants at 20°C reduces brood size *(1)*. *fasn-1(ts); inx-8(+)* homozygotes

show greater embryonic lethality than *fasn-1(ts); (+)/inx-8(rf)* heterozygotes in the

absence *(2)* or presence *(3)* of *Ex[pod-2(+)]*. Averages indicated with dotted

lines. Probability was calculated using Student's 2-tailed *t*-test. **c**) Enzymes

(purple) implicated in germline FAS. Malonyl-CoA transits from soma sheath to

germline through gap junction channels comprised of INX-8/9 (soma) and INX-

14/INX-21 (germline) hemichannels. Germline FASN-1 uses malonyl-CoA to

synthesize palmitate, which may be further extended and modified to produce

other long-chain fatty acids. EMB-8 (NADPH-dependent CYP450 reductase) and

POD-7/8 (CYP450s) are necessary to synthesize the lipid-rich (inner) layer of the

eggshell and support polarity establishment[14,16]; the composition of lipid layer

fatty acids is unknown. *de novo* PUFA synthesis is required in the germline for

embryonic viability, and is implicated in sperm guidance[35,36]. The identities of

other long-chain fatty acids that may be synthesized in the germline are

unknown. PUFAs, polyunsaturated fatty acids; LFAs, long-chain fatty acids;

MUFA, monounsaturated fatty acids; pm, plasma membrane.

Figure 5. Genetic mosaic analyses of *pod-2* reveal a mal-CoA requirement in

somatic gonad sheath. **a**) Abbreviated cell lineage of *C. elegans*. Mosaics

characterized correspond to loss of *sur-5::gfp* expression in the germline (P_4),

intestine (E), or somatic gonad. Five *pod-2(0); Ex[pod-2(+); sur-5::gfp]* animals

lost the array from all somatic gonad sheath cells and produced no viable

embryos. **b**) Eight animals showed differences in array retention in each gonad

arm with phenotypic consequences. Arrays were generated by microinjecting

cosmid W09B6 *[pod-2(+)]* at 4 ng or 20 ng/µl (low/high copy). **c**) Uterus of non-

mosaic *pod-2(0); Ex[pod-2(+)--high copy]* hermaphrodite; developmental

progression of embryos from each gonad arm is comparable. **d**) Mosaic *pod-*

2(0); Ex[pod-2(+)--high copy] with array lost from posterior gonad arm sheath but

retained in a single Sh4 cell of anterior arm. Anterior arm embryos have more

distinct eggshells and develop to later stages. **e-h**) Embryonic development

depends on somatic levels of mal-CoA. Shown are terminal-stage *pod-2(0)*

embryos derived from parents of designated genotypes. **e**) Pretzel-stage embryo

from *+/pod-2(0)*. **f**) 3-fold stage embryo from *pod-2(0); Ex[pod-2(+)--low copy]*,

with array maintained in soma but lost in the germline (P4–). **g**) Pretzel-stage

embryo from P4(–) mosaic *pod-2(0); Ex[pod-2(+)--high copy]*. **h**) Embryo from

mosaic *pod-2(0); Ex[pod-2(+)--low copy]* with uncharacterized somatic loss of

array, but array retained in the germline (P4+). This parent produced exclusively

dead embryos. **i**) Typical terminal undifferentiated *fasn-1(0)* Pod embryo from

fasn-1(0); Ex[fasn-1(+)]. ant ute, anterior uterus; post ute, posterior uterus; v,

vulva; −1, most proximal oocyte. Bar, 20 μm.

Figure 6. Mosaic loss from the intestine of *pod-2*, but not *fasn-1,* leads to an

ARD-like gonad arm. **a**) E(−) genetic mosaic *pod-2(0); Ex[pod-2(+); sur-5::gfp]*

gonad arms (from same animal). Gonad arm on right partially outlined. **b**) Two

focal planes of a presumptive P_1(−) *pod-2(0); Ex[pod-2(+); sur-5::gfp]* mosaic with

constipated intestine. Upper plane (left) with partially outlined intestine; lower

plane (right) shows underlying distal gonad arm. int, intestine; p, pachytene

region; v, vulva. **c**) E(−) genetic mosaic *fasn-1(0); Ex[fasn-1(+); sur-5::gfp]* gonad

arm. −1, most proximal oocyte. Bars, 20 μm.

Table 1

Brood sizes suggest FA synthesis influences embryo output

Genotype	Larvae	Dead embryos	Total (embryo output)
Wild type (N2)	291±29 (n=33)	2±2	293±29
inx-8(rf)	108±40 (n=90)	11±6 (n=30)	119±42
inx-8/9(0); Ex[inx-8(DTC+,Sh-)]	2±2 (n=32)	19±13	20±14
pod-2(0); Ex[pod-2(+)] high copy (20 ng/µl)	164±50 (n=31)	118±41	282±59
pod-2(0); Ex[pod-2(+)] low copy (4 ng/µl)	91±24 (n=41)	156±37	247±50
fasn-1(0); Ex[fasn-1(+)	0.05 (n=80)	276±18 (n=16)	276±18
fasn-1(0);inx-8(rf); Ex[fasn-1(+)	17±15 (n=47)	43±27	60±37

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

637   **Supplementary Information**

639   **Gap junctions deliver malonyl-CoA from soma to germline to support**

640   **embryogenesis in *Caenorhabditis elegans***

642   Todd A. Starich[1,2] and David Greenstein[1]

643   [1]Department of Genetics, Cell Biology and Development, University of

644   Minnesota, Minneapolis, Minnesota 55455 USA

[2]Email: stari001@umn.edu

Contents:
Supplementary Tables: 2
Supplementary Figures: 5
Supplementary Figure Legends
Supplemental References (46-49 in the list in the main text).

Supplementary Table 1

inx-8(rf) exhibits reduced brood size and interacts with FA synthesis pathway genes

Genotype	Brood size	Germ cells per gonad arm
20°C		
N2 (wild type)	291±29 (n=33)	≥1000[46]
inx-8(rf)[1]	108±40 (n=90)	322±18 (n=3; day 2 adult)
22°C		
inx-8(+)[2]	266±40 (n=32)	
+/inx-8(rf)	255±36 (n=32)	
fasn-1(g43ts); inx-8(+)	115±41 (n=11)	
fasn-1(g43ts); +/ inx-8(rf)	114±47 (n=11)	
fasn-1(g43ts); inx-8(rf)	4.5±2.3 (n=12)	
emb-8(hc69ts); inx-8(+)	95±50 (n=15)	
emb-8(hc69ts); +/inx-8(rf)	35±37 (n=15)	
emb-8(hc69ts); inx-8(rf)	11±9 (n=15)	

[1]Full genotype *inx-8(tn1513 tn1555) inx-9(ok1502).* Approximately 4% of animals are infertile in one or both gonad arms (n=168).
[2]*inx-8(+)* provided on the balancer *mIs11* IV.

Supplementary Table 2

Reduced soma-germline gap junction coupling delays gametogenesis

a) *gfp::lin-41* expression at L4 vulval sub-stages

Genotype	L4.4	L4.5	L4.6	L4.7	L4.8	L4.9	L4.9< <molt
gfp::lin-41; inx-8(+)	1/11		3/3	5/6	5/6	4/4	
gfp::lin-41; inx-8(rf)		0/2	0/3	0/6		0/2	16/37
gfp::lin-41; inx-8/9(0); Ex[inx-8(DTC+,Sh-)]	0/11	0/3		0/2	0/1	0/3	6/14

b) sperm appearance at L4 sub-stages

Genotype	L4.4	L4.5	L4.6	L4.7	L4.8	L4.9	L4.9< < molt
N2	0/9	0/2	0/5	4/11	1/2	1/2	6/6
inx-8(rf)	0/3		0/1	0/2	0/3	0/1	1/10

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

653 **Supplementary Figure Legends**
654
Supplementary Figure 1. Predicted topological sites of amino acids mutated in

inx-8(tn1513 tn1555) inx-9(0). **a**) Atomic structure of the *C. elegans* homomeric,

homotypic INX-6 gap junction full channel, determined by Oshima, *et al.*[19] (PDB

I.D. 5H1R, rcsb.org[47], viewed with NGL viewer[48]). Octameric hemichannels

associate via extracellular loops to form an enclosed channel in the region

classically identified by TEM as the "gap" between closely aligned plasma

membranes. cyto, cytoplasmic membrane face; p.m., plasma membrane. **b-f**)

Model for INX-8 hemichannel based on homology alignment to INX-6

(aquaria.ws). **b**) Cytoplasmic view. **c**) Single subunit highlighted. In wild type this

hemichannel is comprised of both INX-8 and INX-9 subunits, and associates with

germline hemichannels comprised of INX-14+INX-21 or INX-14+INX-22. **d**) Side

view of hemichannel. Beta-strands (yellow) in the extracellular loops contribute to

walling off the pore. **e**) Single subunit highlighted. Kinked region (gray arrow)

allows the cytoplasmic structure of the subunits (cytoplasmic loop and C-

terminus) to form a dome at the pore entry. Amino termini (N-term) line the pore

as a funnel; a loop in the amino terminus allows for an interaction with residues in

the dome[19]. D24 in INX-8 (red arrow) lies near the D21 residue predicted by

homology with INX-6 (D25) to interact with the cytoplasmic dome. **f**) INX-8 T239

is predicted to lie adjacent to a beta-strand contributing to hemichannel-

hemichannel association (red arrow). (To expose the extracellular loop region in

relation to "**e**", the aspect of view is rotated towards the viewer and angled up to

the left.)

Supplementary Figure 2. *inx-8(rf)::gfp* can be endocytosed by oocytes. **a**) *inx-*

8(+)::gfp expressed in the somatic gonad sheath is endocytosed during ovulation

and appears as distinct fluorescent puncta in early embryos. **b**) *inx-8(rf)::gfp*

expressed on an extrachromosomal array in *inx-8(0) inx-9(0)* animals is

endocytosed by ovulating oocytes. **c**) *inx-8::gfp* expressed by the DTC in *inx-*

8/9(0); Ex[inx-8(DTC+,Sh-)] is not observed in early embryos. Bar, 20 μm.

Supplementary Figure 3. GFP::POD-2 expression. **a)** Expression in the head

includes hypodermis (carets) and excretory duct cell (arrow). Int, intestine. **b**)

Primary expression in the tail includes hypodermis (carets) and intestine (Int).

Left panel DIC image, right panel GFP fluorescence in live animals. **c)** Dissected

animal stained with anti-GFP antibody shows abundant intestinal expression (Int)

and strong expression in proximal sheath of exposed gonad arm (prox) that

diminishes in the distal arm. No germline expression is seen in the distal arm

where germ cells are revealed due to the absence of sheath coverage. Arrow

indicates CAN neuron. DAPI-staining of DNA in blue. **d**, **e**) Earliest expression in

embryos is evident in developing hypodermis and intestine. **a, b, d, e**) DIC

images in left panels, live GFP fluorescence in right panels. Bars are 20 μm,

except (**c**) 200 μm. **f**) GFP::POD-2 is responsive to starvation. Top row, animals

briefly washed and transferred to bacteria-free NGM plates for 4 hours; lower

row, animals similarly-treated but fed.

Supplementary Figure 4. GFP::FASN-1 expression and *fasn-1(0)* embryonic

phenotypes. **a**) GFP::FASN-1 is expressed throughout the hypodermis (small

arrows). Excretory duct cell expression is also indicated (larger arrow). **b**) Arrow

indicates proximal gonad sheath expression. **c**) Arrow indicates developing

vulva. **d**) Earliest expression in developing embryos is seen in hypodermis. DIC

images in upper panels, live GFP fluorescence in lower panels. **e-h**) Terminal

phenotypes of *fasn-1(0)* embryos or larvae from *+/fasn-1(0)* parents. All bars, 20

μm.

Supplementary Figure 5. Expression of *gfp::pod-2* in the germline fails to

rescue *inx-8(rf)* phenotypes. **a**) Young (first-day) adult *inx-8(rf)* gonad arm. **b, c**)

Young adult *inx-8(rf)* gonad arm expressing *mex-5p::gfp::pod-2::tbb-2-3'UTR*.

The *pod-2* isoform in this construct corresponds to *gfp::pod-2* isoform "a" in Fig.

3. This isoform encompasses sequence homology to all of the enzymatic sites in

other ACC multi-functional enzymes. Sp, spermatheca; -1, most proximal oocyte;

bar, 50 μm.