Enhanced differentiation of functional human T cells in NSGW41 mice with tissue-specific expression of human interleukin-7.

Emilie Coppin[1,2*], Bala Sai Sundarasetty[3*§], Susann Rahmig[1,2], Jonas Blume[4], Nikita A. Verheyden[3], Franz Bahlmann[5], Sarina Ravens[4], Undine Schubert[9,10], Janine Schmid[9,10], Stefan Ludwig[11], Constantin von Kaisenberg[6], Alexander Platz[7], Ronald Naumann[8], Barbara Ludwig[9,10], Immo Prinz[4], Claudia Waskow[1,2,9,12$], Andreas Krueger[3$]

[1] Regeneration in Hematopoiesis, Institute for Immunology, TU Dresden, Dresden, Germany
[2] Immunology of Aging, Leibniz Institute on Aging - Fritz Lipmann Institute, Jena, Germany.
[3] Institute for Molecular Medicine, Goethe University Frankfurt am Main, Germany.
[4] Institute of Immunology, Hannover Medical School, Hannover, Germany.
[5] Department of Obstetrics and Gynecology, Bürgerhospital Frankfurt, Germany.
[6] Department Obstetrics, Gynecology and Reproductive Medicine, Hannover Medical School, Hannover, Germany.
[7] DKMS Cord Blood Bank, Dresden, Germany.
[8] Max-Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany.
[9] Department of Medicine III, Technical University Dresden, Dresden, Germany.
[10] German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany.
[11] Department of Visceral-, Thorax- and Vascular Surgery, Technical University Dresden, Dresden, Germany.
[12] Institute of Biochemistry and Biophysics, Faculty of Biological Sciences, Friedrich-Schiller-University Jena, Jena, Germany.

* same contribution first authors

$ joint supervision

§ Current address: Recombinant Technologies, CSL Behring GmbH, Marburg, Germany

Corresponding author

Claudia Waskow

Claudia.waskow@leibniz-fli.de

Leibniz-Institute on Aging, Beutenbergstrasse 11, 07745 Jena, Germany

Phone: +49 3641 656 707

Key points

1.   Increased intrathymic human T cell differentiation in the absence of pathological lymphoproliferation in humanized NSGW41hIL7 mice.

2.   Increased peripheral human T cell populations in NSGW41hIL7 mice making the analysis of human regulatory T cells in vivo feasible.

Abstract

Humanized mouse models have become increasingly valuable tools to study human hematopoiesis and infectious diseases. However, human T cell differentiation remains inefficient. We generated mice expressing human interleukin (IL-7), a critical growth and survival factor for T cells, under the control of murine IL-7 regulatory elements. After transfer of human cord blood-derived hematopoietic stem and progenitor cells, transgenic mice on the NSGW41 background, termed NSGW41hIL7, showed elevated and prolonged human cellularity in the thymus while maintaining physiological ratios of thymocyte subsets. As a consequence, numbers of functional human T cells in the periphery were increased without evidence for pathological lymphoproliferation or aberrant expansion of effector or memory-like T cells. We conclude that the novel NSGW41hIL7 strain represents an optimized mouse model for humanization to better understand human T cell differentiation *in vivo* and to generate a human immune system with a better approximation of human lymphocyte ratios.

Introduction

Humanized mouse models have emerged as indispensable tools for improving our understanding of human hematopoiesis and the human immune system. Typically, they are generated by transplanting human hematopoietic stem and progenitor cells (HSPCs) into mice (for review see [1]). NSGW41 mice carry the hypomorph W41 allele in the *Kit* gene, harbor the NOD-specific variant of the *Sirpa* gene, are T-, B- and NK-cell deficient based on null mutations in *Prkdc* and *Il2rg* genes, and allow for human donor stem cell engraftment in the absence of preconditioning [2, 3]. NSGW41 mice that are stably engrafted with human hematopoietic stem cell display continuous human hematopoiesis with increased myeloid [2, 4], megakaryocytic and erythroid output compared to irradiated NSG recipient mice [5, 6].

The introduction of human genes to overcome selective deficiencies in human hematopoiesis has resulted in further improvement of humanized mouse models. Thus, mice expressing human cytokines, including a combination of SCF, GM-CSF, and IL-3 (NSG-SGM3) or M-CSF, IL-3, GM-CSF, and thrombopoietin (MISTRG) show improved myelopoiesis [7, 8]. Mice carrying a knock-in of human IL-6 show improved B-cell development and function [9].

Efficient differentiation of human T cells remains a challenge in humanized mice and we have focused on interleukin-7 (IL-7) to improve that situation. Patients with mutations in the *IL7RA* gene suffer from profound T–B+NK+ severe combined immunodeficiency [10]. In mice, loss of the *Il7ra* gene results in combined B and T-cell lymphopenia, pointing towards critical cross-species differences [11, 12]. *In vitro*, murine (m)IL-7 was 100-fold less potent to expand and differentiate human T-cell progenitors when compared to human (h)IL-7 [13]. Given its role as key factor for lymphocyte survival and proliferation [10-13], unrestricted supply of IL-7 might result in unwanted effects including lymphoma generation [14]. Further, excessive amounts of mIL-7 limits T cell differentiation by interfering with Notch signaling [15, 16]. Here, we generated hIL-7 (hIL-7) BAC transgenic NSGW41 mice to improve T cell differentiation in humanized mice while simultaneously avoiding unwanted effects caused by excessive and spatially unrestricted availability of hIL-7.

Results and Discussion

To generate a mouse model with tissue-specific expression of human (h)IL-7, we inserted cDNA encoding *IL7* into a BAC containing regulatory elements of the murine

Il7 locus (Fig. 1a). BAC transgenic mice were generated on the NOD background,
crossed into the NSGW41 strain [2] and termed NSGW41hIL7. NSGW41hIL7 mice
contain three copies of the BAC transgene and expressed hIL-7 mRNA and protein in
BM, spleen and thymus (Fig. 1b,c). Engraftment of human HSPCs and T cell
differentiation was determined at indicated time points after transplantation of CD34^{+}-
enriched cord blood cells into unconditioned NSGW41 or NSGW41hIL7 mice (Fig.
1d). Blood from NSGW41 or NSGW41hIL7 mice contained comparable levels of
human hematopoietic cells peaking at 16-18 weeks after transplantation (Fig. 1e).
However, beginning at week 16 after transplantation frequencies of T cells among
human CD45^{+} cells were significantly increased in NSGW41hIL7 mice compared to
NSGW41 recipients. Correspondingly, ratios of T and B cells increased progressively
over time, with T cells ultimately becoming the predominant lymphocyte population in
blood of NSGW41hIL7 mice (Fig. 1e,f). T/B ratios >1 are also observed in human
blood. Consistently, in spleens, human CD45^{+} leukocytes were increased in
NSGW41hIL7 mice compared to NSGW41 mice, which was mainly attributable to
increased numbers of T cells (Fig. S1a,b). The data indicate that expression of hIL-7
under control of endogenous gene regulatory elements fosters T-cell differentiation in
NSGW41hIL7 mice.

Next, we assessed whether elevated frequencies of peripheral human T cells in
NSGW41hIL7 mice were due to more efficient intrathymic T-cell differentiation.
Human CD45^{+} cell numbers were 3.3-fold, 3.5-fold, and 21.2-fold higher in thymi
from NSGW41hIL7 mice at 15, 18, and 26 weeks after reconstitution, respectively
(Fig. 2a). Ratios of human CD4/CD8 double negative (DN), double positive (DP), and
CD4 and CD8 single positive (SP) thymocytes were comparable in both recipient
lines 15 and 18 weeks after transplantation, indicating that ectopic expression of hIL-
7 did not result in aberrant T cell differentiation (Fig. 2b,c). NSGW41hIL7 but not
NSGW41 thymi predominantly contained DP thymocytes 26-32 weeks after
transplantation. In contrast, DN thymocytes constituted the major population in
NSGW41 mice, suggesting that hIL-7 supports human T cell differentiation for
extended periods of time in NSGW41hIL7 mice. A recently characterized combined
knock-in of hIL-7 and hIL-15 on the NSG background displayed massive skewing
towards CD8 SP cells at the expense of DP thymocytes [17], suggesting that tissue-
specific expression of human cytokines alone is insufficient to promote human T cell

differentiation [18]. Consistent with the T-lineage specific role of hIL-7 in human

hematopoiesis, and confirming specificity of transgenic expression of hIL-7, we

observed no alterations in B-cell differentiation in NSGW41hIL7 mice compared to

NSGW41 (Fig. 2d, S2a,b). We conclude that NSGW41hIL7 mice display improved

and extended intrathymic T cell differentiation from human cord blood-derived

HSPCs, temporally coinciding with a shift in T/B cell ratios in the periphery.

Elevated levels of human T-cell precursors in thymus suggested that the increased

frequencies of T cells in blood and spleen were largely reflecting increased thymic

output. However, hIL-7 might also alter peripheral homeostasis. To test this

possibility we further characterized peripheral T cell subsets in NSGW41hIL7 mice.

CD4$^+$ and CD8$^+$ T cell numbers in the blood were increased to comparable extent

(Fig. 3a). Within CD4$^+$ T cells, frequencies of naive T cells and Recent Thymic

Emigrants (RTEs) and effector T cells increased in NSGW41hIL7 mice compared to

NSGW41 (Fig. 3b,S3). Concomitantly, frequencies of effector memory T cells were

reduced. In CD8$^+$ T cells, a similar increase in naive and RTE and decrease in

effector memory frequencies was observed, whereas frequencies of T effector and

central memory subsets remained comparable to those detected in NSGW41 mice

(Fig. 3b, S3). Together, the relative contribution of different subsets to human T cells

in NSGW41hIL7 mice resembled the contribution in human peripheral blood. Next,

we analyzed T cell receptor (TCR) repertoire diversity using next-generation

sequencing to assess possible post-thymic peripheral expansion of T cells. This

analysis revealed a high frequency of rare T cell clones, which were comparable

between NSGW41 and NSGW41hIL7 mice. Virtually no expanded clones were

observed, indicating the absence of hIL-7-induced lymphoproliferation (Fig. 3c). We

conclude from these data that elevated peripheral frequencies of human T cells in

NSGW41hIL7 mice are largely generated through enhanced intrathymic T-cell

differentiation. Furthermore, this route of differentiation results in a composition of the

peripheral T-cell compartment largely resembling human peripheral blood. Paucity of

peripheral lymph nodes (LN), including mesenteric (m)LN, remains a critical limitation

of current NSG-derived humanized mouse models. In NSGW41hIL7 mice mLN were

increased in number and individual size compared to NSGW41 mice (Fig. 3d). Given

the poor regeneration of lymph nodes in other strains of humanized mice, this

observation suggests that NSGW41hIL7 mice might constitute an improved model for

studying gut-associated immune responses. Next, we characterized the functionality
of conventional human T cells in NSGW41hIL7 mice. To this end, we isolated and
activated *ex vivo* CD3[+] T cells from NSGW41 or NSGW41hIL7 spleens. Upregulation
of the bona fide activation markers CD25 and CD69 indicated similar levels of
activation (Fig. S4a). Finally, NSGW41hIL7-derived CD4[+] and CD8[+] T cells displayed
increased divisions in response to anti-CD3/CD28 or phytohemagglutinin stimulation
(Fig. 3e, S4). We conclude that human T cells differentiated in NSGW41hIL7 mice
respond efficiently to T cell receptor triggering *ex vivo*.

Regulatory T (Treg) cells are central in providing protection from autoreactive T cells
[19]. However, in the context of malignancy Treg-cell mediated immunosuppression
precludes effective anti-tumor immune responses [20]. Within the total T cell pool, Treg
cells constitute a comparatively small population, making their functional analysis
difficult in extant humanized mouse models with low overall T cell numbers. We
observed similar frequencies of regulatory T (Treg) cells in multiple organs from
NSGW41hIL7 or NSGW41 mice (Fig. 4a). However, given the higher overall T cell
numbers in NSGW41hIL7 mice, Treg cell numbers were increased making their
analysis feasible (Fig. 4a). To study the activation of Treg cells in humanized
NSGW41hIL7 mice, we transplanted porcine pancreatic islets into the portal vein of
humanized NSGW41hIL7 mice [4]. 18 hours after xenotransplantation, Treg cells in the
liver but not spleen displayed significantly increased expression of HLA-DR
evidencing site-specific activation through the xenograft (Fig. 4b). We conclude that
humanized NSGW41hIL7 mice represent a good model for the study of human Treg
cells *in vivo*.

Taken together, we have demonstrated that spatially restricted expression of hIL-7 in
NSGW41hIL7 mice results in an improved capacity for T-cell differentiation from
human HSPCs. Thus, we observed T/B cell ratios geared towards a higher
abundance of T cells when compared to NSGW41 and other humanized mouse
models [21]. Notably, increased intrathymic T-cell development promoted the
generation of a peripheral T-cell pool with a high TCR diversity and an overall
composition of T-cell subpopulations reminiscent of human peripheral blood.
Nevertheless, thymus size and cellularity in humanized NSGW41hIL7 mice remained
smaller than that of wild-type mice, suggesting that additional factors, presumably

depending on lymphocyte-stromal cell interactions, are required to generate a
normal-sized thymic microenvironment after humanization. It has been shown that
such a limitation can be partially overcome by co-transplantation of *in vitro*-
differentiated proT cells [22].

We detected no evidence for hIL-7-driven lymphoproliferation. We designed
NSGW41hIL7 mice to express hIL-7 using regulatory elements that allowed to
faithfully identify cells expressing mIL-7 in an earlier study [23]. Our study indicates that
this expression pattern and the observed levels of expression of hIL-7 are able to
strike a balance to overcome the limited potency of mIL-7 on human lymphocytes,
while at the same time avoiding detrimental effects of aberrant expression of hIL-7 [13,
14].

Enhanced T-cell differentiation in NSGW41hIL7 mice allowed for analysis of T cell
subsets, including Treg cells. In addition, size and number of mLN were increased.
Thus, humanized NSGW41hIL7 mice have the potential to foster investigation of rare
human T-populations in vivo as well as gut-associated immunity.

Methods

Generation of NSGW41hIL7 mice: NODhIL7 mice were generated by pronuclear
injection of a BAC containing codon-optimized cDNA of *hIL7* introduced at the 3' end
of the 5'UTR of the *Il7* gene flanked by 96kb upstream, and the entire *Il7* locus plus
an additional 17kb downstream. The BAC was constructed according to a described
strategy [23]. BAC integrity upon integration and copy numbers were determined by
PCR. Offspring showing detectable expression of hIL-7 mRNA was crossed with
NSGW41 mice [2] to generate the NSGW41hIL7 strain. All animal experiments were
performed in accordance with German animal welfare legislation and were approved
by the relevant authorities: Landesdirektion Dresden, the Thüringer Landesamt für
Verbraucherschutz, the Niedersächsisches Landesamt für Verbraucherschutz und
Lebensmittelsicherheit (LAVES), and the Regierungspräsidium Darmstadt.

Human HSPC transplantation: Cord blood samples were obtained from the
Department Obstetrics, Gynecology and Reproductive Medicine, Hannover Medical
School, Hannover, from the Bürgerhospital Frankfurt am Main, and from the DKMS
Cord Blood Bank, Dresden, and were used in accordance with the guidelines
approved by the ethics committees of Hannover Medical School, Frankfurt University
Clinics, Dresden University of Technology, and University Clinics Jena. CD34[+]

HSPCs were isolated using dual magnetic beads enrichment according to the
manufacturer's instructions (Miltenyi Biotech) [2]. Purities >95% were considered
acceptable. Contaminating T cell frequencies were routinely below 1%.

<u>Flow cytometry</u>: Analysis was performed as described before [2]. A full list of antibody
panels is provided in Table S1.

<u>TCR repertoire analysis</u>: After mRNA isolation (Qiagen Micro Kit), cDNA was
generated via the Smarter 5'RACE cDNA amplification kit (Clontech) using 4.5µl
mRNA input and following the recommended protocol. Complementarity-determining
region 3 (CDR3) regions of the human TRB locus were amplified through a gene-
specific primer (2µM final concentration) that targets the constant region of the beta
(β)-chain (GCACACCAGTGTGGCCTTTTGGG) and a primer (1µM final
concentration) binding to the introduced SMARTER oligonucleotide
(CTAATACGACTCACTATAGGGC) using the Advantage 2 PCR kit (Clontech) in a
50µl reaction. Both primer sequences further contain 16S Illumina overhang adapter
sequences. Cycling conditions were as following: 120 s 95°C; 30 times 30 s 95°C, 45
s 64°C, 60 s 72°C; 60 s 72°C. Generated PCR amplicons were agarose gel purified
(Qiagen GelExtract.) Next, PCR samples were indexed with Nextera Illumina Indices
reads using the Advantage 2 PCR kit (Clontech) in a 8 PCR cycle reaction and
purified with Agencourt AMPpure XP beads (Beckman Coulter) according to the
manufacturers protocol. Samples were pooled, denatured and subjected to Illumina
MiSeq analysis using 500 cycles and paired-end sequencing following Illumina
guidelines. Sequencing libraries contained 20% PhIX for library complexity.
Demultiplexed Fastq files were annotated to the human TRB locus via MiXCR
software [18]. Individual CDR3 nucleotide sequences were ranked according to their
abundance within the respective samples and further analyzed using VDJTools [24]
and TcR [25]. TCR repertoire data are available at SRA
(https://www.ncbi.nlm.nih.gov/sra), accession number PRJNA606460.

<u>Islet xenotransplantation</u>: 500 IEQ (Islets Equivalent) adult pig islets or PBS were
transplanted in the portal vein of NSGW41hIL7 mice that were humanized 24 weeks
before. Islets were obtained from Goettingen minipigs (Ellgard) as described before
[26, 27]. Human blood cell chimerism of used mice was 32-85%. Regulatory T cells were
analyzed 18 hours after surgery.

<u>T cell co-stimulation</u>: hCD3[+] T cells were isolated from the spleen of humanized
NSGW41 or NSGW41hIL7 mice 16 to 25 weeks after HSPC transplantation,

enriched using negative depletion (Miltenyi), and CPD labeled (eBioscience). 10^5

hCD3$^+$ T cells were mixed with human T-Activator CD3/CD28 Dynabeads

(Thermofisher, ratio 3:1) or PHA (1µg/mL) in RPMI 10% FCS, 20mM L-glutamine,

10mM Hepes, 1mM Sodium Pyruvate, 50µM ß-mercaptoethanol with recombinant

hIL-2 (30 U/mL) and incubated for 6 days (37°C, 5% CO2).

Statistics: Student's t tests were performed for all statistical analyses using Prism 8

for MacOSX software. In all graphs *p = 0.05–0.01, **p = 0.01–0.001, and ***p =

0.001- 0.0001 and ****p < 0.0001; data represent the mean ± SD. Boxes and

whiskers display the data distribution through their quartiles.

References

1.   Shultz, L.D. *et al.* Humanized mouse models of immunological diseases and

precision medicine. *Mammalian genome : official journal of the International*

Mammalian Genome Society 30, 123-142 (2019).

2.   Cosgun, K.N. *et al.* Kit Regulates HSC Engraftment across the Human-Mouse

Species Barrier. *Cell Stem Cell* 15, 227-238 (2014).

3.   Mende, N. *et al.* CCND1-CDK4-mediated cell cycle progression provides a

competitive advantage for human hematopoietic stem cells in vivo. *J Exp Med*

212, 1171-1183 (2015).

4.   Rahmig, S., Bornstein, S.R., Chavakis, T., Jaeckel, E. & Waskow, C.

Humanized mouse models for type 1 diabetes including pancreatic islet

transplantation. *Hormone and metabolic research = Hormon- und*

Stoffwechselforschung = Hormones et metabolisme 47, 43-47 (2015).

5.   Rahmig, S. *et al.* Improved human erythropoiesis and platelet formation in

humanized NSGW41 mice. *Stem Cell Reports* 7, 591-601 (2016).

6.   Yurino, A. *et al.* Enhanced Reconstitution of Human Erythropoiesis and

Thrombopoiesis in an Immunodeficient Mouse Model with Kit(Wv) Mutations.

Stem Cell Reports 7, 425-438 (2016).

7.   Nicolini, F.E., Cashman, J.D., Hogge, D.E., Humphries, R.K. & Eaves, C.J.

NOD/SCID mice engineered to express human IL-3, GM-CSF and Steel factor

constitutively mobilize engrafted human progenitors and compromise human

stem cell regeneration. *Leukemia* 18, 341-347 (2004).

8.   Rongvaux, A. *et al.* Development and function of human innate immune cells

in a humanized mouse model. *Nat Biotechnol* 32, 364-372 (2014).

9.   Yu, H. *et al.* A novel humanized mouse model with significant improvement of
class-switched, antigen-specific antibody production. *Blood* 129, 959-969
(2017).

10.   Puel, A., Ziegler, S.F., Buckley, R.H. & Leonard, W.J. Defective IL7R
expression in T(-)B(+)NK(+) severe combined immunodeficiency. *Nat Genet*
20, 394-397 (1998).

11.   Peschon, J.J. *et al.* Early lymphocyte expansion is severely impaired in
interleukin 7 receptor-deficient mice. *J Exp Med* 180, 1955-1960. (1994).

12.   von Freeden-Jeffry, U. *et al.* Lymphopenia in interleukin (IL)-7 gene-deleted
mice identifies IL-7 as a nonredundant cytokine. *J Exp Med* 181, 1519-1526.
(1995).

13.   van Lent, A.U. *et al.* IL-7 enhances thymic human T cell development in
"human immune system" Rag2-/-IL-2Rgammac-/- mice without affecting
peripheral T cell homeostasis. *J Immunol* 183, 7645-7655 (2009).

14.   Rich, B.E., Campos-Torres, J., Tepper, R.I., Moreadith, R.W. & Leder, P.
Cutaneous lymphoproliferation and lymphomas in interleukin 7 transgenic
mice. *J Exp Med* 177, 305-316 (1993).

15.   El-Kassar, N. *et al.* High levels of IL-7 cause dysregulation of thymocyte
development. *Int Immunol* 24, 661-671 (2012).

16.   El Kassar, N. *et al.* A dose effect of IL-7 on thymocyte development. *Blood*
104, 1419-1427 (2004).

17.   Matsuda, M. *et al.* Human NK cell development in hIL-7 and hIL-15 knockin
NOD/SCID/IL2rgKO mice. *Life science alliance* 2 (2019).

18.   Bolotin, D.A. *et al.* MiXCR: software for comprehensive adaptive immunity
profiling. *Nat Methods* 12, 380-381 (2015).

19.   Sakaguchi, S. *et al.* Regulatory T Cells and Human Disease. *Annu Rev*
Immunol (2020).

20.   Yano, H., Andrews, L.P., Workman, C.J. & Vignali, D.A.A. Intratumoral
regulatory T cells: markers, subsets and their impact on anti-tumor immunity.
Immunology 157, 232-247 (2019).

21.   Sippel, T.R., Radtke, S., Olsen, T.M., Kiem, H.P. & Rongvaux, A. Human
hematopoietic stem cell maintenance and myeloid cell development in next-
generation humanized mouse models. *Blood advances* 3, 268-274 (2019).

22. Awong, G. *et al.* Human proT-cells generated in vitro facilitate hematopoietic
stem cell-derived T-lymphopoiesis in vivo and restore thymic architecture.
Blood 122, 4210-4219 (2013).

23. Repass, J.F. *et al.* IL7-hCD25 and IL7-Cre BAC transgenic mouse lines: new
tools for analysis of IL-7 expressing cells. *Genesis* 47, 281-287 (2009).

24. Shugay, M. *et al.* VDJtools: Unifying Post-analysis of T Cell Receptor
Repertoires. *PLoS computational biology* 11, e1004503 (2015).

25. Nazarov, V.I. *et al.* tcR: an R package for T cell receptor repertoire advanced
data analysis. *BMC bioinformatics* 16, 175 (2015).

26. Ludwig, B. *et al.* Favorable outcome of experimental islet xenotransplantation
without immunosuppression in a nonhuman primate model of diabetes. *Proc*
Natl Acad Sci U S A 114, 11745-11750 (2017).

27. Steffen, A. *et al.* Production of high-quality islets from goettingen minipigs:
Choice of organ preservation solution, donor pool, and optimal cold ischemia
time. *Xenotransplantation* 24 (2017).

352
353

Acknowledgements

We are grateful to Katrin Witzlau and Esther Imelmann for technical assistance and
management of mouse colonies. We would like to thank Dr. Ellen Richie (MD
Anderson Cancer Center, TX, USA) for providing BAC constructs and advice on
cloning. The work was supported by grants from the German Research Foundation
(DFG, SFB738-A7 and SFB902-B15) (to A.K.) and FOR2033-A03, TRR127-A5,
WA2837/6-1 and WA2837/7-1 (to C.W.).

Authorship contributions

E.C. planned, conducted, interpreted experiments and wrote the paper. B.S. planned,
conducted and interpreted experiments. S.Rah, J.B., and N.V. conducted
experiments. J.B., S.Rav., and I.P. conducted and interpreted experiments on TCR
repertoire. U.S., J.S., S.L. and B.L. isolated and provided porcine islets for
xenotransplanation. F.B., C.v.K, and A.P. provided crucial reagents. R.N. generated
knock-in mice, and C.W. and A.K. conceived the study, planned and interpreted
experiments and wrote the paper.

<u>Disclosure of conflicts of interest</u>

The authors declare that no conflicts of interest exist.

<u>Figure legends</u>

Fig. 1: Enhanced generation of human T cells upon transfer of human HSPCs

in NSGW41hIL7 mice. a, Scheme of BAC construct for the generation of

NSGW41hIL7 mice. **b**, Abundance of hIL7 transcript in bone marrow (BM), spleen

and thymus from humanized NSGW41 or NSGW41hIL7 mice. Each dot represents

an individual mouse. **c**, hIL7 protein levels in thymi isolated from non-humanized

NSGW41 or NSGW41hIL7 mice (top) and in bone marrow, thymus and serum from

NSGW41 or NSGW41hIL7 mice that have received human HPSCs 26-38 weeks

before (bottom). **d**, Scheme of transplantation experiments. **e**, Kinetics of the

appearance of human CD45$^+$ cells (hCD45$^+$, left) and hCD3$^+$ T cells within human

leukocytes (right) in the blood after humanization. Each dot represents an individual

mouse. **f**, Frequencies of T cells, B cells and non-defined other cells of human origin

in the blood of NSGW41 (n=13) or NSGW41hIL7 (n=16) mice 18 weeks after

humanization. Numbers on top indicate T vs. B cell ratios.

Fig. 2: Efficient and prolonged intrathymic T-cell differentiation in humanized

NSGW41hIL7 mice. a, Numbers (top) and fold-change (bottom) of human CD45$^+$

cells in thymi of humanized NSGW41 or NSGW41hIL7 mice at the indicated time

points after humanization. Fold-changes were calculated by dividing human CD45$^+$

thymocyte numbers from humanized NSGW41hIL7 mice through the thymocyte

numbers from humanized NSGW41 mice. This was done separately for each

experiment and the results pooled. **b**, Analysis of CD4 and CD8 expression on

hCD45$^+$ thymocytes from NSGW41 or NSGW41hIL7 mice that have received human

HPSCs 15 weeks (left) or 26 (right) weeks before. **c**, Distribution of thymocyte

subsets in NSGW41 or NSGW41hIL7 mice at the indicated time points after

humanization. **d**, Numbers of B cells subsets in the bone marrow of NSGW41 or

NSGW41hIL7 mice 26 weeks after humanization.

Fig. 3: hIL7-BAC transgene increases functional peripheral T cells in the

absence of excessive lymphoproliferation. a, Human CD4$^+$ and CD8$^+$ T cells in

the blood of humanized NSGW41 or NSGW41hIL7 mice 26 weeks after

humanization. **b**, Composition of blood CD4$^+$ (top) and CD8$^+$ (bottom) T cell subpopulations 26 weeks after humanization: Naïve T cells, central memory (T$_{CM}$), effector memory (T$_{EM}$), T effector (T$_{EFF}$) and recent thymic emigrants (RTE) in NSGW41, NSGW41hIL7 mice, or human blood. **c**, T cell receptor (TCR) repertoire diversity in splenic αβ T cells of NSGW41 or NSGW41hIL7 mice. Clones were binned into rare (0<X≤0.001), small (0.001<X≤0.01), medium (0.01<X≤0.1) and expanded (0.1<X≤1) (n=3 per group). **d**, Photographs of mLN from humanized NSGW41 or NSGW41hIL7 mice isolated 26 weeks after humanization, or C57BL/6 controls (top). mLN number per mouse (bottom left). hCD45$^+$ and hCD3$^+$ cell numbers in mLN from NSGW41 or NSGW41hIL7 mice isolated 26 weeks after humanization (bottom right). **e**, Activation of human T cells from NSGW41 or NSGW41hIL7 mice. Histograms depict division of CDP-labeled spleen hCD3$^+$ T cells 6 days after stimulation with CD3/28 beads or control (w/o, left). Frequencies of non-divided human T cells and T cells that have divided 4, 5 or 6 times 6 days after stimulation with CD3/28 antibody-coated beads or controls (right). Top: CD4$^+$ T cells, bottom: CD8$^+$ T cells.

Fig. 4: Efficient generation of human Treg cells in NSGW41hIL7 mice. a, Representative dot plots analyzing mesenteric lymph nodes (mLN) and blood (left) and numbers (right, top) and frequencies (right, bottom) of Tregs in mLN, blood, spleen and liver of humanized NSGW41 or NSGW41hIL7 mice. **b**, Human activated Tregs after intraportal xenotransplantation of porcine pancreatic islets into NSGW41hIL7 mice 26 weeks after humanization. Representative dot plots of liver and spleen analysis (left). Frequencies of HLA-DR$^+$ FoxP3$^+$ T cells in spleen and liver of humanized NSGW41hIL7 mice (right) 18 hours after transplantation of islets (iTx) or PBS.

Figure 1

Figure 2

Figure 3

a

b

c

d

mLN

NSGW41

NSGW41hIL7

C57BL/6

Medium
Expanded

NSGW41hIL7

w/o   CD3/28

DAPI⁻ hCD45⁺   CD3⁺ CD4⁺

51%   17%

49%   8%

events

CPD

non-divided T cells

T cells after division

% CD4⁺

% CD8⁺

w/o   CD3/28

Naive T_{CM} T_{EM} T_{EFF} RTE

NSGW41   NSGW41hIL7   Human blood

Figure 4

a

Supplemental Material

Enhanced differentiation of functional human T cells in NSGW41 mice with

tissue-specific expression of human interleukin-7.

Emilie Coppin[1,2*], Bala Sai Sundarasetty[3*§], Susann Rahmig[1,2], Jonas Blume[4], Nikita

A. Verheyden[3], Franz Bahlmann[5], Sarina Ravens[4], Undine Schubert[9,10], Janine

Schmid[9,10], Stefan Ludwig[11], Constantin von Kaisenberg[6], Alexander Platz[7], Ronald

Naumann[8], Barbara Ludwig[9,10], Immo Prinz[4], Claudia Waskow[1,2,9,12$], Andreas

Krueger[3$]

Table S1

Antibodies	Clone	Provider
CD3 APC, AF780, APC Cy7	UCHT1, HIT3a	eBioscience
CD4 PE Cy7, Pacific Blue	OKT-4/RPA-T4	eBioscience
CD8 PE Cy5, PerCP, PE Cy7	SK1, HIT8a	eBioscience
CD10 APC Cy7	HI10a	BioLegend
CD14 AF700	M5E2	BD Pharmingen
CD16 PerCP, PE Cy5	3G8	BioLegend, BD Biosciences
CD19 FITC, PC7, APC, FITC	HIB19	eBioscience
CD25 PE	M-A251	BioLegend, BD Biosciences
CD31 PE	MEM-05	Immunotools
CD33 PE, PE Cy7	WM-53	eBioscience
CD34 PE Cy7, Pacific Blue	OKT-4/RPA-T4	eBioscience
mCD45 AF780, AF700	RA3-6B2	eBioscience
hCD45 eF450, FITC, PE Cy7, PerCP	HI30	eBioscience, BioLegend
CD45RA eF450	HI100	eBioscience
CD45RO FITC, PE	UCHL1	BD Biosciences
CD62L Biotin	DREG.55	eBioscience
CD69 PerCP Cy5.5	Y1/82A	BioLegend
CD197 APC, BV510	REA337, G043H7	Miltenyi
CD235 FITC	GA-R2 (HIR2)	eBioscience
Foxp3 APC	PCH101	eBioscience
HLA DR PerCP Cy5.5	4S.B3	BioLegend
IgD PE	IA6-2	BD Pharmingen
IgM Biotin	SA-DA4	BioLegend
Ter119 FITC	11-5921-82	eBioscience
Streptavidin V500, FITC	na	BD Biosciences, BD Pharmingen
Cell Proliferation Dye eF450	na	ebioscience
Fixabe viability dye AF700	na	BD Biosciences

Supplementary figures

Supplementary Figure 1: Human lymphocyte composition in spleen. (**a**) Numbers of hCD45[+] leukocytes (top) and hCD3[+] T cells (bottom) in spleens of NSGW41 or NSGW41hIL7 mice 15, 18 and 26 weeks after humanization. (**b**) Frequencies of hCD3[+], hCD19[+], and other cells within hCD45[+] leukocytes in spleens of NSGW41 or NSGW41hIL7 mice 15, 18 and 26 weeks after humanization. Numbers on top of graphs indicate T vs. B cell ratio.

 Supplementary Figure 2: B lymphopoiesis in NSGW41hIL7 mice. (**a**) Dot plots
32 show the gating strategy of bone marrow human B lineage subsets: pro-B (hCD45$^+$
33 CD19$^+$ CD10$^+$ IgM$^-$ CD34$^+$), pre-B (hCD45$^+$ CD19$^+$ CD10$^+$ IgM$^-$ CD34$^-$), immature B
34 (hCD45$^+$ CD19$^+$ CD10$^+$ IgM$^+$ CD34$^-$) and mature B cells (hCD45$^+$ CD19$^+$ CD10$^-$ IgM$^+$
35 IgD$^+$). (**b**) Frequencies of B cell subsets in bone marrow of humanized mice 26
36 weeks after humanization.

T_{EFF} = Effectors
RTE = Recent Thymic Emigrants
T_{CM} = Central Memory
T_{EM} = Effector Memory

Supplementary Figure 3: Characterization of human T cell subpopulations in the blood. Gating strategy and identification of hCD4$^+$ and hCD8$^+$ T cell subpopulations in blood of humanized mice: T effectors (T_{EFF}, hCD45$^+$ CD3$^+$ CD4$^+$ CCR7$^-$ CD45RA$^+$), central memory (T_{CM}, hCD45$^+$ CD3$^+$ CD4$^+$ CCR7$^+$ CD45RA$^-$), effector memory (T_{EM}, hCD45$^+$ CD3$^+$ CD4$^+$ CCR7$^-$ CD45RA$^-$), naïve T (Naïve, hCD45$^+$ CD3$^+$ CD4$^+$ CCR7$^+$ CD45RA$^+$) and recent thymic emigrants (RTE, hCD45$^+$ CD3$^+$ CD4$^+$ CD31$^+$ CD45RA$^+$). Data was acquired 26 weeks after humanization. The composition of subpopulations in human blood is shown for comparison (bottom).

Supplementary Figure 4: Activation of human T cells from NSGW41 or
NSW41hIL7 mice. (**a**) Representative dot plots showing the expression of hCD25
and hCD69 on hCD3$^+$ T cells isolated from NSGW41 (top) or NSGW41hIL7 (bottom)
spleens after 6 days of CD3/28 or PHA stimulation or non-stimulated controls (left).
Donor mice had received human HSPCs 20 to 26 weeks before. Fold-changes were
calculated by dividing the percentages of stimulated hCD25$^+$ or hCD69$^+$ hCD4$^+$ (top)
or hCD8$^+$ (bottom) splenic T cells by the percentages of non-stimulated cells (w/o)
from NSGW41 or NSGW41hIL7 mice (right). (**b**) Percentages of non-divided T cells
and T cells which have undergone 4, 5 or 6 divisions 6 days after stimulation with
phytohemagglutinin (PHA) or without stimulation (w/o). Top: hCD4$^+$ T cells, bottom:
hCD8$^+$ T cells. Donor mice had received human HSPCs 20 to 26 weeks before.