Title

Heterologous production of 1-tuberculosinyladenosine in *Mycobacterium kansasii* models

pathoevolution towards the transcellular lifestyle of *Mycobacterium tuberculosis*.

Authors

Marwan Ghanem[1,2,3,4]*, Jean-Yves Dubé[1,2,3]*, Joyce Wang[1,2,3,5], Fiona McIntosh[1,2,3], Daniel

Houle[2], Pilar Domenech[1,2,3], Michael B. Reed[1,2,3], Sahadevan Raman[6], Jeffrey Buter[7], Adriaan J.

Minnaard[7], D. Branch Moody[6] and Marcel A. Behr[1,2,3]

Affiliations

1. Department of Microbiology and Immunology, Faculty of Medicine, McGill University,

Montreal, Canada. 2. Infectious Disease and Immunity in Global Health Program, Research

Institute of the McGill University Health Centre, Montreal, Canada. 3. McGill International TB

Centre, Montreal, Canada. 4. Faculty of Medicine, American University of Beirut, Lebanon. 5.

Department of Microbiology and Immunology, University of Michigan Medical School, Michigan,

USA. 6. Division of Rheumatology, Immunity and Inflammation, Brigham and Women's Hospital,

Harvard Medical School, Boston, United States of America. 7. Stratingh Institute for Chemistry,

University of Groningen, 9747AG Groningen, The Netherlands. *These authors contributed

equally.

Running title: 1-TbAd complementation in *M. kansasii*

Author for correspondence: Marcel Behr (marcel.behr@mcgill.ca)

Byline: Ghanem M*, Dube JY*, Wang J, McIntosh F, Houle D, Domenech P, Reed MB, Raman S,

Buter J, Minnaard AJ, Moody DB and Behr MA. 2020. Heterologous production of 1-

tuberculosinyladenosine in *Mycobacterium kansasii* models pathoevolution towards the

transcellular lifestyle of *Mycobacterium tuberculosis*.

ABSTRACT

Mycobacterium kansasii is an environmental non-tuberculous mycobacterium that

causes opportunistic tuberculosis-like disease. It is one of the most closely related species to the

M. tuberculosis complex. Using *M. kansasii* as a proxy for the *M. kansasii-M. tuberculosis*-

common ancestor, we asked whether introducing the *M. tuberculosis*-specific gene pair *Rv3377c-*

Rv3378c into *M. kansasii* affects the course of experimental infection. Expression of these genes

resulted in the production of an adenosine-linked lipid species, known as 1-

tuberculosinyladenosine (1-TbAd), but did not alter growth *in vitro* under standard conditions.

Production of 1-TbAd enhanced growth of *M. kansasii* under acidic conditions through a bacterial

cell-intrinsic mechanism independent of controlling pH in the bulk extracellular and intracellular

spaces. Production of 1-TbAd led to greater burden of *M. kansasii* in the lung of C57Bl/6 mice

during the first 24 hours after infection and *ex vivo* infections of alveolar macrophages

recapitulated this phenotype within the same time frame. However, in long-term infections,

production of 1-TbAd resulted in impaired bacterial survival in both C57Bl/6 mice and *Ccr2-/-*

mice. We have demonstrated that *M. kansasii* is a valid surrogate of *M. tuberculosis* to study

virulence factors acquired by the latter organism, yet shown the challenge inherent to studying

the complex evolution of mycobacterial pathogenicity with isolated gene complementation.

IMPORTANCE

49   This work sheds light on the role of the lipid 1-tuberculosinyladenosine in the evolution

50   of an environmental ancestor to *M. tuberculosis*. On a larger scale, it reinforces the importance

51   of horizontal gene transfer in bacterial evolution and examines novel models and methods to

52   provide a better understanding of the subtle effects of individual *M. tuberculosis*-specific

53   virulence factors in infection settings that are relevant to the pathogen.

INTRODUCTION

M. tuberculosis virulence factors have been established by genetic knock-out and

complementation within the pathogen, producing evidence of an attenuation of virulence in *ex*

vivo or *in vivo* experimental infections (1, 2). While numerous virulence-associated loci have been

identified with this approach, the majority of these are intact in the genomes of non-

transmissible environmental mycobacteria, such as *M. kansasii* (3-6). *M. kansasii* is readily

isolated in clinical settings from pulmonary infections and we have previously shown that it can

be studied in an experimental lung model (3). However, although it causes TB-like disease, *M.*

kansasii infections disproportionately affect patients with underlying pulmonary diseases or

immunosuppression, and there is no evidence supporting its transmission between individuals

(7, 8). The conservation of many virulence factors across *M. tuberculosis* and *M. kansasii*,

including the ESX-1 secretion system, PhoPR 2-component system and DosR/S/T regulon suggest

that they play a role in a broader survival strategy used by mycobacteria (9-11). For example,

some of these factors may be needed for survival of *M. kansasii* within free-living phagocytic

amoeba, but their presence does not provide *M. kansasii* with the pathogenic capabilities of *M.*

tuberculosis inside human hosts (12). Consequently, there is currently an incomplete

understanding of how *M. tuberculosis* emerged as a human-adapted professional pathogen.

It was previously thought that horizontal gene transfer (HGT) does not occur in

mycobacteria. However, evidence accumulated over the past decade has shown that HGT events

have happened during mycobacterial speciation and are associated with the step-wise

emergence of pathogenic species (13-16). Fifty-five genes have been acquired by *M. tuberculosis*

since its divergence from the *M. kansasii-M. tuberculosis* common ancestor (MKMTCA) (13).

Although many of these HGT genes have no postulated function, the *Rv3376-Rv3378c* genomic

island uniquely present in *M. tuberculosis* is known to encode a class II terpene cyclase (*Rv3377c*)

and a tuberculosinyl transferase (*Rv3378c*). Together, the two enzymes are responsible for the

conversion of geranylgeranyl pyrophosphate (GGPP) into the recently identified *M. tuberculosis-*

specific lipid 1-tuberculosinyladenosine (1-TbAd), which is a potential diagnostic molecular

marker for TB disease (17-20). 1-TbAd further undergoes a chemical rearrangement, known as

the Dimroth reaction, to generate N^6-TbAd (18). While GGPP is found in both species and used

as an intermediate in the biosynthesis of 1-TbAd by *M. tuberculosis* (17, 19), it is part of the

biosynthetic pathway for the production of carotenoid pigments of *M. kansasii*, giving its

characteristic yellow colour (21).

We previously reported the important role of 1-TbAd in protecting *M. tuberculosis* from

phagosomal acidification inside macrophages (22). In the present work, we have characterized

the effect of 1-TbAd production in *M. kansasii* complemented with *Rv3377c and Rv3378c*. Here,

we show that *in vitro* growth kinetics and colony morphology in liquid and on solid media are

unaltered by *Rv3377c-Rv3378c* expression. 1-TbAd confers a growth advantage in acidic media

compared to wild-type *M. kansasii*, which we further demonstrated to be independent of

cytosolic and culture medium pH control, suggesting a compartmental mechanism of protection

for the bacterium. *Rv3377c-Rv3378c* provided an early advantage to bacterial replication during

pulmonary infection in mice, consistent with enhanced survival in alveolar macrophages.

However, the *M. kansasii:Rv3377-78c* was outcompeted by the wild-type during long-term

murine infection. This study demonstrates that we can use *M. kansasii* as a proxy of the MKMTCA

to explore the complex evolution of *M. tuberculosis*. It also shows that gene acquisition likely

provided advantages for specific contexts, including a possible and unexpected role in survival

within alveolar macrophages during early stages of infection, despite tradeoffs or challenges

under other circumstances requiring further evolution to overcome.

RESULTS

Expression of *Rv3377c-Rv3378c* in *M. kansasii* leads to 1-TbAd production

We introduced the *M. tuberculosis*-specific gene pair *Rv3377c-Rv3378c* into the *M.*

kansasii genome within an integrative plasmid containing hygromycin resistance to produce *M.*

kansasii::Rv3377-78c (see methods). As a control for subsequent experiments, an integrative

'empty vector' (EV) was employed. After labelling with ^{14}C-adenosine and lipid extraction using

chloroform and methanol, radio-thin-layer chromatography (TLC) was used to detect adenosine-

linked lipids in *M. kansasii:Rv3377-78c* clones for comparison to *M. tuberculosis* strain H37Rv

(Fig. 1a). Conventional molybdenum-based sprays followed by charring broadly detected all lipids

as a loading control (Fig. 1b). Whereas uncomplemented bacterial extracts showed material at

the origin and one weak band in radio-TLC, *Rv3377c-Rv3378c* complementation generated at

least 5 additional lipid species. Three of these novel lipids co-migrated with compounds from *M.*

tuberculosis strain H37Rv. Both results strongly suggested the successful genetic transfer of *M.*

tuberculosis-associated adenosine-linked lipids to *M. kansasii.* 5% phosphomolybdic acid reagent

(PMA) staining showed that similar amounts of total lipids were spotted for each *M. kansasii*

sample (Fig. 1b).

High-performance liquid chromatography-mass spectrometry (HPLC-MS) was used to

chemically identify the compounds produced by *M. kansasii::Rv3377-78c*, in which the expected

retention of 1-TbAd (22.7 min) and N^6-TbAd (5.8 min) were known (Fig. 1c-g). Whereas the *M.*

kansasii::EV control did not release compounds that comigrated with either 1-TbAd or N^6-TbAd,

M. kansasii:Rv3377-78c produced high intensity (6.7-7.0 x10^6 counts) signals (*m/z* 540.354)

matching the expected retention time and mass (*m/z* 540.3544) of the proton adducts of 1-TbAd

and N^6-TbAd. The extractions were performed at a range of pH (4.5 – 7.4) since both the Dimroth

reaction that generates N^6-TbAd and the capture of lysosomotropic agents are sensitive to pH,

as previously explained (18, 22, 23). However, there was no clear impact of altering pre-

extraction pH for two hours to the release nor relative abundance of 1-TbAd and N^6-TbAd,

suggesting that such effects, if existent, did not occur under the tested conditions (Fig. 1c-g).

Similar to patterns observed from *M. tuberculosis*, more 1-TbAd than N^6-TbAd was recovered

from *M. kansasii*::*Rv3377-78c* (Fig. 1f-g).

1-TbAd production does not change visible characteristics of *M. kansasii* in conventional media

To characterize any overt phenotypic effects of 1-TbAd production on *M. kansasii*, we assessed its influence on *in vitro* characteristics of the bacterial culture. *M. kansasii*::*Rv3377-78c* grew similarly to wild-type *M. kansasii* and *M. kansasii*::EV in 7H9 (Fig. 2a) and on 7H10 agar (Fig. 2b). Since the production of 1-TbAd requires the same intermediate GGPP as that of the yellow pigment that *M. kansasii* produces when exposed to light, we then tested *M. kansasii*::*Rv3377-78c*'s ability to turn yellow after light exposure in order to rule out the possibility that a pigment-related phenomenon might affect our outcomes (Fig. 2b). *M. kansasii*::*Rv3377-78c* retained photochromogenic abilities by turning yellow after exposure to light at room temperature within the same timeframe as *M. kansasii*::EV.

Congo Red is an amphiphilic dye that binds to the mycobacterial cell membrane. When grown on Congo Red-containing 7H10 plates, different mycobacteria retain the dye distinctly, and this feature has been associated with differences in the interactions of bacterial cells within the colony (24). In this study, *M. tuberculosis* absorbed the dye and became red, while *M. kansasii* colonies remained white on the plate. Since 1-TbAd is found on the cell surface and has an amphipathic character (17), we tested the effect of its production on intra-colony bacterial interactions of *M. kansasii*::*Rv3377-78c*. Visual inspection of colonies (Fig. 2c) and absorbance at 488nm (Fig. 2d) showed no difference in retention of Congo Red between *M. kansasii*::EV and *M. kansasii*::*Rv3377-78c*.

157 **1-TbAd production enhances growth of *M. kansasii* in acidic media**

158 The ability to survive in mildly acidic environments is a key feature of mycobacteria, both

159 environmental and pathogenic (25, 26). It was recently shown that 1-TbAd production in *M.*

160 *tuberculosis* confers a growth advantage over a pH range (5.0-5.4) comparable to that of an

161 activated phagolysosome, which is not tolerated by most bacteria (22). To confirm whether 1-

162 TbAd production provides protection to *M. kansasii* in acidic 7H9 broth, we inoculated *M.*

163 *kansasii*::EV or *M. kansasii::Rv3377-78c* into fresh pH-adjusted 7H9 media (pH 4.9, 5.0, 5.1, 5.2,

164 5.4, 6.6) and monitored their growth. As expected, *M. kansasii::Rv3377-78c* was able to grow in

165 lower pH than *M. kansasii*::EV (Fig. 3a). This effect was observed consistently in five experiments

166 with a growth differential observed between pH 5.1 and 5.4. In addition, as 1-TbAd can be shed

167 extracellularly to de-acidify the phagosomal environment as seen in *M. tuberculosis*, we

168 hypothesized that the production of 1-TbAd by *M. kansasii::Rv3377-78c* may modulate media pH

169 (22). However, at 8 and 17 days post inoculation, both *M. kansasii*::EV and *M.kansasii::Rv3377-*

170 *78c* slightly increased the pH of the media where there was bacterial growth, and to a similar

171 extent (Fig. 3b). Therefore, although we observed enhanced growth with *M. kansasii::Rv3377-*

172 *78c* compared to *M. kansasii*::EV, we could not demonstrate a causative role for 1-TbAd raising

173 the extracellular pH of the culture media under these conditions.

174

175 **Synthetic 1-TbAd does not directly enhance growth of *M. kansasii***

176 Prior work with *M. tuberculosis* estimated that 1-TbAd might naturally accumulate to μM

177 concentrations in phagosomes, and 5-20 μM 1-TbAd alters lysosomal pH and morphology in

178 human macrophages (22). The proposed lysosomotropic mechanism requires that 1-TbAd access

a low pH compartment where the uncharged conjugate base binds protons to raise pH and

regenerate a concentration gradient that promotes further entry of uncharged conjugate base

to the acidic compartment. Whereas this mechanism can relieve pH stress on the bacterium, the

major alternative, which is not exclusive of lysosomotropism, is that 1-TbAd directly signals for

bacterial growth and division. To distinguish these mechanisms we 'chemically complemented'

WT *M. kanasii* with synthetic 1-TbAd and N^6-TbAd added externally in media. In this experiment

TbAd (already carrying a proton) should not alter pH, but would contact bacteria in high

concentrations. As expected the addition of synthetic 1-TbAd (pK_a ~ 8.5) and N^6-TbAd (pK_a ~ 3.8)

did not alter the pH of the 7H9 media (Sup. Fig. 1a). Next we inoculated *M. kansasii* into pH-

adjusted 7H9 broth (pH 4.0, 4.8, 5.0, 5.2, 5.4, 6.7,) containing 1, 5, 10 or 20 μM TbAd and

monitored growth over 16 days (Fig. 4 and Sup. Fig. 1b). With increasing doses of 1-TbAd or its

isomer N^6-TbAd, we did not observe any promotion of *M. kansasii* growth in normal nor acidic

7H9 broth (Fig. 4 and Sup. Fig 1b). In fact, 1-TbAd partially inhibited growth at 20 μM, the highest

dose tested, at normal pH. These data demonstrated that the protection from low pH in 7H9

culture media afforded by *Rv3377c-Rv3378c* complementation in *M. kansasii* is cell-intrinsic,

promotes growth only at low pH and does not occur with direct exposure to protonated TbAd.

1-TbAd does not maintain bulk cytosolic pH to enhance *M. kansasii* growth in low pH media

We aimed to identify whether 1-TbAd production alters/maintains the pH of the bacterial

cytosol when grown in acidic media. *M. kansasii*::EV or *M.kansasii::Rv3377-78c* were stained

with CFSE to measure their intracellular pH while monitoring their growth in different pH-

adjusted 7H9 broth (pH 4.0, 4.8, 5.0, 5.2, 5.4, 6.0, 6.7, 7.2) overnight (Fig. 5). Lower initial pH of

the media was associated with lowering intracellular pH over the experimental timeframe, as

expected. Importantly, we observed the expected growth advantage with 1-TbAd production at

lower pH (5.0), but there was no apparent intracellular pH difference between *M. kansasii* strains

at this or any pH.   Thus, 1-TbAd production does not aid *M. kansasii* growth at low pH by

maintaining the pH of the bulk intracellular milieu, suggesting the growth advantage is provided

by countering the effect of low pH in a specific region of the cell.

1-TbAd production enhances establishment of *M. kansasii* lung infection and survival in

alveolar macrophages.

1-TbAd is hypothesized to promote virulence of *M. tuberculosis* by countering phagosome

acidification, enhancing survival of the pathogen *in cellulo* (19, 27, 28). We therefore wished to

test the virulence of *M. kansasii*::*Rv3377-78c.* In C57Bl/6 mice, *M. tuberculosis* expands in a

logarithmic scale within the early course of infection. In sharp contrast, *M. kansasii* remains at its

initial levels of infection, suggesting it is a good model for acquisition-of-virulence studies within

the mouse (3). As a measure of virulence, we hypothesized that the pulmonary bacterial load of

M. kansasii in mice would be enhanced with 1-TbAd production. During pilot experiments, we

infected mice with *M. kansasii*::EV and *M. kansasii*::*Rv3377-78c* through aerosolization, and

infections with either strain resulted in a pulmonary burden within one log of the initial infection

up to day 42, with no clear differences between both groups (Sup Fig. 2).

Despite repeated attempts to standardize the inoculum, we consistently noted a higher

number of 1-TbAd-producing *M. kansasii::Rv3377-78c* one day after aerosol infection when

compared with *M. kansasii*::EV (not shown). To test if *Rv3377c-Rv3378c* was altering the dose

administered or instead enhancing establishment or early growth, mice were aerosolized with

M. kansasii::EV or *M. kansasii*::*Rv3377-78c* and their lungs were collected and homogenized

shortly (4 hours) after infection and compared to bacterial counts 24 hours after infection (Fig.

6a-b). Equivalent 4-hour CFU counts were observed for *M. kansasii*::EV and *M. kansasii*::*Rv3377-*

78c; only the latter multiplied successfully 24 hours later (Fig. 6a). *M. kansasii*::*Rv3377-78c*

showed a 1.5-fold increase in numbers from 4 to 24 hours while *M. kansasii*::EV numbers

remained equal (Fig. 6b). The experiment was performed three times, once at a low dose of 100

CFUs/lung and twice at a higher dose of 750 CFU/lung, with *M. kansasii*::*Rv3377-78c* consistently

attaining larger numbers by 24 hours post-infection in all three instances (Sup. Fig. 3).

We hypothesized that 1-TbAd was enhancing proliferation of *M. kansasii* in the lungs by

promoting survival in resident macrophages. First, using bone marrow-derived macrophages

(BMDMs), we infected with *M. kansasii*::EV or *M. kansasii*::*Rv3377-78c* and collected cell lysates

at 4- and 24-hours post infection: the infections proceeded similarly with both bacteria unlike

what we had observed in mouse lungs *in vivo* (Sup. Fig. 4). Alveolar macrophages (AMs) are

resident lung macrophages that phagocytose infectious agents entering the lower airways (29).

We assessed whether fitness would be altered in an *ex vivo* infection of murine AMs. In two

independent experiments, *M. kansasii*::*Rv3377-78c* increased in numbers by CFU counts from 4

to 24 hours post infection, while *M. kansasii*::EV numbers were largely unchanged and inferior

to *M. kansasii*::*Rv3377-78c* after 24 hours (fig. 6c-d). *M. kansasii*::*Rv3377-78c* exhibited a 2.0-fold

increase compared to the 1.1-fold increase seen in *M. kansasii*::EV (Fig. 6e). This is consistent

with our observations in murine lungs and with the conclusion that 1-TbAd provides an advantage

to *M. kansasii* during the early stages of pulmonary infection by specifically promoting survival

or growth in resident alveolar macrophages.

1-TbAd production impairs long-term persistence during mixed lung infections

We validated a translaryngeal infection model wherein WT *M. kansasii* was directly

introduced into the upper respiratory tract (Sup. Fig. 5 and methods) (30). Mice were monitored

for 42 days; bacteria persisted at the same log CFU as the initial infection and the mice did not

become overtly sick (Sup. Fig. 5). To test the effect of 1-TbAd in a high-dose competitive infection,

we used the translaryngeal infection model to generate a mixed infection with 3×10^6 CFUs of a

1:1 WT *M. kansasii* and *M. kansasii*::*Rv3377-78c* (Fig. 7a). Lung homogenates plated on 7H10

plates with and without 50 µg/ml hygromycin revealed a statistically significant decrease in the

M. kansasii::*Rv3377-78c* to WT *M. kansasii* ratio over time, with an initial decrease in the

proportion of *M. kansasii*::*Rv3377-78c* in the bacterial population from 0.48 (week 0) to 0.27

(week 1), stabilizing at that latter proportion over time (Fig. 7a). The competitive setting was

established to identify more subtle effects that the production of 1-TbAd could have on the

producing cells or the whole bacterial population over time. These findings show that such a

beneficial effect does not exist in the tested settings.

C-C chemokine receptor 2 (CCR2) is an essential component for defense in the airways;

Ccr2-/- mice lose the ability to recruit non-tissue resident immune cells and succumb to *M.*

tuberculosis infection (31). We used these mice to test whether the short-term AM phenotype

could be recapitulated in a longer-term *in vivo* setting without the interference of recruited

immune cells in WT mice that might explain the lack of a phenotype in the previous experiment.

We aerosol infected *Ccr2-/-* mice with a mixed 1:1 bacterial suspension of WT *M. kansasii* and *M.*

kansasii::*Rv3377-78c* (Fig. 7b). Interestingly, although WT *M. kansasii* exhibited a slight increase

in numbers over time, *M. kansasii*::*Rv3377-78c* steadily decreased within the same period. A

statistically significant decrease in the *M. kansasii*::*Rv3377-78c* to WT *M. kansasii* ratio over time

was noted, with a steady, non-stabilizing decrease in the proportion of *M. kansasii*::*Rv3377-78c*

in the bacterial population (Fig. 7b). Thus, *M. kansasii*::*Rv3377-78c* was clearly outcompeted by

WT *M. kansasii* in both of these *in vivo* infection models.

DISCUSSION

Our results indicate the feasibility of using *M. kansasii* to study the pathoevolution of *M.*

tuberculosis. The less virulent non-tuberculous mycobacterium (NTM) species is a suitable

surrogate for the expression of *M. tuberculosis*-specific products, such as 1-TbAd, and can readily

be used *in vitro*, and for *ex vivo* and *in vivo* experimental infection models. We showed that 1-

TbAd led to an improved survival during the first 24 hours of infection when tested *in vivo*, and

ex vivo in alveolar macrophages, but the isolated addition of 1-TbAd to *M. kansasii* resulted in

impaired persistence in different murine infection models.

In the current study, we demonstrated that *M. kansasii::Rv3377-78c* produced lipid

species distinct from those seen in *M. kansasii*::EV. Our prior and current data indicated that

transfer of TbAd biosynthesis genes to *M. kansasii* does not promote growth at neutral pH, but

confers increased growth in 7H9 media in the more acidic pH range (5.0-5.4) (22). When added

as a pure compound or produced by *M. tuberculosis*, 1-TbAd detectably raises the pH and swells

lysosomes in human macrophages (18, 22). In broad terms, the mechanisms by which the effects

of 1-TbAd are mediated could occur through direct chemical results of lysosomotropism, or

through signalling.

1-TbAd could act as an amphipathic weak base that penetrates membranes as an

uncharged conjugate base to selectively accumulate in acidic phagolysosomes where proton

capture raises pH and confers a positive charge, trapping the compound and leading to lysosomal

swelling. We did not see clear *Rv3377-Rv3378c*-dependent alkalization of 7H9 broth which might

be explained by compartmentalization: whereas intracellular bacteria are bound in a small

phagosomal compartment of 10^{-15} L, growth in 7H9 media provides a much larger compartment

for 1-TbAd to disperse in if it is physically shed from the bacterium. By a rough estimate, 10^{10}

bacteria-worth of 1-TbAd would be required to change the pH of 1 ml of 7H9 from 5.2 to 5.3,

about 10,000 times the concentration of bacteria we inoculate (see supplementary data

calculation). This result suggested that 1-TbAd production provides a pH-dependent growth

advantage intrinsic to the bacterium, separate from but not exclusive to lysosomal perturbation.

In phagocyte-free systems, we experimentally tested the hypothesis that the pH-

dependent growth advantage of *M. kansasii* producing 1-TbAd might involve direct contact of

the molecules with bacteria. Overall we found that direct exposure to externally added,

protonated 1-TbAd at low and high concentrations did not promote growth as might be expected

of a signaling molecule. Increased growth was selectively observed at low pH (5.1-5.4) when 1-

TbAd was generated through gene transfer and the action of enzymes in the bacterium. While

not fully understood, these divergent outcomes whereby the compartment of origin controls the

protective effect can be explained by the lysosomotropy model. Cytosolic 1-TbAd would be

expected to shed its proton during membrane passage into the periplasm. The lack of pH control

in the cytosol of *M. kansasii::Rv3377-78c* cultured in acidic media suggests that 1-TbAd does not

act as a shield against proton flow into *M. kansasii* cytosol, which is the expected outcome if

membrane penetration is required for the protective effect. These outcomes indicate that the

consistently observed survival advantage could derive from 1-TbAd passage from the cytosolic

membrane into the periplasm, mycolate membrane or surface of *M. kansasii*. 1-TbAd may act

on the bacterial population itself by targeting or protecting specific molecules during exposure

to low pH, stopping damage from occurring.

It is noteworthy that genetic and chemical complementation provide different

information about mycobacterial virulence factors, which in this case might result from the

differential compartmentalization of the molecules. This result also argues that 1-TbAd must

exert its protective effect at a specific location within the bacterial cell or cell wall. Exogenous 1-

TbAd may simply not reach this specific location, or not reach the location in an uncharged

conjugate base state. Together with our genetic complementation data, it is clear that *Rv3377c-*

Rv3378c-dependent metabolites including 1-TbAd do not have a direct growth-promoting effect.

Our data also demonstrate that *Rv3377c-Rv3378c*-dependent metabolites protect against acid

stress *in vitro*, using mechanisms that are independent of macrophage function, including

lysosomes or activating receptors.

Another key finding is that the complemented strain fared better than *M. kansasii*::EV in

the initial stages of *in vivo* infection. We subsequently showed that *M. kansasii*::*Rv3377-78c* was

more fit to thrive inside alveolar macrophages, but not bone marrow-derived macrophages

within that same timeframe, demonstrating that the production of 1-TbAd can subvert the first

lines of host defense encountered by the pathogen. Reasons for the observed increased survival

in alveolar macrophages might include 1-TbAd inducing acid tolerance that enables the bacteria

to survive the conditions inside the phagolysosome.

WT *M. kansasii* eventually outcompeted *M. kansasii*::*Rv3377-78c* in low- and high-dose

mixed infection settings in both WT and *Ccr2-/-* mice. Therefore, although expression of *Rv3377c-*

Rv33778c conferred a survival benefit to *M. kansasii* in specific *in vitro* and short-term infection

contexts, it appears there is a drawback to 1-TbAd expression in the non-adapted mycobacterium

for persisting in the murine host. We hypothesize that the burden of constitutive production of

1-TbAd, which sequesters adenosine molecules, may prevent energy storage in the form of ATP

and have a negative impact on long-term *in vivo* survival for *M. kansasii*. Another consideration

is the extent to which mycobacterial killing is dependent on acid-mediated mechanisms. The

intrinsic antacid properties of 1-TbAd, its tropism for acid compartments, its marked remodeling

of lysosomes and the pH-dependent basis of growth promotion in culture media all point at a

selective role in protection against acid-mediated killing. Therefore, the varied outcomes in the

models examined herein might depend on the extent to which they test acid-dependent killing.

One question that remains unanswered is whether there is a single, predominant mechanism of

action for which 1-TbAd production is mainly conserved in *M. tuberculosis*, or multiple important

functions.

Phenotypes observed after pathogen-specific genetic complementation into non-

pathogenic species provide different information than the more commonly observed loss-of-

function phenotypes observed after deleting genes from pathogens. The latter requires breaking

one link in a causal chain and might have rippling downstream effects, if it is not the final

component of a response cascade. Gain of function is a rare phenomenon that occurs only when

the components of a larger pathogen-specific system are fully recapitulated in the non-pathogen

and then tested under conditions in which this system is essential. In this regard, that

biosynthetic genes for 1-TbAd can promote early stage growth in alveolar macrophages was

unexpected, so these data now point to a new direction for mechanistic studies of these genes

in *M. tuberculosis*.

To date, the established virulence factors of *M. tuberculosis* are largely conserved among

the NTM, with the exception of 1-TbAd, Tuberculosis Necrotizing Toxin (32) and the MoaA1-4

operon (33). Our data indicate that 1-TbAd alone does not confer a long-term *in vivo* benefit

consistent with *in vitro* and *in vivo* phenotypic differences between *M. tuberculosis* and *M.*

kansasii. Therefore, events of acquisition of other *M. tuberculosis*-specific and loss of *M. kansasii*-

specific effectors are likely required to recreate an *M. tuberculosis*-like *M. kansasii* mutant strain,

or alternatively the difference is due to the compounding of multiple subtle effects that

complement one another. The possibility that mycobacterial virulence factors manifest their

phenotype in a cell-dependent fashion is consistent with the known transcellular lifestyle of *M.*

tuberculosis and suggests that different host-cell types should be used to detect undiscovered

virulence determinants.

Viewed in this light, using genetically complemented NTM is therefore useful to single out

the effects of specific elements that contributed to *M. tuberculosis* host adaptation without

producing a clearly hypervirulent NTM. The 1-TbAd family of molecules represents a newly

discovered pathogen-specific collection of compounds that has no clear chemical analog in other

bacterial systems, and their exact mechanisms of action remain elusive. We can conclude from

our study that *Rv3377c-Rv3378c* transfer acts in a eukaryotic cell-free system by a localized

chemical mechanism that involves pH, and that such changes can be determinative of outcomes

in alveolar macrophages at expectedly early time points post-infection. As such, these molecules

may aid in the establishment of infection within the lower respiratory tract.

MATERIALS AND METHODS

Bacterial strains and culture conditions

M. kansasii ATCC 12478 and *M. tuberculosis* H37Rv were grown in Middlebrook 7H9 broth (BD

Difco, MD, USA) as previously described (3). To test the ability of *M. kansasii*::*Rv3377-78c* to

produce yellow pigment, fully-formed colonies were additionally exposed to white light and

incubated at room temperature for 7 days. Where indicated and to ensure single-cell

suspensions, liquid bacterial cultures were de-clumped by slowly passaging through 5x 22G, 5x

25-G and 3x 26-G needles followed by low-speed centrifugation at 50 g for 5 minutes with

passage through a 5-μm filter. To generate *M. kansasii*::*Rv3377-78c*, a 2.4-kb PCR fragment

spanning *Rv3377c-Rv3378c* was generated using primers BamHI-*Rv3377c-Rv3378c*-F and HindIII-

Rv3377c-Rv3378c-R (Sup. Table 1) using high-fidelity Phusion DNA polymerase (New England

Biolabs). The fragment was subsequently digested with BamHI and HindIII (all restriction enzymes

from New England Biolabs) and cloned into the episomal plasmid pMV261 with the constitutive

mycobacterial *hsp60* promoter and a selective apramycin resistance marker. The *hsp60-Rv3377c-*

Rv3378c fragment was shuttled into the integrative vector pMV306 containing a hygromycin

resistance cassette using XbaI and HindIII. All ligations were done using T4 DNA ligase

(Fermentas). The resulting plasmid pMV306::*Hsp60-Rv3377c-Rv3378c* was verified by Sanger

sequencing (Genome Québec) to ensure the absence of frameshift or point mutations during the

cloning process. An unaltered version of pMV306 with a hygromycin resistance cassette was used

to create the empty vector control strain *M. kansasii*::EV. Following electroporation, *M.*

kansasii::EV and *M. kansasii*::*Rv3377-78c* were grown in the presence of 100 μg/mL hygromycin

(Wisent).

Detection of cell filtrate adenosine-linked lipids

M. kansasii::EV, *M. kansasii*::*Rv3377-78c* and *M. tuberculosis* were grown to mid-log phase and

subsequently incubated with 0.25 μCi/mL radiolabeled [8-^{14}C]adenosine (American Radiolabeled

Chemicals) for 14 days. Polar lipid fractions were extracted using $CHCl_3$:CH_3OH:0.3%NaCl

(v/v/v)(22).)(34). Extracted lipids were spotted on a TLC Silica Gel 60 (Millipore Sigma) with

CHCl$_3$:CH$_3$OH:H$_2$O 10:5:1 (v/v/v) used as the mobile phase solvent. The radiolabeled signature

was developed using Storm 840 PhosphorImager (GE Healthcare) to visualize adenosine-linked

lipids in each lane. [8-^{14}C]adenosine 1:100 was used as a no-lipid staining control. The plate was

stained with 5% phosphomolybdic acid reagent (PMA) (Sigma) and heated briefly using an

industrial blow-dryer to visualize the total amounts of lipids loaded in each lane.

HPLC-MS analysis of lipids from cells and supernatant

M. kansasii::Rv3377-78c, *M. kansasii::*Empty vector (EV) and *M. kansasii* parent strain were

grown in 30 ml of 7H9 media supplemented with albumin-dextrose-saline (5% Bovine Serum

Albumin Fraction V, 2% anhydrous dextrose and 0.87% sodium chloride) to late log-phase.

Bacterial cell pellet and supernatant were separated by centrifugation at 5000 rpm for 5 minutes.

The cell pellet was resuspended in 4 ml of PBS at pH 7.4 and distributed equally into four 2 ml

screw-cap tubes. The cells were pelleted by centrifugation and further resuspended in 1 ml PBS

at pH titrated to 7.4, 6.4, 5.5 and 4.5 with hydrochloric acid and incubated for 2 hours at 37°C. At

the end of 2 hours the cell pellet and the PBS supernatant were collected for lipid extraction.

Added 10 volumes (3 ml) of Chloroform/methanol (C/M) at the ratio of 1:2 and extracted for 1

hour at room temperature. A second extraction under similar conditions was performed with 3

ml of C/M at the ratio of 1:1. The extracted fractions were pooled and dried under a stream of

nitrogen gas. Lipids from the 1 ml PBS supernatant was extracted using acidified ethyl acetate by

adding 3 μl of 6N HCl and 1.4 ml of ethyl acetate and mixing for 30 minutes in an Orbitron shaker.

The mixture was centrifuged at 2000 rpm for 15 minutes to collect the upper organic phase and

dried on to glass under a stream of nitrogen gas at room temperature, and total lipids were

weighed using analytical balance. HPLC-MS separations were performed as described (17) using

equal amount of lipid samples from different experimental conditions as determined by weight

on a Mettler balance.

Congo Red uptake assay

Bacterial cultures were grown on Congo Red-containing 7H10 plates for 14 days, scraped into a

15-ml conical tube, washed with water until the supernatant became clear and incubated with 2

ml DMSO for 2 hours (24, 35). Congo Red was measured in the resultant supernatant at A488.

The values were normalized to the dry weight of the pellet to define the Congo Red binding index.

Extracellular pH measurement

For all pH experiments, liquid media was prepared as usual and the pH was equilibrated to 4.0,

4.9, 5.0, 5.1, 5.2, 5.4, 6.0, 6.7 and 7.2 using 2 M HCl or NaOH. OD_{600} was adjusted to 0.34 and 222

µl of mid-log phase de-clumped bacteria were added to 15 ml of freshly prepared, pH-adjusted

7H9 in 150-ml roller bottles (final OD_{600} of 0.005). Triplicate cultures were made per condition

(strain and pH) and incubated at 37°C, rolling in the dark. OD_{600} was measured every 2-3 days

using 2 x 200 µl of culture (technical duplicates) and a Tecan Infinite M200 Pro plate reader. At

days 8 and 17, 1 ml was removed from each culture for centrifugation and recovery of

supernatant, which was stored at 4°C until extracellular pH was read using micro pH combination

electrode (AgCl) (Sigma Aldrich), and Orion Star A111 meter (Thermo Scientific).

Synthetic TbAds and chemical complementation

466 Synthetic 1-TbAd and N^6-TbAd were produced as described previously (36). Bacterial cultures

467 were grown to mid-log phase and de-clumped as described above, then inoculated into 96-well

468 plates in 200 μl pH-adjusted 7H9 containing 1-TbAd, N^6-TbAd or vehicle (DMSO) control as

469 indicated. Plates were incubated at 37°C in the dark and OD_{600} was measured every one to three

470 days with a Tecan Infinite M200 Pro plate reader.

Intracellular pH measurement

Bacterial cultures were grown to mid-log phase and de-clumped as previously described. 5×10^8

CFU were pelleted, the supernatant completely removed, and the cells resuspended in 0.3 ml

PBS containing 100 μM Carboxyfluorescein diacetate succinimidyl ester (CFDA-SE) (CellTrace™

CFSE, ThermoFisher) for 20 minutes at 37°C, shaking at 150 RPM in the dark. Bacteria were next

diluted with 10 ml of 7H9 and incubated for 4 hours at 37°C, rolling in the dark. A portion of

bacteria was taken during this incubation for lysis in normal saline (0.9% NaCl) by beating with

silica beads (MP Biomedicals, FastPrep-24) to extract free protein-CFSE conjugate to generate a

fluorescence-to-pH standard curve. After 4 hours, bacteria were washed and resuspended in

normal saline to an OD_{600} of 0.4. 20 μl of bacteria in saline were sub-cultured into 180 μl of freshly

prepared, pH-adjusted 7H9 in 96-well plates (opaque-black for fluorescence (ThermoScientific

Nunclon Delta Surface), translucent-colourless for absorbance (Falcon). Lysed bacteria were

plated similarly for the pH standard curve. Immediately, plates were placed in plate readers

(Tecan Infinite M200 Pro) at 37°C, shaking and measuring fluorescence or OD_{600} every 30

minutes. 528-nm fluorescence was measured from 490-nm excitation (pH-sensitive), and 520-

nm fluorescence was measured from 450-nm excitation (pH-insensitive). To calculate pH, 7H9

background was subtracted for all data first (pH did not alter 7H9 fluorescence). Next, a standard

curve of 490-excitation/450-excitation in relation to 7H9 pH was created from the CFSE-

containing cell lysate. The 490/450 ratios calculated from the culture wells were applied to the

standard curve to determine intracellular pH.

Murine pulmonary infection

C57Bl/6 and *Ccr2-/-* mice (Jackson Laboratories) of approximately 6 to 20 weeks of age were used

for experiments. All protocols were approved by independent ethics oversight at the RI-MUHC

and followed the guidelines of the Canadian Council on Animal Care (CCAC). C57Bl/6 mice were

infected through aerosolization (ONARES, NJ, USA) of bacterial cultures at OD_{600} 0.4 for 15

minutes as previously described (3). Alternatively, C57Bl/6 mice were infected via trans-laryngeal

intubation using 50 µl of a high-dose mixed bacterial suspension containing both wild-type (WT)

M. kansasii and *M. kansasii*::*Rv3377-78c* at OD_{600} 0.2. C57Bl/6 *Ccr2-/-* (C-C motif chemokine

receptor 2 knockout) mice were infected through aerosolization of a low-dose mixed suspension

at OD_{600} 1.0. Mouse lungs were harvested at 4 or 24 hours (early time-points to measure short-

term bacterial establishment) and 14, 21, 28, 42 or 56 days post infection (later time-points to

measure long-term bacterial persistence) into 1ml 7H9 and homogenized using an Omni Tissue

Homogenizer TH (Omni International) at high speed for 45 seconds. Serial dilutions made in 7H9

liquid media from lung homogenates were plated on 7H10 plates containing PANTA ± 50 µg/ml

hygromycin. Colony-forming units (CFU) were counted 2 weeks post-plating to determine

bacterial burden.

Murine macrophage isolation

Bone marrow was isolated from C57Bl/6 murine tibiae and femora. Bone marrow-derived

macrophage (BMDMs) were differentiated with recombinant M-CSF (100 U/ml) (Peprotech) for

a period of 7 days as previously described (37), after which they were lifted using 4 ml CellStripper

Solution (Corning) and seeded into the appropriate tissue-culture plates. For alveolar

macrophage (AM) isolation, the respiratory tract including the trachea and lungs was isolated

and repeatedly perfused with cold sterile phosphate-buffered saline (PBS) to collect the cells

through bronchoalveolar lavage (BAL). AMs were enriched through adherence purification to

tissue culture-treated 96-well plates over 24 hours, at which point other cells were washed away.

All mammalian cells were cultured in RPMI 1640 media supplemented with non-essential amino

acids, 10 mM HEPES, 10% FBS ± Penicillin/Streptomycin (Wisent).

Macrophage infection

Macrophages were seeded into 96-well plates (100,000 cells/200 µl complete RPMI media

without antibiotics). Bacterial cultures were grown to OD_{600} 0.2-0.5, clumps were removed to

ensure single-cell suspensions and adjusted in complete RPMI media (without antibiotics) to an

OD_{600} 0.01. Macrophages were infected by replacing the media with fresh media containing

bacterial suspension. After 4 hours of infection, the wells were gently washed three times with

PBS to remove extracellular bacteria and fresh complete RPMI media (without antibiotics) was

added to each well. At indicated time points, the plates were spun down at 2,000 g for 5 minutes.

Each well was subjected to PBS containing 1% Triton X-100 for 10 minutes at room temperature

to induce macrophage lysis. Following serial dilution and plating, CFUs were counted on 7H10

plates 2 weeks post-plating to determine bacterial burden.

Statistical analysis

All calculations and statistical analyses were performed using Microsoft Excel or GraphPad Prism.

Calculations included (1) the ratio of individual 24-hr CFU values/mean 4-hr CFU for murine lung

and macrophage infection assays to determine bacterial proliferation [datapoint = (24-hr

CFU)/(mean 4-hr CFU)] and (2) the proportion of *MKAN::Rv33778c /* total (values paired from

individual mice) in competition assays to determine comparative fitness of WT *M. kansasii* vs. *M.*

kansasii::*Rv3377-78c* [datapoint = (CFU on 7H10-Hygromycin)/(CFU on 7H10)].

Acknowledgements

We would like to thank Dr. Maziar Divangahi and Laura Mendonça for transferring their

knowledge on alveolar macrophages and for providing *Ccr2-/-* mice. We would also like to thank

Dr. David Young for assistance with chromatography and figure preparation.

This project was funded through the support of an operating CIHR grant FND-148362.

References

1.   Peirs P, Lefevre P, Boarbi S, Wang XM, Denis O, Braibant M, Pethe K, Locht C, Huygen K,

Content J. 2005. Mycobacterium tuberculosis with disruption in genes encoding the

551  phosphate binding proteins PstS1 and PstS2 is deficient in phosphate uptake and

552  demonstrates reduced in vivo virulence. Infect Immun 73:1898-902.

553 2. Dunphy KY, Senaratne RH, Masuzawa M, Kendall LV, Riley LW. 2010. Attenuation of

554  Mycobacterium tuberculosis functionally disrupted in a fatty acyl-coenzyme A synthetase

555  gene fadD5. J Infect Dis 201:1232-9.

556 3. Wang J, McIntosh F, Radomski N, Dewar K, Simeone R, Enninga J, Brosch R, Rocha EP,

557  Veyrier FJ, Behr MA. 2015. Insights on the emergence of Mycobacterium tuberculosis

558  from the analysis of Mycobacterium kansasii. Genome Biol Evol 7:856-70.

559 4. Bloch KC, Zwerling L, Pletcher MJ, Hahn JA, Gerberding JL, Ostroff SM, Vugia DJ, Reingold

560  AL. 1998. Incidence and clinical implications of isolation of Mycobacterium kansasii:

561  results of a 5-year, population-based study. Ann Intern Med 129:698-704.

562 5. Johnson MM, Odell JA. 2014. Nontuberculous mycobacterial pulmonary infections. J

563  Thorac Dis 6:210-20.

564 6. Sheu LC, Tran TM, Jarlsberg LG, Marras TK, Daley CL, Nahid P. 2015. Non-tuberculous

565  mycobacterial infections at San Francisco General Hospital. Clin Respir J 9:436-42.

566 7. Ricketts WM, O'Shaughnessy TC, van Ingen J. 2014. Human-to-human transmission of

567  Mycobacterium kansasii or victims of a shared source? Eur Respir J 44:1085-7.

568 8. Griffith DE, Aksamit T, Brown-Elliott BA, Catanzaro A, Daley C, Gordin F, Holland SM,

569  Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S, von Reyn CF, Wallace

570  RJ, Jr., Winthrop K, Subcommittee ATSMD, American Thoracic S, Infectious Disease

571  Society of A. 2007. An official ATS/IDSA statement: diagnosis, treatment, and prevention

572  of nontuberculous mycobacterial diseases. Am J Respir Crit Care Med 175:367-416.

9.   Fortune SM, Jaeger A, Sarracino DA, Chase MR, Sassetti CM, Sherman DR, Bloom BR,

Rubin EJ. 2005. Mutually dependent secretion of proteins required for mycobacterial

virulence. Proc Natl Acad Sci U S A 102:10676-81.

10.   Boon C, Dick T. 2002. Mycobacterium bovis BCG response regulator essential for hypoxic

dormancy. J Bacteriol 184:6760-7.

11.   Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I. 2006. The Mycobacterium

tuberculosis PhoPR two-component system regulates genes essential for virulence and

complex lipid biosynthesis. Mol Microbiol 60:312-30.

12.   Salah IB, Ghigo E, Drancourt M. 2009. Free-living amoebae, a training field for

macrophage resistance of mycobacteria. Clin Microbiol Infect 15:894-905.

13.   Veyrier F, Pletzer D, Turenne C, Behr MA. 2009. Phylogenetic detection of horizontal gene

transfer during the step-wise genesis of Mycobacterium tuberculosis. BMC Evol Biol

9:196.

14.   Becq J, Gutierrez MC, Rosas-Magallanes V, Rauzier J, Gicquel B, Neyrolles O, Deschavanne

P. 2007. Contribution of horizontally acquired genomic islands to the evolution of the

tubercle bacilli. Mol Biol Evol 24:1861-71.

15.   Boritsch EC, Khanna V, Pawlik A, HonorÈ N, Navas VH, Ma L, Bouchier C, Seemann T,

Supply P, Stinear TP, Brosch R. 2016. Key experimental evidence of chromosomal DNA

transfer among selected tuberculosis-causing mycobacteria. Proceedings of the National

Academy of Sciences of the United States of America 113:9876-81.

16.   Sapriel G, Brosch R. 2019. Shared pathogenomic patterns characterize a new phylotype,

594   revealing transition towards host-adaptation long before speciation of Mycobacterium

595   tuberculosis. Genome Biol Evol doi:10.1093/gbe/evz162.

596   17.   Layre E, Lee HJ, Young DC, Martinot AJ, Buter J, Minnaard AJ, Annand JW, Fortune SM,

597   Snider BB, Matsunaga I, Rubin EJ, Alber T, Moody DB. 2014. Molecular profiling of

598   Mycobacterium tuberculosis identifies tuberculosinyl nucleoside products of the

599   virulence-associated enzyme Rv3378c. Proc Natl Acad Sci U S A 111:2978-83.

600   18.   Young DC, Layre E, Pan SJ, Tapley A, Adamson J, Seshadri C, Wu Z, Buter J, Minnaard AJ,

601   Coscolla M, Gagneux S, Copin R, Ernst JD, Bishai WR, Snider BB, Moody DB. 2015. In vivo

602   biosynthesis of terpene nucleosides provides unique chemical markers of Mycobacterium

603   tuberculosis infection. Chem Biol 22:516-526.

604   19.   Mann FM, Xu M, Chen X, Fulton DB, Russell DG, Peters RJ. 2009. Edaxadiene: a new

605   bioactive diterpene from Mycobacterium tuberculosis. J Am Chem Soc 131:17526-7.

606   20.   Lau SK, Lam CW, Curreem SO, Lee KC, Lau CC, Chow WN, Ngan AH, To KK, Chan JF, Hung

607   IF, Yam WC, Yuen KY, Woo PC. 2015. Identification of specific metabolites in culture

608   supernatant of Mycobacterium tuberculosis using metabolomics: exploration of potential

609   biomarkers. Emerg Microbes Infect 4:e6.

610   21.   David HL. 1974. Carotenoid pigments of Mycobacterium kansasii. Appl Microbiol 28:696-

611   9.

612   22.   Buter J, Cheng TY, Ghanem M, Grootemaat AE, Raman S, Feng X, Plantijn AR, Ennis T,

613   Wang J, Cotton RN, Layre E, Ramnarine AK, Mayfield JA, Young DC, Jezek Martinot A,

614   Siddiqi N, Wakabayashi S, Botella H, Calderon R, Murray M, Ehrt S, Snider BB, Reed MB,

Oldfield E, Tan S, Rubin EJ, Behr MA, van der Wel NN, Minnaard AJ, Moody DB. 2019.

Mycobacterium tuberculosis releases an antacid that remodels phagosomes. Nat Chem

Biol 15:889-899.

23. de Duve C, Trouet A, Campeneere DD, Baurian R. 1978. Liposomes as lysosomotropic

carriers. Ann N Y Acad Sci 308:226-34.

24. Jankute M, Nataraj V, Lee OY, Wu HHT, Ridell M, Garton NJ, Barer MR, Minnikin DE, Bhatt

A, Besra GS. 2017. The role of hydrophobicity in tuberculosis evolution and pathogenicity.

Sci Rep 7:1315.

25. Portaels F, Pattyn SR. 1982. Growth of mycobacteria in relation to the pH of the medium.

Ann Microbiol (Paris) 133:213-21.

26. Vandal OH, Nathan CF, Ehrt S. 2009. Acid resistance in Mycobacterium tuberculosis. J

Bacteriol 191:4714-21.

27. Pethe K, Swenson DL, Alonso S, Anderson J, Wang C, Russell DG. 2004. Isolation of

Mycobacterium tuberculosis mutants defective in the arrest of phagosome maturation.

Proc Natl Acad Sci U S A 101:13642-7.

28. Stewart GR, Patel J, Robertson BD, Rae A, Young DB. 2005. Mycobacterial mutants with

defective control of phagosomal acidification. PLoS Pathog 1:269-78.

29. Johnson JD, Hand WL, King NL, Hughes CG. 1975. Activation of alveolar macrophages after

lower respiratory tract infection. J Immunol 115:80-4.

30. Guilbault C, Martin P, Houle D, Boghdady ML, Guiot MC, Marion D, Radzioch D. 2005.

Cystic fibrosis lung disease following infection with Pseudomonas aeruginosa in Cftr

636  knockout mice using novel non-invasive direct pulmonary infection technique. Laboratory

637  Animals 39:336-352.

638 31. Peters W, Scott HM, Chambers HF, Flynn JL, Charo IF, Ernst JD. 2001. Chemokine receptor

639  2 serves an early and essential role in resistance to Mycobacterium tuberculosis. Proc Natl

640  Acad Sci U S A 98:7958-63.

641 32. Pajuelo D, Gonzalez-Juarbe N, Tak U, Sun J, Orihuela CJ, Niederweis M. 2018. NAD(+)

642  Depletion Triggers Macrophage Necroptosis, a Cell Death Pathway Exploited by

643  Mycobacterium tuberculosis. Cell Rep 24:429-440.

644 33. Levillain F, Poquet Y, Mallet L, Mazeres S, Marceau M, Brosch R, Bange FC, Supply P,

645  Magalon A, Neyrolles O. 2017. Horizontal acquisition of a hypoxia-responsive

646  molybdenum cofactor biosynthesis pathway contributed to Mycobacterium tuberculosis

647  pathoadaptation. PLoS Pathog 13:e1006752.

648 34. Slayden RA, Barry CE, 3rd. 2001. Analysis of the Lipids of Mycobacterium tuberculosis.

649  Methods Mol Med 54:229-45.

650 35. Cangelosi GA, Palermo CO, Laurent JP, Hamlin AM, Brabant WH. 1999. Colony

651  morphotypes on Congo red agar segregate along species and drug susceptibility lines in

652  the Mycobacterium avium-intracellulare complex. Microbiology 145 (Pt 6):1317-24.

653 36. Buter J, Heijnen D, Wan IC, Bickelhaupt FM, Young DC, Otten E, Moody DB, Minnaard AJ.

654  2016. Stereoselective Synthesis of 1-Tuberculosinyl Adenosine; a Virulence Factor of

655  Mycobacterium tuberculosis. J Org Chem 81:6686-96.

656 37. Zhang X, Goncalves R, Mosser DM. 2008. The isolation and characterization of murine

657  macrophages. Curr Protoc Immunol Chapter 14:Unit 14 1.

Figure 1 – *M. kansasii::*Rv3377-78c produces adenosine-linked lipids 1-TbAd and N6-TbAd. (a) Detection of adenosine-linked lipids extracted from *M. kansasii::*EV (MKAN::EV), *M. kansasii::Rv3377-78c* (MKAN::Rv33778c) and *M. tuberculosis* (H37Rv) through radiolabelling and separation using normal-phase silica thin-layer chromatography. (b) Visualization of migration pattern of total lipids from each sample after staining with 5% phosphomolybdic acid reagent. (c-e) Lipids from *M. kansasii* derived from cell pellets or culture supernatant incubated for two hours at the indicated pH, neutralized and then extracted with organic solvent. Product was analyzed in comparison with a synthetic standard for 1-TbAd. (e) The mass spectrum of lipids extracted from 21-22 min for *M. kansasii* show a m/z value that matches with the measured and expected mass of a 1-TbAd standard (21) (f-g) Total extracted lipids expressed as area under curve from counts versus retention time of the extracted ion chromatogram. 1.0uM of synthetic 1-TbAd was used as the standard.

Figure 2 – 1-TbAd production does not influence the *in vitro* growth characteristics and behaviours of *M. kansasii*. (a) Comparative OD$_{600}$ growth kinetics of wild-type *M. kansasii* (MKAN), *M. kansasii*::EV *(*MKAN::EV) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c) at 37°C in 7H9 broth. The data are presented as the mean of technical triplicates ± SD. The data are representative of three independent experiments. (b) Colony morphology in 2 different incubation settings of *M. kansasii*::EV (MKAN::EV) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c) on 7H10 plates. (c) Colony morphology of *M. kansasii*::EV (MKAN::EV) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c) on 7H10 plates supplemented with Congo Red. (d) Quantitative analysis of Congo Red dye retention by *M. kansasii*::EV (MKAN::EV) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c). DMSO extraction was followed by absorbance at 488nm divided by the weight of the dry culture pellet after washing (in grams). The data are plotted as the mean of technical triplicates ± SD. The data are representative of two independent experiments.

Figure 3 – 1-TbAd production enhances growth at low pH where growth is associated with culture medium alkalization.

*M. kansasii::*EV (MKAN::EV) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c) cultures were inoculated at equal OD_{600} into fresh pH-adjusted 7H9 (using HCl titration) and incubated at 37°C in a rolling incubator over 17 days. The starting pH of the cultures is indicated above each graph. (a) OD_{600} was monitored every 2-3 days. The data are plotted separately for each of three independently growing cultures per strain. The data are representative of 5 independent experiments. (b) The pH of the supernatant was measured at days 8 and 17. The data are presented as the mean of technical triplicates ± SD. The data are representative of three independent experiments.

Figure 4 – Chemical complementation of *M. kansasii* with synthetic TbAd does not promote growth at any pH. (a) *M. kansasii* WT was inoculated into fresh pH-adjusted 7H9 (using HCl titration) containing the indicated concentration of TbAd isomer or DMSO, and then incubated at 37°C in 96-well plates over 16 days. OD_{600} was measured every 1-3 days. The data are presented as the mean of technical quadruplicates ± SD.

Figure 5 – 1-TbAd does not alter the intracellular pH of bacteria - *M. kansasii*::EV (MKAN::EV - solid lines) and *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c - dashed lines) cultures at equal OD_{600} were stained with CFSE, inoculated into fresh pH-adjusted 7H9 and incubated shaking at 37°C in 96-well plates placed in the dark. Growth (grey / OD_{600} measurements) and intracellular pH (black / pH calculated from fluorescence excitation-emission ratios) readings were taken at 30-minute intervals overnight. The data are presented as the median of technical triplicates. Data are representative of 5 independent experiments.

Figure 6 – 1-TbAd enhances the initial establishment of pulmonary infection. (a-b) CFUs were counted from C57Bl/6 mouse lungs isolated at 4- vs. 24-hours post aerosol infection with *M. kansasii*::EV (MKAN::EV) or *M. kansasii*::*Rv3377-78c* (MKAN::Rv33778c) (n=20 lungs/condition/time point). (a) Absolute CFU count data are pooled from two independent experiments with similar initial inoculum. (b) 24-hr CFU/mean 4-hr CFU ratio data were pooled from three independent experiments (n=21-25 lungs/condition/time point). (c-e) CFUs were counted from C57Bl/6 murine-derived AMs at 4- and 24-hours after *ex vivo M. kansasii* infection. (c-d) Absolute CFU count data from two independent experiments (N=5 and 7 replicate wells containing infected AMs, respectively, per condition per timepoint). (e) 24-hr CFU/mean 4-hr CFU ratio data pooled from two independent experiments (N=12 per condition). The data are plotted as the mean ± SD. GraphPad Prism 8.1.2 was used to perform Welch's two-tailed unpaired t-tests where *p<0.05, **p<0.01, ****p<0.0001.

Figure 7 – 1-TbAd production hinders long-term bacterial growth. (a-b) Mixed 1:1 bacterial suspensions of WT *M. kansasii* (MKAN) and *M. kansasii::Rv3377-78c (MKAN::Rv33778c)* were used to infect (a) WT C57Bl/6 mouse lungs isolated at 1, 7, 28 and 56 days post high-dose trans-laryngeal intubation (n=8-10 lung pairs per timepoint) or (b) *Ccr2 -/-* C57Bl/6 mouse lungs isolated at 0, 14 and 29 days post aerosolization (n=5-9 lung pairs per timepoint). CFUs were counted on 7H10 plates + PANTA ± hyg50. The graphs on the right represent the proportion of *M. kansasii::Rv3377-78c* over the total number of bacteria (MKAN + MKAN::Rv33778c) per mouse per timepoint. The raw data (left) and proportions (right) are plotted as individual datapoints (± SD for proportions only). GraphPad Prism 8.1.2 was used to perform the ratio paired t-test (ratio per timepoint/left) and ordinary one-way ANOVA (proportions over time/right).