The role of the meningeal lymphatic system in local inflammation and trigeminal nociception implicated in migraine pain

Nikita Mikhailov[1], Kseniia Koroleva[1,3], Ali Abdollahzadeh[1], Raisa Giniatullina[1], Oleg
Gafurov[3], Tarja Malm[1], Alejandra Sierra[1], Jussi Tohka[1], Francesco Noe[2#], Rashid
Giniatullin[1,3#]

[1]A. I. Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio,
70211, Finland

[2]HiLIFE-Neuroscience Center, University of Helsinki, Helsinki, 00290, Finland

[3]Laboratory of Neurobiology, Kazan Federal University, Kazan 420008, Russia

[#] **Shared last authorship**
*Correspondence:

Francesco Noe, Neuroscience Center, HiLife, Helsinki University, Biomedicum A-wing,
Haartmaninkatu 8, P.O.Box 69, Helsinki 00290, Finland, e-mail: *francesco.noe@helsinki.fi*

Keywords: meningeal lymphatic, neuroinflammation, migraine, pain

Abstract

Background: A system of lymphatic vessels has been recently characterized in the
meninges, with a postulated role in 'cleaning' the brain via cerebral fluid drainage. As
meninges are the origin site of migraine pain, we hypothesized that malfunctioning of the
lymphatic system should affect the local trigeminal nociception. To test this hypothesis, we
studied nociceptive and inflammatory mechanisms in the meninges of K14-VEGFR3-Ig mice
lacking the meningeal lymphatic system.

Methods: We recorded the spiking activity of meningeal afferents and estimated the local
mast cells infiltration, calcitonin gene-related peptide (CGRP) and cytokine levels (basal and
stimulated), as well as the dural trigeminal innervation in freshly-isolated hemiskull
preparations from K14-VEGFR3-Ig (K14) or wild type C57BL/6 mice (WT).

Results: We found that the meningeal level of CGRP and of the pro-inflammatory cytokines
IL12-p70 and TNFα (implicated in migraine) were reduced in the meninges of K14 mice. On
the contrary, in the meninges of K14 mice, we found an increased level of the mast cell
activator MCP-1 and, consistently, a larger number of dural mast cells. The other migraine-
related pro-inflammatory cytokines did not differ between the two genotypes. The patterns of

trigeminal innervation in meninges remained unchanged and we did not observe alterations in
basal or ATP-induced nociceptive firing in the meningeal afferents.

Conclusions: In summary, the lack of meningeal lymphatic system does not induce
migraine-like nociceptive state *per se*, but leads to a new balance between pro- and anti-
inflammatory factors implicated in migraine mechanisms.

Abbreviations:

WT – wild type mice

K14 – K14-VEGFR3-Ig mice

mLVs – meningeal lymphatic vessels

1   Introduction

Migraine is a complex neurological disorder, which originates from the meninges (Bolay et al.,
2002; Olesen et al., 2009). Meninges are densely innervated by somatic trigeminal
(Ramachandran, 2018; Zakharov et al., 2015) and parasympathetic innervation (Delépine and
Aubineau, 1997), which runs along and in contact with meningeal blood vessels (Ebersberger
et al., 2006). It has been proposed that so-called sterile neurogenic inflammation (Moskowitz,
1993, 1984; Moskowitz et al., 1979) in the meninges is an important initiator of the
pathophysiology of migraine. The initial step in neurogenic inflammation is the activation of
the trigeminal nociceptive system in meninges (Moskowitz, 1993). Indeed, trigeminal neurons
release the calcitonin gene-related neuro-peptide (CGRP), which activates dural mast cells
(Ottosson and Edvinsson, 1997), promotes sensitization of trigeminal afferents (Cady et al.,
2011; Giniatullin et al., 2008), and induces local vasodilation (Goadsby et al., 1988).
Vasodilation, in turn, activates mechanosensitive receptors, expressed in trigeminal nerves
within the meninges (Mikhailov et al., 2019). Trigeminal neurons can also be activated by mast
cells-secreted serotonin (Kilinc et al., 2017), and by extracellular ATP, which is released from
endothelial and mast cells, as well as from neurons, upon stimulation of the migraine mediator
CGRP (Koroleva et al., 2019; Suleimanova et al., 2020; Yegutkin et al., 2016). Finally, the
release of acetylcholine (ACh) by parasympathetic fibers results in meningeal mast cell
degranulation (Shelukhina et al., 2017), which activates the trigeminal nociceptive system. In
addition, endothelial (Loberg et al., 2006) and mast cells (Mukai et al., 2018) release various
pro-inflammatory cytokines as potential contributors to local inflammation.

The presence of these diverse endogenous agents in the meninges suggests the existence of
regulatory mechanisms, which modulates the local levels of pro-inflammatory mediators and
minimize the activity of nociceptive fibers. The recently characterized meningeal lymphatic

system (Antila et al., 2017; Aspelund et al., 2015; Louveau et al., 2015) **(Figure 1)** has been
suggested to play a role in the clearance of the central nervous system (CNS) from toxic
molecules, and to be involved in the pathogenesis of different neurodegenerative diseases
(Kwon et al., 2019; Noé and Marchi, 2019; Patel et al., 2019; Rasmussen et al., 2018; Tamura
et al., 2019). We hypothesized that the meningeal lymphatic vessels (mLVs) can play a similar
role also in the meninges, clearing the pro-inflammatory mediators. We also hypothesize that
dysfunction in the lymphatic drainage can trigger migraine.

2   Methods and materials

2.1   Animals

We used adult (30 – 40 g) male and female transgenic K14-VEGFR3-Ig (K14) mice lacking
the meningeal lymphatic system (Aspelund et al., 2015; Mäkinen et al., 2001) and wild-type
C57BL/6J littermates mice (WT). Total number of animal used for this study was 38 WT and
37 TG animals. Animals were housed at the Lab Animal Center of the University of Eastern
Finland under following housing conditions: 12 hours dark/light cycle, grouped housing, given
ad libitum access to food and water, 22°C ambient temperature. All procedures with animals
were conducted in accordance with the European Community Council Directive 2010/63/EU.
Experimental protocols involving the usage of animals were approved by the Animal Care and
Use Committee of the University of Eastern Finland (license EKS-004-2014) and by the
National Animal Experiment Board (ESAVI-2017-008787).

2.2   Hemiskull preparation for ex vivo experiments

For the *in vitro* and *ex vivo* experiments (mast cell staining, CGRP release assay, cytokine
release assay, and electrophysiology), we used hemiskull preparations with preserved
meninges and meningeal trigeminal innervation (Shatillo et al., 2013; Zakharov et al., 2015).
For electrophysiology and CGRP release assay, animals we sacrificed with CO_2 without
anaesthesia. For cytokine release assay and mast cell staining, animals were anesthetized with
intraperitoneal injection of 250 mg/kg tribromoethanol (Avertin, Sigma-Aldrich, USA) and
then underwent transcardial perfusion: for mast cell staining, we perfused with normal saline
(6 minutes at 6 ml/min), followed by 4% paraformaldehyde (PFA) (20 minutes at 3 ml/min);
for cytokine release assay, animals were just perfused with normal saline (4 min at 20 ml/min).
In order to prepare the hemiskulls for subsequent analyses, after decapitation, skin, muscles,
and connective tissues, as well as brain were removed with particular care to not touch and
damage the meninges. Thereafter, hemiskulls were isolated by cutting the skull along the
sagittal axis (*i.e.*, two hemiskulls were prepared from each animal).

2.3   Mast cell staining

Hemiskulls of WT and K14 animals were prepared as described above, washed three times for
20 minutes in PBS (phosphate buffered saline), and stained for 60 minutes with 0.1% toluidine
blue. Stained meninges were immediately imaged, using a Zeiss Axio Zoom.V16
stereomicroscope (Carl Zeiss AG, Germany), at 30x magnification. We imaged two identical
regions-of-interest (ROIs) from each hemiskull: one including the superior sagittal sinus (SSS;
at the border of the image) and the adjoining one, partially overlapping (approximately 10% of
area) with the previous ROI. Size of each ROI was 3.7x2.4 mm^2. In total, four images were
acquired from each mouse. From each ROI, the number of stained mast cells was calculated
using ImageJ 1.51v software (National Institute of Health, USA).

2.4 CGRP release assay

The CGRP release assay was conducted on hemiskulls of WT and K14 animals as previously
described (Mikhailov et al., 2019). We used a CGRP enzyme immunoassay kit (EIA kit,
SPIbio, France) to measure CGRP release from hemiskull preparations. The levels of CGRP
were assessed under control and KCl-treated conditions.
First, fresh hemiskull preparations from WT and K14 animals were perfused for 40 min with
artificial cerebrospinal fluid (aCSF in mM: 119 NaCl, 30 NaHCO$_3$, 11 glucose, 2.5 KCl, 1
MgCl$_2$, 1 NaH$_2$PO$_4$, 2 CaCl$_2$, adjusted pH level = 7.4) at room temperature (RT). Then,
hemiskulls were washed four times with 150 µl aCSF for 15 minutes at 37°C to stabilize the
basal condition. Samples of aCSF (100 µl each) from the last two washes were collected to
assess the basal level of CGRP (control conditions). After the last wash, hemiskulls were
bathed with an aCSF solution containing 30 mM KCl for 15 minutes. Thereafter, a 100 µl
sample was collected to assess CGRP under KCl-treated condition.
Samples were freezed with liquid nitrogen in tubes containing EIA buffer with peptidase
inhibitors. Assay protocol was carried out in accordance with instructions of manufacturer. In
brief, 100 µl of CGRP standard (for calibration curve) or sample were mixed with 100 µl of
anti- CGRP AChE tracer in a well plate, preliminary washed 5 times with washing buffer. After
incubating the plates at 4 °C for 16–20 h, supernatant was removed, and plates were washed 6
times with washing buffer. Finally, 200 µl of Ellman's reagent was added. Optical density was
measured at 405 nm using microplate reader (Wallac VICTOR2™, PerkinElmer, USA).

2.5 Cytokine release assay

For cytokine release assay, we used hemiskull preparations from both WT and K14 mice. One
hemiskull/animal was used to assess the release of cytokine under the control condition, while
the contralateral was used to measure the release of cytokine under treatment condition (100
µM benzoyl ATP, BzATP). After the collection of hemiskulls, all procedures were performed
at 37°C. Hemiskulls were bathed in 150 µl of either aCSF or aCSF + BzATP for three and a
half hours. aCSF samples (50 µl) were taken to assess the level of cytokines in control (treated
with aCSF only) or stimulated (aCSF + BzATP) conditions. The more stable ATP-analogue

BzATP was used for the cytokine release assay, due to the long treatment time (3½ h) and
because of fast degradation of ATP (Yegutkin et al., 2016). The levels of aCSF in the
hemiskulls were monitored throughout the incubation period. If needed, we refilled the
hemiskull preparation with the respective treatment solution to compensate for evaporation.
To assess the level of cytokines (IL-6, IL-10, MCP-1, TNFα, IFNγ, IL-12p70), we used the
CBA Mouse Inflammation Kit (Cytometric Bead Array, BD Biosciences, New Jersey, USA).
All procedures (sample and standard beads preparation) were performed following instructions
provided by the manufacturer. Data were acquired on the CytoFLEX S cytometer (Beckman
Coulter Inc., USA) and analyzed with the Flow Cytometric Analysis Program (FCAP) Array
software (BD Biosciences, USA).
Due to the low concentration of IL-12p70 and IFNγ, in some samples it was not possible to
quantify the exact concentration with the abovementioned method. Thus, for these cytokines
final n was lower than the total number of animals used for the assay.

2.6   Electrophysiology

Direct recording of action potentials from the trigeminal nerve afferents innervating meninges
was conducted as described earlier (Shatillo et al., 2013; Zakharov et al., 2015). Briefly,
hemiskulls of WT and K14 animals were placed in a perfusion chamber with a constant flow
(6 ml/min) of oxygenated aCSF. In order to access the nervus spinosus of the trigeminal nerve,
innervating meninges, we made a small incision in the dura mater. Next, the nervus spinosus
was sucked into a glass recording electrode filled with aCSF. We recorded 10 min of
spontaneous activity to get a baseline and to assess the difference in basal activity between WT
and K14 animals. Then, we applied 100 µM ATP for 10 min to stimulate nociceptive firing in
the trigeminal nerve (Zakharov et al., 2016), followed by 10 min of washout. The recording
was conducted with a digital amplifier (ISO 80, WPI Inc., USA) with bandpass 300 Hz – 3
kHz, gain 10000. Signals were digitized at 125 kHz using a NIPCI 6221 data acquisition board
(National Instruments, USA) and visualized with WinEDR software (Strathclyde University,
UK).

2.7   Meningeal nerve staining for β-tubulin III

For the meningeal nerve staining, meninges from PFA-fixed hemiskulls were isolated as
previously described (Louveau et al., 2015), and stained. Briefly, meningeal whole-mount
preparations were incubated with 1x phosphate-buffered saline (PBS) containing 2% normal
goat serum (NGS), 1% bovine serum albumin (BSA), 0.1% Triton X-100, and 0.05% Tween
20 for 1 hour at RT. After that, meninges were incubated with rabbit anti-β-tubulin III (1:1000,
Cat# T2200, Sigma-Aldrich, USA) overnight at 4°C in PBS containing 1% BSA and 0.5%
Triton X-100. Whole-mounts were washed in PBS at RT (3x) followed by incubation with
Alexa Fluor 488-conjugated goat anti-rabbit IgG antibody (1:500, Cat# A11008, Invitrogen,
USA) for 1 hour at RT in PBS with 1% BSA and 0.5% Triton X-100. After further washing in

PBS (3x) and in phosphate buffer (PB; 2x), meninges were mounted with VECTASHIELD
mounting medium (Vector Laboratories, USA), including DAPI, and coverslipped.
Images were acquired using a Leica TCS SP8 X confocal system (Leica Microsystems,
Germany) using the LAS X software. Images of the region of interest, adjacent to the middle
meningeal artery, were acquired with a 20x objective with 0.75 numerical aperture (NA), with
an in-plane pixel size of $1.14 \times 1.14 \ \mu m^2$ and a z-step of 1 µm. The full images were created by
merging 4×4 tile scans, covering a total area of $2034 \times 2034 \ \mu m^2$.

2.8   Axon segmentation and innervation analysis

We developed a semi-automated segmentation technique to annotate axons and analyze the
innervation in acquired 3D confocal microscopy images. An experienced researcher (AA)
inserts two points along the desired axon on the 2D maximum projection of a 3D confocal
image to define the axonal centerline. Given the two points, a minimal path algorithm
(Benmansour and Cohen, 2011) detects the centerline of that axon. We used active contours
(Chan and Vese, 2001) to segment the desired axon, initiating from the detected axonal
centerline. For each axon, we measured the axonal diameter and the length of the centerlines
in 3D (Abdollahzadeh et al., 2019a). We also quantified an innervation complexity value for
each meningeal tissue by forming a graph from the segmented axons (Abdollahzadeh et al.,
2019b) and reported the number of axonal endpoints, as shown in Fig.7a. We described the
details of axon segmentation and the morphology analysis in Supplementary Materials ("axon
segmentation and innervation analysis" section).

2.9   Statistical analysis

We performed statistical analysis using the GraphPad Prism 8.4.2 software (GraphPad
Software, California, USA). The data are presented as mean ± the standard error of the mean
(for two-way ANOVA) or as median with interquartile range (for Mann-Whitney U test and
linear mixed effect model). We set the alpha-threshold defining the statistical significance as
0.05 for all analyses. Mast cell abundance as well as axon segmentation and innervation
analyses data were analysed using Mann-Whitney U test. CGRP release assay data was
analysed with repeated measures two-way ANOVA. Cytokine release assay data were analysed
by Mann-Whitney U test or by matched two-way ANOVA. A linear mixed effect model was
used to evaluate the differences in the electrophysiology data. Bonferroni correction was used
to adjust p-values in multiple comparison. Please see figure legends for the details of the used
test in each analysis.

3   Results

3.1   Elevated number of mast cells in meninges of mice without lymphatic system

Meningeal mast cells recently emerged as important players in the initiation of a migraine
attack (Giniatullin et al., 2019; Kilinc et al., 2017; Koyuncu Irmak et al., 2019; Levy, 2009).
As the mLVs are important for the movement of immune cells (Ahn et al., 2019; Louveau et
al., 2015), we expected that the lack of mLVs could affect the local accumulation of mast cells.
We used the toluidine blue labelling to characterize the profile of meningeal mast cells.

In WT animals a substantial portion of mast cells was located along the meningeal blood
vessels (**Figure 2A**), where the mLVs are located (Antila et al., 2017; Aspelund et al., 2015).
A similar pattern of mast cell positioning was observed in K14 animals (**Figure 2B**). K14
animals showed a non-significant trend towards increased number of meningeal mast cells
(**Figure 2.C**). Since an increase in the number of mast cells may lead to a more abundant release
of local mediators (Giniatullin et al., 2019; Levy, 2009; Suleimanova et al., 2020), resulting in
meningeal inflammation and in the stimulation of the trigeminal afferents (Kilinc et al., 2017;
Koroleva et al., 2019), we analyzed the levels of the different migraine-associated pro-
inflammatory molecules and the activation of the trigeminal nerves in the K14 and their WT
littermates mice.

3.2 Altered cytokine release in meninges

We first measured the level of cytokines (IL-6, IL-10, IL-12p70, MCP-1, TNFα, and IFNγ)
released into meninges in basal conditions (**Figure 3**). The level of the pro-inflammatory MCP-
1 was significantly higher in K14 animals compared to their WT controls (**Figure 3C**). On the
contrary, the basal level of IL-12p70 (also a pro-inflammatory cytokine) was significantly
decreased in K14 animals compared to WT littermates (**Figure 3E**). No differences in the levels
of all the other measured cytokines were observed in basal conditions between the two
genotypes (**Figure 3**).

To mimic inflammatory conditions in the meanings, we used BzATP, a stable ATP analogue
(Baraldi et al., 2005), to stimulate the pro-inflammatory P2X7 receptors (Takenaka et al., 2016)
in the dural immune cells, including mast cells (Nurkhametova et al., 2019). Then, we
measured the cytokine release from meninges of both WT and K14 animals. Stimulation with
100 µM BzATP significantly increased the levels of IL-10 and TNFα (**Figure 4B, D**).
Interestingly, while the IL-10 increase did not differ between the WT and K14 mice, BzATP-
stimulated release of the pro-inflammatory cytokine TNFα was significant exclusively in the
WT genotype (**Figure 4D**). BzATP stimulation did not alter the levels of other measured
cytokines.

3.3 Decreased CGRP release in meninges of mice lacking lymphatic system

Next, we measured the level of CGRP released in meninges in both basal and stimulated
conditions (**Figure 5**). Basal CGRP levels, as measured in two consecutive samples (baselines

BL), were similar in both genotypes. Interestingly, stimulation with 30 mM KCl, which leads
to CGRP release through the stimulation of peptidergic nerve fibers, induced a significantly
lower increase of CGRP levels in K14 compared to WT hemiskulls (**Figure 5A**).

3.4   Spontaneous and ATP-induced spiking activity in meningeal nerves of mice
242
lacking meningeal lymphatic system

To address the state of trigeminal nociception in animals without the lymphatic system, we
recorded spiking activity from meningeal afferents. First, we recorded spiking activity in
trigeminal nerve fibers in hemiskulls under basal conditions. Thereafter, we stimulated nerve
terminals with ATP, a powerful trigger of local nociception (Koroleva et al., 2019; Yegutkin
et al., 2016) (**Figure 6A**). In both genotypes, we observed a comparable increase in the
nociceptive-related firing after ATP stimulation (**Figure 6B**).

3.5   Unchanged patterns of meningeal innervation
250

To evaluate the density of trigeminal innervation in hemiskull preparations from WT and K14
animals we stained meningeal preparations with the antibodies specific to the neuronal marker
β-tubulin III (**Figure 7**) (Cáceres et al., 1986). After that, we applied a semi-automated
segmentation method to evaluate the following parameters: total innervation length (indirect
evaluation of the innervation density in the meninges) and number of terminal points (*i.e.*,
trigeminal fiber terminal endings, likely releasing CGRP). None of the analyzed parameters
showed a difference between the genotypes.

4   Discussion
258

Our study evaluated the role of the meningeal lymphatic system in the regulation of the
neurochemical profile and functional properties of the meningeal trigeminovascular system
(TVS), implicated in migraine pain. In K14 mice, lacking mLVs, we found that the stimulated
release of CGRP, as well as the release of pro-inflammatory IL-12p70 and TNFα were
decreased. On the contrary, we observed an increase in the levels of the pro-inflammatory
cytokine MCP-1. The density of trigeminal meningeal innervation and nociceptive activity of
nerve terminals were not altered in animals lacking meningeal lymphatic system. The main
findings are summarized in **Figure 8**.

4.1   Dural inflammation responding to the deficiency of the lymphatic system
267

Meninges represent the main site for the generation of migraine pain (Messlinger, 2009;
Pietrobon and Moskowitz, 2013). This brain surrounding membranes comprise blood vessels,
sensory nerve fibers and a high diversity of local immune cells (McIlvried et al., 2017; Olesen
et al., 2009; Pietrobon and Moskowitz, 2013). Among the latter, lymphocytes and mast cells
have been reported to be present in high amount dura mater membranes (McIlvried et al.,

2017). In migraine-like conditions, meninges are involved in so-called 'neurogenic
inflammation', which is induced and maintained by neuropeptides, such as CGRP, released
from trigeminal nerve fibers (Eftekhari et al., 2013; Mason et al., 1984). CGRP is currently
recognized as one of the central migraine mediator, with multiple targets including dural
vessels (Goadsby et al., 1988), meningeal nerve fibers (Cady et al., 2011) and immune cells
(Ottosson and Edvinsson, 1997). Interestingly, it was proposed, that impaired clearance of
CGRP from perivascular space can be a reason for post-traumatic headache after TBI (Piantino
et al., 2019). Surprisingly, here we demonstrate that the release of CGRP in the meninges from
K14 mice lacking mLVs is lower compared to WT control group, suggesting that K14 animals
are less prone to trigger migraine attacks via this signalling pathway. . Noteworthy, CGRP is
vital for lymphatic capillary formation (in particular, alleviating the severity of postoperative
lymphedema) (Matsui et al., 2019), suggesting a potential positive crosstalk between CGRP
signalling and activity of the meningeal lymphatic system. As local trigeminal nerve fibers are
the main source of meningeal CGRP (Eftekhari et al., 2013; Mason et al., 1984), we proposed
that the lymphatic impairment causes either a reduced storage of CGRP in trigeminal nerve
terminals or lowered meningeal innervation by 'peptidergic' nerve fibers (Keller and Marfurt,
1991).
Previous studies on patients have mainly focused on cytokine profiling in the systemic
circulation (Fidan et al., 2006; Khaiboullina et al., 2017; Oliveira et al., 2017; Perini et al.,
2005; Sarchielli et al., 2004; Yücel et al., 2016), whereas the production of cytokines in the
meninges have not been fully characterized. Various immune cells can be identified in the
meninges, including cells transported within the mLVs (McIlvried et al., 2017; Alitalo, 2011;
Hayday et al., 2001). These cells constitute the primary and major source of multiple pro-
inflammatory and anti-inflammatory cytokines. Here we demonstrate that in the meninges the
levels of IL-6, IL-10, MCP-1, TNFα, IL-12p70, and IFNγ are detectable already at basal
conditions. In particular, we found the presence of the pro-inflammatory cytokines TNFα and
IL-6, known to be associated with migraine pathology (Bø et al., 2009; Khaiboullina et al.,
2017; Perini et al., 2005; Sarchielli et al., 2006; Wang et al., 2015; Yücel et al., 2016). Notably,
the level of IL-6 was exceptionally high but did not differ between the two genotypes.
However, we found that the absence of functional mLVs was associated with the increased
levels of the pro-inflammatory cytokine MCP-1 (Wood et al., 2014). MCP-1 is known as the
activator of mast cells, provoking release of mast cells derived components (Lv et al., 2012).
In addition, MCP-1 delays mast cell death (Lopes et al., 2019). This supports our finding of
higher number of mast cells in meninges lacking mLVs. Notably, an increase in MCP-1 was
reported in the cerebrospinal fluid of patients during a migraine attack (Bø et al., 2009). In
contrast, the levels of the pro-inflammatory cytokine (IL-12p70) (Martins et al., 2011), which
is typically released from macrophages and dendritic cells (Müller-Berghaus et al., 2004), were
reduced in the meninges from K14 animals lacking mLVs. It has been shown that IL12-p70
plasma level was increased in migraine patients during the interictal phase (Oliveira et al.,

2017). The reduction of IL12-p70 in K14 animals lacking mLVs probably suggests a decreased
inflammatory state in the meninges. Notably, mast cells can release IL12-p70 (Mukai et al.,
2018) in response to CGRP-induced degranulation (Ottosson and Edvinsson, 1997;
Theoharides et al., 2005). Our data showing decreased levels of CGRP in animals lacking
mLVs may imply that the absence of a functional meningeal lymphatic system leads to a
reduced tendency to develop a migraine through several downstream mechanisms controlled
by this neuropeptide.
Extracellular ATP, acting primarily via P2X7 receptors, is a powerful trigger of inflammatory
reactions involving different dural immune cells, including mast cells (Burnstock, 2016;
Karmakar et al., 2016; Nurkhametova et al., 2019, 2018). Consistent with this view, we found
that the stimulation of meninges with the P2X7 agonist BzATP stimulated the release of the
pro-inflammatory TNFα, a promoter of trigeminal pain (Franceschini et al., 2013). Notably,
ATP receptor stimulation resulted in the significant release of TNFα only in the meninges from
WT animals, but not in the meninges lacking the mLVs. TNFα is the main migraine-related
cytokine (Martami et al., 2018; Oliveira et al., 2017; Perini et al., 2005; Yücel et al., 2016). In
line with this, TNFα has been reported as the agent, which can stimulate CGRP transcription
(Bowen et al., 2006). Thus, the lack of increase in TNFα levels after ATP receptor stimulation
is consistent with our other findings on the reduced level of CGRP in the K14 animals. Taken
together, the lack of meningeal lymphatic system may induce a local shift in the balance of
various humoral factors (i.e., CGRP and pro-inflammatory cytokines such as TNFα),
suggesting a reduction in the neurogenic inflammation signalling.

4.2   Meningeal nociception and morphological properties of nerve terminals

ATP is a powerful trigger of meningeal nociception (Koroleva et al., 2019; Yegutkin et al.,
2016). Notably, ATP can operate either directly via neuronal P2X3 receptors (Yegutkin et al.,
2016), or by degranulating mast cells and release of serotonin, which acts via 5-HT3 receptors
at meningeal nerve terminals (Koroleva et al., 2019). However, despite elevated mast cell levels
found in K14 mice, these mechanisms do not lead to significant differences in spiking activity
between the two genotypes.
Since CGRP is released by trigeminal afferents, we tested if the diminished CGRP release,
observed in K14 meninges, resulted from the reduced density of trigeminal innervation of dura
mater. To investigate this hypothesis, we analyzed several parameters of meningeal
innervation. Meningeal sensory nerve fibers are characterized mainly by nociceptive Aδ and
peptidergic C-fibers (Levy and Strassman, 2002), with the latter considered as the primary
source of CGRP (Eftekhari et al., 2013). Analysis of total length of axonal fibers and of the
number of terminal segments of meningeal afferents in the meninges of K14 mice did not show
any difference compared to WT littermates. This suggests that the lower CGRP released in
K14 mice is not affected by an altered density of meningeal innervation, but rather represents

the different content of neuropeptides or a reduced functionality of the CGRP releasing
machinery.

In conclusion, our data show that innate deficiency in the meningeal lymphatics (as observed
in the K14 mice used in this study) results in the alteration of pro-inflammatory signals in the
meninges, including the increased recruitment of mast cells and, on the other side, a decrease
in the levels of the principal migraine-associated pro-inflammatory cytokines. This novel
phenotype suggests the presence of modulatory (compensatory) mechanisms aimed at
balancing the local inflammation in the meninges and limit the triggering of nociceptive
signals.

4.3   Limitations of the model

To the best of our knowledge, this is the first study investigating the interactions between the
meningeal lymphatic system, neurogenic inflammation and nociception, which can contribute
to generation of migraine pain. However, the model used in our study has some limitations.
First, all our experiments have been conducted in *ex vivo* conditions. In this preparation, the
meningeal lymphatic system is not active, thus affecting the recruitment and trafficking of
immune cells and clearance of migraine mediators. Second, our animals were not sensitized to
mimic migraine attacks. Third, only ATP (although one of most powerful algogen) was tested
as a trigger of the nociceptive response. Forth, it has been previously reported that K14 mice
present alterations in both   systemic (Thomas et al., 2012) and neuro-immunity
(Wojciechowski et al., 2019), which involves mainly the T lymphocytes, which is only a subset
of immune cells known to be involved in migraine (Nurkhametova et al., 2018).

Therefore, in future studies, our results needs to be confirmed in *in vivo* migraine models (e.g.,
nitroglycerin injection or CSD induction) (Demartini et al., 2019; Van Den Maagdenberg et
al., 2004) to better understand the impact of the meningeal lymphatic system on migraine state.

4.4   Pathophysiological implications

Here we explored a possible contribution of mLVs in migraine pathology, given the generally
accepted central role of the meningeal tissues in this disorder. Our main working hypothesis
was that the lack of functional meningeal lymphatic system would affect the clearance of the
dural environment, resulting in the exaggeration of neurogenic inflammation and nociception,
and leading to more severe symptoms of migraine.

Indeed, we found that the lack of the meningeal lymphatic is related to pro-nociceptive changes
in the trigeminovascular system relevant to migraine pathology (*i.e.*, enhanced expression of
the pro-inflammatory cytokine MCP-1). However, we surprisingly also found that the
expression of the key migraine mediator CGRP and of the pro-inflammatory cytokines TNFα
and IL-12p70 was decreased. This suggest that the absence of mLVs caused not uniform, but
rather more complex and eventually functionally opposite, changes in the inflammatory and
nociceptive mechanisms in the meninges. Thus, the lack of mLVs, without specific promoters

of activated genetic factors, is not *per se* the main cause of migraine state, likely due to the
presence of the compensatory mechanisms primerely aimed at balance the local inflammation.
388

5   Conflict of Interest

390 None of the authors has any conflict of interest to disclose. The authors confirm that have read
391 the Journal's position on issues involved in ethical publication and affirm that this report is
392 consistent with those guidelines.

6   Author Contributions

NM, FMN and RashidG originally conceived the study; NM, FMN, RashidG elaborated the
study design; NM, KK, RaisaG performed experiments; NM, AA, OG, AS, JT analyzed the
data; NM, AA, OG, JT performed the statistical analysis; NM, FMN, AA, AS, FNM, RashidG
drafted a significant portion of the manuscript and figures. TM supported the study and edited
the manuscript. All authors contributed to manuscript revision, read, and approved the
submitted version.

7   Funding

401 This study has been supported by the Academy of Finland (Grant 325392 for NM, RG,
402 RashidG; Research Fellowship 309479/2017 for FMN) and by the Doctoral Program in
403 Molecular Medicine of University of Eastern Finland (for NM).

8   Acknowledgements

405 For help with cytokine release assay, we would like to thank Sara Wojciechowski, Dilyara
406 Nurkhametova and Sanna Loppi. For meningeal immunostaining, we would like to thank Maria
407 Vihma.

9   Data Availability

409 The raw data supporting the conclusions of this manuscript will be made available by the
410 corresponding author, upon reasonable request, to any qualified researcher.
411

References

Abdollahzadeh, A., Belevich, I., Jokitalo, E., Tohka, J., Sierra, A., 2019a. Automated 3D Axonal Morphometry of White Matter. Sci. Rep. 9, 1–16. https://doi.org/10.1038/s41598-019-42648-2

Abdollahzadeh, A., Sierra, A., Tohka, J., 2019b. Cylindrical shape decomposition for 3D segmentation of tubular objects. arXiv:1911.00571v2.

Ahn, J.H., Cho, H., Kim, J.H., Kim, S.H., Ham, J.S., Park, I., Suh, S.H., Hong, S.P., Song, J.H., Hong, Y.K., Jeong, Y., Park, S.H., Koh, G.Y., 2019. Meningeal lymphatic vessels at the skull base drain cerebrospinal fluid. Nature 572, 62–66. https://doi.org/10.1038/s41586-019-1419-5

Alitalo, K., 2011. The lymphatic vasculature in disease. Nat. Med. https://doi.org/10.1038/nm.2545

Antila, S., Karaman, S., Nurmi, H., Airavaara, M., Voutilainen, M.H., Mathivet, T., Chilov, D., Li, Z., Koppinen, T., Park, J.H., Fang, S., Aspelund, A., Saarma, M., Eichmann, A., Thomas, J.L., Alitalo, K., 2017. Development and plasticity of meningeal lymphatic vessels. J. Exp. Med. 214, 3645–3667. https://doi.org/10.1084/jem.20170391

Aspelund, A., Antila, S., Proulx, S.T., Karlsen, T.V., Karaman, S., Detmar, M., Wiig, H., Alitalo, K., 2015. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. J. Exp. Med. 212, 991–9. https://doi.org/10.1084/jem.20142290

Baraldi, P., Virgilio, F., Romagnoli, R., 2005. Agonists and Antagonists Acting at P2X7 Receptor. Curr. Top. Med. Chem. 4, 1707–1717. https://doi.org/10.2174/1568026043387223

Benmansour, F., Cohen, L.D., 2011. Tubular structure segmentation based on minimal path method and anisotropic enhancement. Int. J. Comput. Vis. 92, 192–210. https://doi.org/10.1007/s11263-010-0331-0

Bø, S.H., Davidsen, E.M., Gulbrandsen, P., Dietrichs, E., Bovim, G., Stovner, L.J., White, L.R., 2009. Cerebrospinal fluid cytokine levels in migraine, tension-type headache and cervicogenic headache. Cephalalgia 29, 365–372. https://doi.org/10.1111/j.1468-2982.2008.01727.x

Bolay, H., Reuter, U., Dunn, A.K., Huang, Z., Boas, D.A., Moskowitz, M.A., 2002. Intrinsic brain activity triggers trigeminal meningeal afferents in a migraine model. Nat. Med. 8, 136–142. https://doi.org/10.1038/nm0202-136

Bowen, E.J., Schmidt, T.W., Firm, C.S., Russo, A.F., Durham, P.L., 2006. Tumor necrosis
factor-α stimulation of calcitonin gene-related peptide expression and secretion from rat
trigeminal ganglion neurons. J. Neurochem. 96, 65–77. https://doi.org/10.1111/j.1471-
4159.2005.03524.x

Burnstock, G., 2016. P2X ion channel receptors and inflammation. Purinergic Signal. 12, 59–
67. https://doi.org/10.1007/s11302-015-9493-0

Cáceres, A., Banker, G.A., Binder, L., 1986. Immunocytochemical localization of tubulin and
microtubule-associated protein 2 during the development of hippocampal neurons in
culture. J. Neurosci. 6, 714–22.

Cady, R.J., Glenn, J.R., Smith, K.M., Durham, P.L., 2011. Calcitonin gene-related peptide
promotes cellular changes in trigeminal neurons and glia implicated in peripheral and
central sensitization. Mol. Pain 7, 94. https://doi.org/10.1186/1744-8069-7-94

Chan, T.F., Vese, L.A., 2001. Active contours without edges. IEEE Trans. Image Process. 10,
266–277. https://doi.org/10.1109/83.902291

Delépine, L., Aubineau, P., 1997. Plasma Protein Extravasation Induced in the Rat Dura
Mater by Stimulation of the Parasympathetic Sphenopalatine Ganglion. Exp. Neurol.
147, 389–400. https://doi.org/10.1006/exnr.1997.6614

Demartini, C., Greco, R., Zanaboni, A.M., Sances, G., De Icco, R., Borsook, D., Tassorelli,
C., 2019. Nitroglycerin as a comparative experimental model of migraine pain: From
animal to human and back. Prog. Neurobiol.
https://doi.org/10.1016/j.pneurobio.2019.02.002

Ebersberger, A., Takac, H., Richter, F., Schaible, H.-G., 2006. Effect of sympathetic and
parasympathetic mediators on the release of calcitonin gene-related peptide and
prostaglandin E from rat dura mater, in vitro. Cephalalgia 26, 282–289.
https://doi.org/10.1111/j.1468-2982.2005.01035.x

Eftekhari, S., Warfvinge, K., Blixt, F.W., Edvinsson, L., 2013. Differentiation of nerve fibers
storing CGRP and CGRP receptors in the peripheral trigeminovascular system. J. Pain
14, 1289–1303. https://doi.org/10.1016/j.jpain.2013.03.010

Fidan, I., Yüksel, S., Ýmir, T., Irkeç, C., Aksakal, F.N., 2006. The importance of cytokines,
chemokines and nitric oxide in pathophysiology of migraine. J. Neuroimmunol. 171,
184–188. https://doi.org/10.1016/j.jneuroim.2005.10.005

Franceschini, A., Vilotti, S., Ferrari, M.D., van den Maagdenberg, A.M.J.M., Nistri, A.,

Fabbretti, E., 2013. TNFα Levels and Macrophages Expression Reflect an Inflammatory
Potential of Trigeminal Ganglia in a Mouse Model of Familial Hemiplegic Migraine.
PLoS One 8, e52394. https://doi.org/10.1371/journal.pone.0052394

Giniatullin, R., Gupta, K., Theoharides, T., 2019. Editorial: Mast Cells in Itch, Pain and
Neuro-Inflammation. Front. Cell. Neurosci. 13, 521.
https://doi.org/10.3389/fncel.2019.00521

Giniatullin, R., Nistri, A., Fabbretti, E., 2008. Molecular mechanisms of sensitization of pain-
transducing P2X3 receptors by the migraine mediators CGRP and NGF. Mol. Neurobiol.
https://doi.org/10.1007/s12035-008-8020-5

Goadsby, P.J., Edvinsson, L., Ekman, R., 1988. Release of vasoactive peptides in the
extracerebral circulation of humans and the cat during activation of the
trigeminovascular system. Ann. Neurol. 23, 193–196.
https://doi.org/10.1002/ana.410230214

Hayday, A., Theodoridis, E., Ramsburg, E., Shires, J., 2001. Intraepithelial lymphocytes:
Exploring the Third Way in immunology. Nat. Immunol. https://doi.org/10.1038/ni1101-
997

Karmakar, M., Katsnelson, M.A., Dubyak, G.R., Pearlman, E., 2016. Neutrophil P2X7
receptors mediate NLRP3 inflammasome-dependent IL-1β secretion in response to ATP.
Nat. Commun. 7, 10555. https://doi.org/10.1038/ncomms10555

Keller, J.T., Marfurt, C.F., 1991. Peptidergic and serotoninergic innervation of the rat dura
mater. J. Comp. Neurol. 309, 515–534. https://doi.org/10.1002/cne.903090408

Khaiboullina, S.F., Mendelevich, E.G., Shigapova, L.H., Shagimardanova, E., Gazizova, G.,
Nikitin, A., Martynova, E., Davidyuk, Y.N., Bogdanov, E.I., Gusev, O., Van Den
Maagdenberg, A.M.J.M., Giniatullin, R.A., Rizvanov, A.A., 2017. Cerebellar atrophy
and changes in cytokines associated with the CACNA1A R583Q mutation in a Russian
familial hemiplegic migraine type 1 family. Front. Cell. Neurosci. 11, 263.
https://doi.org/10.3389/fncel.2017.00263

Kilinc, E., Guerrero-Toro, C., Zakharov, A., Vitale, C., Gubert-Olive, M., Koroleva, K.,
Timonina, A., Luz, L.L., Shelukhina, I., Giniatullina, R., Tore, F., Safronov, B. V.,
Giniatullin, R., 2017. Serotonergic mechanisms of trigeminal meningeal nociception:
Implications for migraine pain. Neuropharmacology 116, 160–173.
https://doi.org/10.1016/j.neuropharm.2016.12.024

Koroleva, K., Gafurov, O., Guselnikova, V., Nurkhametova, D., Giniatullina, R., Sitdikova,

G., Mattila, O.S., Lindsberg, P.J., Malm, T.M., Giniatullin, R., 2019. Meningeal mast
cells contribute to ATP-induced nociceptive firing in trigeminal nerve terminals: Direct
and indirect purinergic mechanisms triggering migraine pain. Front. Cell. Neurosci. 13.
https://doi.org/10.3389/fncel.2019.00195

Koyuncu Irmak, D., Kilinc, E., Tore, F., 2019. Shared fate of meningeal mast cells and
sensory neurons in migraine. Front. Cell. Neurosci. 13.
https://doi.org/10.3389/fncel.2019.00136

Kwon, S., Moreno-Gonzalez, I., Taylor-Presse, K., Edwards, G., Gamez, N., Calderon, O.,
Zhu, B., Velasquez, F.C., Soto, C., Sevick-Muraca, E.M., 2019. Impaired Peripheral
Lymphatic Function and Cerebrospinal Fluid Outflow in a Mouse Model of Alzheimer's
Disease. J. Alzheimer's Dis. 69, 585–593. https://doi.org/10.3233/JAD-190013

Levy, D., 2009. Migraine pain, meningeal inflammation, and mast cells. Curr. Pain Headache
Rep. 13, 237–40.

Levy, D., Strassman, A.M., 2002. Mechanical response properties of A and C primary
afferent neurons innervating the rat intracranial dura. J Neurophysiol 88, 3021–3031.
https://doi.org/10.1152/jn.00029.2002

Loberg, R.D., Day, L.S.L., Harwood, J., Ying, C., St. John, L.N., Giles, R., Neeley, C.K.,
Pienta, K.J., 2006. CCL2 is a potent regulator of prostate cancer cell migration and
proliferation. Neoplasia 8, 578–586. https://doi.org/10.1593/neo.06280

Lopes, J.P., Stylianou, M., Backman, E., Holmberg, S., Ekoff, M., Nilsson, G., Urban, C.F.,
2019. Cryptococcus neoformans Induces MCP-1 Release and Delays the Death of
Human Mast Cells. Front. Cell. Infect. Microbiol. 9.
https://doi.org/10.3389/fcimb.2019.00289

Louveau, A., Smirnov, I., Keyes, T.J., Eccles, J.D., Rouhani, S.J., Peske, J.D., Derecki, N.C.,
Castle, D., Mandell, J.W., Lee, K.S., Harris, T.H., Kipnis, J., 2015. Structural and
functional features of central nervous system lymphatic vessels. Nature 523, 337–341.
https://doi.org/10.1038/nature14432

Lv, J., Huang, Y., Zhu, S., Yang, G., Zhang, Y., Leng, J., Bo, J., Liu, D., 2012. MCP-1-
Induced Histamine Release from Mast Cells Is Associated with Development of
Interstitial Cystitis/Bladder Pain Syndrome in Rat Models. Mediators Inflamm. 2012, 1–
9. https://doi.org/10.1155/2012/358184

Mäkinen, T., Jussila, L., Veikkola, T., Karpanen, T., Kettunen, M.I., Pulkkanen, K.J.,
Kauppinen, R., Jackson, D.G., Kubo, H., Nishikawa, S.I., Ylä-Herttuala, S., Alitalo, K.,

2001. Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice
expressing soluble VEGF receptor-3. Nat. Med. 7, 199–205.
https://doi.org/10.1038/84651

Martami, F., Razeghi Jahromi, S., Togha, M., Ghorbani, Z., Seifishahpar, M., Saidpour, A.,
2018. The serum level of inflammatory markers in chronic and episodic migraine: a
case-control study. Neurol. Sci. 39, 1741–1749. https://doi.org/10.1007/s10072-018-
3493-0

Martins, T.B., Rose, J.W., Jaskowski, T.D., Wilson, A.R., Husebye, D., Seraj, H.S., Hill,
H.R., 2011. Analysis of proinflammatory and anti-inflammatory cytokine serum
concentrations in patients with multiple sclerosis by using a multiplexed immunoassay.
Am. J. Clin. Pathol. 136, 696–704. https://doi.org/10.1309/AJCP7UBK8IBVMVNR

Mason, R.T., Peterfreund, R.A., Sawchenko, P.E., Corrigan, A.Z., Rivier, J.E., Vale, W.W.,
1984. Release of the predicted calcitonin gene-related peptide from cultured rat
trigeminal ganglion cells. Nature 308, 653–5.

Matsui, S., Tanaka, M., Kamiyoshi, A., Sakurai, T., Ichikawa-Shindo, Y., Kawate, H., Dai,
K., Cui, N., Wei, Y., Tanaka, M., Kakihara, S., Nakamura, K., Yamauchi, A., Ishida, K.,
Tanaka, S., Kawamata, M., Shindo, T., 2019. Endogenous Calcitonin Gene–Related
Peptide Deficiency Exacerbates Postoperative Lymphedema by Suppressing Lymphatic
Capillary Formation and M2 Macrophage Accumulation. Am. J. Pathol. 189, 2487–
2502. https://doi.org/10.1016/j.ajpath.2019.08.011

McIlvried, L.A., Cruz, J.A., Borghesi, L.A., Gold, M.S., 2017. Sex-, stress-, and sympathetic
post-ganglionic-dependent changes in identity and proportions of immune cells in the
dura. Cephalalgia 37, 36–48. https://doi.org/10.1177/0333102416637832

Messlinger, K., 2009. Migraine: Where and how does the pain originate? Exp. Brain Res.
196, 179–193. https://doi.org/10.1007/s00221-009-1756-y

Mikhailov, N., Leskinen, J., Fagerlund, I., Poguzhelskaya, E., Giniatullina, R., Gafurov, O.,
Malm, T., Karjalainen, T., Gröhn, O., Giniatullin, R., 2019. Mechanosensitive meningeal
nociception via Piezo channels: Implications for pulsatile pain in migraine?
Neuropharmacology 149, 113–123.
https://doi.org/10.1016/J.NEUROPHARM.2019.02.015

Moskowitz, M.A., 1993. Neurogenic inflammation in the pathophysiology and treatment of
migraine. Neurology 43, S16–S20.

Moskowitz, M.A., 1984. The neurobiology of vascular head pain. Ann. Neurol. 16, 157–168.

https://doi.org/10.1002/ana.410160202

Moskowitz, M.A., Romero, J., Reinhard, J.F., Melamed, E., Pettibone, D.J., 1979.
NEUROTRANSMITTERS AND THE FIFTH CRANIAL NERVE: IS THERE A
RELATION TO THE HEADACHE PHASE OF MIGRAINE? Lancet 314, 883–885.
https://doi.org/10.1016/S0140-6736(79)92692-8

Mukai, K., Tsai, M., Saito, H., Galli, S.J., 2018. Mast cells as sources of cytokines,
chemokines, and growth factors. Immunol. Rev. https://doi.org/10.1111/imr.12634

Müller-Berghaus, J., Kern, K., Paschen, A., Nguyen, X.D., Klüter, H., Morahan, G.,
Schadendorf, D., 2004. Deficient IL-12p70 secretion by dendritic cells based on IL12B
promoter genotype. Genes Immun. 5, 431–434. https://doi.org/10.1038/sj.gene.6364102

Noé, F.M., Marchi, N., 2019. Central nervous system lymphatic unit, immunity, and
epilepsy: Is there a link? Epilepsia Open. https://doi.org/10.1002/epi4.12302

Nurkhametova, D., Kudryavtsev, I., Guselnikova, V., Serebryakova, M., Giniatullina, R.R.,
Wojciechowski, S., Tore, F., Rizvanov, A., Koistinaho, J., Malm, T., Giniatullin, R.,
2019. Activation of P2X7 receptors in peritoneal and meningeal mast cells detected by
uptake of organic dyes: Possible purinergic triggers of neuroinflammation in meninges.
Front. Cell. Neurosci. 13. https://doi.org/10.3389/fncel.2019.00045

Nurkhametova, D., Kudryavtsev, I., Khayrutdinova, O., Serebryakova, M., Altunbaev, R.,
Malm, T., Giniatullin, R., 2018. Purinergic profiling of regulatory T-cells in patients
with episodic migraine. Front. Cell. Neurosci. 12.
https://doi.org/10.3389/fncel.2018.00326

Olesen, J., Burstein, R., Ashina, M., Tfelt-Hansen, P., 2009. Origin of pain in migraine:
evidence for peripheral sensitisation. Lancet Neurol. https://doi.org/10.1016/S1474-
4422(09)70090-0

Oliveira, A.B., Bachi, A.L.L., Ribeiro, R.T., Mello, M.T., Tufik, S., Peres, M.F.P., 2017.
Unbalanced plasma TNF-α and IL-12/IL-10 profile in women with migraine is
associated with psychological and physiological outcomes. J. Neuroimmunol. 313, 138–
144. https://doi.org/10.1016/j.jneuroim.2017.09.008

Ottosson, A., Edvinsson, L., 1997. Release of histamine from dural mast cells by substance P
and calcitonin gene-related peptide. Cephalalgia 17, 166–174.
https://doi.org/10.1046/j.1468-2982.1997.1703166.x

Patel, T.K., Habimana-Griffin, L., Gao, X., Xu, B., Achilefu, S., Alitalo, K., McKee, C.A.,

Sheehan, P.W., Musiek, E.S., Xiong, C., Coble, D., Holtzman, D.M., 2019. Dural
lymphatics regulate clearance of extracellular tau from the CNS. Mol. Neurodegener. 14.
https://doi.org/10.1186/s13024-019-0312-x

Perini, F., D'Andrea, G., Galloni, E., Pignatelli, F., Billo, G., Alba, S., Bussone, G., Toso, V.,
2005. Plasma cytokine levels in migraineurs and controls. Headache 45, 926–931.
https://doi.org/10.1111/j.1526-4610.2005.05135.x

Piantino, J., Lim, M.M., Newgard, C.D., Iliff, J., 2019. Linking Traumatic Brain Injury, Sleep
Disruption and Post-Traumatic Headache: a Potential Role for Glymphatic Pathway
Dysfunction. Curr. Pain Headache Rep. https://doi.org/10.1007/s11916-019-0799-4

Pietrobon, D., Moskowitz, M.A., 2013. Pathophysiology of Migraine. Annu. Rev. Physiol.
75, 365–391. https://doi.org/10.1146/annurev-physiol-030212-183717

Ramachandran, R., 2018. Neurogenic inflammation and its role in migraine. Semin.
Immunopathol. 40, 301–314. https://doi.org/10.1007/s00281-018-0676-y

Rasmussen, M.K., Mestre, H., Nedergaard, M., 2018. The glymphatic pathway in
neurological disorders. Lancet Neurol. https://doi.org/10.1016/S1474-4422(18)30318-1

Sarchielli, P., Alberti, A., Vaianella, L., Pierguidi, L., Floridi, A., Mazzotta, G., Floridi, A.,
Gallai, V., 2004. Chemokine levels in the jugular venous blood of migraine without aura
patients during attacks. Headache 44, 961–968. https://doi.org/10.1111/j.1526-
4610.2004.04189.x

Shatillo, A., Koroleva, K., Giniatullina, R., Naumenko, N., Slastnikova, A.A., Aliev, R.R.,
Bart, G., Atalay, M., Gu, C., Khazipov, R., Davletov, B., Grohn, O., Giniatullin, R.,
2013. Cortical spreading depression induces oxidative stress in the trigeminal
nociceptive system. Neuroscience 253, 341–349.
https://doi.org/10.1016/j.neuroscience.2013.09.002

Shelukhina, I., Mikhailov, N., Abushik, P., Nurullin, L., Nikolsky, E.E., Giniatullin, R., 2017.
Cholinergic Nociceptive Mechanisms in Rat Meninges and Trigeminal Ganglia:
Potential Implications for Migraine Pain. Front. Neurol. 8, 163.
https://doi.org/10.3389/fneur.2017.00163

Suleimanova, A., Talanov, M., Gafurov, O., Gafarov, F., Koroleva, K., Virenque, A., Noe,
F.M., Mikhailov, N., Nistri, A., Giniatullin, R., 2020. Modeling a Nociceptive Neuro-
Immune Synapse Activated by ATP And 5-HT in Meninges: Novel Clues On
Transduction of Chemical Signals into Persistent or Rhythmic Neuronal Firing. Front.
Cell. Neurosci.

Takenaka, M.C., Robson, S., Quintana, F.J., 2016. Regulation of the T Cell Response by
CD39. Trends Immunol. https://doi.org/10.1016/j.it.2016.04.009

Tamura, R., Yoshida, K., Toda, M., 2019. Current understanding of lymphatic vessels in the
central nervous system. Neurosurg. Rev. https://doi.org/10.1007/s10143-019-01133-0

Theoharides, T.C., Donelan, J., Kandere-Grzybowska, K., Konstantinidou, A., 2005. The role
of mast cells in migraine pathophysiology. Brain Res. Rev.
https://doi.org/10.1016/j.brainresrev.2004.11.006

Thomas, S.N., Rutkowski, J.M., Pasquier, M., Kuan, E.L., Alitalo, K., Randolph, G.J.,
Swartz, M.A., 2012. Impaired Humoral Immunity and Tolerance in K14-VEGFR-3-Ig
Mice That Lack Dermal Lymphatic Drainage . J. Immunol. 189, 2181–2190.
https://doi.org/10.4049/jimmunol.1103545

Van Den Maagdenberg, A.M.J.M., Pietrobon, D., Pizzorusso, T., Kaja, S., Broos, L.A.M.,
Cesetti, T., Van De Ven, R.C.G., Tottene, A., Van Der Kaa, J., Plomp, J.J., Frants, R.R.,
Ferrari, M.D., 2004. A Cacna1a knockin migraine mouse model with increased
susceptibility to cortical spreading depression. Neuron 41, 701–710.
https://doi.org/10.1016/S0896-6273(04)00085-6

Wojciechowski, S., Vihma, M., Galbardi, B., Keuters, M., Antila, S., Koistinaho, J., Noe, F.,
2019. The CNS lymphatic system modulates the adaptive neuro-immune response in the
perilesional cortex after brain trauma. bioRxiv 821645. https://doi.org/10.1101/821645

Wood, S., Jayaraman, V., Huelsmann, E.J., Bonish, B., Burgad, D., Sivaramakrishnan, G.,
Qin, S., DiPietro, L.A., Zloza, A., Zhang, C., Shafikhani, S.H., 2014. Pro-inflammatory
chemokine CCL2 (MCP-1) promotes healing in diabetic wounds by restoring the
macrophage response. PLoS One 9, e91574.
https://doi.org/10.1371/journal.pone.0091574

Yegutkin, G.G., Guerrero-Toro, C., Kilinc, E., Koroleva, K., Ishchenko, Y., Abushik, P.,
Giniatullina, R., Fayuk, D., Giniatullin, R., 2016. Nucleotide homeostasis and purinergic
nociceptive signaling in rat meninges in migraine-like conditions. Purinergic Signal. 12,
561–574. https://doi.org/10.1007/s11302-016-9521-8

Yücel, M., Kotan, D., Çiftçi, G.G., Çiftçi, I.H., Cikriklar, H.I., 2016. Serum levels of
endocan, claudin-5 and cytokines in migraine. Eur. Rev. Med. Pharmacol. Sci. 20, 930–
936.

Zakharov, A., Koroleva, K., Giniatullin, R., 2016. Clustering Analysis for Sorting ATP-
Induced Nociceptive Firing in rat Meninges. Bionanoscience 6, 508–512.

https://doi.org/10.1007/s12668-016-0276-z

Zakharov, A., Vitale, C., Kilinc, E., Koroleva, K., Fayuk, D., Shelukhina, I., Naumenko, N.,
Skorinkin, A., Khazipov, R., Giniatullin, R., 2015. Hunting for origins of migraine pain:
cluster analysis of spontaneous and capsaicin-induced firing in meningeal trigeminal
nerve fibers. Front. Cell. Neurosci. 9, 287. https://doi.org/10.3389/fncel.2015.00287

Figure Legend

Figure 1. Migraine-related cell types and active mediators present in meninges. The main
migraine mediator CGRP is released from the meningeal trigeminal nerve fibers, whereas ATP,
serotonin (5-HT) and cytokines are released from local immune cells, including mast cells. We
propose that the local lymphatic system, which supposedly contributes to meningeal drainage,
maintain local homeostasis preventing the accumulation of pro-inflammatory and pro-
nociceptive compounds.

Figure 2. Dural mast cells in WT and K14 mice lacking the lymphatics. Mast cells localize
in meninges in both WT **(A)** and K14 mice **(B)**. Note the presence of mast cells (red triangles)
aligned along the blood vessels (outlined with black dashed curves). **(C)** Mast cells were
counted in WT and K14 animals. Transgenic animals show a non-significant trend towards a
larger number of mast cells (1055 ± 40 cells in WT *vs.* 1177 ± 56 cells in K14, n =
8/experimental group, Mann-Whitney U = 16.5, p = 0.1089 by Mann-Whitney U test). Scale
bar = 200 µm.

Figure 3. Baseline cytokines release from meninges. Concentrations (pg/ml) of following
cytokines are depicted: IL-6 **(A)**, IL-10 **(B)**, MCP-1 **(C)**, TNFα **(D)**, IL-12p70 **(E)** and IFNγ
(F). In K14 animals, we found a significant increase in the levels of MCP-1 **(C)** (829 ± 48
pg/ml in WT *vs.* 1061 ± 72 pg/ml in K14, Mann-Whitney U = 11, #p = 0.0281, Mann-Whitney
U test), whereas the levels of IL-12p70 **(E)** were significantly lower compared to the one
measured in WT littermates (3.5 ± 0.5 pg/ml in WT *vs.* 1.8 ± 0.2 pg/ml in K14, Mann-Whitney
U = 1.0, ##p = 0.0023, Mann-Whitney U test). No difference was found for other interleukins.
Number of replicates for IL-6, IL-10, MCP-1 and TNFα analysis n = 8/experimental group; for
IL-12p70 analysis: n = 7 and 6 for WT and K14, respectively; for IFNγ analysis: n = 5 and 6
for WT and K14, respectively.

Figure 4. Cytokines release from meninges after BzATP stimulation. Scatter plots reporting
the concentrations of cytokines released from meningeal preparation after stimulation with
BzATP: IL-6 **(A)**, IL-10 **(B)**, MCP-1 **(C)**, TNFα **(D)**, IL-12p70 **(E)** and IFNγ **(F)**. BzATP
induced an increase in the release of IL-10 **(B)** (treatment effect WT- K14-control *vs.* WT-

K14-BzATP, $F_{(1, 14)} = 35.75$, ***$p < 0.0001$ by matched two-way ANOVA) and TNFα **(D)**
(WT- K14-control *vs*. WT- K14-BzATP, $F_{(1, 14)} = 12.43$, **$p = 0.0034$ by matched two-way
ANOVA). Post-hoc comparisons revealed that IL-10 increase did not differ between WT and
K14, whereas BzATP-stimulated release of TNFα was significant only in the WT mice (55 ±
18 pg/ml in WT-control *vs*. 112 ± 23 pg/ml in WT-BzATP, $t_{(14)} = 2.933$, *$p = 0.0218$ by
Bonferroni post-hoc test). Number of replicates for IL-6, IL-10, MCP-1, TNFα analysis: n =
8/experimental group; for IL-12p70 analysis: n = 7 and 6 (control) and n = 5 and 6 (BzATP)
for WT and K14, respectively; for IFNγ analysis: n = 5 and 6 (control) and n = 4 and 5 (BzATP)
for WT and K14, respectively.

Figure 5. CGRP released from trigeminal nerve endings in the meninges, from hemiskull
preparation. (A) CGRP release was not altered in K14 hemiskulls under basal conditions: 24
± 4 pg/ml in WT-BL1 *vs*. 20 ± 3 pg/ml in K14-BL1, and 25 ± 3 pg/ml in WT-BL2 *vs*. 17 ± 2
pg/ml in K14-BL2 (n = 7/experimental group, $p > 0.05$ by matched two-way ANOVA with
Bonferroni post-hoc test). The application of 30 mM KCl triggered a significant increase in the
CGRP level in both genotypes (treatment effect: $F_{(2, 24)} = 103.3$, ***$p < 0.0001$ by matched
two-way ANOVA, n = 7/experimental group). Overall CGRP release was higher in WT
animals (genotype effect: $F_{(1, 12)} = 6.025$, #$p = 0.0303$ by matched two-way ANOVA). Post-
hoc comparisons revealed that the concentration of the released CGRP under the stimulation
was significantly lower in the hemiskulls lacking the lymphatic system compared to the WT
preparations (94 ± 16 pg/ml in K14-KCl *vs*.138 ± 11 pg/ml in WT-KCl; $t_{(36)} = 3.816$, ##$p =$
0.0015 by Bonferroni post-hoc test, n = 7/experimental group). **(B)** We propose two possible
explanations for the altered CGRP release: (1) lower density of trigeminal afferents releasing
CGRP, or (2) lower expression of CGRP.

Figure 6. Direct spikes were recorded in WT and K14 hemiskulls under basal conditions
and ATP stimulation. (A) Sample traces of nociceptive recordings in WT and K14 animals
under basal and stimulated conditions. **(B)** The plot represents number of nociception-related
spikes recorded in 5 minutes (median with interquartile range) under basal conditions, during
the application of 100 µM ATP, and during the following washout. ATP induced a significant
increase in nociception activity in both genotypes (baseline *vs*. ATP, $t_{(50)} = -2.65$, * $p = 0.0107$
by linear mixed-effects model; n = 6/experimental group).

Figure 7. Segmentation analysis of meningeal innervation. (A) A representative image of
meninges stained for β-tubulin III (white), segmented for morphological analyses. Blue lines
represent axons, and yellow filled-circle represent terminal points (*i.e*., CGRP-releasing
trigeminal fiber terminal endings). **(B)** Scatter plot depicting the total axonal length (axons
highlighted with blue on panel **A**): 37 ± 5 mm in WT *vs*. 35 ± 4 mm in K14, Mann-Whitney U
= 47, p = 0.8534 by Mann-Whitney U test test. **(C)** Scatter plot depicting the number of terminal
points (yellow dots on panel **A**): 109 ± 10 in WT *vs*. 127 ± 11 in K14, Mann-Whitney U = 32.5,

p = 0.1972 by Mann-Whitney U test. We analyzed meningeal innervation from the right

hemiskulls of 10 independent animals in each experimental group (WT and K14; one image

per animal).

Figure 8. Experimental findings summary. Dysfunctional lymphatic system triggered

alterations in migraine mediator levels in the meninges. Number of mast cells was increased,

supported by increased level of mast cell-activating cytokine MCP-1. CGRP release was

decreased, supported by decreased level of TNFα, a cytokine triggering CGRP transcription.

The level of the pro-inflammatory IL-12p70 was decreased as well. Trigeminal innervation of

the meninges and trigeminal nociception were not changed. Red indicates pro-inflammatory

and pro-nociceptive changes, green indicates anti-inflammatory and anti-nociceptive changes.

Arrow up (↑) indicates an increase in the parameter, arrow down (↓) – decrease, equals sign

(=) indicates unchanged parameters.