The role of the meningeal lymphatic system in local inflammation and trigeminal nociception implicated in migraine pain

Nikita Mikhailov[1], Kseniia Koroleva[1], Ali Abdollahzadeh[1], Raisa Giniatullina[1], Oleg Gafurov[3], Tarja Malm[1], Alejandra Sierra[1], Jussi Tohka[1], Francesco Noe[2,#], Rashid Giniatullin[1,3,#]

[1]*A. I. Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio, 70211, Finland*
[2]*HiLIFE-Neuroscience Center, University of Helsinki, Helsinki, 00290, Finland*
[3]*Laboratory of Neurobiology, Kazan Federal University, Kazan 420008, Russia*
[#]*shared last authorship*
corresponding author:
Francesco Noe, Neuroscience Center, HiLife, Helsinki University, Biomedicum A-wing, Haartmaninkatu 8, P.O.Box 69, Helsinki 00290, Finland, e-mail: francesco.noe@helsinki.fi

Abstract

Background: A system of lymphatic vessels has been recently characterized in the meninges, with a postulated role in 'cleaning' of the brain via cerebro-spinal fluid drainage. As meninges are an important tissue involved in the origin of migraine pain, we hypothesized that malfunctioning of a 'cleaning' lymphatic system should affect the functional properties of meningeal nociception. To test this hypothesis, we studied migraine-related nociceptive and inflammatory mechanisms in the meninges, in a mouse model of primary lymphedema (K14-VEGFR3-Ig mice) characterized by the lack of functional meningeal lymphatic system.

Methods: To study the migraine-related nociceptive and inflammatory mechanisms we recorded spiking activity from meningeal fibers of the trigeminal nerve, estimated the local mast cells infiltration, calcitonin gene-related peptide (CGRP) and cytokine levels (basal and under stimulating conditions), as well as the dura mater innervation in freshly-isolated hemiskull preparations from K14-VEGFR3-Ig (K14) or wild type C57BL/6 mice (WT).

Results: We found that the level of CGRP and the production of TNFα, which both are implicated in migraine, were reduced in meninges of K14 mice. There was a trend of having a larger number of dural mast cells, consistent with the increased level of the mast cell activator MCP-1 found in these animals. In addition, we found an increased spontaneous and ATP-induced nociceptive firing in the meningeal afferents of mice lacking meningeal lymphatic system. However, the patterns of trigeminal innervation in meninges remained unchanged.

Conclusions: In summary, the lack of meningeal lymphatic system leads to a new balance between pro- and anti-inflammatory mechanisms implicated in peripheral nociception, affecting key cellular and humoral factors implicated in migraine.

Keywords

Meningeal lymphatic, neuroinflammation, migraine, pain

Abbreviations:

WT – wild type mice

K14 – K14-VEGFR3-Ig mice

mLVs – meningeal lymphatic vessels

1. Introduction

Migraine is a complex neurological disorder, which originates from the meninges (Bolay et al., 2002; Olesen et al., 2009). Meninges are densely innervated by somatic trigeminal (Ramachandran, 2018; Zakharov et al., 2015) and parasympathetic innervation (Delépine and Aubineau, 1997), which runs along and in contact with meningeal blood vessels (Ebersberger et al., 2006). It has been proposed that so-called sterile neurogenic inflammation (Moskowitz, 1993, 1984; Moskowitz et al., 1979) in the meninges is an important initiator of the pathophysiology of migraine. The initial step in neurogenic inflammation is the activation of the trigeminal nociceptive system in meninges (Moskowitz, 1993). Indeed, trigeminal neurons release neuropeptide calcitonin gene-related peptide (CGRP), which activates dural mast cells (Ottosson and Edvinsson, 1997), promotes sensitization of trigeminal afferents (Cady et al., 2011; Giniatullin et al., 2008), and induces local vasodilation (Goadsby et al., 1988). Vasodilation, in turn, activates mechanosensitive receptors expressed in trigeminal nerves in meninges (Mikhailov et al., 2019). Trigeminal neurons can also be activated by mast cells-secreted serotonin (Kilinc et al., 2017), and by extracellular ATP, which is released from endothelial and mast cells, as well as from neurons, upon stimulation of the migraine mediator CGRP (Koroleva et al., 2019; Suleimanova et al., 2020; Yegutkin et al., 2016). In addition, endothelial (Loberg et al., 2006) and mast cells (Mukai et al., 2018) release various pro-inflammatory cytokines as potential contributors to local inflammation. Moreover, our previous data show that acetylcholine (ACh, known to be released from parasympathetic fibers) can activate trigeminal afferents and induce meningeal mast cell degranulation (Shelukhina et al., 2017).

The presence of these diverse endogenous agents in the meninges suggests the existence of regulatory mechanisms modulating the local levels of these, mostly pro-inflammatory compounds in meninges

in order to prevent potential inflammation and minimize the activity of nociceptive fibers. In analogy with the suggested role of the recently characterized meningeal lymphatic system (Antila et al., 2017; Aspelund et al., 2015; Louveau et al., 2015) **(Figure 1)** in the clearance of the CNS from toxic molecules in neurodegenerative diseases such as Alzheimer's disease, epilepsy, stroke and TBI (Kwon et al., 2019; Noé and Marchi, 2019; Patel et al., 2019; Rasmussen et al., 2018; Tamura et al., 2019), we hypothesized that the meningeal lymphatic system plays a similar protective/cleaning role also in migraine.

Figure 1. Migraine-related cell types and active compounds presented in meninges. The main migraine mediator CGRP is released from trigeminal neurons, whereas ATP, serotonin (5-HT) and cytokines are released from local immune cells including mast cells. We propose that the local lymphatic system, which is supposed to provide drainage of meninges, prevents the overflooding of these tissues from the pro-inflammatory and pro-nociceptive compounds.

2. Methods and materials

2.1. Animals

We used adult (4 – 20 months old) transgenic K14-VEGFR3-Ig (K14) mice lacking the meningeal lymphatic system (Mäkinen et al., 2001) and wild-type C57BL/6 mice (WT). Total number of animal used for this study was 38 WT and 37 TG animals. Animals were housed at the Lab Animal Center

of the University of Eastern Finland under following housing conditions: 12 hours dark/light cycle, grouped housing, given ad libitum access to food and water, 22°C ambient temperature. All procedures with animals were conducted in accordance with the European Community Council Directive of September 22, 2010 (2010/63/EU). Experimental protocols involving the usage of animals were approved by the Animal Care and Use Committee of the University of Eastern Finland (license EKS-004-2014).

2.2. Hemiskull preparation for ex vivo experiments

For the *in vitro* and *ex vivo* experiments (mast cell staining, CGRP release assay, cytokine release assay, and electrophysiology), we used hemiskull preparations with preserved meninges and meningeal trigeminal innervation (Shatillo et al., 2013; Zakharov et al., 2015). In order to prepare the hemiskulls for subsequent analyses, after decapitation, skin, muscles, and connective tissues, as well as brain were removed with particular care to not touch and damage the meninges. Thereafter, hemiskulls were cut along their sagittal axis (*i.e.*, two hemiskulls were prepared from each animal).

2.3. Mast cell staining

For mast cell staining, WT and K14 animals were transcardially perfused with normal saline (6 minutes at 6 ml/min), followed by 4% paraformaldehyde (PFA) (20 minutes at 3 ml/min). Then hemiskulls were prepared as described above, washed three times for 20 minutes in PBS (phosphate buffered saline), and stained for 60 minutes with 0.1% toluidine blue. Stained meninges were immediately imaged, using a Zeiss Axio Zoom.V16 stereomicroscope (Carl Zeiss AG, Germany), at 30x magnification. We imaged two identical regions-of-interest (ROIs) from each hemiskull: one including the superior sagittal sinus (SSS; at the border of the image) and the next one, which partially overlaps with the previous ROI. In total, four images were acquired from each mouse. From each ROI, the number of stained mast was calculated using ImageJ 1.51v software (National Institute of Health, USA).

2.4. CGRP release assay

The CGRP release assay was conducted as previously described (Mikhailov et al., 2019). Briefly, we used a CGRP enzyme immunoassay kit (EIA kit, SPIbio, France) to measure CGRP release from hemiskull preparations. The levels of CGRP were assessed under control and KCl-treated conditions. First, fresh hemiskull preparations from WT and K14 animals were perfused for 40 minutes with artificial cerebrospinal fluid (aCSF in mM: 119 NaCl, 30 $NaHCO_3$, 11 glucose, 2.5 KCl, 1 $MgCl_2$, 1 NaH_2PO_4, 2 $CaCl_2$, and the level of pH was adjusted to 7.4) at room temperature (RT). Then,

hemiskulls were washed four times with 150 µl aCSF for 15 minutes at 37°C to stabilize the basal condition. Samples of aCSF (100 µl each) from the last two washes were collected to assess the basal level of CGRP (control conditions). After the last wash, hemiskulls were bathed with an aCSF solution containing 30 mM KCl for 15 minutes. Thereafter, a 100 µl sample was collected to assess CGRP under KCl-treated condition.

2.5. Cytokine release assay

For cytokine release assay, we used fresh hemiskull preparations from both WT and K14 mice. One hemiskull/animal was used to assess the release of cytokine under the control condition, while the contralateral was used to measure the the release of cytokine under treatment condition (100 µM BzATP, a more stable analogue of ATP was used because of long treatment time – three and a half hours). After the collection of hemiskulls, all procedures were performed at 37°C. Hemiskulls were bathed in 150 µl of either aCSF or aCSF + BzATP for three and a half hours. aCSF samples (50 µl) were taken to assess the level of cytokines in control (treated with aCSF only) or stimulated (aCSF + BzATP) conditions. The levels of aCSF in the hemiskulls were monitored during all the incubation period and, if needed, refilled with the respective treatment solution to compensate for evaporation. To assess the level of cytokines (Il-6, Il-10, MCP-1, TNFα, INFγ, Il-12p70), we used the CBA Mouse Inflammation Kit (Cytometric Bead Array, BD Biosciences, New Jersey, USA). All procedures (sample and standard beads preparation) were performed following instructions provided by the manufacturer. Data were acquired on the CytoFLEX S cytometer (Beckman Coulter Inc, USA) and analyzed with the Flow Cytometric Analysis Program (FCAP) Array software (BD Biosciences, USA).

Due to the low concentration of IL-12p70 and IFNγ in some samples, it was not possible to quantify the exact concentration with the abovementioned method. Thus, for these cytokines number of n was lower than the number of animals used for the assay.

2.6. Electrophysiology

Direct recording of action potentials in trigeminal nerve afferents innervating meninges was conducted as described earlier (Shatillo et al., 2013; Zakharov et al., 2015). Briefly, hemiskulls of WT and K14 animals were placed in a perfusion chamber with a constant flow of approximately 6 ml/min of oxygenated aCSF. In order to access the nervus spinosus of the mandibular branch of the trigeminal nerve, we made a small incision in the dura mater. Then, the nerve ending was sucked into a glass recording electrode filled with aCSF. We recorded 10 minutes of spontaneous activity to get a baseline and to assess the difference in basal activity between WT and K14 animals. We applied

100 µM ATP solution for 10 minutes to stimulate nociceptive firing in the trigeminal nerve (Zakharov et al., 2016), followed by 10 minutes of washout. The recording was conducted with a digital amplifier (ISO 80, WPI Inc, USA) with bandpass 300 Hz – 3 kHz, gain 10000. Signals were digitized at 125 kHz with NIPCI 6221 data acquisition (DA) board (National Instruments, USA) and visualized with WinEDR software (Strathclyde University, UK).

2.7. Meningeal nerve staining for β-tubulin III

For the meningeal nerve staining, meninges from PFA-fixed hemiskulls were isolated, avoiding damage, and stained as previously described (Louveau et al., 2015). Briefly, meningeal whole-mount preparations were incubated with 1x phosphate-buffered saline (PBS) containing 2% normal goat serum (NGS), 1% bovine serum albumin (BSA), 0.1% Triton X-100, and 0.05% Tween 20 for 1 hour at RT. After that, meninges were incubated with rabbit anti-β-tubulin III (1:1000, Cat# T2200, Sigma-Aldrich, USA) overnight at 4°C in PBS containing 1% BSA and 0.5% Triton X-100. Whole-mounts were washed in PBS at RT (3x) followed by incubation with Alexa Fluor 488-conjugated goat anti-rabbit IgG antibody (1:500, Cat# A11008, Invitrogen, USA) for 1 hour at RT in PBS with 1% BSA and 0.5% Triton X-100. After further washing in PBS (3x) and in phosphate buffer (PB; 2x), meninges were mounted with VECTASHIELD mounting medium (Vector Laboratories, USA), including DAPI, and coverslipped.

Images were acquired using a Leica TCS SP8 X confocal system (Leica Microsystems, Germany) using the LAS X software. Images of the region of interest, adjacent to the middle meningeal artery, were acquired with a 20x objective with 0.75 numerical aperture (NA), with an in-plane pixel size of 1.14×1.14 µm^2 and a z-step of 1 µm. The full images were created by merging 4×4 tile scans, covering a total area of 2034×2034 µm^2.

2.8. Axon segmentation and innervation analysis

We developed a semi-automated segmentation technique to annotate axons and analyze the innervation in acquired 3D confocal microscopy images. To segment an axon, an expert input two points along the desired axon on the maximum projection image. Given the two points, a minimal path algorithm detected the centerline of that axon, based on which an active contour segmented the desired axon. In detail, we formed the 2D maximum intensity projection image along the direction of focal planes, i.e., z-axis, and enhanced axons using Frangi filter (Frangi et al., 1998). We normalized the enhanced images to the range [0, 1] and complemented the intensity of the enhanced images, such that axons appeared darker than their background, denoting the image as P. We used Benmansour and Cohen method (Benmansour and Cohen, 2011) to define the minimal path between two given

points p_1 and p_2 as C_{p_1,p_2}, a path which minimizes an energy functional on P. The energy minimized by the minimal path is $E(\gamma) = \int_\gamma P(\gamma(s))ds$, where γ is a path parametrized along its length by s. To find C_{p_1,p_2}, we computed the minimal action map $U: \Omega \to \mathbb{R}^+$ associated to p_1 on the domain $\Omega \subset \mathbb{R}^2$ which is the minimal energy integrated along a path between p_1 and any point $x \in \Omega$. To compute U, we used the fast marching method (Sethian, 1996) with the speed function $1 - P$. The map U has only one local minimum, the source point p_1. Thus, the minimal path C_{p_1,p_2} can be retrieved with a simple gradient descent on U from p_2 to p_1, solving an ordinary differential equation with a standard numerical Runge-Kutta method. The minimal path C_{p_1,p_2} is the axon centerline, because 1-P was constructed using Frangi filter, admitting higher velocity in the center of the axon.

We used the centerlines to segment the underlying axons using Chan-Vese active contours (Chan and Vese, 2001) (Matlab 2018b, Image Processing Toolbox, activecontour), initialized from the centerlines with the speed function $1-P$. We set the contraction bias equal to -0.2, expanding an initial contour, and the maximum number of iterations to 10 to perform the curve evolution. The initial centerlines were deformed on the speed function to adapt to the shape of axons.

We calculated the diameter of axons along their centerlines, as described in (Abdollahzadeh et al., 2019a). We calculated the length of the centerlines in 3D, by first projecting centerlines from $\mathbb{R}^2$ to $\mathbb{R}^3$, then smoothing the curves with an 11[th]-order one-dimensional median filter and a shape-preserving piecewise cubic polynomial. We reported the sum of the length of all centerlines, excluding centerlines, which corresponded with axon bundles. The underlying segment of an axon bundle has a diameter thicker than a certain threshold, where we set the threshold to 8 μm experimentally. We also quantified the innervation complexity by forming a graph from the union of the extracted centerlines as in (Abdollahzadeh et al., 2019b). We reported the number of graph terminal-vertices (endpoints) with the vertex degree equal to 1 and regular-vertices (branching points) with the vertex degree greater than 2.

2.9. Statistical analysis

We performed statistical analysis using the GraphPad Prism 8.4.2 software (GraphPad Software, California, USA). The data are presented as mean ± the standard error of the mean (SEM). We set the alpha-threshold defining the statistical significance as 0.05 for all analyses. We used paired or unpaired Student t-test and matched two-way ANOVA with Bonferroni post-hoc test for our data.

3. Results

3.1. Elevated number of mast cells in meninges of mice without lymphatic system

Meningeal mast cells recently emerged as important players in the initiation of a migraine attack (Giniatullin et al., 2019; Kilinc et al., 2017; Koyuncu Irmak et al., 2019; Levy, 2009). As the meningeal lymphatic vessels (mLVs) are important for the movement of immune cells (Ahn et al., 2019; Louveau et al., 2015), we expected that the lack of mLVs could affect the local accumulation of mast cells. We took advantage of toluidine blue labelling to characterize the profile of meningeal mast cells and assess whether lack of dural lymphatic system in K14 mice alters the number of meningeal mast cells.

In WT animals a substantial portion of mast cells was located along the meningeal blood vessels (**Figure 2.A**), where allegedly are located the mLVs (Antila et al., 2017; Aspelund et al., 2015). A similar pattern of mast cell positioning was observed in K14 animals (**Figure 2.B**).

Figure 2. Dural mast cells in WT and K14 mice lacking the lymphatics. Mast cells localize in meninges both WT **(A)** and K14 mice **(B)**. Note the presence of mast cells (red triangles) alligned along the blood vessels (outlined with black dashed curves). **(C)** Mast cells were counted in WT and K14 animals. Transgenic animals show a non-significant trend towards a larger number of mast cells (1055 ± 40 cells in WT *vs.* 1177 ± 56 cells in K14, n = 8/experimental group, t(14) = 1.784, p = 0.096 by unpaired Student t-test). Scale bar = 200 µm.

K14 animals showed a non-significant trend towards increased number of meningeal mast cells (**Figure 2.C**). Since an increase in the number of mast cells may lead to a more abundant release of local mediators such as serotonin, histamine and ATP (Giniatullin et al., 2019; Levy, 2009; Suleimanova et al., 2020), resulting in meningeal inflammation, and stimulation of the trigeminal afferents (Kilinc et al., 2017; Koroleva et al., 2019) we analyzed the levels of the different migraine-

associated pro-inflammatory molecules and the ultimate activation of the trigeminal nerves in the K14 and their WT littermates mice.

3.2. Altered cytokine release in meninges

We first measured the level of cytokines (IL-6, IL-10, IL-12p70, MCP-1, TNFα, and IFNγ) released into meninges in basal conditions (**Figure 3**). The levels of the pro-inflammatory MCP-1 were significantly higher in K14 animals compared to their WT controls (**Figure 3.C**). On the contrary, the basal level of IL-12p70, which is also a pro-inflammatory cytokine, was significantly decreased in K14 animals compared to WT littermates (**Figure 3.E**). We failed to detect any differences between the genotypes in the levels of all the other measured cytokines in basal conditions (**Figure 3**).

Figure 3. Baseline cytokines release from meninges. Concentrations (pg/ml) of following cytokines are depicted: IL-6 **(A)**, IL-10 **(B)**, MCP-1 **(C)**, TNFα **(D)**, IL-12p70 **(E)** and IFNγ **(F)**. In K14 animals, we found a significant increase in the levels of MCP-1 **(C)** (829 ± 48 pg/ml in WT *vs.* 1061 ± 72 pg/ml in K14, $t(14) = 2.675$, #p = 0.0181, unpaired Student t-test), whereas the levels of IL-12p70 **(E)** were significantly lower compared to

the one measured in WT littermates (3.5 ± 0.5 pg/ml in WT *vs.* 1.8 ± 0.2 pg/ml in K14, t(11) = 3.283, ##p = 0.0073, unpaired Student t-test). No difference was found for other interleukines. Number of replicates were: for IL-6, IL-10, MCP-1, TNFα n = 8/experimental group; for IL-12p70: n = 7 for WT, n = 6 for K14; for IFNγ: n = 5 for WT, n = 6 for K14.

To mimic inflammatory conditions, we used benzoyl ATP (Baraldi et al., 2005) (BzATP), a stable ATP analogue to stimulate the P2X7 receptors (Takenaka et al., 2016) in dural immune cells, including mast cells (Nurkhametova et al., 2019), and measured the cytokine release from meninges of both WT and K14 animals. Indeed, 100 μM BzATP significantly increased the levels of IL-10 and TNFα **(Figure 4.B, D)**. Interestingly, while the IL-10 increase did not differ between the WT and K14 mice, BzATP-stimulated release of the pro-inflammatory cytokine TNFα was significant only in the WT mice (**Figure 4.D**) and not in K14 animals. BzATP stimulation did not alter the levels of other measured cytokines.

Figure 4. Cytokines release from meninges after BzATP stimulation. Scatter plots reporting the concentrations of cytokines released from meningeal preparation after stimulation with BzATP: IL-6 **(A)**, IL-10 **(B)**, MCP-1 **(C)**, TNFα **(D)**, IL-12p70 **(E)** and IFNγ **(F)**. BzATP induced an increase in the release of IL-10 **(B)** (WT- K14-control *vs.* WT- K14-BzATP, $F(1, 14) = 35.75$, ***$p < 0.0001$ by matched two-way ANOVA) and TNFα **(D)** (WT- K14-control *vs.* WT- K14-BzATP, $F(1, 14) = 12.43$, **$p = 0.0034$ by matched two-way ANOVA). Post-hoc comparisons revealed that IL-10 increase did not differ between WT and K14, whereas BzATP-stimulated release of TNFα was significant only in the WT mice (55 ± 18 pg/ml in WT-control *vs.* 112 ± 23 pg/ml in WT-BzATP, $t(14) = 2.933$, *$p = 0.0218$ by Bonferroni post-hoc test). Number of replicates were: for IL-6, IL-10, MCP-1, TNFα $n = 8$/experimental group for both treatments; for IL-12p70: control $n = 7$ for WT, $n = 6$ for K14, BzATP $n = 5$ for WT, $n = 6$ for K14; for IFNγ: for control $n = 5$ for WT, $n = 6$ for K14, for BzATP $n = 4$ for WT, $n = 5$ for K14.

3.3. Decreased CGRP release in meninges of mice lacking lymphatic system

We measured the level of CGRP released in meninges in both basal and stimulated conditions (**Figure 5**). CGRP levels measured at two consecutive samples collected at basal condition (baselines, BL) were similar in both genotypes. Interestingly, stimulation with 30 mM KCl, which leads to CGRP release through stimulation of peptidergic nerve fibers, in K14 mice shows significantly lower levels of CGRP compared to WT hemiskulls (**Figure 5.B**).

Figure 5. CGRP released from trigeminal nerve endings in the meninges, from hemiskull preparation. (A) Scheme proposing trigeminal nerve innervation and CGRP release in WT and K14 mice. **(B)** CGRP release was not altered in K14 hemiskulls under basal conditions: 24 ± 4 pg/ml in WT-BL1 *vs.* 20 ± 3 pg/ml in K14-BL1, and 25 ± 3 pg/ml in WT-BL2 *vs.* 17 ± 2 pg/ml in K14-BL2 (n = 7/experimental group, p > 0.05 by matched two-way ANOVA with Bonferroni post-hoc test). The application of 30 mM KCl triggered a significant increase in the CGRP level in both genotypes (WT- and K14-KCl *vs.* WT- and K14-BL1/BL2, n = 7/experimental group, $F_{(2, 24)} = 103.3$, ***p < 0.0001 by matched two-way ANOVA). Overall CGRP release was higher in WT animals (WT *vs.* K14, n = 7/experimental group, $F_{(1, 12)} = 6.025$, #p = 0.0303 by matched two-way ANOVA). Post-hoc comparisons revealed that the concentration of the released CGRP under the stimulation was significantly lower in the hemiskulls lacking the lymphatic system compared to the WT preparations (138 ± 11 pg/ml in WT-KCl *vs.* 94 ± 16 pg/ml in K14-KCl; n = 7/experimental group, $t_{(36)} = 3.816$, ##p = 0.0015 by Bonferroni post-hoc test).

3.3. Increased spontaneous and ATP induced spiking activity in meningeal nerves of mice lacking meningeal lymphatic system

To address the state of trigeminal nociception in animals without the lymphatic system, we recorded spiking activity from meningeal afferents. First, we recorded spiking activity in trigeminal nerve fibers in hemiskulls under the basal conditions. Thereafter, we stimulated nerve terminals with ATP, a powerful trigger of local nociception (Koroleva et al., 2019; Yegutkin et al., 2016) (**Figure 6.A**).

ATP induced a significant increase in the nociception in both genotypes. Notably, meninges lacking lymphatic vessels (both in basal conditions and under ATP stimulation), showed a significant increase in nociceptive spiking compared to WT (**Figure 6.B**).

Figure 6. Direct spikes were recorded in WT and K14 hemiskulls under basal conditions and ATP stimulation. (A) Sample traces of nociceptive recordings in WT and K14 animals under basal and stimulated conditions. **(B)** The plot represents an average number of pain spikes recorded in 5 minutes (mean ± SEM) under basal conditions, during the application of 100 μM ATP, and during the following washout. ATP induced a significant increase in nociception activity in both genotypes (baseline *vs.* ATP, WT: n = 8, K14: n = 7, $F_{(3, 39)} = 15.74$, ***$p < 0.0001$ by matched two-way ANOVA). Overall spiking activity was higher in K14 hemiskulls (WT *vs.* K14, WT: n = 8, K14: n = 7, $F_{(1, 13)} = 4.752$, #$p = 0.0482$ by matched two-way ANOVA).

3.5. Unchaged patterns of meningeal innervation

To evaluate the density of trigeminal innervation in hemiskull preparations from WT and K14 animals we stained meningeal preparations with the antibodies specific to the neuronal marker β-tubulin III (**Figure 7**) (Cáceres et al., 1986). After that, we applied a semi-automated segmentation method to evaluate the following parameters: total innervation length (indirect evaluation of the innervation density in the meninges), number of branching points (*i.e.*, the complexity of innervation network) and the number of terminal points (i.e., trigeminal fiber terminal endings, likely releasing CGRP). None of the analyzed parameters showed a difference between the genotypes.

Figure 7. Segmentation analysis of meningeal innervation. (A) A representative image of meninges stained for β-tubulin III (white), segmented for morphological analyses. Blue lines represent axons, magenta filled-circles represent branching points, and yellow filled-circle represent terminal points (*i.e.*, CGRP-releasing trigeminal fiber terminal endings). **(B)** Scatter plot depicting the total axonal length (axons highlighted with blue on panel **A**): 37 ± 5 mm in WT *vs.* 35 ± 4 mm in K14, n = 10/experimental group, t(18) = 0.3498, p = 0.7306 by unpaired Student *t*-test. **(C)** Scatter plot depicting the number of terminal points (yellow dots on panel **A**): 109 ± 10 in WT *vs.* 127 ± 11 in K14, n = 10/experimental group, t(18) = 1.160, p = 0.2614 by unpaired Student t-test. We analyzed meningeal innervation from the right hemiskulls of 10 independent animals in each experimental group (WT and K14; one image per animal).

4. Discussion

Here we evaluated for the first time the role of the meningeal lymphatic system in the regulation of the neurochemical profile and functional properties of the meningeal trigeminovascular system

(TVS), implicated in migraine pain using the K14 mice lacking meningeal lymphatics. Our data show that the level of CGRP is decreased in K14 animals, the sensitivity to the algogen ATP is higher. In addition, in animals lacking meningeal lymphatic system, we found decreased release of pro-inflammatory IL-12p70 and TNFα and increase in pro-inflammatory MCP-1 (summarized in **Figure 8**).

Figure 8. Experimental findings summary. Dysfunctional lymphatic system triggered several alterations in meninges. Number of mast cells was increased, supported by increased level of mast cell-activating cytokine MCP-1. CGRP release was decreased, supported by decreased level of TNFα, cytokine triggering CGRP transcription. Level of pro-inflammatory IL-12p70 was decreased. Trigeminal innervation of the meninges was not changed, however, trigeminal nociception was elevated. Red indicates pro-inflammatory and pro-nociceptive changes, green indicates anti-inflammatory and anti-nociceptive changes. Arrows up (↑) indicate increase in parameters, arrows down (↓) – decrease, equals sign (=) indicates unchanged parameter.

4.1 Dural inflammation responding to the deficiency of the lymphatic system

Meningeal tissues represent the main site for the generation of migraine pain. They comprise of blood vessels, sensory nerve fibers and a high diversity of local immune cells (McIlvried et al., 2017; Olesen et al., 2009; Pietrobon and Moskowitz, 2013). Among the latter, lymphocytes and mast cells have been reported to be present in dura mater membranes (McIlvried et al., 2017). In migraine-like

conditions, meninges are involved in so-called 'neurogenic inflammation,' which is induced and maintained by neuropeptides, such as CGRP, released from trigeminal nerve fibers (Eftekhari et al., 2013; Mason et al., 1984). CGRP is currently recognized as one of the central migraine mediator, with multiple targets including dural vessels (Goadsby et al., 1988), meningeal nerve fibers (Cady et al., 2011) and immune cells (Ottosson and Edvinsson, 1997). Interestingly, it was proposed, that impaired clearance of CGRP from perivascular space can be a reason for post-traumatic headache after TBI (Piantino et al., 2019). Surprisingly, here we demonstrate that the release of CGRP in the meninges from K14 mice lacking mLVs is lower compared to meninges in the WT control group suggesting that K14 animals are less prone to triggering migraine attacks via this particular mechanism. Interestingly, CGRP is vital for lymphatic capillary formation (in particular, alleviating the severity of postoperative lymphedema) (Matsui et al., 2019). These data imply a potential positive crosstalk between CGRP signaling and activity of the meningeal lymphatic system. As the source of meningeal CGRP are local trigeminal nerve fibers (Eftekhari et al., 2013; Mason et al., 1984), we propose that the lymphatic impairment causes either reduced storage of CGRP in trigeminal nerve terminals or lowered meningeal innervation by 'peptidergic' nerve fibers (Keller and Marfurt, 1991). Previous studies on patients have mainly focused on cytokine profiling in the systemic circulation (Fidan et al., 2006; Khaiboullina et al., 2017; Oliveira et al., 2017; Perini et al., 2005; Sarchielli et al., 2004; Yücel et al., 2016), whereas the production of cytokines in the meninges have not been fully characterized. Various immune cells can be identified in the meninges, including cells transported within the mLVs (McIlvried et al., 2017; Alitalo, 2011; Hayday et al., 2001). These cells constitute the primary and major source of multiple proinflammatory and anti-inflammatory cytokines. Here we demonstrate the presence at detectable levels of IL-6, IL-10, MCP-1, TNFα, IL-12p70, and IFNγ even in the basal conditions in meninges. In particular, we found the presence of the pro-inflammatory cytokines TNFα, IL-1β and IL-6 known to be associated with migraine pathology (Bø et al., 2009; Khaiboullina et al., 2017; Perini et al., 2005; Sarchielli et al., 2006; Wang et al., 2015; Yücel et al., 2016). Further, the absence of functional mLVs was associated with the increased levels of the pro-inflammatory cytokine MCP-1 (Wood et al., 2014). MCP-1 is known as the activator of mast cells, provoking release of mast cells derived components (Lv et al., 2012). In addition, MCP-1 delays mast cell death (Lopes et al., 2019). This supports our finding of higher number of mast cells in meninges lacking mLVs. Notably, an increase in MCP-1 was reported in the cerebrospinal fluid of patients during a migraine attack (Bø et al., 2009). In contrast, the levels of the pro-inflammatory cytokine (IL-12p70) (Martins et al., 2011), which is typically released from macrophages and dendritic cells (Müller-Berghaus et al., 2004), were reduced in the meninges from K14 animals lacking mLVs. Interestingly, it has been shown that IL12-p70 plasma level was increased in migraine patients during

the interictal phase (Oliveira et al., 2017). The reduction of IL12-p70 in K14 animals lacking mLVs probably suggests a decreased inflammatory state in the meninges. Notably, mast cells can release IL12-p70 (Mukai et al., 2018) in response to CGRP-induced degranulation (Ottosson and Edvinsson, 1997; Theoharides et al., 2005). Our data showing decreased levels of CGRP in animals lacking mLVs may imply that the absence of a functional meningeal lymphatic system leads to a reduced tendency to develop a migraine through several downstream mechanisms controlled by this neuropeptide.

Apart from CGRP, many immune cells, including mast cells, are controlled by purinergic mechanisms mediated by extracellular ATP which is serving as a powerful trigger of inflammatory reactions acting primarily via P2X7 receptors (Burnstock, 2016; Karmakar et al., 2016; Nurkhametova et al., 2019, 2018). Consistent with this view, we also found that the stimulation of meninges with the P2X7 agonist BzATP stimulated the release of IL-10 and TNFα, demonstrating the pro-inflammatory action of purinergic signaling in meninges. We used BzATP as a stable ATP analogue, because of the long, 3.5 hour treatment time and because of fast degradation of ATP. Notably, ATP receptor stimulation resulted in the significant release of TNFα only in the meninges from WT animals, but not in the meninges lacking the mLVs. TNFα is the main migraine-related pro-inflammatory cytokine (Martami et al., 2018; Oliveira et al., 2017; Perini et al., 2005; Yücel et al., 2016). In line with this, TNFα has been reported as the pro-inflammatory agent, which can stimulate CGRP transcription (Bowen et al., 2006). The lack of increase in TNFα levels after ATP receptor stimulation in the meninges of the K14 mice lacking the lymphatic system is consistent with our other findings on the reduced level of CGRP in these animals K14. Taken together, the lack of meningeal lymphatic system may induce a local shift in the balance of various humoral factors (i.e., CGRP and pro-inflammatory cytokines such as TNFα), suggesting a reduction in the signalling of neurogenic inflammation.

4.3 Meningeal nociception and morphological properties of nerve terminals

Our recordings of nociceptive activity from trigeminal afferents demonstrated a significant increase in nociceptive spiking in response to the powerful trigger of meningeal nociception ATP in K14 mice. Notably, ATP can operate either directly via its own P2X3 receptors at meningeal nerve terminals (Yegutkin et al., 2016), or by degranulating mast cells and release of serotonin, which acts via it own 5-HT3 receptors (Koroleva et al., 2019). Therefeore, one possible explanation for higher spiking frequency in K14 mice could be the involvement of mast cells in ATP-induced nociceptive spiking. Another possibility is the more extensive innervation of meninges, which are heavily innervated by somatic and autonomous nerve terminals, respectively originating from the trigeminal ganglion

(Ramachandran, 2018; Zakharov et al., 2015) and sphenopalatine ganglion (Delépine and Aubineau, 1997). Sensory nerve fibers are characterized mainly by nociceptive Aδ and peptidergic C-fibers (Levy and Strassman, 2002), with the latter considered as the primary source of the key migraine mediator CGRP (Eftekhari et al., 2013). Thus, since CGRP release in meninges without mLVs was lower than in healthy meninges, we speculate that sensitization of dural afferents, which depends on the level of CGRP (Fabbretti et al., 2006), was lower in K14 animals compared to WT animals, leading to a less prominent difference in nociceptive activity.

To clarify if the presence of functional mLVs affects the innervation of dura mater, we analyzed several parameters of meningeal innervation. However, we did not find a difference in the total length of axonal fibers or in the number of terminal segments of meningeal afferents. This suggests that the higher activity of ATP or the lowered level of CGRP released in K14 mice are not due to a reduced number of meningeal innervation or a small number of nerve terminals but rather present the different functionality of local innervation.

4.4 Limitations of the model

Our data for the first time demonstrates the interactions between the meningeal lymphatic system, neurogenic inflammation, and nociception, which determine the predisposition to the generation of the migraine attack. However, the model used in our study has some limitations. First, all our experiments have been conducted in *ex vivo* conditions. Thus, the meningeal lymphatic system was not active in this preparations (e.g., recruitment and trafficking of immune cells and clearance of migraine mediators were not functioning in our samples). Second, our animals were not sensitized to mimic migraine attacks. Third, it has been previously reported that K14-VEGFR3-Ig mice have a defect in systemic immunity (Thomas et al., 2012), involving mainly the CD8+ T lymphocytes, which were not studied here. Third, to study alterations between WT and K14 meninges we used ATP, which is only one of the numerous pro-inflammatory compounds released in meninges. Thus, the impact of the lymphatic system in migraine state in meninges requires further tests with injections of nitroglycerin (Demartini et al., 2019) or induction of CSD representing a model for migraine with aura (Van Den Maagdenberg et al., 2004) and considering other immune cells, such as T cells, known to be involved in migraine (Nurkhametova et al., 2018).

4.5 Pathophysiological implications

The discovery of meningeal lymphatic system revealed new aspects for the functioning of the healthy brain and proposed new mechanisms of several brain diseases such as stroke, epilepsy, TBI, Alzheimer's Disease (Kwon et al., 2019; Noé and Marchi, 2019; Patel et al., 2019; Rasmussen et al.,

2018; Tamura et al., 2019). Here we propose a possible contribution of mLVs also in migraine pathology, given the generally accepted central role of the meningeal tissues in this disorder. Our main working hypothesis was that the lack of the meningeal lymphatic system exaggerates the neurogenic inflammation, leading to more severe symptoms of migraines because such animals are not able to clean up the dural environment. Indeed, we found that the lack of the meningeal lymphatic system produced several pro-nociceptive changes in the trigeminovascular system relevant to migraine pathology. This resulted in the increased nociceptive activity and enhanced expression of the pro-inflammatory MCP-1 cytokine. Yet surprisingly, the expression of key migraine mediator CGRP and other pro-inflammatory TNFα and IL-12p70 cytokines was decreased. This suggest that the absence of meningeal lymphatic system caused not uniform, but rather more complex, and eventually functionally opposite, changes in the nociceptive and inflammatory balance in the meninges.

Our findings highlight the importance of the meningeal lymphatic drainage in the pathophysiology of migraine, suggesting it as a potential target in the search for new therapeutic anti-migraine strategies, or in the re-evaluation of the existing ones.

5. Acknowledgements

This study has been supported by the Academy of Finland (Grant 277442 – NM, RG, RashidG; Academy of Finland research Fellowship #309479/2017 – FMN), Russian Foundation for Basic Research (KOMFI Grant 17-00-00053 – KK, OG) and by the Doctoral Program in Molecular Medicine of University of Eastern Finalnd (NM).

For help with cytokine release assay and staining of meninges for β-tubulin III, we would like to thank Sara Wojciechowski, Dilyara Nurkhametova, Sanna Loppi and Maria Vihma.

6. Author Contributions

NM, FMN and RashidG originally conceived the study; NM, FMN, RashidG elaborated the study design; NM, KK, RG performed experiments; NM, AA, OG, AS, JT analyzed the data; NM, AA, OG, JT performed the statistical analysis; NM, AA, AS, FNM, RashidG drafted a significant portion of the manuscript and figures. TM supported the study and edited the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

7. Conflict of Interest Statement

None of the authors has any conflict of interest to disclose. The authors confirm that have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

8. Data Availability

The raw data supporting the conclusions of this manuscript will be made available by the corresponding author, upon reasonable request, to any qualified researcher.

9. References

Abdollahzadeh, A., Belevich, I., Jokitalo, E., Tohka, J., Sierra, A., 2019a. Automated 3D Axonal Morphometry of White Matter. Sci. Rep. 9, 1–16. https://doi.org/10.1038/s41598-019-42648-2

Abdollahzadeh, A., Sierra, A., Tohka, J., 2019b. Cylindrical shape decomposition for 3D segmentation of tubular objects. arXiv:1911.00571v2.

Ahn, J.H., Cho, H., Kim, J.H., Kim, S.H., Ham, J.S., Park, I., Suh, S.H., Hong, S.P., Song, J.H., Hong, Y.K., Jeong, Y., Park, S.H., Koh, G.Y., 2019. Meningeal lymphatic vessels at the skull base drain cerebrospinal fluid. Nature 572, 62–66. https://doi.org/10.1038/s41586-019-1419-5

Alitalo, K., 2011. The lymphatic vasculature in disease. Nat. Med. https://doi.org/10.1038/nm.2545

Antila, S., Karaman, S., Nurmi, H., Airavaara, M., Voutilainen, M.H., Mathivet, T., Chilov, D., Li, Z., Koppinen, T., Park, J.H., Fang, S., Aspelund, A., Saarma, M., Eichmann, A., Thomas, J.L., Alitalo, K., 2017. Development and plasticity of meningeal lymphatic vessels. J. Exp. Med. 214, 3645–3667. https://doi.org/10.1084/jem.20170391

Aspelund, A., Antila, S., Proulx, S.T., Karlsen, T.V., Karaman, S., Detmar, M., Wiig, H., Alitalo, K., 2015. A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. J. Exp. Med. 212, 991–9. https://doi.org/10.1084/jem.20142290

Baraldi, P., Virgilio, F., Romagnoli, R., 2005. Agonists and Antagonists Acting at P2X7 Receptor. Curr. Top. Med. Chem. 4, 1707–1717. https://doi.org/10.2174/1568026043387223

Benmansour, F., Cohen, L.D., 2011. Tubular structure segmentation based on minimal path method and anisotropic enhancement. Int. J. Comput. Vis. 92, 192–210. https://doi.org/10.1007/s11263-010-0331-0

Bø, S.H., Davidsen, E.M., Gulbrandsen, P., Dietrichs, E., Bovim, G., Stovner, L.J., White, L.R., 2009. Cerebrospinal fluid cytokine levels in migraine, tension-type headache and cervicogenic headache. Cephalalgia 29, 365–372. https://doi.org/10.1111/j.1468-2982.2008.01727.x

Bolay, H., Reuter, U., Dunn, A.K., Huang, Z., Boas, D.A., Moskowitz, M.A., 2002. Intrinsic brain activity triggers trigeminal meningeal afferents in a migraine model. Nat. Med. 8, 136–142. https://doi.org/10.1038/nm0202-136

Bowen, E.J., Schmidt, T.W., Firm, C.S., Russo, A.F., Durham, P.L., 2006. Tumor necrosis factor-α stimulation of calcitonin gene-related peptide expression and secretion from rat trigeminal ganglion neurons. J. Neurochem. 96, 65–77. https://doi.org/10.1111/j.1471-4159.2005.03524.x

Burnstock, G., 2016. P2X ion channel receptors and inflammation. Purinergic Signal. 12, 59–67. https://doi.org/10.1007/s11302-015-9493-0

Cáceres, A., Banker, G.A., Binder, L., 1986. Immunocytochemical localization of tubulin and microtubule-associated protein 2 during the development of hippocampal neurons in culture. J. Neurosci. 6, 714–22.

Cady, R.J., Glenn, J.R., Smith, K.M., Durham, P.L., 2011. Calcitonin gene-related peptide promotes cellular changes in trigeminal neurons and glia implicated in peripheral and central sensitization. Mol. Pain 7, 94. https://doi.org/10.1186/1744-8069-7-94

Chan, T.F., Vese, L.A., 2001. Active contours without edges. IEEE Trans. Image Process. 10, 266–277. https://doi.org/10.1109/83.902291

Delépine, L., Aubineau, P., 1997. Plasma Protein Extravasation Induced in the Rat Dura Mater by Stimulation of the Parasympathetic Sphenopalatine Ganglion. Exp. Neurol. 147, 389–400. https://doi.org/10.1006/exnr.1997.6614

Demartini, C., Greco, R., Zanaboni, A.M., Sances, G., De Icco, R., Borsook, D., Tassorelli, C., 2019. Nitroglycerin as a comparative experimental model of migraine pain: From animal to human and back. Prog. Neurobiol. https://doi.org/10.1016/j.pneurobio.2019.02.002

Ebersberger, A., Takac, H., Richter, F., Schaible, H.-G., 2006. Effect of sympathetic and parasympathetic mediators on the release of calcitonin gene-related peptide and prostaglandin E from rat dura mater, in vitro. Cephalalgia 26, 282–289. https://doi.org/10.1111/j.1468-2982.2005.01035.x

Eftekhari, S., Warfvinge, K., Blixt, F.W., Edvinsson, L., 2013. Differentiation of nerve fibers storing CGRP and CGRP receptors in the peripheral trigeminovascular system. J. Pain 14, 1289–1303. https://doi.org/10.1016/j.jpain.2013.03.010

Fabbretti, E., D'Arco, M., Fabbro, A., Simonetti, M., Nistri, A., Giniatullin, R., 2006. Delayed Upregulation of ATP P2X3 Receptors of Trigeminal Sensory Neurons by Calcitonin Gene-

Related Peptide. J. Neurosci. 26, 6163–6171. https://doi.org/10.1523/JNEUROSCI.0647-06.2006

Fidan, I., Yüksel, S., Ýmir, T., Irkeç, C., Aksakal, F.N., 2006. The importance of cytokines, chemokines and nitric oxide in pathophysiology of migraine. J. Neuroimmunol. 171, 184–188. https://doi.org/10.1016/j.jneuroim.2005.10.005

Frangi, A.F., Niessen, W.J., Vincken, K.L., Viergever, M.A., 1998. Multiscale vessel enhancement filtering, in: Lecture Notes in Computer Science (Including Subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics). Springer Verlag, pp. 130–137. https://doi.org/10.1007/bfb0056195

Giniatullin, R., Gupta, K., Theoharides, T., 2019. Editorial: Mast Cells in Itch, Pain and Neuro-Inflammation. Front. Cell. Neurosci. 13, 521. https://doi.org/10.3389/fncel.2019.00521

Giniatullin, R., Nistri, A., Fabbretti, E., 2008. Molecular mechanisms of sensitization of pain-transducing P2X3 receptors by the migraine mediators CGRP and NGF. Mol. Neurobiol. https://doi.org/10.1007/s12035-008-8020-5

Goadsby, P.J., Edvinsson, L., Ekman, R., 1988. Release of vasoactive peptides in the extracerebral circulation of humans and the cat during activation of the trigeminovascular system. Ann. Neurol. 23, 193–196. https://doi.org/10.1002/ana.410230214

Hayday, A., Theodoridis, E., Ramsburg, E., Shires, J., 2001. Intraepithelial lymphocytes: Exploring the Third Way in immunology. Nat. Immunol. https://doi.org/10.1038/ni1101-997

Karmakar, M., Katsnelson, M.A., Dubyak, G.R., Pearlman, E., 2016. Neutrophil P2X7 receptors mediate NLRP3 inflammasome-dependent IL-1β secretion in response to ATP. Nat. Commun. 7, 10555. https://doi.org/10.1038/ncomms10555

Keller, J.T., Marfurt, C.F., 1991. Peptidergic and serotoninergic innervation of the rat dura mater. J. Comp. Neurol. 309, 515–534. https://doi.org/10.1002/cne.903090408

Khaiboullina, S.F., Mendelevich, E.G., Shigapova, L.H., Shagimardanova, E., Gazizova, G., Nikitin, A., Martynova, E., Davidyuk, Y.N., Bogdanov, E.I., Gusev, O., Van Den Maagdenberg, A.M.J.M., Giniatullin, R.A., Rizvanov, A.A., 2017. Cerebellar atrophy and changes in cytokines associated with the CACNA1A R583Q mutation in a Russian familial

hemiplegic migraine type 1 family. Front. Cell. Neurosci. 11, 263.
https://doi.org/10.3389/fncel.2017.00263

Kilinc, E., Guerrero-Toro, C., Zakharov, A., Vitale, C., Gubert-Olive, M., Koroleva, K., Timonina, A., Luz, L.L., Shelukhina, I., Giniatullina, R., Tore, F., Safronov, B. V., Giniatullin, R., 2017. Serotonergic mechanisms of trigeminal meningeal nociception: Implications for migraine pain. Neuropharmacology 116, 160–173. https://doi.org/10.1016/j.neuropharm.2016.12.024

Koroleva, K., Gafurov, O., Guselnikova, V., Nurkhametova, D., Giniatullina, R., Sitdikova, G., Mattila, O.S., Lindsberg, P.J., Malm, T.M., Giniatullin, R., 2019. Meningeal mast cells contribute to ATP-induced nociceptive firing in trigeminal nerve terminals: Direct and indirect purinergic mechanisms triggering migraine pain. Front. Cell. Neurosci. 13. https://doi.org/10.3389/fncel.2019.00195

Koyuncu Irmak, D., Kilinc, E., Tore, F., 2019. Shared fate of meningeal mast cells and sensory neurons in migraine. Front. Cell. Neurosci. 13. https://doi.org/10.3389/fncel.2019.00136

Kwon, S., Moreno-Gonzalez, I., Taylor-Presse, K., Edwards, G., Gamez, N., Calderon, O., Zhu, B., Velasquez, F.C., Soto, C., Sevick-Muraca, E.M., 2019. Impaired Peripheral Lymphatic Function and Cerebrospinal Fluid Outflow in a Mouse Model of Alzheimer's Disease. J. Alzheimer's Dis. 69, 585–593. https://doi.org/10.3233/JAD-190013

Levy, D., 2009. Migraine pain, meningeal inflammation, and mast cells. Curr. Pain Headache Rep. 13, 237–40.

Levy, D., Strassman, A.M., 2002. Mechanical response properties of A and C primary afferent neurons innervating the rat intracranial dura. J Neurophysiol 88, 3021–3031. https://doi.org/10.1152/jn.00029.2002

Loberg, R.D., Day, L.S.L., Harwood, J., Ying, C., St. John, L.N., Giles, R., Neeley, C.K., Pienta, K.J., 2006. CCL2 is a potent regulator of prostate cancer cell migration and proliferation. Neoplasia 8, 578–586. https://doi.org/10.1593/neo.06280

Lopes, J.P., Stylianou, M., Backman, E., Holmberg, S., Ekoff, M., Nilsson, G., Urban, C.F., 2019. Cryptococcus neoformans Induces MCP-1 Release and Delays the Death of Human Mast Cells. Front. Cell. Infect. Microbiol. 9. https://doi.org/10.3389/fcimb.2019.00289

Louveau, A., Smirnov, I., Keyes, T.J., Eccles, J.D., Rouhani, S.J., Peske, J.D., Derecki, N.C., Castle, D., Mandell, J.W., Lee, K.S., Harris, T.H., Kipnis, J., 2015. Structural and functional features of central nervous system lymphatic vessels. Nature 523, 337–341. https://doi.org/10.1038/nature14432

Lv, J., Huang, Y., Zhu, S., Yang, G., Zhang, Y., Leng, J., Bo, J., Liu, D., 2012. MCP-1-Induced Histamine Release from Mast Cells Is Associated with Development of Interstitial Cystitis/Bladder Pain Syndrome in Rat Models. Mediators Inflamm. 2012, 1–9. https://doi.org/10.1155/2012/358184

Mäkinen, T., Jussila, L., Veikkola, T., Karpanen, T., Kettunen, M.I., Pulkkanen, K.J., Kauppinen, R., Jackson, D.G., Kubo, H., Nishikawa, S.I., Ylä-Herttuala, S., Alitalo, K., 2001. Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble VEGF receptor-3. Nat. Med. 7, 199–205. https://doi.org/10.1038/84651

Martami, F., Razeghi Jahromi, S., Togha, M., Ghorbani, Z., Seifishahpar, M., Saidpour, A., 2018. The serum level of inflammatory markers in chronic and episodic migraine: a case-control study. Neurol. Sci. 39, 1741–1749. https://doi.org/10.1007/s10072-018-3493-0

Martins, T.B., Rose, J.W., Jaskowski, T.D., Wilson, A.R., Husebye, D., Seraj, H.S., Hill, H.R., 2011. Analysis of proinflammatory and anti-inflammatory cytokine serum concentrations in patients with multiple sclerosis by using a multiplexed immunoassay. Am. J. Clin. Pathol. 136, 696–704. https://doi.org/10.1309/AJCP7UBK8IBVMVNR

Mason, R.T., Peterfreund, R.A., Sawchenko, P.E., Corrigan, A.Z., Rivier, J.E., Vale, W.W., 1984. Release of the predicted calcitonin gene-related peptide from cultured rat trigeminal ganglion cells. Nature 308, 653–5.

Matsui, S., Tanaka, M., Kamiyoshi, A., Sakurai, T., Ichikawa-Shindo, Y., Kawate, H., Dai, K., Cui, N., Wei, Y., Tanaka, M., Kakihara, S., Nakamura, K., Yamauchi, A., Ishida, K., Tanaka, S., Kawamata, M., Shindo, T., 2019. Endogenous Calcitonin Gene–Related Peptide Deficiency Exacerbates Postoperative Lymphedema by Suppressing Lymphatic Capillary Formation and M2 Macrophage Accumulation. Am. J. Pathol. 189, 2487–2502. https://doi.org/10.1016/j.ajpath.2019.08.011

McIlvried, L.A., Cruz, J.A., Borghesi, L.A., Gold, M.S., 2017. Sex-, stress-, and sympathetic post-ganglionic-dependent changes in identity and proportions of immune cells in the dura.

Cephalalgia 37, 36–48. https://doi.org/10.1177/0333102416637832

Mikhailov, N., Leskinen, J., Fagerlund, I., Poguzhelskaya, E., Giniatullina, R., Gafurov, O., Malm, T., Karjalainen, T., Gröhn, O., Giniatullin, R., 2019. Mechanosensitive meningeal nociception via Piezo channels: Implications for pulsatile pain in migraine? Neuropharmacology 149, 113–123. https://doi.org/10.1016/J.NEUROPHARM.2019.02.015

Moskowitz, M.A., 1993. Neurogenic inflammation in the pathophysiology and treatment of migraine. Neurology 43, S16–S20.

Moskowitz, M.A., 1984. The neurobiology of vascular head pain. Ann. Neurol. 16, 157–168. https://doi.org/10.1002/ana.410160202

Moskowitz, M.A., Romero, J., Reinhard, J.F., Melamed, E., Pettibone, D.J., 1979. NEUROTRANSMITTERS AND THE FIFTH CRANIAL NERVE: IS THERE A RELATION TO THE HEADACHE PHASE OF MIGRAINE? Lancet 314, 883–885. https://doi.org/10.1016/S0140-6736(79)92692-8

Mukai, K., Tsai, M., Saito, H., Galli, S.J., 2018. Mast cells as sources of cytokines, chemokines, and growth factors. Immunol. Rev. https://doi.org/10.1111/imr.12634

Müller-Berghaus, J., Kern, K., Paschen, A., Nguyen, X.D., Klüter, H., Morahan, G., Schadendorf, D., 2004. Deficient IL-12p70 secretion by dendritic cells based on IL12B promoter genotype. Genes Immun. 5, 431–434. https://doi.org/10.1038/sj.gene.6364102

Noé, F.M., Marchi, N., 2019. Central nervous system lymphatic unit, immunity, and epilepsy: Is there a link? Epilepsia Open. https://doi.org/10.1002/epi4.12302

Nurkhametova, D., Kudryavtsev, I., Guselnikova, V., Serebryakova, M., Giniatullina, R.R., Wojciechowski, S., Tore, F., Rizvanov, A., Koistinaho, J., Malm, T., Giniatullin, R., 2019. Activation of P2X7 receptors in peritoneal and meningeal mast cells detected by uptake of organic dyes: Possible purinergic triggers of neuroinflammation in meninges. Front. Cell. Neurosci. 13. https://doi.org/10.3389/fncel.2019.00045

Nurkhametova, D., Kudryavtsev, I., Khayrutdinova, O., Serebryakova, M., Altunbaev, R., Malm, T., Giniatullin, R., 2018. Purinergic profiling of regulatory T-cells in patients with episodic migraine. Front. Cell. Neurosci. 12. https://doi.org/10.3389/fncel.2018.00326

Olesen, J., Burstein, R., Ashina, M., Tfelt-Hansen, P., 2009. Origin of pain in migraine: evidence for peripheral sensitisation. Lancet Neurol. https://doi.org/10.1016/S1474-4422(09)70090-0

Oliveira, A.B., Bachi, A.L.L., Ribeiro, R.T., Mello, M.T., Tufik, S., Peres, M.F.P., 2017. Unbalanced plasma TNF-α and IL-12/IL-10 profile in women with migraine is associated with psychological and physiological outcomes. J. Neuroimmunol. 313, 138–144. https://doi.org/10.1016/j.jneuroim.2017.09.008

Ottosson, A., Edvinsson, L., 1997. Release of histamine from dural mast cells by substance P and calcitonin gene-related peptide. Cephalalgia 17, 166–174. https://doi.org/10.1046/j.1468-2982.1997.1703166.x

Patel, T.K., Habimana-Griffin, L., Gao, X., Xu, B., Achilefu, S., Alitalo, K., McKee, C.A., Sheehan, P.W., Musiek, E.S., Xiong, C., Coble, D., Holtzman, D.M., 2019. Dural lymphatics regulate clearance of extracellular tau from the CNS. Mol. Neurodegener. 14. https://doi.org/10.1186/s13024-019-0312-x

Perini, F., D'Andrea, G., Galloni, E., Pignatelli, F., Billo, G., Alba, S., Bussone, G., Toso, V., 2005. Plasma cytokine levels in migraineurs and controls. Headache 45, 926–931. https://doi.org/10.1111/j.1526-4610.2005.05135.x

Piantino, J., Lim, M.M., Newgard, C.D., Iliff, J., 2019. Linking Traumatic Brain Injury, Sleep Disruption and Post-Traumatic Headache: a Potential Role for Glymphatic Pathway Dysfunction. Curr. Pain Headache Rep. https://doi.org/10.1007/s11916-019-0799-4

Pietrobon, D., Moskowitz, M.A., 2013. Pathophysiology of Migraine. Annu. Rev. Physiol. 75, 365–391. https://doi.org/10.1146/annurev-physiol-030212-183717

Ramachandran, R., 2018. Neurogenic inflammation and its role in migraine. Semin. Immunopathol. 40, 301–314. https://doi.org/10.1007/s00281-018-0676-y

Rasmussen, M.K., Mestre, H., Nedergaard, M., 2018. The glymphatic pathway in neurological disorders. Lancet Neurol. https://doi.org/10.1016/S1474-4422(18)30318-1

Sarchielli, P., Alberti, A., Vaianella, L., Pierguidi, L., Floridi, A., Mazzotta, G., Floridi, A., Gallai, V., 2004. Chemokine levels in the jugular venous blood of migraine without aura patients during attacks. Headache 44, 961–968. https://doi.org/10.1111/j.1526-4610.2004.04189.x

Sethian, J.A., 1996. A fast marching level set method for monotonically advancing fronts. Proc. Natl. Acad. Sci. U. S. A. 93, 1591–1595. https://doi.org/10.1073/pnas.93.4.1591

Shatillo, A., Koroleva, K., Giniatullina, R., Naumenko, N., Slastnikova, A.A., Aliev, R.R., Bart, G., Atalay, M., Gu, C., Khazipov, R., Davletov, B., Grohn, O., Giniatullin, R., 2013. Cortical spreading depression induces oxidative stress in the trigeminal nociceptive system. Neuroscience 253, 341–349. https://doi.org/10.1016/j.neuroscience.2013.09.002

Shelukhina, I., Mikhailov, N., Abushik, P., Nurullin, L., Nikolsky, E.E., Giniatullin, R., 2017. Cholinergic Nociceptive Mechanisms in Rat Meninges and Trigeminal Ganglia: Potential Implications for Migraine Pain. Front. Neurol. 8, 163. https://doi.org/10.3389/fneur.2017.00163

Suleimanova, A., Talanov, M., Gafurov, O., Gafarov, F., Koroleva, K., Virenque, A., Noe, F.M., Mikhailov, N., Nistri, A., Giniatullin, R., 2020. Modeling a Nociceptive Neuro-Immune Synapse Activated by ATP And 5-HT in Meninges: Novel Clues On Transduction of Chemical Signals into Persistent or Rhythmic Neuronal Firing. Front. Cell. Neurosci.

Takenaka, M.C., Robson, S., Quintana, F.J., 2016. Regulation of the T Cell Response by CD39. Trends Immunol. https://doi.org/10.1016/j.it.2016.04.009

Tamura, R., Yoshida, K., Toda, M., 2019. Current understanding of lymphatic vessels in the central nervous system. Neurosurg. Rev. https://doi.org/10.1007/s10143-019-01133-0

Theoharides, T.C., Donelan, J., Kandere-Grzybowska, K., Konstantinidou, A., 2005. The role of mast cells in migraine pathophysiology. Brain Res. Rev. https://doi.org/10.1016/j.brainresrev.2004.11.006

Thomas, S.N., Rutkowski, J.M., Pasquier, M., Kuan, E.L., Alitalo, K., Randolph, G.J., Swartz, M.A., 2012. Impaired Humoral Immunity and Tolerance in K14-VEGFR-3-Ig Mice That Lack Dermal Lymphatic Drainage . J. Immunol. 189, 2181–2190. https://doi.org/10.4049/jimmunol.1103545

Van Den Maagdenberg, A.M.J.M., Pietrobon, D., Pizzorusso, T., Kaja, S., Broos, L.A.M., Cesetti, T., Van De Ven, R.C.G., Tottene, A., Van Der Kaa, J., Plomp, J.J., Frants, R.R., Ferrari, M.D., 2004. A Cacna1a knockin migraine mouse model with increased susceptibility to cortical spreading depression. Neuron 41, 701–710. https://doi.org/10.1016/S0896-6273(04)00085-6

Wood, S., Jayaraman, V., Huelsmann, E.J., Bonish, B., Burgad, D., Sivaramakrishnan, G., Qin, S., DiPietro, L.A., Zloza, A., Zhang, C., Shafikhani, S.H., 2014. Pro-inflammatory chemokine CCL2 (MCP-1) promotes healing in diabetic wounds by restoring the macrophage response. PLoS One 9, e91574. https://doi.org/10.1371/journal.pone.0091574

Yegutkin, G.G., Guerrero-Toro, C., Kilinc, E., Koroleva, K., Ishchenko, Y., Abushik, P., Giniatullina, R., Fayuk, D., Giniatullin, R., 2016. Nucleotide homeostasis and purinergic nociceptive signaling in rat meninges in migraine-like conditions. Purinergic Signal. 12, 561–574. https://doi.org/10.1007/s11302-016-9521-8

Yücel, M., Kotan, D., Çiftçi, G.G., Çiftçi, I.H., Cikriklar, H.I., 2016. Serum levels of endocan, claudin-5 and cytokines in migraine. Eur. Rev. Med. Pharmacol. Sci. 20, 930–936.

Zakharov, A., Koroleva, K., Giniatullin, R., 2016. Clustering Analysis for Sorting ATP-Induced Nociceptive Firing in rat Meninges. Bionanoscience 6, 508–512. https://doi.org/10.1007/s12668-016-0276-z

Zakharov, A., Vitale, C., Kilinc, E., Koroleva, K., Fayuk, D., Shelukhina, I., Naumenko, N., Skorinkin, A., Khazipov, R., Giniatullin, R., 2015. Hunting for origins of migraine pain: cluster analysis of spontaneous and capsaicin-induced firing in meningeal trigeminal nerve fibers. Front. Cell. Neurosci. 9, 287. https://doi.org/10.3389/fncel.2015.00287