Title:

Novel class of OTU deubiquitinases regulate substrate ubiquitination
upon *Legionella* infection

Authors: Donghyuk Shin[1,2,3], Anshu Bhattacharya[1,2], Yi-Lin Cheng[1,2], Marta Campos Alonso[2],

Ahmad Reza Mehdipour[3], Gerbrand J. van der Heden van Noort[4], Huib Ovaa[4], Gerhard Hummer[3,5]

and Ivan Dikic[1,2,3, *]

Affiliations:

[1] Institute of Biochemistry II, Faculty of Medicine, Goethe University Frankfurt, Theodor-Stern-

Kai 7, 60590 Frankfurt am Main, Germany.

[2] Buchmann Institute for Molecular Life Sciences, Goethe University Frankfurt, Max-von-Laue-

Str. 15, 60438 Frankfurt am Main, Germany.

[3] Max Planck Institute of Biophysics, Max-von-Laue-Str. 3, 60438 Frankfurt am Main, Germany

[4] Oncode Institute and Department of Chemical Immunology, Leiden University Medical Centre,

Einthovenweg 20, 2333 ZC, Leiden, The Netherlands

[5] Institute of Biophysics, Goethe University Frankfurt, Frankfurt am Main, Germany.

*Correspondence & Lead contact: dikic@biochem2.uni-frankfurt.de

Abstract: *Legionella pneumophila* is a gram-negative pathogenic bacterium that causes

Legionaries' disease. The *Legionella* genome codes more than 300 effector proteins able to

modulate host-pathogen interactions during infection. Among them are also enzymes altering the

host-ubiquitination system including bacterial ligases and deubiquitinases. In this study, based on

homology-detection screening on 305 *Legionella* effector proteins, we identified two ***L**egionella*

OTU-like deubiquitinases (**LOT**; LotB (Lpg1621/Ceg23) and LotC (Lpg2529), LotA

(Lpg2248/Lem21) is already known). A crystal structure of LotC catalytic core (LotC$_{14-310}$) was

determined at 2.4 Å and compared with other OTU deubiquitinases, including LotB. Unlike the

classical OTU-family, the structures of *Legionella* OTU-family (LotB and LotC) shows an

extended helical lobe between the Cys-loop and the variable loop, which define a novel class of

OTU-deubiquitinase. Despite structural differences in their helical lobes, both LotB and LotC

interact with ubiquitin. LotB has an additional ubiquitin binding site (S1') enabling specific

cleavage of Lys63-linked poly-ubiquitin chains. By contrast, LotC only contains the S1 site and

cleaves different species of ubiquitin chains. MS analysis of catalytically inactive LotB and LotC

identified different categories of host-substrates for these two related DUBs. Together, our results

provide new structural insights of bacterial OTU deubiquitinases and indicate distinct roles of

bacterial deubiquitinases in host-pathogen interactions.

Keywords: bacterial deubiquitinase, *Legionella pneumophila,* effector proteins, OTU-

deubiquitinases, ubiquitin

INTRODUCTION

Ubiquitination, a well-studied post-translational modification system, regulates the fate of various substrates by tagging them with ubiquitin (Yau and Rape, 2016). A representative example of ubiquitin-mediated cellular process is the ubiquitin-proteasome system (UPS), where misfolded proteins get ubiquitinated and degraded by the proteasome, and recycled (Dikic, 2017). For the larger cellular wastes, such as cellular components (ER, mitochondria etc.), protein aggregates or intracellular bacteria, ubiquitination works together with the autophagy machinery, which includes sequestration of ubiquitinated components and their transfer into the lysosome for degradation (Pohl and Dikic, 2019). To maintain homeostasis in the cell, ubiquitination events are tightly regulated by a reverse process called deubiquitination, where ubiquitin molecules from target substrates are specifically cleaved and recycled by deubiquitinating enzymes (deubiquitinases; DUBs) (Clague et al., 2019).

To date, about 100 different deubiquitinases (DUBs) have been identified from human. They are categorized into seven different classes, based on their structure and mechanism of actions including USP, JAMM (MPN), OTU, MJD (Josepin), UCH and the recently discovered MINDY and ZUFSP (Abdul Rehman et al., 2016; Clague et al., 2019; Haahr et al., 2018; Hermanns et al., 2018; Hewings et al., 2018; Kwasna et al., 2018). Six of them belong to the cysteine protease family (USP, OTU, MJD(Josepin), UCH, MINDY and ZUFSP), while JAMM (MPN) belongs to the zinc-containing metalloproteases. Among them, the OTU-family is distinguished from other DUBs, as they exhibit linkage specificity (Mevissen et al., 2016, 2013; Mevissen and Komander, 2017). For example, Cezanne specifically cleaves Lys11-linked polyubiquitin chains (Bremm et al., 2010) and OTUB1 preferentially cleaves Lys48-linked chains (Edelmann et al., 2009; Wang et al., 2009), while OTULIN exclusively cleaves M1-linked (linear)

chains (Keusekotten et al., 2013). Extensive biochemical and structural studies have provided

general mechanisms of how the diverse linkage specificity can be achieved within the structurally

similar OTU-family, pointing to the presence or absence of additional ubiquitin binding domains

(UBDs), sequence variations on ubiquitinated substrates, or additional S1'/S2 binding sites

(Mevissen et al., 2013).

Considering the importance of ubiquitin-mediated cellular pathways, it is not surprising

that pathogens are armed with various weapons to hijack the host-ubiquitination system. For

instance, *Salmonella typhimurium* encodes HECT type E3 ligase SopA (Diao et al., 2008; Fiskin

et al., 2017; Lin et al., 2012) and *Legionella pneumophila* contains LubX and LegU1, which are

similar to U-box and F-box containing E3 ligases, respectively (Ensminger and Isberg, 2010;

Kubori et al., 2008; Quaile et al., 2015). In addition, bacterial pathogens also possess atypical

ubiquitin ligases that do not belong to any of known E3 ligases, such as IpaH family (*Shigella*) or

SidC/SdCA (*Legionella*) (Hsu et al., 2014; Wasilko et al., 2018). More recently, the SidE family

(SdeA, SdeB, SdeC and SidC) of *Legionella* has been shown to mediate unconventional

phosphoribosyl (PR) serine ubiquitination which is also tightly regulated by the meta-effector SidJ

or PR-ubiquitin specific deubiquitinases (DupA and DupB) (Bhogaraju et al., 2019, 2016; Black

et al., 2019; Kalayil et al., 2018; Qiu et al., 2016; Shin et al., 2020). Pathogenic bacteria encode

not only ubiquitin ligases, but also deubiquitinases (Hermanns and Hofmann, 2019). The most

studied bacterial deubiquitinases are CE-clan proteases, based on the MEROPS database

classification, which cleave either ubiquitin or ubiquitin-like modifiers (SUMO1 or NEDD8)

(Pruneda et al., 2016; Rawlings et al., 2018). In addition to CE-clan DUBs, bacteria and viruses

also encode OTU-like deubiquitinases. Several structures from viral-OTUs revealed that they have

a unique structure compared to those of known OTU family members (Akutsu et al., 2011;

Capodagli et al., 2013; James et al., 2011; Lombardi et al., 2013; van Kasteren et al., 2013). OTUs

from nairovirus Crimean-Congo hemorrhagic fever virus (CCHFV) and Dugbe virus (DUGV)

have an extra β-hairpin in their S1 binding site. Papain-like protease (PLP2) domain of the

artervirus equine arteritis virus (EAV) contains a zinc-finger that also interacts with ubiquitin.

While viral OTUs have been studied, to date only three bacterial OTU-like DUBs have been

identified. ChlaOTU from *Chlamydophila pneumoniae* contains an OTU-domain that cleaves both

K48- and K63-linked polyubiquitin chains (Furtado et al., 2013). LotA (Lpg2248;Lem21),

_L_egionella OTU-like deubiquitinase, contains two OTU-like domains with two catalytic Cys

residues (C13, C303), where both are required for cleaving ubiquitin chains from an LCV

(*Legionella*-containing vacuole) (Kubori et al., 2018). Interestingly, LotA showed K6-linkage

preference that is solely dependent on its first OTU domain (Cys13). Recently, another OTU-like

deubiquitinase from *Legionella* (Lpg1621, Ceg23) has been identified as K63 chain-specific OTU-

deubiquitinase (Ma et al., 2020).

Despite these findings, little is known about the structure and molecular details of bacterial

OTU-like deubiquitinases. Here, we describe the two novel OTU-like deubiquitinaseses in

Legionella (LotB (Lpg1621; Ceg23) and LotC (Lpg2529). Structural analysis on LOT-

deubiqutinases provides insights on how bacterial OTU-deubiquitnases distinguished from the

known OTU members. Furthermore, we also identified the specific host-substrates of LotB and

LotC by mass-spectrometry (MS) analysis using catalytically inactive variants. Collectively, our

findings provide valuable structural insights into bacterial deubiquitinases and their roles in host-

pathogen interactions.

RESULTS

Identification of two novel OTU-like deubiquitinases from *Legionella* effector proteins

To identify putative deubiquitinases amongst *Legionella* effector proteins, we analyzed effector

proteins from *L. pneumophlia* (Lpg genes). Based on the type IV Icm/Dot complex secretion signal

(> 2.0), 305 effector proteins were selected (Burstein et al., 2016). Using pairwise sequence-

structure comparison based on hidden Markov models (HMMs, HHpred suite) (Zimmermann et

al., 2018), revealed four previously uncharacterized proteins as putative deubiquitinases. These

proteins all contain catalytic domains of known deubiquitinases (Fig. 1a). Lpg1621(Ceg23) and

Lpg2529 are found as members of OTU-family, whereas Lpg2411 and Lpg2907 belong to the

UCH and CE-clan, respectively (Fig. 1b, Table 1). An *in vitro* di-ubiquitin cleavage assay with di-

Ub panel (eight different linkage-specific Ub_2 chains(El Oualid et al., 2010)) showed that the OTU-

like DUBs (Lpg1621 and Lpg2529) are capable of cleaving Ub_2 chains with different specificity,

while other candidates (Lpg2411 and Lpg2907) did not show catalytic activity (Supplementary

Fig. 1a). The OTU family deubiquitinases have been shown to have linkage specificity against

certain polyubiquitin chains (Mevissen et al., 2016, 2013). To address whether Lpg1621 and

Lpg2529 follow this fundamental rule, we performed a time-course *in vitro* DUB assay with di-

Ub panel (Fig. 1c-f). Consistent with the recent evidence, Lpg1621 exclusively processed the K63-

linked Ub_2 (Ma et al., 2020), while Lpg2529 showed activity against K6-, K11-, K33-, K48- and

K63- linked Ub_2. Based on the sequence homology and catalytic activity, we now re-named the

Lpg1621 and Lpg2529 as <u>*Legionella*</u> <u>**OT**</u>U-like deubiquitinases (LotB and LotC, respectively).

Biochemical properties of LotB and LotC

The OTU-family belongs to the cysteine protease family, requiring the presence of a catalytic triad

for their activity (Mevissen et al., 2013). Based on the sequence analysis, we identifed the

conserved catalytic triad for both LotB (D27, C29 and H270) and LotC (D17, C24 and H304).

Mutations on either cysteine or histidine completely abolished the catalytic activity of both DUBs,

which suggests that both LotB and LotC follow the general catalytic mechanism of the OTU-

family (Fig. 2a, c). Next, we sought to find whether LotB and LotC require additional ubiquitin

binding sites (S1' or S2). To elicit this information, we used two different types of ubiquitin

activity-based probes (ABP). The propargyl-ubiquitin-ABP (Prg-ABP) contains a highly reactive

propargyl group at the C-terminus of ubiquitin chains, which can form a covalent bond with the

catalytic cysteine (Ekkebus et al., 2013; Flierman et al., 2016; Sommer et al., 2013). The

vinylmethylester-ubiquitin-ABP (VME-ABP) contains VME, which replaces the isopeptide bond

between two ubiquitin moieties in chains, and also forms a covalent bond with the catalytic

cysteine (Borodovsky et al., 2002; Mulder et al., 2014). Both LotB and LotC showed clear shifts

with all Prg-ABPs (mono-, K48- and K63- linked) with different reactivity. LotB shifted after 30

mins, and only partially, as evident by the amount of unreacted species, while LotC has rapidly

reacted with Prg-ABP and was completely conjugated after 30 minutes (Fig. 2d, e). These results

suggest that both LotB and LotC have a primary ubiquitin binding S1 site where the propargyl

group can be located in close proximity to the catalytic cysteine. In contrast with Prg-ABP, only

LotB reacted with $K63-Ub_2$-VME-ABP. This is consistent with the di-Ub panel assay (Fig. 1c-f),

in which LotB showed specificity toward the K63-linkage. The VME-ABp results suggest an extra

S1' ubiquitin binding site on LotB, which helps to properly locate the K63-linked-VME group on

the catalytic site proximal to the catalytic cysteine. LotC lacks this S1' site so that VME group

between two ubiquitin moieties cannot be reached to the catalytic cysteine. Next, we asked whether

both LotB and LotC can also interact with other ubiquitin-like modifiers (SUMO1/2/3, NEDD8)

(Fig. 2f, g, Supplementary Fig. 2a, b). Interestingly, LotB showed modification with both NEDD8-

Prg and ubiquitin-Prg after 30 minutes of reaction, while LotC was modified only with ubiquitin.

It suggests that LotB binds to ubiquitin through the conserved binding interface with NEDD8,

while LotC interacts with ubiquitin through the specific residues present only in ubiquitin

(Supplementary Fig. 2c).

Structural analysis of *Legionella* OTU-like deubiquitinases

Linkage specificity of the OTU family relies on one of the following mechanisms: 1) additional

ubiquitin-binding domains, 2) ubiquitinated sequences in the substrates or 3) defined S1' or S2

ubiquitin-binding sites (Mevissen et al., 2013). To define the minimal OTU domain for

biochemical and structural studies, we designed several constructs and tested their activity against

the di-Ub panel (Fig. 3a, b). While LotC retained its activity with the predicted OTU domain (7-

310), LotB lost its activity after deletion of 50 amino acids (300-350) located at the C-terminus

beyond the predicted OTU domain (11-283). Based on the LotB structure (PDB:6KS5, (Ma et al.,

2020)), we assumed that this extra helical region might be required for the additional ubiquitin

binding site (S1') to accept the distal ubiquitin moiety from K63 Ub_2 (Fig. 3c). To understand the

detailed mechanism of linkage specificity of LotB and LotC at the molecular level, we determined

the crystal structure of the catalytic domain of LotC ($LotC_{14-310}$) at 2.4 Å (Fig. 3d, Supplementary

Table 1, PDB ID: 6YK8). A structural comparison of both LotB and LotC with other OTU-

deubiquitinases predicted by HHpred revealed that both Lot-DUBs have the unique structural

features in the S1 ubiquitin-binding site (Fig. 3c, d and Table 1). Whereas the overall fold of the

catalytic core of LotB and LotC resembles that of other OTU-deubiquitinases, both showed clear

differences in the helical arm region, which has been shown to interact with ubiquitin and it serves

as an S1 binding site (Mevissen et al., 2013). The structure and sequence alignment with other

OTUs clearly showed that both LotB and LotC contain a relatively long insertion between the Cys-

loop and the variable loop, compared to other OTU members (Fig. 3e). The typical length of the

helical lobe of the known OTUs is ranging from 50 to 60 amino acids (except Otubain family

which contain 110-120 amino acids), while LotB and LotC contain 183 and 210 amino acids,

respectively. Based on this observation, we wondered whether LotA, another *Legionella* OTU-

deubiquitinase (Kubori et al., 2018), also contains a longer insertion in the same region. Based on

the catalytic cysteine and histidine residues of the two OTU domains on LotA (Hermanns and

Hofmann, 2019), we analyzed the sequence and found that both OTU domains of LotA also contain

the longer insertion between the Cys loop and the variable loop (179 and 178 amino acids,

respectively; Fig. 3e). Together, our results identify Lot-DUBs as a novel class of the OTU-family

with longer insertions in the helical lobe region (Supplementary Fig.3a).

Novel structural fold of S1-ubiquitin binding sites on *Legionella* OTUs

Both LotB and LotC have extended helices, specifically near the S1 ubiquitin binding site and we

wondered how these regions interact with ubiquitin. To address this, we performed ubiquitin

docking into both LotB and LotC, followed by molecular dynamics (MD) simulations for 600 ns

(Fig. 4a-d). The final models showed that ubiquitin indeed makes contacts with the additional

helical regions of both LotB and LotC. In LotB, Phe143 and Met144 make hydrophobic

interactions with ubiquitin (Phe45 and Ala46). In addition to these interactions, we also found

another hydrophobic patch in LotB (Ile238, Val247, Ala248, Ile264, Ala266) interacting with

ubiquitin (Leu71, Leu73). For LotC, we identified several hydrophobic interactions of the

extended helical region (Tyr119, Tyr149) with ubiquitin (Ile44). During the simulation, C-

terminus of ubiquitin (Arg72, Arg74) formed transient electrostatic interactions with LotC

(Glu153, Glu245). To validate the interactions observed in the simulations, we introduced several

mutations to the binding interface of both LotB and LotC, and performed a ubiquitin-cleavage

assay (Fig.4e, f). Consistent with a recent study(Ma et al., 2020), mutations of both F143 and M144

decreased the catalytic activity of LotB. Interestingly, mutations of the newly identified

hydrophobic patch (I238, A266) also decreased the catalytic activity. For LotC, the mutations in

the hydrophobic patch (Y119, Y149) also affected the catalytic activity of LotC. Remarkably, a

single mutation on E153 completely abolished the catalytic activity, which indicates that the

electrostatic interactions are important for the correct docking of the ubiquitin C-terminus to the

catalytic pocket of LotC. This observation also explains the result observed in ubiquitin-like

protein ABP assays with LotC (Fig. 2g). The NEDD8, which has alanine instead of Arg72 in

ubiquitin, showed no modification towards LotC (Supplementary Fig. 2c). Together, our results

reveal how the extra helical lobes of the Lot-DUBs interact with ubiquitin and how they differ

within the *Legionella* OTU family.

Physiological roles of LotB and LotC

To gain better insights into the physiological roles of LotB and LotC, we decided to identify their

interacting proteins or substrates. First, to enrich for the interacting partners, catalytically inactive

LotB or LotC were expressed in cells and immuno-precipitated from cell lysates. Ubiquitin

(UBA52) is strongly enriched with both catalytically inactive LotB and LotC (Fig. 5a, c). MS

analysis revealed that LotB mainly interacts with membrane protein complexes (COPB1, ATP5B,

ATP5H, COX5A, SEC61B). We also found interactions with some ER-resident proteins (Calnexin

(CANX), DDOST, STT3A). By contrast, most of the enriched proteins from the inactive LotC

pull-down were non-membrane-bound organelle- and ribosome-related proteins (RPS8, RPLP2,

224 RPS27, RPLP1, RPL13) (Fig. 5a, c). To further understand this, we sought to find the cellular

225 localization of both DUBs (Fig. 5b, d). Consistent with the recent publication, LotB specifically

226 co-localized with the ER marker protein Calnexin, but not with other organelle markers (TOMM20

227 and GM130 for Mitochondria and Golgi, respectively, Fig. 5b), and the OTU domain itself failed

228 to localize on the ER (Supplementary Fig. 4a). By contrast, we could not find a specific cellular

229 localization of LotC (Fig. 5d). Next, to gain more insights into the functional roles of LotB and

230 LotC, we decided to explore combinatorial ubiquitination events with other ubiquitin-related

231 *Legionella* effector proteins. To do this, we co-transfected the cells with one of the Lot-DUBs and

232 previously known *Legionella* E3 ligases (SidC or SdcA) (Hsu et al., 2014; Wasilko et al., 2018).

233 We chose these two ligases because their cellular substrates are poorly studied. TMT-labelled

234 samples from cells expressing either SidC or SdcA alone, or together with a catalytically inactive

235 mutant of LotB or LotC, were prepared (four combinations; SidC-LotB, SidC-LotC, SdcA-LotC

236 or SdcA-LotC, Fig. 5e, f, Supplementary Fig. 5a-d). Interestingly, we identified a distinct sub-

237 class of substrates. Overall, a smaller number of proteins were enriched with LotB compared to

238 LotC. We reasoned that LotB specifically interacts with proteins modified with K63-linked

239 ubiquitin chains, while LotC interacts with different types of ubiquitin chains. Intriguingly, we

240 found a significant number of ribosome-structural proteins in LotC:SdcA combination, which were

241 not enriched in SidC background. Together, these results provide fundamental insights into how

242 *Legionella*-derived E3-ligases and DUBs interact with each other.

244 **DISCUSSION**

245  In this study, we have identified two novel bacterial OTU-deubiquitinases from *Legionella*,

246 which we suggest as founding members of a new sub-class. Unlike classical OTU-DUBs, the

Legionella OTU-DUBs (LOT) possess extended helical insertions between the catalytic Cys-loop

and the variable loop. Molecular dynamic simulations with biochemical studies showed that the

helical insertions interact with ubiquitin. As this insertion is unique for *Legionella* OTUs and not

found in other known OTU-family members, *Legionella* OTU-DUBs define a new sub-class of

the OTU-deubiquitinase family. We have also shown that LotB and LotC are specific to certain

ubiquitin chains and have a distinct cellular localization. Moreover, host-protein interactome

studies revealed that LotB and LotC have different sets of host-interacting proteins to LotB and

LotC. Together, these findings establish important guidance on how to screen for more DUBs in

other pathogenic bacteria or viruses, how to characterize their physiological roles during infection.

We also showed that the two *Legionella* OTUs have different ubiquitin-binding modes

that enable them to cleave specific ubiquitin chains. With ubiquitin activity-based probes (Prg-,

VME- probes), we showed that LotB contains an extra ubiquitin-binding site (S1') and is specific

to K63-linked ubiquitin chains, whereas LotC cleaves different types of ubiquitin chains.

Interestingly, we observed a modification of LotB with NEDD8-Prg ABP. Further studies on

neddylated proteins with LotB will give us more insights into dual-activity of LotB. In contrast,

we could not see the modification between NEDD8-ABP and LotC and we reasoned that the Arg72

on ubiquitin, which is replaced by alanine in NEDD8, is important to locate the C-terminus of

ubiquitin to the catalytic site. Indeed, in molecular dynamic simulations of LotC with ubiquitin we

found Arg72 to from transient interactions with Glu153 of LotC. Further structural analysis will

shed light on how different ubiquitin chains bind to LotB or LotC through the different binding

modes should give us further insights.

Deubiquitinases from bacteria and viruses have been shown to alter the host immune

system. For example, papain-like proteases (PLPro) from coronaviruses, such as middle east

respiratory syndrome (MERS) or severe acute respiratory syndrome (SARS) have dual

deubiquitinase and de-ISGylation activities and antagonized type I interferon (IFN-I) response,

which is the primary defense system against viral infections (Davis and Gack, 2015; Devaraj et al.,

2007; Frieman et al., 2009; Sadler and Williams, 2008). Interestingly, we identified OTUD4 as an

interacting partner of LotB. OTUD4 has been shown to deubiquitinate K63-linked chains of

myeoloid differentiation primary response 88 (MYD88) and to downregulate NF-kB-dependent

inflammation. While the recent study on LotB showed that there is no detectable inhibition of NF-

kB reporter expression(Ma et al., 2020), further studies are awaited to show the cross-talk between

LotB and host ubiquitination system.

Legionella pneumophila has been shown to possess multiple genes altering the host

ubiquitination system. However, little is known about functional cross-talk between ubiquitin

ligases and deubiquitinases. To understand the combinatorial effects of *Legionella* ubiquitin

ligases and deubiquitinases, we analyzed the interactome of LotB/LotC with host proteins in the

presence or absence of *Legionella* ligases (SidC and SdcA). We could see clear differences in the

number of enriched proteins in different combinations. Interestingly, we saw the clear difference

in the number of enriched proteins. A significant number of ribosomal proteins were enriched with

LotC in the background of SdcA, but not from SidC background, which has 71 % sequence

similarity to SdcA. This finding suggests distinct physiological roles of SdcA and SidC, and a

putative relationship between LotC and SdcA on regulating translation processes. Since LotC

processes different types of ubiquitin chains, and is mainly localized to the cytosol, a catalytically

dead version of LotC can be used as a standard tool for the identification of specific substrates of

other known *Legionella* ligases. It would be interesting to understand how all these ubiquitin

machineries synergistically work together and alter the host-ubiquitination system at different time

points of infection.

MATERIAL AND METHODS

Protein expression and purification

All proteins used in this study were expressed and purified as previously described (Bhogaraju et

al., 2016; Qiu et al., 2016). Lpg1621 (LotB), Lpg2529 (LotC), Lpg2411 and Lpg2907 were cloned

into either pParallelGST2 or pParallelHis2 vector (Sheffield et al., 1999). T7 express *E. coli*

competent cells (NEB) were transformed with plasmids and grown in LB medium to an OD_{600} of

0.6-0.8 at 37°C. Protein expression was induced by addition of 0.5 mM IPTG (isopropyl D-

thiogalactopyranoside) and the cells were further grown overnight at 18°C and harvested. The cell

pellet was resuspended in lysis buffer (50 mM Tris-HCl pH 7.5 and 150 mM NaCl, 2 mM DTT)

and lysed by sonication and centrifuged at 13,000 rpm to clarify the supernatant. The supernatant

of GST-tagged protein was incubated 1 hour with glutathione-*S*-sepharose pre-equilibrated with

washing buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl, 2 mM DTT) and non-specific proteins

were cleared with washing. GST-proteins were eluted with elution buffer (50 mM Tris-HCl pH

8.0, 50 mM NaCl, 2 mM DTT, 15 mM reduced glutathione) and buffer exchanged to storage buffer

(50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM DTT). For His-tagged proteins, the supernatant

was incubated with Ni-NTA pre-equilibrated with washing buffer (50 mM Tris-HCl, pH 7.5, 500

mM NaCl, 20 mM Imidazole) for 2 hours and eluted with elution buffer (50 mM Tris-HCl, pH 7.5,

500 mM NaCl, 300 mM Imidazole) and the buffer was exchanged to storage buffer. For $LotC_{14-310}$,

instead of using the elution buffer, glutathione beads were incubated with sfGFP-TEV protease

(Wu et al., 2009) overnight at 4°C. Cleaved protein were buffer exchanged to IEX buffer A (20

mM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM DTT) and purified by anion-exchange chromatography

on HitrapQ (GE Healthcare) with gradient elution with IEX buffer B (20 mM Tris-HCl, pH 8.0, 1

M NaCl, 1 mM DTT) and fractions contacting samples were loaded onto size-exclusion column

(Superdex 75 16/60, GE Healthcare) pre-equilibrated with 50 mM Tris-HCl pH 7.5, 50 mM NaCl,

1 mM TCEP. Proteins were concentrated to 20 mg/ml and stored for crystallization.

Di-Ub panel cleavage assay

To activate DUBs, 3 µl of 5 µM of DUBs were mixed with 12 µl of activation buffer (25 mM Tris-
HCl pH 7.5, 150 mM NaCl, 10 mM DTT) and incubated 15 minutes at 25°C. For di-Ubiquitin
samples, 3 µl of di-ubiquitin chains (1 mg/ml) were mixed with 3 µl 10x reaction buffer (500 mM
Tris-HCl pH 7.5, 500 mM NaCl, 50 mM DTT) and 12 µl of ultra-pure water. To initiate the
reaction, the activated DUBs were mixed with di-Ubiquitin and samples were taken at the indicated
time points and the reactions were quenched by the addition of SDS-sample buffer. The samples
were further analyzed by SDS-PAGE and stained with silver-staining kit (Pierce Silver Staining
Kit, Thermo Fischer).
330
Ubiquitin/NEDD8/SUMO-/ISG15/UFM activity-based probes assay
DUBs were diluted (1.5 µM, final concentration) with activation buffer and incubated 10 minutes
at 25°C and the activity-based probes were diluted (50 µM, final concentration) in dilution buffer
(50 mM Tris-HCl 7.5, 150 mM NaCl). The total of 30 ul of reaction mixture was prepared by
mixing 20 µl of activated DUBs (1.5 µM), 3 µl of activity-based probes, 3 µl of reaction buffer
(500 mM Tris-HCl pH 7.5, 500 mM NaCl, 50 mM DTT) and 4 µl of ultra-pure water. Samples
were taken at the indicated time points and the reactions were quenched by the addition of SDS-
sample buffer. Samples were further analyzed by SDS-PAGE and stained with Coomassie staining
solution.
340
Crystallization
The concentrated LotC$_{14-310}$ were screened with sitting drop matrix screens in 96-well plate with
100 nl of protein and 100 nl of precipitant solution at 293K. Initial crystals appeared from solution
containing 25 % PEG3350, 100 mM Tris-HCl pH 8.5, 200 mM NaCl with 18.4 mg/ml protein
concentration. Diffraction-quality crystals were grown in optimized solution containing 19 % PEG
3350, 100 mM Tris-HCl pH 8.5, 150 mM NaCl with 24 mg/ml protein concentration.
347
Data collection, processing and structure determination
To obtain the phase, 0.4 µl of 10 mM K$_2$PtCl$_4$ was added to the drop containing crystals and
incubated for 18 hours. Heavy atom-soaked crystals were cryo-protected using mother liquor
solution supplemented with 15 % (v/v) glycerol. Diffraction data were collected on single frozen
crystal in a nitrogen stream at 100K at beamline PXI as Swiss Light Source, Villigen. Initial data

sets were processed using XDS (Kabsch, 2010), and initial-phases were determined by Autosol in

Phenix (Terwilliger et al., 2009). Structure refinement and manual model building were performed

with Coot and Phenix.Refine (Afonine et al., 2012; Emsley et al., 2010).

Protein-protein docking

We used the Rosetta protein-protein docking method (Gray et al., 2003) to identify low-energy

conformations of the complexes of ubiquitin with LotB and LotC, respectively. Given that the C-

terminus of ubiquitin should interact with the catalytic residue of the OTUs, we used the local

docking approach in which we placed the C-terminal end of ubiquitin (Gly76) near the catalytic

residues in both ligases (Cys29 and His270 for LotB and Cys16 and His296 for LotC). We then

started the docking by optimizing the rigid-body orientation and side-chain conformation sampling.

The program requires two protein structures as inputs, which were prepared by running the

refinement protocol before the docking step. We performed the local docking approach and

generated 100 independent structures for each complex. The complexes in this way were subject

to local refinement to remove remaining small clashes. The complexes were then clustered based

on the distance matrix of Cα atoms between the ligase and ubiquitin using the KMeans method.

The representatives of two major clusters in each case were selected based on the interface score

(I_sc), which represents the energy of the interactions across the interface of two proteins. These

representative complexes were used for MD simulations.

Molecular dynamics simulations

All-atom explicit solvent molecular dynamics (MD) simulations were performed for two docking

results for each ligase. The systems were built using the CHARMM-GUI webserver (Wu et al.,

2014). The systems were hydrated with 150 mM NaCl electrolyte. The all-atom CHARMM36m

force field was used for proteins, lipids and ions, and TIP3P was used for water molecules (Best

et al., 2012). The MD trajectories were analyzed with Visual Molecular Dynamics (VMD)

(Humphrey et al., 1996). The MD simulations were performed using GROMACS 2019 (Abraham

et al., 2015). The starting system was minimized for 5,000 steps with the steepest descent energy

minimization and equilibrated for 6.5 ns of MD simulation first in the NVT ensemble (1.5 ns) and

then in the NPT (5 ns) ensemble, in which all non-hydrogen atoms of the protein were restrained

to the fixed reference positions with progressively reduced force constants, starting at 1000

kJ·mol^{-1}·nm^2. Afterwards, the production runs were carried out in the NPT ensemble for 600 ns

for each setup. To keep the C-terminusof ubiquitin in the catalytic site, 7-Å wall restraints were

placed on the distance between Cα of G76$_{UB}$ and Cys29/His270 in LotB and between Cα of G76$_{UB}$

and Cys16/His296 in LotC. Periodic boundary conditions were used. Particle mesh Ewald (Darden

et al., 1993) with cubic interpolation and 0.12 nm grid spacing for Fast Fourier Transform was

used to treat long-range electrostatic interactions. The time step was 2 fs. The LINCS algorithm

(Hess et al., 1997) was used to fix all bond lengths. Constant temperature (310 K) was set with a

Nosé-Hoover thermostat (Hoover, 1985), with a coupling constant of 1.0 ps. An isotropic

Parrinello-Rahman barostat (Parrinello and Rahman, 1981) with a coupling constant of 5.0 ps was

used to maintain a pressure of 1 bar.

Identification of novel substrates of LotB and LotC

For interactome analysis, HEK 293 cells were transfected with FLAG-LotB WT/C29A or FLAG-

LotC WT/C24A. To identify the substrates modified by *Legionella* derived E3 ligases, GFP-SidC

or GFP-SdcA were co-transfected with FLAG-LotB WT/C29A or FLAF-LotC WT/C24A.

Cells were lysed in ice cold lysis buffer (50mM Tris-Cl, p.H 7.5; 150mM NaCl; 1% Triton x-100)

and equal amount of lysates were incubated with FLAG-M2 beads in IP buffer (Lysis buffer

without detergent). After incubation, IPs were washed with wash buffer (50mM Tris-Cl, pH7.5;

400mM NaCl; 0.5mM EDTA) and the interacting proteins were eluted with 8M urea solution.

After the reduction and alkylation with TCEP and Chloroacetamide, the samples were digested

with 0.5ug Trypsin (Promega) at 37°C overnight after diluting the urea < 2M. Digests were

acidified using trifluoroaceticacid (TFA) to a pH of 2-3, and the peptides were enriched using C18

stage tips (Rappsilber et al., 2003). To get quantitative information, peptides were either labelled

with TMT 10 plex reagent (Thermo fisher) or analyzed label-free. The peptides were separated on

an in-house made C18 column (20 cm length, 75 μm inner diameter, 1.9 μm particle size) by an

easy n-LC 1200 (ThermoFisher) and directly injected in a QExactive-HF or in case of TMT

samples into a Fusion Lumos mass-spectrometer (ThermoFisher) and analyzed in data-dependent

mode. Data analysis was done using Maxquant 1.65 (Cox and Mann, 2008). Fragment spectra

were searched against *Homo sapiens* SwissProt database (TaxID:9606, version 2018). Label-free

quantification was done with MaxLFQ (Cox et al., 2014) method with activated match between

runs. TMT-labelled samples were analysed by using TMT 10 Plex option within the software.

Statistically significant changes between samples were determined using a Two-sample t-test with

a permutation-based FDR of 5% on log2 transformed values in Perseus (Tyanova et al., 2016).

Confocal imaging and image analysis

U2OS cells were transfected with FLAG-LotB or FLAG-LotC by GeneJuice transfection reagent

(Merck) for 24 h and fixed by 4% paraformaldehyde for 20 min. After fixation, cells were

permeabilized and blocked by 0.1% saponin and 1% BSA in PBS for 1 h at room temperature.

Cells were incubated with anti-Flag antibody (Sigma and Cell Signaling), with either anti-calnexin

antibody (Abcam), anti-TOMM20 antibody (Abcam), or anti-GM130 (BD Transduction

Laboratories) at 4 °C overnight. Alexa Fluor 488 and Alexa Fluor 546 (Invitrogen) secondary

antibody were incubated for 1 h at room temperature. Images were acquired by the Zeiss LSM780

microscope system with 63× 1.4 NA oil immersion objective and further analyzed by Zeiss Zen

microscope software.

428
429 **ACKNOWLEDGEMENTS:**
430

We thank Yuxin-Mao for providing SidC and SdcA clones and Stefan Knapp for the advice in

structure determination and sharing synchrotron time. The authors also thank staff at SLS for their

support during crystallographic X-ray diffraction test and data collection. The data collection at

SLS has been supported by the funding from the European Union's Horizon 2020 research and

innovation program under grant agreement number 730872, project CALIPSOplus. This project

was supported by the European Research Council (ERC) under the European Union's Horizon

2020 research and innovation program (ID, grant agreement No 742720), the LOEWE program

DynaMem of the State of Hesse (Germany, Project-ID III L6-519/03/03.001 – (0006)), and

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation (Project-ID 259130777 –

SFB1177; Leibniz-Program to ID; CEF-MC - EXC115/2; SFB 902), the Max Planck Society and

NWO-VIDI grant and Off-rad grand for G.H.

COMPETING INTERSTES: The authors declare no competing interests.

Figure Legends and Tables:
Table1. TOP 5 candidates for putative DUBs from Legionella effector proteins

Legionella Proteins		Target Proteins				
Name	Aligned Region	Name	Aligned Region	Probability (%)	Identities (%)	PDB ID_Chain
Lpg1621	195-274	Viral OTU (CC hemorrhagic fever virus)	69-157	92.59	16	3PHU_B
	195-274	Human OTUD2	63-140	92.52	17	4BOQ_A
	195-279	Human OTUD3	59-142	92.40	13	4BOU_A
	193-278	Human OTUD5	100-183	91.18	21	3PFY_A
	192-279	Viral OTU (Farallon virus)	88-183	91.08	16	6DX5_B
Lpg2529	1-310	Viral OTU (Erve virus)	17-157	96.24	18	5JZE_A
	7-310	Viral OTU (Dera Ghazi Khan orthonairovirus)	25-156	96.15	18	6DX2_B
	20-310	Human Otubain1	50-234	96.02	13	2ZFY_A
	20-310	Human Otubain2	50-233	95.84	13	4FJV_C
	7-310	Viral OTU (Taggert virus)	23-156	95.71	13	6DX3_D
Lpg2411	110-216	Yeast UCH8	152-259	37.94	11	3MHS_A
	183-272	EntA-im (Enterococcus faecium)	7-89	37.06	15	2BL8_B
	33-94	Uncharacterized protein (Corynebacterium diphtheriae)	7-72	36.65	21	3KDQ_D
	120-212	PG0816 (Porphyromonas gingivalis)	53-139	35.31	16	2APL_A
	104-114	PSII reaction center protein K (Cyanidium caldarium)	2-12	32.96	36	4YUU_X2
Lpg2907	117-384	AvrA (Salmonella typhimurium)	59-299	100	11	6BE0_A
	158-398	PopP2 (Arabidopsis thaliana)	99-339	99.93	13	5W3X_C
	115-390	HopZ1a (Pseudomonas syringae)	54-342	99.88	10	5KLP_C
	118-275	XopD (Xanthomonas campestris)	1-148	95.53	11	2OIX_A
	95-276	Human SENP1	2-177	95.21	16	2G4D_A

447   Values are obtained from HHpred server (MPI Bioinformatics Toolkit)
448

Figure 1. Identification of novel deubiqutinases in *Legionella pneumophila.*

(a) Graphical illustration of identification of novel deubiquitinases of Legionella pneumophila Lpg

Genes. **(b)** Predicted deubiquitinase domain of four putative *Legionella* deubiquitinases. **(c, e)**

Time-course Di-ubiquitin panel cleavage assay with Lpg1621 (LotB) and Lpg2529 (LotC),

respectively. **(d, f)** Linkage specificity diagram of Lpg1621 (LotB) and Lpg2529 (LotC).

Percentage of cleaved ubiquitin species at 90 mins were plotted.

Figure 2. Biochemical properties of LotB and LotC

(a) Predicted catalytic residues on LotB and LotC. **(b, c)** Di-Ub cleavage activity assay with wild-

type and catalytic mutants of LotB and LotC, respectively. **(d, e)** Activity based probes (ABPs)

test on LotB and LotC. Propargyl-Ub-ABP (Prg-ABP) and vinylmethylester-ubiquitin-ABP

(VME-ABP) were incubated as indicated time-points with LotB and LotC and analyzed on SDS-

PAGE with Coomassie staining. **(f, g)** Propargyl ubiquitin or ubiquitin-like modifiers reactivity

test on LotB and LotC. Prg-ABPs are incubated with LotB and LotC with indicated time points.

Figure 3. Structural comparison of *Legionella* OTU-DUBs with other OTU-family

(a, b) Minimal domain boundaries of catalytically active LotB and LotC. Different constructs were

cloned based on the predicted OTU-domains and their activity, and were tested with di-Ub$_2$ panel

(c, d) Structural comparison of LotB and LotC with their closest homologues. CCHF- (PDB:

3PHU), Farallon- (PDB: 6DX5), OTUD2 (PDB: 4BOQ), OTUD3 (PDB: 4BOU), Taggert- (PDB:

6DX3), DGK nairo- (PDB: 6DX2), Otubain1 (PDB: 2ZFY), Otubain2 (PDB: 4FJV). **(e)** Sequence

alignment of LotB and LotC with their closest homologues. Catalytic cysteine and histidine are

highlighted in red and conserved residues are highlighted in yellow.

Figure 4. Ubiquitin binding sites on LotB and LotC

(a, b) Molecular docking and simulations of monoubiquitin to LotB and LotC, respectively. Shown

are representative snapshots of the MD simulations. Catalytic cysteine and key residues for the

interaction between ubiquitin and LotB or LorC are depicted as sticks. **(c, d)** Key residues

mediating interactions of ubiquitin with LotB and LotC, respectively. Residues are highlighted in

the structure (left). Side-chain center-of-mass distances are shown as a function of the simulation

time (right). **(e, f)** Di-ubiquitin cleavage assay for mutants of LotB and LotC. The catalytic activity

of LotB or LotC wild-type and its mutants was tested with K63- or K48- linked Ub2, respectively.

Figure 5. Physiological roles of LotB and LotC

(a, c) Proteomic analysis of interacting partners of LotB and LotC, respectively. Catalytically

inactive FLAG-LotB (C29A) and FLAG-LotC (C24A) were transfected and immunoprecipitated.

Co-precipitated interacting proteins were analyzed by mass spectrometry. **(b, d)** Cellular

localization of LotB and LotC. FLAG-tagged LotB and LotC were ectopically expressed in U2OS

cells and immunostained with cellular organelle markers (ER; Calnexin, Mitochondria: TOMM20,

Golgi: GM130). **(e, f)** List of significantly enriched proteins from LotB and LotC with or without

Legionella ligases (SdcA or SidC) as indicated.

REFERENCES:

Abdul Rehman SA, Kristariyanto YA, Choi S-Y, Nkosi PJ, Weidlich S, Labib K, Hofmann K, Kulathu Y. 2016. MINDY-1 Is a Member of an Evolutionarily Conserved and Structurally Distinct New Family of Deubiquitinating Enzymes. *Mol Cell* **63**:146–155. doi:10.1016/j.molcel.2016.05.009

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. 2015. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **1–2**:19–25. doi:10.1016/j.softx.2015.06.001

Afonine P V., Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD. 2012. Towards automated crystallographic structure refinement with phenix.refine. *Acta Crystallogr Sect D Biol Crystallogr* **68**:352–367. doi:10.1107/S0907444912001308

Akutsu M, Ye Y, Virdee S, Chin JW, Komander D. 2011. Molecular basis for ubiquitin and ISG15 cross-reactivity in viral ovarian tumor domains. *Proc Natl Acad Sci* **108**:2228–2233. doi:10.1073/pnas.1015287108

Best RB, Zhu X, Shim J, Lopes PEM, Mittal J, Feig M, MacKerell AD. 2012. Optimization of the Additive CHARMM All-Atom Protein Force Field Targeting Improved Sampling of the Backbone ϕ, ψ and Side-Chain χ 1 and χ 2 Dihedral Angles. *J Chem Theory Comput* **8**:3257–3273. doi:10.1021/ct300400x

Bhogaraju S, Bonn F, Mukherjee R, Adams M, Pfleiderer MM, Galej WP, Matkovic V, Lopez-Mosqueda J, Kalayil S, Shin D, Dikic I. 2019. Inhibition of bacterial ubiquitin ligases by SidJ–calmodulin catalysed glutamylation. *Nature* **572**:382–386. doi:10.1038/s41586-019-1440-8

Bhogaraju S, Kalayil S, Liu Y, Bonn F, Colby T, Matic I, Dikic I. 2016. Phosphoribosylation of Ubiquitin Promotes Serine Ubiquitination and Impairs Conventional Ubiquitination. *Cell* **167**:1636-1649.e13. doi:10.1016/j.cell.2016.11.019

Black MH, Osinski A, Gradowski M, Servage KA, Pawłowski K, Tomchick DR, Tagliabracci VS. 2019. Bacterial pseudokinase catalyzes protein polyglutamylation to inhibit the SidE-family ubiquitin ligases. *Science (80-)* **364**:787–792. doi:10.1126/science.aaw7446

Borodovsky A, Ovaa H, Kolli N, Gan-Erdene T, Wilkinson KD, Ploegh HL, Kessler BM. 2002. Chemistry-Based Functional Proteomics Reveals Novel Members of the Deubiquitinating

Enzyme Family. *Chem Biol* **9**:1149–1159. doi:10.1016/S1074-5521(02)00248-X

Bremm A, Freund SM V, Komander D. 2010. Lys11-linked ubiquitin chains adopt compact

conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne. *Nat Struct*

Mol Biol **17**:939–947. doi:10.1038/nsmb.1873

Burstein D, Amaro F, Zusman T, Lifshitz Z, Cohen O, Gilbert JA, Pupko T, Shuman HA, Segal

G. 2016. Genomic analysis of 38 Legionella species identifies large and diverse effector

repertoires. *Nat Genet* **48**:167–175. doi:10.1038/ng.3481

Capodagli GC, Deaton MK, Baker EA, Lumpkin RJ, Pegan SD. 2013. Diversity of Ubiquitin and

ISG15 Specificity among Nairoviruses' Viral Ovarian Tumor Domain Proteases. *J Virol*

87:3815–3827. doi:10.1128/JVI.03252-12

Clague MJ, Urbé S, Komander D. 2019. Breaking the chains: deubiquitylating enzyme

specificity begets function. *Nat Rev Mol Cell Biol* **20**:338–352. doi:10.1038/s41580-019-

0099-1

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate Proteome-wide Label-

free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction,

Termed MaxLFQ. *Mol Cell Proteomics* **13**:2513–2526. doi:10.1074/mcp.M113.031591

Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized p.p.b.-

range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**:1367–

1372. doi:10.1038/nbt.1511

Darden T, York D, Pedersen L. 1993. Particle mesh Ewald: An $N \cdot \log(N)$ method for Ewald

sums in large systems. *J Chem Phys* **98**:10089–10092. doi:10.1063/1.464397

Davis ME, Gack MU. 2015. Ubiquitination in the antiviral immune response. *Virology* **479–**

480:52–65. doi:10.1016/j.virol.2015.02.033

Devaraj SG, Wang N, Chen Zhongbin, Chen Zihong, Tseng M, Barretto N, Lin R, Peters CJ,

Tseng C-TK, Baker SC, Li K. 2007. Regulation of IRF-3-dependent Innate Immunity by the

Papain-like Protease Domain of the Severe Acute Respiratory Syndrome Coronavirus. *J*

Biol Chem **282**:32208–32221. doi:10.1074/jbc.M704870200

Diao J, Zhang Y, Huibregtse JM, Zhou D, Chen J. 2008. Crystal structure of SopA, a Salmonella

effector protein mimicking a eukaryotic ubiquitin ligase. *Nat Struct Mol Biol* **15**:65–70.

doi:10.1038/nsmb1346

Dikic I. 2017. Proteasomal and Autophagic Degradation Systems. *Annu Rev Biochem* **86**:193–

224. doi:10.1146/annurev-biochem-061516-044908

Edelmann MJ, Iphöfer A, Akutsu M, Altun M, di Gleria K, Kramer HB, Fiebiger E, Dhe-
Paganon S, Kessler BM. 2009. Structural basis and specificity of human otubain 1-mediated
deubiquitination. *Biochem J* **418**:379–390. doi:10.1042/BJ20081318

Ekkebus R, van Kasteren SI, Kulathu Y, Scholten A, Berlin I, Geurink PP, de Jong A, Goerdayal
S, Neefjes J, Heck AJR, Komander D, Ovaa H. 2013. On Terminal Alkynes That Can React
with Active-Site Cysteine Nucleophiles in Proteases. *J Am Chem Soc* **135**:2867–2870.
doi:10.1021/ja309802n

El Oualid F, Merkx R, Ekkebus R, Hameed DS, Smit JJ, de Jong A, Hilkmann H, Sixma TK,
Ovaa H. 2010. Chemical Synthesis of Ubiquitin, Ubiquitin-Based Probes, and Diubiquitin.
Angew Chemie Int Ed **49**:10149–10153. doi:10.1002/anie.201005995

Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot. *Acta
Crystallogr Sect D Biol Crystallogr* **66**:486–501. doi:10.1107/S0907444910007493

Ensminger AW, Isberg RR. 2010. E3 Ubiquitin Ligase Activity and Targeting of BAT3 by
Multiple Legionella pneumophila Translocated Substrates. *Infect Immun* **78**:3905–3919.
doi:10.1128/IAI.00344-10

Fiskin E, Bhogaraju S, Herhaus L, Kalayil S, Hahn M, Dikic I. 2017. Structural basis for the
recognition and degradation of host TRIM proteins by Salmonella effector SopA. *Nat
Commun* **8**:14004. doi:10.1038/ncomms14004

Flierman D, van der Heden van Noort GJ, Ekkebus R, Geurink PP, Mevissen TET, Hospenthal
MK, Komander D, Ovaa H. 2016. Non-hydrolyzable Diubiquitin Probes Reveal Linkage-
Specific Reactivity of Deubiquitylating Enzymes Mediated by S2 Pockets. *Cell Chem Biol*
23:472–482. doi:10.1016/j.chembiol.2016.03.009

Frieman M, Ratia K, Johnston RE, Mesecar AD, Baric RS. 2009. Severe Acute Respiratory
Syndrome Coronavirus Papain-Like Protease Ubiquitin-Like Domain and Catalytic Domain
Regulate Antagonism of IRF3 and NF- B Signaling. *J Virol* **83**:6689–6705.
doi:10.1128/JVI.02220-08

Furtado AR, Essid M, Perrinet S, Balañá ME, Yoder N, Dehoux P, Subtil A. 2013. The
chlamydial OTU domain-containing protein Chla OTU is an early type III secretion effector
targeting ubiquitin and NDP52. *Cell Microbiol* **15**:2064–2079. doi:10.1111/cmi.12171

Gray JJ, Moughon SE, Kortemme T, Schueler-Furman O, Misura KMS, Morozov A V., Baker

584 D. 2003. Protein-protein docking predictions for the CAPRI experiment. *Proteins Struct*

585  *Funct Genet* **52**:118–122. doi:10.1002/prot.10384

586 Haahr P, Borgermann N, Guo X, Typas D, Achuthankutty D, Hoffmann S, Shearer R, Sixma

587  TK, Mailand N. 2018. ZUFSP Deubiquitylates K63-Linked Polyubiquitin Chains to

588  Promote Genome Stability. *Mol Cell* **70**:165-174.e6. doi:10.1016/j.molcel.2018.02.024

589 Hermanns T, Hofmann K. 2019. Bacterial DUBs: deubiquitination beyond the seven classes.

590  *Biochem Soc Trans* **47**:1857–1866. doi:10.1042/BST20190526

591 Hermanns T, Pichlo C, Woiwode I, Klopffleisch K, Witting KF, Ovaa H, Baumann U, Hofmann

592  K. 2018. A family of unconventional deubiquitinases with modular chain specificity

593  determinants. *Nat Commun* **9**:799. doi:10.1038/s41467-018-03148-5

594 Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. 1997. LINCS: A linear constraint solver for

595  molecular simulations. *J Comput Chem* **18**:1463–1472. doi:10.1002/(SICI)1096-

596  987X(199709)18:12<1463::AID-JCC4>3.0.CO;2-H

597 Hewings DS, Heideker J, Ma TP, AhYoung AP, El Oualid F, Amore A, Costakes GT,

598  Kirchhofer D, Brasher B, Pillow T, Popovych N, Maurer T, Schwerdtfeger C, Forrest WF,

599  Yu K, Flygare J, Bogyo M, Wertz IE. 2018. Reactive-site-centric chemoproteomics

600  identifies a distinct class of deubiquitinase enzymes. *Nat Commun* **9**:1162.

601  doi:10.1038/s41467-018-03511-6

602 Hoover WG. 1985. Canonical dynamics: Equilibrium phase-space distributions. *Phys Rev A*

603  **31**:1695–1697. doi:10.1103/PhysRevA.31.1695

604 Hsu F, Luo X, Qiu J, Teng Y-B, Jin J, Smolka MB, Luo Z-Q, Mao Y. 2014. The Legionella

605  effector SidC defines a unique family of ubiquitin ligases important for bacterial

606  phagosomal remodeling. *Proc Natl Acad Sci* **111**:10538–10543.

607  doi:10.1073/pnas.1402605111

608 Humphrey W, Dalke A, Schulten K. 1996. VMD: Visual molecular dynamics. *J Mol Graph*

609  **14**:33–38. doi:10.1016/0263-7855(96)00018-5

610 James TW, Frias-Staheli N, Bacik J-P, Levingston Macleod JM, Khajehpour M, Garcia-Sastre A,

611  Mark BL. 2011. Structural basis for the removal of ubiquitin and interferon-stimulated gene

612  15 by a viral ovarian tumor domain-containing protease. *Proc Natl Acad Sci* **108**:2222–

613  2227. doi:10.1073/pnas.1013388108

614 Kabsch W. 2010. XDS. *Acta Crystallogr Sect D Biol Crystallogr* **66**:125–132.

doi:10.1107/S0907444909047337

Kalayil S, Bhogaraju S, Bonn F, Shin D, Liu Y, Gan N, Basquin J, Grumati P, Luo Z-Q, Dikic I.
2018. Insights into catalysis and function of phosphoribosyl-linked serine ubiquitination.
Nature **557**:734–738. doi:10.1038/s41586-018-0145-8

Keusekotten K, Elliott PR, Glockner L, Fiil BK, Damgaard RB, Kulathu Y, Wauer T,
Hospenthal MK, Gyrd-Hansen M, Krappmann D, Hofmann K, Komander D. 2013.
OTULIN Antagonizes LUBAC Signaling by Specifically Hydrolyzing Met1-Linked
Polyubiquitin. *Cell* **153**:1312–1326. doi:10.1016/j.cell.2013.05.014

Kubori T, Hyakutake A, Nagai H. 2008. Legionella translocates an E3 ubiquitin ligase that has
multiple U-boxes with distinct functions. *Mol Microbiol* **67**:1307–1319. doi:10.1111/j.1365-
2958.2008.06124.x

Kubori T, Kitao T, Ando H, Nagai H. 2018. LotA, a Legionella deubiquitinase, has dual catalytic
activity and contributes to intracellular growth. *Cell Microbiol* **20**:e12840.
doi:10.1111/cmi.12840

Kwasna D, Abdul Rehman SA, Natarajan J, Matthews S, Madden R, De Cesare V, Weidlich S,
Virdee S, Ahel I, Gibbs-Seymour I, Kulathu Y. 2018. Discovery and Characterization of
ZUFSP/ZUP1, a Distinct Deubiquitinase Class Important for Genome Stability. *Mol Cell*
70:150-164.e6. doi:10.1016/j.molcel.2018.02.023

Lin DY -w., Diao J, Chen J. 2012. Crystal structures of two bacterial HECT-like E3 ligases in
complex with a human E2 reveal atomic details of pathogen-host interactions. *Proc Natl*
Acad Sci **109**:1925–1930. doi:10.1073/pnas.1115025109

Lombardi C, Ayach M, Beaurepaire L, Chenon M, Andreani J, Guerois R, Jupin I, Bressanelli S.
2013. A Compact Viral Processing Proteinase/Ubiquitin Hydrolase from the OTU Family.
PLoS Pathog **9**:e1003560. doi:10.1371/journal.ppat.1003560

Ma K, Zhen X, Zhou B, Gan N, Cao Y, Fan C, Ouyang S, Luo Z-Q, Qiu J. 2020. The bacterial
deubiquitinase Ceg23 regulates the association of Lys-63–linked polyubiquitin molecules
on the Legionella phagosome. *J Biol Chem* **295**:1646–1657. doi:10.1074/jbc.RA119.011758

Mevissen TET, Hospenthal MK, Geurink PP, Elliott PR, Akutsu M, Arnaudo N, Ekkebus R,
Kulathu Y, Wauer T, El Oualid F, Freund SMV, Ovaa H, Komander D. 2013. OTU
Deubiquitinases Reveal Mechanisms of Linkage Specificity and Enable Ubiquitin Chain
Restriction Analysis. *Cell* **154**:169–184. doi:10.1016/j.cell.2013.05.046

Mevissen TET, Komander D. 2017. Mechanisms of Deubiquitinase Specificity and Regulation.

Annu Rev Biochem **86**:159–192. doi:10.1146/annurev-biochem-061516-044916

Mevissen TET, Kulathu Y, Mulder MPC, Geurink PP, Maslen SL, Gersch M, Elliott PR, Burke

JE, van Tol BDM, Akutsu M, El Oualid F, Kawasaki M, Freund SMV, Ovaa H, Komander

D. 2016. Molecular basis of Lys11-polyubiquitin specificity in the deubiquitinase Cezanne.

Nature **538**:402–405. doi:10.1038/nature19836

Mulder MPC, El Oualid F, ter Beek J, Ovaa H. 2014. A Native Chemical Ligation Handle that

Enables the Synthesis of Advanced Activity-Based Probes: Diubiquitin as a Case Study.

ChemBioChem **15**:946–949. doi:10.1002/cbic.201402012

Parrinello M, Rahman A. 1981. Polymorphic transitions in single crystals: A new molecular

dynamics method. *J Appl Phys* **52**:7182–7190. doi:10.1063/1.328693

Pohl C, Dikic I. 2019. Cellular quality control by the ubiquitin-proteasome system and

autophagy. *Science (80-)* **366**:818–822. doi:10.1126/science.aax3769

Pruneda JN, Durkin CH, Geurink PP, Ovaa H, Santhanam B, Holden DW, Komander D. 2016.

The Molecular Basis for Ubiquitin and Ubiquitin-like Specificities in Bacterial Effector

Proteases. *Mol Cell* **63**:261–276. doi:10.1016/j.molcel.2016.06.015

Qiu J, Sheedlo MJ, Yu K, Tan Y, Nakayasu ES, Das C, Liu X, Luo Z-Q. 2016. Ubiquitination

independent of E1 and E2 enzymes by bacterial effectors. *Nature* **533**:120–124.

doi:10.1038/nature17657

Quaile AT, Urbanus ML, Stogios PJ, Nocek B, Skarina T, Ensminger AW, Savchenko A. 2015.

Molecular Characterization of LubX: Functional Divergence of the U-Box Fold by

Legionella pneumophila. *Structure* **23**:1459–1469. doi:10.1016/j.str.2015.05.020

Rappsilber J, Ishihama Y, Mann M. 2003. Stop and Go Extraction Tips for Matrix-Assisted

Laser Desorption/Ionization, Nanoelectrospray, and LC/MS Sample Pretreatment in

Proteomics. *Anal Chem* **75**:663–670. doi:10.1021/ac026117i

Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD. 2018. The MEROPS

database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison

with peptidases in the PANTHER database. *Nucleic Acids Res* **46**:D624–D632.

doi:10.1093/nar/gkx1134

Sadler AJ, Williams BRG. 2008. Interferon-inducible antiviral effectors. *Nat Rev Immunol*

8:559–568. doi:10.1038/nri2314

Sheffield P, Garrard S, Derewenda Z. 1999. Overcoming expression and purification problems of

RhoGDI using a family of "parallel" expression vectors. *Protein Expr Purif* **15**:34–9.

doi:10.1006/prep.1998.1003

Shin D, Mukherjee R, Liu Yaobin, Gonzalez A, Bonn F, Liu Yan, Rogov V V., Heinz M, Stolz

A, Hummer G, Dötsch V, Luo Z-Q, Bhogaraju S, Dikic I. 2020. Regulation of

Phosphoribosyl-Linked Serine Ubiquitination by Deubiquitinases DupA and DupB. *Mol*

Cell **77**:164-179.e6. doi:10.1016/j.molcel.2019.10.019

Sommer S, Weikart ND, Linne U, Mootz HD. 2013. Covalent inhibition of SUMO and

ubiquitin-specific cysteine proteases by an in situ thiol–alkyne addition. *Bioorg Med Chem*

21:2511–2517. doi:10.1016/j.bmc.2013.02.039

Terwilliger TC, Adams PD, Read RJ, McCoy AJ, Moriarty NW, Grosse-Kunstleve RW, Afonine

P V., Zwart PH, Hung L-W. 2009. Decision-making in structure solution using Bayesian

estimates of map quality: the PHENIX AutoSol wizard. *Acta Crystallogr Sect D Biol*

Crystallogr **65**:582–601. doi:10.1107/S0907444909012098

Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J. 2016. The

Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat*

Methods **13**:731–740. doi:10.1038/nmeth.3901

van Kasteren PB, Bailey-Elkin BA, James TW, Ninaber DK, Beugeling C, Khajehpour M,

Snijder EJ, Mark BL, Kikkert M. 2013. Deubiquitinase function of arterivirus papain-like

protease 2 suppresses the innate immune response in infected host cells. *Proc Natl Acad Sci*

110:E838–E847. doi:10.1073/pnas.1218464110

Wang T, Yin L, Cooper EM, Lai M-Y, Dickey S, Pickart CM, Fushman D, Wilkinson KD,

Cohen RE, Wolberger C. 2009. Evidence for Bidentate Substrate Binding as the Basis for

the K48 Linkage Specificity of Otubain 1. *J Mol Biol* **386**:1011–1023.

doi:10.1016/j.jmb.2008.12.085

Wasilko DJ, Huang Q, Mao Y. 2018. Insights into the ubiquitin transfer cascade catalyzed by the

Legionella effector SidC. *Elife* **7**:e36154. doi:10.7554/eLife.36154

Wu EL, Cheng X, Jo S, Rui H, Song KC, Dávila-Contreras EM, Qi Y, Lee J, Monje-Galvan V,

Venable RM, Klauda JB, Im W. 2014. CHARMM-GUI Membrane Builder toward realistic

biological membrane simulations. *J Comput Chem* **35**:1997–2004. doi:10.1002/jcc.23702

Wu X, Wu D, Lu Z, Chen W, Hu X, Ding Y. 2009. A Novel Method for High-Level Production

of TEV Protease by Superfolder GFP Tag. *J Biomed Biotechnol* **2009**:1–8.

doi:10.1155/2009/591923

Yau R, Rape M. 2016. The increasing complexity of the ubiquitin code. *Nat Cell Biol* **18**:579–

586. doi:10.1038/ncb3358

Zimmermann L, Stephens A, Nam S-Z, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas

AN, Alva V. 2018. A Completely Reimplemented MPI Bioinformatics Toolkit with a New

HHpred Server at its Core. *J Mol Biol* **430**:2237–2243. doi:10.1016/j.jmb.2017.12.007

Figure 1

a

Effector proteins
(Icm/Dot secretion signal > 2.0) → # of proteins: 305

↓

HHM-HHM prediction
(Uncharacterized proteins) → 4

↓

DUB Activiy validation → 2

↓

Novel OTU-like DUB

b

Icm/Dot
secretion signal

(3.092) **Lpg1621** — OTU (11–283), 439

(7.167) **Lpg2529** — OTU (7–325), 572

(2.121) **Lpg2411** — UCH (110–215), 275

(8.108) **Lpg2907** — CE Clan (117–384), 420

c

Lpg1621 — OTU (11–283), 439

Ub_2
Time: K6 (5 30 90), K11 (5 30 90), K27 (5 30 90), K29 (5 30 90), K33 (5 30 90), K48 (5 30 90), K63 (5 30 90), M1 (5 30 90) (min)

Ub_2
Ub

d

K6, K48, K63

e

Lpg2529 — OTU (7–325), 572

Ub_2
Time: K6 (5 30 90), K11 (5 30 90), K27 (5 30 90), K29 (5 30 90), K33 (5 30 90), K48 (5 30 90), K63 (5 30 90), M1 (5 30 90) (min)

Ub_2
Ub

f

K6, K11, K33, K48, K63

Figure 2

Figure 3

Figure 3 — panels a and b: domain diagrams of LotB and LotC with OTU domains and Ub₂ immunoblot panels for K6, K11, K27, K29, K33, K48, K63, M1.

a LotB 11 OTU 283 439

b LotC 7 OTU 310 572

Ub₂ / Ub blots labeled K6 K11 K27 K29 K33 K48 K63 M1

c LotB — Extra helices — ★ Catalytic triad
CCHF- / Farallon- viral OTU; OTUD2 / OTUD3 human OTUD; helical arm

d LotC — Extra helices — ★ Catalytic triad
Taggert- / DGK Nairo- viral OTU; Otubain1 / Otubain2 human Otubain; helical arm

★ Catalytic triad

e

	Cys loop		Helical lobe		Variable loop		His loop	
LotB	SGAFDNC	29			213 LAKPQVKLSYI		IDALQGHY	271
vCCHF-OTU	QPGDGNC	40			94 LSDNEWGSTLE		LHSGQTHF	152
vFarallon-OTU	VKGDGNC	37			117 ATKEGYWGGTD		IVRCGAHF	175
hOTUD2	VPADNSC	160			213 IKRDDTWGGAI		LIYDGIHY	268
hOTUD3	VPGDGNC	76			127 LAKPGTFAGND		AYRYGEHY	183
LotC	NSGKGNC	24			235 LRREFVWGSEE		NNEGNVHW	305
vTaggert-OTU	QPMDGNC	40			93 IMKPNEWGSSL		IMHRRNHY	151
vDGK-OTU	VPGDGNC	42			94 LRNYYWGGSVE		LKLCQGHI	151
hOtubain1	TRPDGNC	91			208 VEPMCKESDHI		LLYRPGHY	266
hOtubain2	TKGDGNC	51			168 VEPMATECDHI		LLYKTSHY	225
LotA	ATGDGAC	13	Unknown structure		193 AFQRGYSCDEL		LENQEEHW	238
LotA	NPGLGNC	303	Unknown structure		482 ITQDFFWGTHL		TNSNNTHW	536

Figure 4

Figure 5

Supplementary Figure 1

a

b

Whole cell lysate (HEK293)

IB: Ub (P4D1)

IB: GAPDH

Supplementary Figure 2

Supplementary Figure 3

a

Supplementary Figure 4

a

Supplementary Figure 5

Supplmentary Figure 6 - Original blots

Fig. 1c; LotB Di-Ub2

Fig. 1c; LotB Di-Ub2

Fig. 1e; LotC Di-Ub2

Fig. 1e; LotC Di-Ub2

Fig. 2b; LotB catalytic muts

Fig. 2c; LotC catalytic muts

Fig. 2d; LotB ABP

Fig. 2e; LotC ABP

Fig. 2f; LotB ABP

Fig. 2g; LotC ABP

Sup Fig. 2a; LotB ABP

Sup Fig. 2b; LotB ABP

Fig. 3a; LotB WT

Fig. 3a; LotB 300

Fig. 3a; LotB 350

Fig. 3a; LotB 365

Fig. 3b; LotC WT

Fig. 3b; LotC 7

Fig. 3b; LotC 14

Fig. 3b; LotC 21

Fig. 4e; LotB mutants

Fig. 4f; LotC mutants