Cell-free DNA Tissues-of-Origin Profiling to Predict Graft versus Host Disease and Detect Infection after Hematopoietic Cell Transplantation

Authors: Alexandre Pellan Cheng[1], Matthew Pellan Cheng[2,3], Joan Sesing Lenz[1], Kaiwen Chen[2,3], Philip Burnham[4], Kaitlyn Marie Timblin[2,3], José Luis Orejas[2,3], Emily Silverman[2,3], Francisco M. Marty[2,3], Jerome Ritz[2,5], Iwijn De Vlaminck[1],*

Affiliations:
1 Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA
2 Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA
3 Division of Infectious Disease, Brigham and Women's Hospital, Boston, MA, USA
4 Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, USA
5 Department of Medicine, Harvard Medical School, Boston, MA, USA
* Corresponding author (vlaminck@cornell.edu)

ABSTRACT: Allogeneic hematopoietic cell transplantation (HCT) provides effective treatment for hematologic malignancies and immune disorders. Monitoring for immune complications and infection is a critical component of post-HCT therapy, however, current diagnostic options are limited. Here, we propose a blood test that employs genome-wide profiling of methylation marks comprised within circulating cell-free DNA to trace the tissues-of-origin of cell-free DNA (cfDNA), to quantify tissue-specific injury, and to screen for microbial pathogens after allogeneic HCT. We applied this assay to 106 plasma samples collected from 18 HCT recipients at predetermined time points before and after allogeneic HCT. We observe marked dynamics in the composition and abundance of cfDNA from different tissues in response to conditioning chemotherapy and HCT. We find that the abundance of solid-organ derived cfDNA in the blood at one-month after HCT is an early predictor of acute graft-versus-host disease, a frequent immune-related complication of HCT that occurs when donor immune cells attack the patient's own tissues (area under the curve, 0.9, p-value = 0.012). Metagenomic profiling of cfDNA was possible from the same assay and revealed the frequent occurrence of viral and bacterial infection in this patient population. This proof-of-principle study shows that cfDNA has the potential to improve the care of allogeneic HCT recipients by enabling earlier detection and better prediction of immune complications and infection after HCT.

INTRODUCTION

More than 30,000 patients undergo allogeneic hematopoietic cell transplants (HCT) worldwide each year for treatment of a variety of malignant and nonmalignant hematologic diseases[1–3]. However, immune related complications occur frequently after HCT. Up to 50% of patients experience graft-versus-host disease (GVHD) in the first year after transplantation. GVHD occurs when donor immune cells attack the patient's own tissues[2,4–6]. Early and accurate diagnosis of GVHD is critical to inform treatment decisions and to prevent serious long-term complications, including organ failure and death. Unfortunately, there are few, noninvasive diagnostic options that reliably identify patients very early after onset of GVHD symptoms. In current clinical practice, diagnosis of GVHD relies almost entirely on clinical criteria and often requires confirmation with invasive procedures, such as a biopsy of the gastrointestinal tract, skin, or liver[7].

Small fragments of cell-free DNA (cfDNA) circulate in blood. In the absence of disease, cfDNA originates primarily from apoptosis of cells of the hematopoietic lineage[8]. During disease, a significant proportion of cfDNA can be derived from affected tissues[9–15]. In solid-organ transplantation (SOT), we and others have shown that transplant donor derived cfDNA in the blood is a quantitative noninvasive marker of solid organ transplant injury[12,13,16,17]. Here, we sought to investigate the utility of circulating cfDNA as a minimally invasive analyte to detect and quantify tissue injury due to GVHD after HCT. To quantify the contributions of any tissue to the mixture of cfDNA in plasma, and thereby the degree of injury to any vascularized tissue in the setting of HCT, we implemented shallow, whole genome bisulfite sequencing (WGBS) to profile 5-methylcytosine (5mC) marks of cfDNA. These marks are cell, tissue and organ type specific[18], and can inform the tissues-of-origin of cell-free DNA[8,11,19].

We collected serial blood samples from a prospective cohort of allogeneic HCT recipients at predetermined time points before and after HCT. We analyzed a total of 106 plasma samples from 18 HCT recipients with and without GVHD in the first 3 months post HCT. We observed rich dynamics in the tissue-origin of cfDNA in response to pre-transplant conditioning chemotherapy and following HCT. The tissue-origin of cfDNA after transplantation was patient-specific and a function of the manifestation of GVHD. For example, the absolute abundance of organ derived cfDNA at one month after transplantation was significantly elevated for patients diagnosed with GVHD within the first 6 months after HCT, indicating that cfDNA tissues-of-origin profiling can be used to predict GVHD and quantify its severity.

Allogeneic HCT recipients receive therapeutic immunosuppression to manage the risk of GVHD and are consequently at risk of opportunistic infections. Because a wide range of microorganisms can cause disease after HCT, there is a critical need for tools that can broadly inform microbial pathogens. To address this, we investigated the possibility to identify infectious agents via whole-genome bisulfite sequencing of plasma cfDNA, using methods that we previously developed to screen for urinary tract infection[15]. We observe significant increases in viral and bacterial derived cfDNA post HCT. cfDNA from anelloviruses, cytomegalovirus (CMV), herpesvirus 6, Epstein-Barr virus, and polyomavirus was frequently observed in this patient population.

Together, this study provides a proof of principle that cell-free DNA profiling can be used to simultaneously monitor immune and infectious related complications after allogeneic HCT.

RESULTS

We performed a prospective cohort study to evaluate the utility of cfDNA to predict and monitor complications after allogeneic HCT. For this study, we selected 18 adults that underwent allogeneic HCT and assayed a total of 106 serial plasma samples collected at six predetermined time points, including before conditioning chemotherapy, on the day of but before hematopoietic cell infusion, after neutrophil engraftment (>500 neutrophils per microliter), and at one, two, and three months post HCT (**Fig. 1a**). The test cohort included patients with both malignant (n=14) and non-malignant blood disorders (n=4) (**supplementary table 1**). In total, nine patients developed acute GVHD (GVHD+) and nine did not (GVHD-), two developed a bloodstream infection and two developed BK virus disease (see Methods and SI).

We isolated cfDNA from plasma (0.5mL-1.9mL per sample) and implemented whole-genome bisulfite sequencing to profile cytosine methylation marks comprised within cfDNA (**Fig. 1b**). We implemented a single-stranded DNA (ssDNA) library preparation to obtain sequence information after bisulfite conversion[20,21]. This ssDNA library preparation avoids degradation of adapter-bound molecules which is common for WGBS library preparations that rely on ligation of methylated adapters before bisulfite conversion and avoids amplification biases inherent to WGBS library preparations that implement random priming[22]. We obtained 41 ± 15 million paired-end reads per sample, corresponding to 0.9 ± 0.3 fold per-base human genome coverage (**Fig. 1c**) and achieved a high bisulfite conversion efficiency (99.4% $\pm 0.5\%$, **Fig. 1d**). We used paired-end read mapping to characterize the length of bisulfite treated cfDNA at single-nucleotide resolution and to investigate potential degradation of cfDNA due to bisulfite treatment. This analysis revealed a fragmentation profile similar to the fragmentation profile previously reported for plasma cfDNA that was not subjected to bisulfite treatment[23]. The mode of fragments longer than 100bp was 165 bp $\pm$ 7 bp (**Fig. 1e**), and Fourier analysis revealed a 10.4 bp periodicity in the fragment length profile (**Fig. 1e inset**). A second peak at 60-90 bp in the fragment length profile is characteristic of single-stranded library preparation methods and was reported previously[21,24]. Overall, we do not find evidence of significant cfDNA fragmentation due to bisulfite treatment, in line with a recent report[25].

Temporal dynamics in response to conditioning therapy and HCT

To quantify the relative proportion of cfDNA derived from different vascularized tissues and hematologic cell types we analyzed cfDNA methylation profiles against a reference set of methylation profiles of pure cell and tissue types (138 reference tissues, see Methods, **Fig. 1f** and **supplementary dataset 1**)[26–30]. We computed the absolute concentration of tissue-specific cfDNA by multiplying the proportion of tissue-specific cfDNA with the concentration of total host-derived cfDNA (Methods). Figures 1g,h and 2 summarize these measurements for all patients and time points and reveal rich dynamics in tissue-origin of cfDNA in response to both conditioning chemotherapy and HCT (**Fig. 1g-h, Fig. 2**). The most striking features seen in the data include *i*) a decrease in blood-cell specific cfDNA in response to conditioning therapy performed to deplete the patient's own immune cells, as expected (**Fig 1g**, **Fig 2a**), *ii*) an increase in total cfDNA concentration at engraftment (**Fig. 1h**, **Fig. 2b**), *iii*) a decrease in total cfDNA concentration after 60 days for most patients (**Fig. 1h**), and *iv*) an association between tissue-specific cfDNA and the incidence of GVHD (see statistical analysis below).

We next examined these features in more detail to explore the utility of these measurements to monitor immune related complications of HCT (**Fig. 2**). Prior to conditioning, neutrophils, erythrocyte progenitors and monocytes were the major contributors of cfDNA in plasma (23.0%, 12.3% and 11.7%, respectively, average cfDNA concentration 272 $\pm$ 305 ng/mL plasma). A variety of HCT conditioning regimens have been developed with varying degrees of organ toxicity and myelosuppression. The majority of patients in our cohort received reduced intensity conditioning therapy (RIC, n=17), whereas a single patient received myeloablative conditioning therapy (id number 005). Comparison of cfDNA tissues-of-

origin in plasma before and after conditioning showed a significant drop in blood-derived cfDNA as expected from the function of the conditioning therapy (mean proportion of hematopoietic cell cfDNA decreased from 79% ± 10% to 59% ± 20%, p-value=0.0014, **Fig. 2a**). The proportion of blood-derived cfDNA increased to 84% ± 10% at engraftment (p-value = 5.6×10^{-5}, **Fig. 2a**). The most notable effect of stem cell infusion and engraftment was a significant increase in the absolute concentration of cfDNA (mean human-derived cfDNA concentration from 256 ng/mL on day of transplant to 2149 ng/mL at engraftment [p-value = 0.013], **Fig. 2b**).

Performance analysis

We next evaluated the performance of a cfDNA tissue-of-origin measurement to predict GVHD (**Fig. 2c**). We defined GVHD here as the clinical manifestation of any stage of the disease within the first 6 months post HCT (GVHD+, see Methods). We excluded samples collected after GVHD diagnosis, as these patients received additional GVHD treatment. We found that the concentration of solid-organ specific cfDNA was significantly elevated for patients in the GVHD+ group at engraftment, month 1, 2 and 3 (p-values of 0.17, 0.012, 0.0070, 0.020, respectively), but not at the two pre-transplant time points (p = 0.66 prior to conditioning, and p = 0.80 prior to hematopoietic cell infusion). Receiver operating characteristic analysis of the performance of cfDNA as a predictive marker of GVHD yielded an area under the curve (AUC) of 0.7, 0.9, 0.9 and 0.9 at engraftment and months 1, 2, and 3, respectively. These results support the notion that cfDNA predicts GVHD occurrence as early as one month after HCT (mean solid organ cfDNA of 995 and 50 ng/mL plasma for GVHD+ and GVHD-, respectively; AUC = 0.9, p-value < 0.012, **Fig. 2c**).

To evaluate the ability of this assay to pinpoint the site of incidence of GVHD, we quantified the burden of skin-derived cfDNA in the blood of GVHD negative individuals (n=9) and individuals who developed cutaneous GVHD (n=8). We found that plasma samples from individuals with GVHD had a higher burden of skin-derived cfDNA when compared to samples from individuals who did not develop cutaneous GVHD (mean skin cfDNA of 20.6 ng/mL plasma and 3.2ng/mL plasma, respectively, p-value = 0.015 for samples collected post-transplant and pre-diagnosis, **supplementary fig. 1**). The number of samples from patients diagnosed with hepatic and gastrointestinal GVHD was insufficient to test the performance of the assay to pinpoint GVHD related injury to the liver or gut (n = 1 and n = 3, respectively).

We next studied the response to GVHD treatment for three similar patients for which samples and cfDNA tissues-of-origin analyses were available after GVHD diagnosis (male patients with RIC chemotherapy and similar GVHD diagnosis timepoints). These patients were diagnosed with GVHD between days 28 and 39 post HCT and two plasma samples after diagnosis were available for each patient. The first patient was diagnosed with mild GVHD (cutaneous stage 1, overall grade I; resolved day 98), and the tissue-of-origins of cfDNA followed a similar pattern observed for GVHD negative patients (**Fig. 3a,b**). The second patient was diagnosed with moderate GVHD (cutaneous stage 3, overall grade II; resolved day 137). cfDNA tissue-of-origin profiling identified an increase in solid-organ derived cfDNA after diagnosis (36.5 ng/mL at diagnosis, 199.4 ng/mL and 254.1 ng/mL at months 2 and 3, respectively; **Fig. 3c**). The third patient was diagnosed with severe GVHD (cutaneous stage 4, overall grade IV; unresolved; mortality day 91; **Fig. 3d**). cfDNA tissue-of-origin profiling for samples after diagnosis of this patient revealed an increase in solid-organ derived cfDNA in the blood of this patient despite increasingly potent GVHD treatment (233.8 ng/mL at month 1 and 1217.7 ng/mL at month 2; tacrolimus at month 1, and tacrolimus, sirolimus, ruxolitinib, and glucocorticoids at month 2). These three examples illustrate the potential utility of cfDNA tissue-of-origin profiling to monitor GVHD treatment response and outcome.

Plasma infectome after HCT

It is not only human host cells that shed their DNA into the blood; cfDNA from viruses and bacteria
can be detected in the circulation, providing a means to screen for infection via metagenomic cfDNA
sequencing[14,21,31,32]. This may be a particularly powerful approach in the context of HCT, given the high
incidence of infectious complications, and the broad range of microorganisms that can cause disease in
HCT[33]. To test this concept, we mined all data from all patients for microbial derived sequences. In a
previous study, we found close agreement between the abundance of organisms measured by shotgun
sequencing of untreated and bisulfite-treated cfDNA, despite the reduction in sequence complexity inherent
to bisulfite conversion, confirming the possibility to perform metagenomic cfDNA sequencing by WGBS[15].
To identify microbial-derived cfDNA after WGBS, we first identified and removed host related sequences
and we then aligned the remaining unmapped reads to a set of microbial reference genomes ($0.9 \pm 0.4\%$ of
total reads, Materials and Methods). We implemented a background correction algorithm to remove
contributions due to alignment noise and environmental contamination[34].
200
Using this procedure, we found a significant increase in the burden of cfDNA derived from DNA
viruses after HCT (viral cfDNA biomass 1.4×10^{-3} ng/mL and 1.5×10^{-2} ng/mL, day of transplant and at
month 3 respectively, p-value = 0.0082 **Fig. 4a**), but not in bacterial cfDNA (bacterial cfDNA biomass
9.8×10^{-3} ng/mL and 3.3×10^{-2} ng/mL, day of transplant and at month 3 respectively, p-value = 0.50). We and
others have previously reported a link between the abundance in plasma of *Anelloviridae* and the degree of
immunosuppression in solid-organ transplantation, and HCT[30,33]. In line with these observations, the
increase in cfDNA derived from DNA viruses was largely due to an increase in the burden of *Anelloviridae*
cfDNA in the first months after HCT (**Fig. 4b**)[35]. *Herpesviridae* and *Polyomaviridae* frequently establish
latent infection in adults and may reactivate after allogeneic HCT[36]. We identified cfDNA from Human
Herpesviridae and *Polyomaviridae* in 35 of 106 samples from 15 of 18 patients (**Fig. 4c**). In contrast to
Anelloviridae, we did not observe a consistent increase in the burden of cfDNA from these viruses after
HCT (**Fig. 4b**). Our detection of BK polyomavirus is concordant with clinical diagnosis of BK virus disease.
Four individuals in our cohort were diagnosed with BK polyomavirus disease, defined here as the
combination of clinical BK polyomavirus disease symptoms and a positive blood or urine BK PCR test in
the absence of other causes of genitourinary pathology (see Methods). Our metagenomic assay detected
BK polyomavirus cfDNA in the plasma of 4 out of 5 patients with a positive BK blood test (Spearman's
rho 0.79, p-value $<2.2 \times 10^{-16}$), but did not for patients with a negative blood test but a positive urine test,
(n=15).
219
We identified cfDNA from 7 different genera of bacteria (10 species, **supplementary fig. 2**).
Interestingly, all of the identified species are well documented intestinal commensal organisms, in
agreement with results from a recent study by Armstrong *et al.*[37] that suggest a loss of the integrity of the
gut vascular barrier associated with GVHD (**supplementary fig. 2**). For a single patient with unresolved
stage IV skin GVHD we identified a potential bloodstream infection with *Klebsiella pneumoniae*. Two
patients in this cohort developed a clinically diagnosed Streptococcus bloodstream infection within the first
6 months post-transplant. We did not detect Streptococcus cfDNA by metagenomic cfDNA sequencing for
these two patients, potentially because the infection timepoints were at least six days away from the nearest
plasma collection time point and bloodstream infections with Streptococcus species rapidly clear after the
initiation of antimicrobial treatment[38].
230
231   **DISCUSSION**
232
We have described a cfDNA assay with the potential to detect both GVHD related injury and
infection after allogeneic HCT. This work was inspired by recent studies that have shown that transplant
donor specific cfDNA in the blood of solid-organ transplant recipients is a quantitative marker of solid-
organ transplant injury[12,13]. A variety of commercial assays to quantify donor cfDNA in solid organ
transplant recipients are already available[16,39–41], and studies are ongoing to test the utility of donor cfDNA
to screen for allograft rejection, injury and failure after kidney, liver, heart, pancreas and lung

transplantation[42]. We reasoned that cfDNA may also inform injury to vascularized tissues due to GVHD
after HCT. To quantify cfDNA derived from any tissue, we implemented bisulfite sequencing of cfDNA,
to profile cytosine methylation marks that are comprised within cfDNA and that are cell, tissue and organ
type specific. Several other epigenetic marks, including hydroxymethylation[43] and histone modifications[44],
can inform the tissues-of-origin of cfDNA, and profiling of these marks may also be useful to monitor
GVHD after HCT.
245
In recent years, several protein biomarkers have been investigated for the diagnosis of GVHD.
Proteomic approaches have yielded candidate markers that are not directly involved in the pathogenesis of
GVHD, but that are secreted as a result of end-organ damage[46,48]. ST2 and REG3α, which both derive from
the gastrointestinal tract, are two such biomarkers with the strongest predictive power. The cfDNA assay
presented here may provide inherent advantages over protein biomarker technologies. First, because the
concentration of tissue-specific DNA can be directly related to the degree of cellular injury, this assay is
easy to interpret, and offers a measure of injury that can in principle be followed over time. Second, the
cfDNA assay explored here provides a generalizable approach to measure injury to any tissue, whereas
protein injury markers may not be available for all cell and tissue types. Third, this assay is compatible with
a variety of quantitative nucleic acid measurement technologies, including digital and quantitative PCR and
DNA sequencing. Fourth, this assay does not depend on antibodies, which come with challenges of
specificity and reproducibility[49].
258
Whole genome bisulfite sequencing is not only responsive to human host derived cfDNA, but also
to microbial cfDNA that may be present in the blood circulation. Several recent studies have demonstrated
the value of cfDNA metagenomic sequencing to broadly detect infection in a variety of clinical settings,
including urinary tract infection[14,15], sepsis[50] and invasive fungal disease[51]. In HCT, sequencing of cell-free
DNA has been used to identify pathogens in blood before clinical onset of bloodstream infections[52]. We
have previously shown that bisulfite sequencing of cfDNA is compatible with metagenomic analyses,
despite the reduction in sequence complexity due to bisulfite conversion of all unmethylated cytosines[15].
Here, we investigated the potential to screen for microbial and viral derived cfDNA in plasma of HCT
recipients via bisulfite sequencing of cfDNA. BK virus cfDNA was detected using this approach for
samples that were BK virus positive in blood, but not for those that were only BK virus positive in urine.
In addition, while we note increased herpesvirus genomic abundance, during immunosuppression,
differentiating lytic and latent infection and the corresponding host response could be vital in improving
patient care. The assay reported here therefore has the potential to simultaneously inform about GVHD,
from the tissues-of-origin of host cfDNA, and infection, from metagenomic analysis of microbial cfDNA.
Compared to conventional metagenomic sequencing, this assay requires one additional experimental step
to bisulfite convert cfDNA, which can be completed within approximately 2 hours and is compatible with
multiple existing next-generation sequencing workflows.
276
277   This is a proof-of-principle study with several limitations that need to be addressed in future work.
278 The scope of the current study with 18 patients was not powered to detect any association of cfDNA with
279 acute GVHD involving organs other than skin (liver, gut). Our results suggest that cfDNA tissue-of-origin
280 profiling is predictive of acute GVHD, but larger studies will be needed to extend the current observations
281 to other sites of organ damage, and to assess its utility in detecting and diagnosing chronic GVHD.
282
283
284 **MATERIALS AND METHODS**
285
Study cohort. We performed a nested case-control study within a prospective cohort of adult patients
undergoing allogeneic HCT at Dana-Farber Cancer Institute. Patients were followed for 6 months after
HCT. Patients were selected for this study on a rolling basis, and were placed in the GVHD case or control
groups based on clinical manifestation of the disease within the first 6 months after HCT. Individuals were

excluded from the study if they did not provide blood samples for at least 5 of the 6 studied time points
(pre-conditioning, day of transplant, engraftment, months 1, 2 and 3). The study was approved by the Dana-
Farber/Harvard Cancer Center's Office of Human Research Studies. All patients provided written informed
consent.
294
For this study, we used 106 blood samples collected from 18 allogeneic HCT recipients from August 2018-
to April -2019. Baseline patient characteristics were recorded. Covariates of interest included HLA
matching, donor relatedness and donor-recipient sex mismatch (**supplementary table 1**). Date of onset of
GVHD, as well as GVHD prophylaxis and treatment regimens were documented. GVHD was diagnosed
clinically and pathologically. GVHD severity was graded according to the Glucksberg criteria[43]. Other
clinical events of interest included the development of bloodstream infections, BK polyomavirus disease,
and clinical disease from other DNA viruses.
302
303 **Engraftment.** Neutrophil engraftment was considered when blood samples contained an absolute
304 neutrophil count greater or equal than 500 cell per microliter of blood on two separate measurements.
305
306 **BK polyomavirus disease identification.** Patients were identified as BK virus disease positive when they
307 presented BK-related urinary symptoms that correlated with positive BK qPCR test in either urine or blood
308 ($>10^5$ copies/ mL in urine, >0 copies/mL in blood; Viracor BK qPCR test, reference #2500) and did not
309 have evidence of any other cause of genitourinary pathology at the time of symptom onset.
310
Blood sample collection and plasma extraction. Blood samples were collected through standard
venipuncture in EDTA tubes (Becton Dickinson (BD), reference #366643) on admission, before the
beginning of the conditioning chemotherapy; on the day of HCT after the completion of the conditioning
chemotherapy, at engraftment (usually 14 to 21 days after HCT), and at months 1, 2, and 3 post-HCT.
Plasma was extracted through blood centrifugation (2000rpm for 10 minutes using a Beckman Coulter
Allegra 6R centrifuge) and stored in 0.5-2mL aliquots at −80 °C. Plasma samples were shipped from DFCI
to Cornell University on dry ice.
318
319 **Nucleic acid control preparation.** Synthetic oligos were prepared (IDT, **supplementary table 2**), mixed
320 in equal proportions, and diluted at approximately 150 ng/ul. At the time of cfDNA extraction, 8ul of control
321 was added to 1992μL of 1xPBS and processed as a sample in all downstream experiments.
322
323 **Cell-free DNA extraction.** cfDNA was extracted according to manufacturer recommendations (Qiagen
324 Circulating Nucleic Acid Kit, reference #55114, elution volume 45μl). Eluted DNA was quantified using a
325 Qubit 3.0 Fluorometer (using 2μL of eluted DNA). Measured cfDNA concentration was obtained using the
326 following formula:
327

$$cfDNA\ concentration = \frac{(Eluted\ cfDNA\ concentration) * (Elution\ volume)}{(Plasma\ volume)}$$

329
Whole-genome bisulfite sequencing. cfDNA and nucleic acid controls were bisulfite treated according to
manufacturer recommendations (Zymo Methylation Lightning Kit, reference #D5030). Sequencing
libraries were prepared using a previously described single-stranded library preparation protocol[21].
Libraries were quality-controlled through DNA fragment analysis (Agilent Fragment analyzer) and
sequenced on an Illumina NextSeq550 machine using 2x75bp reads. Nucleic acid controls were sequenced
a ~1% of the total sequencing lane.
336
337 **Human genome alignment.** Adapter sequences were trimmed using BBTools[53]. The Bismark[54] alignment
338 tool was used to align reads to the human genome (version hg19), remove PCR duplicates and calculate
339 methylation densities.

Reference tissue methylation profiles and tissue of origin measurement. Reference tissue methylomes were obtained from publicly available databases[26–30] (**supplementary dataset 1**). Genomic coordinates from different sources were normalized and converted to a standard 4 column bed file (columns: chromosome, start, end, methylation fraction) using hg19 assembly coordinates. Methylation profiles were grouped by tissue-type and differentially methylated regions were found using Metilene[55]. Tissues and cell-types of origin were determined using quadratic programming as previously described[15].

Metagenomic alignment and quantification of microbial cfDNA. After WGBS, reads were adapter-trimmed using BBTools[44], and short reads are merged with FLASH[56]. Sequences were aligned to a C-to-T converted genome using Bismark[54]. Unmapped reads were BLASTed[57] using hs-blastn[58] to a list of C-to-T converted microbial reference genomes. A relative abundance of all detected organisms was determined using GRAMMy[59], and relative genomic abundances are measured as previously described[32]. Microbial cfDNA fraction was calculated by dividing the unique number of reads mapping to microbial species (after adjusting for the length of each microbial genome in the reference set) to the total number of sequenced reads. Human fraction is estimated as 1 - microbial fraction. Microbial species were then filtered for environmental contamination and alignment noise using previously described methods (LBBC, PNAS). For viral species identification (**Fig. 4c**), custom genomic reference was generated from representative genomes for BK polyomavirus (NC_001538.1), cytomegalovirus (NC_006273.1), herpesvirus 6A (NC_001664.4) and 6B (NC_00898.1), human polyomavirus 6 (NC_014406.1) and 7 (NC_014407.1) and ebstein-barr virus (NC_007605). Reads that BLASTed to these species were then re-aligned via Bismark, and a threshold of 1 mapped sequence per 40 million total sequenced reads was used to positively identify a viral species within a plasma sample.

cfDNA concentration. cfDNA concentration of a specific tissue or microbe is calculated as follows:

$$Normalized\ cfDNA\ concentration$$
$$= \frac{(cfDNA\ concentration) * (Nucleic\ acid\ control\ input\ mass)}{Nucleic\ acid\ control\ output\ mass}$$

$$Tissue\ specific\ cfDNA\ concentration$$
$$= (Normalized\ cfDNA\ concentration) * (human\ read\ fraction)$$
$$* (tissue\ proportion)$$

$$Microbial\ cfDNA\ concentration$$
$$= (Normalized\ cfDNA\ concentration) * (microbial\ read\ fraction)$$

Depth of coverage. The depth of sequencing was measured by summing the depth of coverage for each mapped base pair on the human genome after duplicate removal, and dividing by the total length of the human genome (hg19, without unknown bases).

Bisulfite conversion efficiency. We estimated bisulfite conversion efficiency by quantifying the rate of C[A/T/C] methylation in human-aligned reads (using MethPipe[60]), which are rarely methylated in mammalian genomes.

Statistical analysis. Statistical analysis was performed in R (version 3.5). All tests were performed using a two-sided Wilcoxon test.

Data availability. The sequencing data generated for this study will be available in the database of Genotypes and Phenotypes (dbGaP). All code used to generate figures and analyze primary data is available at www.github.com/alexpcheng/cfDNAme.

389
390 **CONFLICTS OF INTEREST**
391 The authors declare no conflicts of interest.
392
393 **AUTHOR CONTRIBUTIONS**
394 A.P.C., M.P.C., F.M.M., J.R. and I.D.V. designed the study. A.P.C. and J.S.L. performed experiments.
395 M.P.C., F.M.M., J.R., K.C., K.M.T., J.L.O. and E.S. consented patients and acquired clinical data. A.P.C.,
396 M.P.C., P.B., I.D.V., F.M.M. and J.R. analyzed data. A.P.C., M.P.C., I.D.V., F.M.M. and J.R. wrote the
397 manuscript. All authors reviewed and approved the manuscript.
398
ACKNOWLEDGEMENTS
We would like to thank the Cornell Genomics Center for help with sequencing assays, the Cornell
Bioinformatics facility for computational assistance, The Pasquarello Tissue bank at the Dana-Farber
Cancer Institute for sample processing and cryopreservation and members of the De Vlaminck Lab for
helpful discussions. We thank Francoise Vermeulen of the Cornell Statistical Consulting Unit for helpful
discussion. This work was supported by R01AI146165 (to I.D.V, J.R, M.P.C. and F.M.), R21AI133331 (to
I.D.V.), R21AI124237 (to I.D.V.), DP2AI138242 (to I.D.V.), a National Sciences and Engineering
Research Council of Canada fellowship PGS-D3 (to A.P.C).

REFERENCES

1. Gratwohl, A. *et al.* Hematopoietic stem cell transplantation: a global perspective. *JAMA* **303**, 1617–1624 (2010).

2. McDonald-Hyman, C., Turka, L. A. & Blazar, B. R. Advances and challenges in immunotherapy for solid organ and hematopoietic stem cell transplantation. *Sci. Transl. Med.* **7**, 280rv2 (2015).

3. Niederwieser, D. *et al.* Hematopoietic stem cell transplantation activity worldwide in 2012 and a SWOT analysis of the Worldwide Network for Blood and Marrow Transplantation Group including the global survey. *Bone Marrow Transplant.* **51**, 778–785 (2016).

4. Paczesny, S. Discovery and validation of graft-versus-host disease biomarkers. *Blood* **121**, 585–594 (2013).

5. Ferrara, J. L. M. & Deeg, H. J. Graft-versus-Host Disease. *N. Engl. J. Med.* **324**, 667–674 (1991).

6. Ferrara, J. L. M. & Chaudhry, M. S. GVHD: biology matters. *Blood Adv.* **2**, 3411–3417 (2018).

7. Vogelsang, G. B., Lee, L. & Bensen-Kennedy, D. M. Pathogenesis and Treatment of Graft-Versus-Host Disease After Bone Marrow Transplant. *Annu. Rev. Med.* **54**, 29–52 (2003).

8. Sun, K. *et al.* Plasma DNA tissue mapping by genome-wide methylation sequencing for noninvasive prenatal, cancer, and transplantation assessments. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E5503–E5512 (2015).

9. Mouliere, F. *et al.* Detection of cell-free DNA fragmentation and copy number alterations in cerebrospinal fluid from glioma patients. *EMBO Mol. Med.* **10**, e9323 (2018).

10. Mouliere, F. *et al.* Enhanced detection of circulating tumor DNA by fragment size analysis. *Sci. Transl. Med.* **10**, (2018).

11. Lehmann-Werman, R. *et al.* Identification of tissue-specific cell death using methylation patterns of circulating DNA. *Proc. Natl. Acad. Sci.* **113**, E1826–E1834 (2016).

12. Vlaminck, I. D. *et al.* Circulating Cell-Free DNA Enables Noninvasive Diagnosis of Heart Transplant Rejection. *Sci. Transl. Med.* **6**, 241ra77-241ra77 (2014).

13. De Vlaminck, I. *et al.* Noninvasive monitoring of infection and rejection after lung transplantation. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13336–13341 (2015).

14. Burnham, P. *et al.* Urinary cell-free DNA is a versatile analyte for monitoring infections of the urinary tract. *Nat. Commun.* **9**, 1–10 (2018).

15. Cheng, A. P. *et al.* A cell-free DNA metagenomic sequencing assay that integrates the host injury response to infection. *Proc. Natl. Acad. Sci.* **116**, 18738–18744 (2019).

16. Grskovic, M. *et al.* Validation of a Clinical-Grade Assay to Measure Donor-Derived Cell-Free DNA in Solid Organ Transplant Recipients. *J. Mol. Diagn. JMD* **18**, 890–902 (2016).

17. Sharon, E. *et al.* Quantification of transplant-derived circulating cell-free DNA in absence of a donor genotype. *PLOS Comput. Biol.* **13**, e1005629 (2017).

18. Lokk, K. *et al.* DNA methylome profiling of human tissues identifies global and tissue-specific methylation patterns. *Genome Biol.* **15**, 3248 (2014).

19. Moss, J. *et al.* Comprehensive human cell-type methylation atlas reveals origins of circulating cell-free DNA in health and disease. *Nat. Commun.* **9**, 1–12 (2018).

20. Gansauge, M.-T. & Meyer, M. Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nat. Protoc.* **8**, 737–748 (2013).

21. Burnham, P. *et al.* Single-stranded DNA library preparation uncovers the origin and diversity of ultrashort cell-free DNA in plasma. *Sci. Rep.* **6**, 27859 (2016).

22. Miura, F., Enomoto, Y., Dairiki, R. & Ito, T. Amplification-free whole-genome bisulfite
sequencing by post-bisulfite adaptor tagging. *Nucleic Acids Res.* **40**, e136 (2012).
23. Jiang, P. & Lo, Y. M. D. The Long and Short of Circulating Cell-Free DNA and the Ins and Outs
of Molecular Diagnostics. *Trends Genet.* **32**, 360–371 (2016).
24. Snyder, M. W., Kircher, M., Hill, A. J., Daza, R. M. & Shendure, J. Cell-free DNA comprises an
in vivo nucleosome footprint that informs its tissues-of-origin. *Cell* **164**, 57–68 (2016).
25. Werner, B. *et al.* Circulating cell-free DNA from plasma undergoes less fragmentation during
bisulfite treatment than genomic DNA due to low molecular weight. *PLOS ONE* **14**, e0224338 (2019).
26. ENCODE Project Consortium. The ENCODE (ENCyclopedia Of DNA Elements) Project.
Science **306**, 636–640 (2004).
27. Bujold, D. *et al.* The International Human Epigenome Consortium Data Portal. *Cell Syst.* **3**, 496-
499.e2 (2016).
28. Albrecht, F., List, M., Bock, C. & Lengauer, T. DeepBlue epigenomic data server: programmatic
data retrieval and analysis of epigenome region sets. *Nucleic Acids Res.* **44**, W581–W586 (2016).
29. Bernstein, B. E. *et al.* The NIH Roadmap Epigenomics Mapping Consortium. *Nat. Biotechnol.*
28, 1045–1048 (2010).
30. Fernández, J. M. *et al.* The BLUEPRINT Data Analysis Portal. *Cell Syst.* **3**, 491-495.e5 (2016).
31. Kowarsky, M. *et al.* Numerous uncharacterized and highly divergent microbes which colonize
humans are revealed by circulating cell-free DNA. *Proc. Natl. Acad. Sci.* (2017)
doi:10.1073/pnas.1707009114.
32. De Vlaminck, I. *et al.* Temporal response of the human virome to immunosuppression and
antiviral therapy. *Cell* **155**, 1178–1187 (2013).
33. Tomblyn, M. *et al.* Guidelines for Preventing Infectious Complications among Hematopoietic
Cell Transplant Recipients: A Global Perspective. *Biol. Blood Marrow Transplant. J. Am. Soc. Blood*
Marrow Transplant. **15**, 1143–1238 (2009).
34. Burnham, P. *et al.* Separating the signal from the noise in metagenomic cell-free DNA
sequencing. *bioRxiv* 734756 (2019) doi:10.1101/734756.
35. Schmitz, J. *et al.* The Value of Torque Teno Virus (TTV) as a Marker for the Degree of
Immunosuppression in Adult Patients after Hematopoietic Stem Cell Transplantation (HSCT). *Biol.*
Blood Marrow Transplant. (2019) doi:10.1016/j.bbmt.2019.11.002.
36. Marr, K. A. Delayed opportunistic infections in hematopoietic stem cell transplantation patients:
a surmountable challenge. *Hematol. Educ. Program Am. Soc. Hematol. Am. Soc. Hematol. Educ.*
Program **2012**, 265–270 (2012).
37. Armstrong, A. E. *et al.* Cell-free DNA next-generation sequencing successfully detects infectious
pathogens in pediatric oncology and hematopoietic stem cell transplant patients at risk for invasive fungal
disease. *Pediatr. Blood Cancer* **66**, e27734 (2019).
38. Cheng, M. P. *et al.* Blood Culture Results Before and After Antimicrobial Administration in
Patients With Severe Manifestations of Sepsis: A Diagnostic Study. *Ann. Intern. Med.* (2019)
doi:10.7326/M19-1696.
39. Sigdel, T. K. *et al.* A rapid noninvasive assay for the detection of renal transplant injury.
Transplantation **96**, 97–101 (2013).
40. Beck, J. *et al.* Digital droplet PCR for rapid quantification of donor DNA in the circulation of
transplant recipients as a potential universal biomarker of graft injury. *Clin. Chem.* **59**, 1732–1741 (2013).

494 41. Gielis, E. M. *et al.* Cell-Free DNA: An Upcoming Biomarker in Transplantation. *Am. J.*
495 *Transplant. Off. J. Am. Soc. Transplant. Am. Soc. Transpl. Surg.* **15**, 2541–2551 (2015).

496 42. Knight, S. R., Thorne, A. & Lo Faro, M. L. Donor-specific Cell-free DNA as a Biomarker in
497 Solid Organ Transplantation. A Systematic Review. *Transplantation* **103**, 273–283 (2019).

498 43. Song, C.-X. *et al.* 5-Hydroxymethylcytosine signatures in cell-free DNA provide information
499 about tumor types and stages. *Cell Res.* **27**, 1231–1242 (2017).

500 44. Sadeh, R. *et al.* ChIP-seq of plasma cell-free nucleosomes identifies cell-of-origin gene
501 expression programs. *bioRxiv* 638643 (2019) doi:10.1101/638643.

502 45. Harris, A. C. *et al.* Plasma biomarkers of lower gastrointestinal and liver acute GVHD. *Blood*
503 **119**, 2960 LP – 2963 (2012).

504 46. Chen, Y.-B. & Cutler, C. S. Biomarkers for acute GVHD: can we predict the unpredictable? *Bone*
505 *Marrow Transplant.* **48**, 755–760 (2012).

506 47. Paczesny, S. Discovery and validation of graft-versus-host disease biomarkers. *Blood* **121**, 585
507 LP – 594 (2013).

508 48. Paczesny, S. *et al.* Three Biomarker Panel at Day 7 and 14 Can Predict Development of Grade II-
509 IV Acute Graft-Versus-Host Disease. *Blood* **116**, 675 LP – 675 (2010).

510 49. Voskuil, J. L. A. The challenges with the validation of research antibodies. *F1000Research* **6**,
511 (2017).

512 50. Blauwkamp, T. A. *et al.* Analytical and clinical validation of a microbial cell-free DNA
513 sequencing test for infectious disease. *Nat. Microbiol.* **4**, 663–674 (2019).

514 51. Hong, D. K. *et al.* Liquid biopsy for infectious diseases: sequencing of cell-free plasma to detect
515 pathogen DNA in patients with invasive fungal disease. *Diagn. Microbiol. Infect. Dis.* **92**, 210–213
516 (2018).

517 52. Goggin, K. P. *et al.* Evaluation of Plasma Microbial Cell-Free DNA Sequencing to Predict
518 Bloodstream Infection in Pediatric Patients With Relapsed or Refractory Cancer. *JAMA Oncol.* (2019)
519 doi:10.1001/jamaoncol.2019.4120.

520 53. Bushnell, B., Rood, J. & Singer, E. BBMerge – Accurate paired shotgun read merging via
521 overlap. *PLOS ONE* **12**, e0185056 (2017).

522 54. Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
523 applications. *Bioinformatics* **27**, 1571–1572 (2011).

524 55. Jühling, F. *et al.* metilene: fast and sensitive calling of differentially methylated regions from
525 bisulfite sequencing data. *Genome Res.* **26**, 256–262 (2016).

526 56. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome
527 assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

528 57. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search
529 tool. *J. Mol. Biol.* **215**, 403–410 (1990).

530 58. Chen, Y., Ye, W., Zhang, Y. & Xu, Y. High speed BLASTN: an accelerated MegaBLAST search
531 tool. *Nucleic Acids Res.* **43**, 7762–7768 (2015).

532 59. Xia, L. C., Cram, J. A., Chen, T., Fuhrman, J. A. & Sun, F. Accurate Genome Relative
533 Abundance Estimation Based on Shotgun Metagenomic Reads. *PLoS ONE* **6**, (2011).

534 60. Song, Q. *et al.* A Reference Methylome Database and Analysis Pipeline to Facilitate Integrative
535 and Comparative Epigenomics. *PLOS ONE* **8**, e81148 (2013).

FIGURES and CAPTIONS

Figure 1. Study workflow. **a** Blood samples from hematopoietic cell transplant recipients (n=18) were prospectively collected at six time points. **b** WGBS is performed on cfDNA extracted from patient plasma. Sequenced cfDNA is processed through a custom bioinformatics pipeline. **c,d** hg19 sequence coverage (c) and bisulfite conversion efficiency (d) of sequenced cfDNA (n=106). Red lines indicate the median. **e** Fragment length profiles of 106 cfDNA samples after bisulfite treatment. Inset: Fourier analysis reveals a 10.4 bp periodicity in the fragment length profiles of bisulfite treated cfDNA. **f** UMAP dimensional reduction of cell and tissue methylation profiles. Individual tissues are colored by UMAP coordinates using a linear gradient where each of the four corners is either cyan, magenta, yellow or black. **g,h** Tissue cfDNA fraction (g) and tissue cfDNA concentration (h) of 18 patients with and without aGVHD; tissues are colored by their average color provided by the UMAP coordinate-based color scheme. Vertical dashed lines represent engraftment (yellow) and month 1, 2, and 3 (black) timepoints. Red boxes in h are magnified areas of engraftment and month 1 timepoints. **i** Patient GVHD and BK virus disease. GVHD grades and stages are indicated within the individual boxes. Patient identification numbers are listed on the right. Rows in h,g,i are sorted by mortality, overall GVHD grade and GVHD stage.

555
Figure 2. Host-derived cfDNA dynamics before and after HCT. **a,b** Effect of conditioning and HCT
infusion on cfDNA composition (a) and absolute concentration (b). **c** Solid organ derived cfDNA
concentration in plasma. Top row: dark lines represent mean solid-organ cfDNA and days post transplant
for each patient time point. Error Bars represent standard error of the mean. Bottom row: solid organ cfDNA
by time point. Samples are removed from analysis if plasma was collected after aGVHD diagnosis. * p-
value < 0.05; ** p-value < 0.01
562

565 **Figure 3.** Solid organ cfDNA concentration dynamics. **a** Solid organ cfDNA concentration in GVHD
566 negative individuals. **b,c,d** Solid organ cfDNA concentration in three patients. Blue line represents loess-
567 smoothed solid organ cfDNA in GVHD negative patients.
568

569
570 **Figure 4.** Plasma infectome. **a** Microbial cfDNA concentration by time point. **b** Polyomavirus, annelovirus
571 and human herpesvirus abundance in plasma before and after HCT. **c** Human herpesvirus and polyomavirus
572 species detected per patient (n=15) per time point (patients without detectable herpesvirus or human
573 polyomavirus are not shown (n=3)). Error bars represent standard error of the mean.
574

Supplemental Materials for:

Cell-free DNA Tissues-of-Origin Profiling to Predict Graft versus Host Disease and Detect Infection after Hematopoietic Cell Transplantation

Authors: Alexandre Pellan Cheng[1], Matthew Pellan Cheng[2,3], Joan Sesing Lenz[1], Kaiwen Chen[2,3], Philip Burnham[4], Kaitlyn Marie Timblin[2,3], José Luis Orejas[2,3], Emily Silverman[2,3], Francisco M. Marty[2,3], Jerome Ritz[2,5], Iwijn De Vlaminck[1,]*

Affiliations:
1 Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA
2 Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA
3 Division of Infectious Disease, Brigham and Women's Hospital, Boston, MA, USA
4 Department of Bioengineering, University of Pennsylvania, Philadelphia, PA, USA
5 Department of Medicine, Harvard Medical School, Boston, MA, USA
* Corresponding author (vlaminck@cornell.edu)

Supplementary figure 1. Skin-derived cfDNA in patients with aGVHD. a Skin cfDNA concentration for each individual timepoint. Inset: comparison between GVHD negative and skin GVHD positive individuals (all post-transplant collection timepoints aggregated). **b** Receiving operating characteristic area under the curves for each timepoint between GVHD negative and skin GVHD positive individuals (post-diagnosis samples also considered). * p-value <0.05. Error bars represent standard error of the mean.

Supplementary figure 2. Bacterial species detected in HCT patient plasma through WGBS. Each box represents a single patient at each profiled time point.

Supplementary table 1. Clinical information

Age at enrollment -- median (range) (years)	63 (20-73)	**Relation to donor -- no (%)**	
Female sex -- no (%)	7 (38%)	Unrelated	15 (83%)
Race / Ethnicity -- no (%)		Related	3 (17%)
Caucasian	17 (94%)	**Recipient CMV status -- no (%)**	
American Indian/Alaskan Native	1 (6%)	**R+**	10 (56%)
Reason for hematopoietic cell transplant -- no (%)[1]		**R-**	8 (44%)
Acute myeloid leukemia	4 (22%)	**GVHD status -- no (%)[2]**	
Acute lymphocytic leukemia	4 (22%)	Overall grade I	5 (28%)
Mantle cell lymphoma	2 (11%)	Overall grade II	1 (6%)
Myelodysplastic syndrome	2 (11%)	Overall grade III	1 (6%)
T-Cell lymphoma	2 (11%)	Overall grade IV	3 (17%)
Aplastic anemia	1 (6%)	Skin staging I	3 (17%)
Chronic myelomonocytic leukemia	1 (6%)	Skin staging II	2 (11%)
Chronic lymphocytic leukemia	1 (6%)	Skin staging III	1 (6%)
Myelofibrosis	1 (6%)	Skin staging IV	2 (11%)
Paroxysmal nocturnal hemoglobinuria	1 (6%)	Liver staging I	1 (6%)
Source of HCT -- no (%)		Liver staging II	0 (0%)
Peripheral blood	16 (88%)	Liver staging III	0 (0%)
Umbilical cord	1 (6%)	Liver staging IV	0 (0%)
Bone marrow	1 (6%)	Gut staging I	0 (0%)
HLA matching -- no (%)		Gut staging II	1 (6%)
Match	14 (77%)	Gut staging III	1 (6%)
Mismatch	3 (17%)	Gut staging IV	1 (6%)
Haploidentical	1 (6%)	**Conditioning regimen -- no (%)**	
Conditioning regimen -- no (%)		Reduced intensity	17 (94%)
Busulfan, Fluradabine	10 (56%)	Myeloablative	1 (6%)
Cyclophosmamide, Fludarabine, total body irradiation (TBI)	2 (11%)	**Other characteristics -- no (%)**	
Fludarabine, Melphalan	2 (11%)	Mortality	3 (17%)
Clyclophoamamide, TBI	1 (6%)	Previous HCT	0 (0%)
Clyclophosphamide, Fludarabine, TBI, Mesna	1 (6%)	Time to GVHD onset (median ± std)	95 ± 64 days
Cyclophosphamide, Fludarabine, TBI, anti-thymocyte globulin	1 (6%)	T-cell depletion	0 (0%)
Busulfan, Fluradabine, Melphalan	1 (6%)	**GVHD treatment – no (%)**	
GVHD prophylaxis -- no (%)		Glucocorticoids	6 (67%)
Methotrexate, Tacrolimus, Sirolimus	9 (50%)	Jakafi, Glucocorticoids	3 (33%)
Methotrexate, Tacrolimus	5 (28%)		
Mycophenolate mofetil, Tacrolimus, Post-transplant cyclophosphamide	2 (11%)		

Tacrolimus, Sirolimus	1 (6%)		
Mycophenolate mofetil, Tacrolimus	1 (6%)		

1. One individual received an HCT for two blood disorders
2. One individual had two separate incidences of GVHD

Supplementary table 2. Oligonucleotides comprising nucleic acid control.

	Sequence (5'-3')
oligo1	TTTAACGCATAAACATGCGTTTTGGGTAGTGTTTTTTGGAAACACAGATCCGTGCGCACACCTGGTGGAG
oligo2	ATAAACATGCGTTTTGGGTAGTGTTTTTTGGAAACACAGATCCGTGCGCACACCT
oligo3	GCGTTTTGGGTAGTGTTTTTTGGAAACACAGATCCGTGCG
oligo4	GGTAGTGTTTTTTGGAAACACAGAT